# Excitatory: inhibitory imbalance in Alzheimer's disease is exacerbated by seizures with attenuation after rapamycin treatment in 5XFAD mice.

Barbour AJ[1]*, Gourmaud S[1]*, Lancaster E[1], Li X[1], Stewart D[1,2], Irwin DJ[1,3], Talos DM[1], Jensen FE[1]

**\*These authors contributed equally to this work.**

## Abstract

Approximately 22% of Alzheimer's disease (AD) patients suffer from seizures, and the co-occurrence of seizures and epileptiform activity exacerbate AD pathology and related cognitive deficits. Hence seizures may be a targetable component of AD progression. As epileptogenesis is associated with changes in neuronal excitatory: inhibitory (E:I) balance, we hypothesized that decreased markers of inhibition relative to those of excitation would be present in AD patients and exacerbated further when seizures were a comorbidity. We similarly hypothesized that an E:I imbalance would be present in five times familial AD (5XFAD) mice and augmented following pentylenetetrazol (PTZ) seizure kindling. AD temporal cortical tissue from patients with or without seizure history and brain tissue from 5XFAD mice were examined for changes in several markers of E:I balance, including the inhibitory GABA$_A$ receptor, the chloride cotransporters, sodium potassium chloride cotransporter 1 (NKCC1) and potassium chloride cotransporter 2 (KCC2), and the excitatory NMDA and AMPA type glutamate receptors. We found that AD patients had decreased GABA$_A$ receptor subunits and those with comorbid seizures had worsened cognitive and functional scores, and increased in NKCC1/KCC2 ratios, indicative of depolarizing GABA responses. The E:I imbalance appears to occur early in the disease course, as patch clamp recordings from CA1 neurons in hippocampal slices from prodromal 5XFAD mice showed decreased GABAergic inhibitory transmission and increased intrinsic excitability. In addition, seizure induction in prodromal 5XFAD mice further dysregulated NKCC1/KCC2, and altered the excitatory AMPA glutamate receptor protein expression, with a reduction in GluA2 subunit,

indicative of calcium permeable-receptors. Finally, we found that chronic treatment with the mTORC1 inhibitor, rapamycin, at doses we have previously shown to attenuate seizure-induced β-amyloid pathology and cognitive deficits, could reverse aspects of E:I imbalance in these mice. Our data demonstrate novel mechanisms of interaction between AD and epilepsy and indicate that FDA-approved mTOR inhibitors hold therapeutic promise for AD patients with a seizure history.

**Author affiliations:**

1 Department of Neurology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA;

2 Duke University School of Medicine, Durham NC 27708, USA;

3 Penn Frontotemporal Degeneration Center, University of Pennsylvania, Philadelphia, PA, 19104, USA;

Correspondence:

Frances E. Jensen, MD, FACP

Arthur Knight Asbury MD Professor

Chair, Department of Neurology

Hospital of the University of Pennsylvania, 3400 Spruce Street, Dulles 3

Philadelphia, PA 19104, PA USA

E-mail: frances.jensen@pennmedicine.upenn.edu

# Introduction

Seizures can be a comorbidity in Alzheimer's disease (AD) patients, with 10-22% showing clinically identifiable seizures and up to 64% displaying subclinical epileptiform activity[1]. Notably, the co-occurrence of seizures and epileptiform activity in AD patients enhances disease progression and worsens cognitive performance[1-5]. We and others have demonstrated bidirectional interactions with AD and epilepsy sharing similar pathology and convergence upon common underlying cellular mechanisms. Human temporal lobe epilepsy brain tissue shows increased Aβ plaques and increased hyperphosphorylated tau (pTau)[6-8] and the accumulation of these neuropathological proteins in AD animal models are associated with elevated neuronal hyperexcitability and seizure susceptibility [9-14]. Further, we recently demonstrated that a history of seizures is associated with enhanced Aβ and pTau pathology in human AD[14].

While mechanisms of seizure generation in AD are not fully elucidated, research points to evidence of a synaptic excitatory:inhibitory (E:I) imbalance in chronic epilepsy, including altered expression and function of certain neurotransmitter receptors and ion cotransporters[15-17]. Chronic epilepsy human and animal model studies show evidence of reduced inhibitory neurotransmission, with decreases in the ratio of α1/α3 inhibitory gamma-aminobutyric acid type A receptor (GABA$_A$R) subunits that diminish inhibitory postsynaptic currents (IPSCs)[18-21]. Inhibitory GABAergic transmission is further attenuated by paradoxical depolarizing GABA$_A$R currents due to reversal of the chloride ($Cl^-$) gradient caused by increases in the ratio of the sodium ($Na^+$) potassium ($K^+$) $Cl^-$ co-transporter 1 (NKCC1), which imports $Cl^-$, to that of the potassium $Cl^-$ cotransporter 2 (KCC2), which exports $Cl^-$[22-25]. Commensurate with decreased inhibitory drive in epilepsy, there is an increase in glutamate-receptor mediated excitability, in part due to alterations in α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor (AMPAR) and N-methyl-D-aspartate receptor (NMDAR) subunits expression, with increases in the ratios of GluA1/GluA2 AMPAR subunits and GluN2B/GluN2A NMDAR subunits resulting in GluA2-lacking, $Ca^{2+}$-permeable AMPAR[26-28] and extended decay times and elevated $Ca^{2+}$ influx, respectively[29-31].

Pathologic E:I alterations are not unique to epilepsy, as evidence for neuronal hyperexcitability has been seen in other neurodevelopmental and neurodegenerative disorders[32-35]. In AD patients, altered GABA$_A$R subunit regulation has been observed[36], including decreased α1/α3 expression[37]. Further, evidence for paradoxically depolarizing GABA currents have been

found, with reduced KCC2 in the AD11 mouse model[38] and elevations of NKCC1 after $A\beta_{1-42}$ hippocampal injection[39]. $A\beta_{1-42}$ has also been shown to increase $Ca^{2+}$-permeable AMPARs which can result in memory deficits via reduced long-term potentiation and synapse loss[40-43]. However, given the prevalence of network excitability and seizures in AD, and the evidence that seizures can alter E:I balance in epilepsy, understanding the impact of seizures and the evolution of neuronal hyperexcitability in AD progression may lead to new mitigatable therapeutic E:I targets. To understand the progression of E:I balance, we examined the 5XFAD mouse model at two stages. We also examined E:I balance in human AD with known seizure history and after seizure induction in 5XFAD mice to determine the impact of seizures on E:I balance in AD.

Preclinical and clinical epilepsy research has revealed several candidate signaling pathways that might link E:I imbalance and network hyperexcitability in epilepsy to neurodegeneration. One example is the substantial evidence for the activation of the mammalian target of rapamycin complex 1 (mTORC1) as a key common element between epilepsy and neurodegeneration. mTORC1 mediates pathological tau and APP processing[44-48] and can regulate excitatory and inhibitory receptor and network function[49,50]. Studies of models of Tuberous Sclerosis Complex, characterized by constitutive mTORC1 overactivation commonly resulting in epilepsy[51,52], have demonstrated that blockade of mTORC1 with rapamycin can rescue GABAergic neurotransmission and epilepsy[53-56]. Rapamycin treatment has also rescued neuronal hyperexcitability in models of autism spectrum disorder[57] and neuropathic pain[58]. Together, these results indicate that targeting mTORC1 activity might restore E:I balance across multiple disease models, including AD.

Considering these data and that rapamycin and other rapalogs are approved by the United States Food and Drug Administration for other indications, mTORC1 is a promising target for therapeutic intervention for patients with AD and co-morbid epilepsy. Indeed, we recently found that AD patients with a history of seizure display enhanced tau and $A\beta$ pathology, associated with increased mTORC1 activity[14]. In addition, we observed that the induction of seizures in the 5XFAD mice exacerbates $A\beta$ pathology and cognitive deficits, and that these changes are reversible with chronic low-dose rapamycin treatment.

Given that E:I imbalance may contribute to the cognitive impairments observed in AD, we hypothesized that key components of E:I balance may be dysregulated in AD and that these alterations emerge early in the disease course and may be exacerbated by seizures. First, we

measured established markers of E:I imbalance in largely the same human control and AD temporal neocortex from patients with and without known seizure history as our previous study[14] and indeed found greater alterations in several components of the GABAergic system, along with worsened cognitive and functional performance in AD patients with a seizure history. Next, we examined the 5XFAD mouse model for protein and electrophysiological indications of hyperexcitability at the prodromal stage (~4 months) and directly assessed the effects of seizures induced at this stage on E:I markers at a later, chronic timepoint (7 months of age). Finally, given that we found protective effects on pathology and cognition with rapamycin in this model[14], we utilized the tissue from our prior study to examine whether these therapeutic benefits extend to markers of E:I balance. Overall, we found that E:I imbalance began at a prodromal stage in 5XFAD mice and that seizures significantly exacerbated this imbalance, while chronic rapamycin treatment significantly attenuated these effects.

# Materials and Methods

## Human subjects

All protocols and procedures were approved under the ethical standards of the Institutional Review Board of the University of Pennsylvania (Philadelphia, PA). The AD cognitive study population consisted of 105 patients, selected by querying the integrated neurodegenerative disease database[59] of the Center for Neurodegenerative Disease and Research (CNDR) at the University of Pennsylvania (Philadelphia, PA) for available global clinical dementia rating (CDR) score and CDR Sum of Boxes (CDR-SOB) closest to death. All patients were enrolled in observational research at Penn, with standardized assessments and medical history that includes assessment of seizure history [60]. Of the AD cases, 18 (6 males and 12 females; mean age at death= 77.38 years) had a reported clinical seizure history (AD+Sz). The remaining 87 AD patients (43 males and 44 females; mean age at death= 79.7 years) had no known seizure history (AD-Sz). AD clinical diagnosis was established during life based on the clinical history, neurological and neuropsychological assessment, and confirmed by post-mortem histopathological staging of AD neuropathological markers (A$\beta_{42}$ and tau pathology), according to the National Institute on Aging-Alzheimer's Association (NIA-AA) guidelines[60]. Detailed methods for neurocognitive assessments can be found in Supplementary Materials.

For a subset of the population detailed above, post-mortem brain specimens were acquired from CNDR and consisted of superior/mid-temporal cortex frozen samples (n= 34) and formalin-fixed paraffin-embedded sections (n=15). Detailed clinical characteristics of this AD cohort, obtained from the CNDR INDD database, were previously published[14] and included age at onset of cognitive decline (disease duration), cause of death, brain weight and ordinal rating of gross ventricular enlargement at the time of death, Braak and Thal stages, seizure history and medications. Of the AD cases, 14 (6 males and 8 females; mean age at death= 78.4 years; mean PMI= 11h) had a reported clinical seizure history (AD+Sz). The other 20 AD patients (10 males and 10 females; mean age at death= 73.9 years; mean PMI= 12h) had no known seizure history (AD-Sz). Region-matched control tissue (n=15; 11 males and 4 females; mean age at death= 62.4 years; mean PMI=15.8 hours), received from the CNDR and NIH NeurobioBank, had no known neurologic or psychiatric history. AD patients were significantly older than controls at time of death (p<0.001). PMI did not differ between control and AD groups. Clinical characteristics for all patients included in the cognitive and biochemical studies can be found in Table S1.

## Mice

All animal procedures were approved and performed in accordance with the guidelines of the Institutional Animal Care and Use Committee (IACUC) Office of Animal Welfare of the University of Pennsylvania (Philadelphia, PA). Male and female 5XFAD and WT mice on a mixed B6SJL background second generation (F2) generated from mice originally purchased from Jackson Laboratory (MMRC stock #3840, Bar Harbor ME) were used in these studies. 5XFAD mice harbor the human APP and PSEN1 genes under the Thy1 promoter containing five mutations for aggressive amyloid development (APP: Swedish K670N/M671L, Florida I7516V, and London V717I; PSEN1: M146L and L286V). Genotyping was conducted as specified by Jackson Laboratory. Mice were randomly allocated to treatment groups, which were counterbalanced for sex and litter. No mice were excluded from analysis.

## PTZ kindling

PTZ kindling was performed as previously described [14]. Briefly, mice were administered eight sub-convulsive doses of the $GABA_AR$ antagonist PTZ (i.p., 35 mg/kg of PTZ, Sigma-Aldrich, St Louis, MO) or vehicle (NaCl 0.9%, Sigma-Aldrich) every 48 h for 15 days. Mice were

monitored and video-recorded for one hour after PTZ administration and scored for seizure severity to confirm kindling[14]. One month post-kindling, mice were administered the same dose of PTZ (35 mg/kg) or vehicle to confirm the development of kindling. In our therapeutic cohort, we used a slightly modified kindling procedure to minimize animal loss. Mice were removed from kindling when they reached tonic-clonic seizures on three consecutive days of treatment and were not administered PTZ one month post kindling.

## Rapamycin treatment

Mice were administered encapsulated rapamycin feed dosed at 14mg/kg (5LG6 w/151ppm Encapsulated rapamycin Irr, Emtora Holdings, San Antonio, TX) or control chow (5LG6 w/151ppm Eudragit Irr, Emtora Holdings)[14,61-63] beginning at completion of PTZ kindling (3.5 months of age) until euthanasia at seven months of age. Mice were treated with rapamycin 2.24mg/kg/day by ad libitum consumption of 167 mg/one gram body weight/day.

## Electrophysiology

Mice, 4-5 months of age, were anesthetized with sodium pentobarbital (90 mg/kg, i.p.) and perfused with ice-cold, oxygenated, artificial cerebrospinal fluid (aCSF, in mM, 124 NaCl, 2.5 KCl, 1.2 $NaH_2PO_4$, 24 $NaHCO_3$, 5 HEPES, 2 CaCl2. 2 $MgSO_4$, and 12.5 glucose, pH 7.3-7.4). The brains were removed and cut horizontally (300 μm) in ice-cold, oxygenated, NMDG-aCSF solution (in mM, 93 N-methyl-D-glucamine (NMDG), 2.5 KCl. 1.2 $NaH_2PO_4$, 30 $NaHCO_3$, 20 HEPES, 0.5 $CaCl_2$, 10 $MgSO_4$, 25 glucose, 2 thiourea, 5 sodium ascorbate, and 3 sodium pyruvate, pH 7.3-7.4) using a Leica VT1000s vibratome. Slices were kept in an NMDG-aCSF at 34 °C for 30 min. During this time, incremental amounts of 2M NaCl were added to gradually increase the $Na^+$ concentration in the NMDG-aCSF [64].Then, slices were transferred to aCSF at room temperature and kept until used for recording.

Whole-cell patch clamp recording was performed from hippocampal CA1 pyramidal neurons, identified by a Zeiss Axioscope 2 equipped with IR-DIC. Slices were superfused with room temperature aCSF at a rate of ~ 2ml/min. Signals were amplified and digitized with an Axopatch 200B and a Digidata 1440A. Data were collected and analyzed using pCLAMP 10, Clampfit 10.2 (Molecular Devices) and Mini Analysis 6.0 (Bluecell). For testing action potential (AP) firing properties in current clamp mode, patch pipette electrodes (2-3 MΩ) were filled with

internal solution composed of, in mM: 126 K gluconate, 4 KCl, 10 HEPES, 0.3 EGTA, 0.3 Na-GTP, 3 Mg-ATP, 10 phosphocreatine, pH 7.3, Osm 290. For recording miniature inhibitory synaptic currents (mIPSCs), cells were voltage clamped at -60 mV using electrodes filled with internal solution composed of, in mM; 130 CsCl, 4 NaCl, 10 HEPES, 0.3 EGTA, 1 MgCl2, 0.3 Na-GTP, 3 Mg-ATP, 10 phosphocreatine, pH 7.3, Osm 290. To isolate mIPSCs, TTX (1 μM), NBQX (20 μM), and AP-5 (50 μM) were added to superfusing aCSF. Cells with membrane resistance above 100 MΩ and series resistance less than 10 MΩ were accepted for analysis.

## Western Blot

Mice were anesthetized with 50 mg/kg pentobarbital (Sagent Pharmaceuticals, Schaumburg, IL) and perfused intracardially with cold PBS. Hippocampi and cortices were dissected and immediately frozen. Both human and mouse brain samples were homogenized in a lysis buffer containing 1.1% sucrose, 50 mM Tris HCl pH 7.5, 500 μM $CaCl_2$, 1 mM $MgCl_2$, 1 mM $NaHCO_3$, 1X protease inhibitor, 1mM phenylmethanesulfonyl fluoride (PMSF), and 1X HALT™ (Thermo Fisher Scientific, Carlsbad, CA) on ice and centrifuged at 4,100 g for 10 min at 4° C. Membrane fractions were generated by further centrifugation at 3,000 g for 10 min at 4° C. The resultant supernatant was centrifuged at 13,000 g for 10 min at 4° C. The pellet was reconstituted in lysis buffer. Protein concentrations were determined using the Bradford Protein Assay kit (Bio-Rad, Hercules, CA). Proteins were separated by electrophoresis on 4-20% SDS-PAGE gels (Bio-Rad) and transferred to polyvinylidene difluoride membranes (EMD Millipore, Burlington, MA) following previously used standard protocols[6,14]. Primary antibodies are listed in Table S2. Detected proteins were imaged with Odyssey Imaging System (LI-COR Biosciences) and quantified with Image Lab (Bio-Rad). Each protein was normalized to β-actin expression level.

## Statistics

For cognitive and functional performances, comparisons between AD+Sz and AD-Sz for ranked ordinal data were made by Mann-Whitney test. For the human tissue studies, all AD to control comparisons were made by unpaired t-tests or Mann-Whitney tests for non-normal data. Control, AD-Sz, and AD+Sz comparisons were made by one-way analysis of variance (ANOVA) or Kruskal-Wallis for non-normal data. For the prodromal mouse study, 5XFAD to WT comparisons were made by unpaired t-test. For the nontherapeutic PTZ study, comparisons were

made by two-way ANOVA with genotype and kindling as independent variables. For the therapeutic PTZ study, 5XFAD and WT mice were analyzed separately by two-way ANOVAs with PTZ and rapamycin treatment as independent variables. All ANOVA analyses were followed by Tukey's post hoc test to examine group differences when main effects or interactions were found. Additional multiple linear regressions were performed for all data to determine potential sex effects. Where sex effects were present, data points are displayed as squares (female) or triangles (male) to visualize these differences. Correlations were calculated by simple linear regression. We considered results to be significant at $p < 0.05$. All statistical analyses were performed in GraphPad Prism 9 software (San Diego, CA).

## Data Availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

# Results

## AD patients with seizure history have worsened cognitive and functional performance

To establish potential associations between seizures and cognitive and functional deficits in AD, we examined scores from clinical (global CDR and CDR-SOB) and caretaker questionnaire (DSRS) for subjects with known seizure history. The AD+Sz group showed elevated final global CDR ($p < 0.001$, n=18-87, Figure 1A) and CDR SOB ($p < 0.0001$, n=18-87, Figure 1B) compared to the AD-Sz group. AD+Sz patients also showed significantly worsened overall functional rating (DSRS) compared to those in the AD-Sz group ($p < 0.01$, n=14-19, Figure 1C), including significantly worse ratings in memory, recognition of family members, mobility, orientation to place, social and community activities, and incontinence ($p < 0.05$, n=14-19, Figure S1). These data indicate seizures are associated with worsened cognitive performance and overall function, corroborating previous studies examining epileptiform activity in AD[2-5] and suggesting heightened neuronal dysfunction.

# AD patients with a seizure history exhibit more severe dysregulation of proteins that mediate excitation: inhibition balance

We hypothesized that the worsened functional and cognitive deficits found in AD+Sz patients would be associated with worsening of markers of E:I imbalance. Thus, we examined ratios of $\alpha1/\alpha3$ subunits of the inhibitory GABA$_A$R[19] and Cl$^-$ cotransporters NKCC1/KCC2[22,23,65]. We found a significant decrease in GABA$_A$R$\alpha1$ expression in AD+Sz compared to controls (F=3.52, Tukey's post hoc: p<0.01, Figure 2B), that was not found in AD-Sz, and which drove the reduction in GABA$_A$R$\alpha1$ found across all AD patients (p<0.05, Figure 1B). This alteration also resulted in significantly decreased GABA$_A$R$\alpha1/\alpha3$ ratio in all AD patients when compared to controls (p<0.01, Figure 1B). The ratio of NKCC1/KCC2 was also significantly increased in the AD+Sz group when compared to the AD-Sz group (p<0.05, Figure 2C). This difference was largely attributable to downregulation of KCC2 in the AD+Sz group, compared to the AD-Sz group (F=3.3, Tukey's post hoc: p<0.05, Figure 2C). In addition, sex was a significant predictor of KCC2 dysregulation, with decreased levels in males (p<0.05, [0.0314, 0.230], Figure 2C). Taken together these results suggest AD+Sz patients had greater GABAergic dysfunction due to reduced hyperpolarization and indications of depolarizing GABA$_A$R-mediated signals.

To further explore inhibitory dysfunction, we compared levels of the predominant PV-expressing interneurons[66] in tissue from AD+Sz and AD-Sz, given that there is a well-established association between PV cell loss, cognition[67] and epilepsy[68,69]. Immunohistochemistry revealed that AD+Sz had significantly decreased PV+ cells compared to controls (p<0.01, Figure S2 A-B) that was not found in AD-Sz which drove the significant decrease in PV+ cells found between all AD and controls (p<0.01, Figure S2 A-B).

To determine if neuronal excitation is heightened in AD and whether seizures cause further alterations, we examined the expression of AMPA and NMDA glutamate receptor subunit proteins, namely ratios of GluA1/GluA2 and GluN2B/GluN2A subunits[26,27,29] (Figure 2A-E). While no differences were found between AD+Sz and AD-Sz in AMPAR subunits or subunit ratios (Figure 2D), we found increased expression of GluA2 subunits in AD temporal cortex overall compared to controls (p<0.05, Figure 2D) and the corresponding GluA1/GluA2 ratio was decreased in the AD cases compared to controls (p<0.05, Figure 2D). In examining NMDAR subunits, there was no difference in GluN2B or GluN2A levels nor in the ratio of GluN2B/GluN2A

between the AD+Sz and AD-Sz groups (Figure 2E). However, GluN2A and GluN2B subunits were both expressed at significantly higher levels in AD temporal cortex when compared to controls ($p<0.05$ and $p<0.001$, respectively, Figure 2E).

Finally, we performed linear regressions between clinical outcomes of AD and markers of E:I balance. We found that CDR SOB was inversely correlated with GluN2A levels across all AD subjects ($r=-0.39$, $p<0.05$, Figure 2F) while gross brain atrophy ratings at autopsy was associated with decreased levels of GABA$_A$R$\alpha$1 in AD+Sz ($r= 0.90$, $p<0.01$, Figure 2G), indicating an interaction between these markers of E:I balance and cognitive decline and AD severity, respectively. No correlations were found between markers of E:I balance and pathology previously measured in these samples[14]. These results provide strong evidence of decreased inhibition and increased excitation in AD patients overall, regardless of seizure history, and an exacerbation of selective regulators to increase GABAergic dysfunction in the AD+Sz group.

## Excitation: inhibition imbalance in prodromal stage 5XFAD mice

Given that our human tissue samples are limited by the fact they are from patients at late-stage AD (Table S1), an animal model is ideal to examine earlier stages. Indeed, there is evidence of E:I imbalance reported in several AD mouse models at early stages[70-72]. We thus used the 5XFAD mouse model to determine whether baseline E:I imbalance is present at the prodromal stage (4 months of age), when mild behavioral impairment[73], AD pathology[74], subclinical epileptiform electroencephalographic activity[75], and decreased thresholds for chemoconvulsant induced seizures[14] are present.

Western blot revealed early alterations in markers for GABA inhibition, with 5XFAD mice exhibiting decreased GABA$_A$R$\alpha$1/GABA$_A$R$\alpha$2 ratio in the hippocampus ($p<0.05$, n=14-15), and a trend towards decreased KCC2 in the cortex ($p=0.09$, n=14-15). Notably, no changes in excitatory glutamate AMPAR and NMDAR subunits in hippocampus and cortex remained unchanged at this prodromal stage (Figure 3).

Given that decreases in GABA$_A$R$\alpha$1/$\alpha$2 ratio has been reported to decrease inhibitory synaptic activity in epilepsy models[18-21], we next aimed to determine whether these protein changes also resulted in similar deficits in inhibitory synaptic activity the prodromal 5XFAD mice. We thus measured mIPSCs using whole-cell patch clamp in CA1 pyramidal neurons in hippocampal slices from 4 month old prodromal 5XFAD mice. Indeed, mIPSC amplitudes were

significantly decreased in 5XFAD compared to WT mice (p<0.0001, number of mIPSC events=8316 and 8086, n=16 cells/groups, 2 mice/group, Figure 4A, B, C), consistent with reduced $GABA_AR\alpha1/GABA_AR\alpha2$ ratio[21]. Importantly, there was no change in mIPSC frequency, confirming postsynaptic mechanisms for inhibitory dysfunction. These changes in GABAR function were also associated with increased network excitability in 5XFAD mice as measured by recording evoked action potentials (APs) in current clamp mode using depolarizing step currents of 800 ms duration in the aforementioned CA1 pyramidal neurons (Figure S3). Overall, these data confirmed that the altered GABA expression in hippocampi of prodromal 5XFAD mice was associated with hyperexcitability and may underlie increased seizure susceptibility that we have previously found at this stage in 5XFAD mice[14].

## Kindling during the prodromal stage alters the progressive age-related dysregulation of proteins involved in excitation: inhibition balance in 5XFAD mice

Given the findings of enhanced E:I imbalance in human tissue from AD patients with seizures, we sought to determine how E:I balance changes with disease progression in 5XFAD mice, and whether chronic seizures are associated with further dysregulation of E:I imbalance. We performed PTZ seizure kindling or control protocols in prodromal (3.5 month old) 5XFAD and littermate WT mice and analyzed expression levels of markers of E:I imbalance in the 7 month old (3.5 months post kindling) hippocampus (Figure 5) and cortex (Figure S4). 5XFAD mice had decreased hippocampal $GABA_AR\alpha1/\alpha3$ ratio compared to WT mice (genotype effect: $F_{1,61}$ = 22.76, p<0.0001, Figure 5A), significant decreases in both non-kindled (p<0.05) and PTZ-kindled 5XFAD mice (p<0.01). No differences in $GABA_AR\alpha1/\alpha3$ were found due to kindling. The changes in hippocampal $GABA_AR$ ratio seem to be largely driven by decreases in the $GABA_AR\alpha1$ subunit in 5XFAD mice (genotype effect: $F_{1,61}$=40.09, p<0.0001) (Figure 5A). Decreased $GABA_AR\alpha1/\alpha3$ ratio were also found in the cortex (interaction: $F_{1,43}$ = 4.25, p<0.05), in non-kindled 5XFAD mice compared to WT (p<0.05) (Figure S4A). These changes were similar to patterns seen in human AD brain (Figure 2) and prodromal (4 month) 5XFAD tissue (Figure 4). In addition, sex was a significant predictor of hippocampal $GABA_AR\alpha1/\alpha3$ with greater decreases found in females (p<0.05, [0.0156, -0.1591], Figure 5A). Taken together, these data demonstrate

persistent GABA$_A$R alterations into later disease stages in 5XFAD mice and may contribute to continued heightened seizure susceptibility previously demonstrated in these 5XFAD mice[14].

We next tested whether there were age-related and/or kindling induced alterations to Cl$^-$ cotransporters in 5XFAD mice. As opposed to the prodromal stage, 7-month-old 5XFAD mice showed a decrease in hippocampal expression of the Cl$^-$ exporter, KCC2, regardless of PTZ treatment (genotype effect, $F_{1,61} = 12.35$, $p<0.001$), suggesting worsened inhibitory deficits commensurate with pathological progression. In addition, kindling resulted in an increased NKCC1/KCC2 ratio in the hippocampus of 5XFAD mice (interaction: $F_{1,61}=5.78$, $p<0.05$) compared to WT and non-kindled 5XFAD mice ($p<0.05$. Figure 5B), due to a significant increase in expression of the Cl$^-$ importer, NKCC1 (interaction: $F_{1,61}=7.93$, $p<0.01$; Tukey's post hoc: $p<0.01$) (Figure 5C). In the cortex (Figure S4), we also found a significant increase in NKCC1/KCC2 ratios due to PTZ kindling (kindling effect: $F_{1,43}=4.49$, $p<0.05$), which was largely driven by a reduction of KCC2 in kindled WT mice (interaction: $F_{1,43}= 6.75$, $p<0.05$). Interestingly, there was a decrease in NKCC1 in the non-kindled 5XFAD cortex (genotype effect: $F_{1,43}= 5.2$, $p<0.05$) compared to non-kindled WT mice (Tukey's post hoc: $p<0.05$), which may indicate a compensatory mechanism, although NKCC1/KCC2 ratios were not significantly different (Figure S3B).These data suggest that depolarizing GABA[76] may contribute to the exacerbation of GABAergic dysfunction in kindled 5XFAD mice.

We next sought to determine whether glutamate receptors may be dysregulated as a function of disease progression and also by kindled seizures in 5XFAD mice. In the hippocampus, we found a significant effect to increase GluA1/GluA2 ratios in PTZ kindled hippocampus (genotype x kindling, $F_{1,61}=4.622$, $p<0.05$) (Figure 5C, $p<0.05$). Increased GluA1/GluA2 ratio is a source of E:I imbalance that has been described in both human cases and mouse models of status epilepticus[29]. In addition, both subunits were decreased in the hippocampus of 5XFAD mice regardless of kindling status (GluA1 genotype effect, $F_{1,61} = 7.722$, $p<0.01$; GluA2 genotype effect, $F_{1,61} = 30.28$, $p<0.0001$). Similar alterations in AMPAR subunit expression and ratios were found in the cortex (Figure S4C). The NMDAR GluN2A subunit was also decreased in 5XFAD hippocampus (genotype effect, $F_{1,61}=6.31$, $p<0.05$) (Figure 5D) and cortex ($p<0.05$) (Figure S4D) without change GluN2B and subunit ratios. These data demonstrate dysregulation to glutamate receptor expression beginning by 7 months of age in 5XFAD mice and an exacerbation of this effect due to PTZ kindling.

In addition, we used our retrospective quantitative measures of disease severity in this same cohort of mice[14] and found that NKCC1/KCC2 was positively correlated with A$\beta_{42}$ (r=0.57, p<0.05) and pTau AT100 (Thr212, Ser214)/Tau (r=0.64, p<0.05) in the hippocampus of PTZ kindled 5XFAD mice and with performance on the Y-maze (r=-0.61, p<0.01) (Figure 5F, G, H) which we previously demonstrated was worsened by PTZ kindling. No correlations were found in non-kindled mice nor between GABA$_A$R, AMPAR, or NMDAR in kindled mice. These findings suggest a relationship between inhibitory dysregulation, AD pathology, and cognitive decline.

When taken with the data from prodromal mice (Figure 3), 5XFAD mice show a worsening of markers of inhibitory imbalance, which was associated with worsened pathology and cognitive decline, and the emergence of dysregulated glutamate receptors as pathology progresses from prodrome to later stages. Further, PTZ kindling exacerbated the imbalance in both excitatory and inhibitory markers.

# mTORC1 inhibition modifies select seizure-induced excitation:inhibition markers in the hippocampus of 5XFAD mice

Rapamycin treatment has been shown to rescue neuronal hyperexcitability in various disease models [53-58]. Thus, we sought to determine whether chronic low-dose rapamycin treatment beginning at the cessation of PTZ kindling (prodromal stage ~4 months, 2.24 mg/kg daily), previously used to rescue seizure-induced neuropathology and cognitive deficits in 5XFAD mice[14], could ameliorate PTZ-exacerbated E:I imbalance. The 5XFAD (Figure 6) and WT (Figure S5) mice used in our previous report were analyzed separately by two-way ANOVA with PTZ kindling and rapamycin treatment as independent variables.

Rapamycin showed a strong trend (p=0.058) to increase GABA$_A$R$\alpha$1/$\alpha$3 in non-kindled 5XFAD mice (kindling effect: $F_{1,45}$=6.1, p<0.05; rapamycin effect: $F_{1,45}$=4.1, p<0.05). These shifts corresponded with GABAR$\alpha$1 subunit expression (kindling effect: $F_{1,45}$= 4.2, p<0.05; rapamycin effect: $F_{1,45}$=6.1, p<0.05) (Figure 6A). Rapamycin also reversed the elevation of NKCC1/KCC2 ratio found in PTZ kindled 5XFAD mice (interaction: $F_{1,45}$=9.8, p<0.01) compared to control 5XFAD mice (p<0.05) (Figure 6B). The changes in Cl$^-$ cotransporter ratio was largely driven by changes in the Cl$^-$ importer, NKCC1 (Figure 6B). These results likely reflect a rescue of GABA$_A$R function in kindled 5XFAD mice by rapamycin.

The analysis of AMPAR subunits showed that PTZ increased GluA1/GluA2 ratios in 5XFAD mice (Figure 6C), consistent with the nontherapeutic cohort, and this was reversed by rapamycin treatment (Figure 6C). Rapamycin treatment increased GluA1/GluA2 ratio in non-kindled 5XFAD mice, largely driven by increased GluA1 expression, which reflected similar alterations (interaction: $F_{1,45}$: F=4.3, p<0.05). Additionally, rapamycin treatment reduced GluN2A and GluN2B levels regardless of PTZ treatment (GluN2A rapamycin effect: $F_{1,45}$=5.2, p<0.05; GluN2B rapamycin effect: $F_{1,45}$=4.1, p<0.05) while PTZ kindled, rapamycin treated 5XFAD mice had elevated GluN2B/GluN2A ratio compared to control 5XFAD mice (Tukey's post hoc: p<0.05) (Figure 6D). Overall, these data suggest that rapamycin can ameliorate PTZ-exacerbated inhibitory dysregulation in 5XFAD mice.

In WT mice, no significant differences were found due to PTZ kindling or rapamycin treatment for GABA$_A$Rs, AMPARs, or NMDARs (Figure S5.), although kindling did increase NKCC1 expression in WT mice across rapamycin and vehicle treatments (Kindling effect: $F_{1,47}$=5.2, p<0.05) (Figure S5B).

# Discussion

Here we identified novel mechanisms of neuronal dysfunction that may underlie seizure-exacerbated cognitive and functional deficits in AD. AD temporal cortex showed dysregulation of E:I balance proteins which was worsened in those with seizure history. Similarly, physiologic and protein indications of neuronal E:I imbalance were found in prodromal 5XFAD mice and further perturbations to synaptic protein regulators were seen at later stages and worsened in PTZ-kindled mice (Tables 1 and 2). Together with our recent study demonstrating enhanced AD pathology in AD+Sz patients and PTZ kindled mice[14], our data suggest that E:I imbalance begins at prodromal stages and may play a role in seizure-induced exacerbation of neuropathology and cognitive decline in AD. Specific E:I alterations include decreased GABA$_A$R subunit expression and shifts in the balance between the Cl$^-$ cotransporters NKCC1/KCC2 and the AMPAR GluA1/GluA2 subunit ratios as factors in this interaction between AD and seizures (Tables 1 and 2). Importantly, we found that markers of E:I balance correlated with clinical measures of decline and disease severity in rare autopsy-confirmed tissue samples with prospective collection of seizure history data, justifying follow up studies to further examine clinical outcomes associated with seizure history in AD with more comprehensive statistical modeling that was not possible in the dataset

here. Furthermore, our data indicate that using mTORC1 inhibitors such as rapamycin to therapeutically target the E:I imbalance may prove to be a promising therapy to alleviate AD progression due to neuronal hyperexcitability.

Previous studies have found profound alterations in synaptic GABAergic signaling in AD including downregulation of $GABA_A R$ subunits[36,37,77,78], decreased expression of inhibitory synaptic and perisomatic terminals[79][80], indicative of reduced $GABA_A R$ synapses overall. Here we found decreased $GABA_A R\alpha 1/\alpha 3$ ratios in AD patients as compared to controls, supporting previous reports of reduced GABA sensitivity in membranes isolated from temporal cortex of AD brains[37]. In addition, AD patients with seizure history expressed significantly lower $GABA_A R\alpha 1$ and $GABA_A R\alpha 3$ subunits with $GABA_A R\alpha 1$ level correlating with brain atrophy in AD patients with seizures. NKCC1/KCC2 ratios were increased in AD with a history of seizures as compared to those without, consistent with collapse of neuronal Cl⁻ gradient. In addition, we found a loss of PV containing interneurons in the temporal cortex of AD patients compared to controls, consistent with the previously reported reduction of PV cells within the AD parahippocampal gyrus[81], which was driven by the AD+Sz group. Fast-spiking PV interneurons are major contributors to generation of gamma oscillations, which are disrupted in AD patients[82,83]. Greater loss of this GABAergic population in AD patients with epilepsy may promote network hyperexcitability and contribute to worsened cognitive and functional performance found in these subjects (Figure 1). Taken in the context of prior studies, these data suggest that a history of seizures in AD is associated with diminished GABAergic transmission via reduced $GABA_A R$ synapses, loss of interneurons, and possibly, depolarizing GABA response at those synapses that remain due to elevated intracellular Cl⁻.

Electrophysiological recordings in AD models, and 5XFAD mice in particular, are sparse, especially at stages in which plaque pathology has developed. Our electrophysiological data from the CA1 of prodromal 5XFAD mice demonstrate profound alterations in inhibitory synaptic transmission seen by reductions in mIPSC amplitude associated with increased neuronal excitability. Consistent with alterations in mIPSCs, we found decreased $GABA_A R\alpha 1/ GABA_A R\alpha 2$ ratios in the hippocampus of prodromal 5XFAD mice. Indeed reductions in GABARα1 and α5 transcript were found in patients with mild cognitive impairment suggesting $GABA_A R$ vulnerability at initial disease stages[84].

Non-kindled 7 month old 5XFAD mice also showed decreased expression of $GABA_AR\alpha1$, which we and others previously showed to be associated with decreased PV immunoreactivity[14,85-88], suggesting a broader deficit of inhibition and reflecting our results in AD temporal cortex. Indeed, $GABA_AR$ loss has been linked to seizures and cognitive impairment in mouse models of AD[89-91]. This mechanism may be responsible for such changes in the 5XFAD model and underlie the correlation we found between the loss of $GABA_AR\alpha1$ and increased atrophy in AD patients with seizures. Notably, we found that female sex was a predictor of lower $GABA_AR\alpha1/\alpha3$ ratios, which may be a co-contributor to worsened cognitive outcomes in females[92]. Our data also demonstrate that in 5XFAD mice, induced seizures can exaggerate GABAergic dysfunction increased NKCC1/KCC2 ratios, recapitulating our human data, which were reversed in rapamycin treated mice. Furthermore, our correlations establish a direct link between E:I balance and levels of AD pathology, supporting previous literature positing links between neuronal excitability and AD progression[93].

With respect to glutamatergic neurotransmission, studies of AD hippocampus have highlighted decreased expression of GluA2 compared to controls[94-96], a configuration known to increase AMPAR $Ca^{2+}$ permeability[26,27,29,97,98]. Notably, the human AD temporal cortex we examined was late stage (Table S1) and we instead found an increase in GluA2 with a decrease in corresponding GluA1/GluA2 ratio in AD compared to controls (Figure 1E). These changes may represent a compensatory mechanism to neuronal hyperexcitability, but the significant GluA2 elevation may actually result in further impairments, given that $Ca^{2+}$- permeable AMPAR are critical for the induction of late-phase LTP, essential for the formation of long-term memory[99,100], as seen in neurodevelopmental disorder models[101]. We also found an increase in NMDAR subunits in AD patients, which is consistent with studies of the AD parietal cortex demonstrating increased excitatory to inhibitory synaptic ratio[102]. Elevated GluN2A in AD patients may again represent a compensatory mechanism, as lower GluN2A was associated with worsened CDR-SOB scores across all AD patients in our dataset, consistent with correlations in working memory performance and GluN2A in aged rats[103]. Elevated GluN2B is associated with epileptogenesis[15,29] and extrasynaptic excitotoxic signaling in AD[104]. Further, elevations in AMPAR and NMDAR subunits are suggestive of elevated glutamatergic synapses overall, which may further tip E:I balance towards excitation.

While changes in AMPAR and NMDAR were not found in prodromal 5XFAD mice, we found elevated intrinsic excitability of CA1 neurons, which are the primary output area of hippocampus, together with decreased GABAergic signals suggest increased excitability of the hippocampal network. At later stages, we found increased expression of GluA1/GluA2 AMPAR subunit which was exacerbated by PTZ kindling and ameliorated with rapamycin treatment. Overall, these results suggest that AMPAR and NMDAR subunit composition is dynamic through AD progression and certain changes could be compensatory or neuroprotective from excitotoxicity, as often seen in acute epilepsy models[105,106].

Recent epidemiologic research has underscored strong associations between epilepsy, late onset seizures, and AD[1,107-111]. In the past, seizures were assumed to be an unfortunate biproduct of AD. We and others have provided increasing preclinical and clinical evidence that seizures significantly contribute to neuropathology and cognitive decline and may also be a treatable component of this complex disease. Indeed, in a recent phase 2a clinical trial, levetiracetam, an antiepileptic drug, was shown to improve memory and executive function in AD patients with epileptiform activity[112]. Beyond seizure suppression, our data suggests that targeting specific E:I imbalances also holds therapeutic potential. Notably, a recent study has identified bumetanide, a NKCC1 antagonist, as a top candidate to treat AD, which when applied to AD model mice ameliorated transcriptomic, electrophysiological, and cognitive deficits[113]. Our data demonstrated significant exacerbation of NKCC1/KCC2 in both AD+Sz and seizure kindled mice. NKCC1/KCC2 ratios were also negatively correlated with performance related to spatial working memory in kindled 5XFAD mice, supporting the notion that restoration of the Cl$^-$ gradient, perhaps with bumetanide, is therapeutically relevant, particularly for AD patients with seizure history.

We recently demonstrated that seizures exacerbate mTORC1 in AD patients and 5XFAD mice, and that chronic low dose rapamycin treatment is sufficient to ameliorate AD pathology and cognitive dysfunction in PTZ kindled 5XFAD mice[14]. Rapamycin is FDA approved and has been proven to benefit both AD and epilepsy models[14,114,115]. While rapalogs may affect numerous signaling pathways downstream of mTORC1 to attenuate disease course, including autophagy and neuroinflammation[116,117], growing evidence demonstrates efficacy in regulating neuronal hyperexcitability[53-58]. In the studies presented here, rapamycin reduced NKCC1/KCC2 and GluA1/GluA2 ratios in PTZ kindled 5XFAD mice and showed a trend to increase GABA$_A$Rα1/GABA$_A$Rα3 in non-kindled 5XFAD mice, suggesting reversal of seizure-

exacerbated E:I imbalance, which may indicate restoration of proper circuitry involved in memory in these mice[14]. However, the chronic rapamycin treatment induced small, but significant elevations in GluN2B/GluN2A in PTZ kindled 5XFAD mice and increased GluA1/GluA2 in non-kindled 5XFAD mice. In contrast, studies that used acute rapamycin treatments demonstrated that mTORC1 blockade decreases GluN2B and GluA1[118-121]. Thus, the relative increases in GluN2B and GluA1 that we found may represent a compensatory mechanism in response to chronic administration in 5XFAD mice. Indeed, one prior study demonstrated that chronic rapamycin induced increases in surface GluN2B, and that these changes were associated with amelioration of age-dependent cognitive decline[122]. These data indicate that targeting E:I balance with rapamycin, should be explored for clinical efficacy in AD patients with seizure history.

In summary, the data presented here identified novel mechanisms of neuronal dysfunction in AD that are amplified by seizures and may play a role in worsened cognitive outcomes associated with seizures in AD. These data suggest that targeting E:I imbalance, perhaps with rapamycin as well as other antiepileptic agents, holds therapeutic promise and is particularly relevant for AD patients with comorbid seizures.

# Acknowledgements

The authors thank the CNDR at the University of Pennsylvania for supplying the human tissue utilized in these studies as well as the University of Maryland Brain and Tissue Bank, part of the NIH Neuro Biobank for additional control brain samples. We would also like to thank Dr. Robert Vassar for supplying some of the 4-month 5XFAD mouse tissue.

# Funding

These studies were supported by the National Institutes of Health (NIH) National Institute of Neurological Disorders and Stroke (NINDS): R21NS105437 (FEJ), R01NS101156 (DMT), National Institute on Aging (NIA): R01AG077692 (FEJ & DMT), and Institutional National Research Service Award T32AG000255 (AJB), Alzheimer's Association AARF-22-972333 (AJB). Clinical and postmortem tissue and data from Alzheimer's disease patients was obtained from NIH studies at the University of Pennsylvania including P30-AG-072979 and R01-AG-054519.

# Competing interests

The authors declare no conflicts of interest.

# Authors' contributions

AJB, SG, DMT, FEJ conceived the studies in this manuscript. AJB, SG, XL, DS acquired and analyzed the mouse and human tissue data. AJB, DJI designed the human clinical analysis. EL performed and analyzed electrophysiological experiments. AJB, DJI, FEJ, DMT interpreted the data from mouse and human studies. AJB, DMT, FEJ wrote the manuscript. All authors read and approved the final manuscript.

# Supplementary material.

Supplementary material can be found in a separate PDF file.

# References

Vossel, K. A., Tartaglia, M. C., Nygaard, H. B., Zeman, A. Z. & Miller, B. L. Epileptic activity in Alzheimer's disease: causes and clinical relevance. *Lancet Neurol* **16**, 311-322, doi:10.1016/S1474-4422(17)30044-3 (2017).
Vossel, K. A., Ranasinghe, K. G., Beagle, A. J., Mizuiri, D., Honma, S. M., Dowling, A. F., Darwish, S. M., Van Berlo, V., Barnes, D. E., Mantle, M., Karydas, A. M., Coppola, G., Roberson, E. D., Miller, B. L., Garcia, P. A., Kirsch, H. E., Mucke, L. & Nagarajan, S. S. Incidence and impact of subclinical epileptiform activity in Alzheimer's disease. *Ann Neurol* **80**, 858-870, doi:10.1002/ana.24794 (2016).
Horvath, A. A., Papp, A., Zsuffa, J., Szucs, A., Luckl, J., Radai, F., Nagy, F., Hidasi, Z., Csukly, G., Barcs, G. & Kamondi, A. Subclinical epileptiform activity accelerates the progression of Alzheimer's disease: A long-term EEG study. *Clin Neurophysiol* **132**, 1982-1989, doi:10.1016/j.clinph.2021.03.050 (2021).
Baker, J., Libretto, T., Henley, W. & Zeman, A. A Longitudinal Study of Epileptic Seizures in Alzheimer's Disease. *Front Neurol* **10**, 1266, doi:10.3389/fneur.2019.01266 (2019).
Voglein, J., Ricard, I., Noachtar, S., Kukull, W. A., Dieterich, M., Levin, J. & Danek, A. Seizures in Alzheimer's disease are highly recurrent and associated with a poor disease course. *J Neurol* **267**, 2941-2948, doi:10.1007/s00415-020-09937-7 (2020).
Gourmaud, S., Shou, H., Irwin, D. J., Sansalone, K., Jacobs, L. M., Lucas, T. H., Marsh, E. D., Davis, K. A., Jensen, F. E. & Talos, D. M. Alzheimer-like amyloid and tau alterations

associated with cognitive deficit in temporal lobe epilepsy. *Brain* **143**, 191-209, doi:10.1093/brain/awz381 (2020).

Thom, M., Liu, J. Y., Thompson, P., Phadke, R., Narkiewicz, M., Martinian, L., Marsdon, D., Koepp, M., Caboclo, L., Catarino, C. B. & Sisodiya, S. M. Neurofibrillary tangle pathology and Braak staging in chronic epilepsy in relation to traumatic brain injury and hippocampal sclerosis: a post-mortem study. *Brain* **134**, 2969-2981, doi:10.1093/brain/awr209 (2011).

Tai, X. Y., Koepp, M., Duncan, J. S., Fox, N., Thompson, P., Baxendale, S., Liu, J. Y., Reeves, C., Michalak, Z. & Thom, M. Hyperphosphorylated tau in patients with refractory epilepsy correlates with cognitive decline: a study of temporal lobe resections. *Brain* **139**, 2441-2455, doi:10.1093/brain/aww187 (2016).

Palop, J. J. & Mucke, L. Epilepsy and cognitive impairments in Alzheimer disease. *Arch Neurol* **66**, 435-440, doi:10.1001/archneurol.2009.15 (2009).

Westmark, C. J., Westmark, P. R., Beard, A. M., Hildebrandt, S. M. & Malter, J. S. Seizure susceptibility and mortality in mice that over-express amyloid precursor protein. *Int J Clin Exp Pathol* **1**, 157-168 (2008).

DeVos, S. L., Goncharoff, D. K., Chen, G., Kebodeaux, C. S., Yamada, K., Stewart, F. R., Schuler, D. R., Maloney, S. E., Wozniak, D. F., Rigo, F., Bennett, C. F., Cirrito, J. R., Holtzman, D. M. & Miller, T. M. Antisense reduction of tau in adult mice protects against seizures. *J Neurosci* **33**, 12887-12897, doi:10.1523/JNEUROSCI.2107-13.2013 (2013).

Johnson, E. C. B., Ho, K., Yu, G. Q., Das, M., Sanchez, P. E., Djukic, B., Lopez, I., Yu, X., Gill, M., Zhang, W., Paz, J. T., Palop, J. J. & Mucke, L. Behavioral and neural network abnormalities in human APP transgenic mice resemble those of App knock-in mice and are modulated by familial Alzheimer's disease mutations but not by inhibition of BACE1. *Mol Neurodegener* **15**, 53, doi:10.1186/s13024-020-00393-5 (2020).

Chang, C. W., Evans, M. D., Yu, X., Yu, G. Q. & Mucke, L. Tau reduction affects excitatory and inhibitory neurons differently, reduces excitation/inhibition ratios, and counteracts network hypersynchrony. *Cell Rep* **37**, 109855, doi:10.1016/j.celrep.2021.109855 (2021).

Gourmaud, S., Stewart, D. A., Irwin, D. J., Roberts, N., Barbour, A. J., Eberwine, G., O'Brien, W. T., Vassar, R., Talos, D. M. & Jensen, F. E. The role of mTORC1 activation in seizure-induced exacerbation of Alzheimer's disease. *Brain* **145**, 324-339, doi:10.1093/brain/awab268 (2022).

Rakhade, S. N. & Jensen, F. E. Epileptogenesis in the immature brain: emerging mechanisms. *Nat Rev Neurol* **5**, 380-391, doi:10.1038/nrneurol.2009.80 (2009).

Bakker, J., Basedow, F. J., Dekker, A. D. & Papantoniou, C. Phosphorylation of AMPA-type glutamate receptors: the trigger of epileptogenesis? *J Neurosci* **33**, 5879-5880, doi:10.1523/JNEUROSCI.0496-13.2013 (2013).

Liu, R., Wang, J., Liang, S., Zhang, G. & Yang, X. Role of NKCC1 and KCC2 in Epilepsy: From Expression to Function. *Front Neurol* **10**, 1407, doi:10.3389/fneur.2019.01407 (2019).

Blair, R. E., Sombati, S., Lawrence, D. C., McCay, B. D. & DeLorenzo, R. J. Epileptogenesis causes acute and chronic increases in GABAA receptor endocytosis that contributes to the induction and maintenance of seizures in the hippocampal culture model of

acquired epilepsy. *J Pharmacol Exp Ther* **310**, 871-880, doi:10.1124/jpet.104.068478 (2004).

Noebels, J. L., Avoli, M., Rogawski, M., Olsen, R. & Delgado-Escueta, A. V. "Jasper's Basic Mechanisms of the Epilepsies" Workshop. *Epilepsia* **51 Suppl 5**, 1-5, doi:10.1111/j.1528-1167.2010.02792.x (2010).

Brooks-Kayal, A. R., Shumate, M. D., Jin, H., Rikhter, T. Y. & Coulter, D. A. Selective changes in single cell GABA(A) receptor subunit expression and function in temporal lobe epilepsy. *Nat Med* **4**, 1166-1172, doi:10.1038/2661 (1998).

Zhou, C., Huang, Z., Ding, L., Deel, M. E., Arain, F. M., Murray, C. R., Patel, R. S., Flanagan, C. D. & Gallagher, M. J. Altered cortical GABAA receptor composition, physiology, and endocytosis in a mouse model of a human genetic absence epilepsy syndrome. *J Biol Chem* **288**, 21458-21472, doi:10.1074/jbc.M112.444372 (2013).

Talos, D. M., Sun, H., Kosaras, B., Joseph, A., Folkerth, R. D., Poduri, A., Madsen, J. R., Black, P. M. & Jensen, F. E. Altered inhibition in tuberous sclerosis and type IIb cortical dysplasia. *Ann Neurol* **71**, 539-551, doi:10.1002/ana.22696 (2012).

Palma, E., Amici, M., Sobrero, F., Spinelli, G., Di Angelantonio, S., Ragozzino, D., Mascia, A., Scoppetta, C., Esposito, V., Miledi, R. & Eusebi, F. Anomalous levels of Cl-transporters in the hippocampal subiculum from temporal lobe epilepsy patients make GABA excitatory. *Proc Natl Acad Sci U S A* **103**, 8465-8468, doi:10.1073/pnas.0602979103 (2006).

Barmashenko, G., Hefft, S., Aertsen, A., Kirschstein, T. & Kohling, R. Positive shifts of the GABAA receptor reversal potential due to altered chloride homeostasis is widespread after status epilepticus. *Epilepsia* **52**, 1570-1578, doi:10.1111/j.1528-1167.2011.03247.x (2011).

Kourdougli, N., Pellegrino, C., Renko, J. M., Khirug, S., Chazal, G., Kukko-Lukjanov, T. K., Lauri, S. E., Gaiarsa, J. L., Zhou, L., Peret, A., Castren, E., Tuominen, R. K., Crepel, V. & Rivera, C. Depolarizing gamma-aminobutyric acid contributes to glutamatergic network rewiring in epilepsy. *Ann Neurol* **81**, 251-265, doi:10.1002/ana.24870 (2017).

Lippman-Bell, J. J., Zhou, C., Sun, H., Feske, J. S. & Jensen, F. E. Early-life seizures alter synaptic calcium-permeable AMPA receptor function and plasticity. *Mol Cell Neurosci* **76**, 11-20, doi:10.1016/j.mcn.2016.08.002 (2016).

Rakhade, S. N., Zhou, C., Aujla, P. K., Fishman, R., Sucher, N. J. & Jensen, F. E. Early alterations of AMPA receptors mediate synaptic potentiation induced by neonatal seizures. *J Neurosci* **28**, 7979-7990, doi:10.1523/JNEUROSCI.1734-08.2008 (2008).

Egbenya, D. L., Hussain, S., Lai, Y. C., Xia, J., Anderson, A. E. & Davanger, S. Changes in synaptic AMPA receptor concentration and composition in chronic temporal lobe epilepsy. *Mol Cell Neurosci* **92**, 93-103, doi:10.1016/j.mcn.2018.07.004 (2018).

Loddenkemper, T., Talos, D. M., Cleary, R. T., Joseph, A., Sanchez Fernandez, I., Alexopoulos, A., Kotagal, P., Najm, I. & Jensen, F. E. Subunit composition of glutamate and gamma-aminobutyric acid receptors in status epilepticus. *Epilepsy Res* **108**, 605-615, doi:10.1016/j.eplepsyres.2014.01.015 (2014).

Di Maio, R., Mastroberardino, P. G., Hu, X., Montero, L. & Greenamyre, J. T. Pilocapine alters NMDA receptor expression and function in hippocampal neurons: NADPH oxidase

and ERK1/2 mechanisms. *Neurobiol Dis* **42**, 482-495, doi:10.1016/j.nbd.2011.02.012 (2011).

Di Maio, R., Mastroberardino, P. G., Hu, X., Montero, L. M. & Greenamyre, J. T. Thiol oxidation and altered NR2B/NMDA receptor functions in in vitro and in vivo pilocarpine models: implications for epileptogenesis. *Neurobiol Dis* **49**, 87-98, doi:10.1016/j.nbd.2012.07.013 (2013).

Canitano, R. & Palumbi, R. Excitation/Inhibition Modulators in Autism Spectrum Disorder: Current Clinical Research. *Front Neurosci* **15**, 753274, doi:10.3389/fnins.2021.753274 (2021).

Grigoras, I. F. & Stagg, C. J. Recent advances in the role of excitation-inhibition balance in motor recovery post-stroke. *Fac Rev* **10**, 58, doi:10.12703/r/10-58 (2021).

Lopatina, O. L., Malinovskaya, N. A., Komleva, Y. K., Gorina, Y. V., Shuvaev, A. N., Olovyannikova, R. Y., Belozor, O. S., Belova, O. A., Higashida, H. & Salmina, A. B. Excitation/inhibition imbalance and impaired neurogenesis in neurodevelopmental and neurodegenerative disorders. *Rev Neurosci* **30**, 807-820, doi:10.1515/revneuro-2019-0014 (2019).

Palop, J. J. & Mucke, L. Network abnormalities and interneuron dysfunction in Alzheimer disease. *Nat Rev Neurosci* **17**, 777-792, doi:10.1038/nrn.2016.141 (2016).

Kwakowsky, A., Calvo-Flores Guzman, B., Pandya, M., Turner, C., Waldvogel, H. J. & Faull, R. L. GABAA receptor subunit expression changes in the human Alzheimer's disease hippocampus, subiculum, entorhinal cortex and superior temporal gyrus. *J Neurochem* **145**, 374-392, doi:10.1111/jnc.14325 (2018).

Limon, A., Reyes-Ruiz, J. M. & Miledi, R. Loss of functional GABA(A) receptors in the Alzheimer diseased brain. *Proc Natl Acad Sci U S A* **109**, 10071-10076, doi:10.1073/pnas.1204606109 (2012).

Lagostena, L., Rosato-Siri, M., D'Onofrio, M., Brandi, R., Arisi, I., Capsoni, S., Franzot, J., Cattaneo, A. & Cherubini, E. In the adult hippocampus, chronic nerve growth factor deprivation shifts GABAergic signaling from the hyperpolarizing to the depolarizing direction. *J Neurosci* **30**, 885-893, doi:10.1523/JNEUROSCI.3326-09.2010 (2010).

Lam, P., Vinnakota, C., Guzman, B. C., Newland, J., Peppercorn, K., Tate, W. P., Waldvogel, H. J., Faull, R. L. M. & Kwakowsky, A. Beta-Amyloid (Abeta1-42) Increases the Expression of NKCC1 in the Mouse Hippocampus. *Molecules* **27**, doi:10.3390/molecules27082440 (2022).

Hara, Y., Punsoni, M., Yuk, F., Park, C. S., Janssen, W. G., Rapp, P. R. & Morrison, J. H. Synaptic distributions of GluA2 and PKMzeta in the monkey dentate gyrus and their relationships with aging and memory. *J Neurosci* **32**, 7336-7344, doi:10.1523/JNEUROSCI.0605-12.2012 (2012).

Guntupalli, S., Widagdo, J. & Anggono, V. Amyloid-beta-Induced Dysregulation of AMPA Receptor Trafficking. *Neural Plast* **2016**, 3204519, doi:10.1155/2016/3204519 (2016).

Sanderson, J. L., Freund, R. K., Gorski, J. A. & Dell'Acqua, M. L. beta-Amyloid disruption of LTP/LTD balance is mediated by AKAP150-anchored PKA and Calcineurin regulation of Ca(2+)-permeable AMPA receptors. *Cell Rep* **37**, 109786, doi:10.1016/j.celrep.2021.109786 (2021).

Sun, H., Takesian, A. E., Wang, T. T., Lippman-Bell, J. J., Hensch, T. K. & Jensen, F. E. Early Seizures Prematurely Unsilence Auditory Synapses to Disrupt Thalamocortical Critical Period Plasticity. *Cell Rep* **23**, 2533-2540, doi:10.1016/j.celrep.2018.04.108 (2018).

Caccamo, A., Magri, A., Medina, D. X., Wisely, E. V., Lopez-Aranda, M. F., Silva, A. J. & Oddo, S. mTOR regulates tau phosphorylation and degradation: implications for Alzheimer's disease and other tauopathies. *Aging Cell* **12**, 370-380, doi:10.1111/acel.12057 (2013).

Pei, J. J., An, W. L., Zhou, X. W., Nishimura, T., Norberg, J., Benedikz, E., Gotz, J. & Winblad, B. P70 S6 kinase mediates tau phosphorylation and synthesis. *FEBS Lett* **580**, 107-114, doi:10.1016/j.febslet.2005.11.059 (2006).

Oddo, S. The role of mTOR signaling in Alzheimer disease. *Front Biosci (Schol Ed)* **4**, 941-952, doi:10.2741/s310 (2012).

Cai, Z., Chen, G., He, W., Xiao, M. & Yan, L. J. Activation of mTOR: a culprit of Alzheimer's disease? *Neuropsychiatr Dis Treat* **11**, 1015-1030, doi:10.2147/NDT.S75717 (2015).

Wang, J. Z., Xia, Y. Y., Grundke-Iqbal, I. & Iqbal, K. Abnormal hyperphosphorylation of tau: sites, regulation, and molecular mechanism of neurofibrillary degeneration. *J Alzheimers Dis* **33 Suppl 1**, S123-139, doi:10.3233/JAD-2012-129031 (2013).

McCabe, M. P., Cullen, E. R., Barrows, C. M., Shore, A. N., Tooke, K. I., Laprade, K. A., Stafford, J. M. & Weston, M. C. Genetic inactivation of mTORC1 or mTORC2 in neurons reveals distinct functions in glutamatergic synaptic transmission. *Elife* **9**, doi:10.7554/eLife.51440 (2020).

Kim, S. G., Lee, S., Kim, Y., Park, J., Woo, D., Kim, D., Li, Y., Shin, W., Kang, H., Yook, C., Lee, M., Kim, K., Roh, J. D., Ryu, J., Jung, H., Um, S. M., Yang, E., Kim, H., Han, J., Heo, W. D. & Kim, E. Tanc2-mediated mTOR inhibition balances mTORC1/2 signaling in the developing mouse brain and human neurons. *Nat Commun* **12**, 2695, doi:10.1038/s41467-021-22908-4 (2021).

Talos, D. M., Kwiatkowski, D. J., Cordero, K., Black, P. M. & Jensen, F. E. Cell-specific alterations of glutamate receptor expression in tuberous sclerosis complex cortical tubers. *Ann Neurol* **63**, 454-465, doi:10.1002/ana.21342 (2008).

Wu, X., Sosunov, A. A., Lado, W., Teoh, J. J., Ham, A., Li, H., Al-Dalahmah, O., Gill, B. J. A., Arancio, O., Schevon, C. A., Frankel, W. N., McKhann, G. M., 2nd, Sulzer, D., Goldman, J. E. & Tang, G. Synaptic hyperexcitability of cytomegalic pyramidal neurons contributes to epileptogenesis in tuberous sclerosis complex. *Cell Rep* **40**, 111085, doi:10.1016/j.celrep.2022.111085 (2022).

Amegandjin, C. A., Choudhury, M., Jadhav, V., Carrico, J. N., Quintal, A., Berryer, M., Snapyan, M., Chattopadhyaya, B., Saghatelyan, A. & Di Cristo, G. Sensitive period for rescuing parvalbumin interneurons connectivity and social behavior deficits caused by TSC1 loss. *Nat Commun* **12**, 3653, doi:10.1038/s41467-021-23939-7 (2021).

Bateup, H. S., Johnson, C. A., Denefrio, C. L., Saulnier, J. L., Kornacker, K. & Sabatini, B. L. Excitatory/inhibitory synaptic imbalance leads to hippocampal hyperexcitability in mouse models of tuberous sclerosis. *Neuron* **78**, 510-522, doi:10.1016/j.neuron.2013.03.017 (2013).

Koene, L. M., Niggl, E., Wallaard, I., Proietti-Onori, M., Rotaru, D. C. & Elgersma, Y. Identifying the temporal electrophysiological and molecular changes that contribute to TSC-associated epileptogenesis. *JCI Insight* **6**, doi:10.1172/jci.insight.150120 (2021).

Goto, J., Talos, D. M., Klein, P., Qin, W., Chekaluk, Y. I., Anderl, S., Malinowska, I. A., Di Nardo, A., Bronson, R. T., Chan, J. A., Vinters, H. V., Kernie, S. G., Jensen, F. E., Sahin, M. & Kwiatkowski, D. J. Regulable neural progenitor-specific Tsc1 loss yields giant cells with organellar dysfunction in a model of tuberous sclerosis complex. *Proc Natl Acad Sci U S A* **108**, E1070-1079, doi:10.1073/pnas.1106454108 (2011).

Iezzi, D., Curti, L., Ranieri, G., Gerace, E., Costa, A., Ilari, A., La Rocca, A., Luceri, C., D'Ambrosio, M., Silvestri, L., Scardigli, M., Mannaioni, G. & Masi, A. Acute rapamycin rescues the hyperexcitable phenotype of accumbal medium spiny neurons in the valproic acid rat model of autism spectrum disorder. *Pharmacol Res* **183**, 106401, doi:10.1016/j.phrs.2022.106401 (2022).

Ma, X., Du, W., Wang, W., Luo, L., Huang, M., Wang, H., Lin, R., Li, Z., Shi, H., Yuan, T., Jiang, W., Worley, P. F. & Xu, T. Persistent Rheb-induced mTORC1 activation in spinal cord neurons induces hypersensitivity in neuropathic pain. *Cell Death Dis* **11**, 747, doi:10.1038/s41419-020-02966-0 (2020).

Xie, S. X., Baek, Y., Grossman, M., Arnold, S. E., Karlawish, J., Siderowf, A., Hurtig, H., Elman, L., McCluskey, L., Van Deerlin, V., Lee, V. M. & Trojanowski, J. Q. Building an integrated neurodegenerative disease database at an academic health center. *Alzheimers Dement* **7**, e84-93, doi:10.1016/j.jalz.2010.08.233 (2011).

Hyman, B. T., Phelps, C. H., Beach, T. G., Bigio, E. H., Cairns, N. J., Carrillo, M. C., Dickson, D. W., Duyckaerts, C., Frosch, M. P., Masliah, E., Mirra, S. S., Nelson, P. T., Schneider, J. A., Thal, D. R., Thies, B., Trojanowski, J. Q., Vinters, H. V. & Montine, T. J. National Institute on Aging-Alzheimer's Association guidelines for the neuropathologic assessment of Alzheimer's disease. *Alzheimers Dement* **8**, 1-13, doi:10.1016/j.jalz.2011.10.007 (2012).

Lin, A. L., Zheng, W., Halloran, J. J., Burbank, R. R., Hussong, S. A., Hart, M. J., Javors, M., Shih, Y. Y., Muir, E., Solano Fonseca, R., Strong, R., Richardson, A. G., Lechleiter, J. D., Fox, P. T. & Galvan, V. Chronic rapamycin restores brain vascular integrity and function through NO synthase activation and improves memory in symptomatic mice modeling Alzheimer's disease. *J Cereb Blood Flow Metab* **33**, 1412-1421, doi:10.1038/jcbfm.2013.82 (2013).

Serrano-Pozo, A., Mielke, M. L., Gómez-Isla, T., Betensky, R. A., Growdon, J. H., Frosch, M. P. & Hyman, B. T. Reactive glia not only associates with plaques but also parallels tangles in Alzheimer's disease. *Am J Pathol* **179**, 1373-1384, doi:10.1016/j.ajpath.2011.05.047 (2011).

Lin, A. L., Jahrling, J. B., Zhang, W., DeRosa, N., Bakshi, V., Romero, P., Galvan, V. & Richardson, A. Rapamycin rescues vascular, metabolic and learning deficits in apolipoprotein E4 transgenic mice with pre-symptomatic Alzheimer's disease. *J Cereb Blood Flow Metab* **37**, 217-226, doi:10.1177/0271678x15621575 (2017).

Ting, J. T., Lee, B. R., Chong, P., Soler-Llavina, G., Cobbs, C., Koch, C., Zeng, H. & Lein, E. Preparation of Acute Brain Slices Using an Optimized N-Methyl-D-glucamine Protective Recovery Method. *J Vis Exp*, doi:10.3791/53825 (2018).

Galanopoulou, A. S. GABA(A) receptors in normal development and seizures: friends or foes? *Curr Neuropharmacol* **6**, 1-20, doi:10.2174/157015908783769653 (2008).

Rudy, B., Fishell, G., Lee, S. & Hjerling-Leffler, J. Three groups of interneurons account for nearly 100% of neocortical GABAergic neurons. *Dev Neurobiol* **71**, 45-61, doi:10.1002/dneu.20853 (2011).

Murray, A. J., Woloszynowska-Fraser, M. U., Ansel-Bollepalli, L., Cole, K. L., Foggetti, A., Crouch, B., Riedel, G. & Wulff, P. Parvalbumin-positive interneurons of the prefrontal cortex support working memory and cognitive flexibility. *Sci Rep* **5**, 16778, doi:10.1038/srep16778 (2015).

Drexel, M., Preidt, A. P., Kirchmair, E. & Sperk, G. Parvalbumin interneurons and calretinin fibers arising from the thalamic nucleus reuniens degenerate in the subiculum after kainic acid-induced seizures. *Neuroscience* **189**, 316-329, doi:10.1016/j.neuroscience.2011.05.021 (2011).

Zamecnik, J., Krsek, P., Druga, R., Marusic, P., Benes, V., Tichy, M. & Komarek, V. Densities of parvalbumin-immunoreactive neurons in non-malformed hippocampal sclerosis-temporal neocortex and in cortical dysplasias. *Brain Res Bull* **68**, 474-481, doi:10.1016/j.brainresbull.2005.10.008 (2006).

Kim, K. R., Kim, Y., Jeong, H. J., Kang, J. S., Lee, S. H., Kim, Y., Lee, S. H. & Ho, W. K. Impaired pattern separation in Tg2576 mice is associated with hyperexcitable dentate gyrus caused by Kv4.1 downregulation. *Mol Brain* **14**, 62, doi:10.1186/s13041-021-00774-x (2021).

Kazim, S. F., Chuang, S. C., Zhao, W., Wong, R. K., Bianchi, R. & Iqbal, K. Early-Onset Network Hyperexcitability in Presymptomatic Alzheimer's Disease Transgenic Mice Is Suppressed by Passive Immunization with Anti-Human APP/Abeta Antibody and by mGluR5 Blockade. *Front Aging Neurosci* **9**, 71, doi:10.3389/fnagi.2017.00071 (2017).

Davis, K. E., Fox, S. & Gigg, J. Increased hippocampal excitability in the 3xTgAD mouse model for Alzheimer's disease in vivo. *PLoS One* **9**, e91203, doi:10.1371/journal.pone.0091203 (2014).

Kimura, R. & Ohno, M. Impairments in remote memory stabilization precede hippocampal synaptic and cognitive failures in 5XFAD Alzheimer mouse model. *Neurobiol Dis* **33**, 229-235, doi:10.1016/j.nbd.2008.10.006 (2009).

Oakley, H., Cole, S. L., Logan, S., Maus, E., Shao, P., Craft, J., Guillozet-Bongaarts, A., Ohno, M., Disterhoft, J., Van Eldik, L., Berry, R. & Vassar, R. Intraneuronal beta-amyloid aggregates, neurodegeneration, and neuron loss in transgenic mice with five familial Alzheimer's disease mutations: potential factors in amyloid plaque formation. *J Neurosci* **26**, 10129-10140, doi:10.1523/JNEUROSCI.1202-06.2006 (2006).

Abe, Y., Ikegawa, N., Yoshida, K., Muramatsu, K., Hattori, S., Kawai, K., Murakami, M., Tanaka, T., Goda, W., Goto, M., Yamamoto, T., Hashimoto, T., Yamada, K., Shibata, T., Misawa, H., Mimura, M., Tanaka, K. F., Miyakawa, T., Iwatsubo, T., Hata, J. I., Niikura, T. & Yasui, M. Behavioral and electrophysiological evidence for a neuroprotective role of aquaporin-4 in the 5xFAD transgenic mice model. *Acta Neuropathol Commun* **8**, 67, doi:10.1186/s40478-020-00936-3 (2020).

Schulte, J. T., Wierenga, C. J. & Bruining, H. Chloride transporters and GABA polarity in developmental, neurological and psychiatric conditions. *Neurosci Biobehav Rev* **90**, 260-271, doi:10.1016/j.neubiorev.2018.05.001 (2018).

Govindpani, K., Turner, C., Waldvogel, H. J., Faull, R. L. M. & Kwakowsky, A. Impaired Expression of GABA Signaling Components in the Alzheimer's Disease Middle Temporal Gyrus. *Int J Mol Sci* **21**, doi:10.3390/ijms21228704 (2020).

Rissman, R. A., Mishizen-Eberz, A. J., Carter, T. L., Wolfe, B. B., De Blas, A. L., Miralles, C. P., Ikonomovic, M. D. & Armstrong, D. M. Biochemical analysis of GABA(A) receptor subunits alpha 1, alpha 5, beta 1, beta 2 in the hippocampus of patients with Alzheimer's disease neuropathology. *Neuroscience* **120**, 695-704, doi:10.1016/s0306-4522(03)00030-7 (2003).

Agarwal, S., Tannenberg, R. K. & Dodd, P. R. Reduced expression of the inhibitory synapse scaffolding protein gephyrin in Alzheimer's disease. *J Alzheimers Dis* **14**, 313-321, doi:10.3233/jad-2008-14305 (2008).

Garcia-Marin, V., Blazquez-Llorca, L., Rodriguez, J. R., Boluda, S., Muntane, G., Ferrer, I. & Defelipe, J. Diminished perisomatic GABAergic terminals on cortical neurons adjacent to amyloid plaques. *Front Neuroanat* **3**, 28, doi:10.3389/neuro.05.028.2009 (2009).

Sanchez-Mejias, E., Nunez-Diaz, C., Sanchez-Varo, R., Gomez-Arboledas, A., Garcia-Leon, J. A., Fernandez-Valenzuela, J. J., Mejias-Ortega, M., Trujillo-Estrada, L., Baglietto-Vargas, D., Moreno-Gonzalez, I., Davila, J. C., Vitorica, J. & Gutierrez, A. Distinct disease-sensitive GABAergic neurons in the perirhinal cortex of Alzheimer's mice and patients. *Brain Pathol* **30**, 345-363, doi:10.1111/bpa.12785 (2020).

Stam, C. J., van Cappellen van Walsum, A. M., Pijnenburg, Y. A., Berendse, H. W., de Munck, J. C., Scheltens, P. & van Dijk, B. W. Generalized synchronization of MEG recordings in Alzheimer's Disease: evidence for involvement of the gamma band. *J Clin Neurophysiol* **19**, 562-574, doi:10.1097/00004691-200212000-00010 (2002).

Klein, A. S., Donoso, J. R., Kempter, R., Schmitz, D. & Beed, P. Early Cortical Changes in Gamma Oscillations in Alzheimer's Disease. *Front Syst Neurosci* **10**, 83, doi:10.3389/fnsys.2016.00083 (2016).

Rissman, R. A., Bennett, D. A. & Armstrong, D. M. Subregional analysis of GABA(A) receptor subunit mRNAs in the hippocampus of older persons with and without cognitive impairment. *J Chem Neuroanat* **28**, 17-25, doi:10.1016/j.jchemneu.2004.05.003 (2004).

Ali, F., Baringer, S. L., Neal, A., Choi, E. Y. & Kwan, A. C. Parvalbumin-Positive Neuron Loss and Amyloid-beta Deposits in the Frontal Cortex of Alzheimer's Disease-Related Mice. *J Alzheimers Dis* **72**, 1323-1339, doi:10.3233/JAD-181190 (2019).

Flanigan, T. J., Xue, Y., Kishan Rao, S., Dhanushkodi, A. & McDonald, M. P. Abnormal vibrissa-related behavior and loss of barrel field inhibitory neurons in 5xFAD transgenics. *Genes Brain Behav* **13**, 488-500, doi:10.1111/gbb.12133 (2014).

Giesers, N. K. & Wirths, O. Loss of Hippocampal Calretinin and Parvalbumin Interneurons in the 5XFAD Mouse Model of Alzheimer's Disease. *ASN Neuro* **12**, 1759091420925356, doi:10.1177/1759091420925356 (2020).

Caccavano, A., Bozzelli, P. L., Forcelli, P. A., Pak, D. T. S., Wu, J. Y., Conant, K. & Vicini, S. Inhibitory Parvalbumin Basket Cell Activity is Selectively Reduced during Hippocampal

Sharp Wave Ripples in a Mouse Model of Familial Alzheimer's Disease. *J Neurosci* **40**, 5116-5136, doi:10.1523/JNEUROSCI.0425-20.2020 (2020).

Verret, L., Mann, E. O., Hang, G. B., Barth, A. M., Cobos, I., Ho, K., Devidze, N., Masliah, E., Kreitzer, A. C., Mody, I., Mucke, L. & Palop, J. J. Inhibitory interneuron deficit links altered network activity and cognitive dysfunction in Alzheimer model. *Cell* **149**, 708-721, doi:10.1016/j.cell.2012.02.046 (2012).

Li, Y., Zhu, K., Li, N., Wang, X., Xiao, X., Li, L., Li, L., He, Y., Zhang, J., Wo, J., Cui, Y., Huang, H., Zhang, J., Wang, W., Wang, X. & Zheng, Y. Reversible GABAergic dysfunction involved in hippocampal hyperactivity predicts early-stage Alzheimer disease in a mouse model. *Alzheimers Res Ther* **13**, 114, doi:10.1186/s13195-021-00859-8 (2021).

Ulrich, D. Amyloid-beta Impairs Synaptic Inhibition via GABA(A) Receptor Endocytosis. *J Neurosci* **35**, 9205-9210, doi:10.1523/JNEUROSCI.0950-15.2015 (2015).

Ferretti, M. T., Iulita, M. F., Cavedo, E., Chiesa, P. A., Schumacher Dimech, A., Santuccione Chadha, A., Baracchi, F., Girouard, H., Misoch, S., Giacobini, E., Depypere, H., Hampel, H., Women's Brain, P. & the Alzheimer Precision Medicine, I. Sex differences in Alzheimer disease - the gateway to precision medicine. *Nat Rev Neurol* **14**, 457-469, doi:10.1038/s41582-018-0032-9 (2018).

Targa Dias Anastacio, H., Matosin, N. & Ooi, L. Neuronal hyperexcitability in Alzheimer's disease: what are the drivers behind this aberrant phenotype? *Transl Psychiatry* **12**, 257, doi:10.1038/s41398-022-02024-7 (2022).

Carter, T. L., Rissman, R. A., Mishizen-Eberz, A. J., Wolfe, B. B., Hamilton, R. L., Gandy, S. & Armstrong, D. M. Differential preservation of AMPA receptor subunits in the hippocampi of Alzheimer's disease patients according to Braak stage. *Exp Neurol* **187**, 299-309, doi:10.1016/j.expneurol.2003.12.010 (2004).

Ikonomovic, M. D., Mizukami, K., Davies, P., Hamilton, R., Sheffield, R. & Armstrong, D. M. The loss of GluR2(3) immunoreactivity precedes neurofibrillary tangle formation in the entorhinal cortex and hippocampus of Alzheimer brains. *J Neuropathol Exp Neurol* **56**, 1018-1027, doi:10.1097/00005072-199709000-00007 (1997).

Mohamed, N. E., Zhao, Y., Lee, J. H., Tan, M. G., Esiri, M. M., Wilcock, G. K., Smith, A. D., Wong, P. T., Chen, C. P. & Lai, M. K. Upregulation of AMPA receptor GluR2 (GluA2) subunits in subcortical ischemic vascular dementia is repressed in the presence of Alzheimer's disease. *Neurochem Int* **58**, 820-825, doi:10.1016/j.neuint.2011.03.010 (2011).

Burnashev, N., Monyer, H., Seeburg, P. H. & Sakmann, B. Divalent ion permeability of AMPA receptor channels is dominated by the edited form of a single subunit. *Neuron* **8**, 189-198, doi:10.1016/0896-6273(92)90120-3 (1992).

Rajasekaran, K., Todorovic, M. & Kapur, J. Calcium-permeable AMPA receptors are expressed in a rodent model of status epilepticus. *Ann Neurol* **72**, 91-102, doi:10.1002/ana.23570 (2012).

Park, P., Kang, H., Sanderson, T. M., Bortolotto, Z. A., Georgiou, J., Zhuo, M., Kaang, B. K. & Collingridge, G. L. The Role of Calcium-Permeable AMPARs in Long-Term Potentiation at Principal Neurons in the Rodent Hippocampus. *Front Synaptic Neurosci* **10**, 42, doi:10.3389/fnsyn.2018.00042 (2018).

Pang, P. T. & Lu, B. Regulation of late-phase LTP and long-term memory in normal and aging hippocampus: role of secreted proteins tPA and BDNF. *Ageing Res Rev* **3**, 407-430, doi:10.1016/j.arr.2004.07.002 (2004).

Banke, T. G. & Barria, A. Transient Enhanced GluA2 Expression in Young Hippocampal Neurons of a Fragile X Mouse Model. *Front Synaptic Neurosci* **12**, 588295, doi:10.3389/fnsyn.2020.588295 (2020).

Lauterborn, J. C., Scaduto, P., Cox, C. D., Schulmann, A., Lynch, G., Gall, C. M., Keene, C. D. & Limon, A. Increased excitatory to inhibitory synaptic ratio in parietal cortex samples from individuals with Alzheimer's disease. *Nat Commun* **12**, 2603, doi:10.1038/s41467-021-22742-8 (2021).

McQuail, J. A., Beas, B. S., Kelly, K. B., Simpson, K. L., Frazier, C. J., Setlow, B. & Bizon, J. L. NR2A-Containing NMDARs in the Prefrontal Cortex Are Required for Working Memory and Associated with Age-Related Cognitive Decline. *J Neurosci* **36**, 12537-12548, doi:10.1523/JNEUROSCI.2332-16.2016 (2016).

Babaei, P. NMDA and AMPA receptors dysregulation in Alzheimer's disease. *Eur J Pharmacol* **908**, 174310, doi:10.1016/j.ejphar.2021.174310 (2021).

Russo, I., Bonini, D., Via, L. L., Barlati, S. & Barbon, A. AMPA receptor properties are modulated in the early stages following pilocarpine-induced status epilepticus. *Neuromolecular Med* **15**, 324-338, doi:10.1007/s12017-013-8221-6 (2013).

Leo, A., Giovannini, G., Russo, E. & Meletti, S. The role of AMPA receptors and their antagonists in status epilepticus. *Epilepsia* **59**, 1098-1108, doi:10.1111/epi.14082 (2018).

Zhang, D., Chen, S., Xu, S., Wu, J., Zhuang, Y., Cao, W., Chen, X. & Li, X. The clinical correlation between Alzheimer's disease and epilepsy. *Front Neurol* **13**, 922535, doi:10.3389/fneur.2022.922535 (2022).

Lam, A. D., Sarkis, R. A., Pellerin, K. R., Jing, J., Dworetzky, B. A., Hoch, D. B., Jacobs, C. S., Lee, J. W., Weisholtz, D. S., Zepeda, R., Westover, M. B., Cole, A. J. & Cash, S. S. Association of epileptiform abnormalities and seizures in Alzheimer disease. *Neurology* **95**, e2259-e2270, doi:10.1212/WNL.0000000000010612 (2020).

Lam, A. D., Deck, G., Goldman, A., Eskandar, E. N., Noebels, J. & Cole, A. J. Silent hippocampal seizures and spikes identified by foramen ovale electrodes in Alzheimer's disease. *Nat Med* **23**, 678-680, doi:10.1038/nm.4330 (2017).

Johnson, E. L., Krauss, G. L., Kucharska-Newton, A., Albert, M. S., Brandt, J., Walker, K. A., Yasar, S., Knopman, D. S., Vossel, K. A. & Gottesman, R. F. Dementia in late-onset epilepsy: The Atherosclerosis Risk in Communities study. *Neurology* **95**, e3248-e3256, doi:10.1212/WNL.0000000000011080 (2020).

Kawakami, O., Koike, Y., Ando, T., Sugiura, M., Kato, H., Hiraga, K., Kito, H. & Kondo, H. Incidence of dementia in patients with adult-onset epilepsy of unknown causes. *J Neurol Sci* **395**, 71-76, doi:10.1016/j.jns.2018.09.010 (2018).

Vossel, K., Ranasinghe, K. G., Beagle, A. J., La, A., Ah Pook, K., Castro, M., Mizuiri, D., Honma, S. M., Venkateswaran, N., Koestler, M., Zhang, W., Mucke, L., Howell, M. J., Possin, K. L., Kramer, J. H., Boxer, A. L., Miller, B. L., Nagarajan, S. S. & Kirsch, H. E. Effect of Levetiracetam on Cognition in Patients With Alzheimer Disease With and Without Epileptiform Activity: A Randomized Clinical Trial. *JAMA Neurol* **78**, 1345-1354, doi:10.1001/jamaneurol.2021.3310 (2021).

Taubes, A., Nova, P., Zalocusky, K. A., Kosti, I., Bicak, M., Zilberter, M. Y., Hao, Y., Yoon, S. Y., Oskotsky, T., Pineda, S., Chen, B., Aery Jones, E. A., Choudhary, K., Grone, B., Balestra, M. E., Chaudhry, F., Paranjpe, I., De Freitas, J., Koutsodendris, N., Chen, N., Wang, C., Chang, W., An, A., Glicksberg, B. S., Sirota, M. & Huang, Y. Experimental and real-world evidence supporting the computational repurposing of bumetanide for APOE4-related Alzheimer's disease. *Nature Aging* **1**, 932-947, doi:10.1038/s43587-021-00122-7 (2021).

Kim, J. K. & Lee, J. H. Mechanistic Target of Rapamycin Pathway in Epileptic Disorders. *J Korean Neurosurg Soc* **62**, 272-287, doi:10.3340/jkns.2019.0027 (2019).

Kaeberlein, M. & Galvan, V. Rapamycin and Alzheimer's disease: Time for a clinical trial? *Sci Transl Med* **11**, doi:10.1126/scitranslmed.aar4289 (2019).

Kodali, M., Attaluri, S., Madhu, L. N., Shuai, B., Upadhya, R., Gonzalez, J. J., Rao, X. & Shetty, A. K. Metformin treatment in late middle age improves cognitive function with alleviation of microglial activation and enhancement of autophagy in the hippocampus. *Aging Cell* **20**, e13277, doi:10.1111/acel.13277 (2021).

Zeng, L. H., Xu, L., Gutmann, D. H. & Wong, M. Rapamycin prevents epilepsy in a mouse model of tuberous sclerosis complex. *Ann Neurol* **63**, 444-453, doi:10.1002/ana.21331 (2008).

James, M. H., Quinn, R. K., Ong, L. K., Levi, E. M., Charnley, J. L., Smith, D. W., Dickson, P. W. & Dayas, C. V. mTORC1 inhibition in the nucleus accumbens 'protects' against the expression of drug seeking and 'relapse' and is associated with reductions in GluA1 AMPAR and CAMKIIalpha levels. *Neuropsychopharmacology* **39**, 1694-1702, doi:10.1038/npp.2014.16 (2014).

James, M. H., Quinn, R. K., Ong, L. K., Levi, E. M., Smith, D. W., Dickson, P. W. & Dayas, C. V. Rapamycin reduces motivated responding for cocaine and alters GluA1 expression in the ventral but not dorsal striatum. *Eur J Pharmacol* **784**, 147-154, doi:10.1016/j.ejphar.2016.05.013 (2016).

Calabrese, V., Di Maio, A., Marino, G., Cardinale, A., Natale, G., De Rosa, A., Campanelli, F., Mancini, M., Napolitano, F., Avallone, L., Calabresi, P., Usiello, A., Ghiglieri, V. & Picconi, B. Rapamycin, by Inhibiting mTORC1 Signaling, Prevents the Loss of Striatal Bidirectional Synaptic Plasticity in a Rat Model of L-DOPA-Induced Dyskinesia. *Front Aging Neurosci* **12**, 230, doi:10.3389/fnagi.2020.00230 (2020).

Zhang, Y., Ji, F., Wang, G., He, D., Yang, L. & Zhang, M. BDNF Activates mTOR to Upregulate NR2B Expression in the Rostral Anterior Cingulate Cortex Required for Inflammatory Pain-Related Aversion in Rats. *Neurochem Res* **43**, 681-691, doi:10.1007/s11064-018-2470-6 (2018).

Majumder, S., Caccamo, A., Medina, D. X., Benavides, A. D., Javors, M. A., Kraig, E., Strong, R., Richardson, A. & Oddo, S. Lifelong rapamycin administration ameliorates age-dependent cognitive deficits by reducing IL-1beta and enhancing NMDA signaling. *Aging Cell* **11**, 326-335, doi:10.1111/j.1474-9726.2011.00791.x (2012).

**Figure 1. Seizure history is associated worse cognitive and functional performance in AD.** Comparisons between AD-Sz and AD+Sz in Final global CDR (**A**), CDR sum of boxes (**B**), and DSRS Total (**C**). CDR: n=18 (AD+Sz) – 87 (AD-Sz); DSRS: n=14 (AD+Sz) – 19 (AD-Sz). *p<0.05, **p<0.01.

**Figure 2. Dysregulation of proteins involved in excitatory: inhibitory balance in AD patients with and without seizure history.** (**A**) Representative western blot images for (B-E) showing non-adjacent bands originating from the same blot of the temporal cortex from control cases (Con; n=13) and Alzheimer's disease patients (AD; n=30) split into subgroups, those without (AD-Sz; n=19) or with (AD+Sz; n=11) known seizure history. Semi-quantitative analysis of (**B**) GABA$_A$R subunits GABA$_A$Rα1 and GABA$_A$Rα3 and corresponding ratio GABA$_A$Rα1/GABA$_A$Rα3; (**C**) Cl$^-$ cotransporters NKCC1 and KCC2 and corresponding ratio NKCC1/KCC2; (**D**) AMPAR subunits GluA1 and GluA2 and corresponding ratio GluA1/GluA2; (**E**) NMDAR subunits GluN2A and GluN2B and corresponding ratio GluN2B/GluN2A. *p<0.05, **p<0.01. (**F**) Pearson correlations in all AD showing relationship between GluN2A and CDR SOB score and (**G**) showing the relationship between GABA$_A$Rα1 expression and brain weight at death in grams in AD+Sz patients. Grey areas indicate 95% confidence interval for the two means.

**Figure 3. Excitatory: inhibitory markers in prodromal 5XFAD mice.** Analysis of bands from western blots for E:I markers from the hippocampus (A, C, E, G) and cortex (B, D, F, H) of 5XFAD and WT mice. (**A, B**) GABA$_A$R subunits GABA$_A$Rα1 and GABA$_A$Rα2 and corresponding ratio GABA$_A$Rα1/GABA$_A$Rα2 (**C, D**) Cl$^-$ cotransporters NKCC1 and KCC2 and corresponding NKCC1/KCC2 ratio; (**E, F**) AMPAR subunits GluA1 and GluA2 and corresponding GluA1/GluA2 ratio; (**G, H**) NMDAR subunits GluN2A, GluN2B, and corresponding GluN2B/GluN2A ratio. (**I**) Representative Western blot images for (**A-H**) showing non-adjacent bands originating from the same blot. Error bars represent standard error of the mean. n=11-15 per group. *p<0.05.

**Figure 4. Inhibitory dysfunction in pyramidal neurons from prodromal 5XFAD CA1. (A)** Representative mIPSC traces from WT and 5XFAD. (**B**, **C**) Quantifications of mIPSC amplitudes. n= 16 cells from 2 WT mice, 16 cells from 2 5XFAD mice. ****$p<0.0001$.

**Figure 5. Excitatory:inhibitory imbalance in the hippocampus of 5XFAD mice following induced seizures. (A-E)** Quantification of (**A**) GABA$_A$R subunits GABA$_A$R$\alpha$1 and GABA$_A$R$\alpha$3 and corresponding ratio GABA$_A$R$\alpha$1/GABA$_A$R$\alpha$3; (**B**) Cl⁻cotransporters NKCC1 and KCC2 and corresponding ratio NKCC1/KCC2; (**C**) AMPAR subunits GluA1 and GluA2 and corresponding ratio GluA2/GluA1; (**D**) NMDAR subunits GluN2A and GluN2B and corresponding ratio GluN2B/GluN2A. (**E**) Representative western blot images for (**A-D**) showing non-adjacent bands originating from the same blot. (**I**) Pearson correlation of PTZ kindled 5XFAD mice showing the relationship between NKCC1/KCC2 and A$\beta$ normalized to 5XFAD-Veh, (**F**) pTau AT100(Thr212, Ser14):Tau normalized to WT-Veh (**G**), and % spontaneous alternations in the Y-maze. Group comparisons for A$\beta$ ELISA, pTau western blots and the Y-maze were previously published [14] (**H**). Grey areas indicate 95% confidence interval for the two means. n = 12-19 WT-vehicle, 12-17 WT-PTZ, 12 5XFAD-vehicle and 17 5XFAD-PTZ. *$p<0.05$, **$p<0.01$. Females and males are designated by square and triangle data points, respectively, where sex effects were found.

**Figure 6. Differential effects of mTORC1 inhibition on seizure-induced excitation:inhibition imbalance in 5XFAD mice. (A-D)** Quantification of (**A**) GABA$_A$R$\alpha$1 and GABA$_A$R$\alpha$3 and corresponding GABA$_A$R$\alpha$1/GABA$_A$R$\alpha$3 ratio, (**B**) Cl⁻ cotransporters NKCC1 and KCC2 and corresponding NKCC1/KCC2 ratio, (**C**) AMPAR subunits GluA1 and GluA2 and corresponding ratio GluA2/GluA1, and (**D**) NMDAR subunits GluN2A and GluN2B and corresponding GluN2B/GluN2A ratio. (**E**) Representative Western blot images for (**A-D**) showing non-adjacent bands originating from the same blot. n = 12-13 for each group. *$p<0.05$, **$p<0.01$.

A   GABA$_A$R$\alpha$1   GABA$_A$R$\alpha$3   GABA$_A$R$\alpha$1/$\alpha$3

B   NKCC1   KCC2   NKCC1/KCC2

C   GluA1   GluA2   GluA1/GluA2

D   GluN2A   GluN2B   GluN2B/GluN2A

E

5XFAD

PTZ   -   -   +   +
Rapamycin   -   +   -   +   *kDa*

GABA$_A$R$\alpha$1 — *50*
GABA$_A$R$\alpha$3 — *60*
NKCC1 — *170*
KCC2 — *140*
GluA1 — *100*
GluA2 — *100*
GluN2A — *180*
GluN2B — *166*
$\beta$-actin — *42*

Fold change relative to 5XFAD Vehicle

5XFAD-Veh-Ctrl    5XFAD-Veh-Rapa    5XFAD-PTZ-Ctrl    5XFAD-PTZ-Rapa

| | | Prodromal 5XFAD | 7 month 5XFAD | Human |
|---|---|---|---|---|
| $GABA_AR$ | Hippocampus | —$GABA_AR$ α1<br>—$GABA_AR$ α2<br>⬇$GABA_AR$ α1/α2 | ⬇$GABA_AR$ α1<br>—$GABA_AR$ α3<br>⬇$GABA_AR$ α1/α3 | ⬇$GABA_AR$ α1<br>—$GABA_AR$ α3<br>⬇$GABA_AR$ α1/α3 |
| | Cortex | —$GABA_AR$ α1<br>—$GABA_AR$ α2<br>—$GABA_AR$ α1/α2 | ⬇$GABA_AR$ α1;<br>⬇GABAAR α3;<br>⬇$GABA_AR$ α1/α3 | |
| Cl⁻ cotransporters | Hippocampus | —NKCC1<br>—KCC2<br>—NKCC1/KCC2 | —NKCC1<br>—KCC2<br>—NKCC1/KCC2 | —NKCC1<br>—KCC2<br>—NKCC1/KCC2 |
| | Cortex | —NKCC1<br>—KCC2<br>—NKCC1/KCC2 | ⬇NKCC1<br>—KCC2<br>—NKCC1/KCC2 | |
| AMPAR | Hippocampus | —GluA1<br>—GluA2<br>—GluA1/GluA2 | —GluA1<br>⬇GluA2<br>—GluA1/GluA2 | —GluA1<br>⬆GluA2<br>⬇GluA1/GluA2 |
| | Cortex | —GluA1<br>—GluA2<br>—GluA1/GluA2 | —GluA1<br>⬇GluA2<br>—GluA1/GluA2 | |
| NMDAR | Hippocampus | —GluN2A<br>—GluN2B<br>—GluN2B/GluN2A | ⬇GluN2A<br>—GluN2B<br>—GluN2B/GluN2A | ⬆GluN2A<br>⬆GluN2B<br>—GluN2B/GluN2A |
| | Cortex | —GluN2A<br>—GluN2B<br>—GluN2B/GluN2A | ⬇GluN2A<br>—GluN2B<br>—GluN2B/GluN2A | |

**Table 1.** Effects of AD and 5XFAD genotype on E:I balance. Prodromal = 4 months old.

| | | 7 month 5XFAD | Human AD |
|---|---|---|---|
| GABA$_A$R | Hippocampus | —GABA$_A$R α1<br>—GABA$_A$R α 3<br>—GABA$_A$R α1/α3 | ↓GABA$_A$R α1<br>↓GABA$_A$R α 3<br>—GABA$_A$R α1/α3 |
| | Cortex | —GABA$_A$R α1<br>—GABA$_A$R α 3<br>—GABA$_A$R α1/α3 | |
| Cl$^-$ cotransporters | Hippocampus | ↑NKCC1<br>—KCC2<br>↑NKCC1/KCC2 | —NKCC1<br>↓KCC2<br>↑NKCC1/KCC2 |
| | Cortex | —NKCC1<br>—KCC2<br>—NKCC1/KCC2 | |
| AMPAR | Hippocampus | — GluA1<br>— GluA2<br>↑GluA1/GluA2 | —GluA1<br>—GluA2<br>—GluA1/GluA2 |
| | Cortex | — GluA1<br>— GluA2<br>↑GluA1/GluA2 | |
| NMDAR | Hippocampus | —GluN2A<br>— GluN2B<br>—GluN2B/GluN2A | —GluN2A<br>— GluN2B<br>—GluN2B/GluN2A |
| | Cortex | —GluN2A<br>— GluN2B<br>—GluN2B/GluN2A | |

**Table 2.** Effects of seizures in AD and 5XFAD mice.