Unveiling causal regulatory mechanisms through cell-state parallax

Alexander Po-Yen Wu[†1], Rohit Singh[†*1], Christopher Walsh[2], Bonnie Berger[*1,3]

[†] Contributed equally.

[*] Correspondence: rsingh@mit.edu, bab@mit.edu

Abstract

Genome-wide association studies (GWAS) identify numerous disease-linked genetic variants at noncoding genomic loci, yet therapeutic progress is hampered by the challenge of deciphering the regulatory roles of these loci in tissue-specific contexts. Single-cell multimodal assays that simultaneously profile chromatin accessibility and gene expression could predict tissue-specific causal links between noncoding loci and the genes they affect. However, current computational strategies either neglect the causal relationship between chromatin accessibility and transcription or lack variant-level precision, aggregating data across genomic ranges due to data sparsity. To address this, we introduce GrID-Net, a graph neural network approach that generalizes Granger causal inference to detect new causal locus−gene associations in graph-structured systems such as single-cell trajectories. Inspired by the principles of optical parallax, which reveals object depth from static snapshots, we hypothesized that causal mechanisms could be inferred from static single-cell snapshots by exploiting the time lag between epigenetic and transcriptional cell states, a concept we term "cell-state parallax." Applying GrID-Net to schizophrenia (SCZ) genetic variants, we increased variant coverage by 36% and uncovered noncoding mechanisms that dysregulate 132 genes, including key potassium transporters such as KCNG2 and SLC12A6. Furthermore, we discovered evidence for the prominent role of neural transcription-factor binding disruptions in SCZ etiology. Our work not only provides a strategy

---

[1] Computer Science and Artificial Intelligence Laboratory, MIT, Cambridge, MA 02139, USA
[2] Division of Genetics and Genomics and Howard Hughes Medical Institute, Boston Children's Hospital and Departments of Neurology and Pediatrics, Harvard Medical School, Boston, MA, 02115, USA
[3] Department of Mathematics, MIT, Cambridge, MA 02139, USA

for elucidating the tissue-specific impact of noncoding variants but also underscores the breakthrough potential of cell-state parallax in single-cell multiomics for discovering tissue-specific gene regulatory mechanisms.

Introduction

Dynamic changes in the chromatin states of cis-regulatory genomic elements influence gene expression in many biological processes, including development, differentiation, reprogramming, and perturbation response[1–3]. Correspondingly, genome-wide association studies (GWAS) have implicated thousands of noncoding loci in diseases and complex traits[4,5]. Yet, the challenge of linking these noncoding loci to the genes they regulate has impeded therapeutic advances. The common approach of simply associating a locus to its nearest gene is inadequate, as it overlooks distal locus-gene interactions. This is a critical limitation, as distal interactions are likely to be important for many genes. Meanwhile, experimental approaches to discover such interactions can be difficult to scale. Expression quantitative trait locus (eQTL) analyses require sample collections from large cohorts[6], and CRISPR perturbations have been limited to specific loci[7]. Computational methods based on bulk data have also been proposed, such as Wang et al.'s integration of multiple data types to link variants to genes[8], and Fulco et al.'s activity-by-contact (ABC) model[9,10] that predicts enhancers of a gene based on the activity and 3D chromatin structure around the gene. However, these models focus on the relative spatial positions of enhancers and genes, and are not designed to evaluate whether changes in accessibility at a locus actually influence gene expression.

Recent advances in single-cell multimodal technology, which can simultaneously profile chromatin accessibility (scATAC-seq) and gene expression (scRNA-seq) in the same cell, offer a promising opportunity to uncover tissue-specific, genome-scale, causal relationships between noncoding loci and genes in a single experiment[11–14]. However, current computational methods face substantial hurdles in extracting meaningful locus-gene associations. Existing approaches have diverse and often unrelated objectives: achieving concordance between the modalities[15,16], inferring gene regulatory networks by considering the accessibility of transcription-factor binding motifs[17,18], or using chromatin accessibility to improve RNA velocity inference[19]. Unfortunately,

owing to the extreme sparsity of scATAC-seq data (< 1% non-zero entries per cell [20]), these

approaches aggregate chromatin peaks within a defined range of a gene; for instance, MultiVelo

consolidates chromatin signals into a single, per-gene score[19]. This results in a loss of variant-

and locus-level resolution and also risks ignoring distal peaks. Moreover, these methods are

largely reliant on correlational analyses between scATAC-seq and scRNA-seq, thereby

overlooking the inherent causal time lag in gene regulation (**Figure 1a**). For instance, poised

enhancers, which play essential roles in development, undergo changes in chromatin states and

histone modifications *prior* to activating their target genes and would be missed by a

correlational analysis[21]. Consequently, the appropriate epigenetic and transcriptional readouts

to be cross-referenced may correspond to subtly different cell states.

We refer to this temporal discrepancy as "cell-state parallax," drawing inspiration from

the optical concept of parallax, which describes how different vantage points can unveil new

insights. Cell phone cameras with dual lenses leverage optical parallax by capturing multiple 2D

images of the same object from slightly offset positions to enable 3D depth perception.

Analogously, in single-cell multimodal assays, each modality captures a snapshot of the cell's

state, but at an offset due to the temporal lag between the epigenome and the transcriptome. By

exploiting this offset, causal relationships between the modalities can be unveiled. However, a

fundamental challenge in accounting for these offsets is that single-cell experiments destroy

cells upon measurement, so the temporal ordering of single-cell measurements is often unclear.

Moreover, an inference framework needs to be robust to the sparsity of both scATAC-seq and

scRNA-seq data[22,23]. Some previous methods mitigate sparsity by averaging transcriptionally

similar cells into pseudocells[11], but this comes at the cost of losing cell-level granularity which is

crucial for discerning subtle offsets between distinct modalities. In this work, we operate at a

per-peak and per-cell resolution, addressing sparsity through our machine learning framework.

To resolve single-cell multimodal snapshots into causal regulatory relationships, we

leverage mathematical principles that underlie the dynamics between causes and their effects.

We focus on Granger causality, a statistical framework which exploits the idea that a cause precedes, and is therefore predictive of, its downstream effect. There, a temporally causal relationship between two time-varying processes can be inferred solely by autoregressive analysis of time-resolved observational data. Granger causality has been a powerful tool in biology, econometrics, and other domains for causal discovery without the need to perform interventions.

However, the classical framework of Granger causality is limited to settings where observations can be ordered along a single time-axis. Mathematically, an ordering of observations can be categorized as *total* or *partial*: in the former, any pair of observations can be ordered unambiguously while for the latter only a subset of observations can be so ordered. Partial orderings characterize many dynamical systems, including disinformation cascades in social media, financial transaction networks, and parallel computing systems. In single-cell biology, when a differentiation trajectory has multiple branches, cell states across distinct branches are not directly comparable, necessitating a partial ordering of cells. Classical Granger causal analysis, being limited to total orderings, therefore needs to be adapted for inferring causal single-cell dynamics.

Our key mathematical advance is a generalization of Granger causality to partial orderings. We exploit the equivalence between such orderings and directed acyclic graphs (DAGs) and design a graph neural network framework to infer causality from the dynamics. We complement this with the conceptual advance of cell-state parallax from which we infer a DAG-based representation of single-cell multimodal dynamics. We combine these innovations into GrID-Net (Granger Inference on DAGs), a novel framework for Granger causal inference on DAGs that we apply to single-cell multimodal data for inferring causal noncoding locus-gene relationships (**Figure 1b,c**). Leveraging lagged message-passing on a graph neural network, GrID-Net resolves the cell-state parallax between the epigenome and transcriptome. We demonstrate that it outperforms existing state-of-the-art approaches for accurately identifying

noncoding locus-gene links across a variety of cell types and single-cell multimodal assays. We

then use GrID-Net to link fine-mapped genetic variants associated with schizophrenia (SCZ) to

the genes they affect. We uncover noncoding mechanisms of gene dysregulation that

prominently feature disruptions of neural transcription-factor binding and synaptic pathways as

putative drivers of SCZ disease pathogenesis (**Figure 1d**). Our findings demonstrate the power

of GrID-Net in uncovering tissue-specific causal mechanisms of disease and development,

using data generated from just one sample. Our work also illustrates the utility of multimodal

single-cell studies for investigating the impact of genetic variants on function and phenotype.

The code for GrID-Net can be found at https://github.com/alexw16/gridnet and the full list of

SCZ variant-gene links can be found in **Supplemental Table 1**.

Overview of GrID-Net

In general, inferring true causality can be a challenge as it requires perturbations to confirm

cause-and-effect relationships and validations to rule out potential confounders. Granger

causality is a powerful statistical concept that can *approximate* causality in time-varying

processes solely from observational data. It leans on the principle that a cause precedes its

effect[24,25]. Given two time-dependent variables x (a potential cause) and y (the effect), if the

forecast for y can be improved by accounting for the past values of x, then x is said to Granger

cause y since changes in x temporally precede those in y.

One of our key contributions is to broaden the set of domains to which Granger causal

analyses can be applied. Specifically, we generalize Granger causal inference to dynamical

systems that are characterized by partial orderings. To model the asynchrony between cause

and effect, we modify standard graph neural networks to allow for lagged message passing.

Each node (i.e., a cell state) in the directed acyclic graph accumulates information from its

parent nodes (past states), leveraging this lagged information to forecast the present state in the

context of the graph-structured dynamics. GrID-Net consists of two forecasting models: a

reduced model forecasts an effect y using only the past of y and a *full* model forecasts the

effect y using both the past of y and the past of the candidate causal variable x. The strength of

the Granger causal link between x and y can be evaluated by the statistical F-test which

quantifies the improvement in forecasting accuracy of the *full* model over that of the *reduced*

model.

GrID-Net enables the detection of asynchronous interactions between gene regulatory

elements along single-cell trajectories. For each single-cell multimodal dataset, a DAG of cell

states is constructed from a k-nearest neighbor graph of cells, with edges oriented in

accordance with pseudotime computed after a multimodal integration with Schema[20]. RNA-

velocity based edge orientations that capture local dynamics can also be supplied, so long as

any resulting cycles are broken. We evaluated the Granger causal strength between each gene

and chromatin accessible element ("peak") within one megabase (Mb) of the gene's

transcription start site, ranking the peak-gene pairs by their Granger causal strengths

(**Methods**).

GrID-Net more accurately identifies noncoding locus-gene links

We evaluated GrID-Net on four single-cell multimodal datasets (scATAC-seq and scRNA-seq)

that characterize a range of biological processes, including cell differentiation and drug

treatment response[11,12,26,27]. For each dataset, we sought to validate against expression

quantitative trait (eQTL) loci profiled in matching tissue (**Methods**). As eQTLs represent

associations between genetic variation at a locus and the corresponding changes in a gene's

expression, they serve as useful genome-wide proxies for perturbational validation of causal

interactions between genomic loci and genes.

For each gene, we intersected all ATAC-seq peaks within 1 Mb of its transcription start

site (TSS) with the set of eQTL variants linked to it, associating each peak with the most

significant variant overlapping it. We then evaluated the peak–gene pairs prioritized by GrID-Net

and two correlation-based methods used in previous work[11,27], Pearson correlation and

pseudocell correlation. The latter computes correlations on measurements aggregated across

multiple cells. We also evaluated the ABC model's predictions on bulk data from matching

tissues, comparing it against the predictions from the single-cell datasets (**Methods**).

We evaluated the accuracy of each method in retrieving the peak–gene pairs that

overlap significant eQTLs ($P < 10^{-10}$); our findings were robust to the choice of the eQTL

significance threshold (**Figure S1**). GrID-Net was significantly more accurate than the other

methods across all datasets (**Figure 2a**), exhibiting 7-27% higher area under the receiver

operating characteristic curve (AUROC) in 3 of the 4 datasets. GrID-Net demonstrated the

smallest improvement in AUROC for the sci-CAR dataset, although all methods performed

poorly on this dataset, likely due to the lower data quality of sci-CAR as the earliest single-cell

multimodal assay developed. Meanwhile, the correlation-based methods exhibited near-random

performance across all datasets, consistently underperforming not only GrID-Net but also the

bulk data-based ABC model. The low accuracy of predictions by correlation-based methods,

contrasted with the substantial improvement offered by GrID-Net, demonstrates the potential to

unravel much richer regulatory insights from these multimodal assays than has been possible.

We next assessed the robustness of the single-cell methods to sparsity in single-cell

data. GrID-Net better handled sparse data (**Figure S2**), yielding higher accuracies than the

correlation-based methods for peak–gene predictions involving genes with sparse data (non-

zero transcripts in fewer than 10% of cells) or sparse peaks (under 2% of cells). Notably,

pseudocell correlation failed to consistently outperform standard Pearson correlation on these

sparse peak–gene cases, even though the former is explicitly designed to compensate for the

sparsity of measurements at the level of individual cells. This may be because pseudocell

averaging reduces the effective number of independent observations and thus relinquishes the

statistical advantages of single-cell heterogeneity. Altogether, these analyses indicate not only

GrID-Net's robustness to the current sparsity of single-cell data but also the potential for its

increased effectiveness as the sensitivity of these multimodal assays continues to improve.

To further benchmark GrID-Net using functional indicators of noncoding locus–gene

interactions, we examined three-dimensional chromatin contacts (from bulk Hi-C data) between

gene promoters and noncoding loci. Compared to the two correlation-based methods, we found

that GrID-Net more accurately prioritized peak–gene pairs that were in frequent three-

dimensional contact (**Figure 2b)**. The ABC method uses bulk Hi-C data itself as an input so we

did not evaluate its agreement with chromatin contacts. In future, the GrID-Net framework could

be expanded to also utilize Hi-C data when available— as we describe later, incorporating one-

dimensional sequence proximity as a signal into GrID-Net improves its results. However, Hi-C

data is currently available only for a limited set of tissues[28] and the ability to make regulatory

inferences from just one multimodal assay enables the convenient study of rare cell types,

cancer mutations and drug responses etc.

GrID-Net reveals how chromatin accessibility primes gene activity

To better understand the biological basis of GrID-Net's superior accuracy, we applied it on a

single-cell multimodal dataset profiling human corticogenesis. We then honed in on peak–gene

pairs where GrID-Net and correlation-based methods disagree. Specifically, peak–gene pairs

were ranked according to the difference between their GrID-Net and correlation-based

prioritization scores (**Methods**), with higher ranking pairs being the ones that were more

strongly prioritized by GrID-Net. We then investigated the distinguishing biological

characteristics of these differentially-prioritized pairs.

One of our motivations behind GrID-Net is the observation that regions of chromatin

around lineage-determining genes become accessible prior to gene expression, foreshadowing

lineage choice[26,29]. GrID-Net was designed to resolve this cell-state parallax between chromatin

accessibility and gene expression and we assessed if it was able to do so. We first associated

each peak and each gene with the pseudotime corresponding to its highest value to determine

the lag associated with each peak-gene pair. For both correlation-based methods, we found that

peak–gene pairs differentially prioritized by GrID-Net are marked by significantly longer

pseudotime lags between chromatin accessibility and gene expression ($P < 10^{-26}$, Welch's one-

tailed t-test, **Figure 2c**, **Methods**). This result supports GrID-Net's ability to account for the

asynchrony between noncoding regulatory activity and downstream gene expression control

that are expected to characterize causal interactions between the two. To explore the regulatory

implications of this asynchrony, we obtained data on histone modification states of enhancers in

neural progenitor cells and cross-referenced it with ATAC-seq peaks[30]. In peak–gene pairs

differentially prioritized by GrID-Net, more poised enhancers (H3K4me1 only) overlapped with

the peaks than active enhancers (both H3K4me1 and H3K27ac)[31] (**Figure 2d**, **Methods**). This

was not the case for peak–gene pairs differentially prioritized by correlation-based methods.

Poised enhancers are regulatory sites that can trigger future gene expression changes but have

not yet affected expression[31,32]; in contrast, active enhancers correspond to ongoing

transcription[33]. This suggests that resolving cell-state parallax enables the detection of lagged

relationships between the activities of enhancers and their target genes. GrID-Net can thus

leverage the concept of chromatin regulatory potential[26] to facilitate a deeper understanding of

tissue-specific causal regulatory mechanisms.

To evaluate the biological significance of capturing these putative causal interactions, we

examined the functions of genes implicated in peak–gene pairs differentially prioritized by GrID-

Net. We found these genes to be highly enriched for gene sets associated with neurogenesis

regulation and neural progenitor cells ($FDR < 10^{-7}$, GSEA Preranked[34]), directly relating to the

process of corticogenesis profiled in this dataset. We emphasize that the enrichment score is

calculated only on the peak–gene pairs that GrID-Net prioritizes over the correlation-based

methods. In comparison, the set of genes from peak–gene pairs that *all* methods score highly is

functionally enriched for housekeeping genes rather than neuronal activity, suggesting that

GrID-Net's ability to detect poised enhancers is key to discovering cell-type specific regulatory

relationships. Gene expression control during differentiation has been reported to depend highly

on asynchronous enhancer-gene interactions[35]. The enrichment of these differentiation-related

gene sets thus underscores GrID-Net's ability to construct accurate genome-wide maps of

noncoding locus–gene links that go beyond correlative analyses and towards causality.

Linking variants to schizophrenia disease genes

The ability to link noncoding loci to the genes they regulate enables the discovery of molecular

mechanisms underlying complex diseases. Genome-wide association studies (GWAS) have

identified thousands of loci associated with diseases[36]. However, over 90% of GWAS variants

lie in noncoding regions[37], and interpreting the functions of these noncoding variants involves

the challenge of linking them to the specific genes that they regulate in a cell type-specific

manner. With GrID-Net, we can leverage data from a single-cell assay on matching tissue to

uncover these cell-type-specific variant-gene links and help bridge the gap between genetic

variants and gene dysregulation in disease.

We used GrID-Net to study variants associated with schizophrenia (SCZ), a psychiatric

disorder for which inherited genetic variants are believed to underlie its pathogenesis[38]. SCZ's

high prevalence of around 1%, its substantial morbidity and mortality, and the lack of effective

treatments are largely due to our limited understanding of SCZ pathophysiology[39,40]. We applied

the GrID-Net peak-gene links inferred from human corticogenesis[27], to interpret and analyze

variants reported in the largest GWAS of SCZ to date[41]. We refer to this original study as the

Trubetskoy, Pardiñas *et al*. study.

To boost GrID-Net's accuracy on shorter-range peak–gene interactions, we incorporated

one-dimensional genomic distance as an additional feature for predicting peak–gene

associations. We observed a higher likelihood of peak–gene interaction if the two are located

close to each other, in line with reported patterns of genomic distance-dependent regulatory

effects[42,43]. We calibrated the relative importance of this feature on a dataset profiling mouse

skin differentiation[26] where we fitted a logistic regression model that combines the raw GrID-Net

score with a genomic distance score (**Methods**). We found that integrating genomic distance

with GrID-Net improved predictions of peak–gene links for mouse skin cells. The discriminative

power on short-range peak–gene interactions improved, while the ability to detect long-range

interactions was preserved. Furthermore, this model trained on mouse skin cell data performed

well across species and cell types in human corticogenesis as well (**Figure S3**). This genomic

distance-dependent model's scores were used for all remaining analyses of the human

corticogenesis dataset.

From the SCZ GWAS study, we retrieved all fine-mapped genetic variants with posterior

probability (*PP*) over 5%. We intersected these variants with peaks from the human

corticogenesis dataset. We then generated an initial set of candidate SCZ genes by associating

each variant with up to 3 of its overlapping peak's highest scoring genes, retaining only variant–

gene associations with scores in the top 50% of all peak–gene scores. This approach yielded a

set of 132 genes that not only is enriched for genes implicated in the original SCZ GWAS study

(*FDR* = 0.037, Enrichr[44]) but also includes novel SCZ gene candidates (**Supplemental Table

1**). For example, of the 12 SCZ variants that were prioritized by both GrID-Net and Trubetskoy,

Pardiñas *et al*., 9 were linked to the same target genes by both approaches (**Figure 3a)**. This

overlap included the association between rs324017 and NAB2, a gene implicated in stress-

induced neuroprotective maintenance that also is the downstream target of the SCZ drug

cinnarizine[45,46]. For these same 12 SCZ variants, GrID-Net also proposed 20 additional SCZ

candidate genes, of which 12 (60%) were found to be specifically expressed in brain tissues and

thus potentially relevant in SCZ etiology (**Methods**). Some of these additional candidates

include NXPH4, a neurexophilin involved in synapse function and cerebellar motor control[47],

and EPN2, an epsin linked to neural fate commitment[48]. Moreover, of the 78 SCZ variants with

PP over 50%, only 22 (28%) were linked to genes by the original SCZ GWAS study (**Figure 3b**).

Inclusion of SCZ variants that were linked to genes by GrID-Net increased this total to 30 (38%)

SCZ variants. GrID-Net can, therefore, be used to complement standard strategies for linking

variants to genes, such as those used in the original SCZ GWAS study. It can propose gene

associations for variants with no annotations and also augment existing annotations by

suggesting additional genes that a variant may be affecting.

In particular, GrID-Net helps address a key challenge in GWAS analyses: the common

strategy of assigning GWAS variants to their closest gene[49] misses many biologically

meaningful variant–gene relationships. By leveraging single-cell heterogeneity, GrID-Net can

evaluate longer range interactions. Many of the variant–gene links we identify are characterized

by large genomic distances between the variant and the transcription start site (TSS) of the

linked gene. Moreover, the variants are often linked to genes that are multiple genes over from

their closest gene (**Figure 3c**). For example, DCP1B and NCKAP5L, each of which are

connected to only a single variant, are separated from their respective variants by 18 and 11

genes (235 and 242 kilobases (kb)), respectively. DCP1B has been implicated in SCZ,

hypomania, and memory performance[50,51], while NCKAP5L, a cytoskeletal gene involved in

neuronal activity[52], has previously been associated with SCZ as well as amyotrophic lateral

sclerosis[53], indicating possible roles for these genes in neuron-specific dysregulation in SCZ.

In addition, 52 candidate genes (39.4% of the overall set) were associated with more

than one variant (**Figure 3d**). These genes may be controlled by the coordinated activity of

multiple regulatory elements and co-regulators relevant to SCZ[54]. IGSF9B was linked to 7

noncoding SCZ variants – the most of any gene. Coding variants in IGSF9B have been found to

be associated with major depression and the negative symptoms of SCZ[55,56], underscoring its

role in regulating affective symptoms in SCZ. Similarly, KCNG2, a potassium channel-encoding

gene, was linked to 6 noncoding SCZ variants, and has been proposed as a drug target for SCZ

given the known disruption of KCN-related pathways in SCZ[40,57,58]. Neither of these genes were

prioritized in the original GWAS analysis.

We also assessed the cell-type specificity of the genes linked to SCZ variants. We found

that 104 candidate SCZ genes (78.8% of the overall set) are differentially expressed across the

cell types profiled in the human corticogenesis dataset (Welch's *t*-test with Benjamini-Hochberg

correction *P* < 0.05, **Figure 3e**). Further analysis of the SCZ variants also revealed that 59% of

the variants lie in intronic regions, while only 7% of variants lie in intergenic regions more than 5

kb from any gene (**Figure 3f**, **Methods**). As enhancers with tissue-specific effects are highly

enriched in introns while intergenic enhancers more often regulate housekeeping genes[59], these

findings support the potential cell type-specificity of the regulatory disruptions underlying SCZ

pathogenesis.

Transcription factor binding disruptions are implicated in schizophrenia

Transcription factor (TF) binding at the cis-regulatory elements located in noncoding genomic

loci is a key mechanism of gene regulation. We observed that SCZ variants with high *PP* are

associated with larger TF binding affinity changes (**Figure S4a**). To investigate the

dysregulatory role of SCZ variants prioritized by GrID-Net, we therefore evaluated their effect on

TF binding. For each variant associated with a gene by GrID-Net, we calculated the difference

in the predicted TF binding affinity between the reference allele and the SCZ risk allele

(**Methods**, **Supplemental Table 2**). Of the 115 such SCZ variants, 28 (these were associated

with 49 target genes) were predicted to cause a 20-fold or stronger change in at least one TF's

binding affinity (**Figure 4a**). The TFs we identified as having the most widespread binding

affinity changes are known to be implicated in SCZ, exemplifying GrID-Net's ability to deliver de

novo mechanistic insight. VEZF1, a vasculature-related TF reported to play a role in SCZ

dysregulation[60], displayed diminished binding affinity most frequently (in 5 variants). In addition,

KLF15 and KLF6, two Kruppel-like family (KLF) TFs known to regulate neuronal growth and

linked to SCZ[61,41], were each associated with reduced binding affinity in 4 variants (**Figure S4b**).

Meanwhile, the TFs with the most widespread increase in binding affinity – SP1, SP3, and

KLF16 – have also been reported to be disrupted in SCZ[62,63] (**Figure S4c**). More broadly, of the

TFs with a 20-fold binding affinity change in at least one variant, 57.9% (=33/57) have cell type-

specific expression patterns in corticogenesis (Welch's *t*-test with Benjamini-Hochberg

correction $P < 0.05$). Collectively, these results point to genetic disruption of neuron-specific

regulatory elements as potential drivers of SCZ pathogenesis.

We reasoned that if the TF binding affinity changes at a SCZ variant are phenotypically

meaningful, they should result in expression changes of the corresponding gene(s) (as identified

by GrID-Net). Accordingly, we obtained gene expression data from glial progenitor cells (GPCs)

produced from SCZ patient-derived induced pluripotent cells, as well as matching controls[58,64],

and computed differential expression patterns in SCZ GPCs. Focusing on glial differentiation

genes, we observed that the differential expression of a gene in SCZ GPCs was indeed

significantly correlated with TF binding affinity changes at the corresponding SCZ variant

(Pearson correlation coefficient = 0.59, **Figure 4b**, **Methods**). The consistency between these

independent observations provides support to our overall analytic approach and underscores

the potential to leverage GrID-Net for examining causal, gene-specific regulatory mechanisms.

We drilled down into specific genes reported by our analysis, focusing on those linked to

SCZ variants associated with at least 20-fold TF binding affinity changes (**Figure 4c**). NAB2, the

aforementioned gene implicated in neuroprotective processes[46], was linked to rs324017, which

contributes to a 200-fold decline in binding affinity for the TF PURA. Intriguingly, PURA is also

critical for neural function and development, and its dysfunction is implicated in several

neurological and cognitive impairments[65,66]. In addition, rs12991836 was linked to ZEB2, whose

TSS is positioned 137 kb from the variant. ZEB2 has been previously reported as a target of

long-range regulation, and heterozygous ZEB2 mutations are known to cause Mowat-Wilson

syndrome, which features structural brain abnormalities and intellectual disability[67].

Furthermore, GrID-Net prioritized two potassium transport genes - KCNG2 and

SLC12A6 - a class of genes which has been associated with compromised potassium uptake

when dysregulated[58]. This phenotype is characteristic of not only SCZ but also other neuron-

related diseases, such as Huntington's disease, multiple sclerosis, and hemiplegic migraine[64,68].

KCNG2 was linked to 6 SCZ variants, the second most of any gene as described earlier. Using

TF binding affinity changes to hone in on the most relevant variants, we identified rs8091497 as

the variant with the largest TF binding affinity change. rs8091497 was associated with a 40-fold

decline in binding affinity for SP1, which itself has been linked to behavioral abnormalities in

SCZ[63]. As for SLC12A6, its corresponding variant (rs117799466) increases the binding affinity

for many TFs with repressive capabilities. The three TFs with the greatest binding affinity gains

at this locus – SP1, ZN281, and MAZ – all play a role in neuronal differentiation[63,69,70], and both

SP1 and ZN281 have also shown increased expression levels in SCZ as well[71,72]. Combined

with their increased binding affinity incurred by rs117799466, the upregulation of these TFs may

contribute to the repression of SLC12A6, an outcome that has been previously linked to the

potassium uptake impairments that characterize SCZ glia[58]. Together, these findings suggest a

possible mechanism for SLC12A6 downregulation in SCZ that implicates enhanced binding of

neuron-related repressors to a SLC12A6-targeting enhancer to reduce its expression levels.

We then broadened our analysis to gene modules, exploring if the genes identified by

GrID-Net are systematically involved in SCZ-related pathways. The 14 SCZ variants that cause

at least 40-fold changes in TF binding affinity are linked by GrID-Net to 26 genes, whose

differential expression patterns we examined in SCZ GPCs. We detected 3 distinct gene

modules that are co-expressed with several of these genes (**Figure 4d**, **Methods**). Module A

(SLC12A6, DLX1, and DLX2) is enriched for genes related to noradrenergic neuron

differentiation (*FDR* < 0.05) and morphogenesis (*FDR* < 0.02); Module B (MFAP4, STAT6,

NXPH4) is enriched for extracellular matrix organization (*FDR* < 10^{-14}) and angiogenesis

regulation (*FDR* < 0.02) genes, and Module C (EPN2, P2RX7, ICA1) is enriched for synaptic

transmission (*FDR* < 10^{-4}) and postsynaptic membrane potential (*FDR* < 0.05) (Enrichr[44]). Each

of these pathways has been previously associated with SCZ. Noradrenergic neurons have long

been implicated in disrupted neuromodulation in SCZ[73], neuronal angiogenesis underlies the

vascular abnormalities observed in SCZ[74], and genetic studies have pointed to the critical role of

synaptic function in SCZ[41]. These pathways, therefore, enable the interpretation of SCZ variants

beyond the dysregulation of the specific genes that they directly affect, presenting an

opportunity to discover systems-level mechanisms of SCZ pathogenesis.

Discussion

Most methods for single-cell multimodal data integration have focused on synchronizing

modalities to improve downstream tasks, such as cell type identification. In contrast, we

demonstrate it is also powerful to systematically study the asynchrony between the modalities.

We presented GrID-Net, a method to uncover causal connections between genes and

noncoding genomic loci using co-assayed single-cell RNA-seq and ATAC-seq data. GrID-Net

operates at a genome-wide scale, is tissue-specific, and can effectively detect long-range (up to

1 Mb) cis-regulatory associations. A longstanding challenge with GWAS studies has been to

identify the specific genes that are dysregulated by noncoding disease-linked variants. With

GrID-Net, a single commercially-available assay of the matching tissue can power a genome-

wide estimation of variant-gene relationships. By resolving tissue-specific causal variant-gene

associations, GrID-Net can help identify disease-linked genes as therapeutic targets.

When studying a particular disease, GrID-Net should be paired with a single-cell assay of a

matching tissue undergoing a dynamic biological process, such as development, differentiation,

disease, or drug response. GrID-Net can then translate the results of this single assay to

genome-wide causal hypotheses specific to the disease. We demonstrate this by leveraging a

single-cell assay of human corticogenesis to interpret the functional effects of schizophrenia

(SCZ) variants. GrID-Net expands both the breadth and depth of existing gene prioritization

analyses: it finds novel associations for many unlinked variants while also uncovering additional

causal mechanisms for previously investigated variants. For instance, we identified SCZ

variants causing large (>20-fold) changes in TF binding affinity and connected them to 49

genes. Many of these genes are located distally from the SCZ variants that are predicted to

affect them and would have been neglected by existing GWAS variant–gene associations

analyses that consider only genomic proximity. Furthermore, by correlating TF binding affinity

changes with differential expression in SCZ glial progenitor cells, we predicted possible

mechanisms of TF-driven dysregulation for genes like KCNG2 and SLC12A6. Combined with an

exploration of SCZ-related pathways associated with GrID-Net-identified genes, these analyses

provide a comprehensive strategy for unveiling mechanistic insights about SCZ pathogenesis.

Such analyses can also be applied to investigate other diseases.

Our work rests on two conceptual insights that are broadly applicable. First, when

different modalities are measured in the same cell their causal relationship implies a temporal

offset between them, a phenomenon we refer to as "cell-state parallax". Here, we exploit the

parallax between RNA-seq and ATAC-seq, but such parallax can also be leveraged from other

co-assayed modalities such as chromatin structure or methylation. We view RNA velocity[75] and

protein velocity[76] as specific applications of the broader concept of cell-state parallax, the

distinction being that velocity uses pre-defined mappings between modalities (e.g., spliced vs.

unspliced transcript-counts for a particular gene) to infer the differentiation direction of each cell.

In contrast, such mappings between modalities may be unclear in general cell-state parallax

and require customized inference approaches. Accordingly, our second conceptual insight is to

develop computational frameworks that can accommodate the cross-modality dynamics of cell

differentiation. Much of the existing work on modeling gene expression dynamics utilizes

mathematical and statistical tools designed for time-series analysis[77,78], which make the

unrealistic assumption that a total ordering of cells can capture the underlying biology. We

argue that a partial ordering of cells is necessary to accurately model forks and merges in

trajectories and accordingly formulate the differentiation landscape as a directed acyclic graph

of cells that can represent such an ordering. We expect a promising direction of future research

will be computational inference techniques, like GrID-Net, that can operate under this more

realistic model.

Since GrID-Net operates on observational data, the locus-gene associations it infers are

causal hypotheses that need to be validated by perturbations. In the eQTL-based evaluation, we

sought to mimic such validations by investigating random genetic variants observed in the

population and cross-referencing them to observed expression changes. We also note that such

causal interactions are likely indirect and mediated by additional biophysical events, such as

transcription factor binding. Nonetheless, indirect causality is biologically and therapeutically

valuable, as the underlying mechanisms can still be exploited via interventions to achieve the

desired phenotype.

With the growing availability of single-cell multimodal datasets, our approach is positioned to

serve as a key framework for unlocking new mechanisms of gene regulatory dynamics that

underlie fundamental biology and disease. Moreover, GrID-Net can be applied broadly to other

domains featuring graph-based dynamics, such as uncovering the drivers of social media

disinformation cascades. A key technical advance of this work is generalizing the applicability of

standard Granger causal analysis to a wider set of domains than previously possible. Doing so

required the use of graph-theoretic deep learning techniques to implement the core statistical

intuition of Granger causality for any DAG-structured ordering of observations. Our work, like

others before us[79], thus highlights how classical statistical concepts can be reimagined with

modern deep learning techniques to dramatically widen their applicability.

Methods

GrID-Net formulation

See Supplementary Note.

GrID-Net training details and hyperparameters

All GrID-Net models consisted of $L = 10$ GNN layers and were trained using the Adam optimizer with a learning rate of 0.001 and a mini-batch size of 1024 candidate peak-gene pairs. Models were trained for a maximum of 20 epochs or until the relative change in the loss function was less than $0.1/|P|$ between successive epochs. Here, $|P|$ denotes the number of candidate peak-gene pairs evaluated per dataset. GrID-Net models were implemented in PyTorch and trained on a single NVIDIA Tesla V100 GPU.

Single-cell multimodal datasets

We used four single-cell multimodal datasets that spanned a range of different multimodal profiling techniques: 1) a 10x Multiome dataset profiling human corticogenesis, 2) a SHARE-seq dataset profiling mouse skin differentiation, 3) a SNARE-seq dataset profiling mouse neonatal cerebral cortical cells, 4) a sci-CAR dataset profiling A549 cells treated with dexamethasone. For each of these datasets, we followed recommended practice[80] and applied $log\,(1 + CPM/100)$ and $log\,(1 + CPM/10)$ transformations to the RNA-seq and ATAC-seq counts data, respectively, where CPM indicates counts per million. We then filtered out all genes that were represented in fewer than 1% of cells in the dataset and all peaks that were represented in fewer than 0.1% of cells. We assembled a set of candidate peak-gene pairs by selecting all peaks with centers within 1 Mb from each gene's TSS. For the 10x Multiome dataset, we only considered peak-gene pairs that were separated by 250 kb to cohere with the genomic distance

threshold used in the original study[27]. To consider multiple samples for each dataset, we

generated 5 uniformly sampled sets of 5,000 cells for each of the 10x Multiome, SHARE-seq,

and SNARE-seq datasets, while we generated 5 uniformly sampled sets of 3,000 cells for the

sci-CAR dataset due its smaller quantity of cells (3,081 cells).

Directed acyclic graph (DAG) construction

For each sample of a dataset, we first generated per-cell representations for each of the gene

expression and chromatin accessibility modalities. Gene expression representations were

constructed by (1) scaling the transformed RNA-seq data to unit variance and zero mean, (2)

retaining the top 2000 most highly variable genes, (3) and projecting these normalized

expression values onto their top 100 principal components (excluding components that have

Spearman correlation $\rho > 0.9$ with the total counts per cell. Chromatin accessibility

representations were obtained transforming the ATAC-seq data using tf-idf[81,82] and projecting

the resulting values onto the top principal components as in (3) above. We then used Schema[20]

to learn cell representations that unify information from these gene expression and chromatin

accessibility representations. We constructed a k-nearest neighbor graph ($k = 15$) based on the

Euclidean distances between these cell representations and inferred pseudotime using the

scanpy[83] implementation of diffusion pseudotime[84] with default parameters. We then removed

edges in the k-nearest neighbor graph if they did not align with the direction of increasing

pseudotime, thus ensuring that no cycles exist and the resulting graph is a DAG.

Alternative methods

Pearson correlation. We computed the Pearson correlation between the transformed ATAC-seq

and RNA-seq values for each candidate peak-gene pair, and the absolute value of the

correlation coefficients were used to score the peak-gene pairs.

Pseudocell correlation. For each sample, we identified the 49 nearest neighbors for each cell

based on the Euclidean distances between the joint representations learned from Schema (see

"Directed acyclic graph construction"). We constructed a "pseudocell" from each cell and its 49

nearest neighbors, constructing its RNA-seq and ATAC-seq profiles by summing the raw counts

from each of the modalities for all cells included in the pseudocell. We then applied $log\,(1+$

$CPM/100)$ and $log\,(1+CPM/10)$ transformations to the RNA-seq and ATAC-seq counts data,

respectively. Pearson correlation was applied to the transformed ATAC-seq and RNA-seq

values for each candidate peak-gene pair, and the absolute value of the correlation coefficients

were used to score the peak-gene pairs.

Activity-by-contact (ABC) model. We obtained ABC prediction scores in cell types matching the

cell types profiled in the single-cell multimodal studies. For each single-cell study, we

intersected the candidate peak-gene pairs with the ABC enhancer-gene pairs, assigning each

peak-gene pair the corresponding ABC score. If no ABC enhancer-gene pair overlap was

identified, the peak-gene pair was assigned a value of 0, as the ABC prediction scores were

available only for enhancer-gene pairs that exceeded the threshold of 0.015 defined in the

original study[10]. For single-cell studies that profiled mouse cells, we mapped ABC enhancer-

gene pairs in matching human cell types to the mouse genome due to the strong conservation

of epigenomes between human and mouse[85,86].

eQTL evaluations

We obtained eQTL data from the eQTL Catalogue[87] and used eQTL data specific to the cell

type profiled in each single-cell multimodal study. For mouse tissues, we mapped eQTL results

for matching human tissues to the mouse genome, as human and mouse epigenomes show

strong conservation[85,86].

We first intersected eQTL variant-gene pairs with peak-gene pairs from the single-cell

data and assigned each remaining peak-gene pair the most statistically significant eQTL. We

then binned these peak-gene pairs according to the genomic distance between each gene's

TSS and each peak's center, creating a bin for each of the following ranges of genomic

distances: [1 bp, 1 kb), [1 kb, 10kb), [10kb, 100kb), [100kb, 1 Mb). We identified all peak-gene

pairs that overlapped significant eQTL variant-gene pairs (P < 0.001), and these peak-gene

pairs served as the positive examples. To generate a distance-matched set of negative

examples, we determined the number of positive examples in each genomic distance bin and

selected an equal number of non-significant eQTL variant-gene pairs with the highest P values

for that bin.

Chromatin interactions evaluations

We obtained Hi-C data for cell types matching the 10x Multiome, SNARE-seq, and sci-CAR

datasets[88–90]. We used 5C data for the SHARE-seq dataset[91] due to the unavailability of mouse

skin Hi-C data. Chromatin interactions were mapped to a peak-gene pair by intersecting the

genomic windows in the chromatin interactions data with the peak and the TSS of the gene. For

Hi-C data, peak-gene pairs that were associated with the top 1% of chromatin contact

frequencies were treated as positive examples, while the remaining peak-gene pairs were

considered negative examples. For the 5C dataset, which evaluated the statistical significance

of chromatin loops, peak-gene pairs that were associated with significant looping interactions (P

< 0.001) were treated as positive examples, and the remaining peak-gene pairs were

considered negative examples.

GrID-Net vs. correlation peak-gene prioritization

For the human corticogenesis dataset, we averaged the rankings of all candidate peak-gene

pairs across the five trials for each of the GrID-Net, Pearson correlation, and pseudocell

correlation methods. We then calculated the difference between GrID-Net's average rankings

for each peak-gene pair with those of the Pearson correlation and pseudocell correlation

methods. The peak-gene pairs with the most positive differences represent the peak-gene pairs

that GrID-Net prioritizes over the method against which GrID-Net is being compared. To

evaluate enrichment of gene sets in peak-gene pairs prioritized by GrID-Net, genes were ranked

with respect to these differences and enrichment was computed using GSEA Preranked[34].

Enhancer states

We obtained H3K27ac and H3K4me1 ChIP-seq data for all samples of human neural progenitor

cells and fetal neural cells in ChIP-Atlas [30]. We identified statistically significant ChIP-seq peaks

(MACS2 $Q < 10^{-10}$) that were positioned fully within ATAC-seq peaks from the human

corticogenesis dataset. Active enhancers were defined as ATAC-seq peaks containing

overlapping H3K4me1 and H3K27ac ChIP-seq peaks, while poised enhancers were defined as

ATAC-seq peaks containing H3k4me1 ChIP-seq peaks and no H3K27ac ChIP-seq peaks.

Genomic distance-dependent GrID-Net scores

We constructed a general model that integrated GrID-Net scores with genomic distance to

account for known patterns of genomic distance-dependent transcription factor (TF) regulatory

effects[42,43]. The model assumes that TF regulatory effects exponentially decay with increasing

genomic distance. To learn the decay rate, we trained a logistic regression model to predict

peak-gene pairs overlapping significant skin eQTLs ($P < 10^{-3}$) for the SHARE-seq mouse skin

dataset.

$$\text{logit}(\pi_i) = \log(\frac{\pi_i}{1 - \pi_i}) = \beta_1 \log(s_i) + \beta_2 \hat{d_i}$$

Here, π_i is the probability of peak-gene pair i being a positive example, s_i corresponds to the

mean rankings of the GrID-Net scores across trials for the peak-gene pair, and $\widehat{d_i} =$

$max(log_{10}(d_i) - 4, 0)$, where d_i is the genomic distance in base pairs (bp) between the center

of the peak and the TSS of the gene in the peak-gene pair. This formulation assumes that only

distal interactions at genomic distances greater than 10 kb are affected by the genome

distance-dependent TF regulatory effect.

Positive and negative examples were chosen as described in "eQTL evaluations". We

held out one-third of the example as a test set, and then performed nested 3-fold cross

validation on the remaining examples, selecting the model that yielded the highest AUROC

across the folds. This model increased the AUROC on the test set compared to the predictions

from the averaged GrID-Net rankings (**Figure S3a**). We applied this pre-trained model on the

human corticogenesis dataset and found that this model outperformed the baseline predictions

from the averaged GrID-Net rankings (**Figure S3b**). These results demonstrated the

effectiveness of incorporating genomic distance as an additional feature alongside scores

learned from the single-cell multimodal data.

Linking SCZ GWAS Variants to Genes

We obtained FINEMAP posterior probabilities (*PP*) calculated in the original SCZ GWAS

study[41]. We retained all fine-mapped variants with $PP \geq 0.05$ that overlapped peaks from the

human corticogenesis dataset. We identified the peak that matched each SCZ variant and,

using the genomic distance-dependent GrID-Net scores, we linked the variant to up to 3 of the

peak's highest scoring genes, retaining only peak-gene pairs in the top 50% of all peak-gene

scores.

Brain-specific gene expression

We obtained consensus tissue-specific transcript levels from the Human Protein Atlas[92,93]. For a

given gene, we first calculated the mean and standard deviation of the gene's transcript levels

across all non-brain tissues. The brains-specific tissues were identified to be those from the

basal ganglia, cerebellum, cerebral cortex, medulla oblongata, midbrain, pons, and white

matter. We then converted the transcript level for the gene in each of the brain-specific tissues

into a z-score with respect to the non-brain tissues ($z = \frac{x - \mu_{non-brain}}{\sigma_{non-brain}}$, where x is the transcript

level for the gene in a brain-specific tissue). Genes that had a z-score over 1 in any of the brain-

specific tissues were annotated as having brain-specific gene expression.

TF binding affinity changes

We obtained the reference and SCZ risk alleles for all SCZ variants from the original SCZ

GWAS study[41]. We then flanked each of the alleles with the genomic sequences $\pm$20 bp of the

SCZ variant position using the GRCh38/hg38 reference genome[94], generating a reference

sequence and SCZ variant sequence. We applied FIMO[95] to each of these sequences to predict

TF binding affinities using position weight matrices (PWMs) from HOCOMOCO v11 (Full

Collection)[96], JASPAR CORE (Non-Redundant) v2[97], and CIS-BP v2[98]. We evaluated the effect

of a specific SCZ variant on a TF's binding affinity by comparing the FIMO-calculated *p*-values

for the TF between the reference sequence and SCZ variant sequence.

SCZ GPC gene expression

We obtained bulk RNA-seq data from glial progenitor cells (GPCs) produced from 4 SCZ

patient-derived induced pluripotent cells (iPSCs) and paired controls subejcts[58,64], with at least 3

samples per patient. We applied a $log(1 + CPM)$ transformation to this data.

To evaluate the relationship between TF binding affinity changes caused by SCZ

variants to expression changes in GPCs, we first determined peak-gene pairs that are specific

to glial differentiation. In the human corticogenesis dataset, we identified genes that were

differentially expressed between glial cells and the cycling progenitor cells, which were the root

of the corticogenesis trajectory ($P < 0.05$, Welch's *t*-test). We then used Welch's *t*-test to

compare the expression levels of these glial differentiation-specific genes for each SCZ patient

against all of the control samples. For each gene, we identified the most statistically significant

difference so as to account for all genes that may be differentially expressed in any of the

patients compared to the controls.

SCZ GPC co-expression pathways

We first identified a set of 26 genes linked to SCZ variants causing at least a 40-fold change in

binding affinity for any TF. We identified 3 sets of these SCZ genes that were correlated with

each other (Pearson correlation coefficient ≥ 0.6) in the SCZ GPCs and paired controls

(SLC12A6, DLX1, DLX2; MFAP4, STAT6, NXPH4; EPN2, P2RX7, ICA1). For each of these

sets of SCZ genes, we then determined gene modules that may be implicated in their

dysregulation by identifying genes that are highly co-expressed (Pearson correlation coefficient

≥ 0.8) with any of the SCZ genes in the set. We applied gene set enrichment analysis using

Enrichr to detect biological processes that were enriched in each of these modules.

References

1. Buenrostro, J. D. *et al.* Integrated Single-Cell Analysis Maps the Continuous Regulatory Landscape of Human Hematopoietic Differentiation. *Cell* **173**, 1535–1548.e16 (2018).

2. Papp, B. & Plath, K. Epigenetics of Reprogramming to Induced Pluripotency. *Cell* vol. 152 1324–1343 Preprint at https://doi.org/10.1016/j.cell.2013.02.043 (2013).

3. Pierce, S. E., Granja, J. M. & Greenleaf, W. J. High-throughput single-cell chromatin accessibility CRISPR screens enable unbiased identification of regulatory networks in cancer. *Nat. Commun.* **12**, 2969 (2021).

4. Cano-Gamez, E. & Trynka, G. From GWAS to Function: Using Functional Genomics to Identify the Mechanisms Underlying Complex Diseases. *Front. Genet.* **11**, 424 (2020).

5. Albert, F. W. & Kruglyak, L. The role of regulatory variation in complex traits and disease. *Nature Reviews Genetics* vol. 16 197–212 Preprint at https://doi.org/10.1038/nrg3891 (2015).

6. Nica, A. C. & Dermitzakis, E. T. Expression quantitative trait loci: present and future. *Philosophical Transactions of the Royal Society B: Biological Sciences* vol. 368 20120362 Preprint at https://doi.org/10.1098/rstb.2012.0362 (2013).

7. Shukla, A. & Huangfu, D. Decoding the noncoding genome via large-scale CRISPR screens. *Curr. Opin. Genet. Dev.* **52**, 70–76 (2018).

8. Wang, Q. *et al.* A Bayesian framework that integrates multi-omics data and gene networks predicts risk genes from schizophrenia GWAS data. *Nat. Neurosci.* **22**, 691–699 (2019).

9. Fulco, C. P. *et al.* Activity-by-contact model of enhancer-promoter regulation from thousands of CRISPR perturbations. *Nat. Genet.* **51**, 1664–1669 (2019).

10. Nasser, J. *et al.* Genome-wide enhancer maps link risk variants to disease genes. *Nature* vol. 593 238–243 Preprint at https://doi.org/10.1038/s41586-021-03446-x (2021).

11. Cao, J. *et al.* Joint profiling of chromatin accessibility and gene expression in thousands of

single cells. *Science* **361**, 1380–1385 (2018).

12. Chen, S., Lake, B. B. & Zhang, K. High-throughput sequencing of the transcriptome and chromatin accessibility in the same cell. *Nat. Biotechnol.* **37**, 1452–1457 (2019).

13. Argelaguet, R., Cuomo, A. S. E., Stegle, O. & Marioni, J. C. Computational principles and challenges in single-cell data integration. *Nat. Biotechnol.* **39**, 1202–1215 (2021).

14. Efremova, M. & Teichmann, S. A. Computational methods for single-cell omics across modalities. *Nat. Methods* **17**, 14–17 (2020).

15. Welch, J. D. *et al.* Single-Cell Multi-omic Integration Compares and Contrasts Features of Brain Cell Identity. *Cell* **177**, 1873–1887.e17 (2019).

16. Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888–1902.e21 (2019).

17. Aibar, S. *et al.* SCENIC: single-cell regulatory network inference and clustering. *Nat. Methods* **14**, 1083–1086 (2017).

18. Kamimoto, K., Hoffmann, C. M. & Morris, S. A. CellOracle: Dissecting cell identity via network inference and in silico gene perturbation. *bioRxiv* 2020.02.17.947416 (2020) doi:10.1101/2020.02.17.947416.

19. Li, C., Virgilio, M. C., Collins, K. L. & Welch, J. D. Multi-omic single-cell velocity models epigenome–transcriptome interactions and improves cell fate prediction. *Nat. Biotechnol.* **41**, 387–398 (2022).

20. Singh, R., Hie, B. L., Narayan, A. & Berger, B. Schema: metric learning enables interpretable synthesis of heterogeneous single-cell modalities. *Genome Biol.* **22**, 131 (2021).

21. Crispatzu, G. *et al.* The chromatin, topological and regulatory properties of pluripotency-associated poised enhancers are conserved in vivo. *Nat. Commun.* **12**, 4344 (2021).

22. Chen, H. *et al.* Assessment of computational methods for the analysis of single-cell ATAC-seq data. *Genome Biol.* **20**, 241 (2019).

23. Hou, W., Ji, Z., Ji, H. & Hicks, S. C. A systematic evaluation of single-cell RNA-sequencing imputation methods. *Genome Biol.* **21**, 218 (2020).

24. Granger, C. W. J. Investigating Causal Relations by Econometric Models and Cross-spectral Methods. *Econometrica* vol. 37 424 Preprint at https://doi.org/10.2307/1912791 (1969).

25. Shojaie, A. & Fox, E. B. Granger Causality: A Review and Recent Advances. *Annual Review of Statistics and Its Application* vol. 9 289–319 Preprint at https://doi.org/10.1146/annurev-statistics-040120-010930 (2022).

26. Ma, S. *et al.* Chromatin Potential Identified by Shared Single-Cell Profiling of RNA and Chromatin. *Cell* **183**, 1103–1116.e20 (2020).

27. Trevino, A. E. *et al.* Chromatin and gene-regulatory dynamics of the developing human cerebral cortex at single-cell resolution. *Cell* **184**, 5053–5069.e23 (2021).

28. Singh, R. & Berger, B. Deciphering the species-level structure of topologically associating domains. *bioRxiv* 2021.10.28.466333 (2021) doi:10.1101/2021.10.28.466333.

29. Crispatzu, G. *et al.* The chromatin, topological and regulatory properties of pluripotency-associated poised enhancers are conserved in vivo. *Nat. Commun.* **12**, 4344 (2021).

30. Zou, Z., Ohta, T., Miura, F. & Oki, S. ChIP-Atlas 2021 update: a data-mining suite for exploring epigenomic landscapes by fully integrating ChIP-seq, ATAC-seq and Bisulfite-seq data. *Nucleic Acids Res.* (2022) doi:10.1093/nar/gkac199.

31. Creyghton, M. P. *et al.* Histone H3K27ac separates active from poised enhancers and predicts developmental state. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 21931–21936 (2010).

32. Ostuni, R. *et al.* Latent enhancers activated by stimulation in differentiated cells. *Cell* **152**, 157–171 (2013).

33. Panigrahi, A. & O'Malley, B. W. Mechanisms of enhancer action: the known and the unknown. *Genome Biol.* **22**, 108 (2021).

34. Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for

interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 15545–15550 (2005).

35. Herrmann, J. C., Beagrie, R. A. & Hughes, J. R. Making connections: enhancers in cellular differentiation. *Trends Genet.* **38**, 395–408 (2022).

36. MacArthur, J. *et al.* The new NHGRI-EBI Catalog of published genome-wide association studies (GWAS Catalog). *Nucleic Acids Res.* **45**, D896–D901 (2017).

37. Maurano, M. T. *et al.* Systematic localization of common disease-associated variation in regulatory DNA. *Science* **337**, 1190–1195 (2012).

38. Sullivan, P. F., Kendler, K. S. & Neale, M. C. Schizophrenia as a complex trait: evidence from a meta-analysis of twin studies. *Arch. Gen. Psychiatry* **60**, 1187–1192 (2003).

39. Malan-Müller, S. *et al.* A systematic review of genetic variants associated with metabolic syndrome in patients with schizophrenia. *Schizophr. Res.* **170**, 1–17 (2016).

40. Lago, S. G. & Bahn, S. The druggable schizophrenia genome: from repurposing opportunities to unexplored drug targets. *NPJ Genom Med* **7**, 25 (2022).

41. Trubetskoy, V. *et al.* Mapping genomic loci implicates genes and synaptic biology in schizophrenia. *Nature* **604**, 502–508 (2022).

42. Chen, C.-H. *et al.* Determinants of transcription factor regulatory range. *Nat. Commun.* **11**, 2472 (2020).

43. Wang, S. *et al.* Target analysis by integration of transcriptome and ChIP-seq data with BETA. *Nat. Protoc.* **8**, 2502–2515 (2013).

44. Xie, Z. *et al.* Gene Set Knowledge Discovery with Enrichr. *Curr Protoc* **1**, e90 (2021).

45. Karunakaran, K. B., Chaparala, S. & Ganapathiraju, M. K. Potentially repurposable drugs for schizophrenia identified from its interactome. *Sci. Rep.* **9**, 12682 (2019).

46. Marballi, K. K. & Gallitano, A. L. Immediate Early Genes Anchor a Biological Pathway of Proteins Required for Memory Formation, Long-Term Depression and Risk for Schizophrenia. *Front. Behav. Neurosci.* **12**, 23 (2018).

47. Meng, X. *et al.* Neurexophilin4 is a selectively expressed α-neurexin ligand that modulates specific cerebellar synapses and motor functions. *Elife* **8**, (2019).

48. Cardano, M. *et al.* Epsins Regulate Mouse Embryonic Stem Cell Exit from Pluripotency and Neural Commitment by Controlling Notch Activation. *Stem Cells Int.* **2019**, 4084351 (2019).

49. Mountjoy, E. *et al.* An open approach to systematically prioritize causal variants and genes at all published human GWAS trait-associated loci. *Nat. Genet.* **53**, 1527–1533 (2021).

50. Barkhuizen, W., Pain, O., Dudbridge, F. & Ronald, A. Genetic overlap between psychotic experiences in the community across age and with psychiatric disorders. *Transl. Psychiatry* **10**, 86 (2020).

51. Schröder, J. *et al.* MicroRNA-138 is a potential regulator of memory performance in humans. *Front. Hum. Neurosci.* **8**, 501 (2014).

52. Chahrour, M. H. *et al.* Whole-exome sequencing and homozygosity analysis implicate depolarization-regulated neuronal genes in autism. *PLoS Genet.* **8**, e1002635 (2012).

53. McLaughlin, R. L. *et al.* Genetic correlation between amyotrophic lateral sclerosis and schizophrenia. *Nat. Commun.* **8**, 14774 (2017).

54. Kim, J. *et al.* The co-regulation mechanism of transcription factors in the human gene regulatory network. *Nucleic Acids Res.* **40**, 8849–8861 (2012).

55. Shyn, S. I. *et al.* Novel loci for major depression identified by genome-wide association study of Sequenced Treatment Alternatives to Relieve Depression and meta-analysis of three studies. *Mol. Psychiatry* **16**, 202–215 (2011).

56. Fabbri, C. & Serretti, A. Role of 108 schizophrenia-associated loci in modulating psychopathological dimensions in schizophrenia and bipolar disorder. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* **174**, 757–764 (2017).

57. Devor, A. *et al.* Genetic evidence for role of integration of fast and slow neurotransmission in schizophrenia. *Mol. Psychiatry* **22**, 792–801 (2017).

58. Liu, Z. *et al.* Dysregulated Glial Differentiation in Schizophrenia May Be Relieved by

Suppression of SMAD4- and REST-Dependent Signaling. *Cell Rep.* **27**, 3832–3843.e6 (2019).

59. Borsari, B. *et al.* Enhancers with tissue-specific activity are enriched in intronic regions. *Genome Res.* **31**, 1325–1336 (2021).

60. Chen, R. *et al.* Cell type–specific mechanism of Setd1a heterozygosity in schizophrenia pathogenesis. *Science Advances* vol. 8 Preprint at https://doi.org/10.1126/sciadv.abm1077 (2022).

61. Moore, D. L., Apara, A. & Goldberg, J. L. Krüppel-like transcription factors in the nervous system: novel players in neurite outgrowth and axon regeneration. *Mol. Cell. Neurosci.* **47**, 233–243 (2011).

62. Peyrot, W. J. & Price, A. L. Identifying loci with different allele frequencies among cases of eight psychiatric disorders using CC-GWAS. *Nat. Genet.* **53**, 445–454 (2021).

63. Pinacho, R. *et al.* Analysis of Sp transcription factors in the postmortem brain of chronic schizophrenia: a pilot study of relationship to negative symptoms. *J. Psychiatr. Res.* **47**, 926–934 (2013).

64. Seifert, G., Schilling, K. & Steinhäuser, C. Astrocyte dysfunction in neurological disorders: a molecular perspective. *Nat. Rev. Neurosci.* **7**, 194–206 (2006).

65. Tanaka, A. J. *et al.* De novo mutations in PURA are associated with hypotonia and developmental delay. *Cold Spring Harb Mol Case Stud* **1**, a000356 (2015).

66. Daniel, D. C. & Johnson, E. M. PURA, the gene encoding Pur-alpha, member of an ancient nucleic acid-binding protein family with mammalian neurological functions. *Gene* **643**, 133–143 (2018).

67. Barešić, A., Nash, A. J., Dahoun, T., Howes, O. & Lenhard, B. Understanding the genetics of neuropsychiatric disorders: the potential role of genomic regulatory blocks. *Mol. Psychiatry* **25**, 6–18 (2020).

68. Bøttger, P. *et al.* Glutamate-system defects behind psychiatric manifestations in a familial

hemiplegic migraine type 2 disease-mutation mouse model. *Sci. Rep.* **6**, 22047 (2016).

69. Pieraccioli, M. *et al.* ZNF281 inhibits neuronal differentiation and is a prognostic marker for neuroblastoma. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 7356–7361 (2018).

70. Okamoto, S.-I., Sherman, K., Bai, G. & Lipton, S. A. Effect of the ubiquitous transcription factors, SP1 and MAZ, on NMDA receptor subunit type 1 (NR1) expression during neuronal differentiation. *Brain Res. Mol. Brain Res.* **107**, 89–96 (2002).

71. Pinacho, R. *et al.* Increased SP4 and SP1 transcription factor expression in the postmortem hippocampus of chronic schizophrenia. *J. Psychiatr. Res.* **58**, 189–196 (2014).

72. Wen, Y.-D. *et al.* Genetic Profiles Playing Opposite Roles of Pathogenesis in Schizophrenia and Glioma. *J. Oncol.* **2020**, 3656841 (2020).

73. Mäki-Marttunen, V., Andreassen, O. A. & Espeseth, T. The role of norepinephrine in the pathophysiology of schizophrenia. *Neurosci. Biobehav. Rev.* **118**, 298–314 (2020).

74. Maletic, V., Eramo, A., Gwin, K., Offord, S. J. & Duffy, R. A. The Role of Norepinephrine and Its α-Adrenergic Receptors in the Pathophysiology and Treatment of Major Depressive Disorder and Schizophrenia: A Systematic Review. *Front. Psychiatry* **8**, 42 (2017).

75. La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494–498 (2018).

76. Gorin, G., Svensson, V. & Pachter, L. Protein velocity and acceleration from single-cell multiomics experiments. *Genome Biol.* **21**, 39 (2020).

77. Matsumoto, H. *et al.* SCODE: an efficient regulatory network inference algorithm from single-cell RNA-Seq during differentiation. *Bioinformatics* **33**, 2314–2321 (2017).

78. Deshpande, A., Chu, L.-F., Stewart, R. & Gitter, A. Network inference with Granger causality ensembles on single-cell transcriptomics. *Cell Rep.* **38**, 110333 (2022).

79. Andrew, G., Arora, R., Bilmes, J. & Livescu, K. Deep Canonical Correlation Analysis. in *Proceedings of the 30th International Conference on Machine Learning* (eds. Dasgupta, S. & McAllester, D.) vol. 28 1247–1255 (PMLR, 17--19 Jun 2013).

80. Hie, B. *et al.* Computational Methods for Single-Cell RNA Sequencing. *Annual Review of*

Biomedical Data Science vol. 3 339–364 Preprint at https://doi.org/10.1146/annurev-biodatasci-012220-100601 (2020).

81. Cusanovich, D. A. *et al.* Multiplex single cell profiling of chromatin accessibility by combinatorial cellular indexing. *Science* **348**, 910–914 (2015).

82. Cusanovich, D. A. *et al.* A Single-Cell Atlas of In Vivo Mammalian Chromatin Accessibility. *Cell* **174**, 1309–1324.e18 (2018).

83. Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression data analysis. *Genome Biol.* **19**, 15 (2018).

84. Haghverdi, L., Büttner, M., Wolf, F. A., Buettner, F. & Theis, F. J. Diffusion pseudotime robustly reconstructs lineage branching. *Nat. Methods* **13**, 845–848 (2016).

85. Gjoneska, E. *et al.* Conserved epigenomic signals in mice and humans reveal immune basis of Alzheimer's disease. *Nature* **518**, 365–369 (2015).

86. Xiao, S. *et al.* Comparative epigenomic annotation of regulatory DNA. *Cell* **149**, 1381–1392 (2012).

87. Kerimov, N. *et al.* A compendium of uniformly processed human gene expression and splicing quantitative trait loci. *Nat. Genet.* **53**, 1290–1299 (2021).

88. Won, H. *et al.* Chromosome conformation elucidates regulatory relationships in developing human brain. *Nature* **538**, 523–527 (2016).

89. Dixon, J. R. *et al.* Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature* vol. 485 376–380 Preprint at https://doi.org/10.1038/nature11082 (2012).

90. ENCODE Project Consortium. An integrated encyclopedia of DNA elements in the human genome. *Nature* **489**, 57–74 (2012).

91. Poterlowicz, K. *et al.* 5C analysis of the Epidermal Differentiation Complex locus reveals distinct chromatin interaction networks between gene-rich and gene-poor TADs in skin epithelial cells. *PLoS Genet.* **13**, e1006966 (2017).

92. Uhlén, M. *et al.* Proteomics. Tissue-based map of the human proteome. *Science* **347**, 1260419 (2015).

93. Thul, P. J. & Lindskog, C. The human protein atlas: A spatial map of the human proteome. *Protein Sci.* **27**, 233–244 (2018).

94. Schneider, V. A. *et al.* Evaluation of GRCh38 and de novo haploid genome assemblies demonstrates the enduring quality of the reference assembly. *Genome Res.* **27**, 849–864 (2017).

95. Grant, C. E., Bailey, T. L. & Noble, W. S. FIMO: scanning for occurrences of a given motif. *Bioinformatics* **27**, 1017–1018 (2011).

96. Kulakovskiy, I. V. *et al.* HOCOMOCO: towards a complete collection of transcription factor binding models for human and mouse via large-scale ChIP-Seq analysis. *Nucleic Acids Res.* **46**, D252–D259 (2018).

97. Castro-Mondragon, J. A. *et al.* JASPAR 2022: the 9th release of the open-access database of transcription factor binding profiles. *Nucleic Acids Research* vol. 50 D165–D173 Preprint at https://doi.org/10.1093/nar/gkab1113 (2022).

98. Weirauch, M. T. *et al.* Determination and inference of eukaryotic transcription factor sequence specificity. *Cell* **158**, 1431–1443 (2014).

Figures

634
635

Figure 1. GrID-Net Framework. (a) The causal asynchrony between chromatin accessibility

and gene expression results in "cell-state parallax", where a temporal lag exists between these

two modalities. Leveraging this parallax from single-cell snapshots enables inference of causal

locus-gene associations. **(b)** To identify links between noncoding loci (i.e. peaks) and the genes

they regulate, we apply GrID-Net to single-cell multimodal data in which chromatin accessibility

(ATAC-seq) and gene expression (RNA-seq) have been co-assayed in the same cell. GrID-Net

is an extension of classical Granger causality that can operate on DAG-based representations

of single-cell trajectories. The core of GrID-Net is a graph neural network-based lagged

message passing architecture. **(c)** GrID-Net enables the detection of asynchronous interactions

between regulatory activity at accessible noncoding loci and their downstream effects on gene

expression. **(d)** GrID-Net peak-gene links enable the functional interpretation of genetic variants

associated with disease. We present a framework for studying the functional effects of such

variants both at the level of specific variants and with respect to disease-affected modules.

Figure 2. GrID-Net links noncoding loci to genes from single-cell multimodal dynamics.

(a) Accuracy of single-cell (GrID-Net, Pearson correlation, and pseudocell correlation) and bulk (Activity-By-Contact) methods in predicting expression quantitative trait loci (eQTLs) in matching tissue. In this evaluation, we use eQTLs as proxies for causal interactions between genomic loci and genes. **(b)** Accuracy of methods in predicting high frequency three-dimensional chromatin contacts. The Activity-By-Contact (ABC) model is excluded here since it uses chromatin interaction data as one of its inputs. **(c)** Compared to existing approaches, GrID-Net is better able to identify asynchronous peak-gene associations. On the human corticogenesis dataset, we collected peak-gene pairs that GrID-Net prioritized differently than the correlation-based approaches (top 1%: preferred by GrID-Net; bottom 1%: preferred by the other approach). For each pair, we quantified the asynchrony as the separation between pseudotime values corresponding to the cells where the peak and gene achieved their maximum values,

665   respectively. **(d)** GrID-Net effectively captures instances of chromatin priming. On the human

666   corticogenesis dataset, we show the ratio of poised enhancers to active enhancers for peaks in

667   peak-gene pairs prioritized by GrID-Net vs. each of the correlation-based methods. The

668   estimation of poised-vs.-active enhancers was done using publicly-available histone

669   modification data.

670
671

Figure 3. GrID-Net predictions link schizophrenia variants to affected genes. (a,b)

Schizophrenia (SCZ) variants from Trubetskoy, Pardiñas *et al*., prioritized by GrID-Net. Overall,

195 SCZ variants overlapped with scATAC-seq peaks from the human corticogenesis dataset.

Compared to the original study, GrID-Net enables the annotation of many additional variants in

this subset. Furthermore, as multimodal single-cell assays become common, we expect GrID-

Net's coverage to widen. We also compared specific SCZ variant-gene links for the 12 variants

where the original study also reported an annotation. **(c)** GrID-Net is able to identify long-range

variant gene associations: (Left) genomic distance between an SCZ variant and the

transcription start site (TSS) of its affected gene(s). (Right) number of genes in the genomic

region between an SCZ variant and its affected gene(s). **(d)** Histogram of the number of variants

linked to each SCZ gene. **(e)** Genes linked to SCZ variants are cell type-specific. We show the

heatmap of normalized expression values for all genes linked to SCZ variants in cell types

involved in human corticogenesis. **(f)** Distribution of annotated genomic locations for SCZ

variants linked to genes. A substantial fraction of genes identified by GrID-Net are associated

with intronic variants which have been reported to have tissue-specific regulatory effects.

Figure 4. GrID-Net links transcription factor binding disruptions to gene dysregulation in schizophrenia. (a) TFs with at least a 20-fold change in predicted binding affinity caused by SCZ variants and the genes linked to these variants. Many of these TFs (e.g., KLF and SP families) are implicated in SCZ. **(b)** Relationship between TF binding affinity changes caused by SCZ variants and significance of differential expression in their linked genes between patient-derived SCZ glial progenitor cells and matching controls. **(c)** Examples of SCZ variant-gene links supported by at least 20-fold changes in predicted TF binding affinity. **(d)** Gene modules co-expressed with SCZ genes in SCZ GPCs and matching controls.

Supplementary Figures

703 **Figure S1.** GrID-Net outperforms other methods in eQTL prediction accuracy across different
704 eQTL significance thresholds.

707
708
709 **Figure S2.** GrID-Net is robust to data sparsity in single-cell datasets.
710

712
Figure S3. Genomic distance-dependent model improves peak-gene link prediction accuracy
714 (a) within and (b) across species and cell types.
715
716

717
718

719
720 **Figure S4. (a)** Relationship between FINEMAP posterior probabilities (PP) for a SCZ variant
721 and its corresponding TF binding affinity change (Welch's one-tailed *t*-test). TFs with the most
722 widespread **(b)** increase or **(c)** decrease in binding affinity between the reference and the SCZ
723 risk allele.

724
725