Genome biology and evolution of mating type loci in four cereal rust fungi

Zhenyan Luo[1], Alistair McTaggart[2], and Benjamin Schwessinger[1*]

[1] Research Biology School, Australian National University, Canberra, ACT, Australia

[2] Centre for Horticultural Science, Queensland Alliance for Agriculture and Food Innovation, The University of Queensland, Ecosciences Precinct, Dutton Park, Queensland, Australia

*Corresponding author

E-mail: benjamin.schwessinger@anu.edu.au (BS)

Abstract

Obligate heterozygous loci such as sex- or mating-compatibility regions often display suppression of recombination and signals of genomic degeneration. In Basidiomycota, two distinct gene loci confer mating compatibility. These encode for homeodomain (*HD*) transcription factors and pheromone receptor (*Pra*)-ligand pairs. To date genome level mating type (*MAT*) loci analysis is lacking for obligate biotrophic basidiomycetes in the order *Pucciniales*, which contains many economically important plant pathogens.

Here, we focus on four *Puccinia* cereal rust species, including *P. coronata* f. sp. *avenae*, *Puccinia graminis* f. sp. *tritici*, *P. triticina* and *P. striiformis* f. sp. *tritici*, which infect oat and wheat. *MAT* loci are located on two separate chromosomes supporting previous hypotheses of tetrapolar mating types in the *Pucciniales*. The *HD* locus is multiallelic in all four species while the *PR* locus appears to be biallelic except for *P. graminis* f. sp. *tritici* which displays genetic features of more than two alleles. *HD* loci were largely conserved in their macrosynteny within and between species without strong signals of recombination suppression. *PR* loci proximate regions, however, displayed extensive signs of recombination suppression and genomic degeneration in the three species with a clear biallelic *PR* locus. These observations suggest a correlation between recombination suppression, genomic degeneration and allele status of *MAT* loci which is consistent with recent mathematical modelling and simulations. Finally, we confirm the evolutionary conservation of *MAT* gene expression during the asexual infection cycle of the cereal host which we propose is related to correct nuclear pairing during spore formation. Together, our study provides insights into the evolution of *MAT* loci of key pathogenic *Puccinia* species. This detailed understanding is important to predict possible combinations of nuclear pairs that can arise via sexual

47 reproduction or somatic recombination to enable the evolution of newly virulent isolates of

48 these important plant pathogens.

Author summary

Sex of animals and some plants is determined by sex chromosomes. In fungi, mate

compatibility is determined by mating type (*MAT*) loci, which share some features with sex

chromosomes including recombination suppression around heterozygous loci. Here, we study

the *MAT* loci in fungal pathogens from the order *Pucciniales* that can cause rust diseases on

many economically important plants including wheat and oat. We show that one of the *MAT*

loci is multiallelic, while the other is biallelic. The biallelic locus shows strong signs of

recombination suppression and genetic deterioration with an increase in transposable elements

and gene deserts surrounding the locus. Our findings on the genome biology of *MAT* loci in

four economically important pathogens will lead to a better understanding and prediction of

evolution of novel virulent isolates that can lead to large scale pandemics in agriculture.

Introduction

64 The evolution of 'sex and mating-type' determining loci and chromosomes is a fundamental

65 question in biology. It is widely accepted that X and Y chromosomes were originally

66 homologous, but recombination suppression gradually resulted in genetic degeneration of the

67 Y chromosome (1). Genetic degeneration footprints are alleviated rates of (non-)synonymous

68 substitutions (d_N, d_S), accumulation of transposable elements (TEs), accumulation of

inversions, reduced gene expression and/or reduced gene numbers which are all a consequence of recombination cessation (2-5). In contrast to many animals and plants, antagonistic selection is unlikely to be the evolutionary mechanism that causes recombination cessation in fungal mating-type (*MAT*) loci (6-8). Instead, mathematical modelling and stochastic simulations suggest that recombination suppressors, e.g. inversions or TEs, which are loaded with less deleterious mutations than average initially, are more likely to be fixed at permanently heterozygous *MAT* loci when compared to other autosomal loci. This initial fixation of recombination suppressors at *MAT* loci can lead to a progressive extension of non-recombining regions via the accumulation of additional inversion or TEs over longer evolutionary timeframes (2). Importantly, this model predicts non-recombining regions around *MAT* loci are larger at biallelic loci and in species with smaller effective population size, short haploid phase, outcrossing mating systems, high mutation rates, and extended dikaryotic life stages. Indeed, several fungal species carry extensive non-recombining regions around *MAT* loci. Recombination suppression can range from hundreds of kbp (kilo base pairs) to several mbp (mega base pairs) and lead to genetic degeneration (5, 9), such as in *Neurospora tetrasperma* (10), *Podospora anserina* (11), *Schizothecium tetrasporum* (12) and several *Microbotryum* spp. (8, 13).

Detailed analyses of *MAT* loci and their genome biology has been lacking behind for rust fungi of the order *Pucciniales* in the division Basidiomycota (9). This is despite the fact that *Pucciniales* is the fungal order with the most known plant pathogens with close to 8,000 described species that cause diseases with significant environmental and economic impact for various trees and crops such as poplar, paperbarks, wheat, oat, soybean, and coffee (14-16). Overall, Basidiomycota have evolved unique mating systems to govern nuclear compatibility, mate selection, and completion of the full life cycle (9). From a genetic perspective, two *MAT* loci determine mating-type identity and non-self-recognition in most Basidiomycota. The

pheromone receptor (*PR*) locus contains a pheromone receptor (*Pra*) gene and at least one

mating pheromone peptide precursor gene (*mfa*) (17). The *PR* locus defines pre-mating

compatibility and gamete fusion. The PRA protein is a transmembrane localized G protein-

coupled receptor, which recognizes the processed and posttranslationally modified mature

pheromone peptide MFA encoded by a matching allelic *PR* locus (9). Downstream of initial

gamete fusion, the homeodomain (*HD*) locus determines success in postmating development

(9). The *HD* locus contains two tightly linked homeodomain transcription factors genes, (*bW-*

HD1 and *bE-HD2* in *Pucciniales* (18, 19)*,* also known as *bE-HD1* and *bW-HD2* in other fungi

(9)) which share a promoter and are outwardly transcribed in opposite directions. Their

protein products need to be of different allelic specificity to form heterocomplexes and to

initiate a transcriptional cascade downstream of PRA-MFA activation (9). The *HD*

transcription factors regulate cellular development during mating, maintenance of the

dikaryotic state, and control pathogenicity in some plant pathogens such as smuts (13). The

HD and *PR* loci can be either physically linked or unlinked which means they segregate

together or independently, respectively. The mating system in Basidiomycota influences the

genomic organization and resulting segregation patterns of *MAT* loci. Most basidiomycete

species are heterothallic, where sexual gametes must carry different alleles at both *MAT* loci

and intra-haploid mating is not possible. Hence, heterothallic species can either have bipolar

or tetrapolar mating systems. In bipolar species the *HD* and *PR* loci are physically linked or

one locus has lost its function in mating. In any case, in bipolar species only one *MAT* locus

determines mate compatibility. In tetrapolar species, the *HD* and *PR* loci are unlinked and

both determine mate compatibility. Bipolar mating systems are favorable in inbreeding

populations that undergo selfing. Under selfing a single locus is advantageous as the

likelihood of compatibility amongst gametes derived from the same diploid individual is 50%

(13). In comparison, a tetrapolar mating system has at most a 25% compatibility under the

same selfing conditions. In contrast, outcrossing mating systems benefit from multiallelism at

MAT loci because it increases compatibility at syngamy of haploid gametes from different

diploid individuals (6, 9, 13). In many tetrapolar basidiomycete species, the *HD* locus is

highly polymorphic in the population with ten to hundred known or estimated alleles that are

under negative, frequency dependent selection (13, 20, 21). The tetrapolar mating system is

thought to be the ancestral state in Basidiomycota, yet several species have evolved bipolarity

independently. For example, *Microbotryum* spp. are highly selfing, and multiple independent

evolutionary events have linked the *HD* locus and the *PR* locus into evolutionary strata of

different sizes and different ages in this genus (3, 7, 8, 22, 23). These evolutionary strata show

clear signatures of recombination suppression and genetic degeneration. Similarly, in the

Ustilaginomycotina, several species evolved bipolarity linking both *MAT* loci such as

Ustilago hordei, U. bromivora or *Sporisorium scitamineum* (9). In contrast, a recent study

showed that tetrapolarity has evolved multiple times in the human skin fungus genus

Malassezia, where the ancestral state is pseudobipolar (24).

Studying mating systems in fungal species of the order *Pucciniales* has been difficult because

of their obligate biotrophy and often complex life cycles that prevent culturing outside of their

hosts and often requires multiple hosts to complete their sexual life cycle (9). For example,

macrocyclic and heteroecious rust fungi have five spore stages and require two hosts to

complete their asexual and sexual life cycles. During the asexual infection cycle on the

"primary" host, e.g. wheat and oat, rust fungi are dikaryotic and produce re-infective

urediniospores that harbor two distinct nuclei carrying compatible mating types. The sexual

cycle is initiated with the production of teliospores, which represent the short diploid phase of

rust fungi. Resulting haploid basidiospores infect the "alternate" host, e. g. *Berberis sp.*, and

form haploid infection structures with distinct mating types. Fusion of cell with compatible

mating types generates dikaryotic intercellular mycelia within the plant tissue that gives rise

to dikaryotic aeciospores which are only infective on the "primary" host (25). Infection of the

primary host completes the full life cycle (9, 26). The role of *MAT* genes during the life cycle

of rust fungi is rudimentary. Though it is hypothesized that *MAT* genes regulate appropriate

nuclear paring during dikaryotic spore production and mediate the compatibility of haploid

cell fusions on the "alternate" host (9, 26). Original research suggested that most rust fungal

species are bipolar (26), yet more recent studies on the rust fungi *Melampsora lini* and *P.*

coronata var. *coronata* revealed tetrapolar mating systems (27, 28). In the absence of direct

experimental evidence, genomic insights into *MAT* loci from genome assemblies of the

dikaryotic state can provide hypotheses on the breeding systems. Independent initial genome

analyses of several rust fungi revealed two unlinked *MAT* loci with *PR* and *HD* loci being

located on different contigs, scaffolds or chromosomes including in *Austropuccinia psidii* (29)*,*

P. graminis f. sp. *tritici* (30) and *P. tritici* (18, 19)*.* This genome organization supports the

hypothesis of a tetrapolar mating system in these rust fungi. These rust fungi encode *Pra*

receptor genes belonging to the STE3 family. The likely *bona fide* receptors are encoded by

STE3.2-2 and *STE3.2-3* that display clear signatures of ancient trans-specific polymorphisms

which are likely ancestral to Basidiomycota (31). Here the sequences of a given mating type

is more similar to pheromone receptor alleles of the same mating type in distantly related

fungi than the alternate allele in the same species. In most cases, at least one *mfa* allele has

been found in proximity to the *STE3.2-2/3* locus. The additional STE3 family member

STE3.2-1 is nearly homozygous in studied species such as *P. tritici* and is hypothesized not to

be involved in mating (18). To date, the *PR* locus has been found to be biallelic in available

population level datasets for *P. tritici* and *P. striiformis* f. sp. *tritici* (18, 21, 32), yet

Melampsora larici-populina might be carrying additional alleles and population level

analyses are missing for other rust fungi. In contrast, the *HD* locus is multiallelic in *P.*

triticina and *P. striiformis* f. sp. *tritici* (18, 21, 32) encoding variants of *bW-HD1* and *bE-HD2*

that are highly dissimilar only in the variable N-terminal domain, shown to be essential for functional heterodimerization in other Basidiomycota (33, 34). It is noteworthy that direct biochemical or genetic evidence is lacking for any of these *MAT* genes' involvement in mating in rust fungi. Yet, complementation assays in *U. maydis* and host induced gene silencing on wheat demonstrated the importance of *Pra* and *HD* genes of *P. triticina* for mating in a heterologous system and for spore production during asexual infection (18). In addition, *MAT* genes were expressed during the asexual and/or sexual infection cycle in several rust fungal species (18, 35). These initial studies open up several outstanding knowledge gaps for *MAT* loci in rust fungi: 1) What is the chromosomal organization of *PR* and *HD* loci and how does it compare between closely related species? 2) What is the allelic diversity of *MAT* genes within rust fungal species? 3) What is the composition and organization of *MAT* loci proximal regions, how does it compare between *PR* and *HD* loci, and how does it vary between closely related species? 4) What is the compositional and organizational variation of *MAT* loci within rust fungal species? 5) Is the expression of *MAT* genes during the asexual infection cycle common in rust fungi?

Addressing these knowledge gaps has long been a challenge because initial genome assemblies of rust fungi were fragmented and the structure of the *MAT* loci could not be assessed consistently. For example, clustering of repetitive elements around *MAT* loci (17, 36, 37) is a challenge for studying *MAT* loci in rust fungi with repeat-rich genomes because repetitive sequences cause gaps, fragmentation, and phase-errors in the assembled genome. Here, we overcome this challenge and address important knowledge gaps around *MAT* loci in four cereal rust fungi by providing a detailed comparative analysis of publicly available chromosome scale phased genome assemblies and complementary Illumina short-read datasets. The study species include the oat crown rust fungus *P. coronata* f. sp. *avenae* (38-40), and the three wheat rust fungi, *P. graminis* f. sp. *tritici* (30, 41, 42), *P. triticina* (43-46)

and *P. striiformis* f. sp. *tritici* (47-53), which combined are the biggest threat to wheat production globally causing losses of several billion dollars every year (16). Our detailed comparative analyses set out to address the following specific objectives. 1) To determine the chromosomal organization of *MAT* loci. 2) To assess the allelic diversity of these loci in the four cereal rust fungi. 3) To evaluate synteny, recombination suppression, and genomic deterioration around *PR* and *HD* loci with the prediction that multiallelic loci are more syntenic than biallelic loci and that the latter show stronger footprints of recombination suppression and genomic degeneration. 4) To gage the conservation and/or plasticity of *MAT* loci within a cereal rust fungus species using *P. triticina* as model system. And 5) To determine the expression of *Pra* and *HD* genes during the asexual infection cycle of the cereal hosts.

We show that *HD* genes are multiallelic in all four rust fungal species and *Pra* genes are unequivocal biallelic in three out of four. Only biallelic *MAT* loci show strong signs of recombination suppression and genomic degeneration which confirms with recent mathematical modelling. Our results provide novel insights into the genome biology of *MAT* loci in cereal rust fungi. These are important to consider when making predictions about mating and hybridization events that are possibly linked to the evolution of novel pathogenicity traits.

Results

Genomic organisation and inheritance of mating type genes suggest cereal rust fungi have tetrapolar mating systems

We set out to test previous preliminary observations that suggest cereal rust fungi have a tetrapolar mating system with *PR* and *HD* loci being unlinked (18, 19). We made use of the

seven available chromosome-level genome assemblies of four cereal rust fungi *P. coronata* f.

sp. *avenae* (*Pca*) (38, 40, 54), *P. graminis* f. sp. *tritici* (*Pgt*) (30, 41, 42, 53), *P. triticina* (*Pt*)

(32, 43-46, 55) and *P. striiformis* f. sp. *tritici* (*Pst*) (47, 49-52, 56), in addition to five partially

phased assemblies (Table 1). Our initial analysis focused on one reference genome per species

with *Pca 203*, *Pgt 21-0*, *Pt 76*, and *Pst 134E*.

We used previously characterized *MAT* genes from the *Pt* isolate *BBBD* (Table 2) (18) as

queries to identify orthologs in available genomes and proteomes (Table 1). We identified *HD*

(*bW-HD1* and *bE-HD2*) and *Pra* (*STE3.2-2* and *STE3.2-3*) alleles on chromosome 4 and

chromosome 9 respectively, whereas *STE3.2-1* alleles were located on chromosome 1 (S1

Fig). This is consistent with previous preliminary analysis for *Pgt* and *Pt,* which suggested

HD and *PR* loci are located on two distinct chromosomes (30, 32). We confirmed these

results using five genome resources for cereal rust fungi that are only partially phased and

non-chromosome scale (Table 1). We identified two alleles of *bW-HD1, bE-HD2* and *PRA*

(*STE3.2-2* or *STE3.2-3)* in all five genome assemblies with the exception for *STE3.2-2* in *Pst*

104E and *Pca 12SD80* (38, 48). In the case of *Pst 104E, STE3.2-2* was absent from the

genome assembly but we confirmed its presence in the raw sequencing data by mapping

sequencing reads against the *Pst 134E* phased chromosome scale genome assembly (47).

STE3.2-2 displayed average genome-wide read coverage without any nucleotide variation

confirming that *Pst 104E* was also heterozygous for *Pra*. In the case of *Pca 12SD80,* we

identified two copies of *STE3.2-2* on contig 000183F and 000183F_004. This miss-

characterization of *STE3.2-2* in *Pst 104E* and *Pca 12SD80* is likely caused by early long-read

genome assembly errors using low accuracy PacBio long reads and suboptimal genome

assembly algorithms that struggle with highly repetitive regions. Similar observations have

been made for missing *Pra* genes reported for earlier genome versions of *Melampsora larici-*

populina (57).

We also identified an additional *Pra*-like gene, *STE3.2-1,* on chromosome 1. However, its two

alleles are nearly identical in all dikaryotic genome assemblies and are not associated with

any pheromone peptide precursor genes. Hence, *STE3.2-1* is likely not involved mating

compatibility in rust fungi.

Taken together, the two *MAT* loci, *HD* and *PR*, are unlinked and heterozygous in all twelve

dikaryotic genome assemblies, which is consistent with a genome informed hypothesis of a

tetrapolar mating system for cereal rust fungi.

Mfa pheromone peptide precursor genes are closely linked to *STE3.2-2* but not *STE3.2-3*

We searched the available dikaryotic genomes for putative pheromone peptide precursor

genes, which encode mating factor a (MFA) peptides. *Mfas* are often linked to *Pra* alleles and

predicted to bind to the compatible pheromone receptors encoded at the allelic *PR* locus. We

used the previously identified *mfa*1, *mfa2,* and *mfa3* genes of *Pst*, *Pgt,* and *Pt* as queries (18).

We identified a single *mfa2* pheromone peptide precursor gene in all species. In all cases,

mfa2 and *STE3.2-2* were closely linked, located within 500–1100 bp from each other, and

encoded on the same DNA strand (S2 Fig). The *mfa2* derived peptides were all 34 amino

acids long with a characteristic CAAX motif at the C-terminus, where C is cysteine, A is an

aliphatic amino acid, and X is any amino acid (58) (S3B Fig). The *MFA2* amino acid

sequences were 100% identical at the species level (S3B Fig).

In contrast, the sequence, length, number, and location of *mfa* genes associated with *STE3.2-3*

varied between species (S2, S3A and S3C Figs). *Pca* and *Pst* encoded a single *mfa1* allele.

Pcamfa1 of *Pca 203* encoded a 66 amino acid peptide and was located 0.54 mbp upstream of

STE3.2-3. In *Pst*, *Pstmfa1* encoded a 64 amino acid peptide and was located 10 kbp and 13

kbp away from *STE3.2-3* in *Pst 134E* and *Pst DK0911*, respectively. In contrast, *Pt* carried

two identical copies of *Ptmfa1/3* encoding for 61 amino acid peptides. In the *Pt 76* reference,

Ptmfa1/3 were associated with *STE3.2-3* at a distance of 0.24 mbp and 0.27 mbp, respectively

(S2 Fig). In addition, *Pgt* encoded for two *mfa* genes in close proximity to *STE3.2-3* that were

located upstream and downstream at a distance of 30 kbp and 96 kbp, respectively (S2 Fig).

Yet in contrast to *Pt*, *Pgt mfa1* and *mfa3* encoded for highly distinct pheromone peptide

precursor that were only 26.47% identical at the amino acid level and varied in length (59 vs

57 amino acids, respectively). Based on their amino acid differences, MFA1 and MFA3 were

predicted to be processed into distinct mature MFA peptides that might have different

receptor specificities.

In all four cereal rust fungi, *Pra* alleles were associated with species specific *mfa* genes. Yet,

the number, distance, organization, and sequence of *mfa* genes varied between species.

277 Genealogies elucidate distinct evolutionary histories for the
278 different *MAT* genes

Having identified the *MAT* genes, we investigated the genealogical relationships of each

individual gene and compared them between each other and with the species tree. This

allowed us to test for shared or distinct evolutionary histories, which can provide indications

for recombination within and between loci. For example, so called trans-specific

polymorphism indicate that recombination cessation is older than speciation (31, 59). In the

absence of a robust species tree (60), we first generated a multilocus species tree based on a

multiple sequence alignment of 2284 single orthologs proteins (61) including *P. polysora* f. sp.

zeae as an outgroup (Fig 1A). The species tree identified the three wheat rust fungi being most

closely related to each other when compared to the oat rust fungus *Pca* (Fig 1A). In the wheat

rust clade, *Pgt* was closer related to *Pt, w*hile *Pst* was more distantly related to the two. We

aligned *HD* genes *bW-HD1* and *bE-HD2* separately to construct independent gene trees using

P. polysora f.sp. *zeae* alleles as outgroups (Table 2, Figs 1B and 1C). The gene trees of *bW-*

HD1 and *bE-HD2* group alleles from each species into a species-specific clade (Figs 1B and

1C) and we therefore did not find any evidence for trans-specific polymorphisms for either

HD gene. We identified multiple *bW-HD1* and *bE-HD2* alleles for the four cereal rust fungi

suggesting the *HD* locus is multiallelic. We observed several shared *HD* alleles between

isolates in the case of *Pt* while *HD* alleles in *Pca* and *Pgt* appeared to be more diverse. In the

case of *Pgt*, we were limited to only four alleles, making any meaningful conclusion difficult.

We next tested our null hypothesis that *bW-HD1* and *bE-HD2* have similar evolutionary

histories because they are closely linked and share a promoter. We performed approximately

unbiased (AU) tests (62) within each species to investigate if the tree topologies of *bW-HD1*

and *bE-HD2* are concurrent with each other. In *Pca, Pst,* and *Pt*, we found that the p_{AU}-value

of the AU test was smaller than 0.05 which led us to reject the null hypothesis of concurrent

tree topologies. This suggests distinct evolutionary histories for the *bW-HD1* and *bE-HD2*

genes in these species, which might be caused by recombination within the *HD* locus. In

contrast, the AU-test for *Pgt* suggested similar tree topologies for *bW-HD1* and *bE-HD2*,

which likely is influenced by the low sample number of four alleles.

Next, we explored the evolutionary relationship between *Pra* alleles by building a single gene

tree using *P. polysora* f.sp. *zeae STE3.2-2* and *STE3.2-3* as outgroups. The *Pra* gene tree

formed two obvious clades based on allele identity grouping the *STE3.2-2* alleles of all

species into one clade and all *STE3.2-3* alleles into another clade including respective alleles

of *P. polysora* f.sp. *zeae* (Fig 2). In each clade, there was little to no intra-species variation for

STE3.2-2 and *STE3.2-3* based on the minimal branch lengths that separated allele copies in

each species sub-clade. This indicates that *Pra* might be biallelic in the cereal rust fungi (see

also below). The clear grouping by allele identity indicates strong trans-polymorphisms and a

long-term suppression of recombination at the *PR* locus that predates ancestral species split.

We next investigated if the putative pheromone peptide precursor *mfa* genes followed similar

evolutionary histories compared to their collocated *Pra* genes. Like the *Pra* genealogy, the

gene tree of *mfa* alleles revealed strong trans-specific polymorphisms because *mfa* copies

clustered by allelic and not species identity (S4 Fig). For example, all *Mfa2* alleles grouped

into one large clade according to their physical association with the specific *Pra* allele

STE3.2-2. Similarly, all *Mfa1* alleles grouped together with similar topology as their linked

STE3.3-2 copies. The species specific *Pgtmfa3s*, which are also physically linked to *STE3.2-3,*

were placed as a sister group to the *mfa1* clade, being more closely related to *mfa1* than to

mfa2.

We also investigated the genealogy of the *Pra*-like *STE3.2-1*. We used *P. polysora* f. sp. *zeae*

STE3.2-1 as outgroup for rooting the gene tree (63). The two *STE3.2-1* alleles identified in

each isolate were almost completely identical and grouped by species identity. Overall,

STE3.2-1 showed very little variation within each species (S5 Fig). This further supports our

conclusion that *STE3.2-1* is likely not involved in defining mate compatibility.

HD loci display limited signs of genomic deterioration and their synteny is conserved within species

Different effects on recombination have been reported around *MAT* loci in basidiomycetes (8,

64). Hence, we investigated if we could identify signals of altered recombination, synteny,

and genomic deterioration around the *MAT* loci in rust fungi. First, we analyzed whether there

was a reduction in synteny on chromosome 4 surrounding the *HD* locus as a signature for

long-term recombination suppression. We used mummer (65) to align the haplotypes of

chromosome 4 from *Pca, Pgt*, *Pt,* and *Pst,* against each other (S6 Fig). Overall macro-synteny

of chromosome 4 haplotypes was conserved in all cereal rust fungi with *Pt 76* being the most

syntenic (S6C Fig). Macro-synteny in a 40 kb-sized window around the *HD* locus was mostly

conserved in *Pt 76*, *Pca 203*, and *Pst 134E* while this initial analysis suggested that this locus

was less syntenic in *Pgt 21-0* (S6 Fig).

We investigated the individual allelic divergence along chromosome 4 using synonymous

divergence (d_S) value as an additional measure of footprint of ancient recombination

suppression events (8). This analysis tested if the *HD* and their proximal genes showed an

increase in d_S values, which could be caused by long-term suppression of recombination and

accumulation of independent mutations, as seen in other Basidiomycota (13). We calculated

the pairwise d_S values for all genes across all sister chromosomes in all four species of

Puccinia. This whole genome analysis of all allele pairs on sister chromosomes suggested that

chromosome 4 did not have increased d_S values when compared to all other chromosomes (S7

Fig). The d_S value of alleles for both *HD* genes fell within the upper 95% quantile of d_S values

on chromosome 4. Yet, d_S values of directly adjacent genes were not elevated beyond the

background level when we plotted d_S values for allele pairs along chromosome 4 (Fig 3).

Consistently, regions around *HD* loci did not show any obviously aberrant patterns of gene or

transposon density, which are common features of genetic deterioration (Fig 3). We tested

whether *HD* loci were potentially linked to centromeres as reported in other Basidiomycota

(13, 66). We used previously identified centromere locations for *Pgt* and *Pst* (47, 67) and

identified centromere locations for *Pca* and *Pt* using genome-wide Hi-C heatmaps based on

the bowtie-like Hi-C interaction features known for fungal centromeres (S8-11 Figs) (68).

This analysis revealed that *HD* loci are likely not directly linked to centromeres in any of the

four cereal rust fungi.

Lastly, we investigated overall nucleotide and gene coding regions conservation for the *HD*

locus including proximate regions containing 40 neighboring genes on either side of the locus.

We combined fine scale nucleotide synteny with gene conservation analysis. We specifically

tested if cereal rust fungi contain syntenic blocks with conserved protein coding genes

surrounding *MAT* loci as it was found for species in the genus *Trichosporonales* (64). We

used blastn to identify conserved nucleotide sequences within dikaryotic genomes and

between species. We plotted genes, transposons, nucleotide and protein coding gene

conservation for the *HD* loci proximate regions for all four cereal rust fungi (Fig 4A). Protein

coding genes and their order are mostly conserved within dikaryotic genomes of the same

species with 58/80 genes in *Pca 203*, 40/80 in *Pgt 21-0,* 57/80 in *Pt 76*, and 62/80 in *Pst 134E*

being conserved. Similarly, we observed considerable nucleotide conservation and synteny

within dikaryotic genomes with the exception of *Pgt 21-0*, which is consistent with our initial

mummer-based analysis (S7 Fig). Overall, synteny and protein coding gene conservation was

very limited between species for *HD* loci proximate regions. We could only identify three

conserved genes across all four cereal rust fungi. These three genes code for an integral

membrane protein (PTHR12459), D-2-hydroxyglutarate dehydrogenase (PTHR43716) and

Glucose-6-phosphate 1-sepimerase (IPR025532) with all three clustering downstream of the

HD locus (Fig 4A). Our TE analysis revealed that their coverage was overall consistent at the

order level within species but varied across species (S12 Fig). For example, TIR and other

undetermined class II DNA transposons dominated the *HD* locus of *Pst 134E,* while *HD* loci

of other rust fungi displayed higher coverage of class I RNA transposons including LTRs.

Overall, our analysis of the *HD* locus suggests that it is overall conserved within each

dikaryotic genome on a species level, while conservation has been eroded between species

over longer evolutionary timescales.

PR loci display strong signs of genomic deterioration

To investigate organization and synteny of the *PR* loci, we performed the same approach as

for the *HD* loci. Initial mummer based synteny analysis revealed clear macro- and micro-

synteny breaks around the *PR* loci in all cereal rust fungi (S13 Fig). It also revealed

accumulation of highly repetitive sequences closely associated with *PR* loci, especially in *Pca*

203 and *Pst 134E*. This increase in TEs correlates with a reduction in gene density around *PR*

loci (Figs 4B and 5). The loss in synteny was specific to *PR* loci, as overall macro-synteny of

chromosome 9 was conserved in all species (S13 Fig).

Consistent with the macro-synteny analysis, the d_S values for the *Pra* alleles *STE3.2-2-*

STE3.2-3 belonged to the 99.9% quantile of d_S values of all allele pairs found on chromosome

9 in all species (Fig 5 and S7 Fig). Gene density dropped significantly around *PRA* genes with

STE3.2-2/*STE3.2-3* located within or adjacent to 'gene deserts' that extended over 1 mbps in

the case of *Pst 134E*. Concomitant with gene depletion leading to gene deserts, *PR* loci are

enriched in TEs. One exception was the *PR* locus of *Pgt 21-0*, which might be a biological

feature or a technical artefact (see Discussion). None of the *Pra* alleles was closely linked to

centromeres except for *Pst 134E STE3.2-3,* which was found in close proximity to the

centromere at a distance of 164 kbp, embedded in a > 1 Mb long stretch of TEs depleted of

genes (Fig 5).

Detailed nucleotide synteny and protein coding gene conservation analysis confirmed signals

of extended recombination suppression, loss of synteny and accumulation of transposable

elements around *PR* loci (Fig 4B). We observed little conservation of nucleotide sequences

and protein coding genes in *Pra* proximate regions within dikaryotic genomes or between

species. Indeed, we could not identify any conserved protein-coding gene in *Pra* proximate

regions across the four cereal rust fungi. The TE composition in *Pra* proximate regions varied

within dikaryotic genomes and between species (S12B Fig). This indicates the accumulation

of distinct transposable element families around the *PR* locus in different cereal rust species

(see more detail for *Pt* below).

Overall, the *PR* locus appears highly plastic with little conservation within and between
species.

STE3.2-1 loci are highly conserved within and among cereal rust fungi

415   We analyzed *STE3.2-1* as above for the *MAT* loci. The three independent analyses revealed
416   that *STE3.2-1* and its surrounding regions are highly conserved at the nucleotide and protein
417   coding gene level within dikaryotic genomes while being overall syntenic between species
418   (S14-16 Figs). This clearly sets apart the *STE3.2-1* locus from the two *MAT* loci which further
419   supports the hypothesis that this locus is not involved in mate compatibility in rust fungi.

HD genes are multiallelic in cereal rust fungi while *Pra genes* display little variation at the species level

We aimed to extend our species level analysis of variation in *MAT* genes (Figs 1 and 2) using
publicly available genomic data for all four species (S1 Table). We identified between 12 and
15 representative isolates per species that were assigned to as many distinct genomic lineages
as possible based on previous genealogical studies to avoid repeated sampling from clonal
lineages that are predicted to show no genetic variation at *MAT* genes (30, 40-42, 44-46, 50-
53) (S1 Table). We quality controlled whole genome short read datasets and mapped them
against the respective reference genomes to estimate the genetic variation in *MAT* genes (S17-
20 and 22-25 Figs). We identified high levels of polymorphisms at the *HD* loci for all
analyzed cereal rust fungi. *Pca* showed the highest level of variation of *HD* loci at the species
level (S18 Fig). When mapping Illumina short read data of the *Pca* isolates against the *Pca*
203 reference, the *HD* loci in the *Pca* isolates displayed high levels of polymorphism
including heterozygous SNPs and multiple unaligned regions and gaps. This suggested that

the *HD* loci present in our additionally selected *Pca* isolates were not well reflected in the *Pca*

203 reference assembly. This is likely due to the fact that the selected isolates were highly

diverse sexual populations of *Pca* sampled in the USA (40). We therefore implemented a

novel *de novo* reconstruction approach for *HD* loci and *HD* coding sequences directly from

whole genome Illumina short read datasets (see Methods for detail). We confirmed our

approach by reconstructing the *Pca 203 HD* locus from publicly available Illumina short read

datasets. We confirmed that the two reconstructed *HD* alleles were identical to the ones

derived from the dikaryotic reference genome assembly (Fig 1) via a nucleotide alignment of

the two *bW-HD1* and *bE-HD2* coding sequences to the reconstructed *HD* alleles (S21A Fig).

In total, we were able to reconstruct eleven *bW-HD1* and twelve *bE-HD1* alleles for *Pca*

(S21B Fig). Using this novel approach, we identified extensive genetic variation for *bW-HD1*

and *bE-HD2* in *Pgt* and *Pt* (S21C-S21D Figs) while making use of recently published *bW-*

HD1 and *bE-HD2* alleles in case of *Pst* (21). We identified six, nine, and ten *bW-HD1* and six,

eight, and ten *bE-HD1* alleles for *Pgt, Pt,* and *Pst*, respectively. Overall, this suggests that the

HD genes are multiallelic in all four cereal rust fungi, which is consistent with other recent

reports (21, 32).

Compared to the *HD* genes, the *Pra* genes are reported to be far less polymorphic in many

basidiomycetes (9, 69, 70). Hence, we next focused our analysis on the *Pra* genes using the

same whole genome Illumina short read datasets (S1 Table). As expected, the *Pra* genes were

far less polymorphic within each species compared to *HD* loci (S22-26 Figs), which is

consistent with our initial analysis of *Pra* genes extracted from genome assemblies (Figs 1

and 2). In contrast to the highly polymorphic *HD* genes, we were able to identify only two

SNPs in the coding regions of *STE3.2-2* and *STE3.2-3* in *Pca* with a single SNP being non-

synonymous (S26 Fig). Similarly, *STE3.2-2* and *STE3.2-3* copies of all *Pst* isolates had

identical coding regions. Our short-read mapping analysis of *Pra* in *Pt* identified four SNPs in

coding sequence regions of *STE3.2-2* and *STE3.2-3*. These SNPs gave rise to a single

additional STE3.2-2 variant with two amino acid substitutions close to the C-terminus (S26

Fig). *Pgt* was the most polymorphic for *Pra*. We identified multiple distinct copies of *STE3.2-*

2 and *STE3.2-3* in the global *Pgt* population. We identified one STE3.2-2 variant with several

amino acid changes (S26B Fig). *STE3.2-3* was the most polymorphic in *Pgt* including two

isolates, *TTTSK* and *UVPgt60,* which contain potential nonsense mutations leading to pre-

mature stop codons (S26B Fig). Overall, our analysis suggests that *Pra* is most likely biallelic

in *Pca*, *Pt*, and *Pst* while *Pgt* may have more than two functional *Pra* alleles.

Lastly, we also investigated the nucleotide sequences of all *mfa* alleles at the species level.

Mfa1 and *Mfa2* alleles were fully conserved in *Pca* and *Pst*, while *Pt* had only a single non-

synonymous change in *mfa1/3* (S3 Fig). In contrast, *mfa1* and *mfa3* in *Pgt* had several non-

synonymous variations (S27 Fig). This species level variation in *Pgt* at the *mfa* loci is

consistent with significant variation observed in *STE3.2-2* and *STE3.2-3*.

MAT loci are conserved at the species level in *Puccinia triticina*

while showing transposable element expansion specific to the

STE3.2-3 locus

We made use of four chromosome scale and fully phased genome assemblies for *Pt* (Table 1)

to explore the structural conservation or plasticity of the *MAT* loci within a cereal rust fungal

species. We performed macro-synteny, detailed nucleotide synteny, protein coding gene

conservation and TE analysis for the *HD* and *PR* loci within and between *Pt 76* (43), *Pt 15*

(55), *Pt 19NSW04* and *Pt 20QLD87* (32). A recent study suggested that *Pt 19NSW04* arose

via somatic hybridization between *Pt 76* and *Pt 20QLD87* or close relatives in Australia mid-

2010 (32). Hence, *Pt 19NSW04* and *Pt 76* share one complete nuclear haplotype B, which

contains *STE3.2-2,* and *Pt 19NSW04* and *Pt 20QLD87* share one complete nuclear haplotype

C, which contains *STE3.2-3*. The *Pt15* isolate is unrelated and was sampled in China in 2015.

Macro-synteny plots of the *HD* locus containing chromosomes suggested that the *HD* loci of

shared nuclear haplotypes e.g. *Pt 19NSW04* and *Pt 76* hapB or *Pt 19NSW04* and *Pt 20QLD87*

hapC, are more conserved when compared to *HD* loci contained within each dikaryotic

genome (S6 and S28 Figs). Detailed nucleotide synteny and protein coding gene conservation

analysis revealed that gene order and function is highly conserved in the *HD* locus between

all tested nuclear haplotypes (S29 Fig). To investigate whether any specific TE families

cluster around the *HD* locus, we built a TE database based on the *Pt 76* reference genome and

used it to annotate TEs in all four strains. We identified similar TE coverage at the order level

around the *HD* locus in all dikaryotic genomes (S30A Fig).

We performed identical analysis for the *PR* locus to get a species level understanding of its

evolution in *Pt*. We compared the overall structural relationship between chromosomes

containing *STE3.2-2* or *STE3.2-3* using macro-synteny plots. Like the *HD* locus,

chromosomes of shared nuclear haplotypes, e.g. *Pt 19NSW04* vs *Pt 76* hapB with *STE3.2-2*,

were more similar to each other than non-shared haplotypes, e.g. *Pt 76* hapB vs *Pt15* hapA

with *STE3.2-3* (S13 and S31A Figs). The same applied for *STE3.2-2* (S13 and S31B Figs).

We could clearly observe clustering of repetitive sequences around the *STE3.2-2* and *STE3.2-

3* locus with the latter being more pronounced (S31 Fig). These repetitive sequences appeared

to be allele specific as they were not visible in macro-syntenty plots comparing *STE3.2-2* and

STE3.2-3 containing chromosomes (S13 Fig). Detailed nucleotide synteny and protein coding

gene analysis showed that genes around either *Pra* allele were mostly conserved while some

haplotypes appeared to have inversions downstream of *STE3.2-2* relative to the *Pt 76*

reference (Fig 6A and 6B). Gene synteny was more conserved for genes surrounding *STE3.2-

3* yet gene distance appeared to be especially variable between the two *mfa1/3s* and *STE3.2-3*

(Fig 6B). We explored the TE composition around each *Pra* allele and tested if TE expansions

could be an explanation for the difference in intergenic distances around the *STE3.2-3* locus.

TE composition around *STE3.2-2* and *STE3.2-3* were markedly different with the *STE3.2-3*

locus displaying an increase of LTR retrotransposons (Fig 6C and S30B Fig.). TE coverage

and composition around *STE3.2-2* was overall consistent between the different haplotypes

(S30B Fig). In the case of *STE3.2-3* we found a single LTR *Ty3* (also known as Gypsy (71))

TE family (Ty3_Pt_*STE3.2-3*, Fig 6D) highly expanded at the locus with varying coverage

between haplotypes ranging from 14.7% to 25.4% with *Pt 20QLD87* having the highest

coverage (Fig 6C). The percentage identity relative to the consensus sequence was > 99%

(Fig 6E) and most copies were around 8 kb (Fig 6F) which indicates active transposition of

Ty3_Pt_*STE3.2-3* in recent history. To investigate whether this TE family was specific to the

STE3.2-3 locus, we compared its abundance at the locus with its abundance across the whole

genome. We found that Ty3_Pt_*STE3.2-3* only clustered on *STE3.2-3* containing

chromosomes in all four strains with only a very limited number of single copies found on

other chromosomes (S32 Fig). Indeed, Ty3_Pt_*STE3.2-3* had a strong preference to insert

between *Ptmfa1/3*s and *STE3.2-3* (S33 Fig). To further investigate the structure of

Ty3_Pt_*STE3.2-3*, we used LTR_FINDER (72) to identify 5' and 3' LTR regions and other

features related to retrotransposons. Ty3_Pt_*STE3.2-3* was found to have a short LTR, around

150 bp in all four strains, with target site repeat (TSR) region of 'AAGT', *pol* genes

containing reverse transcriptase (RT), ribonuclease H (RH), integrase (INT). However, we

failed to identify any group-specific antigen (*gag*) in these TEs (Fig 6D).

MAT genes are expressed late in the asexual infection cycle during

urediniospore production

Importance of *MAT* loci in sexual reproduction were well established for *Pt* and *Pst* (18, 73),

yet it is unclear if they are expressed and functional during the asexual reproduction on cereal

hosts as found in *Pt* (18). Hence, we investigated the expression patterns of *MAT* genes in *Pca*,

Pgt and *Pst* using publicly available RNA-seq infection time series of their cereal hosts (38,

41, 48, 74) (S2 Table). We applied the trimmed mean of M-values (TMM) normalization to

read counts and assessed quality of the RNA-seq datasets by multidimensional scaling (MDS)

plots (S34 Fig). The MDS plots confirmed the suitability of the datasets for detailed

expression analysis based on technical replicates clustering closely together and dimension 1

separating samples according to their infection progress. The stable expression of two house

keeping genes for each species across all samples further confirmed the suitability of the

datasets for detailed *MAT* gene expression analysis (S35 Fig.).

STE3.2-2 and *STE3.2-3* were upregulated during the asexual infection process and always

displayed highest expression at the latest time point available, which coincides with

sporulation and production of urediniospores (Fig 7). We confirmed the differential

expression at later infection timepoints with likelihood ratio test with a p-value cut-off of $<$

0.05 (S36 Fig). Similarly, *HD* genes are upregulated during asexual infection of the cereal

host (Fig 7). We did not observe any expression of *STE3.2-1* in any of the samples. Lastly, we

were also interested to see if we could detect RNA of *MAT* genes in cereal rust spores before

infection and in specialized infection structures called haustoria. We used an additional *Pst*

RNA-seq dataset and compared the expression of *MAT* genes in ungerminated spores,

germinated spores, 5 days post infection, 9 days post infection and haustoria enriched samples

(S37 Fig) (48). *HD* genes displayed some expression in spores albeit lower than observed in

later stages of infection during spore production in the asexual cycle. In contrast, we could not

detect expression of *Pra* genes in ungerminated and germinated spores. Yet, *Pra* genes were

expressed and upregulated at later stages of the asexual infection consistent with other

publicly available datasets (Fig 7C and S37 Fig).

Discussion

Our comparative analysis of predicted *MAT* genes in four cereal rust fungi provides novel

insight into the evolution of these genes and their proximal regions within and between

species.

First, we confirmed that *HD* and *PR* loci are unlinked and located on two distinct

chromosomes in all four cereal rust fungi, which confirms previous analysis on more

fragmented genome assemblies or for individual rust fungal species (18, 21, 30). Further, both

loci were heterozygous in all twelve dikaryotic genome assemblies. Taken together, these

observations strongly support the hypothesis of tetrapolar mating types in rust fungi. Though

this genome biology informed hypothesis requires experimental validation via crosses and

genetic analysis of resulting offspring. As previously observed for *Pt* (18), the *Pra* homologs

STE3.2-1 were highly conserved in all cereal rust fungi and had no *mfa* genes closely

associated. This suggests that *STE3.2-1* does not regulate mate compatibility.

In *P. coronata* f. sp. *avenae*, *P. striiformis* f. sp. *graminis* and *P. striiformis* f. sp. *tritici*,

STE3.2-1 is flanked by anaphase promoting complex *APC10/Doc1* which regulates important

events in mitosis (75) and Elongation factor *EF1B* (76). Together, these findings suggest that

STE3.2-1 might play an important role during the cell cycle or different developmental

programs as reported for other basidiomycete fungi (77, 78). However as previously reported

for *P. triticina* (18), expression for *STE3.2-1* was absent or very low during the infection

cycle of the cereal host.

Heterothallic fungi with tetrapolar mating types are most often multiallelic at least at one of

the two *MAT* loci. Multiallelism is beneficial under outcrossing conditions, because it

increases the compatibility between randomly selected mates. At the same time, outcrossing

selects for new and rare alleles because these are compatible with existing and more common

alleles (9). For example, *U. maydis* has many distinct transcription factor alleles at the *HD*

locus (79, 80). We performed extensive analysis of *HD* loci extracted from dikaryotic genome

assemblies and developed a workflow that permitted us to *de novo* reconstruct loci directly

from Illumina short read sequencing data. This enabled us to investigate the allele status of

bW-HD1 and *bE-HD2* in 12-15 additional isolates for each cereal rust fungal species without

the need for long-read genome assemblies or targeted amplicon sequencing. In addition, the

analyzed isolates were selected based on them being the best available representation of

global rust fungal lineages without repeated sampling within clonal lineages to facilitate a

more in-depth analysis of the allele status of *MAT* loci. This extensive analysis demonstrated

that the *HD* locus is also multiallelic in the four cereal rust fungal species. The four species

had between six to twelve alleles of *bW-HD1* and *bE-HD2*. This supports previous analyses

of the allele status of these genes in *P. striiformis* f. sp. *tritici,* which identified nine alleles of

bW-HD1 and *bE-HD2* (21). The number of *bW-HD1* and *bE-HD2* alleles is likely a gross

underestimate giving limited public datasets. The majority of analyzed isolates were retrieved

from the cereal host during the asexual reproduction cycle and from regions that are known to

be deprived of sexually recombining populations. The only exception was *P. coronata* f. sp.

avenae isolates, which are reported to be derived from sexual recombining populations in the

United States of America (40). Indeed, we identified the highest allelic variation of *HD* genes

(eleven *bW-HD1* and twelve *bE-HD2*) in *P. coronata* f. sp. *avenae*. This is similar to previous

studies of isolates from likely sexual populations of *P. striiformis* f. sp. *tritici* in China, India,

and Pakistan, which showed the highest diversity in *HD* alleles compared to other clonal

lineages in Western wheat growing regions (21). In contrast to the multiallelic status of *HD*

genes, the *Pra* and associated *mfa* genes were clearly biallelic in *P. coronata* f. sp. *avenae*, *P.*

triticina and *P. striiformis* f. sp. *tritici*. In all cases, one *Pra* allele was linked to one specific

invariant *mfa* allele whereby *P. triticina* carried two *mfa1/3* genes that were nearly identical in

the sequence of the predicted mature pheromone peptide. This biallelism of *Pra/mfa* is similar

to other tetrapolar dimorphic basidiomycetes like *U. maydis* and *Cryptococcus sp.* (81, 82).

The notable exception was *P. graminis* f. sp. *tritici* which encodes two *mfa* genes (*mfa1* and

mfa3) in proximity to *STE3.2-3*. The two encoded peptides are highly variable and are likely

processed into two distinct mature pheromone peptides based on their sequence divergence,

which might lead to distinct receptor activation specificities. Interestingly, *Pgt STE3.2-3* was

also the most variable *Pra* allele in all species. We identified three *STE3.2-3* variants that

have at least four or more non-synonymous nucleotide changes relative to the most common

Pgt STE3.2-3 variant. Moreover, the *STE3.2-3* variants in *Pgt 126-6711*, *Pgt ME-02*, and *Pgt*

PK-01 shared four distinctive amino acid changes in the C-terminus while being linked to an

mfa1 allele that gives rise to MFA1 with four amino acid substitutions. At the same time all

analyzed *P. graminis* f. sp. *tritici* isolates carried an invariant *STE3.2-2* allele with only one

mfa gene in close proximity and we did not identify a clearly distinct additional *Pra* allele.

Hence the functional significance of the observed variation at the *STE3.2-3* locus is currently

difficult to assess without more extensive sampling of isolates, more phased chromosome

scale genome assemblies and in the absence of direct functional crossing studies in *P.*

graminis f. sp. *tritici*. Yet, the organization of the *STE3.2-3* locus is reminiscent of the *PR*

locus of *Sporisorium reilianum* in the order Ustilaginales. In *S. reilianum,* the *PR* locus is at

least triallelic with each allele encoding one pheromone receptor and two distinct pheromone

peptides. Each of the mature pheromones derived from one allele specifically activates only

one of the receptors encoded by the two other alleles (83). Taken together, more studies are

needed to assess if the *PR* locus in *P. graminis* f. sp. *tritici* is more than biallelic.

We initially predicted that biallelic *MAT* loci would show stronger signatures of

recombination suppression and genomic degeneration than multiallelic loci based on

published mathematical models (2) and observations in other basidiomycete species (9, 84).

Indeed, the multiallelic *HD* loci were mostly syntenic within dikaryotic genomes of the four

cereal rust species. This initial analysis with a limited set of genomes per species is likely to

hold true at the population level. In the case of *P. triticina,* the eight *HD* alleles of four

isolates were all highly syntenic with most of the genes being conserved between the different

alleles. Consistently, the TE composition at the order level was highly similar between *HD*

alleles of the same species. In contrast, *PR* loci showed strong signs of recombination

suppression and genomic degeneration. However, the *PR* loci did not follow the exact same

trends in all four species. The *PR* loci in *P. coronata* f. sp. *avenae*, *P. triticina* and *P.*

striiformis f. sp. *tritici* were the most extended with the *Pra* genes being located at the edge or

within large gene deserts and TE islands of ~0.7 mbp, ~0.7 mbp, and ~1.2 mbp, respectively.

At least some of these repeats were shared between the two *PR* alleles within each dikaryotic

genome of *P. coronata* f. sp. *avenae* and *P. striiformis* f. sp. *tritici* based on whole

chromosome alignments. This was not the case for *PR* alleles in *P. triticina*. In all three cases,

there appeared to be differences in TE composition and coverage at the order level between

the two *PR* alleles. This was most obvious for *P. triticina,* where we analyzed four dikaryotic

genome assemblies and identified the *STE3.2-3* specific TE family Ty3_Pt_*STE3.2-3*.

Ty3_Pt_*STE3.2-3* displayed preferential insertion for the *STE3.2-3* allele and insertions

between the pheromone receptor and the two *mfa* genes. Ty3_Pt_*STE3.2-3* is likely a

currently active TE based on its sequence conservation, length of individual TE copies, and

the suggested nuclear exchange between the studied Australian isolates within the last decade

(32). These initial observations of partial allele specific TE composition and coverage in *P.*

coronata f. sp. *avenae*, *P. triticina* and *P. striiformis* f. sp. *tritici* contrast with observations of

TE analysis of biallelic *MAT* loci in 15 *Microbotryum* spp.. The *MAT* loci in *Microbotryum*

spp. did not display allele specific TE composition patterns, yet were likely reservoirs for TE

families that drove TE expansion at a genome scale at discrete time points associated with the

extension of evolutionary strata around the *MAT* loci (37).

Compared to these observations at biallelic *PR* loci, the *PR* locus in *P. graminis* f. sp. *tritici*

contained fewer TEs, more genes, and was much shorter with ~300 kbp. In addition, the TE

composition and coverage was very similar between the two alleles within the dikaryotic

genome of *Pgt 21-0*. These observations could be biological or a technical artifact of the

assembly process of this specific genome. The *Pgt 21-0* assembly is based on older error

prone PacBio long-read technologies, included extensive manual curation and relied on gene

synteny between contigs and haplotypes for scaffolding. This gene based scaffolding might

have been broken by long-stretches of TEs around the *PR* locus as observed for the three

other cereal rusts and thereby introduced scaffolding errors in *Pgt 21-0* (30). Several of the

other genomes generated with older error prone PacBio long-read technologies had issues

assembling the *Pra* locus correctly. Alternatively, these marked differences of the *P. graminis*

f. sp. *tritici* *PR* locus might reflect biological reality, as models predict that recombination

suppression and genomic degeneration is more pronounced at biallelic loci (2). If the *PR*

locus in *P. graminis* f. sp. *tritici* is not truly biallelic as potentially indicated by the gene-

based analysis of *Pra* and *mfa* alleles, the organization and level of genomic degeneration at

the *PR* locus as observed would fulfill these predictions. However, more and better high-

quality phased chromosome scale genome assemblies are necessary to differentiate these

alternative hypotheses concerning the *PR* locus in *P. graminis* f. sp. *tritici*.

The functions of *MAT* genes in rust fungi are currently unknown, beyond their likely role in

mediating mate compatibility. However, in *P. triticina* and *M. larici-populina* (35, 85), *MAT*

genes are expressed during the sexual and asexual infection cycle. Suppression of *MAT* gene

expression in *P. triticina* during the asexual infection cycle on wheat reduces infection

severity and spore production (18). Here, we show that *MAT* genes are also upregulated

during the late stage of the asexual infection cycle of the cereal host in *P. coronata* f. sp.

avenae, *P. graminis* f. sp. *tritici,* and *P. striiformis* f. sp. *tritici*. This suggests a broader

conservation of *MAT* gene expression during the late stages of the asexual infection cycle of

rust fungi. Rust fungal cells carry more than two nuclei in the same cytosol during the asexual

infection cycle (86). At the same time, ureodiniospores always carry two nuclei of compatible

mating types and never two nuclei of identical mating types. Studies from the early 20[th]

century show that the two nuclei of ureodiniospore mother cells undergo synchronous nuclear

division within the same cytoplasm giving rise to four nuclei within the same ureodiniospore

precursor cells. This is followed by nuclear movement to restore appropriate nuclear pairing

and planar cell division that enables ureodiniospore maturation and reconstitutes the

dikaryotic state (86). The observed upregulation of *MAT* genes aligns with the development of

ureodiniospore primordia development and spore production. Hence, we hypothesize that

MAT genes are involved in the appropriate nuclear pairing at this stage of fungal development.

Understanding mating types in cereal rust fungi has also significant implications for

agriculture and for the prediction of the evolution of new virulent isolates. Recent reports

suggest that nuclear exchange between isolates during the asexual infection cycle (also known

as somatic hybridization (87)), in addition to sexual reproduction, can lead to nuclear

assortment (87). Nuclear exchange and the viability of the resulting offspring is likely

regulated by *MAT* genes because all proposed events have given rise to isolates with

dikaryotic genomes that carry opposing *MAT* gene pairs. We confirmed these observations

with our *MAT* gene analysis in the case of *Pgt Ug99* (*Pgt 21-0* and unknown nuclear donor)

and *Pt 19NSW04* (*Pt 76* and *Pt 20QLD87* as nuclear donors). In reverse, our analysis showed

that nuclear exchange has not occurred for the analyzed isolates of *P. coronata* f. sp. *avenae*,

and *P. striiformis* f. sp. *tritici*, because none of the isolates share identical *HD* and *Pra* alleles

that could be carried in the same nucleus. In cases of nuclear exchange, these nuclei behave

like entire linkage groups with virulence (also known as effectors) allele complements being

tied to specific *MAT* alleles if somatic hybridization occurs in the absence of parasexual

reproduction as currently proposed. This implies that the nuclear *MAT* alleles define the

possible virulence allele combinations that can arise via nuclear exchange. We hence can

predict the most likely novel virulence allele combinations that might arise given circulating

nuclear genomes in existing cereal rust fungal populations on their cereal hosts. This

observation requires us to generate phased and chromosome scale genome assemblies of all

cereal rust fungi at the population level. These will improve disease management strategies

and the durability of characterized resistance loci in the global wheat germplasm leading to

smarter rust resistance breeding programs.

Acknowledgments

We thank Dr. P. Tobias and Dr. M. Moeller for critical comments on manuscript, Dr. M. Bui

for advice in iqtree2 usage, Prof. C. Linde for useful suggestions, and R. Tam for providing a

snakemake pipeline for trimming raw sequencing data and centromere information of *Pst*

134E. We thank J. Lin for suggestions on data visualization. We acknowledge the excellent

and critical suggestions of the editor and peer reviewers that significantly improved the

manuscript.

Material and methods

Assessment of used genomic resources

Detailed information on isolates and genomes used in this study can be found in Table 1. Completeness of genome assemblies were assessed by BUSCO v5.5.0 (88) in the genome mode with a basidiomycota_odb10 set of genes.

Identification of *MAT* genes

MAT genes including the *HD* and *PRA* genes previously identified in *P. triticina* (Table 2) (18) were used as blastp and blastn (89) query to identify orthologs in *P. graminis* f.sp *tritici Ug99* and *21-0* isolates (30), *P. striiformis* f.sp *tritici 134E* (47), *104E* (48) and *DK0911* (49), *P. triticina 76* (43), *19NSW04*, *20QLD87* (32) and *15* (55), *P. polysora GD1913* (63), *P. coronate* f.sp *avenae 203* (39), *12NC29* and *12SD80* (38) isolates. Missed *STE3.2-3* in *Pst 104E* was fixed by mapping short reads to reference genome *Pst 134E* with bwa-mem2 (90). *Mfa genes* were identified by using *mfa* of *Pt BBBD*, *Pgt SCCL* and *Pst 78* (18) as a custom motif database to search similar pattern in all studied species with Geneious Prime v. 2022.1.1 (91), using a function built from fuzznuc (92). Karyograms of *Pca 203*, *Pgt 21-0*, *Pt 76* and *Pst 134E* were made with RIdeogram 0.2.2 (93).

Genealogical analysis of the *HD* and *Pra* genes

To construct a species tree for cereal rust fungi, protein sequences encoded in one nuclear genome from each cereal rust fungus species under study was used as input data. The input data was processed by Orthofinder 2.5.5 (61, 94) to infer maximum likelihood trees from multiple sequence alignments (MSA), fasttree v2.1 (95) was used to calculate likelihood-based local support values. Coding sequences of identified *HD* and *Pra* genes were aligned with MACSE v2.07 (96) with default setting. Coding-sequence based alignments were trimmed by using TrimAl v1.2 (97) with option (-gc 0.9) to remove sequences which have

gaps in more than 10% of samples, trimmed alignments were realigned with MACSE again.

Nucleotide alignments were imported into BEAUti v2.7.6 (98) to generate .xml files,

bModelTest 1.3.3 (99) was used to test the best nucleotide substitution model separately with

a Markov chain length by 15 million. HKY+G was chosen for inferring two *HD* genes,

whereas TN93+I was chosen for *Pra* genes and *STE3.2-1* individually. Then BEAST 2

MCMC runs under a strict molecular clock with the Yule Model tree, a 15 million length

Markov chains was used to provide a sufficient effective sample size (ESS) of independent

samples. Tracer v.1.7.2 (100) was used to check if ESS values of all parameters are above 200

for each run. TreeAnnotator (98) was used to generate a maximum clade credibility tree with

10% burn in. Figtree v.1.4.4 (101) was used to visualize the final output.

Nucleotide sequences of identified *mfa* genes were aligned with MACSE v2.07 (96) with

default settings, amino acid alignment generated was used to construct maximum likelihood

tree via IQ-TREE 2 (version 2.1.4-beta) (102), the best model was estimated by ModelFinder

(-m MFP) (103) based on Akaike Information Corrected Criterion (AICc) score (104), 10000

replicates were set for ultrafast bootstrap (105). JTT+G4 substitution model was chosen by

ModelFinder (103).

To compare topologies of *bW-HD1* and *bE-HD2* gene trees, IQ-TREE 2 (version 2.1.4-beta)

(102) was used to build unrooted maximum likelihood trees for each species and both genes

separately, the best model for each alignment estimated by ModelFinder (-m MFP) (103)

based on AICc score (104), 10000 replicates were set for ultrafast bootstrap (105). Top 100

optimal gene trees of *bW-HD1* and *bE-HD2* were used as alternate topologies after removing

branch length values to perform AU tests (62) with IQ-TREE2 (-keep-ident –zb 10000 -n 0 -

au), p_{AU} <0.05 suggesting topological incongruence between the two *HD* genes.

Estimation of centromeric regions

Hi-C reads of *Pca 203* and *Pt 76* were mapped back to their corresponding haploid genome

assembly with Juicer 1.6 (68) and processed with the 3d-dna (106) pipeline. Juicerbox 1.8.8

(68) was used to visualize Hi-C maps. The approximate positions of centromeric regions were

estimated by zooming in on the Hi-C heatmap in Juicerbox and selecting the region

corresponding to strong interaction signals, which has been validated and widely used in

previous studies (107, 108).

775 d_s value estimation for allele pairs and plotting

Inspired by the approach used by Branco et al. (8) for identifying evolutionary strata on

mating-type chromosomes of *Microbotryum* species, proteinortho 6.0.22 (109) was used to

pair orthologs one-to-one with -synteny and -singles tag. Muscle 3.8.1551 (110) was applied

to generate coding sequence based protein alignments with default settings. PAML 4.9 (111)

was used to calculate synonymous divergence values from alignments generated with Muscle.

Genomes were split in 10 kbp per window, then density of genes and repeats were calculated

with bedtools v 2.29.2 (112). Results were visualized with karyoploteR v3.17 (113).

783 Synteny analysis of *MAT* loci and flanking regions

Self-alignment analyses between two haplotypes of each species were done with Nucmer

(version 3.23) (65) with –maxmatch option, alignments with a size of less than 1000 bp or

less than 90 % identity were removed, filtered data were visualized via Matplotlib (114).

Nucmer (version 3.23) (65) was used to align nucleotide sequences of *MAT* regions.

Blastn 2.10.0 (89) was used to identified conserved genes (identity higher than 70%, e-value

< 0.05) across and within species. Results were visualized with gggenomes 0.9.9.9000 (115).

790 General repeat annotations

REPET v3.0 pipeline (116, 117) was applied to annotate repeats in all genomes used in this

study, together with Repbase v22.05 (118), config files were based on default settings.

TEdenovo was used to predict novel repetitive elements and to construct specific custom

databases from each genome individually as described in the manual instructions. Constructed

custom databases were used for repeat annotation with TEannot. The reference TE database

built for *Pt 76* from TEdenovo with the REPET v3.0 pipeline was used as reference for

annotating repeats in *Pt 15* (55), *Pt 19NSW04* and *Pt 20QLD87* (32) with RepeatMasker 4.1.5

(119) To make analyses consistent, the *Pt 76* genome was reannotated in the same way as the

other three isolates.

To calculate the coverage of each TE family, classification information from TEannot was

used. For TEs with more than one potential classification, hits from Repbase which have more

than 70% of similarity were used as evidence for reclassification.

LTR_FINDER v1.07 (72) with ps_scan (120, 121) was used to identify detailed structures of

LTRs of Ty3_Pt_*STE3.2-3* in default setting. Reverse transcriptase (A0A0C4EV49_PUCT1),

Retrotransposon gag domain (A0A0C4EKM8_PUCT1), Integrase, catalytic domain

(A0A0C4EMG6_PUCT1) and Retropepsins (A0A0C4F2V2_PUCT1) (122), were

additionally used as query to search for motifs in identified LTR retrotransposons using

tblastn 2.12.0+ (89).

Identification of variants in *MAT* loci

Sequence data of *Pca* were published by (40), data of *Pgt* were published by (30, 41, 42, 53)

and under PRJNA39437, data of *Pt* used in this study have been published earlier by (44-46)

and under PRJNA39803 and PRJNA39801, data of *Pst* were published by (50-52) and under

PRJNA60743, all above data were downloaded from NCBI SRA database (URL:

https://www.ncbi.nlm.nih.gov/sra) (Table S1). Four *Pca* isolates, *Pca 203* (39), *Pgt 21-0* (30),

Pt 76 (43) and *Pst 134E* (47), with published chromosome-scale haplotype-phased genome

assemblies were used as reference for mapping. Trimmomatic 0.39.2 (123) was used to

remove adapters from raw reads whereas FastQC v0.11.8 (124) was applied for assessing the

quality before and after trimming. Bwa-mem2 (90) was used to map trimmed reads to

reference genomes, MarkDuplicates (Picard) 1.70 (125) was applied to remove PCR-

generated duplicates. Samtools 1.12 (126) was used to remove mapped reads with mapping

quality lower than 30. IGV 2.16.2 (127) and Qualimap 2 (128) were both used to check

mapping quality.

For reconstruction of *Pra* alleles, freebayes-parallel 1.3.6 (129, 130) was applied to detect

variants, bcftools 1.12 (126) was used to exclude low quality variants with 'QUAL<40'.

Samtools 1.12 (126) was used to filter out regions of interest. The Ensembl Variant Effect

Predictor (VEP) 88.9 (131) was used to annotate sequence variants. Bcftools 1.12 (126) was

used to generate consensus sequences, nucleotide and protein alignments were generated and

visualized with Geneious Prime v. 2022.1.1 (91).

For *de novo* reconstruction of *HD* alleles, Spades (132), a de novo assembler, was used for

building draft genome assemblies from available whole genome Illumina short read datasets

(Table S1). For isolates with average read lengths > 101 bp, k-mer size of 101 was chosen for

building draft genomes. For isolates with average read lengths < 101 bp, k-mer size of 51 bp

was chosen. *HD* genes used in genealogical analysis were used as reference genes, blastn

2.12.0+ (89) was used to search and identify *HD* allele containing contigs of the draft genome

assemblies. KAT (133) was used to compute a k-mer matrix from *de novo* assembled *HD*

contigs and reference *HD* loci sequences with default setting (k-mer size=27). KAT filter was

used to obtain all paired-end reads which contained *HD* k-mers based on the created *HD* k-

mer matrix (-T 0.2). Spades was used to reassemble all *HD* related paired-end reads of the

specific isolate. Final outputs were visualized in Geneious, CDS regions of reconstructed *HD*

alleles were predicted by mapping the CDS of reference *HD* alleles to reconstructed contigs,

ORFs were predicted with Geneious.

Comparison of *MAT* loci among *Pt* isolates

Fully phased genome assemblies of *Pt 15* (55), *Pt 19NSW04* and *Pt 20QLD87* (32) were

downloaded from NCBI SRA database (URL: https://www.ncbi.nlm.nih.gov/sra), annotation

of *Pt 19NSW04* and *Pt 20QLD87* were downloaded from CSIRO Data Access Portal (URL:

https://data.csiro.au/). CDS and protein data of *Pt 19NSW04* and *Pt 20QLD87* were generated

with Gffread (134), *MAT* genes were identified with blastn 2.12.0 (89) using *MAT* genes in *Pt*

76 as query. Nucmer 3.2.3 was applied to make pair-wise comparisons between *Pt* isolates, --

maxmatch tag was used to allow identification of maximum hits, macrosynteny plots were

generated with Matplotlib (114) after removing matches with < 90% identity and shorter than

1000 bp. Blastn 2.10.0 (89) was used to identify orthologs with at least 70 % identity.

Microsynteny plots were generated with gggenome2 (115).

Comparison of expression of *MAT* genes in asexual reproduction

RNA-seq reads of rust fungi on infected plants were obtained from NCBI SRA database

(URL: https://www.ncbi.nlm.nih.gov/sra). The datasets included: RNA-seq data of *Pst* 87/66

on infected wheat (PRJEB12497 (74), *Pca* 12*NC29* on *Brachypodium distachyon*

(PRJNA398546 (38), *Pgt* 21-0 on infected wheat (PRJNA415866 (41), *Pst 104E*

(PRJNA396589 (48). Raw reads were quality checked with FastQC v0.11.8 (124),

trimmomatic 0.50 (123) was used to trim low quality reads with parameters:

ILLUMINACLIP:adapter.fa:2:30:10 LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15

MINLEN:36. Kallisto 0.44.0 (135) was then applied to align and count transcripts with -b 100

parameter for all paired-end reads and -b 50 -l 200 -s 20 --single --single-overhang for single-

end reads. *Pst 104E* cDNA data was used as reference for *Pst 104E* infected tissues. Since no

reference genome of *Pst 87/66* is available, cDNA data of *Pst 134E* was used as reference

instead with *HD* genes of *Pst 87/66* supplemented to obtain better quantification of the

specific *HD* alleles. *Pgt 21-0* and *Pca 12NC29* were used as reference for mapping reads from

Pgt 21-0 and *Pca 12NC29* infected tissues, respectively. Gene names of *Pca 12NC29* and *Pst*

104E were added with funannotate annotate v 1.8.5 (136, 137) with default parameters.

Kallisto outputs were imported to EdgeR 3.17 (138) with tximportdata (139) following user

instruction, followed by normalizing with the TMM method, significant upregulation of *MAT*

genes was assessed with the LRT method and ggplot2 (140) was used to visualize the result.

House keeping genes were identified in each dataset based on the following conditions: *p-*

value>0.1 and log(FC) <0.5, functional annotations of candidate house keeping genes were

retrieved from interproscan (137).

Data Availability

Analysis code used in this study is available at github repository:

https://github.com/ZhenyanLuo/codes-used-for-mating-type. Alignments of genealogical

studies, d_s value of gene pairs in studied cereal rust species, CDS of reconstructed *HD* alleles,

Pra alleles, TE annotation and classification files are available at Dryad

(https://doi.org/10.5061/dryad.w0vt4b8zm).

References
882

1.   Bachtrog D. A dynamic view of sex chromosome evolution. Current opinion in

genetics & development. 2006;16(6):578-85.

2.   Jay P, Tezenas E, Véber A, Giraud T. Sheltering of deleterious mutations explains the

stepwise extension of recombination suppression on sex chromosomes and other supergenes.

PLoS Biology. 2022;20(7):e3001698.

3.   Duhamel M, Carpentier F, Begerow D, Hood ME, Rodríguez de la Vega RC, Giraud

T. Onset and stepwise extensions of recombination suppression are common in mating□type

chromosomes of *Microbotryum anther□smut* fungi. Journal of Evolutionary Biology. 2022.

4.   Carpentier F, Rodríguez de la Vega RC, Jay P, Duhamel M, Shykoff JA, Perlin MH,

et al. Tempo of degeneration across independently evolved nonrecombining regions.

Molecular Biology and Evolution. 2022;39(4):msac060.

5.   Idnurm A, Hood ME, Johannesson H, Giraud T. Contrasted patterns in mating-type

chromosomes in fungi: hotspots versus coldspots of recombination. Fungal Biology Reviews.

2015;29(3-4):220-9.

6.   Hartmann FE, Duhamel M, Carpentier F, Hood ME, Foulongne□Oriol M, Silar P, et

al. Recombination suppression and evolutionary strata around mating□type loci in fungi:

documenting patterns and understanding evolutionary and mechanistic causes. New

Phytologist. 2021;229(5):2470-91.

7.   Bazzicalupo AL, Carpentier F, Otto SP, Giraud T. Little evidence of antagonistic

selection in the evolutionary strata of fungal mating-type chromosomes (*Microbotryum*

lychnidis-dioicae). G3: Genes, Genomes, Genetics. 2019;9(6):1987-98.

8.   Branco S, Badouin H, de la Vega RCR, Gouzy J, Carpentier F, Aguileta G, et al.

Evolutionary strata on young mating-type chromosomes despite the lack of sexual antagonism.

Proceedings of the National Academy of Sciences. 2017;114(27):7067-72.

9.   Coelho MA, Bakkeren G, Sun S, Hood ME, Giraud T. Fungal sex: the Basidiomycota.

The fungal kingdom. 2017:147-75.

10.   Menkis A, Jacobson DJ, Gustafsson T, Johannesson H. The mating-type chromosome

in the filamentous ascomycete *Neurospora tetrasperma* represents a model for early evolution

of sex chromosomes. PLoS Genetics. 2008;4(3):e1000030.

11.   Hartmann FE, Ament-Velásquez SL, Vogan AA, Gautier V, Le Prieur S, Berramdane

M, et al. Size variation of the nonrecombining region on the mating-type chromosomes in the

fungal *Podospora anserina* species complex. Molecular biology and evolution.

2021;38(6):2475-92.

12.   Vittorelli N, Rodríguez de la Vega RC, Snirc A, Levert E, Gautier V, Lalanne C, et al.

Stepwise recombination suppression around the mating-type locus in an ascomycete fungus

with self-fertile spores. PLoS Genetics. 2023;19(2):e1010347.

13.   Carpentier F, de la Vega RCR, Branco S, Snirc A, Coelho MA, Hood ME, et al.

Convergent recombination cessation between mating-type genes and centromeres in selfing

anther-smut fungi. Genome research. 2019;29(6):944-53.

14.   Aime M, McTaggart A. A higher-rank classification for rust fungi, with notes on

genera. Fungal systematics and evolution. 2021;7(1):21-47.

15.   Dean R, Van Kan JA, Pretorius ZA, Hammond Kosack KE, Di Pietro A, Spanu PD,

et al. The Top 10 fungal pathogens in molecular plant pathology. Molecular plant pathology.

2012;13(4):414-30.

16.   Savary S, Willocquet L, Pethybridge SJ, Esker P, McRoberts N, Nelson A. The global

burden of pathogens and pests on major food crops. Nature ecology & evolution.

2019;3(3):430-9.

17.   Fraser JA, Diezmann S, Subaran RL, Allen A, Lengeler KB, Dietrich FS, et al.

Convergent evolution of chromosomal sex-determining regions in the animal and fungal

kingdoms. PLoS biology. 2004;2(12):e384.

18.   Cuomo CA, Bakkeren G, Khalil HB, Panwar V, Joly D, Linning R, et al. Comparative

analysis highlights variable genome content of wheat rusts and divergence of the mating loci.

G3: Genes, Genomes, Genetics. 2017;7(2):361-76.

19.   Wu JQ, Song L, Ding Y, Dong C, Hasan M, Park RF. A chromosome-scale assembly

of the wheat leaf rust pathogen *Puccinia triticina* provides insights into structural variations

and genetic relationships with haplotype resolution. Frontiers in Microbiology. 2021:2180.

20.   Maia TM, Lopes ST, Almeida JMGCF, Rosa LH, Sampaio JP, Gonçalves P, et al.

Evolution of Mating Systems in Basidiomycetes and the Genetic Architecture Underlying

Mating-Type Determination in the Yeast *Leucosporidium scottii*. Genetics. 2015;201(1):75-

89.

21.   Holden S, Bakkeren G, Hubensky J, Bamrah R, Abbasi M, Qutob D, et al. Uncovering

the history of recombination and population structure in western Canadian stripe rust

populations through mating type alleles. BMC Biology. 2023;21(1):233.

22.   Badouin H, Hood ME, Gouzy J, Aguileta G, Siguenza S, Perlin MH, et al. Chaos of

rearrangements in the mating-type chromosomes of the anther-smut fungus *Microbotryum*

lychnidis-dioicae. Genetics. 2015;200(4):1275-84.

23.   Petit E, Giraud T, de Vienne DM, Coelho MA, Aguileta G, Amselem J, et al.

LINKAGE TO THE MATING☐TYPE LOCUS ACROSS THE GENUS *MICROBOTRYUM*:

INSIGHTS INTO NONRECOMBINING CHROMOSOMES. Evolution. 2012;66(11):3519-

33.

24.   Coelho MA, Ianiri G, David-Palma M, Theelen B, Goyal R, Narayanan A, et al.

Frequent transitions in mating-type locus chromosomal organization in *Malassezia* and early

steps in sexual reproduction. Proceedings of the National Academy of Sciences.

2023;120(32):e2305094120.

25.   Bazzicalupo AL, Carpentier F, Otto SP, Giraud T. Little Evidence of Antagonistic

Selection in the Evolutionary Strata of Fungal Mating-Type Chromosomes (*Microbotryum*

lychnidis-dioicae). G3 Genes|Genomes|Genetics. 2019;9(6):1987-98.

26.   Buller AHR. Researches on Fungi, Vol. VII: The Sexual Process in the *Uredinales*:

University of Toronto Press; 1950.

27.   Narisawa K, Yamaoka Y, Katsuya K. Mating type of isolates derived from the

spermogonial state of *Puccinia coronata* var. *coronata*. Mycoscience. 1994;35(2):131-5.

28.   Nemri A, Saunders DG, Anderson C, Upadhyaya NM, Win J, Lawrence G, et al. The

genome sequence and effector complement of the flax rust pathogen *Melampsora lini*.

Frontiers in plant science. 2014;5:98.

29.   Ferrarezi J, McTaggart A, Tobias P, Hayashibara C, Degnan R, Shuey L, et al.

Austropuccinia psidii uses tetrapolar mating and produces meiotic spores in older infections

on *Eucalyptus grandis*. Fungal Genetics and Biology. 2022:103692.

30.   Li F, Upadhyaya NM, Sperschneider J, Matny O, Nguyen-Phuc H, Mago R, et al.

Emergence of the *Ug99* lineage of the wheat stem rust pathogen through somatic

hybridisation. Nature communications. 2019;10(1):1-15.

31.   Devier B, Aguileta G, Hood ME, Giraud T. Ancient trans-specific polymorphism at

pheromone receptor genes in basidiomycetes. Genetics. 2009;181(1):209-23.

32.   Sperschneider J, Hewitt T, Lewis DC, Periyannan S, Milgate AW, Hickey LT, et al.

Nuclear exchange generates population diversity in the wheat leaf rust pathogen *Puccinia*

triticina. Nature Microbiology. 2023:1-12.

33.   Kronstad J, Leong S. The b mating-type locus of *Ustilago maydis* contains variable

and constant regions. Genes & development. 1990;4(8):1384-95.

34.   Casselton LA, Olesnicky NS. Molecular genetics of mating recognition in

basidiomycete fungi. Microbiology and molecular biology reviews. 1998;62(1):55-70.

35.   Duplessis S, Hacquard S, Delaruelle C, Tisserant E, Frey P, Martin F, et al.

Melampsora larici-populina transcript profiling during germination and timecourse infection

of poplar leaves reveals dynamic expression patterns associated with virulence and biotrophy.

Molecular plant-microbe interactions. 2011;24(7):808-18.

36.   Hood ME, Antonovics J, Koskella B. Shared forces of sex chromosome evolution in

haploid-mating and diploid-mating organisms: *Microbotryum violaceum* and other model

organisms. Genetics. 2004;168(1):141-6.

37.   Duhamel M, Hood ME, Rodríguez de la Vega RC, Giraud T. Dynamics of

transposable element accumulation in the non-recombining regions of mating-type

chromosomes in anther-smut fungi. Nature Communications. 2023;14(1):5692.

38.   Miller ME, Zhang Y, Omidvar V, Sperschneider J, Schwessinger B, Raley C, et al. *De*

novo assembly and phasing of dikaryotic genomes from two isolates of *Puccinia coronata* f.

sp. *avenae*, the causal agent of oat crown rust. MBio. 2018;9(1):e01650-17.

39.   Henningsen EC, Hewitt T, Dugyala S, Nazareno ES, Gilbert E, Li F, et al. A

chromosome-level, fully phased genome assembly of the oat crown rust fungus *Puccinia*

coronata f. sp. *avenae*: a resource to enable comparative genomics in the cereal rusts. G3.

2022;12(8):jkac149.

40.   Miller ME, Nazareno ES, Rottschaefer SM, Riddle J, Dos Santos Pereira D, Li F, et al.

Increased virulence of *Puccinia coronata* f. sp. *avenae* populations through allele frequency

changes at multiple putative Avr loci. PLoS genetics. 2020;16(12):e1009291.

41.   Chen J, Upadhyaya NM, Ortiz D, Sperschneider J, Li F, Bouton C, et al. Loss of

AvrSr50 by somatic exchange in stem rust leads to virulence for *Sr50* resistance in wheat.

Science. 2017;358(6370):1607-10.

42.   Lewis CM, Persoons A, Bebber DP, Kigathi RN, Maintz J, Findlay K, et al. Potential

for re-emergence of wheat stem rust in the United Kingdom. Communications biology.

2018;1(1):1-9.

43.   Duan H, Jones AW, Hewitt T, Mackenzie A, Hu Y, Sharp A, et al. Physical separation

of haplotypes in dikaryons allows benchmarking of phasing accuracy in Nanopore and HiFi

assemblies with Hi-C data. Genome biology. 2022;23(1):84.

44.   Wu JQ, Dong C, Song L, Park RF. Long-Read–Based de novo Genome Assembly and

Comparative genomics of the wheat leaf rust pathogen *Puccinia triticina* identifies candidates

for three avirulence genes. Frontiers in genetics. 2020;11:521.

45.   Wu JQ, Sakthikumar S, Dong C, Zhang P, Cuomo CA, Park RF. Comparative

genomics integrated with association analysis identifies candidate effector genes

corresponding to *Lr20* in phenotype-paired *Puccinia triticina* isolates from Australia.

Frontiers in plant science. 2017;8:148.

46.   Fellers JP, Sakthikumar S, He F, McRell K, Bakkeren G, Cuomo CA, et al. Whole-

genome sequencing of multiple isolates of *Puccinia triticina* reveals asexual lineages

evolving by recurrent mutations. G3. 2021;11(9):jkab219.

47.   Schwessinger B, Jones A, Albekaa M, Hu Y, Mackenzie A, Tam R, et al. A

Chromosome Scale Assembly of an Australian *Puccinia striiformis* f. sp. *tritici* Isolate of the

PstS1 Lineage. Molecular Plant-Microbe Interactions. 2022;35(3):293-6.

48.   Schwessinger B, Sperschneider J, Cuddy WS, Garnica DP, Miller ME, Taylor JM, et

al. A near-complete haplotype-phased genome of the dikaryotic wheat stripe rust fungus

Puccinia striiformis f. sp. *tritici* reveals high interhaplotype diversity. MBio.

2018;9(1):e02275-17.

49.   Schwessinger B, Chen Y-J, Tien R, Vogt JK, Sperschneider J, Nagar R, et al. Distinct

life histories impact dikaryotic genome evolution in the rust fungus *Puccinia striiformis*

causing stripe rust in wheat. Genome biology and evolution. 2020;12(5):597-617.

50.   van Schalkwyk HJ, Adams T, Persoons A, Boshoff WH, Wanyera R, Hovmøller MS,

et al. Pathogenomic analyses of *Puccinia striiformis* f. sp. *tritici* supports a close genetic

relationship between South and East Africa. Plant Pathology. 2022;71(2):279-88.

51.   Hubbard A, Lewis CM, Yoshida K, Ramirez-Gonzalez RH, de Vallavieille-Pope C,

Thomas J, et al. Field pathogenomics reveals the emergence of a diverse wheat yellow rust

population. Genome biology. 2015;16(1):1-15.

52.   Ding Y, Cuddy WS, Wellings CR, Zhang P, Thach T, Hovmøller MS, et al. Incursions

of divergent genotypes, evolution of virulence and host jumps shape a continental clonal

population of the stripe rust pathogen *Puccinia striiformis*. Molecular Ecology.

2021;30(24):6566-84.

53.   Upadhyaya NM, Garnica DP, Karaoglu H, Sperschneider J, Nemri A, Xu B, et al.

Comparative genomics of Australian isolates of the wheat stem rust pathogen *Puccinia*

graminis f. sp. *tritici* reveals extensive polymorphism in candidate effector genes. Frontiers in

plant science. 2015;5:759.

54.   Henningsen EC, Hewitt T, Dugyala S, Nazareno E, Gilbert E, Li F, et al. A

chromosome-level, fully phased genome assembly of the oat crown rust fungus *Puccinia*

coronata f. sp. *avenae*: a resource to enable comparative genomics in the cereal rusts. bioRxiv.

2022.

55.   Li C, Qiao L, Lu Y, Xing G, Wang X, Zhang G, et al. Gapless Genome Assembly of

Puccinia triticina Provides Insights into Chromosome Evolution in *Pucciniales*. Microbiology

Spectrum. 2023;11(1):e02828-22.

56.   Schwessinger B, Sperschneider J, Cuddy WS, Garnica DP, Miller ME, Taylor JM, et

al. A near-complete haplotype-phased genome of the dikaryotic wheat stripe rust fungus

Puccinia striiformis f. sp. *tritici* reveals high interhaplotype diversity. MBio. 2018;9(1).

57.   Persoons A, Morin E, Delaruelle C, Payen T, Halkett F, Frey P, et al. Patterns of

genomic variation in the poplar rust fungus *Melampsora larici-populina* identify

pathogenesis-related factors. Frontiers in plant science. 2014;5:450.

58.   Raudaskoski M, Kothe E. Basidiomycete mating type genes and pheromone signaling.

Eukaryotic cell. 2010;9(6):847.

59.   van Diepen LTA, Olson Å, Ihrmark K, Stenlid J, James TY. Extensive Trans-Specific

Polymorphism at the Mating Type Locus of the Root Decay Fungus Heterobasidion.

Molecular Biology and Evolution. 2013;30(10):2286-301.

60.   Aime MC, Bell CD, Wilson AW. Deconstructing the evolutionary complexity

between rust fungi (*Pucciniales*) and their plant hosts. Studies in Mycology. 2018;89(1):143-

52.

61.   Emms DM, Kelly S. STRIDE: species tree root inference from gene duplication

events. Molecular biology and evolution. 2017;34(12):3267-78.

62.   Shimodaira H. An approximately unbiased test of phylogenetic tree selection.

Systematic biology. 2002;51(3):492-508.

63.   Liang J, Li Y, Dodds PN, Figueroa M, Sperschneider J, Han S, et al. Haplotype-

phased and chromosome-level genome assembly of *Puccinia polysora*, a giga-scale fungal

pathogen causing southern corn rust. Molecular Ecology Resources. 2023;23(3):601-20.

64.   Sun S, Coelho MA, Heitman J, Nowrousian M. Convergent evolution of linked

mating-type loci in basidiomycete fungi. PLoS genetics. 2019;15(9):e1008365.

65.   Marçais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A. MUMmer4:

A fast and versatile genome alignment system. PLoS computational biology.

2018;14(1):e1005944.

66.   Hood ME, Scott M, Hwang M. Breaking linkage between mating compatibility factors:

Tetrapolarity in *Microbotryum*. Evolution. 2015;69(10):2561-72.

67.   Sperschneider J, Jones AW, Nasim J, Xu B, Jacques S, Zhong C, et al. The stem rust

fungus *Puccinia graminis* f. sp. *tritici* induces centromeric small RNAs during late infection

that are associated with genome-wide DNA methylation. BMC biology. 2021;19(1):1-25.

68.   Durand NC, Robinson JT, Shamim MS, Machol I, Mesirov JP, Lander ES, et al.

Juicebox provides a visualization system for Hi-C contact maps with unlimited zoom. Cell

systems. 2016;3(1):99-101.

69.   Coelho MA, Sampaio JP, Goncalves P. A deviation from the bipolar-tetrapolar mating

paradigm in an early diverged basidiomycete. PLoS genetics. 2010;6(8):e1001052.

70.   Urban M, Kahmann R, Bölker M. The biallelica mating type locus *maydis*: remnants

of an additional pheromone gene indicate evolution from a multiallelic ancestor. Molecular

and General Genetics MGG. 1996;250(4):414-20.

71.   Wei K, Aldaimalani R, Mai D, Zinshteyn D, Satyaki P, Blumenstiel JP, et al.

Rethinking the "*gypsy*" retrotransposon: A roadmap for community-driven reconsideration of

problematic gene names. 2022.

72.   Xu Z, Wang H. LTR_FINDER: an efficient tool for the prediction of full-length LTR

retrotransposons. Nucleic acids research. 2007;35(suppl_2):W265-W8.

73.   Zhan G, Guo J, Tian Y, Ji F, Bai X, Zhao J, et al. High-throughput RNA sequencing

reveals differences between the transcriptomes of the five spore forms of *Puccinia striiformis*

f. sp. *tritici*, the wheat stripe rust pathogen. Stress Biology. 2023;3(1):29.

74.   Dobon A, Bunting DC, Cabrera-Quio LE, Uauy C, Saunders DG. The host-pathogen

interaction between wheat and yellow rust induces temporally coordinated waves of gene

expression. BMC genomics. 2016;17(1):1-14.

75.   Au SW, Leng X, Harper JW, Barford D. Implications for the ubiquitination reaction of

the anaphase-promoting complex from the crystal structure of the *Doc1/Apc10* subunit.

Journal of molecular biology. 2002;316(4):955-68.

76.   Andersen GR, Nissen P, Nyborg J. Elongation factors in protein biosynthesis. Trends

in biochemical sciences. 2003;28(8):434-41.

77.   Wirth S, Freihorst D, Krause K, Kothe E. What role might non-mating receptors play

in *Schizophyllum commune*? Journal of Fungi. 2021;7(5):399.

78.   Hsueh YP, Xue C, Heitman J. A constitutively active GPCR governs morphogenic

transitions in *Cryptococcus neoformans*. The EMBO journal. 2009;28(9):1220-33.

79.   Zambino P, Groth JV, Lukens L, Garton JR, May G. Variation at the b mating type

locus of *Ustilago maydis*. Phytopathology. 1997;87(12):1233-9.

80.   Silva J. Alleles at the b incompatibility locus in Polish and North American

populations of *Ustilago maydis* (DC) Corda. Physiological Plant Pathology. 1972;2(4):333-7.

81.   Metin B, Findley K, Heitman J. The mating type locus (*MAT*) and sexual reproduction

of *Cryptococcus heveanensis*: insights into the evolution of sex and sex-determining

chromosomal regions in fungi. PLoS Genetics. 2010;6(5):e1000961.

82.   Bölker M, Urban M, Kahmann R. The *a* mating type locus of *U. maydis* specifies cell

signaling components. Cell. 1992;68(3):441-50.

83.   Schirawski J, Heinze B, Wagenknecht M, Kahmann R. Mating type loci of

Sporisorium reilianum: novel pattern with three *a* and multiple *b* specificities. Eukaryotic cell.

2005;4(8):1317-27.

84.   Fontanillas E, Hood ME, Badouin H, Petit E, Barbe V, Gouzy J, et al. Degeneration of

the nonrecombining regions in the mating-type chromosomes of the anther-smut fungi.

Molecular biology and evolution. 2015;32(4):928-43.

85.   Duplessis S, Cuomo CA, Lin Y-C, Aerts A, Tisserant E, Veneault-Fourrey C, et al.

Obligate biotrophy features unraveled by the genomic analysis of rust fungi. Proceedings of

the National Academy of Sciences. 2011;108(22):9166-71.

86.   Savile DBO. Nuclear structure and behavior in species of the *Uredinales*. American

Journal of Botany. 1939:585-609.

87.   Park RF, Wellings CR. Somatic hybridization in the *Uredinales*. Annual review of

Phytopathology. 2012;50:219-39.

88.   Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. BUSCO update: novel

and streamlined workflows along with broader and deeper phylogenetic coverage for scoring

of eukaryotic, prokaryotic, and viral genomes. Molecular biology and evolution.

2021;38(10):4647-54.

89.   Johnson M, Zaretskaya I, Raytselis Y, Merezhuk Y, McGinnis S, Madden TL. NCBI

BLAST: a better web interface. Nucleic acids research. 2008;36(suppl_2):W5-W9.

90.   Li H, Durbin R. Fast and accurate short read alignment with Burrows–Wheeler

transform. bioinformatics. 2009;25(14):1754-60.

91.   Geneious Prime 2022.0.2 [Available from: https://www.geneious.com.

92.   Rice P, Longden I, Bleasby A. EMBOSS: the European molecular biology open

software suite. Elsevier current trends; 2000.

93.   Hao Z, Lv D, Ge Y, Shi J, Weijers D, Yu G, et al. RIdeogram: drawing SVG graphics

to visualize and map genome-wide data on the idiograms. PeerJ Computer Science.

2020;6:e251.

94.   Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative

genomics. Genome biology. 2019;20(1):1-14.

95.   Price MN, Dehal PS, Arkin AP. FastTree 2–approximately maximum-likelihood trees

for large alignments. PloS one. 2010;5(3):e9490.

96.   Ranwez V, Douzery EJ, Cambon C, Chantret N, Delsuc F. MACSE v2: toolkit for the

alignment of coding sequences accounting for frameshifts and stop codons. Molecular biology

and evolution. 2018;35(10):2582-4.

97.   Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. Bioinformatics. 2009;25(15):1972-3.

98.   Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu C-H, Xie D, et al. BEAST 2: a

software platform for Bayesian evolutionary analysis. PLoS computational biology.

2014;10(4):e1003537.

99.   Bouckaert RR, Drummond AJ. bModelTest: Bayesian phylogenetic site model

averaging and model comparison. BMC evolutionary biology. 2017;17:1-11.

100.   Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. Posterior summarization in

Bayesian phylogenetics using Tracer 1.7. Systematic biology. 2018;67(5):901-4.

101.   FigTree 2016 [Available from: http://tree.bio.ed.ac.uk/software/figtree/.

102.   Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von Haeseler A,

et al. IQ-TREE 2: new models and efficient methods for phylogenetic inference in the

genomic era. Molecular biology and evolution. 2020;37(5):1530-4.

103.   Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder:

fast model selection for accurate phylogenetic estimates. Nature Methods. 2017;14(6):587-9.

104.   HURVICH CM, TSAI C-L. Regression and time series model selection in small

samples. Biometrika. 1989;76(2):297-307.

105.   Minh BQ, Nguyen MAT, Von Haeseler A. Ultrafast approximation for phylogenetic

bootstrap. Molecular biology and evolution. 2013;30(5):1188-95.

106.   Dudchenko O, Batra SS, Omer AD, Nyquist SK, Hoeger M, Durand NC, et al. *De*

novo assembly of the *Aedes aegypti* genome using Hi-C yields chromosome-length scaffolds.

Science. 2017;356(6333):92-5.

107.   Yow AG, Zhang Y, Bansal K, Eacker SM, Sullivan S, Liachko I, et al. Genome

sequence of *Monilinia vaccinii-corymbosi* sheds light on mummy berry disease infection of

blueberry and mating type. G3. 2021;11(2):jkaa052.

108.   Schultz DT, Haddock SH, Bredeson JV, Green RE, Simakov O, Rokhsar DS. Ancient

gene linkages support ctenophores as sister to other animals. Nature. 2023:1-8.

109.   Lechner M, Findeiß S, Steiner L, Marz M, Stadler PF, Prohaska SJ. Proteinortho:

detection of (co-) orthologs in large-scale analysis. BMC bioinformatics. 2011;12(1):1-9.

110.   Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Research. 2004;32(5):1792-7.

111.   Yang Z. PAML 4: phylogenetic analysis by maximum likelihood. Molecular biology

and evolution. 2007;24(8):1586-91.

112.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic

features. Bioinformatics. 2010;26(6):841-2.

113.   Gel B, Serra E. karyoploteR: an R/Bioconductor package to plot customizable

genomes displaying arbitrary data. Bioinformatics. 2017;33(19):3088-90.

114.   Hunter JD. Matplotlib: A 2D graphics environment. Computing in science &

engineering. 2007;9(03):90-5.

115.   Thomas Hackl MJA. gggenomes: A Grammar of Graphics for Comparative Genomics

2022 [Available from: https://github.com/thackl/gggenomes.

116.   Flutre T, Duprat E, Feuillet C, Quesneville H. Considering transposable element

diversification in *de novo* annotation approaches. PloS one. 2011;6(1):e16526.

117.   Quesneville H, Bergman CM, Andrieu O, Autard D, Nouaud D, Ashburner M, et al.

Combined evidence annotation of transposable elements in genome sequences. PLoS Comput

Biol. 2005;1(2):e22.

118.   Bao W, Kojima KK, Kohany O. Repbase Update, a database of repetitive elements in

eukaryotic genomes. Mobile Dna. 2015;6(1):11.

119.   Smit A, Hubley R, Green P. RepeatMasker Open-4.0. 2013–2015. 2015.

120.   Gattiker A, Gasteiger E, Bairoch A. ScanProsite: a reference implementation of a

PROSITE scanning tool. Applied bioinformatics. 2002;1(2):107-8.

121.   Sigrist CJ, De Castro E, Cerutti L, Cuche BA, Hulo N, Bridge A, et al. New and

continuing developments at PROSITE. Nucleic acids research. 2012;41(D1):D344-D7.

122.   Consortium U. UniProt: a worldwide hub of protein knowledge. Nucleic acids

research. 2019;47(D1):D506-D15.

123.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics. 2014;30(15):2114-20.

124.   Andrews FK, Anne Segonds-Pichon, Laura Biggins, Christel Krueger, Jo

Montgomery. FastQC: a quality control tool for high throughput sequence data 2010

[Available from: https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

125.   Picard toolkit: Broad Institute; 2019 [Available from:

https://github.com/broadinstitute/picard.

126.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The sequence

alignment/map format and SAMtools. Bioinformatics. 2009;25(16):2078-9.

127.   Thorvaldsdóttir H, Robinson JT, Mesirov JP. Integrative Genomics Viewer (IGV):

high-performance genomics data visualization and exploration. Briefings in bioinformatics.

2013;14(2):178-92.

128.   Okonechnikov K, Conesa A, García-Alcalde F. Qualimap 2: advanced multi-sample

quality control for high-throughput sequencing data. Bioinformatics. 2016;32(2):292-4.

129.   Tange O. GNU parallel 2018: Lulu. com; 2018.

130.   Garrison E, Marth G. Haplotype-based variant detection from short-read sequencing.

arXiv preprint arXiv:12073907. 2012.

131.   McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GR, Thormann A, et al. The ensembl

variant effect predictor. Genome biology. 2016;17(1):1-14.

132.	Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al.

SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J

Comput Biol. 2012;19(5):455-77.

133.	Mapleson D, Garcia Accinelli G, Kettleborough G, Wright J, Clavijo BJ. KAT: a K-

mer analysis toolkit to quality control NGS datasets and genome assemblies. Bioinformatics.

2017;33(4):574-6.

134.	Pertea G, Pertea M. GFF utilities: GffRead and GffCompare. F1000Research. 2020;9.

135.	Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq

quantification. Nature biotechnology. 2016;34(5):525-7.

136.	Jon Palmer JS. nextgenusfs/funannotate: funannotate v1.5.3 Zenodo2019 [Available

from: https://doi.org/10.5281/zenodo.2604804.

137.	Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, et al. InterProScan 5:

genome-scale protein function classification. Bioinformatics. 2014;30(9):1236-40.

138.	Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. Bioinformatics.

2009;26(1):139-40.

139.	Soneson C, Love MI, Robinson MD. Differential analyses for RNA-seq: transcript-

level estimates improve gene-level inferences. F1000Research. 2015;4.

140.	Wickham H. ggplot2: elegant graphics for data analysis: springer; 2016.

1247

Tables

Table 1. Cereal Rust Fungi Genomes used in the present study.

The table provides information on all genome assemblies used in this study, their total dikaryotic genome assembly size ("Genome size"), and completeness as assessed with

BUSCO ("BUSCO (%)"). For the "BUSCO (%)" column the first number presents the

percentage of complete single BUSCO genes identified. The numbers in the brackets provide

the percentage of single copy and of duplicated BUSCO genes in the dikaryotic genome

assembly. Additional presented metadata includes species name ("Species"), strain name

("Strain"), source ("Source"), download date in DD/MM/YYYY ("Download date") and the

initial reference ("Citation"). The [*] genome denotes dikaryotic genome assemblies which are

chromosome scale phased genome. NCBI – National Center for Biotechnology information,

JGI – Joint Genome Institute, CSIRO – Commonwealth Scientific and Industrial Research

Organization.

Species	Strain	Genome size	BUSCO (%)	Source	Download date	Citation
Puccinia graminis f. sp. *tritici*	*Ug99*	176.2 Mb	93.4 [S:7.7, D:85.7]	NCBI	26/05/2021	Li et al. 2019
	21-0[*]	176.9 Mb	94.2 [S:8.3, D:85.9]	NCBI	26/05/2021	
Puccinia striiformis f. sp. *tritici*	*DK0911*	157.8 Mb	91.1 [S:38.4, D:52.7]	JGI	1/06/2021	Schwessinger et al. 2020
	104E	126.5 Mb	92.1 [S:7.9, D:84.2]	JGI	26/05/2021	Schwessinger et al. 2018
	134E[*]	167.7 Mb	92.6 [S:6.8, D:85.8]	Provided by the authors	21/06/2021	Schwessinger et al. 2022
Puccinia coronata f. sp. *avenae*	*12NC29*	105.2 Mb	89.9 [S:18.1, D:71.8]	NCBI	26/05/2021	Miller et al. 2018
	12SD80	99.2 Mb	90.2 [S:33.1, D:57.1]	NCBI	26/05/2021	Miller et al. 2018
	203[*]	208.1 Mb	92.1 [S:6.9, D:85.2]	CSIRO data access portal	20/06/2022	Henningsen et al. 2022
Puccinia triticina	*76[*]*	253.5 Mb	91.0 [S:1.7, D:89.3]	Provided by the authors	28/05/2021	Duan et al. 2022
	15[*]	243.9 Mb	91.1 [S:1.6, D:89.5]	NCBI	02/11/2023	Li et al. 2022
	19NSW04[*]	253 Mb	91.2[S:0.9, D:90.3]	NCBI	10/10/2023	Sperschneider et al. 2023
	20QLD87[*]	248.1 Mb	91.3[S:1.3, D:90.0]	NCBI	10/10/2023	Sperschneider et al. 2023
Puccinia polysora f. sp. *zea*	*GD1913[*]*	1.7 Gb	91.1 [S:1.4, D:89.7]	NCBI	07/14/2023	Liang et al. 2023

1261

Table 2. MAT reference genes used as query and outgroups.

The table provides gene names ("Gene name") and gene identifier ("Gene ID") of *MAT* genes

from *Puccinia triticina* initially used as query sequences and from *Puccinia polysora* f. sp.

zeae used as outgroups. Additional presented metadata includes species name ("Species"),

strain name ("Strain"), source ("Source"), and the initial reference ("Citation"). NCBI –

National Center for Biotechnology information.

Species	Strain	Source	Gene name	Gene ID	Citation
Puccinia graminis f. sp. *tritici*	isolate *CDL 75–36-700–3* (race *SCCL*)	NCBI & Cuomo et al., 2017	*bW-HD1*	PGTG_05143	Cuomo et al. 2017
			bE-HD2	PGTG_05144	
			STE3.2-1	PGTG_00333	
			STE3.2-2	PGTG_19559	
			STE3.2-3	PGTG_01392	
			Pgtmfa1	-	
			Pgtmfa2	-	
			Pgtmfa3	-	
Puccinia triticina	isolate *1-1 / race 1 (BBBD)*	NCBI & Cuomo et al., 2017	*bW-HD1*	PTTG_09683 PTTG_27730	Cuomo et al. 2017
			bE-HD2	PTTG_10928 PTTG_03697	
			STE3.2-1	PTTG_09751	
			STE3.2-2	PTTG_28830	
			STE3.2-3	PTTG_09693	
			Ptmfa1/3	-	
			Ptmfa2	-	
Puccinia striiformis f. sp. *tritici*	*78*	NCBI & Cuomo et al., 2017	*bW-HD1*	PSTG_05919 PSTG_18670	Cuomo et al. 2017
			bE-HD2	PSTG_05918 PSTG_19315	
			STE3.2-1	PSTG_02613	
			STE3.2-2	PSTG_15127	
			STE3.2-3	PSTG_15070	
			Pstmfa1	-	
			Pstmfa2	-	
Puccinia polysora f. sp. *zea*	*GD1913*	NCBI	*bW-HD1*	FUNB_006397	Liang et al. 2023
			bE-HD2	FUNA_006226	

			STE3.2-2	FUNA_023003	
			STE3.2-1	FUNA_001167	
			STE3.2-3	FUNA_013409	

1268

Figure captions

Fig 1. *HD* **genealogies suggest distinct evolutionary histories for** *bW-HD1* **and** *bE-HD2* **in four cereal rust fungi.**

(A) Species tree of *Puccinia polysora* f. sp. *zeae* (*Ppz*), *P. coronata* f. sp. *avenae* (*Pca*), *P. graminis* f. sp. *tritici* (*Pgt*), *P. triticina* (*Pt*) and *P. striiformis* f. sp. *tritici* (*Pst*) inferred from 2284 single-copy orthogroups. Numbers on nodes represent local support value computed by Fasttree. *Puccinia polysora* f. sp. *zeae* (*Ppz)* has been used for rooting the species tree. (B) and (C) Bayesian rooted gene tree built from *bW-HD1* or *bE-HD2* coding-based sequence alignment, respectively. Trees are based on a HKY + G model of molecular evolution. In either case, alleles from the same species are grouped into the same clade. Each node is labelled with its values of posterior probability (PP). PP values above 0.95 are considered to have strong evidence for monophyly of a clade and PP values of identical alleles are not displayed. The scale bar represents the number of nucleotide substitutions per site. * marks an allele with minor variations only outside the variable domain, which means this allele is predicted to be functionally equivalent to its closest neighbor. Alleles of the same species are colored with identical background: *Pca* (yellow), *Pgt* (green), *Pt* (blue), *Pst* (orange).

Fig 2. *Pra* **genealogy displays ancient trans-specific polymorphism in cereal rust fungi.**

Bayesian gene tree built from *Pra (STE3.3-2 and STE3.3-3)* coding-based sequence alignment. The tree is based on a TN93 + I model of molecular evolution. The *Pra* alleles, *STE3.2-2* and *STE3.2-3,* were grouped into two clades by allele identity and not species identity. Each node is labelled with its values of posterior probability (PP). PP values above 0.95 are considered

as strong evidence for monophyly of a clade and PP values of identical alleles are not

displayed. The scale bar represents the number of nucleotide substitutions per site. Species-

specific background coloring is the same as for Fig 1.

Fig 3. Synonymous divergence (d_S) values of *HD* genes but not their immediate

neighbors are slightly elevated on chromosome 4.

Synonymous divergence values (d_S) for all allele pairs are plotted along chromosome 4A for

(A) *P. coronara* f. sp. *avenae* ("*Pca* 203")*,* (B) *P. graminis* f. sp. *tritici* ("*Pgt* 210")*,* (C) *P.*

triticina ("*Pt* 76"), and (D) *P. striiformis* f. sp. *tritici* ("*Pst* 134E"). In each panel, the top track

shows the d_S values ("d_S") of allele pairs along chromosome 4. Each dot corresponds to the d_S

value of a single allele pair. The second and third track shows the averaged TE ("TE") and

gene ("gene") density along chromosome 4 in 10 kbp-sized windows, respectively. The *HD*

genes (*bW-HD1* and *bE-HD2*) are highlighted with a red line and red shading indicates a 0.4

mbp-sized window around the *HD* locus. Predicted centromeric regions are marked with blue

shading. The two lower tracks (d_S values and gene locations) provide a detailed zoomed in

view of red shaded area around the *HD* locus. Species-specific background coloring is the

same as for Fig 1.

Fig 4. *HD* loci are partially conserved within species whereas *PR* loci are highly

heterozygous and display strong signals of genomic degeneration.

Synteny graphs of *HD* locus (A) and (B) *PR* locus including proximal regions in *P. coronara*

f. sp. *avenae* ("*Pca* 203")*, P. graminis* f. sp. *tritici* ("*Pgt* 210")*,* (C) *P. triticina* ("*Pt* 76"), and

(D) *P. striiformis* f. sp. *tritici* ("*Pst* 134E"). Proximal regions are defined as 40 genes

downstream and upstream of *HD* and *Pra* genes, respectively. *HD* loci are relatively syntenic

within each dikaryotic genome but less conserved between species. *PR* loci display little

synteny in *Pgt 21-0, Pt 76* and *Pst 134E* and show strong signs of transposon accumulation

(see also S13 Fig). There is very little conservation of the *PR* loci across species. Red lines

between chromosome sections represent gene pairs with identity higher than 70% and grey

shades represent conserved nucleotide sequences (>=1000 bp and identity >=90%). For

additional annotations please refer to the included legend ("Legend").

Fig 5. *PR* loci display elevated synonymous divergence (d$_S$) values, accumulation of

transposable elements and depletion of genes.

Synonymous divergence values (d$_S$) for all allele pairs are plotted along chromosome 9A for

(A) *P. coronara* f. sp. *avenae* ("*Pca* 203"), (B) *P. graminis* f. sp. *tritici* ("*Pgt* 210"), (C) *P.*

triticina ("*Pt* 76"), and (D) *P. striiformis* f. sp. *tritici* ("*Pst* 134E"). In each panel, the top track

shows the d$_S$ values ("d$_S$") of allele pairs along chromosome 9. Each dot corresponds to the d$_S$

value of a single allele pair. The second and third track shows the averaged TE ("TE") and

gene ("gene") density along chromosome 9 in 10 kbp-sized windows, respectively. The *Pra*

alleles (*STE3.2-2* and *STE3.2-3*) are highlighted with a red line and red shading indicates a 0.4

mbp-sized window around the *PR* locus. Predicted centromeric regions are marked with blue

shading. The two lower tracks (d$_S$ values and gene locations) provide a detailed zoomed in

view of red shaded area around the *PR* locus. Species-specific background coloring is the

same as for Fig 1.

Fig 6. The *PR* locus shows an *STE3.2-3* allele specific accumulation of a unique Ty3-

transposon family in *Puccinia triticina*

Synteny graphs of (A) *STE3.2-2* and (B) *STE3.2-3* including proximal regions in four

different *P. triticina* isolates *Pt 15*, *Pt 19NSW04*, *Pt 20QLD87,* and *Pt 76*. Both loci are

overall syntenic while the *STE3.2-3* locus displays an extension of the intergenic distance

especially between the pheromone receptor and the two *mfa* genes. Red lines between

chromosome sections represent gene pairs with identity higher than 70% and grey shades

represent conserved nucleotide sequences (>=1000 bp and identity >=90%). For additional annotations please refer to the included legend ("Legend"). (C) Transposable element coverage at the order level at the *STE3.2-3* locus in the four different *P. triticina* isolates. (D) Structure of a representative Ty3_Pt_*STE3.2-3* copy. LTR – Long Terminal Repeat, RT – Reverse transcriptase, RH – RNAse H, and INT – Integrase. Distributions of (E) percentage identity relative to the consensus sequence and (F) lengths of individual Ty3_Pt_*STE3.2-3* insertions at the *STE3.2-3* locus in four different *P. triticina* isolates.

Fig 7. *MAT* gene expression is upregulated in the late stage of asexual infection cycle of the cereal host.

(A) Trimmed mean of M-values (TMM)-normalized values of *MAT* genes in *P. coronara* f. sp. *avenae* ("*Pca 12NC29*") at 48 and 120 hour post infection (hpi). (B) TMM-normalized values of *MAT* genes in *P. graminis* f. sp. *tritici* ("*Pgt 21-0*") at 48, 72, 96, 120, 144 and 168 hpi. (C) TMM-normalized values of *MAT* genes in *P. striiformis* f. sp. *tritici* ("*Pst 87/66*") at 24, 48, 72, 120, 168, 216 and 264 hpi.

Supporting information captions

Table S1 List of whole genome Sequence Read Archive data used in the present study.

The table provides information on all whole genome Sequence Read Archive (SRA) data used in the present study. Metadata includes species abbreviation ("Species"), isolate name ("Isolate"), SRA identifier ("SRA ID"), and the initial reference ("Citation"). *Pca - P. coronata* f. sp. *avenae*, *Pgt - Puccinia graminis* f. sp. *tritici*, *Pt - P. triticina* and *Pst - P. striiformis* f. sp. *tritici*.

Table S2 List of RNAseq Sequence Read Archive data used in the present study.

The table provides information on all RNAseq Sequence Read Archive (SRA) data used in

the present study. Metadata includes species abbreviation ("Species"), isolate name

("Isolate"), timepoint of infection (hpi, hours post infection) or sample type ("Sample Type"),

timepoint/status of spores ("Time point/Status"), replicate number ("Replicate"), SRA

identifier ("SRA ID"), and the initial reference ("Citation"). *Pca - P. coronata* f. sp. *avenae*,

Pgt - Puccinia graminis f. sp. *tritici*, and *Pst - P. striiformis* f. sp. *tritici.*

S1 Fig. Two unlinked *MAT* loci suggest tetrapolar mating types in four *Puccinia spp.*

Karyograms of *P. coronata* f. sp. *avenae* ("*Pca 203*"), *P. graminis* f. sp. *tritici* ("*Pgt 21-0*"),

P. triticina ("*Pt 76*") and *P. striiformis* f. sp. *tritici* ("*Pst 134 E*") with the positions of *HD,*

PR and *STE3.2-1* loci marked by black arrow heads. *HD*, *PR* and *STE3.2-1,* located on

chromosome 4, chromosome 9 and chromosome 1, respectively, suggest tetrapolar mating

types in these four species.

S2 Fig. Genetic distance between pheromone precursor genes (*mfa*) and pheromone

receptor (*Pra*) genes varies from 100 bp to 100 kb. The diagrams display the location of

STE3.2-2 and *STE3.2-3* and their linked *mfa* genes on chromosome 9A and 9B in *P. coronata*

f. sp. *avenae* ("*Pca 203*"), *P. graminis* f. sp. *tritici* ("*Pgt 21-0*"), *P. triticina* ("*Pt 76*") and *P.*

striiformis f. sp. *tritici* ("*Pst 134 E*"). The grey bar represents chromosome subsections

containing the genes of interest with the numbers indicating the absolute location in mega

base pairs on chromosome 9A and 9B. The genetic distance between *mfa1/mfa3* and *STE3.2-3*

are highly variable between species, whereas *STE3.2-2* and *mfa2* are tightly linked in all

species.

S3 Fig. MFA protein alignments indicate mating factor a proteins are overall conserved

within species. (A) Alignment of MFA1 protein sequences from four different rust fungal

species. The MFA1, which is linked to *STE3.2-3,* is mostly conserved within species. Two

near identical MFA1/3 copies can be identified in four *P. triticina* isolates. In *P. graminis* f.

sp. *tritici* MFA1 has four amino acid substitutions in *Pgt SCCL* versus *Pgt 21-0*. (B)

Alignment of MFA2 protein sequences from four different rust fungal species. MFA2 is fully

conserved within each species. (C) Alignment of MFA3 protein sequences from three *P.*

graminis f. sp. *tritici* isolates. *Mfa3* is only present *P. graminis* f. sp. *tritici* in close proximity

to *STE3.2-3*.

S4 Fig. Pheromone peptide mating factor a (MFA) proteins group according to their

proximate *Pra* alleles and display strong signals of trans-specific polymorphisms.

Maximum likelihood tree of MFA identified in four cereal rust fungi including multiple

isolates per species. Tips are labelled with the species abbreviation, protein names, and isolate

names are provided in parentheses. Branch support was assessed by 1000 replicates. The scale

bar represents 0.4 substitutions per site. *Pca - P. coronata* f. sp. *avenae*, *Pgt - Puccinia*

graminis f. sp. *tritici*, *Pt - P. triticina* and *Pst - P. striiformis* f. sp. *tritici*.

S5 Fig. Genealogy of *STE3.2-1* in rust fungi indicates sequence of *STE3.2-1* are

conserved within species. Bayesian rooted gene tree built from *STE3.2-1* coding-based

sequence alignment from four cereal rust fungi: *P. coronata* f. sp. *avenae* (*Pca*), *P. graminis* f.

sp. *tritici* (*Pgt*), *P. triticina* (*Pt*) and *P. striiformis* f. sp. *tritici* (*Pst*) and *P. polysora* f. sp. *zeae*

(*Ppz*) *GD1913* was included as outgroup. Trees are based on a TN93 + I model of molecular

evolution. Each node is labelled with its values of posterior probability (PP). PP values above

0.95 are considered to have strong evidence for monophyly of a clade and PP values of

identical alleles are not displayed. The scale bar represents the number of nucleotide

substitutions per site. Alleles of the same species are colored with identical background: *Pca*

(yellow), *Pgt* (green), *Pt* (blue), *Pst* (orange).

S6 Fig. Whole chromosome alignments of *HD* loci containing chromosomes between the two haplotypes of each dikaryotic genome exhibit slightly reduced synteny around *HD* locus. The figure shows dots plots of whole chromosome alignments between the two *HD* loci containing chromosomes from dikaryotic genome assemblies. Each panel consists of dot plots of the whole chromosome and subset dot plot zooming into the *HD* locus. The *HD* locus is labelled and line colors show the nucleotide percentage identity and nucleotide orientation as indicated in the figure legend. Subfigure A to D show *P. coronata* f. sp. *avenae* ("*Pca 203*"), *P. graminis* f. sp. *tritici* ("*Pgt 21-0*"), *P. triticina* ("*Pt 76*") and *P. striiformis* f. sp. *tritici* ("*Pst 134 E*"), respectively.

S7 Fig. Allele pairs on chromosomes 4 and 9 are not overall more diverged when compared to other chromosomes. The plots show the distribution of d_S values of allele pairs on sister chromosomes in four cereal rust fungal species. Bar plots show the distribution of d_S values up to the 99% quantile. Red lines represent threshold of d_S values of 95% of all alleles. Blue lines represent threshold of d_S values of 90% of all alleles. Black points show d_S values of individual allele pairs. *HD* and *STE3* allele pairs are highlighted as red points. Subfigures A to D show *P. coronata* f. sp. *avenae* ("*Pca 203*"), *P. graminis* f. sp. *tritici* ("*Pgt 21-0*"), *P. triticina* ("*Pt 76*") and *P. striiformis* f. sp. *tritici* ("*Pst 134 E*"), respectively.

S8 Fig. Hi-C heatmap of *P. coronata* f. sp. *avenae* ("*Pca 203*") hapA.

S9 Fig. Hi-C heatmap of *P. coronata* f. sp. *avenae* ("*Pca 203*") hapB.

S10 Fig. Hi-C heatmap of *P. triticina* ("*Pt 76*") hapA.

S11 Fig. Hi-C heatmap of *P. triticina* ("*Pt 76*") hapB.

S12 Fig. Coverage of different transposable element orders at the *HD*, *PR*, and *STE3.2-1* locus. The plots show the percentage of nucleotides covered by different transposable element orders at (A) *HD* locus (B) *PR* locus (C) *STE3.2-1* locus. Each subfigure A to C shows the

coverage in each haplotype of the dikaryotic genomes of *P. coronata* f. sp. *avenae* ("*Pca 203*"), *P. graminis* f. sp. *tritici* ("*Pgt 21-0*"), *P. triticina* ("*Pt 76*") and *P. striiformis* f. sp. *tritici* ("*Pst 134 E*"). Different TE orders are color coded as shown in the legend. TEs with no assigned class are labelled "Undetermined". TEs with no assigned order but belonging to Class I (RNA retrotransposons) or Class II (DNA transposons) are labelled "Undetermined Class I" or "Undetermined Class II", respectively.

S13 Fig. Whole chromosome alignments of *PR* loci containing chromosomes between two haplotypes of each dikaryotic genome exhibit strong signs of synteny loss. The figure shows dot plots of whole chromosome alignments between the two *PR* loci containing chromosomes from the dikaryotic genome assemblies. Each panel consists of dot plots of the whole chromosome and subset dot plots zooming into the *PR* locus. The *PR* locus is labelled and line colors show the nucleotide percentage identity and nucleotide orientation as indicated in the figure legend. Subfigures A to D show *P. coronata* f. sp. *avenae* ("*Pca 203*"), *P. graminis* f. sp. *tritici* ("*Pgt 21-0*"), *P. triticina* ("*Pt 76*") and *P. striiformis* f. sp. *tritici* ("*Pst 134 E*"), respectively.

S14 Fig. Whole chromosome alignments of *STE3.2-1* loci containing chromosomes between two haplotypes of each dikaryotic genome suggest high conservation of *STE3.2-1*. The figure shows dot plots of whole chromosome alignments between the two *STE3.2-1* loci containing chromosomes from dikaryotic genome assemblies. Each panel consists of dot plots of the whole chromosome and subset dot plots zooming into the *STE3.2-1* locus. The *STE3.2-1* locus is labelled and line colors show the nucleotide percentage identity and nucleotide orientation as indicated in the figure legend. Subfigures A to D show *P. coronata* f. sp. *avenae* ("*Pca 203*"), *P. graminis* f. sp. *tritici* ("*Pgt 21-0*"), *P. triticina* ("*Pt 76*") and *P. striiformis* f. sp. *tritici* ("*Pst 134E*"), respectively.

S15 Fig. Synteny analysis of *STE3.2-1* loci and their flanking regions reveal no evidence

of genetic degeneration. Synteny graphs of the *STE3.2-1* locus including proximal regions in

(A) *P. coronara* f. sp. *avenae* ("*Pca 203*"), *P. graminis* f. sp. *tritici* ("*Pgt 21-0*"), *P. triticina*

("*Pt 76*"), and *P. striiformis* f. sp. *tritici* ("*Pst 134E*"). Proximal regions are defined as 40

genes downstream and upstream of the *STE3.2-1* alleles, respectively. *STE3.2-1* loci are

highly syntenic within each dikaryotic genome and conserved between species. Red lines

between chromosome sections represent gene pairs with sequence identity higher than 70%

and grey shades represent conserved nucleotide sequences (>=1000 bp and identity >=90%).

For additional annotations please refer to the provided legend ("Legend").

S16 Fig. Synonymous divergence (d_S) values between *STE3.2-1* alleles along

chromosome 1A suggest conservation of this locus. Synonymous divergence values (d_S) for

all allele pairs are plotted along chromosome 1A for (A) *P. coronara* f. sp. *avenae* ("*Pca*

203"), (B) *P. graminis* f. sp. *tritici* ("*Pgt 21-0*"), (C) *P. triticina* ("*Pt 76*"), and (D) *P.*

striiformis f. sp. *tritici* ("*Pst 134E*"). In each panel, the top track shows the d_S values ("$d_S^{"}$") of

allele pairs along chromosome 1. Each dot corresponds to the d_S value of a single allele pair.

The second and third track show the averaged TE ("TE") and gene ("gene") density along

chromosome 1 in 10 kbp-sized windows, respectively. The *STE3.2-1* alleles are highlighted

with a red line and red shading indicates a 0.4 mbp-sized window around the *STE3.2-1* locus.

Predicted centromeric regions are marked with blue shading. The two lower tracks (d_s values

and gene locations) provide a detailed zoomed in view of red shaded area around the *STE3.2-*

1 locus. Species-specific background coloring is the same as for Fig 1.

S17 Fig. IGV screen shots of short read Illumina mapping of various *P. coronata* f. sp.

***avenae* isolates against the *HD* locus in the *Pca 203* reference.**

(A) shows mapping against the *HD* locus on chromosome 4A and (B) chromosome 4B,

respectively.

S18 Fig. IGV screen shot of short read Illumina mapping of various *P. graminis* f. sp.

***tritici* isolates against the *HD* locus in the *Pgt 21-0* reference.**

(A) shows mapping against the *HD* locus on chromosome 4A and (B) chromosome 4B,

respectively.

S19 Fig. IGV screen shots of short read Illumina mapping of various *P. triticina* against

the *HD* isolates locus in the *Pt 76* reference.

(A) shows mapping against the *HD* locus on chromosome 4A and (B) chromosome 4B,

respectively.

S20 Fig. IGV screen shots of short read Illumina mapping of various *P. striiformis* f. sp.

***tritici* isolates against the *HD* locus in the *Pst 134E* reference.**

(A) shows mapping against the *HD* locus on chromosome 4A and (B) chromosome 4B,

respectively.

S21 Fig. Nucleotide alignments of the *HD* gene coding sequence in four cereal rust fungi

indicate *bW-HD1* and *bE-HD2* are multiallelic in each species. (A) Nucleotide alignment

of the *de novo* reconstructed *HD* locus from *Pca 203* Illumina short read data with the coding

sequences of *bW-HD1* and *bE-HD2* alleles from *Pca 203* dikaryotic reference genome. (B) to

(D) multiple sequence alignments of *de novo* reconstructed *HD* gene coding sequences, (E)

multiple sequence alignments of *Pst bW-HD1* and *bE-HD2* alleles (21). In each subfigure (B)

to (E), the top two track shows the consensus sequence length and relative sequence identity,

respectively. Subfigure (B) to (E) show *P. coronata* f. sp. *avenae* ("*Pca*"), *P. graminis* f. sp.

tritici ("*Pgt*"), *P. triticina* ("*Pt*") and *P. striiformis* f. sp. *tritici* ("*Pst*"), respectively. The *bW-*

HD1 and *bE-HD2* are numbered in accordance with Fig 1.

S22 Fig. IGV screen shots of short read Illumina mapping of various *P. coronata* f. sp.

***avenae* isolates against the *PR* locus in the *Pca 203* reference.**

(A) shows mapping against the *PR* locus on chromosome 9A and (B) chromosome 9B,

respectively.

S23 Fig. IGV screen shot of short read Illumina mapping of various *P. graminis* f. sp.

***tritici* isolates against the *PR* locus in the *Pgt 21-0* reference.**

(A) shows mapping against the *PR* locus on chromosome 9A and (B) chromosome 9B,

respectively.

S24 Fig. IGV screen shots of short read Illumina mapping of various *P. triticina* isolates

against the *PR* locus in the *Pt 76* reference.

(A) shows mapping against the *PR* locus on chromosome 9A and (B) chromosome 9B,

respectively.

S25 Fig. IGV screen shots of short read Illumina mapping of various *P. striiformis* f. sp.

***tritici* isolates against the *PR* locus in the *Pst 134E* reference.**

(A) shows mapping against the *PR* locus on chromosome 9A and (B) chromosome 9B,

respectively.

S26 Fig. Amino acid alignments of STE3.2-2 and STE3.2-3 alleles of four cereal rust

fungi. Multiple sequence alignment of *de novo* reconstructed STE3.2-2 and STE3.2-3 protein

sequences. Subfigure contains MSAs; one for STE3.2-2 and one for STE3.2-3. Subfigure A to

D show *P. coronata* f. sp. *avenae* ("*Pca*"), *P. graminis* f. sp. *tritici* ("*Pgt*"), *P. triticina* ("*Pt*")

and *P. striiformis* f. sp. *tritici* ("*Pst*"), respectively.

S27 Fig. Amino acid alignments of the three MFA alleles from various *P. graminis* f. sp.

***tritici* isolates.** Amino acid substitutions are highlighted by color.

S28 Fig. Whole chromosome alignments of *HD* genes containing chromosomes between

haplotypes of four different *P. triticina* isolates. The figure shows dots plots of whole

chromosome alignments of *HD* loci containing chromosomes derived from distinct dikaryotic

genomes of four different *P. triticina* isolates including *Pt 15*, *Pt 19NSW04*, *Pt 20QLD87*

against *Pt 76*. Each panel consists of a dot plot of the whole chromosome and a subset dot

plot zooming into the *HD* locus. The *HD* locus is labelled and line colors show the nucleotide

percentage identity and nucleotide orientation as indicated in the figure legend. (A)

Comparison of nucleotide sequence of chromosome 4s of *Pt 15* and *Pt 76*. (B) Comparison of

nucleotide sequence of chromosome 4s of *Pt 19NSW04* and *Pt 76*. (C) Comparison of

nucleotide sequence of chromosome 4s of *Pt 20QLD87* and *Pt 76*. (D) Comparison of

nucleotide sequence of chromosome 4s of *Pt 19NSW04* and *Pt 20QLD87*.

S29 Fig. The *HD* locus is highly conserved in four *P. triticina* isolates. Synteny graphs of

HD loci including proximal regions in the four *P. triticina* isolates *Pt 15*, *Pt 19NSW04*, *Pt*

20QLD87 and *Pt 76*. Red lines between chromosome sections represent gene pairs with

nucleotide sequence identity higher than 70% and grey shades between conserved nucleotide

sequences (>=1000 bp and identity >=90%). For additional annotations please refer to the

provide legend ("Legend").

S30 Fig. Coverage of different transposable element orders at the *HD* and *STE3.2-2*

locus in four *P. triticina* isolates. The plots show the percentage of nucleotides covered by

different transposable element orders at the (A) *HD* locus and (B) *STE3.2-2* locus. Each

subfigure A and B shows the coverage in each haplotype of the dikaryotic genomes of *P.*

triticina isolates *Pt 15*, *Pt 19NSW04*, *Pt 20QLD87* and *Pt 76*. Different TE orders are color

coded as shown in the legend. TEs with no assigned class are labelled "Undetermined". TEs

with no assigned order but belonging to Class I (RNA retrotransposons) or Class II (DNA

transposons) are labelled "Undetermined Class I" or "Undetermined Class II", respectively.

S31 Fig. Whole chromosome alignments of *Pra* gene containing chromosomes between

haplotypes of four different *P. triticina* isolates. The figure shows dot plots of whole

chromosome alignments of *STE3.2-2* or *STE3.2-3* containing chromosomes derived from

distinct dikaryotic genomes of four different *P. triticina* isolates including *Pt 15*, *Pt 19NSW04*,

Pt 20QLD87 against *Pt 76*. Each panel consists of a dot plot of the whole chromosome and a

subset dot plot zooming into the *STE3.2-2* or *STE3.2-3* locus. The *STE3.2-2* or *STE3.2-* locus

is labelled and line colors show the nucleotide percentage identity and nucleotide orientation

as indicated in the figure legend. (A) Comparison of nucleotide sequence of chromosome 9s

containing *STE3.2-2* loci (B) Comparison of nucleotide sequence of chromosome 9s

containing *STE3.2-3*.

S32. Nucleotide coverage distribution of Ty3_Pt_*STE3.2-3* on the chromosomes of four

***P. triticina* isolates**

The plots show the percentage of nucleotides covered by the transposable element family

Ty3_Pt_*STE3.2-3* on each chromosome of four *P. triticina* isolates *Pt 15* (A), *Pt 19NSW04*

(B), *Pt 20QLD87* (C) and *Pt 76* (D). Chromosomes carrying *STE3.2-3* are highlighted in red.

S33. Schematic illustration of the location of Ty3_Pt_*STE3.2-3* copies at the *STE3.2-3*

locus in four *P. triticina* isolates.

S34 Fig. Multidimensional scaling (MDS) plot of RNA dataset used in this study. MDS

plots were made with TMM normalized counts for quality control, each dot represents a

single sample, and replicates were color coded as indicated in each subfigure ledgend. (A)

MDS plot of TMM-normalized value of *Pca* at 48 hour post infection (hpi) and 120 hpi. (B)

MDS plot of TMM-normalized value of *Pgt* at 48, 72, 96, 120, 148 and 168 hpi. (C) MDS

plot of TMM-normalized value of *Pst* at 24, 48, 72, 120, 168, 216 and 264 hpi. (D) MDS plot

of TMM-normalized value of ungerminated spore (US), germinated spores (GS) stages, 144

hpi, 216 hpi and haustoria enriched samples (HE) of *Pst*.

S35 Fig. Expression of housekeeping genes in four RNAseq datasets. (A) TMM-

normalized value of housekeeping genes in *P. coronata* f. sp. *avenae* ("*Pca 12NC29*"), *P.*

graminis f. sp. *tritici* ("*Pgt 21-0*"), and *P. striiformis* f. sp. *tritici* ("*Pst 87/66*" and "*Pst*

104E"). (B) Likelihood ratio test (LRT) method was applied to test significant upregulation of

housekeeping genes between timepoints, none of the housekeeping genes show significant

upregulation between timepoints. red dashed lines indicate logFC=0.5 and logFC=-0.5

respectively, stars above or below bars indicate statistically significant differences between

the two adjacent time points: *$p<0.05$, **$p<0.01$, **$p<0.001$. Genes are labelled with

different colors. Transcription elongation factor TFIIS (*TFIIS),* Actin-related protein 3

(*ARP3),* Actin/actin-like protein 2 (*ACTIN2),* Ubiquitin carboxyl-terminal hydrolase 6

(*UBP6),* Conserved oligomeric Golgi complex subunit 3 (*COG3),* Ctr copper transporter 2

(*CTR2).*

S36 Fig. Upregulation of *MAT* genes in late stages of the asexual life cycle. Likelihood

ratio test (LRT) method was applied to test significant upregulation of *MAT* genes between

timepoints, red dashed lines indicate logFC=0.5 and logFC=-0.5 respectively, stars above or

below bars indicate statistically significant differences between the two adjacent time points:

*$p<0.05$, **$p<0.01$, ***$p<0.001$. *MAT* genes were labelled with different colors. (A) The

expression levels of *MAT* genes in *P. coronata* f. sp. *avenae* ("*Pca 12NC29*") were compared

between 120 hours post infection (hpi) and 48 hpi. (B) The expression levels of *MAT* genes in

P. graminis f. sp. *tritici* ("*Pgt 21-0*") were compared between 72 hpi and 48 hpi, 96 hpi and

72 hpi, 120 hpi and 96 hpi, 144 hpi and 120 hpi, 168 hpi and 144 hpi. (C) The expression

levels of *MAT* genes in *P. striiformis* f. sp. *tritici* ("*Pst 87/66*") were compared between 48

hpi and 24 hpi, 72 hpi and 48 hpi, 120 hpi and 72 hpi, 168 hpi and 120 hpi, 216 hpi and 168

hpi, 264 hpi and 216 hpi. (D) The expression levels of *MAT* genes in *P. striiformis* f. sp. *tritici*

("*Pst 104E*") were compared between germinated spores (GS) and ungerminated spores (US),

144 hpi and GS, 216 hpi and 144 hpi, haustoria enriched samples (HE) and 216 hpi.

S37 Fig. *MAT* genes are upregulated in the late asexual infection stage of *P. striiformis* f.

sp. *tritici* ("*Pst 104E*"). TMM-normalized value of *MAT* genes in ungerminated spores (US),

germinated spores (GS) stages, 144 hpi, 216 hpi and in haustoria enriched samples (HE) of

Pst. Genes are labelled with different colors.

1606

Legend

Symbol	Description
Yellow triangle	*bW-HD1 / bE-HD2*
Purple triangle	*STE3.2-3*
Cyan triangle	*STE3.2-2*
Red triangle	Phermone peptide gene
Tan arrow	Genes
Dark teal bar	Repetitive elements
Red/pink band	Conserved genes (Identity > 70%)
Grey band	Conserved nucleotide sequences