Canonical and ETT-mediated auxin signalling pathways synergistically promote

gynoecium development in *Arabidopsis thaliana*

Heather M. McLaughlin[1,2], Bhavani Natarajan[1], Lars Østergaard[1]

[1] John Innes Centre, Norwich Research Park, Norwich, NR4 7UH, UK,

[2] The Sainsbury Laboratory, The University of Cambridge, 47 Bateman Street, Cambridge,

CB2 1LR, UK

Correspondence

Lars Østergaard (Lars.Ostergaard@jic.ac.uk)

Abstract

Auxin is a phytohormone involved in a wide range of developmental processes. Whilst auxin-

mediated signalling is now known to occur through a range of pathways, the transduction of

auxin signalling within the carpels has mainly been studied in the context of the ETTIN (ETT)

mediated pathway. Several lines of evidence suggest a role for the canonical TIR1/AFB2-

mediated pathway in carpel development. However, the relationship between canonical and

ETT-mediated mechanisms has not been investigated. Here, *tir1-1 afb2-3 ett-3* triple mutants

were generated in *Arabidopsis*, and phenotypic analysis showed synergism between the

canonical and ETT-mediated pathways in gynoecium development. Furthermore, ETT shares

partial redundancy with ARF4, a close paralog which interacts with the canonical pathway

AUX/IAAs. However, the role of ARF4 in the regulation of auxin-responsive gene expression

has not been studied in the gynoecium. Here comparative transcriptomics were utilized to

assess the roles of ETT and ARF4 in gynoecium development, and the mis-regulation of the

auxin response in *ett-3 arf4-2* mutants. The data suggest that the auxin independent mis-

regulation of *YABBY2*, *YABBY5,* and *CRABSCLAW* contributes to the abaxial-adaxial identity

defects observed in gynoecia and leaves of the double mutant. Auxin sensitive transcriptomics

analyses suggest that ETT and ARF4 maintain the auxin insensitivity of both distinct and

overlapping targets, many of which become upregulated by auxin in their absence. This is

consistent with previously proposed models of A-B ARF antagonism, and suggests that the

maintenance of auxin insensitivity at target loci may play an equally important role as the

auxin-sensitive induction of genes for a range of developmental processes.

Introduction

The role of auxin in plant development has been widely studied, and auxin signalling is now known to be transduced by several distinct pathways (Dharmasiri et al., 2005; Kepinski & Leyser, 2005; Simonini et al., 2016; Fendrych et al., 2018; Cao et al., 2019; Kuhn et al., 2020; Friml et al., 2022). The canonical auxin signalling pathway induces the expression of auxin sensitive genes in the nucleus (Paponov et al., 2008), and is composed of three main components: Auxin/Indole-3-Acetic Acid repressor proteins (AUX/IAAs), Transport Inhibitor Response1/Arabidopsis F-box (TIR1/AFB) receptors, and Auxin Response Factors (ARFs). In the absence of auxin, ARFs are bound by AUX/IAAs, which recruit the TOPLESS (TPL) co-repressor, preventing the expression of auxin responsive genes (Szemenyei et al., 2008). Auxin acts as a molecular glue that increases the affinity of the TIR1/AFBs for AUX/IAAs. AUX/IAAs are subsequently degraded by the proteasome, allowing ARFs to promote auxin responsive gene expression (Dharmasiri et al., 2005; Kepinski & Leyser, 2005).

ARFs are classified into three distinct clades: A-ARFs, B-ARFs, and C-ARFs (Ulmasov et al., 1999; Tiwari et al., 2003; Finet et al., 2013). A-ARFs are transcriptional activators which readily interact with AUX/IAAs, whilst B and C ARFs are repressive, and interact with AUX/IAAs less readily (Ulmasov et al., 1999; Tiwari et al., 2003; Finet et al., 2013; Vernoux et al., 2011; reviewed in Guilfoyle & Hagen, 2012).

The ETTIN (ETT/ARF3)-mediated pathway was elucidated more recently in the context of its role in gynoecium development (Simonini et al., 2016; Kuhn et al., 2020). ETT is a B-class repressive ARF which lacks the C-terminal domain required for AUX/IAA interaction (Simonini et al., 2016). ETT recruits a TPL-HISTONE DEACETYLASE 19 (HDA19) complex to prevent the transcription of target loci in the absence of auxin. ETT interacts with auxin through its C-terminal ETTIN-SPECIFIC (ES) domain and undergoes a conformational change which breaks the ETT-TPL-HDA19 complex. Histones then become acetylated, and genes can be expressed once more (Kuhn et al., 2020).

ett mutants exhibit severe gynoecium defects which phenocopy aberrations produced upon treatment with the auxin transport inhibitor N-1-Naphthylphthalamic Acid (NPA), highlighting the importance of ETT-mediated signalling for auxin related processes in carpel development (Nemhauser et al., 2000). Gynoecium morphogenesis is dependent on the dynamic distribution of auxin (Moubayidin and Østergaard, 2014). However, studies on auxin signal transduction in the carpels have mainly focused on the role of the ETT-mediated pathway (Simonini et al., 2016; Simonini et al., 2017; Kuhn et al., 2020). An understanding of the relative contribution of canonical and ETT-mediated signalling to gynoecium development remains lacking, although several lines of evidence implicate the canonical pathway in this process. Four of the canonical auxin receptors - *TIR1*, *AFB1*, *AFB2*, and *AFB3* - are expressed in the style of the developing *Arabidopsis* gynoecium (Dharmasiri et al., 2005; Cecchetti et al., 2008).

Higher order *tir1afb1245* mutants exhibit curled siliques and reduced fertility indicative of
gynoecium defects, and infrequently produce siliques with only one valve (Prigge et al., 2020).
AUX/IAA interacting ARFs, which are therefore implicated in canonical auxin signalling
(Vernoux et al., 2011; Piya et al., 2014), are also involved in the regulation of gynoecium
development (Nagpal et al., 2005; Wang et al., 2005; Pekker et al., 2005; Kelley et al., 2012),
further supporting a role for the canonical pathway.
ETT shares partial redundancy with its paralogous ARF, ARF4, which contains a C-terminal
PB1-domain and is capable of AUX/IAA interaction (Pekker et al., 2005; Kelley et al., 2012;
Finet et al., 2010; Vernoux et al., 2011). However, whilst the transcriptional targets of ETT are
well described (Simonini et al., 2017; Andres-Robin et al., 2020), those of ARF4 are not.
Here, *tir1-1 afb2-3 ett*-3 mutants were generated in *Arabidopsis.* The triple mutant showed
exacerbation of the *ett-3* phenotype suggesting that the canonical and ETT-mediated
pathways function synergistically in gynoecium development. As little is known about the
global transcriptional regulation of genes by ARF4, a comparative transcriptomics analysis
was conducted to elucidate how ETT and ARF4 regulate targets, and to what extent these
targets overlap. The data presented suggest that the regulation of gynoecium development by
ETT and ARF4 is two-fold. Firstly, ETT and ARF4 redundantly regulate targets involved in cell
fate specification independently of auxin, and the mis-regulation of these genes correlates with
phenotypes observed in the double mutant. Secondly, ETT and ARF4 redundantly maintain
auxin insensitivity at target loci involved in the regulation of pectin, the mis-regulation of which
has previously been linked to morphogenesis defects within the *ett* gynoecium (Andres-Robin
et al., 2018). Most redundantly regulated targets became downregulated in response to auxin
in the *ett-3 arf4-2* mutant, whilst a subset of targets became upregulated. ETT and ARF4
maintain the auxin insensitivity of a range of targets, which became upregulated in response
to auxin upon their mutation, suggesting that A-B ARF antagonism previously observed in
Marchantia (Kato et al., 2020) and *Physcomitrium* (Lavy et al., 2016) may also regulate gene
expression in *Arabidopsis.* Together, the findings presented reveal crosstalk between the
canonical and ETT-mediated pathway and suggest that the maintenance of auxin insensitivity
at target loci plays an important role in a range of processes.
103
104 **Results**
105
Phenotypic analyses suggest synergism between canonical and ETT-mediated
pathways
To understand the relationship between the canonical and ETT-mediated auxin signalling
pathways during gynoecium development, *tir1-1 afb2-3 ett-3* triple mutants were generated in
Arabidopsis, and the resulting phenotypes were assessed (Figure 1, Supplemental Figure 1).

The *ett-3* mutant exhibited reduced valves, a loss of radialization in the style, and over-
proliferation of stigmatic tissues at later developmental stages, whilst the *arf4-2* and *tir1-2*
afb2-3 mutants resembled Col-0 (Figure 1, Supplemental Figure 1). The *ett-3* phenotype was
exacerbated in both *ett-3 arf4-2* and *tir1-1 afb2-3 ett-3*. Valves in *ett-3 arf4-2* gynoecia were
significantly reduced, producing an open ovary with ectopic ovules and few stigmatic papillae
(Figure 1, Supplemental Figure 1). The *ett-3 arf4-2* gynoecia lacked a style and split into two
distinct lobes with stigmatic papillae at their tips. Similarly, *tir1-1 afb2-3 ett-3* mutants exhibited
enhanced valve reduction relative to *ett-3* mutants, producing a partially open ovary with
ectopic ovules, and no discernable style or replum (Figure 1, Supplemental Figure 1). The
stigmatic papillae in *tir1-1 afb2-3 ett-3* gynoecia also developed atop two lobes produced at
the apex, and exhibited over proliferation of stigmatic tissues at later developmental stages
comparable to those observed in *ett-3* (Figure 1, Supplemental Figure 1).

The exacerbation of the *ett-3* phenotype by *tir1-1 afb2-3* mutation demonstrates that the
canonical and ETT-mediated pathways function synergistically to promote gynoecium
development. Furthermore, the phenotypic similarities between *ett-3 arf4-2* and *tir1-1 afb2-3*
ett-3 mutants, especially at the early stages of gynoecium development, suggest that
ETT/ARF4 and TIR1/AFB2 regulate similar processes in gynoecium development.

Comparative transcriptomics analysis of auxin independent and auxin sensitive
ETT/ARF4 targets

ETT is known to function partially redundantly with ARF4 in the regulation of gynoecium
development (Pekker et al., 2005; Finet et al., 2010; Kelley et al., 2012). Whilst ETT-mediated
signalling and the regulation of ETT-target genes are relatively well studied (Simonini et al.,
2016; Simonini et al., 2017; Kuhn et al., 2020), a global understanding of how ARF4 regulates
its targets, and which genes it modulates in conjunction with ETT, is lacking. ARF4 is known
to physically interact with AUX/IAAs (Vernoux et al., 2011), implicating it in the canonical
pathway.

To better understand how ARF4 regulates its targets both alone and together with ETT, a
comparative transcriptomic analysis (RNA sequencing, from here on RNA-Seq) was
conducted utilizing whole inflorescence tissues of Col-0, *ett-3, arf4-2, and ett-3 arf4-2* lines
treated with 100 μM Indole-3-Acetic Acid (IAA) and 10 μM N-1-Naphthylphthalamic Acid (NPA)
or mock treatments for one hour.

A principal component analysis revealed that the lines differed more by genotype than by
auxin treatment (Supplemental Figure 2), and many genes which are reportedly auxin
responsive (Paponov et al., 2008) were modulated in response to auxin in wildtype plants,
indicating that the auxin treatment was effective (Supplemental Table 1). *PINOID*, a previously
well-characterized ETT target (Simonini et al., 2016; Simonini et al., 2017; Kuhn et al., 2019)

which is upregulated in wildtype in response to auxin (Kuhn et al., 2020) was not identified as
an auxin sensitive gene within this study. Previous transcriptomics analyses aiming to identify
ETT targets utilized different schedules of auxin treatment (Kuhn et al., 2020; Simonini et al.,
2017) which may explain the observed discrepancies. Significant overlap was however
observed between genes mis-regulated in *ett-3* mutants between this study and Simonini et
al., 2017 (Supplemental Figure 3, Supplemental Table 2 and 3).
Two analyses were conducted on the resulting RNA-Seq data: an auxin independent analysis
which aimed to assess the developmental roles of ETT and ARF4 independently of auxin, and
an auxin sensitive analysis which assessed how the auxin responses vary from wildtype in the
mutants.
158
ETT and ARF4 are implicated in the regulation of cell fate commitment independently
of auxin
The auxin independent analysis was conducted in accordance with the schema in
Supplemental Figure 4. Two datasets each consisting of three lists of genes were used for
this analysis. The first dataset comprised lists of genes which were differentially expressed
relative to wildtype in *ett-3*, *arf4-2,* and *ett-3 arf4-2* after mock treatment, whilst the second
dataset comprised the same after IAA treatment. These lists were compared in Venny 2.1 to
generate two three-way Venn diagrams, one for each dataset. The data were then compared
across mock and auxin treatments to identify genes which are mis-regulated in the same
manner in both datasets. Any auxin sensitive genes within these lists were removed, leaving
genes which are regulated by ETT and ARF4 independently of auxin. Based on where these
genes positioned within a three-way Venn diagram, models of gene behavior were devised
which may explain the observed patterns of statistical significance (Figure 2 A and B).
172
Heatmaps were generated to assess whether genes followed the proposed models of
regulation, and Gene Ontology (GO) term analyses using PANTHER statistical
under/overrepresentation tests were conducted to identify biological processes which are
enriched within each target category (Figure 3).
The transcriptomics analysis identified 98 genes (53 up, 45 down) which were significantly
mis-regulated in *ett-3 arf4-2* relative to wildtype in accordance with their redundant regulation
by ETT and ARF4 (Figure 2B and C, Supplemental Figure 5). GO term analysis indicated
significant enrichment within this list for genes involved in cell fate commitment (Figure 3),
including *CLAVATA3/ESR-RELATED 19* (*CLE19*), *NO TRANSMITTING TRACT* (*NTT*), and
FAMA, which were significantly upregulated; and *YABBY2* (*YAB2*) and *YAB5* which were
significantly downregulated in *ett-3 arf4-2* (Figure 4). The mis-regulation of *YABBY* genes in

ett-3 arf4-2 is consistent with the abaxial-adaxial axis defects present within the leaves and

gynoecia of this line (Figure 1, Supplemental Figure 6).

Whilst *FAMA* was upregulated in *ett-3 arf4-2*, *TOO MANY MOUTHS* (*TMM*), another regulator

of the stomatal lineage, was constitutively downregulated (Supplemental Figure 5). Along with

the mis-regulation of stomatal lineage genes in *ett-3 arf4-2*, stomatal defects, such as stomatal

clustering on laminal outgrowths on the abaxial leaf surface, were also observed

(Supplemental Figure 6). The constitutive upregulation of the WUSCHEL related homeobox

transcription factor *WOX9*, and the CLV3 related peptide encoding gene *CLE19* (Figure 4,

Supplemental Figure 5), were observed in *ett-3 arf4-2*, along with variability in the numbers of

floral organs produced within flowers (Supplemental Figure 6). Finally, the auxin biosynthesis

gene *TAA1*, which promotes the production of auxin through the IPA pathway, was

constitutively upregulated in the double mutant (Supplemental Figure 5), suggesting that auxin

biosynthesis is also regulated redundantly by ETT and ARF4 independently of auxin.

Together, these data correlate with phenotypes observed in the *ett-3 arf4-2* mutant and

suggest that ETT and ARF4 may redundantly regulate cell fate specification independently of

auxin in a range of biological contexts, including the abaxial-adaxial axis of lateral organs and

the stomatal cell lineage.

Targets which became similarly mis-regulated in all mutant backgrounds, consistent with their

regulation by ETT-ARF4 heterodimers (151 genes, 56 up, 95 down; Supplemental Figure 7)

showed significant enrichment for involvement in gluco- and glycosinolate related secondary

metabolic processes (Figure 2, A, B and C; Figure 3). However, glucosinolate-related

phenotypes were not assessed within this study. These data suggest a role for ETT and ARF4

in the regulation of metabolic processes independently of auxin, and suggest that ETT and

ARF4 may regulate genes as heterodimers. Consistent with this hypothesis, interactions were

observed between ETT and ARF4 in yeast two hybrid assays. This interaction required the N-

terminal DNA Binding domain of ARF4, and was insensitive to auxin (Supplemental Figure 8).

The 76 targets which became similarly mis-regulated in both *ett-3* and *ett-3 arf4-2*, consistent

with their ETT-specific regulation, included *INDEHISCENT* (*IND*), *JAGGED* (*JAG*), and

CRABSCLAW (*CRC*), which have previously been implicated in gynoecium development

(Figure 5).

Whilst the *arf4-2* mutant appeared as wildtype (Figure 1), 49 ARF4-specific genes were

identified (Supplemental Figure 9). This list was not enriched for any specific biological

processes and did not contain any known regulators of gynoecium development, suggesting

that ARF4 mainly functions in conjunction with ETT to elicit its auxin-independent effects in

the carpels.

Together, these data suggest that ETT and ARF4 regulate distinct and overlapping targets

independently of auxin to promote gynoecium development.

Analysis of auxin sensitivity in ETT/ARF4 targets

Gynoecium morphogenesis is dependent on the dynamic distribution of auxin (Moubayidin &
Østergaard, 2014), and gynoecium defects of *ett-3 arf4-2* mutants suggest a role for ETT and
ARF4 in the transduction of auxin signalling (Pekker et al., 2005). A second analysis was
therefore conducted to assess how the auxin response varies from wildtype in the mutants.
Genes which were significantly auxin responsive in wildtype were compared to genes which
were significantly auxin responsive in each mutant background (*ett-3*, *arf4-2,* and *ett-3 arf4-*
2). Models of gene behaviour were devised, which may explain the patterns of statistical
significance observed for genes that fall into specific segments of a 4-way Venn diagram
(Figure 6). Evidence was found in the experimental data to support the following models of
gene regulation: genes are redundantly regulated by ETT and ARF4 to prevent their auxin
responsiveness (Figure 6B3); genes are regulated by ETT-ARF4 heterodimers which promote
(Figure 6B4) and prevent (Figure 6B10) their auxin responsiveness; the auxin insensitivity of
genes is specifically promoted by ARF4 (Figure 6B6); and genes are regulated by auxin
independently of ETT and ARF4 (Figure 6B8).

ETT and ARF4 redundantly prevent the auxin sensitivity of target genes

The auxin sensitive analysis identified 179 genes which were auxin sensitive only in the *ett-3*
arf4-2 mutant background (44 up and 135 down upon auxin treatment; Figure 6A, B3, C,
Figure 7). This suggests that these genes are redundantly regulated by ETT and ARF4 to
maintain their auxin insensitivity. PANTHER statistical over/under-representation analysis
showed that this list was enriched for genes involved in the regulation of pH, pectin catabolism,
and cell wall organization (Supplemental Figure 10). As the regulation of cell wall organization
has previously been linked to auxin-mediated cell expansion (Majda and Robert, 2018) and
organogenesis (Traas, 2019), the expression profiles of these genes were analysed in more
detail (Figure 8). Targets implicated in pectin metabolism and cell wall organization included
the *PECTIN METHYLESTERASES* (*PMEs*) *PME23*, *PME28*, *PME48*, *PME49*, *PPME1,* and
VGDH2 (VANGUARD 1 HOMOLOG 2); and six members of the pectin lyase superfamily
AT1G05660, *AT3G07850*, *AT5G48140*, *AT3G07830*, *AT3G07840*, *AT3G14040* (Figure 8). In
all cases, these genes became significantly downregulated in response to auxin only in the
ett-3 arf4-2 background, suggesting that ETT and ARF4 redundantly promote their expression
in the presence of auxin to prevent their downregulation (Figure 8).

ETT/ARF4 targets conform to models of auxin sensitive heterodimer regulation

Genes were identified within the auxin sensitive analysis which conformed to the heterodimer-mediated model of regulation (Figure 6, A, B4 and B10, C). 46 genes (36 up and 10 downregulated upon auxin treatment) were auxin sensitive in wildtype, and auxin insensitive in all mutant backgrounds, consistent with a model in which ETT-ARF4 heterodimers promote their auxin sensitivity (Figure 6 A, B4, C, Supplemental Figure 11). This list was significantly enriched for genes involved in the regulation of secondary metabolic processes relating glucosinolates (Supplemental Figure 12). A further 143 genes were auxin insensitive in wildtype and upregulated in response to auxin in all mutant backgrounds, suggesting that ETT-ARF4 dimers maintain auxin insensitivity at these loci (Figure 6, A, B10, C Supplemental Figure 13). The upregulation of these genes in response to auxin in all mutant backgrounds is consistent with A-B ARF antagonism previously observed in *Marchantia polymorpha* and *Physcomitirium patens* (Kato et al., 2020; Lavy et al., 2016). Consistent with this hypothesis, *ARF5*, *ARF6,* and *ARF8*, which are A-class activators, are expressed in the gynoecium along with ETT and ARF4 at various stages of development (Supplemental Figure 14). The list of targets whose auxin insensitivity is maintained by ETT-ARF4 dimers was enriched for targets involved in the negative regulation of defence responses (Supplemental Figure 12). No known regulators of gynoecium development were found within the lists of genes which conformed to heterodimer mediated regulation, suggesting that this mode of gene activity does not directly regulate auxin-mediated carpel development, but may be involved in the regulation of other biological processes.

ARF4 prevents the auxin sensitivity of specific targets independently of ETT

199 genes were identified which were similarly mis-regulated in *arf4-2* and *ett-3 arf4-2* mutants (five down, 194 upregulated upon auxin treatment; Figure 6A, B6, C, Figure 9), consistent with their ARF4-specific regulation. These genes were auxin insensitive in wildtype and became auxin sensitive in both mutant lines, implying that ARF4 maintains their auxin insensitivity. The upregulation of these targets in response to auxin in *arf4-2* and *ett-3 arf4-2* is consistent with previously proposed models of A-B ARF antagonism (Kato et al., 2020; Lavy et al., 2016). The list of ARF4 specific targets is significantly enriched for genes involved in the regulation of fruit ripening and processes relating to ethylene biosynthesis (Supplemental Figure 15).

Discussion

Synergism between canonical and ETT-mediated signalling

Prior to this study, several lines of evidence implicated the canonical pathway in the regulation of gynoecium development, including carpel defects observed in higher order *tir1/afb* mutants and class A-ARF mutants (Prigge et al., 2020; Nagpal et al., 2005); and the expression of canonical pathway receptor genes in the gynoecium (Cechetti et al., 2008; Dharmasiri et al., 2005). However, the relationship between the canonical and ETT-mediated pathways had not been dissected. The data presented here indicate synergism between the canonical and ETT-mediated pathway and suggest that ETT functions with ARF4 to regulate both distinct and overlapping targets.

ARF4 has previously been shown to interact with AUX/IAA proteins (Vernoux et al., 2011; Piya et al., 2014), implicating ARF4 in the canonical pathway. However, whether such interactions have a biological relevance *in planta* remains unknown. In *Marchantia*, the overexpression of repressive ARFs suppresses the constitutive auxin response phenotypes observed in *aux/iaa* mutants (Lavy et al., 2016), suggesting that repressive ARFs can transcriptionally regulate genes without AUX/IAA interactors. Whilst this is certainly true for *Arabidopsis* ETT, which lacks the C-terminal domain required for AUX/IAA interaction, the C-terminal AUX/IAA interacting domain of ARF4 is necessary for ARF4 functionality in the gynoecium (Finet et al., 2010).

In the canonical model of auxin signalling, AUX/IAAs allow the recruitment of TOPLESS related co-repressors to loci to switch off their expression in the absence of auxin (Causier et al., 2012). However, ETT and ARF4 contain repressor motifs which facilitate direct co-repressor recruitment (Lokerse and Weijers, 2009). The auxin sensitive interaction between ETT and TPL underpins the repression of genes in the absence of auxin in the ETT-mediated pathway (Kuhn et al., 2020). Future work will be required to unpick the relevance of ARF4-AUX/IAA and direct ARF4-TPL interactions to gain a better understanding of the mechanisms regulating ARF4 activity.

Comparative transcriptomics correlate with phenotypic defects observed in *ett-3 arf4-2* mutants

The auxin sensitive analysis identified distinct and overlapping targets of ETT and ARF4. Targets under the auxin-independent redundant regulation of ETT and ARF4 included the YABBY TFs *YAB2* and *YAB5*, whilst *CRC* was mis-regulated in both *ett-3* and *ett-3 arf4-2*, consistent with its ETT-specific regulation. YABBY proteins specify abaxial identity along with

the KANADI family of transcription factors (Eshed et al., 2004). ETT and ARF4 function co-
operatively with KANADI proteins to promote abaxial cell fate of lateral organs (Pekker et al.,
2005). Laminal outgrowths from the abaxial leaf surface have previously been observed in
kan1 kan2 loss of function mutants. This phenotype is suppressed in *yab1 yab3 yab5* (Eshed
et al., 2004) and *yuc1 yuc2 yuc4* backgrounds (Wang et al., 2011), suggesting that the
production of outgrowths is dependent on YABBYs and auxin biosynthesis. Similar outgrowths
were observed on the abaxial surface of *ett-3 arf4-2* mutant leaves in this study, consistent
with previously proposed roles for ETT/ARF4 in KANADI-mediated signalling (Pekker et al.,
2005). However, given that *YAB2*, *YAB5*, and *CRC* were significantly downregulated in the
ett-3 arf4-2 mutant background, it is unclear if these outgrowths are YABBY dependent. Whilst
yabby single mutants lack obvious abaxial-adaxial identity defects, higher order mutants show
significant alterations in abaxial-adaxial polarity (Sarojam et al., 2010; Siegfried et al., 1999).
The downregulation of multiple *YAB* transcription factors within the *ett-3 arf4-2* mutant is
therefore consistent with the abaxial-adaxial identity defects observed in the leaves and
gynoecia of the double mutant in this study.

Previous studies reported reduced valve margin definition in *ind* (Liljegren et al., 2004), and
apical gynoecium defects in *jag* and *crc* mutants (Bowman & Smyth, 1999; Ohno et al., 2004).
Similar gynoecium defects were exhibited in this study by *ett-3* and *ett-3 arf4-2* (Figure 1),
along with the constitutive downregulation of *IND, JAG,* and *CRC,* which is consistent with the
previously proposed roles of these genes in gynoecium development. Future studies should
aim to determine which of the targets identified within this study are direct/indirect ETT/ARF4
targets.

349
350
351

Auxin Sensitive analysis suggests ETT and ARF4 maintain auxin insensitivity at target
loci
Previous studies in *Marchantia polymorpha* and *Phiscomitrium patens* found antagonism
between A- and B-ARFs to regulate the auxin sensitivity of tissues (Kato et al., 2020; Lavy et
al., 2016), with the auxin response being promoted by A-ARFs and inhibited by B-ARFs.
Consistent with this model of B-ARF function, ETT and ARF4 were previously shown to
repress auxin responses in an *Arabidopsis* root hair system (Choi et al., 2018). The
transcriptomics data presented here suggest that ETT and ARF4 largely prevent the auxin
sensitivity of distinct and overlapping targets, many of which become upregulated in response
to auxin in various mutant backgrounds, consistent with models of A/B-ARF antagonism.
However, many of their redundantly regulated targets, including genes involved in pectin
methylesterification, pectin catabolism, and cell wall organisation become downregulated in

response to auxin. Auxin-mediated repression has been observed in TMK-mediated
signalling, where atypical AUX/IAAs are stabilized in the presence of auxin (Cao et al., 2019).
However, it is unclear if such a mechanism functions in the gynoecium. Previous studies have
implicated PME family members in the regulation of cell wall stiffness during gynoecium
development in *ett* mutants and showed that ETT promotes PME expression in the carpels
(Andres-Robin et al., 2018; Andres-Robin et al., 2020). Furthermore, the overexpression of
PECTIN METHYLESTERASE INHIBITOR 3 (*PMEI3*), which antagonises PME activity,
phenocopied the valve reductions observed in *ett* mutants (Andres-Robin et al., 2018). Taken
together with these previous studies, the data presented here suggest that ETT and ARF4
may redundantly promote the expression of *PMEs* to prevent their downregulation in response
to auxin, and the mis-regulation of these genes may contribute to the gynoecium defects
observed in *ett-3 arf4-2*.
The data presented suggest that ETT functions alongside ARF4 to prevent the expression of
genes which would otherwise become auxin sensitive in its absence, possibly through the
activity of A-ARFs with which they are co-expressed, or due to the activity of TMK-related
pathways in the case of auxin-downregulated genes. The data also suggest that the
maintenance of genes in an auxin insensitive state may be equally as important as the
promotion of auxin sensitivity at other loci to facilitate normal growth and development. Future
studies should continue to dissect the relationships between distinct auxin signalling pathways
to gain a greater understanding of the wider auxin signalling network, allowing for a more
global perspective of auxin-mediated plant development.
385
Materials and Methods

Plant materials and growth conditions

Seeds were sown onto damp soil (Levington F2 compost with Intercept and grit at a 6:1 ratio)
and stratified for 3 days at 4°C in the dark before transferal to a controlled environment room
(CER) and growth in long day conditions (16 hours light/8 hours dark, 22°C). All mutant lines
were generated in a Col-0 background.

Mutant and transgenic lines used in this work which have been previously published are: *ett-*
3 (Sessions et al., 1997; Simonini et al., 2016), *arf4-2* (SALK_070506, Alonso et al., 2003)
and *tir1-1 afb2-3* (Parry et al., 2009). The *ett-3 arf4-2* line was generated by crossing the *ett-*
3 line and *arf4-2* (SALK_070506). The *tir1-1 afb2-3 ett-3* line was generated by crossing *ett-*
3 with *tir1-1 afb2-3*. The PCR conditions and genotyping primers used for these lines are
listed in Supplemental Table 4 and Supplemental Table 5. Additional information relating to

the genotyping of each line can also be found in the supplemental material and methods.
pARF3/ETT: GFP, *pARF4: GFP*, *pARF5: GFP*, *pARF6: GFP* and *pARF8: GFP* are
previously published (Rademacher *et al.*, 2011).

Scanning Electron Microscopy

Whole inflorescences were harvested and incubated in fixative (3.7% formaldehyde, 3% acetic
acid and 50% ethanol) for 16 hours at room temperature on a shaker. The fixative was
aspirated, and the tissues were washed in an ethanol series (70%, 80%, 90% and 100%) at
room temperature on a shaker. Each wash was incubated for 30 minutes. The samples were
then washed twice for 30 minutes in 100% dry ethanol and critical point dried with carbon
dioxide using a Leica CPD300. Gynoecia were dissected onto stubs and gold coated using an
Agar high resolution sputter coater. Imaging was conducted with a Zeiss Supra 55VP Field
Emission Scanning Electron Microscope using a 3kV acceleration voltage.

RNA sequencing Analysis

Whole inflorescences were harvested from Col-0, *ett-3*, *arf4-2,* and *ett-3 arf4-2* lines treated
with either mock (dH2O, 0.1% tween, 0.1% ethanol) or 100 µM IAA and 10µM NPA (dissolved
in ethanol), 0.1% tween for one hour. Three biological replicates were harvested for each
genotype one hour after treatment. Each replicate contained pooled material from three plants.
The tissue was snap frozen in liquid nitrogen and RNA extraction was conducted using a
Qiagen RNeasy Plant Mini Kit with on-column DNase treatment, in accordance with the
manufacturer's guidelines. The RNA quality was assessed by gel electrophoresis, and the
RNA concentration was measured using a nanodrop (ThermoFisher), and the RNA integrity
was tested using a Bioanalyzer 2100 (Agilent). Novogene (Hong Kong), conducted cDNA
library preparation (250-300 bp insert cDNA library), and sequencing (20M pair reads), the
details of which can be found in the supplemental materials and methods.

Principle Component Analysis

Principle component analysis was conducted in iDEP 9.1 (http://ge- lab.org/idep/, (Ge et al.,
2018)) on expression values normalised to fragments per kilobase of transcript per million
mapped reads (FPKM).

Analysis of auxin independent ETT/ARF4 targets

This analysis was conducted using the differentially expressed gene lists provided by Novogene (Hong Kong). Six lists of differentially expressed genes were used in this analysis: (1) *arf4-2* vs Col-0 mock; (2) *ett-3* vs Col-0 mock; (3) *ett-3 arf4-2* vs Col-0 mock; (4) *arf4-2* vs Col-0 IAA; (5) *ett-3* vs Col-0 IAA; and (6) *ett- 3 arf4-2* vs Col-0 IAA. Only genes which were at least 2-fold up or downregulated in at least one background were included in this analysis (log2fold value of >1 or < -1). Analyses were conducted with Venny (Version 2.1; https://bioinfogp.cnb.csic.es/tools/venny/index.html; Oliveros, 2007) to identify genes mis-regulated only in one background, mis-regulated only in two backgrounds, or mis-regulated in all three backgrounds relative to wildtype. The mock and auxin treated datasets were independently assessed using Venny, and then compared to identify genes which were similarly differentially expressed in both the presence and absence of auxin. Finally, any genes which were significantly differentially expressed (2-fold up or down-regulated in at least one background) in response to auxin were removed from the lists, leaving only those genes which were regulated independently of auxin.

Analysis of auxin sensitive ETT/ARF4 targets

The auxin sensitive analysis utilized DEGs from the following lists generated by Novogene (Hong Kong): (1) Col-0 IAA vs Col-0 mock; (2) *ett-3* IAA vs *ett-3* mock; (3) *arf4-2* IAA vs *arf4-2* mock; and (4) *ett-3 arf4-2* IAA vs *ett-3 arf4-2* mock. The genes which were differentially expressed in each background in response to auxin were then compared to reveal how the auxin response in each mutant differed from WT. Venny (Version 2.1) was utilized to generate a four-way Venn diagram.

Heatmap Analysis

Heatmapper (Babicki et al., 2016) was utilized to generate heatmaps showing the average FPKM for genes in each category in both the auxin independent and auxin sensitive analyses. The settings used in Heatmapper were as follows: Scale type was set to row; colour brightness to 50; number of shades = 50; colour scheme red/green; clustering method average linkage; distance measurement method Spearman rank correlation; clustering applied to rows; and dendrograms were shown for rows.

Go Term Analysis

Gene Ontology (GO) term analysis was conducted on the lists of genes which fell into each
category of the Venn diagrams in both the auxin independent and auxin sensitive analyses to
identify whether these lists were enriched for genes involved in specific biological processes.
This GO term analysis was conducted using a statistical under/over-representation test, which
utilizes Fisher's Exact Test for p value estimation (p<0.05). The GO term analysis utilized the
PANTHER classification system (Mi et al., 2019).

Yeast 2 Hybrid

The full-length coding sequence of *ETT* and *ARF4* and truncated *ARF4* fragments were
amplified by PCR using primers designed for gateway cloning (Supplemental Table 6). The
resulting PCR products were purified using a Qiagen QIAquick PCR purification kit and
introduced in the gateway entry vector pDONR207 (Life Technologies) by BP clonase
reactions, in accordance with the Thermo-Fisher Gateway protocol. The products of this
reaction were transformed into DH5α cells (Invitrogen) according to the manufacturer's
guidelines and grown on LB agar plates containing gentamycin (10 µg/ml) overnight at 37 $^\circ$C.
Constructs were validated by colony PCR and sequencing. The ARF sequences within
pDONR207 were then recombined into the pGAD-T7 and pGBK-T7 (Clontech) expression
vectors by LR clonase reactions, transformed into *E. coli*, and once more validated by colony
PCR and sequencing. The expression vectors were then co-transformed into AH109
Saccharomyces cerevisiae (Clontech), by lithium acetate mediated transformation (Gietz and
Schiestl, 2007). Different combinations of AD (pGAD-T7) and BD (pGBK-T7) vectors, or empty
vector controls, were introduced in pairs to competent yeast cells to determine whether ARF4
and ETT interact, and through which domains of ARF4. Transformants were plated onto YSD
-W-L media and grown at 28°C for 72 hours, before serial dilution (10^0, 10^1, 10^2, and 10^3) and
plated onto YSD -W-L-H-A plates to assess interactions. After plating on selective media,
yeast colonies were photographed after growth at 28°C for 7 days. Figures were created using
images from the 10^1 dilution in all cases. Auxin sensitivity was assessed by adding 100 µM
IAA (Sigma) to the medium.

Confocal Microscopy

Stage 8-14 gynoecia were dissected from the plant and placed onto slides in water containing 0.015% Silwet L-77 (De Sangosse Ltd.) to prevent the formation of bubbles. Cover slips were then applied, and the gynoecia were imaged using a Leica SP5 confocal microscope (HyD detector, Smart Gain 30% 200hz, 4x Line Average) with a 40x water immersion lens. GFP was excited at a wavelength of 488 nm and detected at 498–530 nm. Images were taken as Z-stacks, and then processed in Image J 1.48 to form composite 3D projections, which were then converted to RGB images.

Author contribution

 H.M.M. and L.Ø. devised the experiments. H.M.M. and B.N. conducted experiments. H.M.M. wrote the manuscript.

Acknowledgements

Apologies to any colleagues whose work we have not cited due to space limitations. Special thanks to Yang Dong (IBCAS) for his continued input and collaboration on this project; to Henrik Jönsson (SLCU) for supporting the completion of the project; and to James McLaughlin (EMBL-EBI) for bioinformatics support. H.M.M. was funded by the John Innes Foundation (at JIC) and the Gatsby Charitable Foundation (at SLCU, GAT3395-PR4B), B.N. by BBSRC (BB/S002901/1) and L.Ø. by BBSRC (BB/P013511/1).

References

References

Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, Stevenson DK, Zimmerman
J, Barajas P, Cheuk R, Gadrinab C, Heller C, Jeske A, Koesema E, Meyers CC, Parker
H, Prednis L, Ansari Y, Choy N, Deen H, Geralt M, Hazari N, Hom E, Karnes M,
Mulholland C, Ndubaku R, Schmidt I, Guzman P, Aguilar-Henonin L, Schmid M, Weigel
D, Carter DE, Marchand T, Risseeuw E, Brogden D, Zeko A, Crosby WL, Berry CC, Ecker
JR. 2003. Genome-wide insertional mutagenesis of Arabidopsis thaliana. *Science (1979)*
301. doi:10.1126/science.1086391

Andres-Robin A, Reymond MC, Brunoud G, Martin-Magniette ML, Monéger F, Scutt CP. 2020.
Immediate targets of ETTIN suggest a key role for pectin methylesterase inhibitors in the
control of Arabidopsis gynecium development. *Plant Signal Behav* **15**.
doi:10.1080/15592324.2020.1771937

Andres-Robin A, Reymond MC, Dupire A, Battu V, Dubrulle N, Mouille G, Lefebvre V, Pelloux
J, Boudaoud A, Traas J, Scutt CP, Monéger F. 2018. Evidence for the regulation of
gynoecium morphogenesis by ETTIN via cell wall dynamics. *Plant Physiol* **178**.
doi:10.1104/pp.18.00745

Babicki S, Arndt D, Marcu A, Liang Y, Grant JR, Maciejewski A, Wishart DS. 2016.
Heatmapper: web-enabled heat mapping for all. *Nucleic Acids Res* **44**.
doi:10.1093/NAR/GKW419

Cao M, Chen R, Li P, Yu Y, Zheng R, Ge D, Zheng W, Wang X, Gu Y, Gelová Z, Friml J,
Zhang H, Liu R, He J, Xu T. 2019. TMK1-mediated auxin signalling regulates differential
growth of the apical hook. *Nature* **568**. doi:10.1038/s41586-019-1069-7

Cecchetti V, Altamura MM, Falasca G, Costantino P, Cardarelli M. 2008. Auxin regulates
Arabidopsis anther dehiscence, pollen maturation, and filament elongation. *Plant Cell*
20:1760–1774. doi:10.1105/tpc.107.057570

Choi HS, Seo M, Cho HT. 2018. Two TPL-binding motifs of ARF2 are involved in repression
of auxin responses. *Front Plant Sci* **9**. doi:10.3389/fpls.2018.00372

Dharmasiri N, Dharmasiri S, Weijers D, Lechner E, Yamada M, Hobbie L, Ehrismann JS,
Jürgens G, Estelle M. 2005. Plant development is regulated by a family of auxin receptor
F box proteins. *Dev Cell* **9**:109–119.

Eshed Y, Izhaki A, Baum SF, Floyd SK, Bowman JL. 2004. Asymmetric leaf development and
blade expansion in Arabidopsis are mediated by KANADI and YABBY activities.
Development **131**. doi:10.1242/dev.01186

Fendrych M, Akhmanova M, Merrin J, Glanc M, Hagihara S, Takahashi K, Uchida N, Torii KU,
Friml J. 2018. Rapid and reversible root growth inhibition by TIR1 auxin signalling. *Nat*
Plants **4**. doi:10.1038/s41477-018-0190-1

Finet C, Berne-Dedieu A, Scutt CP, Marlétaz F. 2013. Evolution of the ARF gene family in
land plants: Old domains, new tricks. *Mol Biol Evol* **30**:45–56.
doi:10.1093/molbev/mss220

Finet C, Fourquin C, Vinauger M, Berne-Dedieu A, Chambrier P, Paindavoine S, Scutt CP.
2010. Parallel structural evolution of auxin response factors in the angiosperms. *Plant*
Journal **63**. doi:10.1111/j.1365-313X.2010.04292.x

Friml J, Gallei M, Gelová Z, Johnson A, Mazur E, Monzer A, Rodriguez L, Roosjen M,
Verstraeten I, Živanović BD, Zou M, Fiedler L, Giannini C, Grones P, Hrtyan M, Kaufmann
WA, Kuhn A, Narasimhan M, Randuch M, Rýdza N, Takahashi K, Tan S, Teplova A,
Kinoshita T, Weijers D, Rakusová H. 2022. ABP1–TMK auxin perception for global
phosphorylation and auxin canalization. *Nature* **609**:575–581. doi:10.1038/s41586-022-
05187-x

Gietz RD, Schiestl RH. 2007. High-efficiency yeast transformation using the LiAc/SS carrier
DNA/PEG method. *Nat Protoc* **2**. doi:10.1038/nprot.2007.13

Guilfoyle TJ, Hagen G. 2012. Getting a grasp on domain III/IV responsible for Auxin Response
Factor-IAA protein interactions. *Plant Science*. doi:10.1016/j.plantsci.2012.04.003

Kato H, Mutte SK, Suzuki H, Crespo I, Das S, Radoeva T, Fontana M, Yoshitake Y, Hainiwa
E, van den Berg W, Lindhoud S, Ishizaki K, Hohlbein J, Borst JW, Boer DR, Nishihama
R, Kohchi T, Weijers D. 2020. Design principles of a minimal auxin response system. *Nat*
Plants **6**:473–482. doi:10.1038/s41477-020-0662-y

Kelley DR, Arreola A, Gallagher TL, Gasser CS. 2012. ETTIN (ARF3) physically interacts with
KANADI proteins to form a functional complex essential for integument development and
polarity determination in Arabidopsis. *Development* **139**. doi:10.1242/dev.067918

Kepinski S, Leyser O. 2005. The Arabidopsis F-box protein TIR1 is an auxin receptor. *Nature*
435. doi:10.1038/nature03542

Kuhn A, Harborough SR, McLaughlin HM, Natarajan B, Verstraeten I, Friml J, Kepinski S,
Østergaard L. 2020. Direct ETTIN-auxin interaction controls chromatin states in
gynoecium development. *Elife* **9**. doi:10.7554/eLife.51787

Kuhn A, Runciman B, Tasker-Brown W, Østergaard L. 2019. Two auxin response elements
fine-tune pinoid expression during gynoecium development in arabidopsis thaliana.
Biomolecules **9**. doi:10.3390/biom9100526

Lavy M, Prigge MJ, Tao S, Shain S, Kuo A, Kirchsteiger K, Estelle M. 2016. Constitutive auxin
response in Physcomitrella reveals complex interactions between Aux/IAA and ARF
proteins. *Elife*. doi:https://doi.org/10.7554/eLife.13325

Liljegren SJ, Roeder AHK, Kempin SA, Gremski K, Østergaard L, Guimil S, Reyes DK,
Yanofsky MF. 2004. Control of fruit patterning in Arabidopsis by INDEHISCENT. *Cell* **116**.
doi:10.1016/S0092-8674(04)00217-X

Lokerse AS, Weijers D. 2009. Auxin enters the matrix - assembly of response machineries for
specific outputs. *Curr Opin Plant Biol*. doi:10.1016/j.pbi.2009.07.007

Majda M, Robert S. 2018. The role of auxin in cell wall expansion. *Int J Mol Sci*.
doi:10.3390/ijms19040951

Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD. 2019. PANTHER version 14: More
genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools.
Nucleic Acids Res **47**. doi:10.1093/nar/gky1038

Moubayidin L, Østergaard L. 2014. Dynamic control of auxin distribution imposes a bilateral-
to-radial symmetry switch during gynoecium development. *Current Biology* **24**.
doi:10.1016/j.cub.2014.09.080

Nagpal P, Ellis CM, Weber H, Ploense SE, Barkawi LS, Guilfoyle TJ, Hagen G, Alonso JM,
Cohen JD, Farmer EE, Ecker JR, Reed JW. 2005. Auxin response factors ARF6 and
ARF8 promote jasmonic acid production and flower maturation. *Development* **132**.
doi:10.1242/dev.01955

Nemhauser JL, Feldman LJ, Zambryski PC. 2000. Auxin and ETTIN in Arabidopsis gynoecium
morphogenesis. *Development* **127**. doi:10.1242/dev.127.18.3877

Oliveros JC. 2007. VENNY. An interactive tool for comparing lists with Venn Diagrams.
http://bioinfogp.cnb.csic.es/tools/venny/index.html.
BioinfogpCnbCsicEs/Tools/Venny/IndexHtml.

Paponov IA, Paponov M, Teale W, Menges M, Chakrabortee S, Murray JAH, Palme K. 2008.
Comprehensive transcriptome analysis of auxin responses in Arabidopsis. *Mol Plant*
1:321–337. doi:10.1093/mp/ssm021

Parry G, Calderon-Villalobos LI, Prigge M, Peret B, Dharmasiri S, Itoh H, Lechner E, Gray
WM, Bennett M, Estelle M. 2009. Complex regulation of the TIR1/AFB family of auxin
receptors.

Pekker I, Alvarez JP, Eshed Y. 2005. Auxin response factors mediate Arabidopsis organ
asymmetry via modulation of KANADI activity. *Plant Cell* **17**. doi:10.1105/tpc.105.034876

Piya S, Shrestha SK, Binder B, Neal Stewart C, Hewezi T. 2014. Protein-protein interaction
and gene co-expression maps of ARFs and Aux/IAAs in Arabidopsis. *Front Plant Sci* **5**:1–
9. doi:10.3389/fpls.2014.00744

Prigge MJ, Platre M, Kadakia N, Zhang Y, Greenham K, Szutu W, Pandey BK, Bhosale RA,
Bennett MJ, Busch W, Estelle M. 2020. Genetic analysis of the arabidopsis TIR1/AFB
auxin receptors reveals both overlapping and specialized functions. *Elife* **9**.
doi:10.7554/eLife.54740

Sarojam R, Sappl PG, Goldshmidt A, Efroni I, Floyd SK, Eshed Y, Bowmana JL. 2010.
Differentiating Arabidopsis shoots from leaves by combined YABBY activities. *Plant Cell*
22. doi:10.1105/tpc.110.075853

Sessions A, Nemhauser JL, McColl A, Roe JL, Feldmann KA, Zambryski PC. 1997. ETTIN
patterns the Arabidopsis floral meristem and reproductive organs. *Development* **124**.
doi:10.1242/dev.124.22.4481

Siegfried KR, Eshed Y, Baum SF, Otsuga D, Drews GN, Bowman JL. 1999. Members of the
YABBY gene family specify abaxial cell fate in Arabidopsis. *Development* **126**.
doi:10.1242/dev.126.18.4117

Simonini S, Bencivenga S, Trick M, Østergaard L. 2017. Auxin-induced modulation of ETTIN
activity orchestrates gene expression in arabidopsis. *Plant Cell* **29**.
doi:10.1105/tpc.17.00389

Simonini S, Deb J, Moubayidin L, Stephenson P, Valluru M, Freire-Rios A, Sorefan K, Weijers
D, Friml J, Østergaard L. 2016. A noncanonical auxin-sensing mechanism is required for
organ morphogenesis in Arabidopsis. *Genes Dev* **30**. doi:10.1101/gad.285361.116

Szemenyei H, Hannon M, Long JA. 2008. TOPLESS mediates auxin-dependent
transcriptional repression during Arabidopsis embryogenesis. *Science (1979)* **319**.
doi:10.1126/science.1151461

Tiwari SB, Hagen G, Guilfoyle T. 2003. The roles of auxin response factor domains in auxin-
responsive transcription. *Plant Cell* **15**:533–543. doi:10.1105/tpc.008417

Traas J. 2019. Organogenesis at the shoot apical meristem. *Plants*.
doi:10.3390/plants8010006

Ulmasov T, Hagen G, Guilfoyle TJ. 1999. Activation and repression of transcription by auxin-
response factors. *Proc Natl Acad Sci U S A* **96**. doi:10.1073/pnas.96.10.5844

Vernoux T, Brunoud G, Farcot E, Morin V, van den Daele H, Legrand J, Oliva M, Das P,
Larrieu A, Wells D, Guédon Y, Armitage L, Picard F, Guyomarc'H S, Cellier C, Parry G,
Koumproglou R, Doonan JH, Estelle M, Godin C, Kepinski S, Bennett M, de Veylder L,
Traas J. 2011. The auxin signalling network translates dynamic input into robust
patterning at the shoot apex. *Mol Syst Biol* **7**. doi:10.1038/msb.2011.39

Wang JW, Wang LJ, Mao YB, Cai WJ, Xue HW, Chen XY. 2005. Control of root cap formation
by MicroRNA-targeted auxin response factors in Arabidopsis. *Plant Cell* **17**:2204–2216.
doi:10.1105/tpc.105.033076

Wang W, Xu B, Wang H, Li J, Huang H, Xu L. 2011. YUCCA genes are expressed in response
to leaf adaxial-abaxial juxtaposition and are required for leaf margin development. *Plant*
Physiol **157**. doi:10.1104/pp.111.186395

Figures

689 Figure 1: Phenotypes of stage 10-12 and 12-14 gynoecia in scanning electron micrographs

690 (with backgrounds removed for clarity, original images can be seen in Supplemental Figure

691 1), and photographs of stage 16-18 siliques in Col-0, *ett-3*, *arf4-2*, *tir1-1 afb2-3*, *ett-3 arf4-2*

692 and *tir1-1 afb2-3 ett-3* lines. Scale bars in Scanning electron micrographs show 200 μm in

693 photographs, 3 cm.

Figure 2: Summary of the findings of the auxin independent analysis A) Venn diagram of the proposed relationships between ETT, ARF4 and downstream targets which are regulated independently of auxin. B) Models of the behaviour of genes which fall into each of these categories, ns, not significant, * significant, -, mock, +, IAA treatment. Whilst the bar charts display an example of genes becoming upregulated in these models, in the datasets analysed they could be up or down regulated. C) Numbers of genes that fall in each sector of the Venn diagram, which are significantly differentially expressed relative to Col-0 in the mutant backgrounds.

753 Figure 3: Graphical representation of the significantly enriched GO terms (biological process)

754 identified in PANTHER over/under-representation analysis of the auxin independent dataset.

Figure 4: The relative expression (FPKM) of genes involved in cell fate commitment in each genotype after mock (-) or auxin (+) treatment. Statistical analysis was conducted using read count normalization, negative binomial distribution analysis and FDR correction using the BH procedure. The error bars show +/- standard deviation, ns, not significant * = padjust<0.05.

812 Figure 5: the relative expression (FPKM) of ETT-specific targets with a known role in
813 gynoecium development after mock (-) or auxin (+) treatment. Statistical analysis was
814 conducted using read count normalization, negative binomial distribution analysis and FDR
815 correction using the BH procedure (conducted by Novogene). Error bars show +/- standard
816 deviation, ns, not significant, * = padjust<0.05.

Figure 6: Summary of the findings of the auxin sensitive analysis A) a summary of how the genes in each section of the Venn diagram are predicted to behave, and B) models of gene behaviour corresponding to the segments shown in A after mock (-) or auxin (+) treatment, where * denotes statistical significance. Whilst the examples show genes becoming significantly upregulated in response to auxin in the figure, this is just to demonstrate the patterns of significance expected; in the experimental data the genes could become significantly up or downregulated. C) The number of genes identified which fall into each category of the Venn diagram.

Figure 7: Heatmap showing the genes which become auxin sensitive in the *ett-3 arf4-2* background suggesting that ETT and ARF4 redundantly prevent their auxin responsiveness. Mock and auxin treatment are shown by – and + respectively.

Figure 8: The relative expression (FPKM) of genes involved in pectin catabolism and cell wall organization identified through GO term (biological process) analysis. Statistical analysis was conducted using read count normalization, negative binomial distribution analysis and FDR correction using the BH procedure. The error bars show +/- standard deviation, ns, not significant, * = padjust<0.05. Mock and auxin treatments are shown by – and + respectively.

Figure 9**:** Heatmap of genes whose auxin sensitivity is prevented specifically by ARF4. Genes were clustered by average linkage using Spearman's rank correlation. Genes which are highly expressed are shown in green, whilst low expression is shown in red, and mock and auxin treatments are shown by – and + respectively.