# Scale dependence in agricultural drivers of field margin plant communities: influence of spatial resolution and extent

**Isis Poinas[a,b,*], Guillaume Fried[b], Laura Henckel[c], Christine N Meynard[a]**

[a] CBGP, INRAE, CIRAD, Institut Agro, IRD, Univ Montpellier, Montferrier-sur-Lez, France

[b] Anses, Laboratoire de la Santé des Végétaux, Unité Entomologie et Plantes invasives, Montferrier-sur-Lez, France

[c] INRAE Dijon, UMR Agro-écologie, Dijon, France

**Authorship statement:** I.P., G.F. and C.M. planned and designed the research; L.H. contributed to data processing; I.P. analyzed the data and wrote the first draft of the manuscript; G.F., L.H. and C.M. contributed substantially to revisions. All authors gave final approval for publication.

**Data accessibility statement:** Data available via the Data INRAE Repository, at

https://doi.org/10.57745/NMIPCY

**Number of words in abstract:** 300

**Number of words in main text:** 6991

**Number of cited references:** 47

**Number of tables:** 4

**Number of figures:** 3

---

[*] Corresponding author. Tel: +33 6 86 53 41 80. Corresponding address: 755 avenue du campus Agropolis, CS 30016, 34988 Montferrier-sur-Lez Cedex, France
E-mail address: isis.poinas@inrae.fr.

**Abstract**

In recent decades, agricultural intensification has led to a strong decline in biodiversity. Agricultural

field margins, defined as the uncultivated herbaceous vegetation located between the cultivated strip

and an adjacent habitat, usually constitute an important refuge for biodiversity in highly disturbed

landscapes, and are critical to the preservation of ecosystem services. However, field margins are also

strongly impacted by agricultural practices in their immediate vicinity. Agricultural impacts are often

studied at local to landscape scales, and rarely include national or larger spatial extents, where one of

the challenges is the lack of availability of standardized monitoring efforts including both biodiversity

and agricultural practices. Here, we take advantage of one such monitoring effort to assess the effects

of agricultural practices on field margin flora at different extents and resolutions in continental France.

We used spatial simultaneous autoregressive and generalized dissimilarity models to assess the

response of species richness and community composition to climatic, soil, landscape, agricultural

(fertilization, herbicides) and margin management factors. Analyses were repeated at the plot-level,

as well as at 25, 40, 60 and 75 km resolutions, and at regional and national extents. We found that

species richness is better explained by agricultural practices at the plot-level, and by climate and crop

diversity at coarse resolutions, whereas composition responds more strongly to climate at the plot-

level, and to fertilization and crop diversity at coarse resolutions. There was a strong variation in the

explanatory percentage among regions, but climate effects tended to be weaker within biogeographic

units compared to the national level. This allows a better understanding of landscape and management

effects, and reveals interactions between local and regional effects. We suggest that providing an

agricultural regulatory framework at the scale of biogeographic subdivisions could help provide

pertinent measures to conserve biodiversity that are better adapted to the regional context.

**Keywords:** field margin, arable plants, plant assembly, agroecosystem, spatial scale, spatial resolution,

spatial extent, species pool, crop diversity, agricultural practices

**Introduction**

Since the second half of the 20th century, agricultural intensification has resulted in steady declines in biodiversity across a wide range of taxa and habitats (Emmerson et al., 2016). The mechanization of agriculture and the resulting aggregation of cropped fields has contributed to the loss of semi-natural areas, such as herbaceous field margins, defined here as uncultivated vegetated areas located between the cultivated strip and the adjacent habitat. Despite their modest area in France (less than 1%), field margins play an important ecological role in agroecosystems, as refuge islands and food sources supporting endangered species (Storkey et al., 2011), pollinators (Zamorano et al., 2020) and pest predators (Woodcock et al., 2016). Within landscapes, they act as natural corridors linking habitats and supporting the maintenance of populations of plants, insects, mammals and birds (Haddaway et al., 2018). Locally, they buffer pesticide and fertilizer drift into adjacent terrestrial and aquatic habitats (Haddaway et al., 2018). Their importance has been increasingly recognized in recent European agricultural policies, for example through the conditioning of a green payment provided that margin strips are preserved (Matthews, 2013). Field margins are ecotones and, as such, reflect a mixture of weed ruderal species well adapted to agricultural disturbances, with species that are typical of grasslands and forests (Aavik & Liira, 2009). Many of these species are not directly targeted by agricultural practices, but their habitat is the most directly exposed. Therefore, they are of great interest for studying the unintended impacts of agriculture on biodiversity.

At the local scale, agricultural practices play a key role in shaping the species distribution and diversity of plants from field margins (Marshall & Moonen, 2002). Each crop type has its own characteristic combination of practices (e.g., sowing date, fertilization level, herbicide use), that shapes the floristic community structure within agricultural fields (Bourgeois et al., 2020), and may have spillover effects into the field margins. Chemical inputs (e.g., pesticides and fertilizers) drift to the margins and directly affect plant composition and richness (Fried et al., 2018). Finally, direct management of margins further selects for species adapted to these disturbances (Bassa et al., 2011). Agricultural practices also

aggregate at larger spatial scales and impact landscape dispersal dynamics (Tscharntke et al., 2005). At

the landscape scale, habitat composition is an important driver of community structure. The

proportion and diversity of non-crop habitats in the landscape generally has a positive effect on field

margin biodiversity (Rader et al., 2014), along with crop diversity (Sirami et al., 2019). At the regional

and national scale, cropping systems are defined by the sequence of crop successions along with the

cultural practices associated to each crop (Leenhardt et al., 2010); they form stable uniform

agricultural units which are called agricultural regions in France (Richard-Schott, 2009).  However, the

influence of these agricultural regions on regional species pools remains to be clarified. While it is well

established that soil and climate shape the distribution of species across regions forming cohesive

pools of species (Cornell & Harrison, 2014), the challenge remains to characterize how these wide

assemblages respond to large-scale structured agricultural practices.

In ecology, spatial scale is usually defined by two attributes: resolution (or grain) and extent, both of

them having a strong importance on the processes unveiled. Spatial grain is the size of the unit of

analysis. It is often assumed that coarser resolutions allow ignoring local stochasticity to focus on the

main patterns structuring communities (Chase, 2014). For example, landscape is expected to have the

greatest influence when the grain size is closest to the average area of the different habitats. However,

using excessively coarse resolutions may lead to uninformative patterns (Viana & Chase, 2019). Spatial

extent is the total area covered by the sampling. It determines the range of variability in environmental

gradients, as well as the potential of inclusion of dispersal processes (Viana & Chase, 2019). Naturally,

local filtering effects are easier to detect at small extents, and within biogeographic regions, where

different study sites are more likely to represent homogeneous conditions for other predictors (e.g.

climate) (Viana et al., 2016).

Very few studies have explored issues related to the integration of several spatial scales in

agroecosystems. Some suggest that agricultural effects could prevail at fine spatial grains. For example,

weed species' richness within winter wheat crops in Spain was found to decrease linearly with field-

level intensification, while remaining constant at the landscape-level (Guerrero et al., 2014). However,

field margin communities may be subjected to a greater influence of landscape than agricultural

practices (as found by Berquer et al. (2021) at a scale lower than 1 km), in contrast to what happens

within agricultural fields. Concepción et al. (2012) revealed that extensive farming has a weaker

positive impact on weed diversity in field center and inner field edge when the landscape is too simple

within a 500 m radius. This suggests that landscape-scale simplification or intensification may result in

a depletion of the regional species pool, which then becomes primarily composed of species adapted

to agricultural disturbances, therefore altering the diversity and composition of local assemblages in

the long term. We can thus assume that, in some instances, the structure of agricultural communities

could be better explained at a landscape or regional spatial grain. Regarding spatial extent, although

some studies have addressed national (Rader et al., 2014), continental (Billeter et al., 2008) or regional

scales (Nascimbene et al., 2012), these usually included a small sample size, several proxies instead of

direct measures of the targeted agricultural practices, and a lack of standard survey procedures across

plots. The development of larger-scale studies with standardized sampling efforts is required to

thoroughly address the issue of spatial scale on agricultural impacts.

Here we aim at teasing out the effects of spatial resolution and extent, for a better understanding of

agricultural practices on plant diversity and composition in field margins. To this end, we rely on a

standardized national monitoring effort in 462 agricultural field margins covering France, including

monitoring of plant communities as well as agricultural practices. Initial analyses of this dataset at a

plot-level showed that vegetation composition is primarily structured along a landscape and soil

gradient, and secondarily, along an agricultural intensification gradient (Andrade et al., 2021; Fried et

al., 2018). Composition is driven by margin management and fertilization, while species richness

responds to the level of intensification (herbicide applications). Building on these first findings, we

investigate further by varying the spatial extent and resolution of analysis. Regarding spatial resolution,

our first hypothesis is that local agricultural practices will be easier to detect at finer grains because

field margin flora is primarily affected by local factors. Our second hypothesis is that landscape and

regional aggregation effects, which influence species filtering at larger spatial grains, will predominate

over local practices at broader resolutions. Regarding spatial extent, our hypothesis is that soil and

climate will dominate at the national extent and become less important within biogeographic regions,

providing a clearer picture of the influence of agricultural practices at the regional extent.

**Materials and methods**

*Vegetation survey*

We used vegetation data from a national monitoring effort, the 500-ENI network, which is funded by

the French Ministry of Agriculture (see details in Andrade et al., 2021). Agricultural field margins

covering continental France, were surveyed yearly since 2013, representing a total of 543 unique plots

between 2013 and 2019 (including some plot turnover between years, see **Fig. SA.1**). These survey

plots are located in field margins representing four main crop types:  annual crops (with winter wheat

or maize as the main crop production, but including other crops in the rotation), market gardening

crops (mainly lettuce) and vineyards. The proportion of plots under organic farming is roughly 20%,

but agricultural practices cover a wide range of pesticide application, fertilizers and soil management.

We selected a subset of plots that had at least five years of botanical data between 2013 and 2019,

leaving a total of 462 field margins. Within each survey plot, plant species were identified in ten 1 m²

quadrats along the field margin: the ten quadrats were divided into two sets of five contiguous

quadrats of 0.25 x 2 m² separated by 30 m (**Fig. SA.2**). The transect line was equidistant from the

cultivated strip and the adjacent habitat, and the presence-absence of each species was recorded for

each quadrat. Surveys were performed at peak flowering (between the end of April and the beginning

of August, depending on the region). At the national scale, this represented 3079 observations (year x

plot), leading to the identification of 689 species. Since we were interested in spatial rather than

temporal variations, we summarized plant surveys by averaging the presence-absence of the species

detected across years, resulting in one data point per field margin, therefore representing a frequency

of occurrence varying continuously from 0 to 1 (0 = was never detected in that plot; 1 = was detected

throughout all years).

*Explanatory variables*

We gathered two types of explanatory variables: the first came directly from the 500-ENI network and

reflects agricultural practices assessed directly on the plots; the second one was retrieved from

external open access databases. These include soil, climate and landscape data.

Agricultural practices were reported yearly from interviews of farmers into an online database

according to a national standardized acquisition protocol. Data collected mainly relate to fertilization,

pesticide use, tillage and boundary management (**Fig. SA.3**). Soil characteristics (in the layer 0-5 cm)

were downloaded from SoilGrids at a 250 m resolution (Hengl et al., 2014), while climatic data (means

and variability by year and season for the period 1979-2013) were downloaded from Chelsa Climate

database at a 1 km resolution (Karger et al., 2017). We extracted landscape descriptors from OSO land

cover maps at a 10 m resolution for 2016 (Inglada et al., 2017). This is a median year with respect to

the temporal period covered by our sampling, and we assumed that landscape cover did not change

dramatically between 2013 and 2019. We considered percent cover of eight landscape classes (**Fig.

SA.3**) within buffers of 1 km radius, and used Shannon diversity index (SHDI) (Turner & Gardner, 2015)

to characterize compositional landscape heterogeneity from these landscape metrics. A Shannon's

diversity index was also estimated for crop landscape diversity within a buffer of 1 km, considering 17

types of crops (**Fig. SA.3**) retrieved from external databases (OSO, Graphical Parcel Register and the

database TOPO).

Our dataset contained missing values on some of the predictors (from 0 to 21.5% missing values at the

level of observations), which were completed with a multivariate imputation method, according to a

random forest algorithm (package $_{MICE}$ of R v.4.0.0, Buuren & Groothuis-Oudshoorn, 2011). As for

biodiversity data, we used mean values over the analysis period for all predictors for each plot; these

means by plot were not substantially changed by the imputation method (**Appendix B**).

For compositional analyses, all explanatory variables were standardized to have equal mean and

standard deviation. We limited the number of explanatory variables to seven (**Table 1**) so that we could

include the same ones at different resolutions (see section on data analysis below). We aimed to

represent the essential variables in different categories (climate, soil, landscape and practices) and

ensure they were not highly correlated (Pearson correlation coefficient < 0.6). We retained mean

annual temperature and soil pH, since they are tightly related to floristic composition and richness in

agroecosystems (Fried et al., 2008). Landscape and crop diversity were selected as indicators of

landscape heterogeneity (Rader et al., 2014; Sirami et al., 2019). Finally, we included the herbicide

treatment frequency index, the nitrogen dose applied in fertilizers (within adjacent agricultural fields)

and the number of margin management events. These variables have consistently been reported to

have a significant effect on field margin communities (Aavik & Liira, 2009; Bassa et al., 2011; Fried et

al., 2018).

*Data analysis*

To analyze the relationships between floristic species richness, composition and environment

(including agricultural practices) in relation to spatial scale, we built on two statistical methods, each

repeated at the national extent using different resolutions, and then at different spatial extents using

the plot resolution. We used Gaussian spatial simultaneous autoregressive models (SAR) with species

richness as response variable, to take into account spatial autocorrelation, and used generalized

dissimilarity models (GDM) to study species composition while dealing with nonlinear relationships

between environmental distance and species turnover (Ferrier et al., 2007).

SARs were implemented (package SPDEP, function *errorsarlm*) using a correlated spatial error term,

which is added as a predictor and is a function of the spatial neighborhood of the response variable

(**Appendix C**) (Cressie, 2015). We can thus model linear relationships that take into account spatial

autocorrelation in the data, i.e. the tendency of nearby points to have more similar values than

expected by chance. Hereafter, we will refer to spatial structuring as spatial autocorrelation, that can

result from an effect of the spatially structured environment or from an internal process of the

community (Borcard et al., 2018). Species richness was transformed with a square root to comply with

normality. We assessed the residual autocorrelation of each model with a Moran's I test (with a p-

value threshold of 0.05) (Thioulouse et al., 2018), based on the same spatial weight matrix included in

the models. None of the models displayed significant residual autocorrelation. In order to quantify the

importance of each variable and the spatial component, we computed partial regression using the

Nagelkerke pseudo-$R^2$ (Lichstein et al., 2002; **Appendix C**).

The use of GDMs (R package ₉ₐₘ, function *gdm*) to model species composition is appropriate when

species turnover rate varies along the environmental gradients and hence when the response is non-

linear. The Bray-Curtis dissimilarity among all pairs of plots was modeled as a function of

environmental pairwise distance (computed for each factor) and geographical distance (Ferrier et al.,

2007). The geographical distances account for the spatial structuring into communities. Since the same

plots are used multiple times in the calculation of biological and environmental distances, the

environmental table was permuted 500 times to assess variable significance. We then evaluated the

importance of each variable in the model with the same procedure as above (partial regressions based

on the explained deviance; Maestri et al., 2017).

The different analyses detailed in the above sections were reiterated at different spatial scales. The

first approach was to increase the study resolution by aggregating data into grid cells. We opted for

the following spatial resolutions: 25, 40, 60 and 75 km (**Fig. 1A, Fig SD.1**). Only cells containing at least

two plots were retained. Cells for which all plots were less than 5 km apart were also removed to avoid

autocorrelation issues that may lead to extreme values, as were cells in the land - sea borders with

more than half of their surface covered by water. Overall, this led to some analyses at finer resolutions

that contained less grid cells than coarser analyses. Since not all scales have the same number of grid

cells or plots by cell (**Table 2**), the potential influence of these variations was investigated by

resampling (**Appendix D**). Whenever possible, explanatory data were directly extracted at the scale of

interest (climate, soil and landscape). All other factors were averaged by cell, including the frequency

of occurrence by plot. Species richness in each grid cell was obtained by aggregating plot-level species

lists. The coordinates of the grid cell centers were used in all spatial analyses. In order to take into

consideration the arbitrary position of the grid on the map, we replicated analyses by shifting the grid

position three times to check for consistency in analyses.

Lastly, we proceeded with the same analysis at two different extents: national and regional. For this

purpose, plots were grouped within ten biogeographical regions standing for the wide pedo-climatic

and floristic gradients in France (**Fig. 1B**, based on the VégétalLocal map; Office français de la

biodiversité, 2021). The following four regions were dropped because the number of plots was too low

(<40): Alps, Corsica, Massif Central and Rhone-Saone-Jura basin.

**Results**

*Influence of agricultural and environmental variables on community structure at different spatial*

*resolutions*

At the plot-level resolution, the predictors explained 27.8% of the variance in species richness.

Herbicides and nitrogen dose were the main drivers and explained respectively 1.5 and 1.3% of the

total variance in species richness, both having a slight negative effect on richness. With respect to plant

community composition, the GDM explained 29.5% of deviance, the main drivers being mean annual

temperature (6.9%) and soil pH (1.2%).

With increasing spatial resolution, the relative importance of farming practices decreased in favor of

an increased influence of climate and landscape on richness (except at intermediate resolution, **Fig.**

**2A-E**). Temperature had a positive effect on richness, while soil pH had no significant influence. Crop

diversity was the predominant landscape factor at all aggregated scales, showing a consistent positive

influence on richness, but only significant at the coarsest resolutions (60-75 km, **Table 3A**). On the

contrary, landscape diversity did not seem to have any impact (**Table 3A**). Both herbicides and nitrogen

dose reduced the number of species at the plot-level, but these effects were not robust to the grid and

resolution studied (**Table 3A**). Similarly, the weak positive effect of margin management on species

richness was mainly apparent at the plot-resolution (**Table 3A**). Richness variation attributed to spatial

structure was higher at the plot resolution (11.4%) compared to coarser resolutions where it was non-

significant (**Fig. 2A-E**).

With regard to composition, the explanatory power of the models increased at coarser spatial grains

(**Fig. SE.1**). Climate and landscape (mainly through crop diversity) together contributed to overall

differences in species composition among plots, much more strongly than farming practices and soil

at all scales (**Fig. 2F-J**). In contrast to richness, climate had less influence on composition at coarser

resolutions while the influence of landscape increased. Fertilization structured the species distribution

more strongly at the coarsest resolution (4.3%) than at the finest, for which it was not significant (**Fig.**

**2F-J, Table 3B**). While herbicides decreased species richness, they did not influence species

composition. Also, we observed that the importance of spatial structure was relatively high at all

aggregated scales (**Fig. 2F-J**).

*Influence of agricultural and environmental variables on community structure at different spatial*

*extents*

The influence of environmental filters on species richness and composition varied greatly among

regions (**Fig. 3**). For example, the high explanatory power of the SAR for the Mediterranean zone (ZM)

(approximately 38.8%) contrasted with the lower one for the Armorican Massif (MA) (15.1%).

Some landscape effects emerged as more significant than at the national extent. For example,

landscape diversity had an important influence (6.5%) on richness of the northern Paris Basin (BPN),

an effect that was not visible at the national extent with all regions confounded (**Table 4A**). Agricultural

effects on richness were often more predominant at the regional extent (**Fig. 3A**). The effects of

nitrogen fertilization, herbicides and margin management were more clearly revealed at this scale. The

direction of the relationships between richness and predictors remained consistent with analyses at

the national level (except for the south-western France (ZSO), which depicted a negative relationship

of species richness with temperature, **Table 4A**).

With regard to composition, climatic conditions appeared to have a stronger influence at a national

rather than regional extent (**Fig. 3B**). As with richness, new impacts of landscape and practices were

highlighted within regions, such as the influence of landscape diversity on composition in ZM (2.2%),

and the number of margin management events in ZM (5.6%) and MA (6.9%) (**Fig. 3B**, **Table 4B**).

**Discussion**

In line with our initial hypotheses, the effects of agricultural practices on field margin flora, although

detected at all resolutions, were more prominent within biogeographic regions. In general terms,

herbicides and fertilization were important to explain species richness at the plot-level, whereas

fertilization stood out in explaining species composition at coarse resolutions. Conversely, climate and

landscape variables always predominated when data were aggregated at coarser resolutions and large

(i.e. national) extent. Climate also explained a large percentage of variation in species composition at

the plot-level. As the botanical diversity of field margins is currently declining, these results are of great

interest to protect these habitats, which sometimes represent the last reservoirs in the most intensive

landscapes (Marshall & Moonen, 2002).

*Influence of agricultural and environmental variables on community structure at different spatial*

*resolutions*

Our results provided evidence that composition was better explained by large-scale agricultural

effects, while richness was mostly impacted by local land use intensification. This is consistent with

some studies reporting that the effects of agricultural practices on field margin plant communities are

at least as important to landscape effects (Bassa et al., 2011; Gabriel et al., 2006; but see Martin et al.,

2020). The negative impacts of herbicides and nitrogen on local richness of field margins have already

been reported many times (Aavik & Liira, 2009; Fried et al., 2018). Here, nitrogen fertilization affected

both richness and composition, while herbicides only had an impact on richness. This is in agreement

with Fried et al. (2018), who reported that herbicides decreased species richness, but had no effect on

functional composition, as opposed to fertilization. Our results revealed a large-scale effect of nitrogen

use, previously evidenced on species richness of natural, semi-natural and agricultural habitats (Billeter

et al., 2008; Kleijn et al., 2009). Storkey et al. (2011) demonstrated that large-scale eutrophication is a

major explanatory factor of the threat status of arable species in Europe and even more than herbicide

use, although the two factors are difficult to disentangle and likely in interaction. However, the

importance of this effect on composition and not on richness contrasted with studies conducted so

far.

The fact that composition was better explained at coarser resolutions, while richness was not (**Fig.**

**SE.1**) may be explained by a distinct effect of the aggregation of plots in the grid cells. Composition can

be better explained at coarser resolutions because rare species, specialists of particular conditions, can

easily be missing at the plot-level but have higher probabilities to be present in larger grid cells that

combine several plots. Coarser resolutions also buffer stochastic effects related to the dispersal of

generalist species that can attenuate the weight of explanatory variables at the plot-level. However,

at the scale of coarser grid cells, richness may depend more on the variability (complementarity) of the

practices of each plot included in the cell than on their average values as computed here. While

considering only local agricultural effects may be sufficient to explain species richness of field margins,

a full understanding of community assembly will only be achieved by including the influence of species

pools and their responses to regional intensification.

Regarding landscape variables, crop diversity was the best predictive factor of richness and

composition and was particularly determinant at large-scales. While the response of plant species

richness to Shannon crop diversity is inconsistent among studies and dependent on the level of

intensification (Martin et al., 2020; Sirami et al., 2019), we found here a consistently positive effect

regardless of the scale. This can be viewed as landscape complementation for species that are

specialists of a single crop and not encountered in any other type of habitat (Storkey et al., 2011).

Another explanation would be that crop diversity is negatively related to field size (Martin et al., 2020),

and this latter might have an additional effect on richness (Fahrig et al., 2015). In contrast, our analysis

did not reveal any effects of habitat diversity in the landscape on richness, but an effect on composition

and only at the plot-level. Although a diverse landscape does not necessarily support more species, it

may favor functionally more diverse communities (dispersing from different neighboring habitats),

supporting the idea that species richness alone cannot capture all aspects of biodiversity (Aavik et al.,

2008; Fried et al., 2018). The effect of crop diversity on composition increased with resolution, while

this was not the case for landscape diversity, likely because landscapes became more homogeneous

in terms of habitat diversity, while they still remained different in terms of crops (**Fig. SF.1**).

*Influence of agricultural and environmental variables on community structure at different spatial*

*extents*

Our results pointed to strong regional specificities, underlining complex interactions between local and

regional scales. For example, the influence of landscape diversity on richness was very high in the BPN,

a region mainly composed of large cereal open-fields (**Fig. SF.2-3**). The direction of this effect was

unexpected, since landscape diversity decreased the number of species in field margins. It is possible

that diverse landscapes have fewer croplands and more natural habitats, so that species adapted to

agricultural disturbances are less prevalent in the landscape species pool. Given the high level of

intensification in this region, the species from natural habitats are too vulnerable to establish locally.

The discrepancy between the low level of landscape disturbance and the high level of local disturbance

ultimately results in a local loss of species. This apparently counter-intuitive finding could thus be

explained by interactions between the regional pool and local conditions (Cornell & Harrison, 2014;

Myers & Harms, 2009). Likewise, the detrimental effect of nitrogen fertilization on richness was only

significant in the Mediterranean Zone (ZM). In this region, soils had low nitrogen content (**Fig. SF.3**),

and the species pool is mostly composed of stress-tolerant species adapted to nutrient-poor soil

conditions. Due to its regional composition, Mediterranean flora would thus be more sensitive to soil

enrichment. This supports the results of Kleijn et al. (2009) who concluded that the impact of nitrogen

is greater in less intensive regions. Biogeographic regions differed not only in their agricultural impacts

but also in the diversity of their regional pools. ZM and ZSO were slightly more diverse than MA (**Fig.**

**SF.3**), which reflected in their percentages of total explained richness. When species richness is higher,

species turnover between plots also has the potential to be higher, thereby providing new insights into

some agricultural effects. Clarifying regional specificities should thus be a strong priority for future

agroecological studies, as it paves the way for a deeper comprehension of interactions among scales,

environmental and agricultural factors.

*Limits and implications for upcoming studies and management in agricultural context*

Our study presented a number of shortcomings that needed to be considered. First, at coarser

resolutions and at the regional extent, the number of units available for analysis decreased with

respect to the plot-level at the national scale. This constrained our analysis to a limited number of

variables, forcing us to prioritize over other factors that are also recognized to have an influence on

field margin vegetation (e.g. rainfall, soil texture, Fried et al., 2008). Nevertheless, our purpose was

not to determine the most influential descriptors at all scales, but rather to assess the congruence of

the results by considering the same set of variables at different spatial scales. Furthermore, we tested

the bias that emerged from averaging over coarser scales on different number of plots by cell by

bootstrapping and observed that it caused little variation of the model parameters (**Fig. SD.2**). We are

therefore fairly confident that our results are robust in this respect. Another potential shortcoming

was that the means computed within aggregated grid cells might not fully represent the regional pool

or agricultural region. Indeed, the monitoring plots were stratified to represent the main crops and

practices at the national level and not at the landscape or regional level. Checking the

representativeness of the plots with the regional context is thus key to assess the relevance of the

results.

Our results have not only methodological but also practical implications for crop management. It

turned out that crop diversity had a major positive impact on plant diversity in field margins, whose

importance in public policies has just begun to be recognized (Galán-Martín et al., 2015). Promoting a

wide range of crops at the landscape level could slow the decline of some highly specialized arable

plant species, particularly in landscapes with large semi-natural cover including field margins (Sirami

et al., 2019). In this respect, the obligation of crop rotation as part of the enhanced conditionality of

subsidies in the new CAP in Europe can be seen as very beneficial for field margins. Fertilization

significantly altered floristic assemblages and is more impactful on a large scale than herbicides, which

is of major interest for national agricultural policies. Finally, measures should also be adapted to each

region. Indeed, our findings suggest that some decisions could be very costly in some regions and

provide little benefit, while less effort would yield better outcomes in another region (Kleijn et al.,

2009). For instance, a drastic reduction of fertilizers in highly intensive regions mainly composed of

nitrophilous species will have little impact on plant communities of field margins if this action is not

coupled with landscape restoration. In contrast, diversified areas will be more susceptible to respond

to less costly mitigation measures and are of higher conservation concern (Stevens et al., 2010). In

conclusion, we encourage agricultural stakeholders and scientists involved in monitoring programs to

further evaluate the changes in field margin flora and its protection measures across different spatial

scales, and implement regulatory measures that incorporate landscape-level management planning

adapted to each biogeographic / agricultural region.

**Acknowledgements**

The 500-ENI network is developed by the French Ministry of Agriculture under the Ecophyto

framework with funding from the French Biodiversity Agency (Office Français de la Biodiversité). We

Funding: This work was supported by an INRAE-ANSES thesis fellowship and two Ecophyto II+ projects: GT STEP 500 ENI and GTP 500 ENI (OFB-21-1642).

**Appendix A-F. Supplementary data**

Supplementary data associated with this article can be found, in the online version, at XXXXX.

**References**

Aavik, T., Augenstein, I., Bailey, D., Herzog, F., Zobel, M., & Liira, J. (2008). What is the role of local landscape structure in the vegetation composition of field boundaries? *Applied Vegetation Science*, *11*(3), 375–386. https://doi.org/10.3170/2008-7-18486

Aavik, T., & Liira, J. (2009). Agrotolerant and high nature-value species—Plant biodiversity indicator groups in agroecosystems. *Ecological Indicators*, *9*(5), 892–901. https://doi.org/10.1016/j.ecolind.2008.10.006

Andrade, C., Villers, A., Balent, G., Bar-Hen, A., Chadoeuf, J., Cylly, D., Cluzeau, D., Fried, G., Guillocheau, S., Pillon, O., Porcher, E., Tressou, J., Yamada, O., Lenne, N., Jullien, J., & Monestiez, P. (2021). A real-world implementation of a nationwide, long-term monitoring program to assess the impact of agrochemicals and agricultural practices on biodiversity. *Ecology and Evolution*, *11*(9), 3771–3793. https://doi.org/10.1002/ece3.6459

Bassa, M., Boutin, C., Chamorro, L., & Sans, F. X. (2011). Effects of farming management and landscape heterogeneity on plant species composition of Mediterranean field boundaries. *Agriculture, Ecosystems & Environment*, *141*(3–4), 455–460. https://doi.org/10.1016/j.agee.2011.04.012

Berquer, A., Martin, O., & Gaba, S. (2021). Landscape Is the Main Driver of Weed Assemblages in Field

Margins but Is Outperformed by Crop Competition in Field Cores. *Plants*, *10*(10), 2131.

https://doi.org/10.3390/plants10102131

Billeter, R., Liira, J., Bailey, D., Bugter, R., Arens, P., Augenstein, I., Aviron, S., Baudry, J., Bukacek, R.,

Burel, F., Cerny, M., Blust, G. D., Cock, R. D., Diekötter, T., Dietz, H., Dirksen, J., Dormann, C.,

Durka, W., Frenzel, M., … Speelmans..., M. (2008). Indicators for Biodiversity in Agricultural

Landscapes: A Pan-European Study. *Journal of Applied Ecology*, *45*(1), 141–150.

https://doi.org/10.1111/j.1365-2664.2007.01393.x

Borcard, D., Gillet, F., & Legendre, P. (2018). *Numerical Ecology with R*. Springer.

Bourgeois, B., Gaba, S., Plumejeaud, C., & Bretagnolle, V. (2020). Weed diversity is driven by complex

interplay between multi-scale dispersal and local filtering. *Proceedings of the Royal Society B:*

*Biological Sciences*, *287*(1930), 20201118. https://doi.org/10.1098/rspb.2020.1118

Buuren, S. van, & Groothuis-Oudshoorn, K. (2011). mice: Multivariate Imputation by Chained Equations

in R. *Journal of Statistical Software*, *45*(1), 1–67. https://doi.org/10.18637/jss.v045.i03

Chase, J. M. (2014). Spatial scale resolves the niche versus neutral theory debate. *Journal of Vegetation*

*Science*, *25*(2), 319–322. https://doi.org/10.1111/jvs.12159

Concepción, E. D., Díaz, M., Kleijn, D., Báldi, A., Batáry, P., Clough, Y., Gabriel, D., Herzog, F., Holzschuh,

A., Knop, E., Marshall, E. J. P., Tscharntke, T., & Verhulst, J. (2012). Interactive effects of

landscape context constrain the effectiveness of local agri-environmental management:

Landscape constrains the effectiveness of local management. *Journal of Applied Ecology*, *49*(3),

695–705. https://doi.org/10.1111/j.1365-2664.2012.02131.x

Cornell, H. V., & Harrison, S. P. (2014). What Are Species Pools and When Are They Important? *Annual*

*Review of Ecology, Evolution, and Systematics*, *45*(1), 45–67. https://doi.org/10.1146/annurev-

ecolsys-120213-091759

Cressie, N. (2015). *Statistics for Spatial Data*. John Wiley & Sons.

Emmerson, M., Morales, M. B., Oñate, J. J., Batáry, P., Berendse, F., Liira, J., Aavik, T., Guerrero, I.,

Bommarco, R., Eggers, S., Pärt, T., Tscharntke, T., Weisser, W., Clement, L., & Bengtsson, J.

(2016). How Agricultural Intensification Affects Biodiversity and Ecosystem Services. In

*Advances in Ecological Research* (Vol. 55, pp. 43–97). Elsevier.

https://linkinghub.elsevier.com/retrieve/pii/S0065250416300204

Fahrig, L., Girard, J., Duro, D., Pasher, J., Smith, A., Javorek, S., King, D., Lindsay, K. F., Mitchell, S., &

Tischendorf, L. (2015). Farmlands with smaller crop fields have higher within-field biodiversity.

*Agriculture, Ecosystems & Environment*, *200*, 219–234.

https://doi.org/10.1016/j.agee.2014.11.018

Ferrier, S., Manion, G., Elith, J., & Richardson, K. (2007). Using generalized dissimilarity modelling to

analyse and predict patterns of beta diversity in regional biodiversity assessment. *Diversity and*

*Distributions*, *13*(3), 252–264. https://doi.org/10.1111/j.1472-4642.2007.00341.x

Fried, G., Norton, L. R., & Reboud, X. (2008). Environmental and management factors determining

weed species composition and diversity in France. *Agriculture, Ecosystems & Environment*,

*128*(1–2), 68–76. https://doi.org/10.1016/j.agee.2008.05.003

Fried, G., Villers, A., & Porcher, E. (2018). Assessing non-intended effects of farming practices on field

margin vegetation with a functional approach. *Agriculture, Ecosystems & Environment*, *261*,

33–44. https://doi.org/10.1016/j.agee.2018.03.021

Gabriel, D., Roschewitz, I., Tscharntke, T., & Thies, C. (2006). Beta diversity at different spatial scales:

Plant communities in organic and conventional agriculture. *Ecological Applications*, *16*(5),

2011–2021. https://doi.org/10.1890/1051-0761(2006)016[2011:BDADSS]2.0.CO;2

Galán-Martín, Á., Pozo, C., Guillén-Gosálbez, G., Antón Vallejo, A., & Jiménez Esteller, L. (2015). Multi-

stage linear programming model for optimizing cropping plan decisions under the new

Common Agricultural Policy. *Land Use Policy*, *48*, 515–524.

https://doi.org/10.1016/j.landusepol.2015.06.022

Guerrero, I., Carmona, C. P., Morales, M. B., Oñate, J. J., & Peco, B. (2014). Non-linear responses of

functional diversity and redundancy to agricultural intensification at the field scale in

Mediterranean arable plant communities. *Agriculture, Ecosystems & Environment*, *195*, 36–

43. https://doi.org/10.1016/j.agee.2014.05.021

Haddaway, N. R., Brown, C., Eales, J., Eggers, S., Josefsson, J., Kronvang, B., Randall, N. P., & Uusi-

Kämppä, J. (2018). The multifunctional roles of vegetated strips around and within agricultural

fields. *Environmental Evidence*, *7*(1), 14. https://doi.org/10.1186/s13750-018-0126-2

Hengl, T., Jesus, J. M. de, MacMillan, R. A., Batjes, N. H., Heuvelink, G. B. M., Ribeiro, E., Samuel-Rosa,

A., Kempen, B., Leenaars, J. G. B., Walsh, M. G., & Gonzalez, M. R. (2014). SoilGrids1km—Global

Soil   Information   Based   on   Automated   Mapping.   *PLOS   ONE*,   *9*(8),   e105992.

https://doi.org/10.1371/journal.pone.0105992

Inglada, J., Vincent, A., Arias, M., Tardy, B., Morin, D., & Rodes, I. (2017). Operational High Resolution

Land Cover Map Production at the Country Scale Using Satellite Image Time Series. *Remote

Sensing*, *9*(1), 95. https://doi.org/10.3390/rs9010095

Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N. E., Linder,

H. P., & Kessler, M. (2017). Climatologies at high resolution for the earth's land surface areas.

*Scientific Data*, *4*(1), 170122. https://doi.org/10.1038/sdata.2017.122

Kleijn, D., Kohler, F., Báldi, A., Batáry, P., Concepción, E. d, Clough, Y., Díaz, M., Gabriel, D., Holzschuh,

A., Knop, E., Kovács, A., Marshall, E. j. p, Tscharntke, T., & Verhulst, J. (2009). On the

relationship between farmland biodiversity and land-use intensity in Europe. *Proceedings of

the   Royal   Society   B:   Biological   Sciences*,   *276*(1658),   903–909.

https://doi.org/10.1098/rspb.2008.1509

Leenhardt, D., Angevin, F., Biarnès, A., Colbach, N., & Mignolet, C. (2010). Describing and locating

cropping systems on a regional scale. A review. *Agronomy for Sustainable Development*, *30*(1),

131–138. https://doi.org/10.1051/agro/2009002

Lichstein, J. W., Simons, T. R., Shriner, S. A., & Franzreb, K. E. (2002). Spatial autocorrelation and

autoregressive models in ecology. *Ecological Monographs*, *72*(3), 19.

https://doi.org/10.1890/0012-9615(2002)072[0445:SAAAMI]2.0.CO;2

Maestri, R., Shenbrot, G. I., & Krasnov, B. R. (2017). Parasite beta diversity, host beta diversity and

environment: Application of two approaches to reveal patterns of flea species turnover in

Mongolia. *Journal of Biogeography*, *44*(8), 1880–1890. https://doi.org/10.1111/jbi.13025

Marshall, E. J. P., & Moonen, A. C. (2002). Field margins in northern Europe: Their functions and

interactions with agriculture. *Agriculture, Ecosystems & Environment*, *89*(1–2), 5–21.

https://doi.org/10.1016/S0167-8809(01)00315-2

Martin, A. E., Collins, S. J., Crowe, S., Girard, J., Naujokaitis-Lewis, I., Smith, A. C., Lindsay, K., Mitchell,

S., & Fahrig, L. (2020). Effects of farmland heterogeneity on biodiversity are similar to—Or even

larger than—The effects of farming practices. *Agriculture, Ecosystems & Environment*, *288*,

106698. https://doi.org/10.1016/j.agee.2019.106698

Matthews, A. (2013). Greening agricultural payments in the EU's Common Agricultural Policy. *Bio-*

*Based and Applied Economics*, *2*(1), 1–27.

Myers, J. A., & Harms, K. E. (2009). Seed arrival, ecological filters, and plant species richness: A meta-

analysis. *Ecology Letters*, *12*(11), 1250–1260. https://doi.org/10.1111/j.1461-

0248.2009.01373.x

Nascimbene, J., Marini, L., & Paoletti, M. G. (2012). Organic Farming Benefits Local Plant Diversity in

Vineyard Farms Located in Intensive Agricultural Landscapes. *Environmental Management*,

*49*(5), 1054–1060. https://doi.org/10.1007/s00267-012-9834-5

Office français de la biodiversité. (2021). *Carte des régions d'origine—Référentiel technique de la*

*marque Végétal local—Version 2021*. www.vegetal-local.fr

Rader, R., Birkhofer, K., Schmucki, R., Smith, H. G., Stjernman, M., & Lindborg, R. (2014). Organic

farming and heterogeneous landscapes positively affect different measures of plant diversity.

*Journal of Applied Ecology*, *51*(6), 1544–1553. https://doi.org/10.1111/1365-2664.12344

Richard-Schott, F. (2009). Le Recensement Général de l'Agriculture de 1955, une référence pour les

géographes ? *Géocarrefour*, *2009*(4), Article Vol. 84/4.

https://doi.org/10.4000/geocarrefour.7567

Sirami, C., Gross, N., Baillod, A. B., Bertrand, C., Carrié, R., Hass, A., Henckel, L., Miguet, P., Vuillot, C.,

Alignier, A., Girard, J., Batáry, P., Clough, Y., Violle, C., Giralt, D., Bota, G., Badenhausser, I.,

Lefebvre, G., Gauffre, B., … Fahrig, L. (2019). Increasing crop heterogeneity enhances

multitrophic diversity across agricultural regions. *Proceedings of the National Academy of*

*Sciences*, *116*(33), 16442–16447. https://doi.org/10.1073/pnas.1906419116

Stevens, C. J., Duprè, C., Dorland, E., Gaudnik, C., Gowing, D. J. G., Bleeker, A., Diekmann, M., Alard, D.,

Bobbink, R., Fowler, D., Corcket, E., Mountford, J. O., Vandvik, V., Aarrestad, P. A., Muller, S.,

& Dise, N. B. (2010). Nitrogen deposition threatens species richness of grasslands across

Europe. *Environmental Pollution*, *158*(9), 2940–2945.

https://doi.org/10.1016/j.envpol.2010.06.006

Storkey, J., Meyer, S., Still, K. S., & Leuschner, C. (2011). The impact of agricultural intensification and

land-use change on the European arable flora. *Proceedings of the Royal Society B: Biological*

*Sciences*, *279*(1732), 1421–1429. https://doi.org/10.1098/rspb.2011.1686

Thioulouse, J., Dray, S., Dufour, A.-B., Siberchicot, A., Jombart, T., & Pavoine, S. (2018). Analysing

Spatial Structures. In J. Thioulouse, S. Dray, A.-B. Dufour, A. Siberchicot, T. Jombart, & S.

Pavoine (Eds.), *Multivariate Analysis of Ecological Data with ade4* (pp. 239–260). Springer.

https://doi.org/10.1007/978-1-4939-8850-1_12

Tscharntke, T., Klein, A., Kruess, A., Steffan-Dewenter, I., & Thies, C. (2005). Landscape perspectives on

agricultural intensification and biodiversity—Ecosystem service management. *Ecology Letters*,

*8*(8), 857–874. https://doi.org/10.1111/j.1461-0248.2005.00782.x

Turner, M. G., & Gardner, R. H. (2015). Landscape Metrics. In M. G. Turner & R. H. Gardner (Eds.),

*Landscape Ecology in Theory and Practice: Pattern and Process* (pp. 97–142). Springer.

https://doi.org/10.1007/978-1-4939-2794-4_4

Viana, D. S., & Chase, J. M. (2019). Spatial scale modulates the inference of metacommunity assembly

processes. *Ecology*, *100*(2), e02576. https://doi.org/10.1002/ecy.2576

Viana, D. S., Figuerola, J., Schwenk, K., Manca, M., Hobæk, A., Mjelde, M., Preston, C. D., Gornall, R. J.,

Croft, J. M., King, R. A., Green, A. J., & Santamaría, L. (2016). Assembly mechanisms

determining high species turnover in aquatic communities over regional and continental

scales. *Ecography*, *39*(3), 281–288. https://doi.org/10.1111/ecog.01231

Woodcock, B. A., Bullock, J. M., McCracken, M., Chapman, R. E., Ball, S. L., Edwards, M. E., Nowakowski,

M., & Pywell, R. F. (2016). Spill-over of pest control and pollination services into arable crops.

*Agriculture,        Ecosystems        &        Environment*,        *231*,        15–23.

https://doi.org/10.1016/j.agee.2016.06.023

Zamorano, J., Bartomeus, I., Grez, A. A., & Garibaldi, L. A. (2020). Field margin floral enhancements

increase pollinator diversity at the field edge but show no consistent spillover into the crop

field: A meta-analysis. *Insect Conservation and Diversity*, *13*(6), 519–531.

https://doi.org/10.1111/icad.12454

**Figures**

**Fig. 1.** Distribution maps of the field margins (colored points) surveyed in France distributed in (A) a

grid of 25 km and (B) in different biogeographic regions. Only cells selected for analyses are presented

in A). A = « Alps », BPN = « Northern Paris Basin », BPS = « Southern Paris Basin », BRSJ = « Rhone-

Saone-Jura Basin », C = « Corsica », MA = « Armorican Massif », MC = « Massif Central », ZM =

« Mediterranean Zone », ZNE = « North-Eastern France » and ZSO = « South-Western France ». The

regions C, A, MC and BRSJ have been removed because of their small number of plots.

**Fig. 2.** Percentage of explained variance of species richness (SAR) and composition (GDM) by each factor for each spatial resolution. (A, F) 0 km; (B, G) 25 km; (C, H) 40 km; (D, I) 60 km; (E, J) 75 km. Standard deviations are computed from the three resamplings of grids. Percentages of explained variance are reported in **Table SE.1**.

**Fig. 3.** Percentage of explained variance for (A) richness (SAR) and (B) composition (GDM) for each biogeographic region and national extent. Regions are ordered by area from left to right. Percentages of explained variance are reported in **Table SE.2**.

**Tables**

576 **Table 1.** Description of explanatory factors used in analyses, with their abbreviations and scale of

577 computation. The variables computed by observation and then averaged by plot are underlined.

| Factors | Abbreviations | Units | Calculation | Type of factor | Data source | Scale of computation |
|---|---|---|---|---|---|---|
| **Mean annual temperature** | TEMP | °C x 10 | | Climate | Chelsa Climate | 1 km |
| **Soil pH (water)** | PH | pH x 10 | | Soil | SoilGrids | 250 m |
| **Shannon's habitat diversity index** | SHDI | | $-\sum_{i=1}^{n} p_i \ln p_i$<br>With $p_i$ the proportion of the habitat i in the landscape | Landscape | OSO | 1 km |
| **<u>Shannon's crop diversity index</u>** | SHDI_C | | $-\sum_{i=1}^{n} p_i \ln p_i$<br>With $p_i$ the proportion of crop i in the agricultural area | Landscape | RPG | 1 km |
| **<u>Dose of nitrogen (fertilization and amendments)</u>** | N_DOSE | kg/ha | $\dfrac{Quantity \times N\ of\ the\ formulation}{100}$ | Agricultural practices | 500-ENI | Plot |
| **<u>Treatment Frequency Index of herbicides</u>** | HERB | | $\dfrac{Applied\ dose\ of\ used\ products \times treated\ area}{certified\ dose \times plot\ area}$<br>For herbicides in vineyards, the treated area accounts for one third of the plot area if the treatment is done on the rows and two thirds if it is done on the inter-rows. In case of missing information, the treatment is supposed to involves only the rows (i.e. the most frequent practice). | Agricultural practices | 500-ENI | Plot |
| **<u>Number of management events</u>** | MAN | | All types of management (mowing, grazing…) | Margin management | 500-ENI | Field margin |
| **Spatial structure** | SPAT | m (Lambert 93) | Based on geographical coordinates, see **Appendix C** for more details | Spatial | 500-ENI | Plot |

579  **Table 2.** Details concerning the sampling for each resolution. Means and standard deviations were

580  computed based on the three positions of grids at each resolution.

| Spatial resolution (km) | 0 | 25 | 40 | 60 | 75 |
|---|---|---|---|---|---|
| **Mean number of plots by cell** (rounded to the hundredth) | | 2,68 ± 0,07 | 3,30 ± 0,09 | 4,63 ± 0,09 | 5,78 ± 0,07 |
| **Mean number of plots** (rounded to the unit) | 462 | 306 ± 10 | 387 ± 9 | 418 ± 12 | 428 ± 1 |
| **Mean number of cells** (rounded to the unit) | | 87 ± 7 | 105 ± 3 | 86± 2 | 71 ± 2 |

582 **Table 3.** Significance of each predictor (in row) for each spatial resolution (in column, in km). For

583 species richness in A), the direction of the relationship is indicated by +/-. The color gradient indicates

584 the number of times the predictor was significant over the three grid shifts.

585 **A) Species richness**

| | 0 | 25 | 40 | 60 | 75 |
|---|---|---|---|---|---|
| **Temperature** | + | + | + | + | + |
| **Ph** | + | - | - | - | - |
| **Landscape diversity** | - | - | - | - | - |
| **Crop diversity** | + | + | + | + | + |
| **Herbicides** | - | - | - | - | - |
| **Nitrogen dose** | - | - | - | + | - |
| **Margin management** | + | + | + | + | + |
| **Spatial structure** | | | | | |

Legend:
- Significant in all grids
- Significant in 2 grids
- Significant in 1 grid
- Never significant

588 **B) Composition**

| | 0 | 25 | 40 | 60 | 75 |
|---|---|---|---|---|---|
| **Temperature** | | | | | |
| **Ph** | | | | | |
| **Landscape diversity** | | | | | |
| **Crop diversity** | | | | | |
| **Herbicides** | | | | | |
| **Nitrogen dose** | | | | | |
| **Margin management** | | | | | |
| **Spatial structure** | | | | | |

**Table 4.** Significance of each predictor (in row) for each biogeographic region and for national extent (in column, see **Fig. 1** for abbreviations of region names). For species richness in A), the direction of the relationship is indicated by +/-. Cells are colored in green when the relationship is significant.

**A) Species richness**

| | BPN | BPS | MA | ZM | ZNE | ZSO | France |
|---|---|---|---|---|---|---|---|
| Temperature | + | + | + | - | + | - *(green)* | + *(green)* |
| Ph | - | + | - | + | + | - *(green)* | + |
| Landscape diversity | - *(green)* | - | - | - | + | + | - |
| Crop diversity | + | + | + | + *(green)* | + | + | + |
| Herbicides | - *(green)* | - | - | - | - | - | - *(green)* |
| Nitrogen dose | - | - | - | - *(green)* | - | - | - *(green)* |
| Margin management | + *(green)* | + | - | + *(green)* | + | + *(green)* | + *(green)* |
| Spatial structure | | | | | *(green)* | *(green)* | *(green)* |

**B) Composition**

| | BPN | BPS | MA | ZM | ZNE | ZSO | France |
|---|---|---|---|---|---|---|---|
| Temperature | | | | *(green)* | | | *(green)* |
| Ph | | | | | | | *(green)* |
| Landscape diversity | | | | *(green)* | | | *(green)* |
| Crop diversity | | | | *(green)* | | | *(green)* |
| Herbicides | | | | | | | |
| Nitrogen dose | | | | | | | |
| Margin management | | | *(green)* | *(green)* | *(green)* | | |
| Spatial structure | *(green)* | *(green)* | *(green)* | *(green)* | *(green)* | *(green)* | *(green)* |