Feedback control of organ size precision is mediated by *BMP2*-regulated apoptosis in the *Drosophila* eye

T. Navarro[1], A. Iannini[1], M. Neto[1,5], A. Campoy[1,2], Javier Muñoz[3], S. Ares[4]*, F. Casares [1]*

1 – CABD, CSIC/Universidad Pablo de Olavide. 41013 Seville, SPAIN.
2 – ALMIA, CABD, CSIC/Universidad Pablo de Olavide. 41013 Seville, SPAIN.
3 – Universidad Carlos III, 28911 Madrid, SPAIN.
4 – CNB, CSIC/UAM, 28049 Madrid, SPAIN.
5 – Present address: Instituto de Neurociencia, CSIC/Universidad Miguel Hernandez, 03550 Sant Joan d'Alacant, SPAIN.
* Corresponding authors: fcasfer@upo.es; saul.ares@csic.es

SUMMARY

Biological processes are intrinsically noisy and yet, the result of development –like the species-specific size and shape of organs– is usually remarkably precise. This precision suggests the existence of mechanisms of feedback control that ensure that deviations from a target size are minimized. Still, we have very limited understanding of how these mechanisms operate. Here, we investigate the problem of organ size precision using the *Drosophila* eye. The size of the adult eye depends on the rates at which eye progenitor cells grow and differentiate. We first find that the progenitor net growth rate results from the balance between their proliferation and apoptosis, with this latter contributing to determining both final eye size and its variability. In turn, apoptosis of progenitor cells is hampered by Dpp, a BMP2/4 signaling molecule transiently produced by early differentiating retinal cells. Our genetic and computational experiments show how the status of retinal differentiation is communicated to progenitors through the differentiation-dependent production of Dpp which, by adjusting the rate of apoptosis, exerts a feedback control over the net growth of progenitors to reduce final eye size variability.

INTRODUCTION

The size of some organs is remarkably precise (Boulan and Leopold, 2021). Although this precision is a functional need, it is not easily explained (Lander, 2011; Wolpert, 2010). Most organs are specified as small primordia whose growth is exponential. However, exponential growth is prone to amplify the noise that is inescapable to biological processes. This fact has led to the proposal that organ development must be subject to mechanisms of feedback control (Lander, 2011) which, in the engineering sense, reduce their internal "developmental instability" (Waddington, 1942) and increase the precision with which organs reach their species-specific size. Some organs that undergo constant tissue renewal, like the mammalian olfactory epithelium, maintain their size homeostasis through integral feedback (Lander et al., 2009). Some other organs, though, grow to a size that, once reached, cannot be further modulated. This is the case of insects. After their last molt, the adult organs cease growth and then become covered in a rigid chitin exoskeleton that precludes further size regulation. The eye of *Drosophila* fruitflies serves as a good example of this type of organ growth: In *Drosophila,* the end point of eye development is reached when all retinal progenitor

cells differentiate, as differentiation is accompanied by the exit of the cell cycle. Three
features should have resulted in a strong evolutionary pressure to maximize the
precision in eye size: first, size impacts vision directly, as image resolution and contrast
sensitivity is proportional to the number of light sensing units in the eye (Casares and
McGregor, 2020); second, making and maintaining the eyes is energetically very
expensive (Niven and Laughlin, 2008) so there is a need to match eye size to vision
needs; and third, left and right eyes must survey a symmetrical part of the space, so
eye asymmetry, which could be driven by developmental noise, should be minimized.
In this paper we investigate the mechanism by which the *Drosophila* eye ensures its
size precision as an example of organ size control.
57
The development of the *Drosophila* eye takes place as a wave of differentiation
crosses the eye primordium during the last larval stage (Figure 1a,b). As the wave
moves, proliferative progenitors anterior to the wave are recruited as differentiating
retinal cells that exit the cell cycle. Final eye size is attained when the differentiation
wave reaches the anterior-most region of the primordium: with no remaining
progenitors, which are the source of growth, the process ends. The movement of the
wave is driven primarily by two secreted signaling molecules. The first of them is Hh
(*hedgehog*). Hh's major role is the induction of differentiation of progenitor ("G") cell into
retinal ("R") cells (Heberlein and Moses, 1995), a role that is shared by its homologue
Shh in the differentiation of the ganglion cells of the vertebrate retina (Neumann and
Nuesslein-Volhard, 2000). Hh, initially expressed adjacent to the eye primordium,
contributes to initiate R differentiation. Once initiated, R cells produce Hh themselves
(Lee et al., 1992; Ma et al., 1993), resulting in the self-sustained progression of a
differentiation wave (Rogers et al., 2005). The second signal is a BMP2/4 homologue,
Dpp (*decapentaplegic*). *dpp* expression is induced by Hh in the newly-recruited retinal
cells ("Rn") (Chanut and Heberlein, 1995; Heberlein et al., 1993). Dpp, by repressing
the progenitor transcription factor Hth (*homothorax*) increases the propensity of
progenitors to enter differentiation (Bessa et al., 2002; Lopes and Casares, 2010).
However, and in contrast with Hh, whose expression is maintained in differentiated R
cells, Dpp is expressed only transiently: as newly recruited Rn cells progress along
their differentiation to Rs, *dpp* expression is turned-off (Blackman et al., 1991;
Heberlein et al., 1993; Masucci et al., 1990). Thus, a stripe of Dpp marks the front of
the differentiation wave (Figure1a). This process is schematically represented in Figure
1c.

Figure 1. The eye primordium ((a) "E", outlined) develops into the adult eye (b) through the action of a differentiation wave that sweeps across it (arrow). (a) eye primordium ("E") of a *Hh:GFP*; *dpp-Z* larva. The differentiation wavefront expresses *dpp-Z*. Posterior to it further differentiated photoreceptors lose *dpp* expression and express Hh. (b) Lateral view of an adult *Drosophila* eye. (c) The recruitment of proliferative progenitor cells ("G") into early differentiating (*dpp*-expressing) retinal cells ("Rn") and then into terminally differentiating (non-*dpp*-expressing) retinal cells ("R") is controlled by two signaling molecules, Hh and Dpp. (e-h) Frontal views of female heads of the indicated genotypes, with the eyes colored for better visualization. (e) Control, *optix>+*, (f) *optix>tkvRNAi,* (g) *optix>tkvRNAi + dadRNAi,* (h) *Bar (B) mutant*. See main text for details.

RESULTS

To ensure that the target, species-specific final size is reached with precision, the rates of growth and of differentiation need to be coordinated within the eye primordium (Casares and McGregor, 2020; Fried et al., 2016; Lobo-Cabrera et al., 2021). For example, a transient halt of the differentiation wave would produce some excess of progenitor cells and a concomitant deviation from the target size -unless this halt were communicated to the progenitors as a negative feedback on their rate of growth. This feedback could be implemented either by slowing down progenitor proliferation or by curving the excess of progenitor cells through cell death (we do not consider a change in cell size as this change would alter the size of the ommatidia and thus their optical properties). We noticed that, in wild-type control eye primordia, the apoptosis signal detected using an antibody against the activated form of caspase-3 is usually very low. However, if an antibody against the terminal caspase Dcp-1 (Song et al., 1997) is used instead, we detected a stripe of apoptotic cells just anterior to the differentiation wave-front (Suppl. Fig1. to Fig. 1). Likely, the activation of cas-3 is more brief than the expression of dcp-1 during the apoptotic process, making detection of apoptosis easier with Dcp-1 as a marker.

In any case, we were intrigued by this localized apoptosis happening in normal eye primordia and asked what its role could be. To answer this, we blocked apoptosis by driving, specifically anterior to the differentiation wave-front, a polycistronic RNAi against *reaper, hid* and *grim*, the three major pro-apoptotic genes in *Drosophila* (*optix>Reaper,Hid,Grim-RNAi,* abbreviated *optix>RHGRI* (Siegrist et al., 2010)). These flies showed two major phenotypic changes: first, the median eye size ("rE" = average intraindividual eye area relative to head area expressed as percentage) was larger than those of controls (*optix>+)* and, second, the distribution of eye sizes was also significantly broader (Figure 2a and Supp. table 1 to Figure 2). When we measured the apoptotic index of *optix>RHGRI* it was smaller than that of *optix>+* and very close to zero (Figure 3). The increased dispersion in the distribution of eye size is indicative of a loss in size precision. The fluctuating asymmetry index (FAi), which measures the disparity between left and right organs within the same individual, has also been frequently used as a measure of precision as it captures the sensitivity of the process to developmental noise (Parsons, 1992). The FAi is normally calculated as the absolute value of the normalized difference between the size of the left (L) and right (R) organs from the same individual. Here, we use the *signed* FAi ("sFAi") to have into account the sign of the asymmetry, $(L - R)/(L + R)$, which allows to compare the asymmetry across different genotypes based on the amplitude of the distribution of this metric for each of them (Figure 2b and Suppl. table 1 to Figure 2). In apoptosis-blocked individuals the variance of sFAi increased significantly as well. To control for potential non-specific effects on eye size and its variability of RNAi overexpression, we drove a "neutral" RNAi against the fluorescent protein Cherry (*optix>CherryRI*) and observed that it did not affect neither of the two parameters when compared to control (*optix>+*) or wild type (*Oregon-R*) flies (see Suppl. Fig. 1 to Figure 2).

Figure 2. Effects of different genetic manipulations on the distribution of eye size ("rE") (a) and signed Fluctuating Asymmetry index ("sFAi", b). Each violin plot includes the male (left) and female (right) distributions, which were not significantly different from each other in any genotype. The median of each distribution is indicated (solid and dashed line, male and female, respectively). The predicted rE median for each genotype and the statistical model from which they are derived can be found in Suppl. Table 1 to Figure2. The variances of each genotype were compared against the variance in the control group (*optix>+*) in rE and sFAi (Suppl. table_1 to Figure2). Eye area is measured relative to the total head area and expressed as a percentage in this and subsequent figures.

These results pointed to a role of developmental apoptosis in controlling eye size and its precision. But in order to be involved in such control, the apoptotic rate in progenitors should be coupled to the rate of eye differentiation. This communication between progenitors and differentiating cells might be exerted by either of the two major signals in eye development: Hh and Dpp.

Intuitively, if one of these two signals were to be used to convey feedback information to progenitors about the status of retinal differentiation, Dpp would seem a better candidate. Although both Hh and Dpp regulate progenitor gene expression (Greenwood and Struhl, 1999), Dpp, by being expressed transiently, can be better used as readout of the *rate* of retinal differentiation. In contrast, Hh is constantly produced in R cells, regardless of whether the wave is moving or not. If Dpp were used in a feedback control mechanism, it would be expected that the attenuation of its signaling pathway in progenitors should lead to a decrease in final eye size precision. In addition, prior work on *dpp* mutant alleles had shown an increase in apoptosis in the mutant eye primordia (Bryant, 1988; Masucci et al., 1990), reinforcing the idea that Dpp could be this coupling signal.

To test this point, we attenuated the Dpp pathway specifically in eye progenitors during development, by targeting the expression of an RNAi against the Dpp receptor *tkv* (*thickveins*) (Brummel et al., 1994) in these cells (*optix>tkvRI*). This treatment caused two effects: First, the median eye size was reduced (Figures 1e,f and 2a and Suppl. table1 to Figure 2). This effect -eye reduction- had been described previously for mutant alleles of *dpp* (Spencer et al., 1982) and explained as a consequence of Dpp's requirement in progenitor cell recruitment (Chanut and Heberlein, 1997; Greenwood and Struhl, 1999). The eye size reduction was concomitant to a dramatic increase of both the dispersion of the eye size distribution and of its sFAi (Figure 2b and Suppl. table1 to Figure 2), something expected if Dpp were involved in feedback control. To confirm that this phenotype resulted from the attenuation of the Dpp pathway, we simultaneously drove expression of RNAis against *tkv* and against *dad* (*daughters against-dpp*), a Smad molecule that acts as a feedback inhibitor of the pathway (Inoue et al., 1998; Tsuneizumi et al., 1997). In individuals of this genotype (*optix>tkvRI+DadRI*) the eyes almost recovered their control size (Figure 1e,f and 2a, Suppl. table1 to Figure 2). Similarly, the dispersion of eye size distribution as well as the sFAi decreased towards values closer to control, although not fully (Figure 2b, Suppl. table1 to Figure 2). The attenuation of the *tkv-RNAi* effect cannot be attributed to its weaker expression caused by the introduction of a second UAS sequence (*UAS-Dad-RNAi*) in the genotype (Suppl. Fig. 2 to Figure 2).

Next, we confirmed that attenuation of the Dpp pathway (*optix>tkvRI*) caused a dramatic quantitative increase in dcp-1 apoptotic signal (Figure 3 and Suppl. Table 1 to Figure 3), while the mitotic rate of progenitors remained unaffected (Suppl. Fig. 1 to Figure 3), and that this increase in apoptosis was rescued by coexpression of the Dad-RNAi (*optix>tkvRI+DadRI*) (Figure 3). The expression of *UAS-RHGRI* alone or in combination resulted in the reduction of apoptosis almost completely (Figure 3).

Figure 3. Eye size reduction and apoptosis induced by attenuation of the Dpp signaling pathway can be reverted by blocking Dad. Lateral views of adult eyes (a-c) and L3 eye discs, stained with Dcp-1 (and counterstained with rhodamine phalloidin, "actin") (a'-c') of the indicated genotypes. (d) Distribution of apoptosis intensity in L3 discs of the indicated genotypes. Blocking apoptosis driving the UAS-RHG-RNAi reduces the apoptosis caused by *tkv* attenuation (*optix>tkvRI + RHGRI*). The statistical analysis corresponding to these data can be found in Suppl. Table 1 to Figure 3.

Therefore, we expected that any perturbation affecting the coupling between retina and progenitor apoptosis would not be able to fully rescue the size precision observed in control flies (both the dispersion of the eye size distribution or the sFAi). This was the case in all the experiments in which the feedback between retina-produced Dpp and apoptosis was disrupted, either by artificially upregulating the Dpp pathway (expressing *dad-RNAi*) or by blocking apoptosis (expressing the RHG-RNAi) (Figures 2a,b and Suppl. table 1 to Figure 2).

To further substantiate our results, we decided to analyze the phenomenon of eye size control in another mutant situation in which the *dpp* pathway was affected. We chose the *Bar ("bar-eyed")* mutant. *Bar*-mutant eyes are reduced (Tice, 1914)(Figure 1e,h)

and this size reduction is associated with the control of *dpp* expression and an increased apoptosis during larval development (Kang et al., 2014). In *Bar* eye primordia we detected a fragmented *dpp* expression, monitored by the *dpp-Z* transcriptional reporter, and an increased dcp-1-positive apoptosis (Figure 4a,b).

Figure 4. Increased cell death and fragmented *dpp* expression in *Bar* discs. L3 *dpp-Z* discs (a,b), stained for apoptosis (dcp-1, green), *dpp* transcription (beta-galactosidase, red) and the retinal differentiation marker Elav (blue). (a) control ("+": Oregon-R) and (b) *Bar* mutant discs. The ellipse approximately outlines the eye region. "*a*" marks the antennal primordium.

Bar eyes are smaller and more variable (Figure 2a) as well as more asymmetric (Figure 2b) than control flies (see also Suppl. table 1 to Figure 2). Similarly to what happened in our previous experiments, in which the Dpp pathway was attenuated directly by targeting the Dpp receptor with an RNAi, either blocking apoptosis (*Bar; optix>RHGRI*) or derepressing the Dpp pathway (*Bar; optix>DadRI*) partly rescued eye size (Figure 2a) and reduced both eye dispersion (Figure 2a) and their fluctuating asymmetry (Figure 2b; Suppl. table 1 to Figure 2).

The results obtained so far indicated that a major role of Dpp during eye development was the regulation of developmental apoptosis. Losing this regulation, by either blocking apoptosis directly or by "disconnecting" retina and progenitors by the attenuation of Dpp signal (by *tkvRI* expression or in *Bar* mutants), led to a decreased size precision. Therefore, not only the size of the eye, but its precision is under the control of the Dpp pathway.

The fact that the effects on eye size and precision caused by *dpp* signaling attenuation were almost fully rescued by preventing apoptosis pointed to the regulation of this process as a key phenomenon. In order to explore further how the regulation of apoptosis impinges on eye size, we built and analyzed a mathematical model of the process. With this model, we first explored whether our description of the biological phenomenon is complete by testing if the model recapitulates the biological results when parameterized using experimental measurements.

Our model comprises G (progenitors), Rn (newly differentiated retinal cells, expressing *dpp*) and R (differentiating retinal cells, no longer expressing *dpp* but producing Hh) cells as variables, as well as their transformations: proliferation of progenitors ("self-

renewal": G → G + G), apoptosis of progenitors (G → Φ), initial differentiation of
progenitors into *dpp*-expressing early retinal cells (G → Rn) and final differentiation of
these cells into *hh*-expressing photoreceptors (Rn → R) (Figure 5a,b), and includes
noise in all four differentiation steps. Its full description is available as Supplementary
material.
Conceptually, our model expresses the rates at which the G, Rn and R cells change
over time as in Figure 5c, where the variables influencing each of the steps (e.g.
"proliferation") are within the parentheses (e.g. "G, Rn").
256
Figure 5. Overview of the model of the G→Rn→R including apoptosis (Φ). (a) Schematics
of the process. R cells stimulate the differentiation process (by expressing Hh). Rn cells
negatively regulate apoptosis (by producing Dpp). G cells can divide, die or progress further into
differentiation by transiting to Rn. (b) This process, schematized in space. MF: morphogenetic
furrow, is the differentiation wavefront that moves (arrow) from posterior (right) to anterior (left).
L_A, L_P are the maximum distances anterior and posterior to the MF, respectively. The L_p can be
used as a proxy for developmental time: as the MF moves with constant speed, L_p increases as
time passes. L_A corresponds to G cells. L_p comprises Rn (orange bar) + R cells. (c) "Qualitative"
model. (d) Typical temporal profiles of all populations as predicted by the model. (e) The model
fits experimental data (L_{AP} as minor axis of elliptical fits of eye primordia (blue points) and L_A =
L_{AP} -L_p (orange points)) very well (see main text and supplementary material for details).

268
The key processes are modeled as follows: apoptosis is a decreasing function of the

Dpp-producing Rn cells, because the more Dpp the less apoptosis, and is proportional
to the number of progenitors (G), such that $apoptosis = a(Rn)G$. All G cells not
entering apoptosis will be proliferating with $proliferation = 1 - a(Rn)G$. As for the rate
of G to Rn differentiation ("early differentiation") we have considered two factors. First,
the activation of differentiation would depend on an increasing function of the Hh-
producing R cells, $h(R)$. Second, the induction of G cells to differentiate is limited to the
range of action of Dpp produced by Rn. Therefore, since beyond a certain point having
more G cells will not result in increased differentiation (that is, those G cells out of the
range of action of the signals will not be recruited), we modeled the function describing
G growth as a saturating sigmoidal function $f(G)$. Imposing $f(0) = 0$, this function also
ensures that once the G population has been exhausted, no further differentiation can
occur. Thus, $initial\ differentiation = f(G)h(R)$. Finally, the final differentiation, or
transition from Rn into R, can be expressed as being proportional to Rn, as in
$final\ differentiation = b * h(R)Rn$, where b is a constant. With this expression we
indicate that *Rn* differentiates "automatically" into R cells with a constant pace,
simplifying a complex process of successive cell induction leading to the formation of
the ommatidial cell types (Roignant and Treisman, 2009). However, eye size does not
depend on this process, as the number of ommatidia and therefore, the size of the eye,
is directly related to the number of G cells that are recruited as Rn cells. The
expressions of the relevant functions $a(Rn), f(G)$ and $h(R)$ can be found in the full
description on the model in the supplementary material. These functions allow us to
model the RNAi treatments as well.
292
With this model, we can calculate the evolution of all these cell populations over time
(Figure 5d). First, we fitted the model to an experimentally determined growth curve of
the eye primordium (Figure 5e; data from (Vollmer et al., 2016)). Since the
differentiation velocity is about constant during most of development (Vollmer et al.,
2016; Wartlick et al., 2014) the width of the R + Rn domain (the distance between the
posterior margin of the primordium to the differentiation wave-front, called "posterior
length", or "Lp") is a proxy for elapsed developmental time, and we used it as a time
axis (as in (Vollmer et al., 2016)). The fitting demanded that the growth rate of G cells
declined over developmental time, something that had been shown experimentally
(Vollmer et al., 2017; Wartlick et al., 2014). With these parameters, the fitting of the
model to the data was very good, indicating that the model captured the biology
underneath the control of eye size and its variability (Figure 5e and Suppl. Fig. 1 to
Figure 5).
Since the model was able to recapitulate well the dynamics of eye differentiation, we
asked which conditions would allow the system to reach robustly a final size. This can
be done by carrying out a linear stability analysis of the system. The "set point" that
needs to be stable is the final eye size, R*. Note that this analysis says nothing about
what the actual final eye size R* is (this depends on the specific rates and initial
conditions), just how stable it is. The finalization of the process is attained when G=0
and Rn=0: without any G and Rn cells left, the eye has reached R*. This R* state is
unstable, though. Any fluctuation (noise) in the number of G or Rn cells would alter the
value of R*. However, the linear stability analysis (see Linear stability analysis of the
deterministic model in supplementary materials) indicates that there is a factor that

strongly favors stability: the "basal apoptotic probability", a_0, which is the G cells'
tendency to enter apoptosis in the absence of Dpp signaling, with high apoptotic rates
favoring stability, since apoptosis promotes the depletion of the G pool and therefore
the termination of growth. Specifically, with an a_0 = 0,4 (i.e. 40% of apoptosis
determined experimentally in *optix>tkvRI*), this apoptosis rate would be close to
ensuring the R* stability condition on its own (see supplementary Linear stability
analysis). Therefore, high propensity of apoptosis in progenitor G cells seems to be a
requirement for robust size. Next, we used our model to predict the effects that noise
would have on eye size in different genotypes with different rates of apoptosis.

We reasoned at the onset that regulating organ size would rely on the differentiating
retina feeding information about its status on progenitor growth rates (something we
have shown to occur through the modulation of apoptosis). Therefore, we simulated the
effects of increasing the noise in the recruitment of Rn cells ("initial differentiation"),
which are the source of Dpp, on eye size and its variation in three genotypes: two in
which apoptosis is possible (control (*optix>+*) and *dpp* signaling attenuated (*optix> tkv-*
RNAi)) and one in which apoptosis is blocked (*optix>RHG-RNAi*) (Figure 6). While
increasing noise in Rn affects very little the MEDIAN eye size in the two genotypes
where apoptosis is allowed, it results in a steady increase of eye size if apoptosis is
blocked (until the noise levels are very high, when allowing apoptosis or not becomes
irrelevant). This computational result reinforces the idea that the *dpp*-sensitive
apoptosis of progenitors is instrumental in maintaining the developmental stability of
eye size.

Figure 6. Effects of increasing noise strength on median eye size (rE). If apoptosis is
blocked (by setting to zero the apoptotic rate of progenitors in the computational model,
equivalent to *optix>RHG-RI*) eye size increases more rapidly than in the other two genotypes, in
which apoptosis is allowed, until noise is too high (beyond 10^2), point at which allowing
apoptosis or not becomes irrelevant. Computational results. Data for each point is from 200
simulations *per* genotype.

DISCUSSION

For organs to attain a final size with low variability, mechanisms of control must exist to reduce the effects of developmental noise. In this paper, we describe one such mechanism, in which the processes of eye growth and differentiation are coupled through a feedback control. In this feedback, the discrepancies between the rates of growth and differentiation of progenitors are adjusted through the modulation of the progenitor cells' propensity to enter apoptosis, so that any production of progenitors in excess is curved, as a proportion of these progenitors will not receive sufficient Dpp signaling to overcome apoptosis. In this way, it is the Dpp production which, acting as a surrogate of the process of differentiation, feeds back onto the growth rate of progenitors, a rate that depends on their combined proliferation and apoptosis (this latter Dpp-signaling dependent). When analyzing our data, comparing the actual FAi versus the fluctuating asymmetry of randomized pairs generated from the same experimental data (FAn), we observed that FAn was slightly larger than FAi, and this difference (FAn>FAi) was invariant across genotypes. This indicated a small, but consistent effect of the individual. Therefore, although most variation in size can be attributed to organ-autonomous variation, the two eyes from an individual are slightly correlated (Suppl. Fig. 3 to Figure 2). This is expected though, as it has been shown that, besides organ-autonomous mechanisms of growth control, the growth discrepancies among organs in an individual affect the systemic growth of the organism through neuroendocrine communication mediated by insulin-like peptides (Colombani et al., 2012; Garelli et al., 2012). In addition, each larva might be exposed to small microenvironmental variations during its growth (Morgante et al., 2015). Still, our results indicate that the feedback mechanism we have described operates mostly in an organ-autonomous manner.

Our experiments show that a major role of *dpp* signaling during eye development is the regulation of apoptosis, since the effects on eye size caused by attenuating this pathway are almost fully recovered just by blocking it. This role is very different from the function in cell differentiation that has normally been attributed to *dpp* during eye development (Chanut and Heberlein, 1997). We note, however, that in our experiments, by using RNAi, we are attenuating, rather than fully abrogating, *dpp* signaling. This could mean that the low levels of signaling presumably remaining could still participate in cell fate specification. This would be similar to the situation found in the wing disc, with different signaling thresholds being important for distinct functions, with high *dpp* signaling being necessary for wing patterning, while lower levels sufficing for tissue growth (Barrio and Milan, 2017; Bosch et al., 2017; Matsuda and Affolter, 2017). Recently, the Kumar laboratory has shown that the main role of the Pax6 gene *eyeless (ey)*, a gene essential for normal eye development (Hunt, 1971), is the activation of *dpp* transcription (Weasner and Kumar, 2022). Interestingly, loss of *ey* function causes abundant cell death in eye progenitors (Zhu et al., 2017). The role for the *dpp* pathway in preventing progenitor cell death we have discovered would, in principle, connect *ey* function to cell death. At present we do not have a mechanistic explanation for why eye progenitor cells are prone to die (a tendency akin to the "death by default" state of cells proposed by Raff (Raff, 1992)), or how *dpp* signaling prevents it, but these are very relevant questions worth pursuing.

393
Is this control mechanism, based on regulating the rate of cell death of proliferating
progenitors, working beyond the eyes? Although not tested formally, the result from
Neto-Silva and co-workers, showing that wing asymmetry increased dramatically in
mutants for the apoptosis regulator *hid* (Neto-Silva et al., 2009), suggests that
apoptosis regulation may be involved in the control of developmental stability also of
wings too. Beyond *Drosophila,* developmental apoptosis is very frequent (Fuchs and
Steller, 2011) and, in some cases, associated to the regulation of cell number, with the
mammalian immune cells (Rathmel, 10.1111/j.0105-2896.2004.0096.x) and the
nervous systems (Raff et al., 1993) being striking examples. Therefore, it seems very
possible to us that mechanisms similar to the one we have described here will use the
feedback modulation of apoptosis as a means not only to controlling organ size, but
also size robustness in other developmental systems as well.
406
407
MATERIALS AND METHODS
RESOURCE AVAILABILITY
Lead contact: Further information and requests for resources and reagents should be
directed to and will be fulfilled by the lead contacts, Fernando Casares
(fcasfer@upo.es) and Saul Ares (saul.ares@csic.es).
Materials availability: this study did not generate new unique reagents.
Data and code availability: images (as .tiff flies) and quantifications of all genotypes
analyzed that might be required to reanalyze the data reported in this paper are
available from the lead contacts upon request. The macros for the quantification of
apoptosis/proliferation signal from confocal images is provided as supplementary
material. Any additional information required to reanalyze the data
reported in this paper is available from the lead contacts upon request.
420
METHOD DETAILS
***Drosophila* strains and genetic manipulations**
The *optix2/3-GAL4* driver (abbreviated "*optix>*") is expressed in
undifferentiated eye progenitors (Neto et al., 2017). Other strains used were: *UAS-*
tkvRNAi (VDRC #102319), *UAS-DadRNAi* (VDRC #108340), *UAS-Dad* (Tsuneizumi et
al., 1997) and *UAS-RHGRNAi* (Siegrist et al., 2010). The *optix2/3-GAL4, UAS-*
tkvRNAi/CyO recombinant strain was generated by standard genetic methods. Other
strains used were the *dpp* transcriptional reporter *dpp-Z* (Masucci and Hoffmann,
1993), the *Bar* mutant on the FM7c balancer chromosome (Flybase: FBba0000009)
and the wild type strain *Oregon-R (Or-R)*. The *UAS-mCherry* RNAi line (Bloomington
stock centre, reference BDSC_35785) was used as a neutral UAS-RNAi line.
432
Immunofluorescence and confocal imaging:
Fixation and immunofluorescence of eye primordia was done as in (Neto et al., 2016).
Imaging was carried out on a Leica SPE confocal setup. Images were then processed
using ImageJ. Primary antibodies used were: rabbit *anti*-phospho-histone H3 (pH3)
used at 1:1000 (Sigma), rabbit *anti*-Dcp 1 (Dcp-1) used at 1:200 (Cell Signaling
Technology). Fluorescence secondary antibodies were from Molecular Probes and

used at 1:1000. Rhodamine-phalloidin used at 1:400 (Life technologies), was used in
some experiments to counterstain the tissue. All primary and secondary antibodies
were diluted in PBT (PBS + 0,1% Triton X-100).
442
QUANTIFICATION AND STATISTICAL ANALYSIS
Quantification of adult eye size
Heads of 1-3 day old adult males and females were dissected in PBS, cleared with
15% Hydrogen Peroxide and mounted in Hoyer's medium: lactic acid (1:1) (Casares
and Mann, 2000). Frontal and occipital planes of each head were photographed using
a 10X objective on a Leica DM5000B microscope and a Leica DFC490 digital
camera. The areas of the eyes and head were measured using the polygonal tool of
ImageJ (Rueden et al., 2017) and expressed as arbitrary units. The Supplementary
figure to Materials and Methods illustrates the areas as measured. The area of each
eye was obtained as the sum of the area of the eye in the frontal and occipital planes.
Quantification of mitotic and apoptotic rates
To quantify the apoptosis (detected with anti-Dcp-1) and mitosis (detected with anti-
PH3) rates as the Dcp-1/PH3 signal relative to the area anterior to the morphogenetic
furrow (i.e. differentiation wave front), we wrote a plug-in for ImageJ (included as
Supplementary material). Briefly, it works as follows: First, for each sample, the images
of a 3D confocal stack are projected along the Z-axis using the maximum pixel
intensity. Then, a Region of Interest (ROI) is drawn by hand to define the region of the
eye disc anterior to the morphogenetic furrow, which contains the undifferentiated eye
cells. A threshold for pixel intensity is chosen manually on the green channel (Dcp-
1/PH3), making sure that the thresholded area includes accurately all the signal. This
threshold level is kept for every sample of the series, which were acquired with exactly
the same microscope configuration. Finally, the thresholded area is projected onto the
original image for the Mean Intensity Value and Integrated Density to be measured.
Statistical analysis
The statistical analyses were supported on *Rsoftware* (R-Core-Team, 2013) and
hypothesis decisions were made under 95% confidence threshold in all cases.
Modeling for intergroups comparison
The metrics of interest were modeled depending on genoype group and sex factors by
fitting a median regression model (quantile regression: *rq* function from *quantreg* library
for factors) where normality and homoscedasticity assumptions are not required
(Howard, 2018). This model allows estimating the effect of the different covariables and
factors in the median of the response variable, thus allowing the prediction of the
median response variable for each group as a sum of the corresponding effects
(Beyerlein, 2014). In our context, the *optix>+* group was set as reference level and
genotype group was coded as multiple dummy factors of presence/absence of each
genotype. Thus, the effect of simultaneous downregulation of two genes is also
evaluated as an interaction term in the model. Models considering or not sex as factor
were compared by calculating F-statistic of the ratio of variances explained by each
fitted model (*anova* function). Sex effect turned out to be non-significant regardless of
the response variable evaluated, thus, the general model is denoted by the following
expression:
484

$$y_i = \beta_{0;0.5}(optix) + \beta_{1;0.5}\,RHGRI_i + \beta_{2;0.5}\,tkvRI_i + \beta_{3;0.5}\,tkvRI_i\,DadRI_i$$
$$+ \beta_{4;0.5}\,DadRI_i + \beta_{5;0.5}\,tkvRI_i\,RHGRI_i + \beta_{6;0.5}\,Bar_i$$
$$+ \beta_{7;0.5}\,Bar_i\,RHGRI_i + \beta_{8;0.5}\,Bar_i\,DadRI_i + u_{i,0.5}$$

where:

y_i is the i-th value of rE or %Apop variables;

$\beta_{j,0.5}$ is the j-th value of the estimated conditioned coefficient to the 0.5 quantile;

$u_{i,0.5}$ is the i-th value of the error variable conditioned to the 0.5 quantile.

Comparison of sFAi variances of each genotype versus control

The signed Fluctuation Asymmetry (sFAi) is a intraindividual measure of eye size
precision. It is calculated as the relative differences between intraindividual (Left and
Right) eye areas:

$$sFA_i = (L - R)/(L + R)$$

We compared intergroup sFAi through their variances since sFAi distributions are
theoretically centered on 0. First, we assume there are no intragroup variance
differences due to sex (Levene test median centered; *LeveneTest function* from *car*
package). Second, median was subtracted for each group-sex combination to perform
a sex-independent comparison. Finally, Levene test median centered was performed
for pairwise comparisons of each group against the control one (*optix>+)*. P.values
were adjusted by Benjamini-Hochberg method (*LeveneTest* function from *car* package
and *p.adj* function*, method= "BH")*.

Comparison of rE variances of each genotype versus control

rE variance is a interindividual measure of size precision in eye development. First, we
assumed there were no intragroup variance differences due to sex (Levene test median
centered; *LeveneTest function* from *car* package). Second, median was subtracted for
each group-sex combination to perform a sex-independent comparison. Finally, Levene
test median centered was performed for pairwise comparisons of each group against
the control one (*optix>+)*. P.values were adjusted by Benjamini-Hochberg method
(*LeveneTest* function from *car* package and *p.adj* function*, method= "BH")*.

REFERENCES
Barrio, L., Milan, M., 2017. Boundary Dpp promotes growth of medial and lateral
regions of the Drosophila wing. Elife 6.
Bessa, J., Gebelein, B., Pichaud, F., Casares, F., Mann, R.S., 2002. Combinatorial
control of Drosophila eye development by eyeless, homothorax, and teashirt. Genes
Dev 16, 2415-2427.
Beyerlein, A., 2014. Quantile Regression—Opportunities and Challenges From a
User's Perspective. American Journal of Epidemiology 180, 330-331.
Blackman, R.K., Sanicola, M., Raftery, L.A., Gillevet, T., Gelbart, W.M., 1991. An
extensive 3' cis-regulatory region directs the imaginal disk expression of
decapentaplegic, a member of the TGF-beta family in Drosophila. Development 111,
657-666.
Bosch, P.S., Ziukaite, R., Alexandre, C., Basler, K., Vincent, J.P., 2017. Dpp controls
growth and patterning in Drosophila wing precursors through distinct modes of action.
Elife 6.

Boulan, L., Leopold, P., 2021. What determines organ size during development and
regeneration? Development 148.
Brummel, T.J., Twombly, V., Marques, G., Wrana, J.L., Newfeld, S.J., Attisano, L.,
Massague, J., O'Connor, M.B., Gelbart, W.M., 1994. Characterization and relationship
of Dpp receptors encoded by the saxophone and thick veins genes in Drosophila. Cell
78, 251-261.
Bryant, P.J., 1988. Localized cell death caused by mutations in a Drosophila gene
coding for a transforming growth factor-beta homolog. Dev Biol 128, 386-395.
Casares, F., Mann, R.S., 2000. A dual role for homothorax in inhibiting wing blade
development and specifying proximal wing identities in Drosophila. Development 127,
1499-1508.
Casares, F., McGregor, A.P., 2020. The evolution and development of eye size in flies.
Wiley interdisciplinary reviews. Developmental biology, e380.
Chanut, F., Heberlein, U., 1995. Role of the morphogenetic furrow in establishing
polarity in the Drosophila eye. Development 121, 4085-4094.
Chanut, F., Heberlein, U., 1997. Role of decapentaplegic in initiation and progression
of the morphogenetic furrow in the developing Drosophila retina. Development 124,
559-567.
Colombani, J., Andersen, D.S., Leopold, P., 2012. Secreted peptide Dilp8 coordinates
Drosophila tissue growth with developmental timing. Science 336, 582-585.
Fried, P., Sanchez-Aragon, M., Aguilar-Hidalgo, D., Lehtinen, B., Casares, F., Iber, D.,
2016. A Model of the Spatio-temporal Dynamics of Drosophila Eye Disc Development.
PLoS Comput Biol 12, e1005052.
Fuchs, Y., Steller, H., 2011. Programmed cell death in animal development and
disease. Cell 147, 742-758.
Garelli, A., Gontijo, A.M., Miguela, V., Caparros, E., Dominguez, M., 2012. Imaginal
discs secrete insulin-like peptide 8 to mediate plasticity of growth and maturation.
Science 336, 579-582.
Greenwood, S., Struhl, G., 1999. Progression of the morphogenetic furrow in the
Drosophila eye: the roles of Hedgehog, Decapentaplegic and the Raf pathway.
Development 126, 5795-5808.
Heberlein, U., Moses, K., 1995. Mechanisms of Drosophila retinal morphogenesis: the
virtues of being progressive. Cell 81, 987-990.
Heberlein, U., Wolff, T., Rubin, G.M., 1993. The TGF beta homolog dpp and the
segment polarity gene hedgehog are required for propagation of a morphogenetic
wave in the Drosophila retina. Cell 75, 913-926.
Howard, M., 2018. Comparison of the Performance of Simple Linear Regression and
Quantile Regression with Non-Normal Data: A simulation study, Mathematics.
University of South Carolina.
Hunt, D.M., 1971. The physiological control of gene action in the eyeless and eyegone
mutants of Drosophila melanogaster. Genet Res 17, 195-208.
Inoue, H., Imamura, T., Ishidou, Y., Takase, M., Udagawa, Y., Oka, Y., Tsuneizumi, K.,
Tabata, T., Miyazono, K., Kawabata, M., 1998. Interplay of signal mediators of
decapentaplegic (Dpp): molecular characterization of mothers against dpp, Medea, and
daughters against dpp. Mol Biol Cell 9, 2145-2156.
Kang, J., Yeom, E., Lim, J., Choi, K.W., 2014. Bar represses dPax2 and
decapentaplegic to regulate cell fate and morphogenetic cell death in Drosophila eye.
PLoS One 9, e88171.
Lander, A.D., 2011. Pattern, growth, and control. Cell 144, 955-969.
Lander, A.D., Gokoffski, K.K., Wan, F.Y., Nie, Q., Calof, A.L., 2009. Cell lineages and
the logic of proliferative control. PLoS Biol 7, e15.
Lee, J.J., von Kessler, D.P., Parks, S., Beachy, P.A., 1992. Secretion and localized
transcription suggest a role in positional signaling for products of the segmentation
gene hedgehog. Cell 71, 33-50.

Lobo-Cabrera, F.J., Navarro, T., Iannini, A., Casares, F., Cuetos, A., 2021. Quantitative
Relationships Between Growth, Differentiation, and Shape That Control Drosophila
Eye Development and Its Variation. Front Cell Dev Biol 9, 681933.
Lopes, C.S., Casares, F., 2010. hth maintains the pool of eye progenitors and its
downregulation by Dpp and Hh couples retinal fate acquisition with cell cycle exit. Dev
Biol 339, 78-88.
Ma, C., Zhou, Y., Beachy, P.A., Moses, K., 1993. The segment polarity gene hedgehog
is required for progression of the morphogenetic furrow in the developing Drosophila
eye. Cell 75, 927-938.
Masucci, J.D., Hoffmann, F.M., 1993. Identification of two regions from the Drosophila
decapentaplegic gene required for embryonic midgut development and larval viability.
Dev Biol 159, 276-287.
Masucci, J.D., Miltenberger, R.J., Hoffmann, F.M., 1990. Pattern-specific expression of
the Drosophila decapentaplegic gene in imaginal disks is regulated by 3' cis-regulatory
elements. Genes Dev 4, 2011-2023.
Matsuda, S., Affolter, M., 2017. Dpp from the anterior stripe of cells is crucial for the
growth of the Drosophila wing disc. Elife 6.
Morgante, F., Sorensen, P., Sorensen, D.A., Maltecca, C., Mackay, T.F., 2015. Genetic
Architecture of Micro-Environmental Plasticity in Drosophila melanogaster. Scientific
reports 5, 9785.
Neto, M., Aguilar-Hidalgo, D., Casares, F., 2016. Increased avidity for Dpp/BMP2
maintains the proliferation of progenitors-like cells in the Drosophila eye. Dev Biol 418,
98-107.
Neto, M., Naval-Sanchez, M., Potier, D., Pereira, P.S., Geerts, D., Aerts, S., Casares,
F., 2017. Nuclear receptors connect progenitor transcription factors to cell cycle
control. Scientific reports 7, 4845.
Neto-Silva, R.M., Wells, B.S., Johnston, L.A., 2009. Mechanisms of growth and
homeostasis in the Drosophila wing. Annu Rev Cell Dev Biol 25, 197-220.
Neumann, C.J., Nuesslein-Volhard, C., 2000. Patterning of the zebrafish retina by a
wave of sonic hedgehog activity. Science 289, 2137-2139.
Niven, J.E., Laughlin, S.B., 2008. Energy limitation as a selective pressure on the
evolution of sensory systems. J Exp Biol 211, 1792-1804.
Parsons, P.A., 1992. Fluctuating asymmetry: a biological monitor of environmental and
genomic stress. Heredity (Edinb) 68 (Pt 4), 361-364.
R-Core-Team, 2013. R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria.
Raff, M.C., 1992. Social controls on cell survival and cell death. Nature 356, 397-400.
Raff, M.C., Barres, B.A., Burne, J.F., Coles, H.S., Ishizaki, Y., Jacobson, M.D., 1993.
Programmed cell death and the control of cell survival: lessons from the nervous
system. Science 262, 695-700.
Rogers, E.M., Brennan, C.A., Mortimer, N.T., Cook, S., Morris, A.R., Moses, K., 2005.
Pointed regulates an eye-specific transcriptional enhancer in the Drosophila hedgehog
gene, which is required for the movement of the morphogenetic furrow. Development
132, 4833-4843.
Roignant, J.Y., Treisman, J.E., 2009. Pattern formation in the Drosophila eye disc. Int J
Dev Biol 53, 795-804.
Rueden, C.T., Schindelin, J., Hiner, M.C., DeZonia, B.E., Walter, A.E., Arena, E.T.,
Eliceiri, K.W., 2017. ImageJ2: ImageJ for the next generation of scientific image data.
BMC Bioinformatics 18, 529.
Siegrist, S.E., Haque, N.S., Chen, C.H., Hay, B.A., Hariharan, I.K., 2010. Inactivation of
both Foxo and reaper promotes long-term adult neurogenesis in Drosophila. Curr Biol
20, 643-648.
Song, Z., McCall, K., Steller, H., 1997. DCP-1, a Drosophila cell death protease
essential for development. Science 275, 536-540.

Spencer, F.A., Hoffmann, F.M., Gelbart, W.M., 1982. Decapentaplegic: a gene
complex affecting morphogenesis in Drosophila melanogaster. Cell 28, 451-461.
Tice, S.C., 1914. A new sex-linked character in Drosophila. Biol. Bull. 26, 221-230.
Tsuneizumi, K., Nakayama, T., Kamoshida, Y., Kornberg, T.B., Christian, J.L., Tabata,
T., 1997. Daughters against dpp modulates dpp organizing activity in Drosophila wing
development. Nature 389, 627-631.
Vollmer, J., Fried, P., Aguilar-Hidalgo, D., Sanchez-Aragon, M., Iannini, A., Casares,
F., Iber, D., 2017. Growth control in the Drosophila eye disc by the cytokine Unpaired.
Development 144, 837-843.
Vollmer, J., Fried, P., Sanchez-Aragon, M., Lopes, C.S., Casares, F., Iber, D., 2016. A
quantitative analysis of growth control in the Drosophila eye disc. Development 143,
1482-1490.
Waddington, C.H., 1942. Canalization of development and the inheritance of acquired
characters. Nature 150, 563–565.
Wartlick, O., Julicher, F., Gonzalez-Gaitan, M., 2014. Growth control by a moving
morphogen gradient during Drosophila eye development. Development 141, 1884-
1893.
Weasner, B.M., Kumar, J.P., 2022. The timing of cell fate decisions is crucial for
initiating pattern formation in the Drosophila eye. Development 149.
Wolpert, L., 2010. Arms and the man: the problem of symmetric growth. PLoS Biol 8.
Zhu, J., Palliyil, S., Ran, C., Kumar, J.P., 2017. Drosophila Pax6 promotes
development of the entire eye-antennal disc, thereby ensuring proper adult head
formation. Proc Natl Acad Sci U S A 114, 5846-5853.
659