**Done in 65 ms: Express visuomotor responses in upper limb muscles in Rhesus Macaques**

**Running head: Express visuomotor responses in macaques**

Aaron L. Cecala[1,2], Rebecca A. Kozak[3], J. Andrew Pruszynski[1,2], Brian D. Corneil[1,2,4]

[1]Department of Physiology and Pharmacology, Western University, London, Ontario, Canada, N6A 5B7
[2]Robarts Research Institute, 1151 Richmond St. N, London, Ontario, Canada, N6A 5B7
[3]Graduate Program in Neuroscience, Western University, London, Ontario, Canada, N6A 5B7
[4]Department of Psychology, Western University, London, Ontario, Canada, N6A 5B7

**Corresponding Author:** Brian D. Corneil: Robarts Research Institute, 1151 Richmond Street North, Western University, London, ON, Canada, N6A 5B7; E-mail: bcorneil@uwo.ca

Number of pages: 31
Number of figures: 5
Number of tables: 0
Number of words:
       Abstract: 240
       Introduction: 690
       Discussion: 1874

**Conflicts of interest:** The authors declare no competing financial interests.

**Keywords:** rhesus monkey; visually-guided reaching; electromyography; sensorimotor transformation

**Acknowledgements**: The authors would like to thank Katherine Faubert, Rhonda Kersten, Kristy Gibbs, Darren Pitre, and Matthew Reeves for their technical assistance and animal husbandry support, and Kevin Barker for his apparatus design and production expertise. We thank Tim Carrol, Sam Contemori, Rechu Divakar, Sebastian Lehmann, and Gerald Loeb for feedback on earlier versions of this manuscript.

**Grants:** This work was supported by operating grants to BDC from the Canadian Institutes of Health Research (CIHR) [MOP-93796; MOP-142317; PJT-180279] and the Natural Sciences and Engineering Research Council of Canada (NSERC) [RGPIN-311680; RGPIN 04394-2021], as well as by the Canada First Research Excellence Fund to BrainsCAN. RAK was supported by an Ontario Graduate Scholarship. JAP received a salary award from the Canada Research Chairs Program.

**Abstract**

41
How rapidly can the brain transform vision into action? Work in humans has established that the
transformation for visually-guided reaching can be remarkably rapid, with the first phase of upper
limb muscle recruitment, the *express visuomotor response*, beginning within less than 100 ms of
visual target presentation. Such short-latency responses limit the opportunities for extensive
cortical processing, leading to the hypothesis that they are generated via the subcortical
tectoreticulospinal pathway. Here, we examine if non-human primates (NHPs) exhibit express
visuomotor responses. Two male macaques made visually-guided reaches in a behavioral
paradigm known to elicit express visuomotor responses in humans, while we acquired
intramuscular recordings from the deltoid muscle. Across several variants of this paradigm,
express visuomotor responses began within 65 ms (range 48–91 ms) of target presentation.
Although the timing of the express visuomotor response did not co-vary with reaction time, larger
express visuomotor responses tended to precede shorter latency reaches. Finally, the magnitude of
the express visuomotor response was muted on trials where NHPs withheld a reach to one stimulus
in order to move to a stimulus appearing 34 ms later in the opposite direction. Overall, the response
properties and contextual control of express visuomotor responses in NHPs resemble those in
humans. Our results establish a new benchmark for visuomotor transformations underlying
visually-guided reaches, setting the stage for experiments that can directly test the comparative
role of cortical and subcortical areas in reaching when time is of the essence.

**Significance statement:**
Express visuomotor responses in upper limb muscles are brief periods of recruitment preceding
visually-guided reaches. Such responses begin ~90 ms after visual target presentation in humans,
and potentially arise from signaling along the tecto-reticulo-spinal pathway. Here, we show that
express visuomotor responses in macaques upper limb muscles resemble those in humans,
excepting that they evolve ~65 ms after target onset, consistent with shorter responses latencies in
macaques versus humans. Our results clock the completion of the visuomotor transformation for
rapid reaching, and set the stage for experiments to directly test the underlying substrates.

69
70
71

## Introduction

When required, the brain has a remarkable capacity to rapidly transform sensory input into motor output. Expedited responses can consist of eye movements to evaluate a potential threat, or mid-flight adjustments when reaching to a slipping cell phone. In some cases, the brain relies on subcortical circuits that provide a short-latency link between sensation and action, which are nested within broader fronto-parietal circuits mediating more deliberative actions. This appears to be the case for express saccades, which rely on the integrity of the midbrain superior colliculus (SC) and are potentiated by, but not critically reliant on, input from fronto-parietal circuits (Schiller et al., 1987; Edelman and Keller, 1996; Dorris et al., 1997; Dash et al., 2018, 2020). Although the SC has been implicated in mid-flight reach corrections (Day and Brown, 2001; Pruszynski et al., 2016; Reschechtko and Pruszynski, 2020; Suzuki et al., 2022) and interceptions of rapidly moving targets (Perfiliev et al., 2010), debates remain about the comparative role of subcortical versus cortical areas in expedited visually-guided reaching (Fautrelle et al., 2010; Gaveau et al., 2014; Novembre and Iannetti, 2021).

To better understand the neural mechanisms underlying expedited reaches, we and others have studied the timing and response properties of the first change of upper limb muscle recruitment elicited by visual stimulus onset. Doing so establishes the benchmark for how rapidly premotor circuits transform vision into action, and inferences can be drawn by comparing the timing and magnitude of muscle activity to response properties in candidate premotor areas. Such work has revealed that the first wave of upper limb muscle recruitment driven by stimulus presentation evolves within less than 100 ms in humans, regardless of the ensuing reaction time (Pruszynski et al., 2010; Wood et al., 2015; Gu et al., 2016; Kozak et al., 2019; Contemori et al., 2021a). This phenomenon has been termed the stimulus-locked response (Pruszynski et al., 2010) or fast or rapid visuomotor response (Glover and Baker, 2019; Kozak and Corneil, 2021), but lately we have adopted the term express visuomotor response (Contemori et al., 2021a, 2021b) to emphasize similarities with both express saccades and the timing and response properties of visual bursts on premotor cells in the motor layers of the SC (Corneil and Munoz, 2014). The timing and magnitude of such SC visual responses, express saccades, and express visuomotor responses on upper limb muscles are all potentiated by high-contrast, low spatial frequency stimuli (Marino et al., 2012,

2015; Wood et al., 2015; Chen et al., 2018; Kozak et al., 2019; Kozak and Corneil, 2021), and by

a variety of cognitive processes promoting premotor preparation (Rohrer and Sparks, 1993; Paré

and Munoz, 1996; Munoz et al., 2000; Edelman et al., 2007; Boehnke and Munoz, 2008; Li and

Basso, 2008; Bibi and Edelman, 2009; Gu et al., 2018; Contemori et al., 2021b, 2022a).

Here we aim to establish whether express visuomotor responses can be observed in non-human

primates (NHPs). We examine three response properties. First, express saccades and the related

phenomenon of express neck muscle responses occur ~20 ms earlier in NHPs than humans (Fischer

and Boch, 1983; Fischer and Ramsperger, 1984; Corneil et al., 2004; Goonetilleke et al., 2015). If

evoked by a similar mechanism, express visuomotor responses in arm muscles in NHPs should

evolve ~60-80 ms after stimulus onset. Second, although more time-locked to stimulus rather than

movement onset, larger express visuomotor responses in humans consistently precede shorter

reaction times (Pruszynski et al., 2010; Gu et al., 2016; Contemori et al., 2021a); we predict a

similar inverse relationship between recruitment magnitude and reaction time in NHPs. Third, the

magnitude of express visuomotor responses in humans is dampened if subjects know to move

away from or occasionally not respond to the stimulus (Wood et al., 2015; Gu et al., 2016; Atsma

et al., 2018; Kozak et al., 2020; Contemori et al., 2021b, 2022a). Such dampening is a signature of

top-down cortical control of a subcortical circuit, and we predict that similar top-down control can

be instantiated in NHPs. Overall, we find that NHPs readily generate express visuomotor responses

consistent with all these properties, establishing the NHP as a suitable animal model to study

underlying neural mechanisms.

**Methods**

*Subjects and Surgical Procedures*

Two male rhesus monkeys (*Macaca mulatta;* monkey G and monkey B), weighing 8.5 and 13.1

kg respectively, were used in these experiments. All training, surgical, and experimental

procedures were in accordance with the Canadian Council on Animal Care policy on the use of

laboratory animals (Olfert et al., 1993) and approved by the Western University Animal Use

Subcommittee. We monitored the monkeys' weights daily and their health was under the close

supervision of the university veterinarians.

Each animal underwent an aseptic procedure in which a head post was attached to the skull by way

of a dental acrylic pedestal prior to behavioral training. During training and experimental sessions,

animals were seated with the head-restrained in a customized primate chair that restricted

movements of the left arm, torso and hips while permitting unrestricted movements of the right

arm in a three-dimensional workspace. Testing was performed in a dimly illuminated room and

the monkeys were monitored via a video camera. Visual stimuli, behavioral control, and data

acquisition were controlled by MonkeyLogic (Hwang et al., 2019). All visual stimuli were

presented on a 42-in. color touch-sensitive monitor (4202L Elo Touch Solutions, Inc., Milpitas,

CA) positioned 33 cm in front of the monkeys. The monitor had a spatial resolution of 1920 ×

1080 pixels and a refresh rate of 60 Hz.

*The Emerging Target Task*

Monkeys generated visually-guided reaches to stimuli presented within the context of an emerging

target task (Kozak et al., 2020). The task involves a moving stimulus that temporarily disappears

behind an occluder, and then emerges in motion at either the left or right outlet. This task was

chosen as it engenders express visuomotor responses in most human subjects (Contemori et al.,

2021a, 2021b; Kozak and Corneil, 2021; Kearsley et al., 2022).

Data was collected from three variants of the emerging target task (**Fig. 1**). All trials started with

a red or green 'start position' stimulus that appeared at a central location below a gray rectangular

occluder. The monkeys were required to touch and hold the location of this start position stimulus

within a computer-defined window of 7 cm. Following this, a red or green 'target' stimulus that

matched the color of the start position stimulus appeared above the occluder. Providing the monkey

kept in contact with the start position stimulus for 500 ms, this target stimulus dropped vertically

at a constant velocity of 20 cm/s, and then disappeared behind the occluder. After a fixed latency

of 300 ms, a red target then emerged in motion below the occluder at either the right or left outlet.

At the same time, the start position stimulus disappeared and a secondary visual stimulus that lay

under a photodiode (and hence was unseen by the monkey) was presented for data alignment

purposes. On double target trials, a green target emerged in motion at the other outlet 34 ms later.

To obtain a reward, the monkeys had to maintain contact on the start position until target

emergence below the barrier, and then reach to touch the colored target (within a 11 cm radius)

that matched the color of the stimuli presented at the start of the trial before it disappeared off the

screen (i.e., reach to the red target on single target red (**Fig. 1A**) and double target red (**Fig. 1B**)

trials, or to the green target on double target green trials **(Fig. 1C)**). Thus, correct performance on

double target green trials required the monkeys to not reach towards the red target that emerged

slightly (34 ms) earlier. The moving target that appeared below the barrier started 15 cm to the

right or left of the start position stimulus, and then moved in an lateral-downward direction (45

deg below horizontal) at 20 cm/s. Trials where the monkey touched or moved toward the wrongly

colored target were tabulated as incorrect reaches, and retained for further analysis.

Across different experimental sessions, some blocks of trials consisted of only single-target red

trials, while other blocks of trials consisted of all task variants. Within a typical experimental day,

we first collected data in a block of only single-target red trials, prioritizing such data as EMG

recording electrodes occasionally were removed as animals performed the task. Within a block of

trials, all trial conditions (defined by target location and, when used, task variant) were presented

with equal frequency but randomly shuffled. Across all experimental sessions, we obtained muscle

recordings from a minimum of 380 to a maximum of 1470 trials.

**Figure 1. A-C.** Emerging Target Task. At the start of each trial, either a red (**A, B**) or green (**C**) stimulus appears at the start position (red or green squares), below an occluder (gray box). The monkey touches and holds the screen at this start position. A target of the same color then appears above the occluder and moves down the chute before disappearing briefly behind the occluder for 300 ms. In Single Target Red trials (**A**), a single red target emerges in motion below the occluder at one of two outlets. In the Double Target Red (**B**) and Double Target Green (**C**) trials, appearance of the red target in one outlet was followed 34 ms later by appearance of a green target at the other outlet. The monkey was rewarded for reaching to the target that matched the color of the stimuli presented at the start of the trial. **D.** Movement-aligned activity for the anterior, middle, and posterior heads of deltoids for the two monkeys for single-target red trials, showing greater recruitment for rightward than leftward reaches. Data shows the average of the mean recruitment from each session, normalized to the peak movement-aligned activity from each session. Contours subtend the mean +/- stderr.

*EMG electrode insertion and data collection*

Prior to a data collection session, monkeys were lightly sedated before intramuscular, fine-wire

EMG electrodes were acutely inserted into the posterior, medial, and anterior heads of the deltoid

muscle. Sedation followed an established procedure (Corneil et al., 2012), involving induction

with a low-dose of medetomidine, and reversal after electrode insertion with atipamazole. After

reversal, monkeys were given ~20 minutes to recover prior to beginning data collection. For each

intramuscular recording, we inserted two electrodes into the muscle belly staggered by ~1 cm to

enable bipolar recordings of bulk muscle activity. Each electrode was fabricated in-house by

threading PFA-coated 7-strand stainless steel wire (Type 316, A-M Systems, Sequim, WA) into a

191 27 gauge cannula, and then stripping the final ~3 mm of insulation. Anatomically, the different

192 heads of the deltoid were identified by manually palpating the head of the humerus. A ground

193 electrode was positioned on the dorsal neck approximately between the scapulae. Intramuscular

194 EMG activity was differentially amplified and recorded with a Myopac Junior system (Run

195 Technologies, low-pass filter modified to 2 kHz) and digitized at a rate of 1,000 Hz onto a Plexon

196 Omniplex system, which also received the photodiode signal.  Off-line, any DC offset was

197 removed and EMG data were full-wave rectified and smoothed with a 7-point running average.

198 Our sampling frequency was limited to 1,000 Hz, which undersamples intramuscular signals. We

199 confirmed that such undersampling had only a minor influence on our results by collecting data

200 from one additional session on a Ripple Grapevine system that sampled EMG data at 30,000 Hz.

201 This allowed us to compare results obtained from 3 simultaneously recorded muscles processed

202 offline by either downsampling the EMG signal to 1,000 Hz versus that obtained by first full-wave

203 rectifying the 30 kHz signal followed by integration into 1 ms bins (Loeb and Gans, 1986; Elsley

204 et al., 2007). For all three muscles recorded at 30 kHz, these different processing steps changed

205 the discrimination latency of the express visuomotor response by only 1 ms. Thus, while our

206 undersampled data would not be appropriate for identifying single motor-unit action potentials, it

207 does accurately capture the timing of changes in multi-unit recruitment after visual target

208 presentation.

210 *Data Analysis*

211 All data analyses were conducted in MATLAB® (R2021a; Natick, MA, USA). We rejected trials

212 where the monkey either did not respond or failed to respond within 500 ms, as well as trials where

213 movements began within less than 80 ms of target emergence below the barrier. Reach RT was

214 calculated as the time from target emergence below the barrier (indicated by the photo-diode

215 signal) to the time point where the monkey's hand was either no longer in contact with the

216 touchscreen at the start position, or when the monkey's hand started deviating toward the emerging

217 target. Offline, we further inspected all trials within a graphical user interface, allowing us to adjust

218 the movement onset mark at a resolution of 1 kHz, and flag trials where the initial movement was

219 in the opposite direction of the correct response. Monkey G moved more rapidly and tended to

220 remain in contact with the screen during the reach (i.e., 'smearing' their hand across the screen),

221 whereas Monkey B moved more slowly and had a greater tendency to lift their finger off the screen

when reaching. Thus, incorrect reaches in Monkey G included trials where the monkey first moved

in the incorrect direction before correcting their reach in mid-flight. In contrast, incorrect reaches

in Monkey B were sometimes only identified when the monkey touched the wrong location. On

other occasions, an incorrect reach in Monkey B was identified by a deflection in the horizontal

touch screen position in the wrong direction, as identified in the user interface.

The different heads of the deltoid muscle are known to have different preferred reaching directions

in the rhesus macaque (Kurtzer et al., 2006). Note however that the reaching task in this study

required that the monkeys reach forward and maintain contact with the touchscreen at the start

point, and then upon target emergence move their arm to the right or left to obtain a reward. In the

context of this bidirectional task, as shown in **Fig. 1D**, all heads of deltoid exhibited greater

recruitment before rightward versus leftward reaches on the touchscreen. Therefore, for the

purpose of analyzing the express visuomotor response, the rightward direction was deemed to be

the preferred direction of all muscles.

As with previous work (Pruszynski et al., 2010; Gu et al., 2016; Contemori et al., 2021a), we used

a time-series receiver-operating characteristic (ROC) analysis to define the presence and latency

of the express visuomotor response. Data recorded from each muscle was treated as an independent

sample, and data was segregated by side of target emergence and task variant. For blocks with

multiple interleaved task variants, separate time-series ROCs were run on data for each task

variant, comparing EMG activity after left or right target emergence. The ROC value (i.e., the area

under the ROC curve) was calculated and plotted at every time sample (1 ms) from 100 ms before

to 500 ms after target emergence (e.g., **Fig. 2C**); this value indicates the likelihood of

discriminating the side of target emergence based on EMG activity alone. A ROC value of 0.5

indicates chance discrimination, whereas values of 1 or 0 indicate perfectly correct or incorrect

discrimination, respectively. Although time-series ROCs often fluctuated around 0.5 in the 100 ms

surrounding target emergence, we often noted the presence of linear trends in this interval as well

as a high degree of variance on task variants with fewer overall trials. These precluded the use of

fixed thresholds to determine the latency of any express response. Accordingly, we first detrended

the time-series ROC based on a linear regression run on the time-series ROC over a baseline

interval spanning from 100 ms before to 45 ms after target emergence. We then set a discrimination

threshold to be the first of 8 of 10 points where the time-series ROC exceeded 3 standard deviations
above the mean of the detrended baseline time-series ROC. If this time was between 45 to 100 ms
after target emergence, then the muscle was deemed to be exhibiting an express visuomotor
response, and the latency was taken to be the discrimination time. The magnitude of the express
visuomotor response was determined by the integral of EMG activity measured over the next 30
ms. All EMG data is reported in source microvolts, and was not subjected to any normalization
unless otherwise noted.

*Experimental design and statistical analyses*

All statistical analyses were performed in MATLAB® (R2021a; Natick, MA, USA). Comparisons

of the magnitude and latencies of express visuomotor responses and of reaction times across the

various trial types were performed via *t*-tests or one-way ANOVAs, using repeated measures

where appropriate. Post-hoc comparisons relied on the Tukey–Kramer critical value, as well as

paired and unpaired *t*-tests when appropriate. Correlational analyses were performed on latency,

magnitude, and RT measures, as described below. Comparisons of error rates across different task

variants were performed using a chi-squared test.

**Results**

Two rhesus macaques reached to red or green targets presented in the context of the emerging

target task (**Fig. 1A-C**). This task readily elicits express visuomotor responses in humans (Kozak

et al., 2020; Contemori et al., 2021a; Kozak and Corneil, 2021). Our goal was to establish the

phenomenon of express visuomotor responses in monkey, by: i) describing the response latencies

and consistency of such responses both within and across experimental sessions, ii) investigating

the relationships with ensuring reach reaction times (RTs), and iii) establishing whether the express

visuomotor response to the red stimulus would be decreased when monkeys were instructed to

reach in the opposite direction to a delayed emerging green stimulus. Our dataset consists of EMG

recordings made in 8 sessions in monkey G (7, 8, and 7 successful recordings from anterior,

medial, and posterior deltoid respectively) and 10 sessions in monkey B (9, 8, and 8 from anterior,

medial, and posterior deltoid respectively). As all muscles exhibited greater recruitment for

rightward vs leftward reaches on the touchscreen (**Fig. 1D**), and since we noted no differences in

the express visuomotor responses from the three heads of deltoid, we treated each recording as an

independent sample of multiple motor units, and pooled data within a given monkey.

*Stimulus emergence evokes short-latency express visuomotor responses*

We begin by describing the behavior and patterns of muscle recruitment immediately following

target emergence in single target red trials. On such trials, Monkey G reached within 179 +/- 61

ms and Monkey B reached within 223 +/- 47 ms (**Fig. 2A**). Both animals made some reaching

errors when only a single red target emerged (6% for Monkey G, 5% for Monkey B), but more

commonly anticipated target emergence (8.6% for Monkey G, 12.1% for Monkey B).

Representative examples of EMG data recorded from each monkey are shown in **Fig. 2B** and **C**.

As animals held their arm against gravity in order to touch the central starting position, there was

a modest level of recruitment prior to and immediately after target emergence. Very soon after

target emergence, muscle recruitment increased or decreased following target emergence to the

right or left, respectively. Such lateralization is a hallmark of express visuomotor responses in

humans (Pruszynski et al., 2010; Wood et al., 2015), and emphasizes that express visuomotor

responses are tuned to stimulus location rather than being a generic startle or preparatory response

(Gu et al., 2019; Selen et al., 2019). Another defining feature of express visuomotor responses in

humans is the trial-by-trial consistency in which this initial change in muscle recruitment is more

locked to target emergence rather than movement onset. This indeed was the case in the

representative examples shown in **Fig. 2B** (see red rectangles): note how this initial lateralized

change in muscle activity occurred ~60-70 ms after target emergence, well before the hand moved

or lifted away from the starting location. Following the express visuomotor response, rightward

reaches were preceded by a gradually increasing level of recruitment that peaked just before reach

initiation, whereas muscle recruitment before leftward reaches persisted near or below the baseline

level of activity prior to target emergence. The data shown for Monkey B exhibited a second target-

locked decrease in muscle activity ~130-140 ms after left target emergence, resembling reports of

oscillations in muscle recruitment initiated by a visual stimulus in humans (Wood et al., 2015).

Such double-dips were seen in about half of the recordings in Monkey B but were not observed in

Monkey G. We suspect this is because monkey G had generally shorter RTs, so that the recruitment

aligned with reach onset overlapped with any subsequent activity following the express

visuomotor response.

**Figure 2. A.** Frequncy histograms of reaction times (20 ms bins) for Monkey G (top row, n = 1303 trials) and Monkey B (bottom row, n = 2648 trials) across all experimental sessions. **B.** Representative examples of express visuomotor responses in Monkey G (top row) and Monkey B (bottom row), depicting trial-by-trial EMG recruitment aligned to target emergence to the left or right. In these plots, trials are sorted by movement onset (black squares), and the color conveys the magnitude of recruitment. **C.** For each monkey, the top row depicts mean EMG activity subtended by the standard error for left (light blue) and right (dark blue) reaches, and bottom row shows the time-series ROC (AUC = area under curve). The red rectangles in **B** and grey rectangles in **C** show the 30 ms interval following the detection of the express visuomotor response.

To analyze the timing of express visuomotor responses, we performed a time-series receiver operating characteristics (ROC) analysis. This analysis determines the timing at which information following leftward or rightward target emergence influenced muscle recruitment. For the examples shown in **Fig. 2C**, the express visuomotor response began 68 ms (Monkey G) or 55 ms (Monkey B) after target emergence. In both examples, the time-series ROC rose quickly after this discrimination time, and remained elevated from the next ~200 ms.

We repeated this analysis across our sample of deltoid recordings. Across our sample, an express visuomotor response, which was defined as occurring within 45-100 ms, was detected in 77% (17 of 22) recordings in monkey G and in all 25 recordings in monkey B. When present, express visuomotor responses began very soon after stimulus emergence, with discrimination latencies being 65 +/- 8 ms (range: 52 to 81 ms) for monkey G and 64 +/- 11 ms (range: 48 to 91 ms for monkey B; **Fig. 3A**; no difference between monkeys, *t*-test, $t(40) = 0.34$, $p = 0.72$). A potential concern about these short discrimination latencies is that they may arise only because of recruitment associated with short-RT reaches. We therefore repeated the time-series ROC analysis

across our sample only on the slower half or all trials. Excluding the shorter-than-average RT trials

in each session increased reach reaction times by ~25 ms to 205 +/- 63 ms for Monkey G and 253

+/- 37 ms for Monkey B. Express visuomotor responses persisted in 72% (16 of 22) of recordings

in monkey G and in all 25 recordings in monkey B, with average discrimination latencies being at

69 +/- 7 ms for monkey G and 68 +/- 12 ms for monkey B (**Fig. 3A**). Although significant in both

monkeys (paired *t*-test, monkey G: *t*-test, $t(15) = -6.9$, $p < 10^{-5}$; monkey B: $t(24) = -3.10$, $p = 0.005$),

the ~4 ms increase in discrimination latency after excluding shorter-than-average RTs was less

than the ~25 ms increase in mean RT. We suspect that slightly longer discrimination latencies

reflects lower magnitude express visuomotor responses associated with longer RT trials (see next

section), as reported previously (Contemori et al., 2021a, 2021b).

**Figure 3.** Properties of express visuomotor responses. **A.** Latencies of express visuomotor responses for both monkeys, showing latencies when all trials are considered regardless of reaction time, and after including only the longer-than-average reaction times. Grey lines connect observations from the same muscle from a given experimental day (grey + marks denote representative data shown in Fig. 2). Black lines and squares show mean +/- standard error. Asterisks show significant differences of paired t-test (p < 0.05). **B.** Calculation of the proportional magnitude of the express visuomotor response, using the representative data from Monkey B. We integrate the difference in the mean EMG curve for right versus left responses over the 30 ms after the onset of the express visuomotor responses, and divide this by the integral of the 30 ms interval surrounding peak EMG activity aligned to right reach onset. Same

general format as **A;** asterisk showing significant results of a student's t-test (p < 0.05).

Previous work in humans has shown that express visuomotor responses observed in the emerging

target task can be quite large in magnitude, occasionally equalling if not exceeded 'voluntary'

recruitment magnitude achieved when recruitment is aligned to reach onset reach onset (Contemori

et al., 2021a; Kozak and Corneil, 2021). To establish the proportional magnitude of the express

visuomotor response, we expressed it relative to a 30 ms interval surrounding the peak magnitude

of movement-aligned recruitment (**Fig. 3B**). Expressed in this fashion, the express visuomotor

response reached 0.37 +/- 0.21 (range: 0.09 to 0.85) for monkey G and 0.26 +/- 0.10 (range: 0.03

to 0.47) for monkey B (proportional magnitudes were significantly larger in monkey G than

monkey B; *t*-test, $t(40) = 2.2$, $p = 0.03$). Thus, on average, the recruitment magnitude of the express

visuomotor response was about one quarter to one third of the magnitude of reach-related

recruitment.

*Express visuomotor response magnitude inversely correlates with RT*

Although the express visuomotor response may be modest in magnitude, previous work shows that

larger express visuomotor responses tend to precede short RTs (Pruszynski et al., 2010; Gu et al.,

2016; Contemori et al., 2021a). To investigate whether a similar inverse correlation is seen in

NHPs, we calculated the trial-by-trial magnitude of EMG recruitment during an express

visuomotor response window (integrated over 30 ms after the express visuomotor response was

detected), and plotted it as a function of RT. **Figure 4** shows these plots for the representative data

from monkey G (**Fig. 4A**) and monkey B (**Fig. 4B**), and both show the inverse correlation between

magnitude and RT expected from previous studies (Monkey G: $r$ = -0.57, $p < 10^{-10}$; Monkey B: $r$

= -0.40, $p < 10^{-7}$). We repeated this analysis for every session in our sample, and for both monkeys,

observed the expected negative correlation (**Fig. 4C**; Monkey G: mean $r$ = -0.48 +/- 0.26. Monkey

B: mean $r$ = -0.31 +/- 0.17). The distributions for both monkeys lay significantly below zero ($t$-

test; Monkey G: $t(16)$ = -7.7, $p < 10^{-6}$. Monkey B: $t(24)$ = -9.2, $p < 10^{-8}$), and they were significantly

different from each other ($t$-test, $t(40)$ = -2.61, $p$ = 0.01).

**Figure 4.** Larger express visuomotor responses tended to precede shorter reach reation times (RTs). **A, B**. Representative data plotting magnitude of express visuomotor response as a function of reach reaction time, from the representative sessions shown in Figure 2. Each data point represents a single trial, and the black diagonal line shows the linear regression. The correlation coefficient (*r*) is shown in the top-right of each subplot. **C**. Histogram of correlation coefficients across all sessions in which an express visuomotor response was detected. Vertical dash line indicates 0. Solid or grey lines or bins denote data for Monkey G or B, respectively. Vertical arrows show mean *r*-values.

*Contextual control influences the magnitude but not timing of the express visuomotor response*

Having established that macaque upper limb muscle recruitment following the emergence of a

single target resembles that observed in humans, we now turn to data recorded when two targets

emerged below the barrier. The sequence of target emergence was the exact same on double-target

trials; the red stimulus always appeared 500 ms after the initial descending stimulus disappeared

behind the barrier, and the red stimulus always appeared 32 ms before the green stimulus. Monkeys

reached to the emerging stimulus that matched the color of the central hold stimulus and dropping

stimulus (**Fig. 1B,C**). Double target green trials therefore required monkeys to not reach to the red

emerging stimulus in order to reach to the green stimulus that subsequently emerged on the

opposite side. Thus, monkeys had to exert a different degree of contextual control in how they

responded to the red emerging stimulus on double target red than double target green trials.

We first compared reaching behavior on double target red vs double target green trials. As

expected, the RTs for correct responses (timed to the emergence of the red stimulus) were longer

for double target green trials than double target red trials (**Fig. 5A**, Monkey G: double target red =

188 +/- 70 ms, n = 572; double target green = 209 +/- 97 ms, n = 206; $t(776) = -3.36$, $p < 0.001$.

Monkey B: double target red = 235 +/- 50 ms, n = 1607; double target green = 255 +/- 49 ms, n =

962; $t(2579) = -9.63$, $p < 10^{-10}$). Both monkeys made reaching errors at a significantly higher rate

on double target green than double target red trials (Monkey G: chi square-test, $X^2 (1) = 33.9$, $p <$

10-8; Monkey B: $X^2 (1) = 11.25$; $p < 10^{-3}$). When generated, such errors started earlier than correct

responses (Monkey G: error RTs on double target green = 172 +/- 45 ms, n = 207; $t$-test against

RTs of correct responses on double target green trials $t(411) = 5.01$, $p < 10^{-7}$; Monkey B: error

RTs on double target green = 186 +/- 50 ms, n = 409; $t(1376) = 23.38$, $p < 10^{-10}$). While Monkey

G made a higher proportion of errors on double target green trials than Monkey B (chi-square-test,

$X^2 (1) = 5.21$; $p = 0.02$), an analysis of hand trajectories showed that Monkey G often moved

initially toward the red stimulus before rapidly reversing reach direction to reach toward the green

stimulus (**Fig. 5B**). Overall, these behavioral patterns are consistent with the animals trying, and

occasionally failing, to exert more contextual control on double target green trials.

**Figure 5.** Contextual control on Double Target trials. **A.** Histograms across all sessions of reaction times (20 ms bins) relative to the emergence of the red stimulus for Monkey G (top row) and Monkey B (bottom row), for Double Target Red trials (left column, Monkey G: 632 trials, error rate = 9.4%; Monkey B: 1788 trials, error rate = 9.8%) or Double Target Green trials (right column, Monkey G: 413 trials, error rate = 50.1%; Monkey B: 1378 trials, error rate = 29.6%). Upward/downward histograms represent correct/incorrect responses. **B.** Trajectory of horizontal touchscreen position averaged across all trials for when the monkey should have reached right on correct Double Target Red trials (red contours), correct Double Target Green trials (green contours) and incorrect Double Target Green trials (black contours). Contours denote mean +/- stderr. **C.** Depiction of mean EMG activity +/- stderr for when the red stimulus emerged to the right. Grey rectangles denote the 30 ms interval following the express visuomotor response, over which magnitude was calculated. **D,E.** The latency (**D**) or magnitude (**E**) of the express visuomotor response, by trial type and/or performance, across 29 occurrences where an express visuomotor response could be detected on both Double Target Red and Double Target Green trials. Grey lines connect observations from the same muscle from a given experimental day. Black lines and squares show mean +/- standard error. Asterisks show significant differences of Bonferroni corrected paired t-test (p < 0.05).

We next asked the degree to which contextual control influenced the express visuomotor response to the red stimulus. In humans, express visuomotor responses evolve at the same time but are dampened in magnitude when subjects humans move away from rather than toward a peripheral stimulus (Gu et al., 2016; Kozak et al., 2020), or exert some other degree of contextual control (Wood et al., 2015; Atsma et al., 2018; Contemori et al., 2021b). **Figure 5C** shows the mean +/- stderr representations of EMG activity from the same representative sessions as shown in **Figure**

**2**, depicting right middle deltoid recruitment when the red stimulus appeared to right. The monkeys

reached to the right on double target red trials (red contours in **Fig. 5C**), and the express visuomotor

response resembled that seen on the single target red trials (**Fig. 2B,C**). On double target green

trials, the correct response was to withhold the rightward reach, and reach to the left. When the

animals did this correctly, the initial express visuomotor response to the red stimulus was muted

(green contours in **Fig. 5C**). In contrast, when the monkeys incorrectly reached to the red stimulus

on double target green trials, initial muscle recruitment during the express response interval

resembled that on double target red trials (compare black and red contours in **Fig. 5C**), whereas

later phases of recruitment resembled correctly performed double target green trials, as the animal

corrected their reach. Thus, for these examples, the express visuomotor response to the earlier

appearing red stimulus depended both on the context of the trial (double target red vs double target

green), and on performance on double target green trials.

Across our sample, we found that contextual control influenced the magnitude but not timing of

the express visuomotor response to the red stimulus. To quantify this, we first compared the timing

of the express visuomotor response, when such a response was detected on both double-target red

and double-target green trials. We found no influence of context on express visuomotor response

latency, with the latency being unchanged across double-target red trials and double-target green

trials (**Fig. 5D**. Monkey G: double-target red = 66 +/- 9 ms; double-target green = 70 +/- 14 ms;

paired *t*-test; $t(9) = -0.75$, $p = 0.46$; Monkey B: double-target red = 67 +/- 13 ms; double-target

green = 67 +/- 10 ms; $t(18) = 0.02$, $p = 0.98$). We then calculated the magnitude of the express

visuomotor response over the next 30 ms on double target red or double target green trials,

subdividing the latter based on whether monkeys correctly reached to the green stimulus or

incorrectly reached to the red stimulus. To do this, we normalized all express visuomotor response

magnitudes to the red stimulus to the average value across all double target trials. A one-way

repeated-measures ANOVA showed that the express visuomotor response magnitude to the red

stimulus varied as a function of instruction and performance (**Fig. 5E**, one-way repeated-measures

ANOVA, $F_{2,56} = 10.15$, $p < 0.0005$). Post-hoc analyses revealed significantly larger responses on

double target red versus correct double target green trials (paired *t*-test, $t(28) = 2.90$, $p = 0.01$), and

significantly larger responses on incorrect vs correct double target green trials ($t(28) = 4.22$, $p <$

0.001). Thus, as in humans, the magnitude but not latency of the express visuomotor response

changed with the context of the task (REFS). Furthermore, as with anti-reaches (Gu et al., 2016),

larger express visuomotor responses preceded incorrect responses.

**Discussion**

We focused on the initial recruitment of upper limb muscles in monkeys reaching towards

emerging targets, using a paradigm which readily elicits express visuomotor responses in humans

(Kozak et al., 2020). Express visuomotor responses in monkeys exhibited the response properties

known in humans: its timing is more time-locked to target emergence than movement onset, and

its magnitude is both inversely related to RT and liable to dampening via contextual control. Our

results establish how rapidly the visual-to-motor is completed in the rhesus macaque brain, place

constraints on the timing of premotor events, and set the stage for neurophysiological experiments

that can directly compare the characteristics of such premotor events to those seen in the motor

periphery.

*Comparisons of express responses in humans and monkeys*

Rhesus macaques offer an excellent animal model for understanding rapid visual-to-motor

transformations, due to the many homologies in foveate vision, implicated brain networks, and the

musculoskeletal anatomy of the eyes, neck, and upper limb. Humans and macaques generate a

variety of expedited visuomotor responses in the lab, including express saccades (Fischer and

Weber, 1993), microsaccades (Hafed and Ignashchenkova, 2013; Tian et al., 2016), express

visuomotor responses in neck muscle (Goonetilleke et al., 2015), reaches to rapidly moving stimuli

(Perfiliev et al., 2010), and ocular following responses (Miles et al., 1986; Gellman et al., 1990).

A common finding is that such responses evolve ~20-30 ms earlier in macaques than humans,

which is broadly consistent with the 3/5th rule (Schroeder et al., 2004) for quicker sensory

processing in macaques versus humans. Consistent with this, express visuomotor responses in

macaques started at ~65 ms, compared to 80-100 ms in humans.

A notable exception for earlier responses in macaques vs. humans is when subjects adjust reaching

trajectories in mid-flight to a displaced visual target. Humans can initiate such on-line corrections

within ~150-200 ms (Gaveau et al., 2014), even without conscious perception (Goodale et al.,

1986). The response latencies for on-line corrections in macaques are generally longer than in

humans, ranging between 200-250 ms (Georgopoulos et al., 1983; Archambault et al., 2009, 2011;
Pastor-Bernier et al., 2012; Dickey et al., 2013), although latencies comparable with humans have
been reported (Ames et al., 2019). The generally slower responses in macaques likely arises due
to an task-imposed emphasis of accuracy over speed, so that animals were not responding as
quickly as possible. Work by Perfiliev and colleagues (2010) compared 'interceptive' reaching
responses to rapidly moving targets of humans, monkeys, and cats, and clearly showed that all
three species can initiate reaches in less than 150 ms, with limb muscle recruitment in humans
starting at 90-100 ms. Such recruitment onsets are within the range of express visuomotor
responses in humans, but Perfiliev and colleagues did not record EMG in cats or monkeys.
475
*Methodological considerations*
A study by Lara and colleagues (Lara et al., 2018) recorded upper limb muscle activity in monkeys
in a 'quasi-automatic' task, where monkeys intercepted a target moving radially on a touch screen.
This task expedited RTs to ~200 ms on average, which is similar to the reaction times we observed.
As with the emerging target paradigm, the quasi-automatic task features a suddenly appearing
moving visual target. Lara and colleagues recorded from multiple heads of deltoid in addition to
other upper limb muscles, reporting the earliest changes in average muscle recruitment at ~100
ms, with single-trial onsets as low as 90 ms. Although these response latencies are short, they are
still ~25-35 ms longer than those reported here. The failure for the quasi-automatic task to elicit
express visuomotor responses may be due to a lack of precise temporal certainty about target
presentation or the absence of implied motion, both of which are critical in the emerging target
task (Kozak et al., 2020; Contemori et al., 2021a; Kozak and Corneil, 2021). The quasi-automatic
task in the Lara study was also intermingled with other trial types that engender longer RTs, which
may also lower the probability of express visuomotor responses.
Caution is always required when comparing responses latencies across different laboratories.
Stimulus luminance and spatio-temporal frequency impact the latencies of on-line reach
corrections (Veerman et al., 2008; Kozak et al., 2019), the timing and magnitude of express muscle
responses (Wood et al., 2015; Kozak et al., 2019; Kozak and Corneil, 2021), and on the timing and
vigor of sensorimotor responses in the superior colliculus (SC) (Li and Basso, 2008; Marino et al.,
2012; Chen et al., 2018). Determining whether limb muscle recruitment meets an "express"
criterion should involve an assessment of trial-by-trial muscle recruitment over a range of RTs,

lest averaged recruitment profiles be biased by trials with the shortest RTs. The trial-by-trial
representations of EMG activity presented in the Lara study did not show consistently short
recruitment latencies across trials, strengthening our contention that express visuomotor responses
were not consistently elicited in their quasi-automatic task.

The emerging target paradigm provides a framework in which to study the most rapid visual-to-
motor transformations for reaching, and our results show that the underlying phenomenon is
distributed to many deltoid motor units in the macaque. The magnitudes of express responses
compared to later waves of activity are more modest in monkeys than in humans. This may be
because the deltoid muscles targeted here may not be prime movers when monkeys interact with
a touchscreen. Future work recording from more muscles of the upper limb may also find stronger
express visuomotor responses, and experimental setups with a robotic manipulandum will allow
loading of selected muscle, as done routinely in humans. Work in NHPs may also benefit from
various endogenous or exogenous cueing strategies that potentiate the phenomenon of express arm
responses in the human arm (Contemori et al., 2021b, 2022a) or neck (Corneil et al., 2008). While
the express visuomotor responses we recorded in NHPs are modest, similarly-sized express
visuomotor responses induce force transients or small movements toward the stimulus (Wood et
al., 2015; Gu et al., 2016; Atsma et al., 2018). We did observe small movements of the hand toward
the earlier appearing red target in Monkey G on double target green trials, which we surmise arose
from both the express visuomotor responses and subsequent periods of recruitment. The
assessment of limb motion via the touchscreen is coarse, hence setups using robotic manipulanda
or motor tracking will permit more precise assessment of the kinematic consequences of express
visuomotor responses in monkeys.

*Implications of our results*
Express visuomotor responses are thought to arise from a subcortical tecto-reticulo-spinal circuit
that runs in parallel with corticospinal circuits (Boehnke and Munoz, 2008; Pruszynski et al., 2010;
Corneil and Munoz, 2014; Contemori et al., 2022b). This hypothesis has never been directly tested
in monkeys, as the lone reaching study inactivating the SC during a reaching task (Song et al.,
2011) did not record limb EMG and used a delayed response task that would not engender express
visuomotor responses. Considerable circumstantial evidence implicates the visual response within

the intermediate and deep layers of the superior colliculus (dSC) as being key in driving express
responses. The timing and vigor of the visual response, when summed with pre-existing
preparatory activity, drives express saccades (Edelman and Keller, 1996; Dorris et al., 1997;
Sparks et al., 2000; Dash et al., 2018). Visual responses in the dSC are also correlated with the
magnitude of express visuomotor responses in the neck, leading such responses by an appropriate
efferent lag (Rezvani and Corneil, 2008). Reach-related neurons in the dSC display short-latency
visual responses (Werner et al., 1997a, 1997b) and correlate with upper limb muscle activity
(Werner et al., 1997a; Stuphorn et al., 1999), although the properties of visual responses in reach-
related neurons have not been systematically investigated. Our speculation of a subcortical route
initiating visually-guided reaches to a punctate target is consistent with recent stimulation results
in humans (Divakar et al., 2022), and parallels a growing appreciation for the role of subcortical
pathways in manual following responses to whole-field motion (Saijo et al., 2005) or flashes
(Nakajima et al., 2021).

While we acknowledge the shortcomings in comparing results across laboratories and
experimental tasks, the express visuomotor responses reported here on arm muscles occur at an
interval when neural activity in the primary motor cortex is thought to be in a null space that is not
tied to motor output (Lara et al., 2018). Indeed, one of the main conclusions for the Lara study
mentioned above is that a mandatory transition of neural activity within motor cortex from a null-
space to a movement-potent subspace precedes motor output by ~20-30 ms. Our results present
two alternatives which can frame future experiments.

First, if our hypothesis is correct, a subcortical route through the dSC can drive express visuomotor
responses when M1 is in a null space. An intriguing corollary to this alternative, given ascending
connections from the SC to thalamic areas that themselves project to motor cortices (Benevento
and Fallon, 1975; Matelli and Luppino, 1996), is that it is signaling from the SC to the motor cortex
that drives the transition of cortical activity from null to movement potent subspaces (Ames et al.,
2019), which presumably could allow cortical activity to drive subsequent muscle recruitment that
is informed by the consequences of the preceding express visuomotor response. Ascending signals
from the SC through the thalamus have been long established as being critical for relaying
efference copy signals for eye movements (Sommer and Wurtz, 2002), and implicated in

prioritizing spatial locations for target selection for reaching (Song et al., 2011) and for higher

order processing (Lovejoy and Krauzlis, 2010; Bogadhi et al., 2021; Peysakhovich et al., 2023).

Alternatively, could the transition of neural activity from a null to movement potent subspace in

M1 simply occur earlier in the emerging target paradigm? This line of reasoning surmises that

signaling along a cortico-spinal or cortico-reticulo-spinal route drives express visuomotor

responses. If so, visually-related activity would have to evolve in motor cortex ~10-20 ms earlier

than the express visuomotor responses we report, so at around 50 ms. There are reports of visual

responses in the macaque primary motor cortex at such latencies (Murphy et al., 1982; Lamarre et

al., 1983; Crammond and Kalaska, 1994; Riehle et al., 1997; Cisek and Kalaska, 2005; Ames et

al., 2019), but comparisons to these reports are again hindered by the use of tasks or visual stimuli

that in retrospect may not have been optimized to elicit express responses, and various filtering

settings or analysis schemes that complicate timing comparisons. Examination of the rasters of

single-unit data from such reports, when available, also do not always reveal the trial-by-trial

consistency that we observed in the motor periphery. One intriguing exception comes from the

work of Reimer and colleagues (Reimer and Hatsopoulos, 2010), who studied motor cortical

activity while NHPs adjusted the trajectory of ongoing limb movements to jumped visual targets.

When one accounts for an acknowledged timing error, motor cortex activity changed consistently

~50 ms after the target jump. While this theoretically shows that signaling related to visual signal

onset could arrive in time within motor cortices, we note that express visuomotor responses in

humans evolve ~10-15 ms earlier (and hence ~80 ms or less) when evoked when the arm is already

in motion (Kozak et al., 2019), presumably due to disengagement of a control policy for postural

stability (Cluff and Scott, 2016).

 **Figure Legends**

**Figure 1. A-C.** Emerging Target Task. At the start of each trial, either a red (**A, B**) or green (**C**)
stimulus appears at the start position (red or green squares), below an occluder (gray box). The
monkey touches and holds the screen at this start position. A target of the same color then appears
above the occluder and moves down the chute before disappearing briefly behind the occluder for
300 ms. In Single Target Red trials (**A**), a single red target emerges in motion below the occluder
at one of two outlets. In the Double Target Red (**B**) and Double Target Green (**C**) trials, appearance
of the red target in one outlet was followed 34 ms later by appearance of a green target at the other
outlet. The monkey was rewarded for reaching to the target that matched the color of the stimuli
presented at the start of the trial. **D.** Movement-aligned activity for the anterior, middle, and
posterior heads of deltoids for the two monkeys for single-target red trials, showing greater
recruitment for rightward than leftward reaches. Data shows the average of the mean recruitment
from each session, normalized to the peak movement-aligned activity from each session. Contours
subtend the mean +/- stderr.

**Figure 2. A.** Frequency histograms of reaction times (20 ms bins) for Monkey G (top row, n =
1303 trials) and Monkey B (bottom row, n = 2648 trials) across all experimental sessions. **B.**
Representative examples of express visuomotor responses in Monkey G (top row) and Monkey B
(bottom row), depicting trial-by-trial EMG recruitment aligned to target emergence to the left or
right. In these plots, trials are sorted by movement onset (black squares), and the color conveys
the magnitude of recruitment. **C.** For each monkey, the top row depicts mean EMG activity
subtended by the standard error for left (light blue) and right (dark blue) reaches, and bottom row
shows the time-series ROC (AUC = area under curve). The red rectangles in **B** and grey rectangles
in **C** show the 30 ms interval following the detection of the express visuomotor response.

**Figure 3.** Properties of express visuomotor responses. **A.** Latencies of express visuomotor
responses for both monkeys, showing latencies when all trials are considered regardless of reaction
time, and after including only the longer-than-average reaction times. Grey lines connect
observations from the same muscle from a given experimental day (grey + marks denote
representative data shown in Fig. 2). Black lines and squares show mean +/- standard error.

Asterisks show significant differences of paired t-test ($p < 0.05$). **B**. Calculation of the proportional

magnitude of the express visuomotor response, using the representative data from Monkey B. We

integrate the difference in the mean EMG curve for right versus left responses over the 30 ms after

the onset of the express visuomotor responses, and divide this by the integral of the 30 ms interval

surrounding peak EMG activity aligned to right reach onset. Same general format as **A**; asterisk

showing significant results of a student's t-test ($p < 0.05$).

**Figure 4.** Larger express visuomotor responses tended to precede shorter reach reaction times

(RTs). **A, B**. Representative data plotting magnitude of express visuomotor response as a function

of reach reaction time, from the representative sessions shown in Figure 2. Each data point

represents a single trial, and the black diagonal line shows the linear regression. The correlation

coefficient (r) is shown in the top-right of each subplot. **C**. Histogram of correlation coefficients

across all sessions in which an express visuomotor response was detected. Vertical dash line

indicates 0. Solid or grey lines or bins denote data for Monkey G or B, respectively. Vertical arrows

show mean r-values.

**Figure 5.** Contextual control on Double Target trials. **A.** Histograms across all sessions of reaction

times (20 ms bins) relative to the emergence of the red stimulus for Monkey G (top row) and

Monkey B (bottom row), for Double Target Red trials (left column, Monkey G: 632 trials, error

rate = 9.4%; Monkey B: 1788 trials, error rate = 9.8%) or Double Target Green trials (right column,

Monkey G: 413 trials, error rate = 50.1%; Monkey B: 1378 trials, error rate = 29.6%).

Upward/downward histograms represent correct/incorrect responses. **B**. Trajectory of horizontal

touchscreen position averaged across all trials for when the monkey should have reached right on

correct Double Target Red trials (red contours), correct Double Target Green trials (green

contours) and incorrect Double Target Green trials (black contours). Contours denote mean +/-

stderr. **C.** Depiction of mean EMG activity +/- stderr for when the red stimulus emerged to the

right. Grey rectangles denote the 30 ms interval following the express visuomotor response, over

which magnitude was calculated. **D,E**. The latency (**D**) or magnitude (**E**) of the express visuomotor

response, by trial type and/or performance, across 29 occurrences where an express visuomotor

response could be detected on both Double Target Red and Double Target Green trials. Grey lines

connect observations from the same muscle from a given experimental day. Black lines and

squares show mean +/- standard error. Asterisks show significant differences of Bonferroni

corrected paired t-test ($p < 0.05$).

**References**

Ames KC, Ryu SI, Shenoy KV (2019) Simultaneous motor preparation and execution in a last-
moment reach correction task. Nat Commun 10:2718.

Archambault PS, Caminiti R, Battaglia-Mayer A (2009) Cortical Mechanisms for Online Control
of Hand Movement Trajectory: The Role of the Posterior Parietal Cortex. Cerebral Cortex
19:2848–2864 Available at: http://dx.doi.org/10.1093/cercor/bhp058.

Archambault PS, Ferrari-Toniolo S, Battaglia-Mayer A (2011) Online control of hand trajectory
and evolution of motor intention in the parietofrontal system. Journal of Neuroscience
31:742–752.

Atsma J, Maij F, Gu C, Medendorp WP, Corneil BD (2018) Active Braking of Whole-Arm
Reaching Movements Provides Single-Trial Neuromuscular Measures of Movement
Cancellation. J Neurosci 38:4367–4382.

Benevento LA, Fallon JH (1975) The ascending projections of the superior colliculus in the
rhesus monkey (Macaca mulatta). J Comp Neurol 160:339–361.

Bibi R, Edelman JA (2009) The influence of motor training on human express saccade
production. J Neurophysiol 102:3101–3110.

Boehnke SE, Munoz DP (2008) On the importance of the transient visual response in the
superior colliculus. Curr Opin Neurobiol 18:544–551.

Bogadhi AR, Katz LN, Bollimunta A, Leopold DA, Krauzlis RJ (2021) Midbrain activity shapes
high-level visual properties in the primate temporal cortex. Neuron 109:690–699.e5.

Chen C-Y, Sonnenberg L, Weller S, Witschel T, Hafed ZM (2018) Spatial frequency sensitivity
in macaque midbrain. Nat Commun 9:1–13.

Cisek P, Kalaska JF (2005) Neural correlates of reaching decisions in dorsal premotor cortex:
specification of multiple direction choices and final selection of action. Neuron 45:801–814.

Cluff T, Scott SH (2016) Online Corrections are Faster Because Movement Initiation Must
Disengage Postural Control. Motor Control 20:162–170.

Contemori S, Loeb GE, Corneil BD, Wallis G, Carroll TJ (2021a) The influence of temporal
predictability on express visuomotor responses. J Neurophysiol 125:731–747.

Contemori S, Loeb GE, Corneil BD, Wallis G, Carroll TJ (2021b) Trial-by-trial modulation of
express visuomotor responses induced by symbolic or barely detectable cues. J
Neurophysiol 126:1507–1523.

Contemori S, Loeb GE, Corneil BD, Wallis G, Carroll TJ (2022a) Symbolic cues enhance
express visuomotor responses in human arm muscles at the motor planning rather than the
visuospatial processing stage. Journal of Neurophysiology 128:494–510 Available at:

683   http://dx.doi.org/10.1152/jn.00136.2022.

684 Contemori S, Loeb GE, Corneil BD, Wallis G, Carroll TJ (2022b) Express visuomotor responses
685   reflect knowledge of both target location and conscious intent during reaches of different
686   amplitudes. bioRxiv:2022.11.06.515363 Available at:
687   https://www.biorxiv.org/content/biorxiv/early/2022/11/06/2022.11.06.515363 [Accessed
688   March 1, 2023].

689 Corneil BD, Goonetilleke SC, Peel TR, Green KA, Welch ID (2012) Ultrasound-guided insertion
690   of intramuscular electrodes into suboccipital muscles in the non-human primate. J
691   Electromyogr Kinesiol 22:553–559.

692 Corneil BD, Munoz DP (2014) Overt responses during covert orienting. Neuron 82:1230–1243.

693 Corneil BD, Munoz DP, Chapman BB, Admans T, Cushing SL (2008) Neuromuscular
694   consequences of reflexive covert orienting. Nat Neurosci 11:13–15.

695 Corneil BD, Olivier E, Munoz DP (2004) Visual responses on neck muscles reveal selective
696   gating that prevents express saccades. Neuron 42:831–841.

697 Crammond DJ, Kalaska JF (1994) Modulation of preparatory neuronal activity in dorsal
698   premotor cortex due to stimulus-response compatibility. J Neurophysiol 71:1281–1284.

699 Dash S, Peel TR, Lomber SG, Corneil BD (2018) Frontal eye field inactivation reduces saccade
700   preparation in the superior colliculus but does not alter how preparatory activity relates to
701   saccades of a given latency. eNeuro 5 Available at:
702   https://www.eneuro.org/lookup/doi/10.1523/ENEURO.0024-18.2018.

703 Dash S, Peel TR, Lomber SG, Corneil BD (2020) Impairment but not abolishment of express
704   saccades after unilateral or bilateral inactivation of the frontal eye fields. J Neurophysiol
705   123:1907–1919.

706 Day BL, Brown P (2001) Evidence for subcortical involvement in the visual control of human
707   reaching. Brain 124:1832–1840.

708 Dickey AS, Amit Y, Hatsopoulos NG (2013) Heterogeneous neural coding of corrective
709   movements in motor cortex. Front Neural Circuits 7:51.

710 Divakar R, Loeb GE, Corneil BD, Wallis G, Carroll TJ (2022) A subcortical origin for rapid,
711   target-oriented corticospinal excitability changes during visually guided reaching.
712   bioRxiv:2022.12.13.520229 Available at:
713   https://www.biorxiv.org/content/biorxiv/early/2022/12/13/2022.12.13.520229 [Accessed
714   March 1, 2023].

715 Dorris MC, Paré M, Munoz DP (1997) Neuronal activity in monkey superior colliculus related to
716   the initiation of saccadic eye movements. J Neurosci 17:8566–8579.

717 Edelman JA, Keller EL (1996) Activity of visuomotor burst neurons in the superior colliculus

accompanying express saccades. J Neurophysiol 76:908–926.

Edelman JA, Kristjánsson A, Nakayama K (2007) The influence of object-relative visuomotor
set on express saccades. J Vis 7:12.

Elsley JK, Nagy B, Cushing SL, Corneil BD (2007) Widespread presaccadic recruitment of neck
muscles by stimulation of the primate frontal eye fields. J Neurophysiol 98:1333–1354.

Fautrelle L, Prablanc C, Berret B, Ballay Y, Bonnetblanc F (2010) Pointing to double-step visual
stimuli from a standing position: very short latency (express) corrections are observed in
upper and lower limbs and may not require cortical involvement. Neuroscience 169:697–
705.

Fischer B, Boch R (1983) Saccadic eye movements after extremely short reaction times in the
monkey. Brain Res 260:21–26.

Fischer B, Ramsperger E (1984) Human express saccades: extremely short reaction times of goal
directed eye movements. Exp Brain Res 57:191–195.

Fischer B, Weber H (1993) Express saccades and visual attention. Behav Brain Sci 16:553–567.

Gaveau V, Pisella L, Priot A-E, Fukui T, Rossetti Y, Pélisson D, Prablanc C (2014) Automatic
online control of motor adjustments in reaching and grasping. Neuropsychologia 55:25–40.

Gellman RS, Carl JR, Miles FA (1990) Short latency ocular-following responses in man. Vis
Neurosci 5:107–122.

Georgopoulos AP, Kalaska JF, Caminiti R, Massey JT (1983) Interruption of motor cortical
discharge subserving aimed arm movements. Exp Brain Res 49:327–340.

Glover IS, Baker SN (2019) Multimodal stimuli modulate rapid visual responses during
reaching. J Neurophysiol 122:1894–1908.

Goodale MA, Pelisson D, Prablanc C (1986) Large adjustments in visually guided reaching do
not depend on vision of the hand or perception of target displacement. Nature 320:748–750.

Goonetilleke SC, Katz L, Wood DK, Gu C, Huk AC, Corneil BD (2015) Cross-species
comparison of anticipatory and stimulus-driven neck muscle activity well before saccadic
gaze shifts in humans and nonhuman primates. J Neurophysiol 114:902–913.

Gu C, Andrew Pruszynski J, Gribble PL, Corneil BD (2018) Done in 100 ms: path-dependent
visuomotor transformation in the human upper limb. Journal of Neurophysiology
119:1319–1328 Available at: http://dx.doi.org/10.1152/jn.00839.2017.

Gu C, Pruszynski JA, Gribble PL, Corneil BD (2019) A rapid visuomotor response on the human
upper limb is selectively influenced by implicit motor learning. J Neurophysiol 121:85–95.

Gu C, Wood DK, Gribble PL, Corneil BD (2016) A Trial-by-Trial Window into Sensorimotor

Transformations in the Human Motor Periphery. J Neurosci 36:8273–8282.

Hafed ZM, Ignashchenkova A (2013) On the dissociation between microsaccade rate and
direction after peripheral cues: microsaccadic inhibition revisited. J Neurosci 33:16220–
16235.

Hwang J, Mitz AR, Murray EA (2019) NIMH MonkeyLogic: Behavioral control and data
acquisition in MATLAB. J Neurosci Methods 323:13–21.

Kearsley SL, Cecala AL, Kozak RA, Corneil BD (2022) Express arm responses appear
bilaterally on upper-limb muscles in an arm choice reaching task. J Neurophysiol 127:969–
983.

Kozak RA, Cecala AL, Corneil BD (2020) An Emerging Target Paradigm to Evoke Fast
Visuomotor Responses on Human Upper Limb Muscles. J Vis Exp Available at:
http://dx.doi.org/10.3791/61428.

Kozak RA, Corneil BD (2021) High-contrast, moving targets in an emerging target paradigm
promote fast visuomotor responses during visually guided reaching. J Neurophysiol
126:68–81.

Kozak RA, Kreyenmeier P, Gu C, Johnston K, Corneil BD (2019) Stimulus-Locked Responses
on Human Upper Limb Muscles and Corrective Reaches Are Preferentially Evoked by Low
Spatial Frequencies. eneuro 6:ENEURO.0301–0319.2019 Available at:
http://dx.doi.org/10.1523/eneuro.0301-19.2019.

Kurtzer I, Herter TM, Scott SH (2006) Nonuniform distribution of reach-related and torque-
related activity in upper arm muscles and neurons of primary motor cortex. J Neurophysiol
96:3220–3230.

Lamarre Y, Busby L, Spidalieri G (1983) Fast ballistic arm movements triggered by visual,
auditory, and somesthetic stimuli in the monkey. I. Activity of precentral cortical neurons. J
Neurophysiol 50:1343–1358.

Lara AH, Elsayed GF, Zimnik AJ, Cunningham JP, Churchland MM (2018) Conservation of
preparatory neural events in monkey motor cortex regardless of how movement is initiated.
Elife 7 Available at: http://dx.doi.org/10.7554/eLife.31826.

Li X, Basso MA (2008) Preparing to Move Increases the Sensitivity of Superior Colliculus
Neurons. Journal of Neuroscience 28:4561–4577 Available at:
http://dx.doi.org/10.1523/jneurosci.5683-07.2008.

Loeb GE, Gans C (1986) Electromyography for Experimentalists. University of Chicago Press.

Lovejoy LP, Krauzlis RJ (2010) Inactivation of primate superior colliculus impairs covert
selection of signals for perceptual judgments. Nat Neurosci 13:261–266.

Marino RA, Levy R, Boehnke S, White BJ, Itti L, Munoz DP (2012) Linking visual response

properties in the superior colliculus to saccade behavior. Eur J Neurosci 35:1738–1752.

Marino RA, Levy R, Munoz DP (2015) Linking express saccade occurance to stimulus
properties and sensorimotor integration in the superior colliculus. J Neurophysiol 114:879–
892.

Matelli M, Luppino G (1996) Thalamic input to mesial and superior area 6 in the macaque
monkey. The Journal of Comparative Neurology 372:59–87 Available at:
http://dx.doi.org/10.1002/(sici)1096-9861(19960812)372:1<59::aid-cne6>3.0.co;2-l.

Miles FA, Kawano K, Optican LM (1986) Short-latency ocular following responses of monkey.
I. Dependence on temporospatial properties of visual input. J Neurophysiol 56:1321–1354.

Munoz DP, Dorris MC, Paré M, Everling S (2000) On your mark, get set: brainstem circuitry
underlying saccadic initiation. Can J Physiol Pharmacol 78:934–944.

Murphy JT, Kwan HC, MacKay WA, Wong YC (1982) Precentral unit activity correlated with
angular components of a compound arm movement. Brain Res 246:141–145.

Nakajima T, Ohtsuka H, Irie S, Suzuki S, Ariyasu R, Komiyama T, Ohki Y (2021) Visual
information increases the indirect corticospinal excitation via cervical interneurons in
humans. J Neurophysiol 125:828–842.

Novembre G, Iannetti GD (2021) Towards a unified neural mechanism for reactive adaptive
behaviour. Prog Neurobiol 204:102115.

Olfert ED, Cross BM, McWilliam AA, Others (1993) Guide to the care and use of experimental
animals. Canadian Council on Animal Care Ottawa.

Paré M, Munoz DP (1996) Saccadic reaction time in the monkey: advanced preparation of
oculomotor programs is primarily responsible for express saccade occurrence. J
Neurophysiol 76:3666–3681.

Pastor-Bernier A, Tremblay E, Cisek P (2012) Dorsal premotor cortex is involved in switching
motor plans. Front Neuroeng 5:5.

Perfiliev S, Isa T, Johnels B, Steg G, Wessberg J (2010) Reflexive limb selection and control of
reach direction to moving targets in cats, monkeys, and humans. J Neurophysiol 104:2423–
2432.

Peysakhovich B, Tetrick SM, Silva AA, Li S, Zhu O, Ibos G, Johnston WJ, Freedman DJ (2023)
Primate superior colliculus is engaged in abstract higher-order cognition. bioRxiv Available
at: http://dx.doi.org/10.1101/2023.01.17.524416.

Pruszynski JA, Johansson RS, Flanagan JR (2016) A Rapid Tactile-Motor Reflex Automatically
Guides Reaching toward Handheld Objects. Curr Biol 26:788–792.

Pruszynski JA, King GL, Boisse L, Scott SH, Flanagan JR, Munoz DP (2010) Stimulus-locked

responses on human arm muscles reveal a rapid neural pathway linking visual input to arm
motor output. Eur J Neurosci 32:1049–1057.

Reimer J, Hatsopoulos NG (2010) Periodicity and evoked responses in motor cortex. J Neurosci
30:11506–11515.

Reschechtko S, Pruszynski JA (2020) Voluntary modification of rapid tactile-motor responses
during reaching differs from its visuomotor counterpart. J Neurophysiol 124:284–294.

Rezvani S, Corneil BD (2008) Recruitment of a head-turning synergy by low-frequency activity
in the primate superior colliculus. J Neurophysiol 100:397–411.

Riehle A, Kornblum S, Requin J (1997) Neuronal correlates of sensorimotor association in
stimulus–response compatibility. J Exp Psychol Hum Percept Perform 23:1708–1726.

Rohrer WH, Sparks DL (1993) Express saccades: the effects of spatial and temporal uncertainty.
Vision Res 33:2447–2460.

Saijo N, Murakami I, Nishida S 'ya, Gomi H (2005) Large-field visual motion directly induces
an involuntary rapid manual following response. J Neurosci 25:4941–4951.

Schiller PH, Sandell JH, Maunsell JH (1987) The effect of frontal eye field and superior
colliculus lesions on saccadic latencies in the rhesus monkey. J Neurophysiol 57:1033–
1049.

Schroeder CE, Molhom S, Lakatos P, Ritter W, Foxe JJ (2004) Human–simian correspondence
in the early cortical processing of multisensory cues. Cogn Process 5:140–151.

Selen LPJ, Corneil BD, Medendorp WP (2019) Single-trial dynamics of competing reach plans
in the human motor periphery. bioRxiv Available at:
https://www.biorxiv.org/content/10.1101/765180.abstract.

Sommer MA, Wurtz RH (2002) A Pathway in Primate Brain for Internal Monitoring of
Movements. Science 296:1480–1482 Available at:
http://dx.doi.org/10.1126/science.1069590.

Song J-H, Rafal RD, McPeek RM (2011) Deficits in reach target selection during inactivation of
the midbrain superior colliculus. Proc Natl Acad Sci U S A 108:E1433–E1440.

Sparks D, Rohrer WH, Zhang Y (2000) The role of the superior colliculus in saccade initiation: a
study of express saccades and the gap effect. Vision Res 40:2763–2777.

Stuphorn V, Hoffmann KP, Miller LE (1999) Correlation of primate superior colliculus and
reticular formation discharge with proximal limb muscle activity. J Neurophysiol 81:1978–
1982.

Suzuki S, Nakajima T, Irie S, Ariyasu R, Ohtsuka H, Komiyama T, Ohki Y (2022) Subcortical
Contribution of Corticospinal Transmission during Visually Guided Switching Movements

of the Arm. Cereb Cortex 32:380–396.

Tian X, Yoshida M, Hafed ZM (2016) A Microsaccadic Account of Attentional Capture and
Inhibition of Return in Posner Cueing. Front Syst Neurosci 10:23.

Veerman MM, Brenner E, Smeets JBJ (2008) The latency for correcting a movement depends on
the visual attribute that defines the target. Exp Brain Res 187:219–228.

Werner W, Dannenberg S, Hoffmann KP (1997a) Arm-movement-related neurons in the primate
superior colliculus and underlying reticular formation: comparison of neuronal activity with
EMGs of muscles of the shoulder, arm and trunk during reaching. Exp Brain Res 115:191–
205.

Werner W, Hoffmann KP, Dannenberg S (1997b) Anatomical distribution of arm-movement-
related neurons in the primate superior colliculus and underlying reticular formation in
comparison with visual and saccadic cells. Exp Brain Res 115:206–216.

Wood DK, Gu C, Corneil BD, Gribble PL, Goodale MA (2015) Transient visual responses reset
the phase of low-frequency oscillations in the skeletomotor periphery. European Journal of
Neuroscience 42:1919–1932 Available at: http://dx.doi.org/10.1111/ejn.12976.