Done in 65 ms: Express visuomotor responses in upper limb muscles in Rhesus Macaques

Running head: Express visuomotor responses in macaques

Aaron L. Cecala[1,2], Rebecca A. Kozak[3], J. Andrew Pruszynski[1,2], Brian D. Corneil[1,2,4]

[1]Department of Physiology and Pharmacology, Western University, London, ON, Canada, N6A 5B7
[2]Robarts Research Institute, 1151 Richmond St. N, London, ON, Canada, N6A 5B7
[3]Graduate Program in Neuroscience, Western University, London, ON, Canada, N6A 5B7
[4]Department of Psychology, Western University, London, ON, Canada, N6A 5B7

Corresponding Author: Brian D. Corneil: Robarts Research Institute, 1151 Richmond Street North, Western University, London, ON, Canada, N6A 5B7; E-mail: bcorneil@uwo.ca

Number of pages: 36
Number of figures: 6
Number of tables: 0
Number of words:
    Abstract: 239
    Introduction: 690
    Discussion: 2285

Conflicts of interest: The authors declare no competing financial interests.

Keywords: rhesus monkey; visually-guided reaching; electromyography; sensorimotor transformation

Acknowledgements: The authors thank Katherine Faubert, Rhonda Kersten, Kristy Gibbs, Darren Pitre, and Matthew Reeves for their technical assistance and animal husbandry support, and Kevin Barker for his apparatus design and production expertise. We thank Tim Carrol, Sam Contemori, Rechu Divakar, Sebastian Lehmann, and Gerald Loeb for feedback on earlier versions of this manuscript.

Grants: This work was supported by operating grants to BDC from the Canadian Institutes of Health Research (CIHR) [MOP-93796, -142317; PJT-180279] and the Natural Sciences and Engineering Research Council of Canada (NSERC) [RGPIN-311680, -04394-2021], and by the Canada First Research Excellence Fund to BrainsCAN. RAK was supported by an Ontario Graduate Scholarship. JAP received a salary award from the Canada Research Chairs Program.

Abstract

41
How rapidly can the brain transform vision into action? Work in humans has established that the
transformation for visually-guided reaching can be remarkably rapid, with the first phase of upper
limb muscle recruitment, the *express visuomotor response*, beginning within less than 100 ms of
visual target presentation. Such short-latency responses limit the opportunities for extensive
cortical processing, leading to the hypothesis that they are generated via the subcortical
tectoreticulospinal pathway. Here, we examine if non-human primates (NHPs) exhibit express
visuomotor responses. Two male macaques made visually-guided reaches in a behavioral
paradigm known to elicit express visuomotor responses in humans, while we acquired
intramuscular recordings from the deltoid muscle. Across several variants of this paradigm,
express visuomotor responses began within 65 ms (range 48–91 ms) of target presentation.
Although the timing of the express visuomotor response did not co-vary with reaction time, larger
express visuomotor responses tended to precede shorter latency reaches. Finally, the magnitude of
the express visuomotor response was muted on trials where NHPs withheld a reach to one stimulus
in order to move to a stimulus appearing 34 ms later in the opposite direction. Overall, the response
properties and contextual control of express visuomotor responses in NHPs resemble those in
humans. Our results establish a new benchmark for visuomotor transformations underlying
visually-guided reaches, setting the stage for experiments that can directly compare the role of
cortical and subcortical areas in reaching when time is of the essence.

Significance statement:

Express visuomotor responses in upper limb muscles are brief periods of recruitment preceding
visually-guided reaches. Such responses begin ~90 ms after visual target presentation in humans,
and potentially arise from signaling along the tecto-reticulo-spinal pathway. Here, we show that
express visuomotor responses in macaques upper limb muscles resemble those in humans,
excepting that they evolve ~65 ms after target onset, consistent with shorter responses latencies in
macaques versus humans. Our results clock the completion of the visuomotor transformation for
rapid reaching, and set the stage for experiments to directly test the underlying substrates.

69
70
71

Introduction

When required, the brain has a remarkable capacity to rapidly transform sensory input into motor output. Expedited responses can consist of eye movements to evaluate a potential threat, or mid-flight adjustments when reaching to a slipping cell phone. In some cases, the brain relies on subcortical circuits that provide a short-latency link between sensation and action, which are nested within broader fronto-parietal circuits mediating more deliberative actions. This appears to be the case for express saccades, which rely on the integrity of the midbrain superior colliculus (SC) and are potentiated by, but not critically reliant on, input from fronto-parietal circuits (Schiller et al., 1987; Edelman and Keller, 1996; Dorris et al., 1997; Dash et al., 2018, 2020). Although the SC has been implicated in mid-flight reach corrections (Day and Brown, 2001; Pruszynski et al., 2016; Reschechtko and Pruszynski, 2020; Suzuki et al., 2022) and interceptions of rapidly moving targets (Perfiliev et al., 2010), debates remain about the comparative role of subcortical versus cortical areas in expedited visually-guided reaching (Fautrelle et al., 2010; Gaveau et al., 2014; Novembre and Iannetti, 2021).

To better understand the neural mechanisms underlying expedited reaches, we and others have studied the timing and response properties of the first change of upper limb muscle recruitment elicited by visual stimulus onset. Doing so establishes the benchmark for how rapidly premotor circuits transform vision into action, and inferences can be drawn by comparing the timing and magnitude of muscle activity to response properties in candidate premotor areas. Such work has revealed that the first wave of upper limb muscle recruitment driven by stimulus presentation evolves within less than 100 ms in humans, regardless of the ensuing reaction time (Pruszynski et al., 2010; Wood et al., 2015; Gu et al., 2016; Kozak et al., 2019; Contemori et al., 2021a). This phenomenon has been termed the stimulus-locked response (Pruszynski et al., 2010) or fast or rapid visuomotor response (Glover and Baker, 2019; Kozak and Corneil, 2021), but lately we have adopted the term express visuomotor response (Contemori et al., 2021a, 2021b) to emphasize similarities with both express saccades and the timing and response properties of visual bursts on premotor cells in the motor layers of the SC (Corneil and Munoz, 2014). The timing and magnitude of such SC visual responses, express saccades, and express visuomotor responses on upper limb muscles are all potentiated by high-contrast, low spatial frequency stimuli (Marino et al., 2012,

2015; Wood et al., 2015; Chen et al., 2018; Kozak et al., 2019; Kozak and Corneil, 2021), and by

a variety of cognitive processes promoting premotor preparation (Rohrer and Sparks, 1993; Paré

and Munoz, 1996; Munoz et al., 2000; Edelman et al., 2007; Boehnke and Munoz, 2008; Li and

Basso, 2008; Bibi and Edelman, 2009; Gu et al., 2018; Contemori et al., 2021b, 2022a).

Here we aim to establish whether express visuomotor responses can be observed in non-human

primates (NHPs). We examine three response properties. First, express saccades and the related

phenomenon of express neck muscle responses occur ~20 ms earlier in NHPs than humans (Fischer

and Boch, 1983; Fischer and Ramsperger, 1984; Corneil et al., 2004; Goonetilleke et al., 2015). If

evoked by a similar mechanism, express visuomotor responses in arm muscles in NHPs should

evolve ~60-80 ms after stimulus onset. Second, although more time-locked to stimulus rather than

movement onset, larger express visuomotor responses in humans consistently precede shorter

reaction times (Pruszynski et al., 2010; Gu et al., 2016; Contemori et al., 2021a); we predict a

similar inverse relationship between recruitment magnitude and reaction time in NHPs. Third, the

magnitude of express visuomotor responses in humans is dampened if subjects know to move

away from or occasionally not respond to the stimulus (Wood et al., 2015; Gu et al., 2016; Atsma

et al., 2018; Kozak et al., 2020; Contemori et al., 2021b, 2022a). Such dampening is a signature of

top-down cortical control of a subcortical circuit, and we predict that similar top-down control can

be instantiated in NHPs. Overall, we find that NHPs readily generate express visuomotor responses

consistent with all these properties, establishing the NHP as a suitable animal model to study

underlying neural mechanisms.

Methods

Subjects and Surgical Procedures

Two male rhesus monkeys (*Macaca mulatta;* monkey G and monkey B), weighing 8.5 and 13.1

kg respectively, were used in these experiments. All training, surgical, and experimental

procedures were in accordance with the Canadian Council on Animal Care policy on the use of

laboratory animals (Olfert et al., 1993) and approved by the Western University Animal Use

Subcommittee. We monitored the monkeys' weights daily and their health was under the close

supervision of the university veterinarians.

Each animal underwent an aseptic procedure in which a head post was attached to the skull by way
of a dental acrylic pedestal prior to behavioral training. During training and experimental sessions,
animals were seated with the head-restrained in a customized primate chair that restricted
movements of the left arm, torso and hips while permitting unrestricted movements of the right
arm in a three-dimensional workspace. Testing was performed in a dimly illuminated room and
the monkeys were monitored via a video camera. Visual stimuli, behavioral control, and data
acquisition were controlled by MonkeyLogic (Hwang et al., 2019). All visual stimuli were
presented on a 42-in. color touch-sensitive monitor (4202L Elo Touch Solutions, Inc., Milpitas,
CA) positioned 33 cm in front of the monkeys. The monitor had a spatial resolution of 1920 ×
1080 pixels and a refresh rate of 60 Hz.

The Emerging Target Task
Monkeys generated visually-guided reaches to stimuli presented within the context of an emerging
target task (Kozak et al., 2020). The task involves a moving stimulus that temporarily disappears
behind an occluder, and then emerges in motion at either the left or right outlet. This task was
chosen as it engenders express visuomotor responses in most human subjects (Contemori et al.,
2021a, 2021b; Kozak and Corneil, 2021; Kearsley et al., 2022), likely due to the combination of
salient visual stimulus presentation following a fixed period of implied motion.

Data was collected from three variants of the emerging target task (**Fig. 1**). All trials started with
a red or green 'start position' stimulus that appeared at a central location below a gray rectangular
occluder. The monkeys were required to touch and hold the location of this start position stimulus
within a computer-defined window of 7 cm. Following this, a red or green 'target' stimulus that
matched the color of the start position stimulus appeared above the occluder. Providing the monkey
kept in contact with the start position stimulus for 500 ms, this target stimulus dropped vertically
at a constant velocity of 20 cm/s, and then disappeared behind the occluder. After a fixed latency
of 300 ms, a red target then emerged in motion below the occluder at either the right or left outlet.
At the same time, the start position stimulus disappeared and a secondary visual stimulus that lay
under a photodiode (and hence was unseen by the monkey) was presented for data alignment
purposes. On double target trials, a green target emerged in motion at the other outlet 34 ms later.
To obtain a reward, the monkeys had to maintain contact on the start position until target

emergence below the barrier, and then reach to touch the colored target that matched the color of
the stimuli presented at the start of the trial before it disappeared off the screen (i.e., reach to the
red target on single target red (**Fig. 1A**) and double target red (**Fig. 1B**) trials, or to the green target
on double target green trials **(Fig. 1C))**. Thus, correct performance on double target green trials
required the monkeys to not reach towards the red target that emerged slightly (34 ms) earlier. The
moving target that appeared below the barrier started 15 cm to the right or left of the start position
stimulus, and then moved in an lateral-downward direction (45 deg below horizontal) at 20 cm/s.
Trials where the monkey touched or moved toward the wrongly colored target were tabulated as
incorrect reaches, and retained for further analysis. We decided to emphasize response speed over
accuracy, and hence used large acceptance windows (radius of 11 cm) around the emerging target.
An analysis of endpoints showed that both monkeys tended to reach more accurately than this
window, with the standard deviations of horizontal or vertical endpoints being 2.5 and 1.9 cm for
monkey G and 1.5 and 1.0 cm for monkey B. Recent work in humans (Contemori et al., 2022b)
shows that larger express visuomotor responses are observed when subjects land accurately on
target, as opposed to deliberately overshooting or undershooting the target. Thus, we do not believe
our results are influenced by our choice to emphasize response speed over accuracy.

Within a given block of trials, all trial conditions (defined by target location and, when used, task
variant) were presented with equal frequency but randomly shuffled. Some datasets consisted of
only single-target red trials (2 sessions for monkey G, 1 session for monkey B). For the other days,
collected data in a block of ~100 single-target red trials, prioritizing such data as EMG recording
electrodes occasionally were removed as animals performed the task (this happened in one day for
monkey G). After these 100 single-target red trials, we then intermixed the other task variants.
Across all experimental sessions, we obtained muscle recordings from a minimum of 380 to a
maximum of 1470 trials.

Figure 1. A-C. Emerging Target Task. At the start of each trial, either a red (**A, B**) or green (**C**) stimulus appears at the start position (red or green squares), below an occluder (gray box). The monkey touches and holds the screen at this start position. A target of the same color then appears above the occluder and moves down the chute before disappearing briefly behind the occluder for 300 ms. In Single Target Red trials (**A**), a single red target emerges in motion below the occluder at one of two outlets. In the Double Target Red (**B**) and Double Target Green (**C**) trials, appearance of the red target in one outlet was followed 34 ms later by appearance of a green target at the other outlet. The monkey was rewarded for reaching to the target that matched the color of the stimuli presented at the start of the trial. **D**. Movement-aligned activity for the anterior, middle, and posterior heads of deltoids for the two monkeys for single-target red trials, showing greater recruitment for rightward than leftward reaches. Data shows the average of the mean recruitment from each session, normalized to the peak movement-aligned activity from each session. Contours subtend the mean +/- stderr.

EMG electrode insertion and data collection

Prior to a data collection session, monkeys were lightly sedated before intramuscular, fine-wire

EMG electrodes were acutely inserted into the posterior, medial, and anterior heads of the deltoid

muscle. Sedation followed an established procedure (Corneil et al., 2012), involving induction

with a low-dose of medetomidine, and reversal after electrode insertion with atipamazole. After

reversal, monkeys were given ~20 minutes to recover prior to beginning data collection. For each

intramuscular recording, we inserted two electrodes into the muscle belly staggered by ~1 cm to

enable bipolar recordings of bulk muscle activity. Each electrode was fabricated in-house by

threading PFA-coated 7-strand stainless steel wire (Type 316, A-M Systems, Sequim, WA) into a

27 gauge cannula, and then stripping the final ~3 mm of insulation. Anatomically, the different

heads of the deltoid were identified by manually palpating the head of the humerus. A ground

electrode was positioned on the dorsal neck approximately between the scapulae. Intramuscular

EMG activity was differentially amplified and recorded with a Myopac Junior system (Run

Technologies, low-pass filter modified to 2 kHz) and digitized at a rate of 1,000 Hz onto a Plexon

Omniplex system, which also received the photodiode signal. Off-line, any DC offset was

removed and EMG data were full-wave rectified and smoothed with a 7-point running average.

Our sampling frequency was limited to 1,000 Hz, which undersamples intramuscular signals. We

confirmed that such undersampling had only a minor influence on our results by collecting data

from one additional session on a Ripple Grapevine system that sampled EMG data at 30,000 Hz.

This allowed us to compare results obtained from 3 simultaneously recorded muscles processed

offline by either downsampling the EMG signal to 1,000 Hz versus that obtained by first full-wave

rectifying the 30 kHz signal followed by integration into 1 ms bins (Loeb and Gans, 1986; Elsley

et al., 2007). For all three muscles recorded at 30 kHz, these different processing steps changed

the discrimination latency of the express visuomotor response by only 1 ms. Thus, while our

undersampled data would not be appropriate for identifying single motor-unit action potentials, it

does accurately capture the timing of changes in multi-unit recruitment after visual target

presentation.

Data Analysis

All data analyses were conducted in MATLAB® (R2021a; Natick, MA, USA). We rejected trials

where the monkey either did not respond or failed to respond within 500 ms, as well as trials where

movements began within less than 80 ms of target emergence below the barrier. Reach RT was

calculated as the time from target emergence below the barrier (indicated by the photo-diode

signal) to the time point where the monkey's hand was either no longer in contact with the

touchscreen at the start position, or when the monkey's hand started deviating toward the emerging

target. Offline, we further inspected all trials within a graphical user interface, allowing us to adjust

the movement onset mark at a resolution of 1 kHz, and flag trials where the initial movement was

in the opposite direction of the correct response. Monkey G moved more rapidly and tended to

remain in contact with the screen during the reach (i.e., 'smearing' their hand across the screen),

whereas Monkey B moved more slowly and had a greater tendency to lift their finger off the screen

when reaching. Thus, incorrect reaches in Monkey G included trials where the monkey first moved

in the incorrect direction before correcting their reach in mid-flight. In contrast, incorrect reaches

in Monkey B were sometimes only identified when the monkey touched the wrong location. On

other occasions, an incorrect reach in Monkey B was identified by a deflection in the horizontal

touch screen position in the wrong direction, as identified in the user interface.

The different heads of the deltoid muscle are known to have different preferred reaching directions

in the rhesus macaque (Kurtzer et al., 2006). Note however that the reaching task in this study

required that the monkeys reach forward and maintain contact with the touchscreen at the start

point, and then upon target emergence move their arm to the right or left to obtain a reward. In the

context of this bidirectional task, as shown in **Fig. 1D**, all heads of deltoid exhibited greater

recruitment before rightward versus leftward reaches on the touchscreen. Therefore, for the

purpose of analyzing the express visuomotor response, the rightward direction was deemed to be

the preferred direction of all muscles.

As with previous work (Pruszynski et al., 2010; Gu et al., 2016; Contemori et al., 2021a), we used

a time-series receiver-operating characteristic (ROC) analysis to define the presence and latency

of the express visuomotor response. This analysis method is suitable when targets can appear at

one of two diametrically opposite locations, although we note that express visuomotor responses

have been studied with other methods with different potential target configurations (Gu et al.,

2019; Kearsley et al., 2022; Selen et al., 2023), and with single-trial analyses (Contemori et al.,

2021b, 2022a). Data recorded from each muscle was treated as an independent sample, and data

was segregated by side of target emergence and task variant. For blocks with multiple interleaved

task variants, separate time-series ROCs were run on data for each task variant, comparing EMG

activity after left or right target emergence. The ROC value (i.e., the area under the ROC curve)

was calculated and plotted at every time sample (1 ms) from 100 ms before to 500 ms after target

emergence (e.g., **Fig. 2C**); this value indicates the likelihood of discriminating the side of target

emergence based on EMG activity alone. A ROC value of 0.5 indicates chance discrimination,

whereas values of 1 or 0 indicate perfectly correct or incorrect discrimination, respectively.

Although time-series ROCs often fluctuated around 0.5 in the 100 ms surrounding target

emergence, we often noted the presence of linear trends in this interval as well as a high degree of

variance on task variants with fewer overall trials. Such trends have been observed previously in

previous studies (Goonetilleke et al., 2015), and we suspect they arise from trials with short

reaction times where the animal was guessing, which are included or excluded if the animal

guessed correctly, or not. These linear trends precluded the use of fixed thresholds to determine

the latency of any express response. Accordingly, we first detrended the time-series ROC based

on a linear regression run on the time-series ROC over a baseline interval spanning from 100 ms

before to 45 ms after target emergence. We then set a discrimination threshold to be the first of 8

of 10 points where the time-series ROC exceeded 3 standard deviations above the mean of the

detrended baseline time-series ROC. If this time was between 45 to 100 ms after target emergence,

then the muscle was deemed to be exhibiting an express visuomotor response, and the latency was

taken to be the discrimination time. The magnitude of the express visuomotor response was

determined by the integral of EMG activity measured over the next 30 ms. All EMG data are

reported in source microvolts, and were not subjected to any normalization unless otherwise noted.

Experimental design and statistical analyses

All statistical analyses were performed in MATLAB® (R2021a; Natick, MA, USA). Comparisons

of the magnitude and latencies of express visuomotor responses and of reaction times across the

various trial types were performed via *t*-tests, using paired and unpaired *t*-tests when appropriate.

Correlational analyses were performed on latency, magnitude, and RT measures, as described

below. Comparisons of error rates across different task variants were performed using a chi-

squared test.

Results

Two rhesus macaques reached to red or green targets presented in the context of the emerging

target task (**Fig. 1A-C**). This task readily elicits express visuomotor responses in humans (Kozak

et al., 2020; Contemori et al., 2021a; Kozak and Corneil, 2021). Our goal was to establish the

phenomenon of express visuomotor responses in monkey, by: i) describing the response latencies

and consistency of such responses both within and across experimental sessions, ii) investigating

the relationships with ensuing reach reaction times (RTs), and iii) establishing whether the express

visuomotor response to the red stimulus would be decreased when monkeys were instructed to

reach in the opposite direction to a delayed emerging green stimulus. Our dataset consists of EMG

recordings made in 8 sessions in monkey G (7, 8, and 7 successful recordings from anterior,
medial, and posterior deltoid respectively) and 10 sessions in monkey B (9, 8, and 8 from anterior,
medial, and posterior deltoid respectively). As all muscles exhibited greater recruitment for
rightward vs leftward reaches on the touchscreen (**Fig. 1D**), and since we noted no differences in
the express visuomotor responses from the three heads of deltoid, we treated each recording as an
independent sample of multiple motor units, and pooled data within a given monkey.
299
Stimulus emergence evokes short-latency express visuomotor responses
We begin by describing the behavior and patterns of muscle recruitment immediately following
target emergence in single target red trials. On such trials, Monkey G reached within 179 ± 61 ms
(median = 170 ms) and Monkey B reached within 223 ± 47 ms (median = 220 ms) (**Fig. 2A**). Both
animals made some reaching errors when only a single red target emerged (6% for Monkey G, 5%
for Monkey B), but more commonly anticipated target emergence (8.6% for Monkey G, 12.1% for
Monkey B). Representative examples of EMG data recorded from each monkey are shown in **Fig.**
2B and **C**. As animals held their arm against gravity in order to touch the central starting position,
there was a modest level of recruitment prior to and immediately after target emergence. Very
soon after target emergence, muscle recruitment increased or decreased following target
emergence to the right or left, respectively. Such lateralization is a hallmark of express visuomotor
responses in humans (Pruszynski et al., 2010; Wood et al., 2015), and emphasizes that express
visuomotor responses are tuned to stimulus location rather than being a generic startle or
preparatory response (Gu et al., 2019; Selen et al., 2023). Another defining feature of express
visuomotor responses in humans is the trial-by-trial consistency in which this initial change in
muscle recruitment is more locked to target emergence rather than movement onset. This indeed
was the case in the representative examples shown in **Fig. 2B** (see red rectangles): note how this
initial lateralized change in muscle activity occurred ~60-70 ms after target emergence, well before
the hand moved or lifted away from the starting location. Following the express visuomotor
response, rightward reaches were preceded by a gradually increasing level of recruitment that
peaked just before reach initiation, whereas muscle recruitment before leftward reaches persisted
near or below the baseline level of activity prior to target emergence. The data shown for Monkey
B exhibited a second target-locked decrease in muscle activity ~130-140 ms after left target
emergence, resembling reports of oscillations in muscle recruitment initiated by a visual stimulus

in humans (Wood et al., 2015). Such double-dips were seen in about half of the recordings in Monkey B but were not observed in Monkey G. We suspect this is because monkey G had generally shorter RTs, so that the recruitment aligned with reach onset overlapped with any subsequent activity following the express visuomotor response.

Figure 2. A. Frequency histograms of reaction times (20 ms bins) for Monkey G (top row, n = 1303 trials) and Monkey B (bottom row, n = 2648 trials) across all experimental sessions. Solid/ dashed lines depict means/medians **B.** Representative examples of express visuomotor responses in Monkey G (top row) and Monkey B (bottom row), depicting trial-by-trial EMG recruitment aligned to target emergence to the left or right. In these plots, trials are sorted by movement onset (black squares), and the color conveys the magnitude of recruitment. **C.** For each monkey, the top row depicts mean EMG activity subtended by the standard error for left (light blue) and right (dark blue) reaches, and bottom row shows the time-series ROC (AUC = area under curve). Solid vertical lines depict mean reaction time for the representative examples. The red rectangles in **B** and grey rectangles in **C** show the 25 ms interval following the detection of the express visuomotor response.

To analyze the timing of express visuomotor responses, we performed a time-series receiver operating characteristics (ROC) analysis. This analysis determines the timing at which information following leftward or rightward target emergence influenced muscle recruitment. For the examples shown in **Fig. 2C**, the express visuomotor response began 68 ms (Monkey G) or 55 ms (Monkey B) after target emergence. In both examples, the time-series ROC rose quickly after this discrimination time, and remained elevated for the next ~200 ms.

We repeated this analysis across our sample of deltoid recordings. Across our sample, an express visuomotor response, which was defined as occurring within 45-100 ms, was detected in 77% (17 of 22; 7, 8, and 7 recordings from anterior, middle, and posterior deltoid) recordings in monkey G and in all 25 recordings in monkey B (9, 8, and 8 recordings from anterior, middle, and posterior

deltoid). When present, express visuomotor responses began very soon after stimulus emergence,

with discrimination latencies being 65 ± 8 ms (range: 52 to 81 ms) for monkey G and 64 ± 11 ms

(range: 48 to 91 ms for monkey B; **Fig. 3A**; no difference between monkeys, *t*-test, $t(40) = 0.34$, *p*

$= 0.72$). A potential concern about these short discrimination latencies is that they may arise only

because of recruitment associated with short-RT reaches. We therefore repeated the time-series

ROC analysis across our sample only on the slower half or all trials. Excluding the shorter-than-

average RT trials in each session increased reach reaction times by ~25 ms to 205 ± 63 ms for

Monkey G and 253 ± 37 ms for Monkey B. Express visuomotor responses persisted in 72% (16 of

22) of recordings in monkey G and in all 25 recordings in monkey B, with average discrimination

latencies being at 69 ± 7 ms for monkey G and 68 ± 12 ms for monkey B (**Fig. 3A**). Although

significant in both monkeys (paired *t*-test, monkey G: *t*-test, $t(15) = -6.9, p < 10^{-5}$; monkey B: $t(24)$

$= -3.10, p = 0.005$), the ~4 ms increase in discrimination latency after excluding shorter-than-

average RTs was less than the ~25 ms increase in mean RT. We suspect that slightly longer

discrimination latencies reflects lower magnitude express visuomotor responses associated with

longer RT trials (see next section), as reported previously (Contemori et al., 2021a, 2021b).

Figure 3. Properties of express visuomotor responses. **A.** Latencies of express visuomotor responses for both monkeys, showing latencies when all trials are considered regardless of reaction time, and after including only the longer-than-average reaction times. Gray lines connect observations from the same muscle from a given experimental day (gray + marks denote representative data shown in Fig. 2). Black lines and squares show mean +/- standard error. Asterisks show significant differences of paired t-test (p < 0.05). **B.** Calculation of the proportional magnitude of the express visuomotor response, using the representative data from Monkey B. Using data normalized to the peak of movement-aligned activity, we integrate the difference in the mean EMG curve for right versus left responses over the 25 ms after the onset of the express visuomotor responses, and divide this by the integral of the 25 ms interval surrounding peak EMG activity aligned to right reach onset. Same general format as **A;** asterisk showing significant results of a student's t-test (p < 0.05).

Previous work in humans has shown that express visuomotor responses observed in the emerging

target task can be quite large in magnitude, occasionally equalling if not exceeded 'voluntary'

recruitment magnitude achieved when recruitment is aligned to reach onset (Contemori et al.,

2021a; Kozak and Corneil, 2021). To establish the proportional magnitude of the express

visuomotor response, we expressed it relative to a 25 ms interval surrounding the peak magnitude

of movement-aligned recruitment (**Fig. 3B**). Expressed in this fashion, the express visuomotor

response reached 0.33 ± 0.21 (range: 0.08 to 0.81) for monkey G and 0.25 ± 0.11 (range: 0.03 to

0.48) for monkey B (proportional magnitudes were not significantly larger in monkey G than

monkey B; *t*-test, $t(40) = 1.76$, $p = 0.08$). Thus, on average, the recruitment magnitude of the

express visuomotor response was about one quarter to one third of the magnitude of reach-related

recruitment.

Express visuomotor response magnitude inversely correlates with RT

Although the express visuomotor response may be modest in magnitude, previous work shows that

larger express visuomotor responses tend to precede short RTs (Pruszynski et al., 2010; Gu et al.,

2016; Contemori et al., 2021a). To investigate whether a similar inverse correlation is seen in

NHPs, we calculated the trial-by-trial magnitude of EMG recruitment during an express

visuomotor response window, integrated over 25 ms after the express visuomotor response was

detected (Pruszynski et al., 2010), and plotted it as a function of RT. **Figure 4** shows these plots

for the representative data from monkey G (**Fig. 4A**) and monkey B (**Fig. 4B**), and both show the

inverse correlation between magnitude and RT expected from previous studies (Monkey G: $r = -

0.56$, $p < 10^{-9}$; Monkey B: $r = -0.37$, $p < 10^{-6}$). We repeated this analysis for every session in our

sample, and for both monkeys, observed the expected negative correlation (**Fig. 4C**; Monkey G:

mean $r = -0.45 \pm 0.26$. Monkey B: mean $r = -0.29 \pm 0.16$). The distributions for both monkeys lay

significantly below zero (*t*-test; Monkey G: $t(16) = -7.3$, $p < 10^{-5}$. Monkey B: $t(24) = -9.4$, $p < 10^{-}$

[8]).

Figure 4. Larger express visuomotor responses tended to precede shorter reach reaction times (RTs). **A, B**. Representative data plotting magnitude of express visuomotor response as a function of reach reaction time, from the representative sessions shown in Figure 2. Each data point represents a single trial, and the black diagonal line shows the linear regression. The correlation coefficient (*r*) is shown in the top-right of each subplot. **C.** Histogram of correlation coefficients across all sessions in which an express visuomotor response was detected. Vertical dash line indicates 0. Solid or gray lines or bins denote data for Monkey G or B, respectively. Vertical arrows show mean *r*-values.

Contextual control influences the magnitude but not timing of the express visuomotor response

Having established that macaque upper limb muscle recruitment following the emergence of a

single target resembles that observed in humans, we now turn to data recorded when two targets

emerged below the barrier. The sequence of target emergence was the exact same on double-target

trials; the red stimulus always appeared 300 ms after the initial descending stimulus disappeared

behind the barrier, and the green stimulus always appeared 32 ms after the red stimulus. Monkeys

reached to the emerging stimulus that matched the color of the central hold stimulus and dropping

stimulus (**Fig. 1B,C**). Double target green trials therefore required monkeys to not reach to the red

emerging stimulus in order to reach to the green stimulus that subsequently emerged on the

opposite side. Thus, monkeys had to exert a different degree of contextual control in how they

responded to the red emerging stimulus on double target red than double target green trials.

We first compared reaching behavior on double target red vs double target green trials. RTs for

correct responses (timed to the emergence of the red stimulus) were longer for double target green

trials than double target red trials (**Fig. 5A**, Monkey G: double target red = 188 ± 70 ms, n = 572;

double target green = 209 ± 97 ms, n = 206; $t(776) = -3.36$, $p < 0.001$. Monkey B: double target

red = 235 ± 50 ms, n = 1607; double target green = 255 ± 49 ms, n = 962; $t(2579) = -9.63$, $p < 10^{-10}$

^{10}). Both monkeys made reaching errors at a significantly higher rate on double target green than

double target red trials (Monkey G: chi square-test, $X^2 (1) = 33.9$, $p < 10^{-8}$; Monkey B: $X^2 (1) =$

11.25; $p < 10^{-3}$), and Monkey G generated more errors than Monkey B on double target green trials

(chi-square-test, $X^2 (1) = 5.21$; $p = 0.02$). When generated, such errors started earlier than correct

responses on either double target green or double target red trials (Monkey G: error RTs on double

target green = 172 ± 45 ms, n = 207; t-test against RTs of correct responses on double target green

trials $t(411) = 5.01$, $p < 10^{-7}$; t-test against RTs of correct responses on double target red trials

$t(777) = 3.06$, $p < 0.005$; Monkey B: error RTs on double target green = 186 ± 50 ms, n = 409; t-

test against RTs of correct responses on double target green trials $t(1376) = 23.38$, $p < 10^{-10}$; t-test

against RTs of correct responses on double target red trials $t(2019) = 17.62$, $p < 10^{-10}$).

Figure 5. Contextual control on Double Target trials. **A.** Histograms across all sessions of reaction times (20 ms bins) relative to the emergence of the red stimulus for Monkey G (top row) and Monkey B (bottom row), for Double Target Red trials (left column, Monkey G: 632 trials, error rate = 9.4%; Monkey B: 1788 trials, error rate = 9.8%) or Double Target Green trials (right column, Monkey G: 413 trials, error rate = 50.1%; Monkey B: 1378 trials, error rate = 29.6%). Upward/downward histograms represent correct/incorrect responses. Solid/dashed lines depict histogram means/medians. **B.** Trajectory of horizontal touchscreen position averaged across all trials for Double Target Red trials to the right or left (red contours), and Double Target Green trials, showing correct trials when the monkey should have reached left (green contours), and incorrect trials when the monkey should have reached right (black contours). Note how incorrect trials initially go left before being corrected mid-flight to go to the right. Contours show mean +/- stderr. Solid colored vertical lines in B depict mean reaction time for corresponding trial type. **C.** Same data, depicting vertical position as a function of horizontal position. Insets in **B** and **C** for Monkey B show zoomed-in representation of early reach trajectory, given the modest movement in the wrong direction.

Both monkeys generated a high proportion of errors on double target trials, leading us to wonder

about the degree to which they were exerting contextual control on these trials. First, we did find

that the RTs on double target red trials were significantly longer than the RTs on single target red

trials, suggesting the presence of intermixed trials slowed the animals down (Monkey G:

t(1873)=2.75, $p < 0.01$; Monkey B: t(4258)=8.58, $p < 10^{-10}$). Second, while the RTs of both correct

and incorrect trials pooled together did not differ significantly on double target green versus double

target red trials in either monkey (Monkey G: t(1043)= -1.41, $p = 0.16$; Monkey B: t(3015)= -1.63,

$p = 0.10$), incorrect reaches were quite small, and an analysis of the kinematics of reaching revealed

notable differences in performance on double target red vs double target green trials. Recall that

Monkey G tended to smear his hand on the touchscreen, which allowed us to visualize the

trajectory of the reaching response. The grouped trajectories for a subset of reaching responses on

double target trials are shown for Monkey G in **Fig. 5B** and **C** (top row). Correct leftward reaches

on double target green trials (green traces) follow the trajectory of correct leftward reaches on

double target red trials (red traces). In contrast, while incorrect right reaches on double target green

trials initially follow a similar trajectory as these leftward reaches (black trace), they are rapidly

reversed in direction, and end up at similar locations to correct rightward reaches on double target

red trials. Thus, even though Monkey G initiated correct and incorrect movements at similar

latencies, later parts of incorrect reaches were rapidly re-directed in the appropriate direction.

Similar tendencies are observed in Monkey B, although to a much more subtle degree since

Monkey B tended to lift his hand off the touchscreen (see insets zooming in on the early parts of

the reach trajectories). These behavioral patterns are consistent with the animals trying, and often

failing, to exert more contextual control on double target green trials. However, later parts of the

reach trajectory on double target green trials confirmed that both animals had consolidated the task

instruction to reach to the correct location.

We next asked the degree to which contextual control influenced the express visuomotor response

to the red stimulus. In humans, express visuomotor responses evolve at the same time but are

dampened in magnitude when subjects move away from rather than toward a peripheral stimulus

(Gu et al., 2016; Kozak et al., 2020), or exert some other degree of contextual control (Wood et

al., 2015; Atsma et al., 2018; Contemori et al., 2021b). An analysis of contextual control on the

express visuomotor response requires that we compare muscle recruitment on trials where the red

stimulus appears at the same location on either double target red or double target green trials.

Figure 6A shows the mean ± stderr representations of EMG activity from the same representative

sessions as shown in **Figure 2**, depicting right middle deltoid recruitment when the red stimulus

appeared to right. The monkeys reached to the right on double target red trials (red contours in

Fig. 6A), and the express visuomotor response resembled that seen on the single target red trials

(**Fig. 2B,C**). On double target green trials, the correct response was to withhold the rightward

reach, and reach to the left. When the animals did this correctly, the initial express visuomotor

response to the red stimulus was muted (green contours in **Fig. 6A**). In contrast, when the monkeys

incorrectly reached to the red stimulus on double target green trials, initial muscle recruitment

during the express response interval resembled that on double target red trials (compare black and

red contours in **Fig. 6A**), whereas later phases of recruitment resembled correctly-performed

double target green trials, as the animal corrected their reach. Thus, for these examples, the express

visuomotor response to the earlier appearing red stimulus depended both on the context of the trial

(double target red vs double target green), and on performance on double target green trials.

Figure 6. Contextual control on Express Visuomotor Response (EVR) on Double Target trials. **A.** Depiction of mean EMG activity +/- stderr from a representative session when the red stimulus emerged to the right. Gray rectangles denote the 25 ms interval following the express visuomotor response, over which magnitude was calculated. Solid colored vertical lines in **A** depict mean reaction time for corresponding trial type from representative sessions. **B,C.** The latency (**B**) or magnitude (**C**) of the express visuomotor response, by trial type and/or performance, across 29 occurrences where an express visuomotor response could be detected on both Double Target Red and Double Target Green trials. Magnitudes are expressed as a modulation index (see equation), for a variety of contrasts. Gray lines connect observations from the same muscle from a given experimental day. Black lines and squares show mean +/- standard error. Crosses denote distributions significantly different from zero (t-test, p < 0.05). Asterisks show significant differences of Bonferroni corrected paired t-test (p < 0.05).

Across our sample, we found that contextual control influenced the magnitude but not timing of

the express visuomotor response to the red stimulus. To quantify this, we first compared the timing

of the express visuomotor response, when such a response was detected on both double-target red

and double-target green trials. We found no influence of context on express visuomotor response

latency, with the latency being unchanged across double-target red trials and double-target green

trials (**Fig. 6B**. Monkey G: double-target red = 66 ± 9 ms; double-target green = 70 ± 14 ms; paired

t-test; $t(9) = -0.75$, $p = 0.46$; Monkey B: double-target red = 67 ± 13 ms; double-target green = 67

± 10 ms; $t(18) = 0.02$, $p = 0.98$).

Next, we determined the comparative magnitude of the express visuomotor response over the next

25 ms on double target red or double target green trials, expressing the change in magnitude as a

modulation index (see equation in **Fig. 6C**). Calculated this way, a value of 0 arises when there is

no modulation of the express visuomotor response magnitude for different trial types, whereas

values greater than (less than) 0 arise when the magnitude of the express visuomotor response is

larger (smaller) on double target red than double target green trials. Our first calculation

(comparison "A" in **Fig. 6C**) compared double target red trials to all double target green trials,

regardless of whether double target green trials were performed correctly or not. While there was

considerable variance, modulation indices across our sample averaged 0.09 ± 0.24, and were

significantly distributed above zero (*t*-test, $t(31) = 2.14$, $p < 0.05$). Thus, on average, express

visuomotor responses tended to be ~10% smaller on double target green than double target red

trials.

We then subdivided double target green trials based on whether monkeys correctly reached to the

green stimulus or incorrectly reached to the red stimulus. There were 29 comparisons with

sufficient numbers of both correct and incorrect double target green trials. The modulation index

for comparison B compared double target red to correctly performed double target green trials;

across our sample this modulation index averaged 0.17 ± 0.29 and was significantly distributed

above zero (*t*-test Bonferroni-corrected for multiple comparisons, $t(28) = 3.15$, $p < 0.005$). On

average, express visuomotor responses tended to be ~20% smaller on correctly performed double

target green than double target red trials. In contrast, comparison C compared double target red to

incorrectly performed double target green trials; across our sample this modulation index averaged

-0.06 ± 0.23 and was not significantly different from zero (Bonferroni-corrected *t*-test, $t(28) = -$

1.4, $p = 0.34$). Finally, since comparisons B and C could be derived within the same session, we

found that the modulation indices for comparison B tended to be significantly greater than for

comparison C (Bonferroni-corrected paired *t*-test, $t(28) = 4.22$, $p < 0.001$). In other words, we

observed a greater degree of contextual modulation on correct vs incorrect double target green

trials. Overall, and as in humans, we found that the magnitude but not latency of the express

visuomotor response changed with the context of the task (Gu et al., 2016; Contemori et al., 2021b,

2022a). Furthermore, as with anti-reaches (Gu et al., 2016), larger express visuomotor responses

preceded incorrect responses.

Discussion

We focused on the initial recruitment of upper limb muscles in monkeys reaching towards

emerging targets, using a paradigm which readily elicits express visuomotor responses in humans

(Kozak et al., 2020). Express visuomotor responses in monkeys exhibited the response properties

known in humans: its timing is more time-locked to target emergence than movement onset, and

its magnitude is both inversely related to RT and liable to dampening via contextual control. Our

results establish how rapidly the visual-to-motor transformation is completed in the rhesus

macaque brain, place constraints on the timing of premotor events, and set the stage for

neurophysiological experiments that can directly compare the characteristics of such premotor

events to those seen in the motor periphery.

Comparisons of express responses in humans and monkeys

Rhesus macaques offer an excellent animal model for understanding rapid visual-to-motor

transformations, due to the many homologies in foveate vision, implicated brain networks, and the

musculoskeletal anatomy of the eyes, neck, and upper limb. Humans and macaques generate a

variety of expedited visuomotor responses in the lab, including express saccades (Fischer and

Weber, 1993), microsaccades (Hafed and Ignashchenkova, 2013; Tian et al., 2016), express

visuomotor responses in neck muscle (Goonetilleke et al., 2015), reaches to rapidly moving stimuli

(Perfiliev et al., 2010), and ocular following responses (Miles et al., 1986; Gellman et al., 1990).

A common finding is that such responses evolve ~20-30 ms earlier in macaques than humans,

which is broadly consistent with the 3/5th rule (Schroeder et al., 2004) for quicker sensory

processing in macaques versus humans. Consistent with this, express visuomotor responses in

macaques started at ~65 ms, compared to 80-100 ms in humans.

A notable exception for earlier responses in macaques vs. humans is when subjects adjust reaching
trajectories in mid-flight to a displaced visual target. Humans can initiate such on-line corrections
within ~150-200 ms (Gaveau et al., 2014), even without conscious perception (Goodale et al.,
1986). The response latencies for on-line corrections in macaques are generally longer than in
humans, ranging between 200-250 ms (Georgopoulos et al., 1983; Archambault et al., 2009, 2011;
Pastor-Bernier et al., 2012; Dickey et al., 2013), although latencies comparable with humans have
been reported (Ames et al., 2019). The generally slower responses in macaques likely arises due
to an task-imposed emphasis of accuracy over speed, so that animals were not responding as
quickly as possible. Work by Perfiliev and colleagues (2010) compared 'interceptive' reaching
responses to rapidly moving targets of humans, monkeys, and cats, and clearly showed that all
three species can initiate reaches in less than 150 ms, with limb muscle recruitment in humans
starting at 90-100 ms. Such recruitment onsets are within the range of express visuomotor
responses in humans, but Perfiliev and colleagues did not record EMG in cats or monkeys.

Methodological considerations

A study by Lara and colleagues (Lara et al., 2018) recorded upper limb muscle activity in monkeys
in a 'quasi-automatic' task, where monkeys intercepted a target moving radially on a touch screen.
This task expedited RTs to ~200 ms on average, which is similar to the reaction times we observed.
As with the emerging target paradigm, the quasi-automatic task features a suddenly appearing
moving visual target. Lara and colleagues recorded from multiple heads of deltoid in addition to
other upper limb muscles, reporting the earliest changes in average muscle recruitment at ~100
ms, with single-trial onsets as low as 90 ms. Although these response latencies are short, they are
still ~25-35 ms longer than those reported here. The failure for the quasi-automatic task to elicit
express visuomotor responses may be due to a lack of precise temporal certainty about target
presentation or the absence of implied motion, both of which are critical in the emerging target
task (Kozak et al., 2020; Contemori et al., 2021a; Kozak and Corneil, 2021). The quasi-automatic
task in the Lara study was also intermingled with other trial types that engender longer RTs, which
may also lower the probability of express visuomotor responses. Finally, the Lara study employed
eight potential target locations, whereas we employed two. We do not think this difference explains
the absence of express visuomotor responses in the Lara study, as recent studies in humans have

reported express visuomotor responses in paradigms with a similar or higher number of potential
target locations (Gu et al., 2019; Kearsley et al., 2022; Selen et al., 2023) locations.

Caution is always required when comparing responses latencies across different laboratories.
Stimulus luminance and spatio-temporal frequency impact the latencies of on-line reach
corrections (Veerman et al., 2008; Kozak et al., 2019), the timing and magnitude of express muscle
responses (Wood et al., 2015; Kozak et al., 2019; Kozak and Corneil, 2021), and on the timing and
vigor of sensorimotor responses in the superior colliculus (SC) (Li and Basso, 2008; Marino et al.,
2012; Chen et al., 2018). Determining whether limb muscle recruitment meets an "express"
criterion should involve an assessment of trial-by-trial muscle recruitment over a range of RTs,
lest averaged recruitment profiles be biased by trials with the shortest RTs. The trial-by-trial
representations of EMG activity presented in the Lara study did not show consistently short
recruitment latencies across trials, strengthening our contention that express visuomotor responses
were not consistently elicited in their quasi-automatic task.

The emerging target paradigm provides a framework in which to study the most rapid visual-to-
motor transformations for reaching, and our results show that the underlying phenomenon is
distributed to many deltoid motor units in the macaque. The magnitudes of express responses
compared to later waves of activity are more modest in monkeys than in humans. However,
similarly-sized express visuomotor responses induce force transients or small movements toward
the stimulus (Wood et al., 2015; Gu et al., 2016; Atsma et al., 2018). We speculate that the modest
levels of recruitment may be because the deltoid muscles targeted here may not be prime movers
when monkeys interact with a touchscreen. Future work recording from more muscles of the upper
limb may find stronger express visuomotor responses, and experimental setups with a robotic
manipulandum will allow loading of selected muscle, as done routinely in humans. We did observe
small movements of the hand toward the earlier appearing red target on double target green trials,
which we surmise arose from both the express visuomotor responses and subsequent periods of
recruitment. The assessment of limb motion via the touchscreen is coarse, hence setups using
robotic manipulanda or motor tracking will permit more precise assessment of the kinematic
consequences of express visuomotor responses in monkeys. Future studies should also track eye
movements. While human studies have shown that the eyes are stable at the time of stimulus

presentation in the emerging target paradigm (Contemori et al., 2021a), what the eyes were doing
at stimulus emergence in our study is not known. It is possible that the modest magnitude of
express visuomotor responses could be due to saccades made around the time of target emergence,
as such contemporaneous eye movements can decrease the vigor of visual responses in movement-
related neurons in the superior colliculus (Chen and Hafed, 2017).

Double target green trials required the monkeys to suppress the tendency to look to the earlier-
appearing red stimulus. We expected that RTs for such trials would be longer than the RTs for
double target red trials, but somewhat surprisingly this was not the case if both correct and
incorrect trials were pooled together. Future variants of this task may benefit from longer delays
between the emergence of the red or green stimuli, and on-line identification of incorrect reaches.
In our data, the incorrect reaches themselves were often quite small in amplitude, hence the animals
could be rewarded on trials where they initially moved in the wrong direction so long as they
eventually reached the target. That being said, arm movements have substantial inertia, and in such
systems establishing the criteria used to distinguish "incorrect" from "correct" movements
necessitate a degree of subjectivity. Determining such criteria is all the more pernicious given
evidence that the express visuomotor response itself, whether on upper limb or neck muscles,
imparts sufficient forces to subtly move the arm or head (Corneil et al., 2008; Gu et al., 2016).
Regardless, reach trajectories (**Fig. 5C**) show that both animals rapidly corrected incorrect
movements in mid-flight, which is consistent with them consolidating task instruction. The impact
of task context on the express visuomotor response to the red stimulus was both modest and
variable (**Fig. 6**), but the average magnitude resembled that seen on monkey neck muscles
performing pro- or anti-saccades (Chapman and Corneil, 2011). Contextual modulation of the
express visuomotor responses is likely something that can be optimized in future experiments, for
example by employing various endogenous or exogenous cueing strategies that potentiate the
phenomenon of express visuomotor responses in the human arm (Contemori et al., 2021b, 2022a)
or monkey neck (Corneil et al., 2008).

Implications of our results
Express visuomotor responses are thought to arise from a subcortical tecto-reticulo-spinal circuit
that runs in parallel with corticospinal circuits (Boehnke and Munoz, 2008; Pruszynski et al., 2010;

Corneil and Munoz, 2014; Contemori et al., 2022b). This hypothesis has never been directly tested

in monkeys, as the lone reaching study inactivating the SC during a reaching task (Song et al.,

2011) did not record limb EMG and used a delayed response task that would not engender express

visuomotor responses. Considerable circumstantial evidence implicates the visual response within

the intermediate and deep layers of the superior colliculus (dSC) as being key in driving express

responses. The timing and vigor of the visual response, when summed with pre-existing

preparatory activity, drives express saccades (Edelman and Keller, 1996; Dorris et al., 1997;

Sparks et al., 2000; Dash et al., 2018). Visual responses in the dSC are also correlated with the

magnitude of express visuomotor responses in the neck, leading such responses by an appropriate

efferent lag (Rezvani and Corneil, 2008). Reach-related neurons in the dSC display short-latency

visual responses (Werner et al., 1997a, 1997b) and correlate with upper limb muscle activity

(Werner et al., 1997a; Stuphorn et al., 1999), although the properties of visual responses in reach-

related neurons have not been systematically investigated. Our speculation of a subcortical route

initiating visually-guided reaches to a punctate target is consistent with recent stimulation results

in humans (Divakar et al., 2022), and parallels a growing appreciation for the role of subcortical

pathways in manual following responses to whole-field motion (Saijo et al., 2005) or flashes

(Nakajima et al., 2021).

While we acknowledge the shortcomings in comparing results across laboratories and

experimental tasks, the express visuomotor responses reported here on arm muscles occur at an

interval when neural activity in the primary motor cortex is thought to be in a null space that is not

tied to motor output (Lara et al., 2018). Indeed, one of the main conclusions for the Lara study

mentioned above is that a mandatory transition of neural activity within motor cortex from a null-

space to a movement-potent subspace precedes motor output by ~20-30 ms. Our results present

two alternatives which can frame future experiments.

First, if our hypothesis is correct, a subcortical route through the dSC can drive express visuomotor

responses when M1 is in a null space. An intriguing corollary to this alternative, given ascending

connections from the SC to thalamic areas that themselves project to motor cortices (Benevento

and Fallon, 1975; Matelli and Luppino, 1996), is that it is signaling from the SC to the motor cortex

that drives the transition of cortical activity from null to movement potent subspaces (Ames et al.,

2019), which presumably could allow cortical activity to drive subsequent muscle recruitment that

is informed by the consequences of the preceding express visuomotor response. Ascending signals

from the SC through the thalamus have been long established as being critical for relaying

efference copy signals for eye movements (Sommer and Wurtz, 2002), and implicated in

prioritizing spatial locations for target selection for reaching (Song et al., 2011) and for higher

order processing (Lovejoy and Krauzlis, 2010; Bogadhi et al., 2021; Peysakhovich et al., 2023).

Alternatively, could the transition of neural activity from a null to movement potent subspace in

M1 simply occur earlier in the emerging target paradigm? This line of reasoning surmises that

signaling along a cortico-spinal or cortico-reticulo-spinal route drives express visuomotor

responses. If so, visually-related activity would have to evolve in motor cortex ~10-20 ms earlier

than the express visuomotor responses we report, so at around 50 ms. There are reports of visual

responses in the macaque primary motor cortex at such latencies (Murphy et al., 1982; Lamarre et

al., 1983; Crammond and Kalaska, 1994; Riehle et al., 1997; Cisek and Kalaska, 2005; Ames et

al., 2019), but comparisons to these reports are again hindered by the use of tasks or visual stimuli

that in retrospect may not have been optimized to elicit express responses, and various filtering

settings or analysis schemes that complicate timing comparisons. Examination of the rasters of

single-unit data from such reports, when available, also do not always reveal the trial-by-trial

consistency that we observed in the motor periphery. One intriguing exception comes from the

work of Reimer and colleagues (Reimer and Hatsopoulos, 2010), who studied motor cortical

activity while NHPs adjusted the trajectory of ongoing limb movements to jumped visual targets.

When one accounts for an acknowledged timing error, motor cortex activity changed consistently

~50 ms after the target jump. While this theoretically shows that signaling related to visual signal

onset could arrive in time within motor cortices, we note that express visuomotor responses in

humans evolve ~10-15 ms earlier (and hence ~80 ms or less) when evoked when the arm is already

in motion (Kozak et al., 2019), presumably due to disengagement of a control policy for postural

stability (Cluff and Scott, 2016).

 Figure Legends

 Figure 1. A-C. Emerging Target Task. At the start of each trial, either a red (**A, B**) or green (**C**)

 stimulus appears at the start position (red or green squares), below an occluder (gray box). The

 monkey touches and holds the screen at this start position. A target of the same color then appears

 above the occluder and moves down the chute before disappearing briefly behind the occluder for

 300 ms. In Single Target Red trials (**A**), a single red target emerges in motion below the occluder

 at one of two outlets. In the Double Target Red (**B**) and Double Target Green (**C**) trials, appearance

 of the red target in one outlet was followed 34 ms later by appearance of a green target at the other

 outlet. The monkey was rewarded for reaching to the target that matched the color of the stimuli

 presented at the start of the trial. **D.** Movement-aligned activity for the anterior, middle, and

 posterior heads of deltoids for the two monkeys for single-target red trials, showing greater

 recruitment for rightward than leftward reaches. Data shows the average of the mean recruitment

 from each session, normalized to the peak movement-aligned activity from each session. Contours

 subtend the mean ± stderr.

 Figure 2. A. Frequency histograms of reaction times (20 ms bins) for Monkey G (top row, n =

 1303 trials) and Monkey B (bottom row, n = 2648 trials) across all experimental sessions.

 Solid/dashed lines depict means/medians. **B.** Representative examples of express visuomotor

 responses in Monkey G (top row) and Monkey B (bottom row), depicting trial-by-trial EMG

 recruitment aligned to target emergence to the left or right. In these plots, trials are sorted by

 movement onset (black squares), and the color conveys the magnitude of recruitment. **C.** For each

 monkey, the top row depicts mean EMG activity subtended by the standard error for left (light

 blue) and right (dark blue) reaches, and bottom row shows the time-series ROC (AUC = area under

 curve). Solid vertical lines depict mean reaction time for the representative examples. The red

 rectangles in **B** and gray rectangles in **C** show the 25 ms interval following the detection of the

 express visuomotor response.

 Figure 3. Properties of express visuomotor responses. **A.** Latencies of express visuomotor

 responses for both monkeys, showing latencies when all trials are considered regardless of reaction

 time, and after including only the longer-than-average reaction times. Gray lines connect

observations from the same muscle from a given experimental day (grey + marks denote

representative data shown in Fig. 2). Black lines and squares show mean ± standard error.

Asterisks show significant differences of paired t-test ($p < 0.05$). **B**. Depiction of the calculation

of the proportional magnitude of the express visuomotor response, using the representative data

from Monkey B. Using data normalized to the peak of movement-aligned activity, we integrate

the difference in the mean EMG curve for right versus left responses over the 25 ms after the onset

of the express visuomotor responses, and divide this by the integral of the 25 ms interval

surrounding peak EMG activity aligned to right reach onset. Same general format as **A**; asterisk

showing significant results of a student's t-test ($p < 0.05$).

Figure 4. Larger express visuomotor responses tended to precede shorter reach reaction times

(RTs). **A, B**. Representative data plotting magnitude of express visuomotor response as a function

of reach reaction time, from the representative sessions shown in Figure 2. Each data point

represents a single trial, and the black diagonal line shows the linear regression. The correlation

coefficient (*r*) is shown in the top-right of each subplot. **C**. Histogram of correlation coefficients

across all sessions in which an express visuomotor response was detected. Vertical dash line

indicates 0. Solid or gray lines or bins denote data for Monkey G or B, respectively. Vertical arrows

show mean r-values.

Figure 5. Contextual control on Double Target trials. **A.** Histograms across all sessions of reaction

times (20 ms bins) relative to the emergence of the red stimulus for Monkey G (top row) and

Monkey B (bottom row), for Double Target Red trials (left column, Monkey G: 632 trials, error

rate = 9.4%; Monkey B: 1788 trials, error rate = 9.8%) or Double Target Green trials (right column,

Monkey G: 413 trials, error rate = 50.1%; Monkey B: 1378 trials, error rate = 29.6%).

Upward/downward histograms represent correct/incorrect responses. Solid/dashed lines depict

histogram means/medians. **B.** Trajectory of horizontal touchscreen position averaged across all

trials for Double Target Red trials to the right or left (red contours), and Double Target Green

trials, showing correct trials when the monkey should have reached left (green contours), and

incorrect trials when the monkey should have reached right (black contours). Note how incorrect

trials initially go left before being corrected mid-flight to go to the right. Contours show mean ±

stderr. Solid colored vertical lines in B depict mean reaction time for corresponding trial type. **C.**

Same data, depicting vertical position as a function of horizontal position. Insets in **B** and **C** for
Monkey B show zoomed-in representation of early reach trajectory, given the modest movement
in the wrong direction.
740
Figure 6. Contextual control on Express Visuomotor Response (EVR) on Double Target trials. **A.**
Depiction of mean EMG activity +/- stderr from representative sessions when the red stimulus
emerged to the right. Gray rectangles denote the 25 ms interval following the express visuomotor
response, over which magnitude was calculated. Solid colored vertical lines in **A** depict mean
reaction time for corresponding trial type from representative sessions. **B,C**. The latency (**B**) or
magnitude (**C**) of the express visuomotor response, by trial type and/or performance, across 29
occurrences where an express visuomotor response could be detected on both Double Target Red
and Double Target Green trials. Magnitudes are expressed as a modulation index (see equation),
for a variety of contrasts. Gray lines connect observations from the same muscle from a given
experimental day. Black lines and squares show mean +/- standard error. Crosses denote
distributions significantly different from zero (t-test, $p < 0.05$). Asterisks show significant
differences of Bonferroni corrected paired t-test ($p < 0.05$).
753
754

References

References

Ames KC, Ryu SI, Shenoy KV (2019) Simultaneous motor preparation and execution in a last-
moment reach correction task. Nat Commun 10:2718.

Archambault PS, Caminiti R, Battaglia-Mayer A (2009) Cortical Mechanisms for Online Control
of Hand Movement Trajectory: The Role of the Posterior Parietal Cortex. Cereb Cortex
19:2848–2864.

Archambault PS, Ferrari-Toniolo S, Battaglia-Mayer A (2011) Online control of hand trajectory
and evolution of motor intention in the parietofrontal system. J Neurosci 31:742–752.

Atsma J, Maij F, Gu C, Medendorp WP, Corneil BD (2018) Active Braking of Whole-Arm
Reaching Movements Provides Single-Trial Neuromuscular Measures of Movement
Cancellation. J Neurosci 38:4367–4382.

Benevento LA, Fallon JH (1975) The ascending projections of the superior colliculus in the
rhesus monkey (Macaca mulatta). J Comp Neurol 160:339–361.

Bibi R, Edelman JA (2009) The influence of motor training on human express saccade
production. J Neurophysiol 102:3101–3110.

Boehnke SE, Munoz DP (2008) On the importance of the transient visual response in the
superior colliculus. Curr Opin Neurobiol 18:544–551.

Bogadhi AR, Katz LN, Bollimunta A, Leopold DA, Krauzlis RJ (2021) Midbrain activity shapes
high-level visual properties in the primate temporal cortex. Neuron 109:690–699.e5.

Chapman BB, Corneil BD (2011) Neuromuscular recruitment related to stimulus presentation
and task instruction during the anti-saccade task. Eur J Neurosci 33:349–360.

Chen C-Y, Hafed ZM (2017) A neural locus for spatial-frequency specific saccadic suppression
in visual-motor neurons of the primate superior colliculus. J Neurophysiol 117:1657–1673.

Chen C-Y, Sonnenberg L, Weller S, Witschel T, Hafed ZM (2018) Spatial frequency sensitivity
in macaque midbrain. Nat Commun 9:1–13.

Cisek P, Kalaska JF (2005) Neural correlates of reaching decisions in dorsal premotor cortex:
specification of multiple direction choices and final selection of action. Neuron 45:801–814.

Cluff T, Scott SH (2016) Online Corrections are Faster Because Movement Initiation Must
Disengage Postural Control. Motor Control 20:162–170.

Contemori S, Loeb GE, Corneil BD, Wallis G, Carroll TJ (2021a) The influence of temporal
predictability on express visuomotor responses. J Neurophysiol 125:731–747.

Contemori S, Loeb GE, Corneil BD, Wallis G, Carroll TJ (2021b) Trial-by-trial modulation of
express visuomotor responses induced by symbolic or barely detectable cues. J

Neurophysiol 126:1507–1523.

Contemori S, Loeb GE, Corneil BD, Wallis G, Carroll TJ (2022a) Symbolic cues enhance
express visuomotor responses in human arm muscles at the motor planning rather than the
visuospatial processing stage. J Neurophysiol 128:494–510.

Contemori S, Loeb GE, Corneil BD, Wallis G, Carroll TJ (2022b) Express visuomotor responses
reflect knowledge of both target location and conscious intent during reaches of different
amplitudes. bioRxiv:2022.11.06.515363 Available at:
https://www.biorxiv.org/content/biorxiv/early/2022/11/06/2022.11.06.515363 [Accessed
March 1, 2023].

Corneil BD, Goonetilleke SC, Peel TR, Green KA, Welch ID (2012) Ultrasound-guided insertion
of intramuscular electrodes into suboccipital muscles in the non-human primate. J
Electromyogr Kinesiol 22:553–559.

Corneil BD, Munoz DP (2014) Overt responses during covert orienting. Neuron 82:1230–1243.

Corneil BD, Munoz DP, Chapman BB, Admans T, Cushing SL (2008) Neuromuscular
consequences of reflexive covert orienting. Nat Neurosci 11:13–15.

Corneil BD, Olivier E, Munoz DP (2004) Visual responses on neck muscles reveal selective
gating that prevents express saccades. Neuron 42:831–841.

Crammond DJ, Kalaska JF (1994) Modulation of preparatory neuronal activity in dorsal
premotor cortex due to stimulus-response compatibility. J Neurophysiol 71:1281–1284.

Dash S, Peel TR, Lomber SG, Corneil BD (2018) Frontal eye field inactivation reduces saccade
preparation in the superior colliculus but does not alter how preparatory activity relates to
saccades of a given latency. eNeuro 5 Available at:
https://www.eneuro.org/lookup/doi/10.1523/ENEURO.0024-18.2018.

Dash S, Peel TR, Lomber SG, Corneil BD (2020) Impairment but not abolishment of express
saccades after unilateral or bilateral inactivation of the frontal eye fields. J Neurophysiol
123:1907–1919.

Day BL, Brown P (2001) Evidence for subcortical involvement in the visual control of human
reaching. Brain 124:1832–1840.

Dickey AS, Amit Y, Hatsopoulos NG (2013) Heterogeneous neural coding of corrective
movements in motor cortex. Front Neural Circuits 7:51.

Divakar R, Loeb GE, Corneil BD, Wallis G, Carroll TJ (2022) A subcortical origin for rapid,
target-oriented corticospinal excitability changes during visually guided reaching.
bioRxiv:2022.12.13.520229 Available at:
https://www.biorxiv.org/content/biorxiv/early/2022/12/13/2022.12.13.520229 [Accessed
March 1, 2023].

Dorris MC, Paré M, Munoz DP (1997) Neuronal activity in monkey superior colliculus related to
the initiation of saccadic eye movements. J Neurosci 17:8566–8579.

Edelman JA, Keller EL (1996) Activity of visuomotor burst neurons in the superior colliculus
accompanying express saccades. J Neurophysiol 76:908–926.

Edelman JA, Kristjánsson A, Nakayama K (2007) The influence of object-relative visuomotor
set on express saccades. J Vis 7:12.

Elsley JK, Nagy B, Cushing SL, Corneil BD (2007) Widespread presaccadic recruitment of neck
muscles by stimulation of the primate frontal eye fields. J Neurophysiol 98:1333–1354.

Fautrelle L, Prablanc C, Berret B, Ballay Y, Bonnetblanc F (2010) Pointing to double-step visual
stimuli from a standing position: very short latency (express) corrections are observed in
upper and lower limbs and may not require cortical involvement. Neuroscience 169:697–
705.

Fischer B, Boch R (1983) Saccadic eye movements after extremely short reaction times in the
monkey. Brain Res 260:21–26.

Fischer B, Ramsperger E (1984) Human express saccades: extremely short reaction times of goal
directed eye movements. Exp Brain Res 57:191–195.

Fischer B, Weber H (1993) Express saccades and visual attention. Behav Brain Sci 16:553–567.

Gaveau V, Pisella L, Priot A-E, Fukui T, Rossetti Y, Pélisson D, Prablanc C (2014) Automatic
online control of motor adjustments in reaching and grasping. Neuropsychologia 55:25–40.

Gellman RS, Carl JR, Miles FA (1990) Short latency ocular-following responses in man. Vis
Neurosci 5:107–122.

Georgopoulos AP, Kalaska JF, Caminiti R, Massey JT (1983) Interruption of motor cortical
discharge subserving aimed arm movements. Exp Brain Res 49:327–340.

Glover IS, Baker SN (2019) Multimodal stimuli modulate rapid visual responses during
reaching. J Neurophysiol 122:1894–1908.

Goodale MA, Pelisson D, Prablanc C (1986) Large adjustments in visually guided reaching do
not depend on vision of the hand or perception of target displacement. Nature 320:748–750.

Goonetilleke SC, Katz L, Wood DK, Gu C, Huk AC, Corneil BD (2015) Cross-species
comparison of anticipatory and stimulus-driven neck muscle activity well before saccadic
gaze shifts in humans and nonhuman primates. J Neurophysiol 114:902–913.

Gu C, Andrew Pruszynski J, Gribble PL, Corneil BD (2018) Done in 100 ms: path-dependent
visuomotor transformation in the human upper limb. Journal of Neurophysiology
119:1319–1328.

Gu C, Pruszynski JA, Gribble PL, Corneil BD (2019) A rapid visuomotor response on the human
upper limb is selectively influenced by implicit motor learning. J Neurophysiol 121:85–95.

Gu C, Wood DK, Gribble PL, Corneil BD (2016) A Trial-by-Trial Window into Sensorimotor
Transformations in the Human Motor Periphery. J Neurosci 36:8273–8282.

Hafed ZM, Ignashchenkova A (2013) On the dissociation between microsaccade rate and
direction after peripheral cues: microsaccadic inhibition revisited. J Neurosci 33:16220–
16235.

Hwang J, Mitz AR, Murray EA (2019) NIMH MonkeyLogic: Behavioral control and data
acquisition in MATLAB. J Neurosci Methods 323:13–21.

Kearsley SL, Cecala AL, Kozak RA, Corneil BD (2022) Express arm responses appear
bilaterally on upper-limb muscles in an arm choice reaching task. J Neurophysiol 127:969–
983.

Kozak RA, Cecala AL, Corneil BD (2020) An Emerging Target Paradigm to Evoke Fast
Visuomotor Responses on Human Upper Limb Muscles. J Vis Exp Available at:
http://dx.doi.org/10.3791/61428.

Kozak RA, Corneil BD (2021) High-contrast, moving targets in an emerging target paradigm
promote fast visuomotor responses during visually guided reaching. J Neurophysiol
126:68–81.

Kozak RA, Kreyenmeier P, Gu C, Johnston K, Corneil BD (2019) Stimulus-Locked Responses
on Human Upper Limb Muscles and Corrective Reaches Are Preferentially Evoked by Low
Spatial Frequencies. eneuro 6:ENEURO.0301–0319.2019 Available at:
http://dx.doi.org/10.1523/eneuro.0301-19.2019.

Kurtzer I, Herter TM, Scott SH (2006) Nonuniform distribution of reach-related and torque-
related activity in upper arm muscles and neurons of primary motor cortex. J Neurophysiol
96:3220–3230.

Lamarre Y, Busby L, Spidalieri G (1983) Fast ballistic arm movements triggered by visual,
auditory, and somesthetic stimuli in the monkey. I. Activity of precentral cortical neurons. J
Neurophysiol 50:1343–1358.

Lara AH, Elsayed GF, Zimnik AJ, Cunningham JP, Churchland MM (2018) Conservation of
preparatory neural events in monkey motor cortex regardless of how movement is initiated.
Elife 7 Available at: http://dx.doi.org/10.7554/eLife.31826.

Li X, Basso MA (2008) Preparing to Move Increases the Sensitivity of Superior Colliculus
Neurons. Journal of Neuroscience 28:4561–4577.

Loeb GE, Gans C (1986) Electromyography for Experimentalists. University of Chicago Press.

Lovejoy LP, Krauzlis RJ (2010) Inactivation of primate superior colliculus impairs covert

891   selection of signals for perceptual judgments. Nat Neurosci 13:261–266.

892 Marino RA, Levy R, Boehnke S, White BJ, Itti L, Munoz DP (2012) Linking visual response
893   properties in the superior colliculus to saccade behavior. Eur J Neurosci 35:1738–1752.

894 Marino RA, Levy R, Munoz DP (2015) Linking express saccade occurance to stimulus
895   properties and sensorimotor integration in the superior colliculus. J Neurophysiol 114:879–
896   892.

897 Matelli M, Luppino G (1996) Thalamic input to mesial and superior area 6 in the macaque
898   monkey. The Journal of Comparative Neurology 372:59–87.

899 Miles FA, Kawano K, Optican LM (1986) Short-latency ocular following responses of monkey.
900   I. Dependence on temporospatial properties of visual input. J Neurophysiol 56:1321–1354.

901 Munoz DP, Dorris MC, Paré M, Everling S (2000) On your mark, get set: brainstem circuitry
902   underlying saccadic initiation. Can J Physiol Pharmacol 78:934–944.

903 Murphy JT, Kwan HC, MacKay WA, Wong YC (1982) Precentral unit activity correlated with
904   angular components of a compound arm movement. Brain Res 246:141–145.

905 Nakajima T, Ohtsuka H, Irie S, Suzuki S, Ariyasu R, Komiyama T, Ohki Y (2021) Visual
906   information increases the indirect corticospinal excitation via cervical interneurons in
907   humans. J Neurophysiol 125:828–842.

908 Novembre G, Iannetti GD (2021) Towards a unified neural mechanism for reactive adaptive
909   behaviour. Prog Neurobiol 204:102115.

910 Olfert ED, Cross BM, McWilliam AA, Others (1993) Guide to the care and use of experimental
911   animals. Canadian Council on Animal Care Ottawa.

912 Paré M, Munoz DP (1996) Saccadic reaction time in the monkey: advanced preparation of
913   oculomotor programs is primarily responsible for express saccade occurrence. J
914   Neurophysiol 76:3666–3681.

915 Pastor-Bernier A, Tremblay E, Cisek P (2012) Dorsal premotor cortex is involved in switching
916   motor plans. Front Neuroeng 5:5.

917 Perfiliev S, Isa T, Johnels B, Steg G, Wessberg J (2010) Reflexive limb selection and control of
918   reach direction to moving targets in cats, monkeys, and humans. J Neurophysiol 104:2423–
919   2432.

920 Peysakhovich B, Tetrick SM, Silva AA, Li S, Zhu O, Ibos G, Johnston WJ, Freedman DJ (2023)
921   Primate superior colliculus is engaged in abstract higher-order cognition. bioRxiv Available
922   at: http://dx.doi.org/10.1101/2023.01.17.524416.

923 Pruszynski JA, Johansson RS, Flanagan JR (2016) A Rapid Tactile-Motor Reflex Automatically
924   Guides Reaching toward Handheld Objects. Curr Biol 26:788–792.

Pruszynski JA, King GL, Boisse L, Scott SH, Flanagan JR, Munoz DP (2010) Stimulus-locked
responses on human arm muscles reveal a rapid neural pathway linking visual input to arm
motor output. Eur J Neurosci 32:1049–1057.

Reimer J, Hatsopoulos NG (2010) Periodicity and evoked responses in motor cortex. J Neurosci
30:11506–11515.

Reschechtko S, Pruszynski JA (2020) Voluntary modification of rapid tactile-motor responses
during reaching differs from its visuomotor counterpart. J Neurophysiol 124:284–294.

Rezvani S, Corneil BD (2008) Recruitment of a head-turning synergy by low-frequency activity
in the primate superior colliculus. J Neurophysiol 100:397–411.

Riehle A, Kornblum S, Requin J (1997) Neuronal correlates of sensorimotor association in
stimulus–response compatibility. J Exp Psychol Hum Percept Perform 23:1708–1726.

Rohrer WH, Sparks DL (1993) Express saccades: the effects of spatial and temporal uncertainty.
Vision Res 33:2447–2460.

Saijo N, Murakami I, Nishida S 'ya, Gomi H (2005) Large-field visual motion directly induces
an involuntary rapid manual following response. J Neurosci 25:4941–4951.

Schiller PH, Sandell JH, Maunsell JH (1987) The effect of frontal eye field and superior
colliculus lesions on saccadic latencies in the rhesus monkey. J Neurophysiol 57:1033–
1049.

Schroeder CE, Molhom S, Lakatos P, Ritter W, Foxe JJ (2004) Human–simian correspondence
in the early cortical processing of multisensory cues. Cogn Process 5:140–151.

Selen LPJ, Corneil BD, Medendorp WP (2023) Single-Trial Dynamics of Competing Reach
Plans in the Human Motor Periphery. J Neurosci 43:2782–2793.

Sommer MA, Wurtz RH (2002) A Pathway in Primate Brain for Internal Monitoring of
Movements. Science 296:1480–1482.

Song J-H, Rafal RD, McPeek RM (2011) Deficits in reach target selection during inactivation of
the midbrain superior colliculus. Proc Natl Acad Sci U S A 108:E1433–E1440.

Sparks D, Rohrer WH, Zhang Y (2000) The role of the superior colliculus in saccade initiation: a
study of express saccades and the gap effect. Vision Res 40:2763–2777.

Stuphorn V, Hoffmann KP, Miller LE (1999) Correlation of primate superior colliculus and
reticular formation discharge with proximal limb muscle activity. J Neurophysiol 81:1978–
1982.

Suzuki S, Nakajima T, Irie S, Ariyasu R, Ohtsuka H, Komiyama T, Ohki Y (2022) Subcortical
Contribution of Corticospinal Transmission during Visually Guided Switching Movements
of the Arm. Cereb Cortex 32:380–396.

Tian X, Yoshida M, Hafed ZM (2016) A Microsaccadic Account of Attentional Capture and
Inhibition of Return in Posner Cueing. Front Syst Neurosci 10:23.

Veerman MM, Brenner E, Smeets JBJ (2008) The latency for correcting a movement depends on
the visual attribute that defines the target. Exp Brain Res 187:219–228.

Werner W, Dannenberg S, Hoffmann KP (1997a) Arm-movement-related neurons in the primate
superior colliculus and underlying reticular formation: comparison of neuronal activity with
EMGs of muscles of the shoulder, arm and trunk during reaching. Exp Brain Res 115:191–
205.

Werner W, Hoffmann KP, Dannenberg S (1997b) Anatomical distribution of arm-movement-
related neurons in the primate superior colliculus and underlying reticular formation in
comparison with visual and saccadic cells. Exp Brain Res 115:206–216.

Wood DK, Gu C, Corneil BD, Gribble PL, Goodale MA (2015) Transient visual responses reset
the phase of low-frequency oscillations in the skeletomotor periphery. European Journal of
Neuroscience 42:1919–1932.