1

Comparison of enrichment methods for efficient nitrogen fixation on a biocathode

Axel Rous[1], Gaëlle Santa-Catalina[1], Elie Desmond-Le Quéméner[1], Eric Trably[1], Nicolas Bernet[1]

[1] INRAE, Université de Montpellier, LBE, 102 avenue des Étangs, 11100 Narbonne, France

*Corresponding author

Correspondence:nicolas.bernet@inrae.fr

ABSTRACT

The production of nitrogen fertilizers in modern agriculture is mostly based on the Haber-Bosch process, representing nearly 2% of the total energy consumed in the world. Low-energy bioelectrochemical fixation of N_2 to microbial biomass was previously observed on a biocathode but the microbial interactions in N_2-fixing electroactive biofilms are still poorly understood. The present study aims to develop a method of enrichment of autotrophic and diazotrophic bacteria from soil samples for further characterization. The enrichment method was based on a first classical step of selection of N_2-fixing bacteria from soil samples. Then, a polarized cathode was used for the enrichment of autotrophic bacteria using H_2 (hydrogenotrophic) or the cathode as energy sources. This enrichment was compared with an enrichment of diazotrophic hydrogenotrophic bacteria without the use of the microbial electrochemical system. Both methods showed comparable results for N_2 fixation rates at day 340 of the enrichment, with an estimated average of approximately 0.2 mgN_{fixed}/L.d. Current densities up to -15 A/m² were observed in the polarized cathode enrichments and a significant increase of the microbial biomass on the cathode was shown between 132 and 214 days of enrichment. These results confirm the enrichment of scrautotrophic, electrotrophic and diazotrophic bacteria in the polarized cathode

enrichments. Finally, the analysis of the enriched communities suggest that *Desulforamulus*
ruminis mediated microbial interactions between autotrophic anaerobic and heterotrophic
aerobic bacteria in polarized cathode enrichment. These interactions could play a key role in
the development of biomass in these systems and on N_2 fixation. Based on these findings, a
conceptual model on the functioning of mixed cultures N_2-fixing electroactive biofilms was
proposed.

Keywords: Nitrogen fixation, Microbial electrochemical system, Biomass electrostimulation,
Enrichment method

41
42

                                                Introduction

Nitrogen is one of the essential elements for the growth of all living organisms, especially for cellular
protein synthesis. In modern agriculture, ammonia is often used as a nitrogen source for plants (Bagali,
2012; Burris & Roberts, 1993; Masclaux-Daubresse et al., 2010). This compound is produced at industrial
scale by the Haber-Bosch process which allows the reduction of N_2 to NH_3 at the expanse of large
quantities of H_2 and energy (Kandemir et al., 2013; Martín et al., 2019). This process is associated with
significant CO_2 emissions, due to the source of H_2 obtained either by methane steam reforming or coal
gasification (Martín et al., 2019). Alternatives for green H_2 production, such as water electrolysis, are
therefore nowadays considered to feed the Haber-Bosch process that is also contributing to a high
energy demand of the process (Cherkasov et al., 2015). A more direct alternative is to reduce N_2 directly
on a cathode by chemical catalysts. However, these catalysts are not renewable and are currently not
sufficiently selective regarding the hydrogen evolution reaction at ambient conditions (Deng et al., 2018;
A. Liu et al., 2020).
Recently, some authors proposed to use N_2-fixing bacteria in association with electrochemical
systems for N_2 reduction at low energy cost (Liu et al., 2017; Rago et al., 2019). This idea was inspired by
previous observations of microbial CO_2 fixation on microbial cathodes in a process known as microbial
electrosynthesis (Liu et al., 2020; Logan et al., 2019). The microbial fixation of both N_2 and CO_2 with a
polarized cathode was demonstrated in two recent studies (C. Liu et al., 2017; Rago et al., 2019). First,
(Liu et al. (2017) demonstrated the growth of *Xanthobacter autotrophicus* in a hybrid organic-inorganic
electrochemical system in the absence of nitrogen sources other than N_2. Then, Rago et al. (2019)
demonstrated N_2 fixation through microbial electrosynthesis (MES) with a mixed microbial community.
This new type of process had the potential to produce biomass from N_2 and CO_2 with only electricity as
energy input that could be provided by small production units such as solar panels (Rago et al., 2019).
This biomass could then be used as a fertilizer with a low impact on the environment (Chakraborty &
Akhtar, 2021; Rago et al., 2019; Hafeez et al., 2006). Although the proof of concept was made for this
process, the microbial interactions supporting N_2 fixation in this microbial electrochemical systems are
still poorly understood. Different N_2 fixation scenarios are indeed possible, such as: (i) fixation by a single
population capable of fixing N_2 and CO_2 using the electrode as sole energy source, (ii) fixation by
heterotrophic diazotrophic bacteria that can utilize the organic carbon produced by electro-autotrophic
bacteria, (iii) fixation through an interaction between methanogenic archaea and methanotrophs that
could use CH_4 as an energy source for N_2 fixation and (iv) fixation through an interaction mediated by
direct interspecies electron transfer (DIET) between electro-autotrophic bacteria and diazotrophic
bacteria (Rago et al., 2019). A better understanding of these interactions is essential to optimize N_2
fixation in microbial electrochemical system.
In nature, biological N_2 fixation is a key mechanism of the nitrogen cycle where atmospheric nitrogen
is uptaken by living organisms (Bagali, 2012). It is carried out by so-called diazotrophic bacteria
responsible for the transformation of N_2 into NH_3 (Kim & Rees, 1994). Some of these bacteria can be
found on the roots of plants where they are living in symbiosis (Burris & Roberts, 1993; Franche et al.,
2009). These microorganisms are able to fix N_2 from the air, making it assimilable in the form of NH_4^+ or
amino acids (L-glutamine, L-glutamate) which are further used in plants for protein or DNA synthesis
(Burris & Roberts, 1993). In exchange, the bacteria use the organic matter of root exudates produced by
the plants as carbon and energy sources. Among the diazotrophic bacteria, the genera *Frankia* and
Rhizobium spp. are often associated with leguminous plant roots (Burris & Roberts, 1993; Peoples &
Craswell, 1992). In contrast, other free-living N_2-fixing bacteria such as *Azospirillum*, are able to fix N_2
without interacting with plants and can use organic or inorganic materials to produce their own energy
(Tilak et al., 1986).
The fixed N is then used for the synthesis of proteins and in particular for the synthesis of L-glutamate
and L-glutamine which are inhibitors of nitrogenase synthesis (Moreno-Vivián et al., 1989; Y. Zhang et al.,
2022). Interestingly, nitrogenases are not fully specific and are able to catalyze other reactions such as
the production of H_2 and the reduction of ethylene to acetylene which require less energy (Burgess &
Lowe, 1996). The latter reaction is used to indirectly measure N_2 fixation rate by the acetylene reduction
assay (ARA) (Bergersen, 1970; Soper et al., 2021). This measurement has particularly contributed to the

better understanding of the microbial mechanisms supporting N_2 fixation (Bergersen, 1970; Saiz et al.,
2019; Soper et al., 2021).
In order to better understand the microbial mechanisms supporting N_2 fixation in polarized cathode
enrichment, this work aims at developing an enrichment method of cathodic biofilms for direct fixation
of CO_2 and N_2. For this, soil samples were used as sources of N_2-fixing bacteria, and successive
enrichments in autotrophic bacteria in polarized cathode enrichment (PCE) were performed to select a
electroactive biofilm capable of fixing N and C with a cathode as sole energy source. The enriched biofilm
was compared with a classical enrichment of N_2-fixing hydrogen-oxidizing bacteria (HOB) in flasks (named
H_2 enrichment, H_2E) (Hu et al., 2020; Liu et al., 2017).

105   **Methods**

Inoculum
Soil samples from a forest, agricultural crop field and a commercial compost were used as microbial
inoculum sources. Samples were collected from forest and agricultural soils in the Haute Vallée de l'Aude,
France. These sources were selected based on their assumed abundance of N_2-fixing bacteria and their
theoretical C/N ratio (Khan et al., 2016). One to two mg of each samples were used as inoculum in 50mL
of medium for preliminary enrichment culture.

Culture media
The culture media were both formulated on the basis of H3 medium (81 DSMZ) used for enrichment
of soil autotrophic bacteria. The medium consisted of 2.3g KH_2PO_4 and 2.9g Na_2HPO_4 $2H_2O$ per liter as
buffer, 0.5g $MgSO_4$ $7H_2O$, 0.01g $CaCl_2$ $2H_2O$, 0.005g $MnCl_2$ $4H_2O$, 0.005g $NaVO_3$ H_2O, and 5mL of SL-6
trace element solution per liter of medium, with 5mL of vitamin solution. The vitamin solution consisted
of 0.1g $ZnSO_4$ $7H_2O$, 0.03g $MnCl_2$ $4H_2O$, 0.3g H_3BO_3 0.2 $CoCl_2$ $6H_2O$, 0.01g $CuCl_2$ $2H_2O$, 0.02g $NiCl_2$ $6H_2O$,
and 0.03g Na_2MoO_4 $2H_2O$ per liter of solution. Iron citrate was added to the enrichment bottles at a
concentration of 0.05 g/L but not to the microbial bioelectrochemical systems in which the cathode was
used as sole electron source. An organic carbon solution (organic C) was composed of 2g/L D-glucose,
1g/L yeast extract, 1g/L Na-acetate, 1g/L DL-malic acid, 1g/L Na-lactate, 1g/L Na-pyruvate, and 1g/L D-
mannitol and used when indicated. NH_4Cl was added at 1 g/L only when indicated. All enrichment
procedures were maintained at 30°C and the pH was adjusted to 6.8 with $NaHCO_3$ in the microbial
bioelectrochemical systems and the enrichment cultures without organic C addition. When the organic C
solution was used, the pH was adjusted between 6.3 and 6.5.

Design of microbial electrochemical system
The electrochemical system used for our enrichment were composed of two chambers separated by
an anion exchange membrane (AEM) (fumasep ® FAB-PK-130). The AEM was used to avoid the migration
of NH_4^+ ions to the anodic chamber. Each chamber had a total volume of one liter. The pH of the reactors
was adjusted to 6.8 at the beginning of each batch experiment. Each system had a 25 cm^2 square carbon
felt working electrode with a thickness of 0.7 cm and a 16 cm^2 square Pt-Ir grid as a counter electrode.
The carbon felt electrodes were conditioned using chemical treatment with HCl , a flush with ethanol and
a heat treatment at +400°C as described elsewhere by Paul et al. (2018). The systems were inoculated
with the flask enrichments used for N_2 fixation in presence of organic C (N-free medium). After
inoculation, the organic C supply was reduced to 10% of the initial supply in all reactors. The organic C
supply was then totally replaced by CO_2 supply. Two of the systems were polarized and two other
reactors were used as controls with open current voltage (non-polarized cathode enrichment, nPCE). The
working electrodes of the polarized cathode enrichment (PCE) were poised at a potential of -0.940 V vs.
saturated calomel electrode (SCE) used as a reference. The system was connected to a VMP3.0
potentiostat (BioLogic). The current was measured over time by a chronoamperometry method and cyclic
voltammetry was performed every 3-7 days at the beginning of enrichment periods. The monitoring of
the current intensity was used to monitor the efficiency in enriching the biofilms in autotrophic and/or
electrotrophic bacteria (Zaybak et al., 2013). The current density J was calculated using the surface of the
working electrode, i.e. 25 cm^2.

Before inoculation of the reactors, an initial chronoamperometry measurement was performed along
the first 2 days of operation with only organic carbon in the medium to determine the basal current
density in absence of bacteria. Two other abiotic reactors for 15 days were then implemented to
measure the current density in a medium without organic C.
The energy required for the production of microbial metabolites and for biomass growth was then
used to calculate the Coulombic efficiency of the PCE according to the equations 1-5:
151
152 Eq. 1 $2.1\,H_2 + CO_2 + 0.2NH_4^+ \rightarrow CH_{1.8}O_{0.5}N_{0.2} + 1.5H_2O + 0.2H^+$ for biomass production
153 (21 $mol_{e^-}/mol_{Nbiomass}$) (Wresta et al., 2021)
154
155 Eq. 2 $N_2 + 3H_2 \rightarrow 2NH_3$ for nitrogen fixation (3 $mol_e/mol_{Nfixé}$) (L. Zhang et al., 2022)
156
157 Eq. 3 $2CO_2 + 4H_2 \rightarrow CH_3COO^- + H^+ + 2H_2O$ for acetate production (8 $mol_e/mol_{CH3COO-}$)
158 (Wresta et al., 2021)
159
160 Eq. 4 $2H^+ + 2e^- \rightarrow H_2$ for H$_2$ production (2 mol_e/mol_{H2})
161
162 Eq. 5 $CE = \dfrac{n_e \times F \times n_{product}}{\int i\, dt}$
163
With $_{CE}$: the Coulombic efficiency in percentage of electron recovery in the circuit, n_e the number of
moles of electrons per moles of product ($mol_e/mol_{produit}$) calculated from the stoichiometric equations, F
the Faraday constant (96485 C.mol^{-1}), $n_{product}$ the number of $mol_{product}$ and i the current intensity.

Enrichment procedures
Two enrichment procedures based on a sequential method by enriching separately N$_2$ fixation and
the use of an inorganic energy source were carried out.
The first enrichment method was performed in three steps:
- The first step was performed in a 120 mL bottle with N-free medium supplemented with organic C
source. The headspace was composed of an O$_2$/N$_2$ mixture (10/90) at 0.5 bar (absolute pressure).
Cultures were carried out in bottles containing 50 mL of liquid and 70 mL of headspace. Subcultures of
these enrichments were performed every 7 days for 6 weeks. The time between 2 subcultures was then
reduced to 3-4 days using 10% of the volume of the previous culture (5mL/50mL).
- After 55 days of enrichment in 10 successive batches, the enriched cultures were used to inoculate
the cathodic chambers of the polarized microbial electrochemical system (Polarized Cathode Enrichment,
PCE) and the non-polarized microbial electrochemical system (non-Polarized Cathode Enrichment nPCE).
The same inorganic medium supplemented with 10% of the organic C source was fed each week to start
the enrichment of autotrophic bacteria. 80% of the medium was renewed every second week to promote
biofilm growth on the cathode. Headspace composition was monitored by GC and flushed with N$_2$ if pO$_2$
exceeded 10% of the gas volume. Organic C supply was stopped when a significant current density was
measured in the polarized systems with regard to the controls.
- In the third and final enrichment step, CO$_2$ was used as sole carbon source. A 80/20 (v:v) CO$_2$/N$_2$
atmosphere was set up in the headspace with trace amounts of O$_2$ (< 5%). The medium was replaced
every 15-30 days. The gas recycling vessel was filled with CO$_2$ and was replaced with a new one when O$_2$
exceeded 5% of the volume due to gas volume depletion. The nPCE controls were operated in the same
conditions as the polarized systems but without monitoring the current density. The only available energy
source in the nPCE controls was the organic C fed at the beginning of the enrichment.
In the second enrichment method, diazotrophic autotrophic bacteria were enriched in inorganic
medium supplemented with H$_2$ as sole energy source. These enrichments on H$_2$ were obtained by pre-
enriching in strict autotrophic bacteria using 50 ml of inorganic medium in a 120 ml bottle. The
headspace consisted of a mixture of 75/15/8/2 (v:v:v:v) H$_2$/CO$_2$/N$_2$/O$_2$ at an initial pressure of 1.5 bar
(absolute). Two two-weeks batches (30 days) were performed with NH$_4$Cl (1g/L) as nitrogen source in the
first stage of this enrichment. This nitrogen source was then replaced by N$_2$ as only N source to enrich N$_2$-
fixing bacteria in the second stage of enrichment. Centrifugation (10min, 7500 RPM, ~7500g) of 80% of

the initial medium (40mL) was done at each subculture every 15-20 days. The pellets obtained after
centrifugation were suspended in 5mL of sterile medium before being used for subculturing.

Medium analyses
NH_4^+, NO_2^- and NO_3^- concentrations were measured using a Gallery+ sequential analyzer (Thermo
Fisher Scientific). Two mL of samples were centrifuged at 13500 RPM (~12300g) and then filtered to
0.2μm with nylon membranes before being stored at 4°C. The remaining pellets were stored at -20°C and
used for community analysis. VFAs and other carbon compounds were measured on Clarus 580 GC
equipped with FID and a Dionex UltiMate 3000 HPLC as described elsewhere (Carmona-Martínez et al.,
2015; Moscoviz et al., 2019).
Total nitrogen was measured using a CHNS Flashsmart elemental analyzer (Thermo Fisher Scientific).
The sample (approximatively 2.5 mg) was weighed and was introduced into the oxidation/reduction
chamber of the analyzer. 200 mL of medium were sampled at each medium change. These samples were
dried for 4-5 days at 60°C. The samples were then freeze-dried and then grounded with a mortar. Two to
four mg of each sample was used in the CHNS analyzer. The nitrogen content was then compared to the
dry weight measured before freeze-drying to determine the mass of nitrogen in the medium. No CHNS
analysis was performed on H_2 –based enrichment due to a low culture volume (50mL). A C/N ratio was
then calculated to evaluate the state of the biomass and possible inhibitions of N_2 fixation (Khan et al.,
2016; W. Zhang et al., 2020).
Nitrogen present in the biomass (biofilm and planktonic) was estimated by calculation from 16S rDNA
qPCR. The rrnDB-5.7 database was used to estimate the actual bacterial amount from 16S rDNA qPCR
using theoretical 16S rDNA copies per strain, genus or family (Stoddard et al., 2015). Then, the theoretical
number of bacteria was used to determine the nitrogen content in the biomass using the theoretical
average dry mass of an *Escherichia coli* cell of 216×10^{-15} g/bacterium and with a theoretical relative mass
of nitrogen in microbial biomass, ie. 11.4% according to the average biomass formula $CH_{1.8}O_{0.5}N_{0.2}$ (Heldal
et al., 1985; Loferer-Krößbacher et al., 1998). The nitrogen present in the biomass was therefore
estimated using the formula below:
223
224 Eq. 6 $N\_(in\ biomass) = theoretical\ bacteria\ count \times mass\ E.coli \times relative\ mass\ of\ N\ in\ biomass$
225
With $N_{in\ biomass}$ the nitrogen concentration in the medium from biomass in mg_N/L, the *theoretical*
bacteria count based on the bacteria concentration in each enrichment (number of bacteria/L), *mass E.*
coli a constant of $2.16\ 10^{-10}$ $mg/cell_{E\ coli}$ and *relative mass of N in bacteria* is 11.4% of the dry mass.

Gas analysis and acetylene reduction assay (ARA)
CO_2, H_2 and N_2 used in the headspace of the enrichments were provided at laboratory grade. Pure
ethylene was also supplied for calibration of the gas chromatography ethylene measurement.
Headspace compositions and pressures were analyzed every 1-2 days. The pressure was manually
measured with a Keller LEO 2 manometer (KELLER AG, Wintherthur, Switzerland). Gas analyses were
carried out on a Perkin Elmer Clarus 580 GC equipped with RT-Q-Bond and RT-Msieve 5Å columns with a
TCD allowing the quantification of H_2, CO_2, N_2, O_2 and CH_4 with Ar as carrier gas as described by A.
Carmona-Martínez (Carmona-Martínez et al., 2015). Acetylene and ethylene were measured using a
Perkin Elmer Clarus 480 GC equipped with RT-U-Bond and RT-Msieve 5Å columns with TCD and He as
carrier gas as described in a previous work (Carmona-Martínez et al., 2015).
The acetylene reduction assay (ARA) was performed during enrichment to quantify the rate of N_2
fixation using the ability of nitrogenases to reduce acetylene to ethylene. This reaction occurs at a rate
proportional to the rate of N_2 fixation according to the theoretical ratio $C_2H_2:N_2$ (3:1) (Bergersen, 1970).
The ARA was performed only after 18 batch cycles for H_2 enrichment and 11 batch cycles in polarized
cathode enrichment fed with CO_2 (340 days). A specific N_2-fixing activity was then calculated with the N_2-
fixing bacteria measured by quantification of the *nifH* gene used as a marker for these bacteria. The
acetylene used for the acetylene reduction assay (ARA) was obtained by adding calcium carbide (CaC_2) in
water and recovered in a bag. The acetylene concentration in the bag was then measured by gas
chromatography. Gas from the bag was added to each enrichment to reach a composition of 10% V/V
acetylene and an equivalent amount of gas was removed from the headspaces. Ethylene production was

then daily monitored for 7 days in the microbial electrolysis systems (PCE and nPCE) and 15 in the H_2
bottles by the Perkin Elmer Clarus 480 GC with TCD. To ensure separation of ethylene from CO_2 on the
RT-U-Bond column, a CO_2 trap with sodium hydroxide (6M NaOH) was used at the time of sampling. After
the ARA method was completed, the headspaces were flushed with N_2 and the gas recycling system was
changed.

Community sequencing and biomass quantification
The microbial communities were quantified using the 16S rDNA qPCR to determine the total bacterial
concentration and *nifH* gene qPCR for N_2-fixing bacteria. In parallel, 16S rDNA sequencing was performed
to identify major members of each community. This sequencing was also necessary to convert the
amount of 16S rDNA to total bacteria using the rrnDB-5.7 database. To analyze the communities present
in suspension, 1.8 mL of sample were collected for qPCR. For the cathodes, 1 cm^2 was recovered at
several times. The piece of carbon felt was then chopped with a sterile scalpel before being immersed in
20 mL of sterile inorganic media. 1.8 mL was then recovered after shaking 20 mL of medium to resuspend
as much biomass as possible. These samples were then centrifuged 10 min 13500 RPM (12340g). The
supernatant was discarded and the pellets retained for DNA extractions. After qPCR, the concentrations
measured in the electrode samples were expressed considering the volume of medium.
Genomic DNA was extracted using the PowerSoil™ DNA Isolation Sampling Kit (MoBio Laboratories,
Inc., Carlsbad, CA, USA) according to the manufacturer's instructions. The qPCR amplification program
was performed in a BioRad CFX96 Real-Time Systems C1000 Touch thermal cycler (Bio-Rad Laboratories,
USA). For the analysis of total bacteria, primers 330F (ACGGTCCAGACTCCTACGGG) and 500R
(TTACCGCGGCTGCTGGCAC) were used. For the bacteria qPCR mix: SsoAdvanced™ Universal SYBR Green
Supermix (Bio-rad Laboratories, USA), primer 330F (214 nM), primer 500R (214 nM), 2 µL of DNA, and
water were used to a volume of 12 µL. The qPCR cycle was as follows: incubation for 2 min at 95°C and 40
cycles of dissociation (95°C, 10 s) and elongation (61°C, 20 s) steps. The results were then compared to a
standard curve to obtain the copy number of the target in the sample. Both the 16S rDNA concentration
of the PCE media and the cathodes are considered in the calculation of the total 16S rDNA concentration
of the PCE. These concentrations are used as an indicator of the biomass present and the use of a
database of the number of 16S operons per bacterial genome was used to estimate the actual amount of
bacteria.
The presence of N_2-fixing bacteria was monitored by qPCR of the *nifH* gene of the Fe-Fe subunit of
nitrogenases (Dos Santos et al., 2012; Gaby & Buckley, 2012). All qPCR amplification programs were
performed in a BioRad CFX96 Real-Time Systems C1000 Touch thermal cycler (Bio-Rad Laboratories,
USA). The primers PolF-TGCGAYCCSAARGCBGACTC and PolRmodify reverse-AGSGCCATCATYTCRCCGGA
were used (Poly et al., 2001). The mixture: 6µl SsoAdvanced™ Universal SYBR Green Supermix (Bio-rad
Laboratories, USA), F primer (500 nM), R primer (500 nM), 2 µL of DNA and water was used up to a
volume of 12 µL. The qPCR cycle was as follows: incubation for 2 min at 95°C and 40 cycles of dissociation
(95°C, 30 s) and elongation (60°C, 30 s) steps. Then, the results were compared to a standard curve to
obtain the number of copies of the target in the sample. These two quantifications allow us to calculate
the ratios of N_2-fixing bacteria per total bacteria of the enrichments at different points to track the
enrichment of N_2-fixing bacteria. This ratio can also help us to derive hypotheses on the functioning of
our communities that can be completed by the analysis of the communities during sequencing.
After quantification, our enriched communities were sequenced according to their 16S rDNA and the
results are available on NCBI repository PRJNA838547, Biosample SAMN28447998-SAMN28448066. The
V3-V4 region of the 16S rDNA was amplified using universal primers as reported elsewhere (Carmona-
Martínez et al., 2015). The PCR mixture consisted of MTP Taq DNA Polymerase (Sigma-Aldrich, Germany)
(0.05 u/µL) with enzyme buffer, forward and reverse primers (0.5 mM), dNTPs (0.2 mM), sample DNA (5-
10 ng/µL), and water to a final volume of 60 µL. 30 cycles of denaturation (95°C, 1 min), annealing (65°C,
1 min), and elongation (72°C, 1 min) were performed in a Mastercycler thermal cycler (Eppendorf,
Germany). A final extension step was added for 10 min at 72 °C at the end of the 30th amplification cycle.
PCR amplifications were verified by the 2100 Bioanalyzer (Agilent, USA). The GenoToul platform
(Toulouse, France http://www.genotoul.fr) used an Illumina Miseq sequencer (2 x 340 bp pair-end run)
for the sequencing reaction. The raw sequences obtained were analyzed using bioinfomatic tools.

Mothur version 1.39.5 was used for cleaning, assembly and quality control of the reads. Alignment was
performed with SILVA version 128 (the latter was also used as a taxonomic contour).

Data analysis
All results were analyzed using R (4.2.0) and Rstudio (2022.07.1) for calculations and graphics. The
Tidyverse package was used for data manipulation(*Tidyverse*, n.d.). The packages ggplot2, ggpubr, scales,
cowplot, corrplot and palettetown were used for the graphical representations. Visual representation of
bacterial relative abundances was performed with the phyloseq package (McMurdie, 2011/2023).
Inkscape software was also used to edit the graphs when necessary. The uncertainties shown for the
values presented are standard deviations. All data and scripts used here are available online (Rous, 2023).
310

<div align="center">**Results and discussion**</div>

Nitrogen fixation after 340 days of enrichment
N_2 fixation was quantified at the day 340 of the enrichment using acetylene reduction assays (ARA).
This assay was performed in H_2-fed bottle enrichments (named 'H$_2$E'), in polarized cathode enrichment
(named 'PCE') and in the non-polarized cathode enrichments as controls (named 'nPCE'). As shown in
Figure 1, the ARA results confirmed the N_2 fixation capacity of the enriched communities (Bergersen,
1970). This indicates that the cathode and/or H_2 was used as energy sources for N_2 fixation both in PCE
and in H_2E bottles. The average rates were similar in both enrichment methods with 32±17 µmolC$_2$H$_4$/L.d
in PCE and 36±13 µmolC$_2$H$_4$/L.d in H_2E bottles. The PCE corresponding N_2 fixation rates ranged from 0.12
mg$_{Nfixed}$/L.d (minimum) to 0.51 mg$_{Nfixed}$/L.d (maximum), which is consistent with the rate of 0.2
mg$_{Nfixed}$/L.d estimated by Rago et al., (2019) and also with N_2 fixation rates reported for soil bacteria
(Hardy et al., 1973; Kifle & Laing, 2016). Despite these significant N_2 fixation rates and the long duration
of the experiments, the rate of ammonium production in solution remained lower than 0.07 mg$_N$/L.d at
the day 340 (Table 1), indicating that most of the fixed N_2 was probably rapidly used by bacteria.
325

327   **Figure 1** - Reduction rate of acetylene in µmol C$_2$H$_4$/L.d in the different reactors after 340 days of operation . For H$_2$
328 condition, 3 bottles were used for the acetylene reduction assay with one injection in each bottle, i.e. 3 measurements. For
329 PCE, 2 reactors were used and 4 injections were made to validate the repeatability of the measurements, which gives 8
330 measurements for each of these two conditions. For nPCE, 2 reactors were used which gives 2 measurements for each
331   conditions

The ability of the microbial communities to fix N_2 was also assessed by qPCR of the *nifH* gene (Dos Santos et al., 2012; Pogoreutz et al., 2017). The amounts of N_2-fixing bacteria after 340 days of enrichment are reported in Table 1**Error! Reference source not found.**. The average *nifH* gene concentration in PCE was estimated at 7.8 10^7 copies$_{nifH}$/mL, two orders of magnitude higher than the average concentration of 8.3 10^5 copies$_{nifH}$/mL measured in the H_2 enrichment bottles. This observation was surprising since N_2-fixation rates were similar in both configurations (Figure 1). This suggests that the fixation rate per *nifH* copy was much higher in H_2E than in PCE. The estimated specific activities per *nifH* copy were indeed of 0.2±0.3 $\mu mol_{C2H4}/10^8$ copies$_{nifH}$.d in the PCE and 2.1±0.7 $\mu mol_{C2H4}/10^8$ copies$_{nifH}$.d in H_2 enrichment bottles (Table 1). Furthermore, the comparison between *nifH* gene and 16S rRNA gene copy numbers gives an idea of the proportions of nitrogen fixing bacteria in each microbial community. Interestingly, this proportion was 18% in H_2 enrichment bottles that was four times higher than the value of 4.6% in PCE (Table 1**Error! Reference source not found.**). Therefore, N_2-fixing bacteria constituted a smaller proportion of the bacterial populations in PCE than in H_2E and only a small proportion of the bacteria participated to N_2 fixation in the PCE. In nPCE controls, the biomass was higher than in H_2 enrichment and the *nifH*/16S ratio lower (Table 1). The lack of N_2 fixation observed in nPCE indicated that the microbial biomass maintained along the experiment without the need of fixing N_2. It is assumed that the microbial biomass was partly degraded and used as a source of C and N by other bacteria. Thus, the presence of NH_4^+ in the nPCE was very likely related to cell lysis since no measurable fixation was detected by ARA.

Table 1 - Measurements of average ammonium, nifH gene, 16S gene, and nifH/16S ratio concentrations averaged at 340 days for the three experimental configurations. Averages measured on one batch for the two polarized cathode enrichment (PCE), the two non-polarized Cathode Enrichments (nPCE) and the six H_2 enrichment bottles (H_2E)

	PCE	nPCE	H_2E
N-NH_4^+ *mg/L.d*	0.07±0.01	0.07±0.01	0.04±0.03
nifH gene copies *copies/mL*	7.8±9.5 10^7	2.5±2.8 10^6	8.3±2.2 10^5
16S rDNA gene copies *copies/mL*	1.7±1.9 10^9	1.4±0.2 10^8	4.6±1.5 10^6
nifH/16S	3.8 %	1.7 %	19.0 %

Current density and Autotrophic/electrotrophic enrichment in polarized cathode enrichment

The average current density for the two PCE over experimental time is shown in Figure 1. As the current measured at the cathode was negative by convention, a more negative current corresponds to a higher reduction activity. Regarding the current densities in the abiotic systems, the average current densities were measured at -0.75 A/m² for two times four days with an organic C source and -1.1 A/m² for 16 days with only CO_2 as the carbon source.

The average current density measured in the PCE was not different from the current density measured in the abiotic control (approximately -1 A/m²) for the first 45 days of operation (days 55 to 100 of the enrrichment). The medium was then changed for a medium with no organic C. An sharp increase in current density to -5 A/m² was observed in the PCE with regards to the average current in the abiotic controls and standard deviation up to -2 A/m² (Figure 1). The higher current density was assumed to be associated to the electroactive activity of electron uptake by enriched bacteria. The current density in the PCE then continuously increased until day 250 of the enrichment to reach a value of -15 A/m².

The high current consumption observed after 250 days of enrichment strongly suggests a specific enrichment in electroactive bacteria. Indeed, current consumption was 5 to 10 times higher than in average for the abiotic controls, and was likely due to the direct use of electrons by bacteria in a cathodic biofilm as proposed by Z. Zaybak et al (Zaybak et al., 2013). Compared to the current densities obtained Rago et al. (Rago et al., 2019) in the order of magnitude of -10 mA/m² at the same potential (-0.7 vs SHE), the current densities observed here (-5 to -10 A/m²) were about 1000 times higher. These current density

levels are close to those measured by Zhang et al. (L. Zhang et al., 2022) who reported a maximum of -10
A/m² at the same applied potential.
After 230 days, power failures occurred, interrupting temporarily the current supply to the cathodes.
An important decrease of the current density was observed afterwards, down to -5 A/m² after 260 days
and -3 A/m² after 320 days. The lower current density reflected a change in the functioning of the
microbial communities, leading to less electron exchange with the cathode.
380

Figure 1 - Mean current density measured for the two PCE (blue line). Levels shown in red correspond to theoretical
mean current density and standard deviation estimated from two abiotic electrochemical systems current densities. For
abiotic electrochemical system, one batch of 2 days were made with Organic C and one batch of 16 days with CO_2. The
peaks observed are due to the batch operation of the PCE with disturbances each time the medium was renewed. Power
failures occurred at 230 days and 260 days. The time indicated on the x axis corresponds to the experimental time starting
at day 0 of the enrichment where soil samples were first introduced in the bottles with a medium containing organic C. Day
55 corresponds to the start of microbial electrochemical system with the precultured communities. The dashed line on day
110 corresponds to the passage on CO_2 as the sole carbon source in PCE

Biomass quantification
Bacterial biomass was monitored in the enrichments by measuring 16S rDNA concentrations by qPCR
(Figure 2). At day 131 (18 days after switching from organic C to CO_2), the average 16S rDNA
concentrations measured in the polarized cathode enrichment (PCE) and in the non-polarized cathode
enrichment (nPCE) controls were $4.6\pm0.4 \ 10^9$ and $4.2\pm1.2 \ 10^9$ copies16SrDNA/mL, respectively. These
concentrations corresponded to $9.4\pm1.0 \ 10^8$ and $8.0\pm1.5 \ 10^8$ bacteria/mL, respectively, as presented in
Figure 2. These bacteria concentrations resulted from the first enrichment phase with organic C. During
this phase, organic substrates were used as carbon and energy sources for biomass growth in both
configurations (PCE and nPCE). At day 214, the 16S rDNA concentration in the PCE was estimated at
$3.3\pm2.1 \ 10^9$ bacteria/mL, corresponding to a biomass increase by a factor of 3.5 between 131 and 214
days (Figure 2). At the same time, the bacterial concentration dropped in nPCE from $8.0\pm1.5 \ 10^8$
bacteria/mL to $3.0\pm0.8 \ 10^8$ bacteria/mL. This drop was explained by the lack of available energy source
for growth, which led to a sharp decrease of the bacterial populations. At day 214, the 16S rDNA
concentration in PCE was therefore 11-fold higher than in nPCE controls. This difference is consistent
with the difference reported by Rago *et al.* (Rago et al., 2019) between polarized and non-polarized
conditions, with electroactive biocathodes enriched in autotrophic diazotrophic bacteria. These results
suggested that the enriched microbial communities were able to use the electrodes polarized at -0.7 V vs.
SHE as sole energy sources to grow while fixing N_2.
In the H_2-fed enrichment (H_2E) bottles, the 16S rDNA concentration steadily decreased along the
experiment. The concentration decreased from $1.1\pm1.3 \ 10^9$ bacteria/ml at the beginning of the
enrichment down to $1.3\pm0.3 \ 10^6$ bacteria/ml after 340 days (Figure 2). These concentrations appear
lower than the biomass concentrations observed in the PCE medium at the same time of enrichment, ie.
$4.5\pm5.9 \ 10^6$ bacteria/mL at day 244 and $8.6\pm8.8 \ 10^6$ bacteria/mL at day 340. These results confirm that
bacterial growth was higher on the cathodes than in an H_2 supplied environment. It was therefore

concluded that the PCE provided more favorable environment for biomass growth than H_2 fed bottles as
the surface provided by the electrode was likely favorable for biofilm growth.

We also calculated the *nifH*/16S ratio representing the part of bacteria able to fix N_2 among the total
bacteria. A ratio of 0.06% of *nifH* gene copies per 16S rDNA copy was measured for samples at the very
beginning of enrichment, both for H_2E and PCE. After 131 days, corresponding to the switch to CO_2 as
sole C-source, this level increased to 3.1% in PCE and 1.5% in nPCE control. These results are consistent
with an enrichment in diazotrophic bacteria during the enrichment phase on organic carbon (Bowers et
al., 2008). The bacterial enrichment in nPCE was likely possible as the organic C was used by the bacteria
as energy source. After 214 days, the level decreased to 1.9% in PCE but remained higher than the level
at the beginning of enrichment. This variation suggests interactions within the community that favored
the growth of non-N_2 fixing bacteria after the shift to CO_2 as sole C-source. After 340 days, the ratio of
nifH to 16S rDNA was 3.8% as presented in Table 1. In parallel, a ratio of *nifH* to 16S rDNA of 90% was
measured for H_2E at 244 days. Therefore, most of the bacteria were able to fix N_2 in H_2E bottles,
confirming the efficient enrichment in diazotrophic bacteria (Bowers et al., 2008). Given the loss of
biomass observed in H_2E during the experiment (Figure 2), this high ratio corresponded likely to the
surviving bacteria that were selected on their ability to fix N_2. The ratio measured in these H_2E then
decreased down to 19%, suggesting a decrease in N_2-fixing bacteria biomass.

As previously mentioned, after 230 days, power failures occurred and interrupted the polarization of
the electrodes. These interruptions impacted the microbial communities with a decrease in biomass
concentration to $8.1 \pm 7.6 \ 10^8$ bacteria/mL at 244 days and $5.5 \pm 6.0 \ 10^8$ bacteria/mL after 340 days
compared to the concentration of $3.3 \ 10^9$ bacteria/mL measured at 214 days. At the same time, the
nifH/16S rDNA ratio increased up to 5%, indicating that N_2-fixing bacteria were more resistant.
Nevertheless, a decrease was observed in *nifH* quantities, from $2.3 \ 10^8$ copies$_{nifH}$/mL after 214 days to 7.8
10^7 copies$_{nifH}$/mL after 340 days.

Figure 2 - Bacteria concentrations over time in the different enrichments calculated from 16S rDNA qPCR
quantifications in bulk and biofilm. Green disks correspond to H_2 enrichments in bottles, blue triangles correspond to
polarized cathode enrichment (PCE), red squares correspond to controls in non-polarized cathode enrichment (nPCE). The
partially purple symbols marked Org correspond to the first phases of enrichment with organic C for the PCE and nPCE. The
arrow indicates the transition from bottle enrichments to cathode enrichments in microbial electrochemical systems for
445   PCE and nPCE. Error bars correspond to the calculated standard deviation.

N quantification and coulombic efficiency

Total N contents of the different experiments are shown in Figure 3a. The total N corresponded to the
sum of the nitrogen measured in the liquid phase by N-ion concentration analysis (N-NH_4^+, N-NO_3^-, N-
NO_2^-), in the medium by CHNS elemental analysis for PCE and nPCE, in the suspended biomass from qPCR
results only for H_2E where the dry mass was not measured, and on the electrode based on the bacterial
concentrations. The total N concentration was estimated after 131 days of enrichment (ie. before the

shift to CO_2 as sole C source) at 25.5±0.4 mg_N/L and 25.8±4.7 mg_N/L in the PCE and in the nPCE controls,
respectively. After 214 days of enrichment, total N increased up to 87.6±56.1 mg_N/L in the PCE with
regards to the low value of 14.8±7.0 mg_N/L in the nPCE controls. Ammonium represented only a small
fraction of the total N in the PCE. The maximum ammonium concentration observed at a batch end in the
PCE was 1.5±0.3 mg_N/L at the day 340 of the enrichment in comparison with the maximum value of
4.5±0.4 mg_N/L found in nPCE control after 131 days (1.4 mg_N/L at the day 340) .(Figure 3b). The average N
fraction in the form of ammonium was therefore of 12% in nPCE and only 1.6% in PCE. It was thus
assumed that the higher level of ammonium in nPCE was related to the decay of biomass in absence of
energy sources. In counterpart, the ammonium produced by N_2 fixation in the PCE was likely directly used
for protein synthesis as suggested elsewhere (Temple et al., 1998). In addition, nPCE $N-NO_3^-$
concentrations were 5 to 15 times lower than $N-NH_4^+$ concentrations with maximum values of 0.9 mg_N/L
$N-NO_3^-$ in the nPCE control and 0.2 mg_N/L $N-NO_3^-$ in PCE. An average concentration of 0.1±0.2 mg_N/L N-
NO_3^- in H_2E bottles was also observed. NO_2^- concentrations were negligible.

The C/N ratios are shown in Figure 3. C/N ratios in the PCE were around 10 after 131 and 214 days in
comparison with ratio of 5 to less than 1 in the nPCE control. These ratios are consistent with the
theoretical ratio of 8 to 10 assumed for microbial biomass (Goldman et al., 1987; Heuck et al., 2015;
Yaghoubi Khanghahi et al., 2021). In contrast, the low ratios in nPCE indicated that the dry mass was most
probably not only composed of microbial cells. Interestingly, the level of C/N reached 30/1 in the PCE
after 244 days of enrichment. Such increase could be related to the release of carbon molecules such as
exopolysaccharides. the C/N ratio finally dropped down to 1.8 after 340 days, confirming cell lysis as
previously suggested.

The H_2 enrichment bottles (H_2E) showed an average $N-NH_4^+$ accumulation of 35.6±36 $\mu g_N/L$ after 131
days and an average of 0.1±0.2 mg_N/L over the duration of enrichment. This concentration represented a
small fraction of the total N at the very beginning of the enrichment, with an estimated concentration of
29.4±28 mg_N/L. The N concentration decreased during the enrichment, which is consistent with the
biomass loss as shown in Figure 2. H_2E were therefore less efficient for N_2 accumulation than polarized
cathode enrichment, with a lower microbial biomass production.

Figure 3 - (a) Total N concentration based on N measured in dry weight from medium samples for polarized cathode
enrichment (PCE) and non-polarized cathode enrichment (nPCE) and estimated from biomass for H_2-fed enrichment (H_2E),
on ions ($N-NH_4^+$, $N-NO_3$, $N-NO_2$) concentrations, and on N estimated form the biomass in biofilm for PCE and nPCE, (b) N-

Current densities and rates of acetate production, N_2 fixation and biomass growth are shown in Table
1. Coulombic efficiencies associated with each reaction were estimated based on these results (Table 1).
During the first 214 days of enrichment, 0.6 to 3.3% of the electrons were used for N_2 fixation in the two
PCE. In comparison, efficiencies of 0.5% and 20% for NH_4^+ synthesis was reported in two recent works
carried out under similar conditions (Yadav et al., 2022; L. Zhang et al., 2022). As the amount of fixed N
was highly dependent on biomass accumulation, negative results were obtained at day 244 when
biomass started to decrease. The electrons used at the cathode for biomass synthesis during the first
period (131 to 214 days) accounted for 2.8% and 17.3% in the two PCE. These high coulombic efficiencies
were probably also associated to microaerophilic conditions. As acetogenic bacteria do not tolerate the
presence of oxygen, dissolved O_2 was very probably consumed in some part of the biofilm, leaving other
parts in strict anaerobic conditions more favorable for acetogenic bacteria growth.
The H_2 recovered in the headspace of the PCE accounted for 12 to 22% of the electrons supplied to
the cathode as presented in Table 1. Therefore, H_2 was not related to the biological activity and mostly
resulted from an abiotic reaction at the cathode.
Interestingly, a significant production of acetate was also observed. An average rate of 149.1
µmol/L.d and 421.6 µmol/L.d were measured in both PCE for the period from day 131 to day 214, as
presented in Table 1. Acetate production almost stopped with the power failures with acetate measured
only on one to two batches per PCE. This decrease correspond to rate of 61.0 µmol/L.d and 57.4 µmol/L.d
of acetate in both PCE. Acetate production accounted for 7.9 and 39% of the cathodic electrons during
the first period (up to 214 days) and for less than 5% after power failures. To explain the decrease in
acetate production, biomass growth and power consumption, it was hypothesized that acetate
production might have been due to a specific loss of members able to fix CO_2, and more especially
electrotrophic bacteria, within the enriched community. Thus, with acetate no longer being produced,
heterotrophic bacteria did not have enough organic C to sustain their growth, causing their decrease.
Therefore, electrotrophic bacteria responsible for CO_2 fixation and heterotrophic bacteria that could also
be H_2 dependent for N_2 fixation were greatly affected, leading to a decrease in the reduction reactions at
the cathode and in current density. In addition, acetate was not found in the H_2E bottles, indicating the
absence of acetogenesis and an important difference in microbial pathways and/or communities.

Table 1 - Current densities, rates and coulombic efficiencies for the two polarized cathode enrichment (PCE) over two
different periods of current consumption with CO_2 as sole carbon source. During the first period (131-214 days) current
consumption increased, whereas during the second period (>215) current consumption decreased after several power
outages (see Figure 1).

		131-214 days		> 215 days	
		PCE 1	PCE 2	PCE 1	PCE 2
J A/m²	Mean	7.7±3.1	5.9±2.4	6.2±4.1	3.8±2.2
	Max	28.2	11.3	19.5	9.9
Acetate µmol/L.d	Mean	149.1±203.1	421.6±216.7	57.4±151.8	61.0±106.5
N µmol/L.d	N-NH_4^+	0.108±0.225	0.155±0.249	2.0±1.4	1.5±1.9
	N Bulk Dry weigth	10.3±6.0	27.5±13.4	9.0±6.4	12.1±6.4
	N Biomass theoretical (16S/bact)	16.7	80.4	-3.0	-7
Biomass bact/L.j	Bulk	$0.4 \pm 0.5\ 10^9$	$3.5 \pm 2.0\ 10^9$	$0.4 \pm 0.5\ 10^9$	$0.2 \pm 0.1\ 10^9$

	Cathode	$0.9\ 10^{10}$	$4.6\ 10^{10}$	$-2.6\pm3.5\ 10^{10}$	$-5.9\pm7.8\ 10^{10}$
Coulombic efficiency %	CO_2 to Acetate	7.9 ± 9.5	30.9 ± 8.9	2.1 ± 5.6	5.0 ± 6.6
	H^+ to H_2	22.3 ± 15.4	12.1 ± 8.8	9.4 ± 13.5	14.4 ± 14.4
	N_2 fixation	0.6 ± 0.2	3.3 ± 0.8	0.2 ± 1.0	-0.5 ± 2.2
	Biomass growth	2.8 ± 0.5	17.3 ± 5.3	-2.5 ± 4.0	-8.3 ± 14.2
	Total	33.6 ± 14.4	63.6 ± 9.0	9.2 ± 16.5	10.5 ± 18.8

Microbial communities

519
16S rDNA sequencing was performed at the beginning of the culture, and at 214 or 232 days of
enrichment in polarized cathode enrichment (PCE) and in H_2-fed enrichment bottles, respectively. The
sampling days were selected because they were associated to a high microbial activity (high current
densities and high biomass concentrations). In H_2E, the *nifH*/16S abundance ratio was also maximum
(90%) at day 232. Averages of relative abundances of the bacterial families are shown in Figure 5.
At the beginning of the PCE in bottles fed with organic C, communities were strongly dominated by
only four families accounting for more than 96% of total sequences: *Clostridiaceae* (14%),
Enterobacteriaceae (65%), *Enterococcaceae* (10%) and *Lachnospiraceae* (7%). A dominance of the
Enterobacteriaceae family (mainly of the *Citrobacter* genus) was observed. All these families, and in
particular *Clostridiaceae* and *Enterobacteriaceae* families are supposed to possess numerous genera
capable of N_2 fixation and possessing the *nifH* genes (Huang et al., 2019; Lin et al., 2012; Minamisawa et
al., 2004). This observation indicated a rapid selection of bacteria having the capability of N_2-fixation at
the early stage of enrichment.

Figure 4 - Average relative abundance of bacterial families in H_2E and PCE at the beginning of enrichment (top barplots) and after 214 (for PCE) and 232 days (for H_2E) of enrichment (bottom barplots). Only families with an average relative abundance above 5% in the replicates of one condition are shown.

In the H_2-based enrichments, and in contrast with the enrichments in PCE, *Pseudomonadaceae* (45%) family was mostly dominant. *Nocardiaceae* (17%), *Flavobacteriales* (14%), *Xanthomonadaceae* (8%) and *Comamonadaceae* (7%) were also present. These families, with the exception of *Flavobacteriales*, are also known to have members possessing the set of genes necessary for N_2 fixation (Dos Santos et al., 2012; Ghodhbane-Gtari et al., 2019; Huda et al., 2022). These families accounted for 77% of the sequences which is high compared to the *nifH*/16S rDNA ratio of less than 1% at the same time point. This suggests that either the *nifH* primers were not adapted to these specific species or that the species found at this point did not possess the genes for nitrogenases. As the H_2E cultures started on a medium containing NH_4Cl, the presence of this source of nitrogen was likely favorable to the growth of non-diazotrophic bacteria.

After 214 days of enrichment, PCE major families found in the PCE were affiliated to *Peptococcaceae* (29%), *Xanthomonadaceae* (18%), *Rhodospirillaceae* (6%, *Azospirillum*) and *Comamonadaceae* (10%). A clear shift in microbial communities from the beginning of the experiment was therefore observed. Only *Enterobacteriaceae* (6%) family was maintained although at minor relative abundance. Members of *Clostridiales incertae sedis* absent from original inoculum also appeared on polarized cathode. These families are known to exhibit the role of plant growth promoting bacteria (PGPB). These communities could thus be beneficial when used as living fertilizers (Cassan & García de Salamone, 2008; Rojas-Tapias et al., 2012; Singh et al., n.d.). The *Comamonadaceae* as well as the *Enterobacteriaceae* families mostly include heterotrophic species, which would be consistent with our hypotheses about the existence of interactions between heterotrophic and electrotrophic populations (F. Liu et al., 2011; Wu et al., 2018). More precisely, the *Peptococcaceae* sequences were affiliated to species *Desulforamulus ruminis* (>98%). This species was already described for their ability to fix N_2 (Postgate, 1970). The *Desulforamulus* and *Desulfotomaculum* genera have also several species able to grow with H_2 and CO_2 as energy and C sources (Aullo et al., 2013; Klemps et al., 1985; Zaybak et al., 2013). They were previously reported to be able to produce acetate by CO_2 reduction through the Calvin cycle (Klemps et al., 1985), and some were already found in microbial electrochemical system on a biocathode producing acetate (Zaybak et al., 2013). The other main family, *Xanthomonadaceae*, was represented by several genera with a majority of *Pseudoxanthomonas*. In this genus, some members were identified as N_2 fixers with a need of external organic C source, exhibiting a mixotrophic metabolism depending on the environmental conditions (J. Hu et al., 2022; Ryan et al., 2009). Sequences associated to the *Rhodospirillaceae* family were mainly affiliated to the species *Azospirillum lipoferum* which is able to grow in autotrophy with H_2, CO_2 and N_2 (Tilak et al., 1986). This soil bacterium is also known for its role as a PGPB with a capacity to solubilize phosphates, making it a good candidate as a fertilizer (Cassan & García de Salamone, 2008; Tilak et al., 1986). Interestingly, many of the identified bacteria in the polarized cathode enrichment were previously described to possess the N_2-fixing genes and capability. This supports the fact that the primers were not able to amplify the full diversity of *nifH* genes from these communities.

E The *Xanthobacteraceae* (51%), *Peptococcaceae* (17%, identified as *Desulforamulus*) and *Chitinophagaceae* (8%) families were found to be dominant in H_2E bottles at day 214. The *Xanthobacteraceae* family was mostly represented by the species *Xanthobacter autotrophicus* which is known as N_2-fixing HOB (Wiegel, 2005). This species was already been used for N_2 fixation by Liu *et al.* (C. Liu et al., 2017) in an hybrid system using the H_2 produced by a cathode. *Xanthobacter autotrophicus* was also found in the medium of the polarized cathode enrichment but in lower abundance (< 5%). Therefore, the community enriched with H_2 was mostly composed of N_2-fixing bacteria, as also supported by the high *nifH*/16S ratio (90%). After 214 days of enrichment, diazotrophic HOB were specifically selected.

The presence of mixotrophic and heterotrophic bacteria in the PCE suggested that carbon-based interactions could have occurred. Acetate was the only abundant soluble carbon metabolite found in these enrichments (see Table 1). Therefore, acetate was assumed to be used as intermediate for carbon and electron transfer between electrotrophic or hydrogenotrophic homoacetogens, *e.g. Desulforamulus rumnis,* and heterotrophic bacteria such as *Comamonas sp.*).

Furthermore the low concentration of $N-NH_4^+$ in the PCE before day 210 (Table 1) was probably due
to its rapid consumption for bacteria growth. Considering these hypothesis, a conceptual scheme of
microbial interactions between the main bacterial families in the PCE was proposed and is presented in
Figure 6. The presence of heterotrophic bacteria and their potential use of O_2 as a final electron acceptor
was also considered. The concentration of dissolved O_2 would have decreased in a deep layer of the
biofilm due to its use by heterotrophic bacteria. A structure of the biofilm in two layers could then be
proposed with a first layer composed mainly of homoacetogens fixed on the cathode and reducing CO_2 to
acetate, and a second layer composed mainly of heterotrophic bacteria using acetate and dissolved O_2 to
sustain their growth. It was assumed that bacteria in the first layer would not access to N_2 that would be
mostly fixed by organisms of the second layer.

Figure 5 - Conceptual scheme of microbial interactions occurring in polarized cathode enrichment (PCE) after
enrichment for N_2 fixation with inorganic energy and carbon sources

Conclusion

Enrichment cultures of N_2-fixing bacteria were successfully carried out in H_2-fed bottles (H_2E) and in
polarized cathode enrichment (PCE). Both methods showed significant N_2 fixation after 340 days of
enrichment. The microbial communities selected were able to fix N_2 with CO_2 as sole carbon source and
H_2 or cathodic electrons as sole energy sources. Biomass growth on the cathode up to 4.6 10^{10}
bacteria/L.d is another evidence of autotrophic growth in the PCE while bacterial growth was much lower
in the H_2E. Current consumption confirmed the activity of electrotrophic bacteria in the PCE. As the
coulombic efficiency of N_2 fixation was low with a maximum of 3.3% and considering the low
concentrations of NH_4^+, it was concluded that the major part of the nitrogen was incorporated into
microbial biomass during the enrichment procedure. Interestingly, acetate was also produced in the PCE
corresponding to a coulombic efficiency of 27%. The related microbial communities found in both
enrichments had some bacterial families in common, but the communities found in the PCE appeared
metabolically more diverse, suggesting probable rich microbial interactions with exchanges of electrons,
carbon and nitrogen between autotrophic, heterotrophic and mixotrophic populations. Several members
of the enriched communities were furthermore reported as plant growth promoting bacteria (PGPB)
which could be interesting for the production of environment friendly fertilizers.To summarize, a
conceptual model of microbial interactions between the main bacterial families found in the
bioelectrochemical system was proposed, suggesting a key role of each autotrophic, heterotrophic and
mixotrophic population in the process of N_2 fixation by cathodic biofilms.

619   **Acknowledgements**

620   The authors would like to thank the INRAE Bio2E Facility (Bio2E, INRAE,2018. Environmental
621   Biotechnology and Biorefinery Facility, https://doi.org/10.15454/1.557234103446854E12) for
622   experimental support.

623   **Data and scripts availability**

624   Data and scripts are available online on: https://doi.org/10.57745/ONNGWZ (Recherche Data gouv).

625   **Conflict of interest disclosure**

626   The authors declare that they comply with the PCI rule of having no financial conflicts of interest in
627   relation to the content of the article.

628   **Funding**

629   This work was funded by the French National Research Agency (ANR, ANR-19-CE43-0013 Cathomix).

630   **References**

Aullo T, Ranchou-Peyruse A, Ollivier B, Magot, M (2013) *Desulfotomaculum* spp. and related gram-
positive sulfate-reducing bacteria in deep subsurface environments. *Front. Microbiol., 4, 362*.
https://www.frontiersin.org/article/10.3389/fmicb.2013.00362
Bagali S (2012) Review: Nitrogen fixing microorganisms. *Int. J. Microbiol. Res., 3*, 46–52.
https://doi.org/10.5829/idosi.ijmr.2012.3.1.61103
Bergersen F J (1970) The quantitative relationship between nitrogen fixation and the acetylene-reduction
assay. *Aust. J. Biol. Sci., 23*(4), 1015-1026. https://doi.org/10.1071/bi9701015
Bowers TH, Reid NM, Lloyd-Jones G (2008) Composition of *nifH* in a wastewater treatment system reliant
on N_2 fixation. *Appl. Microbiol. Biotechnol., 79*(5), 811–818. https://doi.org/10.1007/s00253-008-
1486-2
Burgess BK, Lowe DJ (1996) Mechanism of molybdenum nitrogenase. *Chem. Rev., 96*(7), 2983-3012.
https://doi.org/10.1021/cr950055x
Burris RH, Roberts GP (1993) Biological nitrogen fixation. *Annu. Rev. Nutr., 13*(1), 317-335.
https://doi.org/10.1146/annurev.nu.13.070193.001533
Carmona-Martínez AA, Trably E, Milferstedt K, Lacroix R, Etcheverry L, Bernet N (2015) Long-term
continuous production of H_2 in a microbial electrolysis cell (MEC) treating saline wastewater. *Water
Res., 81*, 149–156. https://doi.org/10.1016/j.watres.2015.05.041
Cassan F, García de Salamone I (2008) *Azospirillum: Cell physiology, plant response, agronomic and
environmental research in Argentina*. ISBN 978-987-98475-8-9.
Chakraborty T, Akhtar N (2021) Biofertilizers: Prospects and Challenges for Future. In *Biofertilizers* (pp.
575–590). John Wiley & Sons, Ltd. https://doi.org/10.1002/9781119724995.ch20
Cherkasov N, Ibhadon AO, Fitzpatrick P (2015) A review of the existing and alternative methods for
greener nitrogen fixation. *Chem. Eng. Process., 90*, 24–33. https://doi.org/10.1016/j.cep.2015.02.004
Deng J, Iñiguez JA, Liu C (2018) Electrocatalytic nitrogen reduction at low temperature. *Joule, 2*(5), 846–
856. https://doi.org/10.1016/j.joule.2018.04.014
Dos Santos PC, Fang Z, Mason SW, Setubal JC, Dixon R (2012) Distribution of nitrogen fixation and
nitrogenase-like sequences amongst microbial genomes. *BMC Genomics, 13*, 162.
https://doi.org/10.1186/1471-2164-13-162
Franche C, Lindström K, Elmerich C (2009) Nitrogen-fixing bacteria associated with leguminous and non-
leguminous plants. *Plant Soil, 321*, 35–59. https://doi.org/10.1007/s11104-008-9833-8

Gaby JC, Buckley DH (2012) A comprehensive evaluation of PCR primers to amplify the *nifH* gene of
nitrogenase. *PLoS ONE, 7*(7), e42149. https://doi.org/10.1371/journal.pone.0042149
Ghodhbane-Gtari F, Nouioui I, Hezbri K, Lundstedt E, D'Angelo T, McNutt Z, Laplaze L, Gherbi H, Vaissayre
V, Svistoonoff S, ben Ahmed H, Boudabous A, Tisa LS (2019) The plant-growth-promoting
actinobacteria of the genus *Nocardia* induces root nodule formation in *Casuarina glauca. Antonie van*
Leeuwenhoek, 112, 75–90. https://doi.org/10.1007/s10482-018-1147-0
Goldman JC, Caron DA, Dennett MR (1987) Regulation of gross growth efficiency and ammonium
regeneration in bacteria by substrate C: N ratio1. *Limnol. Oceanogr., 32*(6), 1239–1252.
https://doi.org/10.4319/lo.1987.32.6.1239
Hardy RWF, Burns RC, Holsten RD (1973) Applications of the acetylene-ethylene assay for measurement
of nitrogen fixation. *Soil Biol. Biochem., 5*(1), 47–81. https://doi.org/10.1016/0038-0717(73)90093-X
Heldal M, Norland S, Tumyr O (1985). X-ray microanalytic method for measurement of dry matter and
elemental content of individual bacteria. *Appl. Environ. Microbiol., 50*(5), 1251–1257.
https://doi.org/10.1128/aem.50.5.1251-1257.1985
Heuck C, Weig A, Spohn M (2015) Soil microbial biomass C:N:P stoichiometry and microbial use of organic
phosphorus. *Soil Biol. Biochem., 85*, 119–129. https://doi.org/10.1016/j.soilbio.2015.02.029
Hu J, Tang H, Wang YZ, Yang C, Gao M, Tsang YF, Li J (2022) Effect of dissolved solids released from
biochar on soil microbial metabolism. *Environ. Sci.: Processes Impacts, 24*, 598–608.
https://doi.org/10.1039/D2EM00036A
Hu X, Kerckhof FM, Ghesquière J, Bernaerts K, Boeckx P, Clauwaert P, Boon N (2020) Microbial protein
out of thin air: fixation of nitrogen gas by an autotrophic hydrogen-oxidizing bacterial enrichment.
Environ. Sci. Technol., 54(6), 3609-3617. https://doi.org/10.1021/acs.est.9b06755
Huang X, Liu L, Zhao J, Zhang J, Cai Z (2019) The families *Ruminococcaceae, Lachnospiraceae*, and
Clostridiaceae are the dominant bacterial groups during reductive soil disinfestation with
incorporated plant residues. *Appl. Soil Ecol., 135*, 65–72. https://doi.org/10.1016/j.apsoil.2018.11.011
Huda N u, Tanvir R, Badar J, Ali I, Rehman Y (2022) Arsenic-resistant plant growth promoting
Pseudoxanthomonas mexicana S254 and *Stenotrophomonas maltophilia* S255 isolated from
agriculture soil contaminated by industrial effluent. *Sustainability, 14*(17), 10697.
https://doi.org/10.3390/su141710697
Kandemir T, Schuster ME, Senyshyn A, Behrens M, Schlögl R (2013) The Haber–Bosch process revisited:
on the real structure and stability of "ammonia iron" under working conditions. *Angew. Chem., Int.*
Ed. Engl., 52(48), 12723-12726. https://doi.org/10.1002/anie.201305812
Khan KS, Mack R, Castillo X, Kaiser M, Joergensen RG (2016) Microbial biomass, fungal and bacterial
residues, and their relationships to the soil organic matter C/N/P/S ratios. *Geoderma, 271*, 115–123.
https://doi.org/10.1016/j.geoderma.2016.02.019
Kifle MH, Laing MD (2016) Isolation and screening of bacteria for their diazotrophic potential and their
influence on growth promotion of maize seedlings in greenhouses. *Front. Plant Sci., 6*, 1225.
https://doi.org/10.3389/fpls.2015.01225
Kim J, Rees DC (1994) Nitrogenase and biological nitrogen fixation. *Biochemistry, 33*, 389-397.
https://doi.org/10.1021/bi00168a001
Klemps R, Cypionka H, Widdel F, Pfennig N (1985) Growth with hydrogen, and further physiological
characteristics of *Desulfotomaculum* species. *Arch. Microbiol., 143*, 203–208.
https://doi.org/10.1007/BF00411048
Lin L, Li Z, Hu C, Zhang X, Chang S. Yang L, Li Y, An Q (2012) Plant growth-promoting nitrogen-fixing
enterobacteria are in association with sugarcane plants growing in Guangxi, China. *Microbes Environ.,*
27(4), 391–398. https://doi.org/10.1264/jsme2.me11275
Liu A, Yang Y, Ren X, Zhao Q, Gao M, Guan W, Meng F, Gao L, Yang Q, Liang X, Ma T (2020) Current
progress of electrocatalysts for ammonia synthesis through electrochemical nitrogen reduction under
ambient conditions. *ChemSusChem, 13*(15), 3766–3788. https://doi.org/10.1002/cssc.202000487
Liu C, Sakimoto KK, Colón BC, Silver PA, Nocera DG (2017) Ambient nitrogen reduction cycle using a
hybrid inorganic–biological system. *Proc. Natl. Acad. Sci., 114*(25), 6450-6455.
https://doi.org/10.1073/pnas.1706371114

Liu F, Zhao C, Xia L, Yang F, Chang X, Wang Y (2011) Biofouling characteristics and identification of
preponderant bacteria at different nutrient levels in batch tests of a recirculating cooling water
system. *Environ. Technol.*, *32*(8), 901–910. https://doi.org/10.1080/09593330.2010.517220
Loferer-Krößbacher M, Klima J, Psenner R. (1998) Determination of bacterial cell dry mass by
transmission electron microscopy and densitometric image analysis. *Appl. Environ. Microbiol.*, *64*(2),
688–694. https://doi.org/10.1128/AEM.64.2.688-694.1998
Logan BE, Rossi R, Ragab A, Saikaly PE (2019) Electroactive microorganisms in bioelectrochemical
systems. *Nat. Rev. Microbiol.*, *17*, 307-319. https://doi.org/10.1038/s41579-019-0173-x
Martín A J, Shinagawa T, Pérez-Ramírez J. (2019) Electrocatalytic reduction of nitrogen: from Haber-Bosch
to ammonia artificial leaf. *Chem*, *5*(2), 263-283. https://doi.org/10.1016/j.chempr.2018.10.010
Masclaux-Daubresse C, Daniel-Vedele F, Dechorgnat J, Chardon F, Gaufichon L, Suzuki A. (2010) Nitrogen
uptake, assimilation and remobilization in plants: Challenges for sustainable and productive
agriculture. *Ann. Bot.*, *105*(7), 1141–1157. https://doi.org/10.1093/aob/mcq028
McMurdie PJ (2023) *Phyloseq* [R]. https://github.com/joey711/phyloseq (Original work published 2011)
Minamisawa K, Nishioka K, Miyaki T, Ye B, Miyamoto T, You M, Saito A, Saito M, Barraquio WL,
Teaumroong N, Sein T, Sato T (2004) Anaerobic nitrogen-fixing consortia consisting of *Clostridia*
isolated from gramineous plants. *Appl. Environ. Microbiol.*, *70*(5), 3096-3102.
https://doi.org/10.1128/AEM.70.5.3096-3102.2004
Moreno-Vivián C, Caballero FJ, Cárdenas J, Castillo F (1989) Effect of the C/N balance on the regulation of
nitrogen fixation in *Rhodobacter capsulatus* E1F1. *Biochim. Biophys. Acta Bioenerg.*, *977*(3), 297–300.
https://doi.org/10.1016/S0005-2728(89)80083-0
Moscoviz R, Desmond-Le Quéméner E, Trably E, Bernet N (2019) Bioelectrochemical Systems for the
Valorization of Organic Residues. In J.-R. Bastidas-Oyanedel & J. E. Schmidt (Eds.), *Biorefinery:*
Integrated Sustainable Processes for Biomass Conversion to Biomaterials, Biofuels, and Fertilizers (pp.
511–534). Springer International Publishing. https://doi.org/10.1007/978-3-030-10961-5_21
Paul D, Noori MT, Rajesh PP, Ghangrekar MM, Mitra A (2018) Modification of carbon felt anode with
graphene oxide-zeolite composite for enhancing the performance of microbial fuel cell. *Sustain.*
Energy Technol. Assess., *26*, 77–82. https://doi.org/10.1016/j.seta.2017.10.001
Peoples MB, Craswell ET (1992) Biological nitrogen fixation: Investments, expectations and actual
contributions to agriculture. *Plant Soil*, *141*(1), 13–39. https://doi.org/10.1007/BF00011308
Pogoreutz C, Rädecker N, Cárdenas A, Gärdes A, Wild C, Voolstra CR (2017) Nitrogen fixation aligns with
nifH abundance and expression in two coral trophic functional groups. *Front. Microbiol.*, *8*.
https://www.frontiersin.org/articles/10.3389/fmicb.2017.01187
Poly F, Ranjard L, Nazaret S, Gourbière F, Monrozier L J (2001) Comparison of *nifH* gene pools in soils and
soil microenvironments with contrasting properties. *Appl. Environ. Microbiol.*, *67*(5), 2255–2262.
https://doi.org/10.1128/AEM.67.5.2255-2262.2001
Postgate JR (1970) Nitrogen fixation by sporulating sulphate-reducing bacteria including rumen strains. *J.*
Gen. Microbiol, *63*(1), 137–139. https://doi.org/10.1099/00221287-63-1-137
Rago L, Zecchin S, Villa F, Goglio A, Corsini A, Cavalca L, Schievano A (2019) Bioelectrochemical nitrogen
fixation (e-BNF): electro-stimulation of enriched biofilm communities drives autotrophic nitrogen and
carbon fixation. *Bioelectrochem.*, *125*, 105–115. https://doi.org/10.1016/j.bioelechem.2018.10.002
Rojas-Tapias D, Moreno-Galván A, Pardo-Díaz S, Obando M, Rivera D, Bonilla R (2012) Effect of
inoculation with plant growth-promoting bacteria (PGPB) on amelioration of saline stress in maize
(*Zea mays*). *Appl. Soil Ecol.*, *61*, 264–272. https://doi.org/10.1016/j.apsoil.2012.01.006
Rous, A. (2023). *Script and data used for the article 'Comparison of enrichment methods for efficient*
nitrogen fixation on a biocathode' (V1 ed.). Recherche Data Gouv. https://doi.org/10.57745/ONNGWZ
Ryan RP, Monchy S, Cardinale M, Taghavi S, Crossman L, Avison MB, Berg G, van der Lelie D, Dow JM
(2009) The versatility and adaptation of bacteria from the genus *Stenotrophomonas*. *Nature Reviews*
Microbiology, *7*, 5114-525. https://doi.org/10.1038/nrmicro2163
Saiz E, Sgouridis F, Drijfhout FP, Ullah S (2019) Biological nitrogen fixation in peatlands: Comparison
between acetylene reduction assay and $^{15}N_2$ assimilation methods. *Soil Biol. Biochem.*, *131*, 157–165.
https://doi.org/10.1016/j.soilbio.2019.01.011
Singh RK, Singh P, Li HB, Guo DJ, Song QQ, Yang T, Malviya MK, Song XP, Li YR. (n.d.). Plant-PGPR
interaction study of plant growth-promoting diazotrophs *Kosakonia radicincitans* BA1

and *Stenotrophomonas maltophilia* COA2 to enhance growth and stress-related gene expression
in *Saccharum* spp. *J. Plant Interact.*,*15(1)*, 427-445. https://doi.org/10.1080/17429145.2020.1857857
Soper FM, Simon C, Jauss V (2021) Measuring nitrogen fixation by the acetylene reduction assay (ARA): Is
3 the magic ratio? *Biogeochemistry*, *152*(2), 345–351. https://doi.org/10.1007/s10533-021-00761-3
Stoddard SF, Smith BJ, Hein R, Roller BRK, Schmidt TM (2015) *rrnDB*: Improved tools for interpreting rRNA
gene abundance in bacteria and archaea and a new foundation for future development. *Nucleic Acids*
Res., *43*(D1), D593–D598. https://doi.org/10.1093/nar/gku1201
Temple SJ, Vance CP, Stephen Gantt J (1998) Glutamate synthase and nitrogen assimilation. *Trends Plant*
Sci., *3*(2), 51–56. https://doi.org/10.1016/S1360-1385(97)01159-X
Tidyverse. (n.d.). Retrieved 9 February 2023, from https://www.tidyverse.org/
Tilak KVBR, Schneider K, Schlegel HG (1986) Autotrophic growth of nitrogen-fixing *Azospirillum* species
and partial characterization of hydrogenase from strain CC. *Current Microbiol.*, *13*(6), 291–297.
https://doi.org/10.1007/BF01577194
Wiegel J (2006). The Genus *Xanthobacter* . In: Dworkin, M., Falkow, S., Rosenberg, E., Schleifer, KH.,
Stackebrandt, E. (eds) The Prokaryotes. Springer, New York, NY. https://doi.org/10.1007/0-387-
30745-1_16
Wresta A, Widyarani R, Boopathy R, Setiadi T (2021) Thermodynamic approach to estimating reactions
and stoichiometric coefficients of anaerobic glucose and hydrogen utilization. *Eng. Rep.*, *3*(6), e12347.
https://doi.org/10.1002/eng2.12347
Wu Y, Zaiden N, Cao B (2018) The core- and pan-genomic analyses of the genus *Comamonas*: from
environmental adaptation to potential virulence. *Front. Microbiol.*, *9*, 3096.
https://doi.org/10.3389/fmicb.2018.03096
Hafeez FY, Yasmin S, Ariani D, Rahman Mu, Zafar Y, Malik KA (2006). Plant growth-promoting bacteria as
biofertilizer. *Agron. Sustain. Dev.*, *26*(2), 143–150. https://doi.org/10.1051/agro:2006007
Yadav R, Chiranjeevi P, Yadav S, Singh R, Patil, SA (2022) Electricity-driven bioproduction from CO_2 and N_2
feedstocks using enriched mixed microbial culture. *J. CO_2 Util.*, *60*, 101997.
https://doi.org/10.1016/j.jcou.2022.101997
Yaghoubi Khanghahi M, Strafella S, Allegretta I, Crecchio C (2021) Isolation of bacteria with potential
plant-promoting traits and optimization of their growth conditions. *Current Microbiol.*, *78*(2), 464–
478. https://doi.org/10.1007/s00284-020-02303-w
Zaybak Z, Pisciotta JM, Tokash JC, Logan BE (2013) Enhanced start-up of anaerobic facultatively
autotrophic biocathodes in bioelectrochemical systems. *J. Biotechnol.*, *168*(4), 478–485.
https://doi.org/10.1016/j.jbiotec.2013.10.001
Zhang L, Tian C, Wang H, Gu W, Zheng D, Cui M, Wang X, He X, Zhan G, Li D (2022). Improving
electroautotrophic ammonium production from nitrogen gas by simultaneous carbon dioxide fixation
in a dual-chamber microbial electrolysis cell. *Bioelectrochem.*, *144*, 108044.
https://doi.org/10.1016/j.bioelechem.2021.108044
Zhang W, Yu C, Wang X, Hai L (2020) Increased abundance of nitrogen transforming bacteria by higher
C/N ratio reduces the total losses of N and C in chicken manure and corn stover mix composting.
Biores. Technol., *297*, 122410. https://doi.org/10.1016/j.biortech.2019.122410
Zhang Y, Hu T, Wang H, Jin H, Liu Q, Chen Z, Xie Z (2022). Nitrogen content and C/N ratio in straw are the
key to affect biological nitrogen fixation in a paddy field. *Plant Soil*, 481, 535-546.
https://doi.org/10.1007/s11104-022-05654-4