Identification of novel buffering mechanisms in aortic arch artery development and congenital

heart disease

AnnJosette Ramirez[1,2], Karl Degenhardt[4], and Sophie Astrof[1,2,3*]

1. Department of Cell Biology and Molecular Medicine, New Jersey Medical School, Rutgers

Biomedical and Health Sciences, Newark, NJ, 07103

2. Multidisciplinary Ph.D. Program in Biomedical Sciences: Cell Biology, Neuroscience and

Physiology Track, New Jersey Medical School, Rutgers Biomedical and Health Sciences,

Newark, NJ, 07103

3. Multidisciplinary Ph.D. Program in Biomedical Sciences: Molecular Biology, Genetics, and

Cancer Track, New Jersey Medical School, Rutgers Biomedical and Health Sciences, Newark,

NJ, 07103

4. Children's Hospital of Pennsylvania, University of Pennsylvania, Philadelphia, PA 19107

* Author for correspondence

Sophie Astrof (sophie.astrof@rutgers.edu)

Summary

The remarkable robustness of embryonic development to genetic and environmental perturbations has been long appreciated; however, little is known about buffering mechanisms underlying the constancy of development. Yet, aberrations in developmental programs can lead to lethal outcomes, exemplified by congenital heart disease (CHD). Defective morphogenesis of the aortic arch is a lethal CHD phenotype common in 22q11 deletion syndrome (22q11DS). Previously, we found that the second heart field (SHF) gives rise to the pharyngeal arch artery (PAA) endothelium. Here, we show that robust PAA development is dependent on both a threshold and an optimum number of SHF-derived endothelial cells (ECs) and that the optimum number is regulated by *VEGFR2* and *Tbx1* gene dosage. We discovered that when the number of SHF-derived ECs falls below the optimum but remains within the threshold, PAA development is rescued by a compensatory endothelium. To determine the source of compensating endothelium and the mechanisms regulating its recruitment, we used *in toto* confocal imaging to investigate EC connectivity, fate, and gene expression at single-cell and single-molecule resolutions. Our studies demonstrate that the loss of SHF-derived ECs leads to alterations in the pharyngeal tissue microenvironment, resulting in increased synthesis of *VEGFA*, eliciting ectopic angiogenesis from the nearby veins but not arteries. We show that this rescuing mechanism is regulated by *Tbx1*, a major 22q11DS disease gene; the absence of a single allele of *Tbx1* prevents this angiogenic response leading to unpredictable arch artery morphogenesis and lethal CHD in *Tbx1*$^{+/-}$ animals. Together, our work uncovers a novel buffering mechanism underlying the fidelity of arch artery development and provides insights into the pathogenesis of CHD.

Introduction

How complex developmental processes almost invariably lead to predictable phenotypes has remained a conundrum for many years. In 1942, C.H. Waddington proposed the concept of canalization of development, which refers to the tendency of developmental processes to follow a specific trajectory despite genetic and environmental perturbations (Waddington, 1942). Noting that genetically "wild type" organisms have much less developmental variability than organisms with genetic mutations, Waddington argued for a buffering mechanism(s) that allow constancy in developmental outcomes despite minor genetic or environmental perturbations. One example of canalized development is the organogenesis of the cardiovascular system. The aortic arch is a major vessel that circulates oxygenated blood throughout the body. In wild-type mice and humans, this vessel forms a leftward arch to connect the heart and the descending aorta. However, mutations affecting pharyngeal mesoderm or cardiac neural crest development can result in unpredictable defects in arch artery development and remodeling; the aortic arch can form a rightward arch, form two arches, or be absent completely (Stoller and Epstein, 2005). Although rare, congenital defects offer an opportunity to understand buffering mechanisms underlying the constancy of developmental outcomes and their alterations resulting in birth defects.

Congenital heart disease (CHD) is the most prevalent congenital anomaly, affecting approximately 8 per 1000 live births in the United States (Benjamin et al., 2018; Tsao et al., 2022). Defects in the development of the aorta, aortic arch arteries (AAAs), and pulmonary trunk give rise to the most severe forms of CHD. The aortic arch and AAAs are derived from three bilateral pairs of pharyngeal arch arteries (PAAs), namely the 3rd, 4th, and 6th. During development, nascent PAAs undergo asymmetric remodeling to give rise to mature AAAs (Kau et al., 2007; Stoller and Epstein, 2005). Defective formation and remodeling of the left 4th PAA leads to the lethal CHD phenotype interrupted aortic arch type B (IAA-B), resulting in the loss of luminal continuity between the ascending and descending aorta and interrupting the circulation of the oxygenated blood throughout the body (Celoria and Patton, 1959). Improper formation and remodeling of the right 4th PAA results in the non-lethal CHD phenotype retro-esophageal right subclavian artery (RERSA), where the right subclavian artery branches from the descending aorta. Although a non-lethal CHD phenotype, RERSA can lead to dysphagia lusoria (Levitt and Richter, 2007).

Chromosome 22q11 deletion syndrome (22q11 DS) is one of the most prevalent chromosomal
microdeletion syndromes, with an incidence of 1 in 3,000-6,000 live births (McDonald-McGinn
and Sullivan, 2011; McDonald-McGinn et al., 2015). Characterized by multiple anomalies, 80%
of 22q11DS cases present with congenital heart disease (McDonald-McGinn et al., 2015;
Momma, 2010). Further, approximately 50-60% of patients with interrupted aortic arch have
22q11DS, with IAA-B being the most prevalent subtype (Collins-Nakai et al., 1976; Goldmuntz et
al., 1993; Moerman et al., 1987; Van Mierop and Kutsche, 1986b). *Tbx1* is among the genes
deleted in 22q11DS, and knockout studies demonstrated that heterozygosity of *Tbx1* in mice
results in aberrant 4[th] PAA development and phenocopies 4[th] PAA-derived aortic arch defects in
22q11DS patients (Jerome and Papaioannou, 2001; Lindsay et al., 2001; Merscher et al., 2001).

The PAAs develop within pharyngeal arches, a series of bulges on the lateral surface of the
thoracic region of the embryo (Graham, 2003). PAAs form by vasculogenesis, where mesoderm-
derived vascular progenitors populate the pharyngeal arches and differentiate into endothelial
cells (ECs). These endothelial cells form a primary network of small vessels which then coalesce
into a patent artery (Li et al., 2012; Nagelberg et al., 2015; Paffett-Lugassy et al., 2013; Wang et
al., 2017; Warkala et al., 2021). We and others have previously shown that PAA endothelium,
particularly the 4[th] and 6[th] PAAs, is mainly derived from a subpopulation of splanchnic mesoderm
known as the Second Heart Field (SHF) (Paffett-Lugassy et al., 2013; Wang et al., 2017; Warkala
et al., 2021). However, the role of SHF-derived vascular progenitors in PAA morphogenesis, the
mechanisms regulating their contribution to the PAAs, and the impact of *Tbx1* haploinsufficiency
on this process are unknown.

To investigate the developmental mechanisms regulating PAA formation and remodeling, we
used transgenic mouse models to manipulate the expression of *VEGFR2* and *Tbx1*. Our findings
demonstrate that the correct *VEGFR2* gene dosage in the SHF is crucial for the contribution of
SHF-derived ECs to the PAA endothelium. We show that robust arch artery development requires
both a threshold and an optimum number of SHF-derived endothelial cells (ECs). The expression
of *VEGFR2* and *Tbx1* regulates this optimum number in a dosage-dependent manner.
Furthermore, we discovered that when the number of SHF-derived endothelium falls below the
optimum, the arch artery can be rebuilt from endothelial cells that arise from nearby veins via
angiogenesis. This angiogenic response is suppressed when the optimum number of SHF-
derived endothelial cells is present in the pharyngeal arches. However, when the number of SHF-
derived ECs falls below the optimum, the pharyngeal tissue microenvironment is altered, resulting
in increased synthesis of VEGFA in the pharyngeal arches. Of significant importance for human

disease, the buffering capacity during arch artery development is regulated by *Tbx1* in a dosage-
dependent manner, which mediates the recruitment of venous endothelium by regulating
angiogenic sprouting from veins. Altogether, this work provides insights into buffering
mechanisms underlying robustness and plasticity in the development of the aortic arch.

Results

The expression of VEGFR2 in the SHF is required for the specification of pharyngeal arch
endothelium.

To determine whether the SHF is a principal source of the PAA endothelium, we ablated VEGFR2
from the SHF using the *Isl1^{Cre}* knock-in strain by crossing *VEGFR2$^{+/-}$;Isl1$^{Cre/+}$* mice with
VEGFR2$^{f/f}$;Rosa$^{mTmG/mTmG}$ mice,   generating   *VEGFR2$^{flox/-}$;Isl1$^{Cre/+}$*   (*VEGFR2$^{Isl1-KO}$*),
VEGFR2$^{flox/+}$;Isl1$^{Cre/+}$ (*VEGFR2$^{Isl1-Het}$*), and *VEGFR2$^{flox/+}$;Isl1$^{+/+}$* (*VEGFR2$^{Isl1-WT}$*) embryos. Embryos
were examined at embryonic day (E) 10.5 when PAA formation was nearly complete (34 – 36s)
(Anderson and Bamforth, 2022; Wang et al., 2017; Warkala et al., 2021). Vasculature was
visualized using whole-mount immunofluorescent staining (WM-IF) and antibodies against
VEGFR2, followed by confocal imaging of the entire pharyngeal region and three-dimensional
(3D) reconstruction of optical sections. We hypothesized that the deletion of VEGFR2 from the
SHF would result in the absence of PAA endothelium and that no PAAs would form. Our analyses
showed   that   PAA   formation   in   *VEGFR2$^{Isl1-Het}$*   and   *VEGFR2$^{Isl1-KO}$*   embryos   was   defective,
compared to *VEGFR2$^{Isl1-WT}$* (**Fig. S1**); the 6th PAA of *VEGFR2$^{Isl1-Het}$* embryos was hypoplastic
(compare **Fig. S1A-B** with **Fig. S1C-D**), whereas the 4th and 6th PAAs of *VEGFR2$^{Isl1-KO}$* embryos
were   either   hypoplastic   or   aplastic   (**Fig. S1E-F, Table S1**).   Surprisingly,   despite   the
downregulation of VEGFR2 from the SHF (**Fig. S1G-I, outlined**), VEGFR2+ endothelium was
present in all three pharyngeal arches of *VEGFR2$^{Isl1-KO}$* embryos (**Fig. S1E-F**). These results
suggest that either VEGFR2 expression is not essential for the differentiation of SHF-derived
vascular progenitors into PAA ECs or that the endothelium in the mutant embryos was derived
from a source other than the SHF.

To test these hypotheses and determine the fate of SHF-derived mesoderm in mutant embryos,
we used the *mef2c-AHF-Cre* and *Rosa^{tdTomato}* strains. For these experiments, *VEGFR2$^{+/-}$;mef2c-*
AHF-Cre mice were crossed with *VEGFR2$^{flox/flox}$;Rosa$^{tdTom/tdTom}$* mice to generate *VEGFR2$^{flox/-}$*
;mef2c-AHF-Cre+;Rosa$^{tdTom/+}$ (*VEGFR2^{SHF-KO}*)   embryos   and   *VEGFR2$^{flox/+}$;mef2c-AHF-*
Cre+;Rosa$^{tdTom/+}$ (*VEGFR2$^{SHF-Het}$*) embryos. *VEGFR2$^{+/-}$;mef2c-AHF-Cre* mice were also crossed

with $Rosa^{tdTom/tdTom}$ mice to generate $VEGFR2^{+/+};mef2c\text{-}AHF\text{-}Cre+;Rosa^{tdTom/+}$ ($VEGFR2^{SHF\text{-}WT}$)

embryos. To verify the ablation of VEGFR2 in the *mef2c-AHF*-lineage, we performed WM-IF and

used antibodies against VEGFR2 and tdTomato, encoded by the Cre-Reporter. In $VEGFR2^{SHF\text{-}}$

^{WT} embryos, Cre-reporter+ cells in the anterior and posterior poles of the dorsal pericardial wall

and pharyngeal arches expressed VEGFR2 (**Fig. 1A-I**). In $VEGFR2^{SHF\text{-}KO}$ embryos, VEGFR2

expression was absent in the dorsal pericardial wall (**Fig. 1J-N, outlined**), and >80% of Cre-

reporter+ cells in the pharyngeal arches did not express VEGFR2 (**Figs. 1O-R arrows, Table**

S2). These experiments demonstrated the successful ablation of VEGFR2 protein from the

mef2c-AHF-lineage in $VEGFR2^{SHF\text{-}KO}$ embryos.

To determine whether the expression of VEGFR2 in the *mef2c-AHF*-lineage was required for PAA

development, we examined the formation of the PAAs in $VEGFR2^{SHF\text{-}WT}$, $VEGFR2^{SHF\text{-}Het}$ and

$VEGFR2^{SHF\text{-}KO}$ embryos at E10.5. Embryos were stained by WM-IF using antibodies against

VEGFR2, the ETS transcription factor ERG, and the Cre-reporter tdTomato. As with the $Isl1^{Cre/+}$

strain, we observed defective PAA formation in $VEGFR2^{SHF\text{-}Het}$ and $VEGFR2^{SHF\text{-}KO}$ embryos (**Fig.**

2A-F, Table S1). $VEGFR2^{SHF\text{-}Het}$ embryos had hypoplastic 6th PAAs (**Fig. 2C-D**). PAA defects in

$VEGFR2^{SHF\text{-}KO}$ mutants were more severe than in the $VEGFR2^{SHF\text{-}Het}$: both 4th and 6th PAAs were

mostly aplastic, with pharyngeal endothelium remaining in the primordial plexus form without

forming a central artery (**Fig. 2E-F**).

Quantitative analyses showed that the number of SHF-derived cells and the number of endothelial

cells were similar across all the genotypes (**Fig. 2G-H, Table S2-3**). However, the proportion of

pharyngeal arch ECs derived from the SHF was decreased in $VEGFR2^{SHF\text{-}Het}$ and $VEGFR2^{SHF\text{-}KO}$

embryos (**Fig. 2I, Table S3**). This indicates that VEGFR2 is not required for the presence of SHF-

derived cells in the pharyngeal arch mesenchyme but is necessary for their differentiation into

ECs. Since we observed similar phenotypes when ablating VEGFR2 using either the $Isl1^{Cre}$ or

mef2c-AHF-Cre strains, these experiments indicate that the expression of *VEGFR2* in the SHF,

the common expression domain of the Cre recombinase in these strains, regulates the

differentiation of vascular progenitors.

Compared with controls, the proportion of SHF-derived ECs in $VEGFR2^{SHF\text{-}Het}$ was reduced about

2-fold in all pharyngeal arches (**Fig. 2I, Table S3**). These data indicate that the expression of one

copy of VEGFR2 was not sufficient for SHF-derived cells to adopt EC fate. In $VEGFR2^{SHF\text{-}KO}$, the

majority of pharyngeal arch ECs were not derived from the *mef2c-AHF* lineage; the proportions

of SHF-derived ECs were decreased by 3-, 10-, and 20-fold relative to controls in the 3rd, 4th, and

6th pharyngeal arches, respectively **(Fig. 2I, Table S3)**. Together, these results reveal that an

alternative vascular source compensates for the reduction in SHF-derived ECs in $VEGFR2^{SHF\text{-}Het}$

and $VEGFR2^{SHF\text{-}KO}$ embryos and rescues the numbers of ECs in pharyngeal arches to control

levels.

Compensatory endothelium rescues arch artery morphogenesis in $VEGFR2^{SHF\text{-}Het}$

embryos.

Defective PAA formation leads to abnormal morphogenesis of the aortic arch and its branches

(Hutson and Kirby, 2007). To determine the role of SHF-derived endothelium in the remodeling

of PAAs into the aortic arch arteries (AAA), we examined embryos heterozygous or null for

VEGFR2 in either the *mef2c-AHF-Cre* or $Isl1^{Cre}$ lineages after E14.5, when the remodeling of the

AAAs is complete. A Mendelian distribution of each genotype was observed between E14.5 and

P0, indicating that loss of SHF-derived vascular progenitors does not cause prenatal lethality

(**Table S4**). To identify defects in AAA development, embryos were analyzed at either E14.5 or

E18.5 by micro-computed tomography or conventional histology and H&E staining. 80% of

embryos null for VEGFR2 in the SHF from either the $Isl1^{Cre}$ or *mef2c-AHF-Cre* lines developed

AAA defects (**Fig. 3; Table S5**). Despite a significantly decreased number of SHF-derived ECs in

the PAAs, only 1 out of 18 (5.6%) embryos heterozygous for VEGFR2 in the SHF had aberrant

AAA patterning, displaying RERSA (**Fig. 3A, Fig. S2, Table S6**). The AAA defects observed in

80% of conditional VEGFR2-null and 5.6% of conditional VEGFR2-heterozygous embryos

(**Tables S5-6**) were due to defective development and remodeling of the 4th PAAs (Stoller and

Epstein, 2005). This indicates that relative to the vessels arising from the 3rd and 6th PAAs, the

development of 4th PAA-derived vessels is more sensitive to the decrease in the numbers of SHF-

derived ECs.

Although $VEGFR2^{SHF\text{-}Het}$ and $VEGFR2^{SHF\text{-}KO}$ embryos had similar total numbers of ECs in each

pharyngeal arch (**Fig. 2H**), $VEGFR2^{SHF\text{-}KO}$ embryos displayed more severe defects in PAA

formation and subsequent AAA remodeling (**Fig. 3A, Table S5-6**). A notable difference between

these genotypic classes is the significant decrease in the proportion of SHF-derived cells in the

pharyngeal arches of $VEGFR2^{SHF\text{-}KO}$ embryos (**Fig. 2I, black bars**). These findings indicate that

SHF-derived ECs play a crucial role in PAA development and that a threshold number of SHF-

derived ECs is necessary for proper AAA morphogenesis.

Tbx1 regulates the recruitment of compensatory endothelium and the rescue of PAA development.

Fourth PAA-derived defects, such as IAA-B, are observed in about 19% of patients with 22q11DS (Momma, 2010; Yagi et al., 2003). These patients have one functional copy of the _Tbx1_ gene. Meta-analyses of studies on different _Tbx1_$^{+/-}$ strains and df(16)1/+ mice, which model 22q11DS, indicate that _Tbx1_ heterozygosity results in defects in 4th PAA formation with an average penetrance of 43% and defects in AAA development with an average penetrance of 36% (**Table S7 – 8**) (Aggarwal et al., 2006; Calmont et al., 2009; Guris et al., 2006; Hasten and Morrow, 2019; Jerome and Papaioannou, 2001; Lindsay et al., 1999; Lindsay and Baldini, 2001; Lindsay et al., 2001; Merscher et al., 2001; Papangeli and Scambler, 2013; Phillips et al., 2019; Randall et al., 2009; Ryckebusch et al., 2010; Stothard et al., 2020; Taddei et al., 2001; Vitelli et al., 2002; Zhang and Baldini, 2008).

To investigate the role of _Tbx1_ in PAA development and in the contribution of SHF-derived progenitors to the PAA endothelium, we crossed _Tbx1_$^{+/-}$;_mef2c-AHF-Cre_ mice with _Rosa_$^{tdTom/tdTom}$ mice. _Tbx1_$^{+/-}$ (_Tbx1_$^{+/-}$;_mef2c-AHF-Cre)_ and control (_mef2c-AHF-Cre_) embryos were collected at E10.5, and the total EC number and the number of SHF-derived ECs were quantified in the 4th pharyngeal arches. We found that the 4th PAAs in _Tbx1_$^{+/-}$ embryos were either hypoplastic or aplastic (**Fig. 4A-B, Table S1**). Quantitative analyses of EC populations indicated that unlike in _VEGFR2_$^{SHF-Het}$ embryos, the total number of ECs in the 4th pharyngeal arches of _Tbx1_$^{+/-}$ mice was reduced by 30% relative to controls (p < 0.0001, **Fig. 4C**). Our analyses of EC numbers indicated that the average number of total ECs in the 4th pharyngeal arch was constant across five different control genotypes examined in our lab in the past decade and is independent of the investigator performing the analyses (**Fig. S3**) (Warkala et al., 2021). We also found that a 25-30% decrease in the total number of ECs in the 4th pharyngeal arch, as seen in _Tbx1_$^{+/-}$ mice, leads to defective formation of the 4th PAAs, resulting in IAA-B or RERSA (Warkala et al., 2021). Together, these data are consistent with the idea that the ~476+/-81 ECs is the optimum EC number required for the robust formation and remodeling of the 4th PAAs.

Although the number of SHF-derived ECs was similar between _Tbx1_$^{+/-}$ and _VEGFR2_$^{SHF-Het}$ embryos (**Fig. 4D**), the decrease in the SHF-derived EC population in _VEGFR2_$^{SHF-Het}$ was complemented by the increase in the number of non-SHF derived ECs (**Fig. 4E, Table S3**). In contrast, no such increase in non-SHF-derived ECs was observed in _Tbx1_$^{+/-}$ embryos (**Fig. 4E**).

Despite having a comparable number of SHF-derived ECs between $Tbx1^{+/-}$ and $VEGFR2^{SHF-Het}$

embryos, $Tbx1^{+/-}$ embryos develop AAA defects at a much higher penetrance (36% on average,

Table S8). These data indicate the recruitment of compensatory ECs rescues PAA and AAA

morphogenesis in $VEGFR2^{SHF-Het}$ embryos (**Tables S1** and **S6**). Therefore, it is crucial to

understand the source(s) of the compensatory ECs and the mechanisms that regulate their

recruitment.

Ectopic sprouts in the pharyngeal region of $VEGFR2^{SHF-KO}$ embryos are absent in $Tbx1^{+/-}$

embryos.

To determine the origin of compensatory endothelium in $VEGFR2^{SHF-KO}$ embryos, we analyzed

the 3D connectivity of vessels surrounding the pharyngeal arches in confocal images of E9.5

$VEGFR2^{SHF-WT}$ and $VEGFR2^{SHF-KO}$ embryos (27 – 29s). Located laterally to the PAAs is a network

of venous vasculature, including the cardinal vein (CV) and pharyngeal vein (PV, also known as

the ventral vein) (**Fig. 5A**). In contrast to the $VEGFR2^{SHF-WT}$ embryos (**Fig. 5B**), we observed

ectopic vascular sprouts from CV and PV connecting with the pharyngeal arch endothelium in

$VEGFR2^{SHF-KO}$ embryos, indicating a possible source of compensatory ECs (**Fig. 5C, yellow**

bracket, and arrows). These aberrant vascular connections were absent in $Tbx1^{+/-}$ embryos,

supporting the role of $Tbx1$ in the recruitment of the rescuing endothelium (**Fig. 5D, red**

asterisks).

Angiogenic sprouting from veins gives rise to the compensatory endothelium rescuing

PAA morphogenesis.

To investigate whether non-SHF-derived ECs in $VEGFR2^{SHF-KO}$ pharyngeal arches originate from

the surrounding veins, we used lineage tracing to label venous endothelium in $VEGFR2^{SHF-KO}$

embryos. First, we employed the $ApjCreERT2$ transgenic strain (Chen et al., 2014) and tamoxifen

to induce Cre-mediated recombination at E9.5 and observed extensive labeling of venous

vasculature, including the PV, CV, and venous intersomitic branches in wild-type embryos (**Fig.**

S4A); arterial structures, including the dorsal aorta, arterial intersomitic branches, and PAAs were

not labeled except for one or two cells (**Fig. S4B**), indicating specificity for venous lineage tracing.

To label venous vasculature in $VEGFR2^{SHF-KO}$ embryos, we generated the $Apj^{DreERT2/+}$ knock-in

strain and obtained $VEGFR2^{SHF-WT};Apj^{DreERT2/+}$ or $VEGFR2^{SHF-KO};Apj^{DreERT2/+}$ embryos. Lineage

tracing of the venous vasculature revealed a significant increase in Apj-derived ECs in the 4th PAA

of $VEGFR2^{SHF-KO}$ compared to $VEGFR2^{SHF-WT}$ embryos (**Fig. 6A-E**). Importantly, we did not

observe any labeling of the dorsal aorta or its angiogenic sprouts, indicating high specificity of
venous lineage labeling (**Fig. 6B and D, white arrows**). Immunofluorescence staining and single-
molecule (sm) RNA-FISH experiments confirmed an increase in the number of ECs expressing
venous markers COUP-TFII and *Apj* in the 4th pharyngeal arches of *VEGFR2^{SHF-KO}* embryos (**Fig.**
6F-K). Our results indicate that angiogenesis from the venous endothelium generates
compensatory ECs, restoring the total number of pharyngeal arch ECs in *VEGFR2^{SHF-KO}* embryos
to control levels.

Increased *VEGFA* mRNA expression drives ectopic angiogenesis from veins.

Next, we aimed to determine mechanisms regulating the recruitment of compensatory venous
endothelium. We hypothesized that the loss of VEGFR2+ SHF-derived vascular progenitors in
pharyngeal arches would result in increased expression of proangiogenic factors, such as
VEGFA. To test this, we performed smRNA-FISH to examine *VEGFA* mRNA at early E9.5 before
the 4th PAA forms. Quantification showed a significant increase of *VEGFA* mRNA levels in the
SHF and pharyngeal endoderm of *VEGFR2^{SHF-KO}* embryos, compared to *VEGFR2^{SHF-WT}* (**Fig. 7**).
These results suggest that the loss of VEGFR2 in the SHF and the consequent inability of SHF-
derived ECs to contribute to the PAAs results in an increased expression of *VEGFA*, which in turn
drives ectopic angiogenesis from the surrounding venous vasculature to rescue pharyngeal arch
artery development.

Discussion

Proper development and remodeling of the PAAs, particularly the left 4th PAA, is crucial for
establishing appropriate blood circulation after birth. Elucidating the mechanisms underlying arch
artery formation and the response of the pharyngeal arch microenvironment to genetic
perturbations is essential for understanding how genetic mutations lead to CHD phenotypes, such
as IAA-B. Through our investigations, we have identified several critical factors that contribute to
proper arch artery morphogenesis and uncovered a previously unknown ability of the developing
mammalian embryo to restore and regenerate the PAA endothelium.

The phenomenon of development remaining relatively constant despite genetic and
environmental disturbances has captivated and confounded researchers. In 1942, Waddington
proposed the concept of buffering to explain this remarkable characteristic of multicellular
organisms. He further hypothesized that this buffering capacity is based on an optimum and a

threshold stimuli, beyond which the optimal response can no longer be achieved (Waddington,
1942). However, despite a few notable studies (Abrams et al., 2015; Jelier et al., 2016; Sharma
et al., 2017), our understanding of buffering processes and their underlying mechanisms still
needs to be improved.

Our study uncovers a heretofore unknown capacity of the mammalian embryo to repair and
regenerate the aortic arch artery development against genetic perturbations. Our findings suggest
that there exists both a threshold and an optimum number of SHF-derived ECs needed for robust
arch artery development. The expression of *VEGFR2* and *Tbx1* regulates this optimum number
in a dosage-dependent manner. Our studies suggest that crosstalk between SHF-derived ECs
and the surrounding pharyngeal tissues regulates the addition of compensating endothelium to
the PAAs. In the presence of the optimal number of SHF-derived ECs, ectopic angiogenesis is
suppressed. We hypothesize that there may be an evolutionary advantage for the formation of
the PAAs from progenitors in the SHF. However, when the number of SHF-derived ECs falls
below the optimum, the pharyngeal tissue microenvironment is modified, resulting in increased
synthesis of *VEGFA*, eliciting an angiogenic response from the veins. 3D vessel tracing, gene
expression, and lineage mapping indicate that the angiogenic response from the venous
endothelium provides compensatory endothelium rebuilding the PAAs. Our studies also reveal
the existence of a threshold number of SHF-derived ECs, below which arch artery development
becomes unpredictable, as in $VEGFR2^{SHF-KO}$ embryos.

Most patients with 4th PAA-derived CHDs, including IAA-B, are diagnosed with 22q11DS
(Moerman et al., 1987; Van Mierop and Kutsche, 1986a). Studies using $Tbx1^{+/-}$ mice have
demonstrated that these defects are attributed to haploinsufficiency in the *Tbx1* gene. The
expression of *Tbx1* in splanchnic mesoderm and pharyngeal epithelia is required for PAA
development (Jerome and Papaioannou, 2001; Lindsay et al., 2001; Merscher et al., 2001; Zhang,
2006; Zhang et al., 2005). Our research shows that *Tbx1* is a critical regulator of the contribution
of SHF-derived progenitors to the PAA endothelium and that the heterozygosity of *Tbx1* results
in a penetrant phenotype consisting of aplastic and hypoplastic PAAs. Interestingly, the number
of SHF-derived ECs in $Tbx1^{+/-}$ embryos is comparable to that in $VEGFR2^{SHF-Het}$ embryos; yet
$Tbx1^{+/-}$ embryos lack the compensating endothelium and develop AAA defects with a penetrance
of approximately 36% (**Table S8**). In contrast, the incidence of AAA defects in $VEGFR2^{SHF-Het}$
mice, where compensatory endothelium is recruited, is much lower at approximately 6%.
Altogether, these studies indicate that the threshold number of SHF-derived endothelial cells is

insufficient to support proper PAA formation and aortic arch morphogenesis in the absence of the
compensatory endothelium.

Our studies suggest that the absence of compensatory endothelium in *Tbx1*[+/-] embryos impacts
the ability of aplastic PAAs to recover and properly remodel into the AAAs. Thus, *Tbx1* regulates
the buffering capacity during arch artery development in a dosage-dependent manner. Our
experiments show that *Tbx1* mediates the recruitment of the venous endothelium by regulating
angiogenic sprouting. Taken together, these findings uncover the buffering mechanism underlying
robustness and plasticity in the development of the aortic arch. Further understanding of how
Tbx1 impacts the rescue of the PAA endothelium can lead to the prevention or treatment of CHD
phenotypes associated with 22q11DS.

Methods: Detailed methods are provided in a separate file

Author contributions: AJR and SA conceived and designed the experiments; AJR performed
all the experiments, KD processed embryos for microCT imaging, and helped with the analyses
and interpretation of congenital heart defects. AJR and SA analyzed the data and wrote the paper.

Acknowledgments: We thank Sam Russo and Bogdana Mimic for their technical assistance. Xia
Wang, for teaching all the experimental techniques. Caolán O'Donnell, Brianna Alexander, Cecilia
Arriagada, Gideon Obeng, Michael Warkala, and Huaning Zhao for reading the paper and for their
helpful suggestions throughout this work. Brittany Ramirez for her artistic expertise. This work
was supported by funding from the National Heart, Lung, and Blood Institute of the National
Institutes of Health grants R01 HL103920, R01 HL134935, and R01 HL158049, and a
Transformational Project Award from the American Heart Association 20TPA35490074 to S.A.
AJR was supported by National Heart, Lung, and Blood Institute of the National Institutes of Health
pre-doctoral fellowships F31 HL150949, HL103920-08S1, and by the National Institute of Arthritis
and Musculoskeletal and Skin Diseases training grant T32052273-11.

Figure Legends

Main Figures

Figure 1. Efficient ablation of VEGFR2 in the second heart field (SHF) using *mef2c-AHF-
***Cre* mice.** Cre-lox recombination and immunofluorescent staining were used to trace *mef2c-AHF-*
Cre lineage cells (cyan) when VEGFR2 (magenta) is conditionally ablated from the SHF in

embryos. **(A-E)** Optical confocal sections of E9.5 *VEGFR2^{SHF-WT}* embryos show VEGFR2+ cells

in the anterior **(B-C)** and posterior **(D-E)** dorsal pericardial wall outlined in yellow. **(F-I)** PAA ECs

in E10.5 *VEGFR2^{SHF-WT}* embryos are derived from the SHF. The box in F is expanded in **(G-I)**.

Arrowheads point to the co-expression of VEGFR2 (magenta), ERG (orange), and Cre-reporter

(cyan) in the 4th pharyngeal arch endothelium. **(J-N)** Optical sections of E9.5 *VEGFR2^{SHF-KO}*

embryos show the absence of VEGFR2 expression in the anterior **(K-L)** and posterior SHF dorsal

pericardial wall **(M-N),** outlined in yellow. **(O-R)** PAA ECs in E10.5 *VEGFR2^{SHF-KO}* embryos are

derived from a non-SHF endothelium. The box in **O** is expanded in **(P-R).** Arrowheads point to

Cre-reporter negative endothelium. Arrows point to Cre-reporter+ VEGFR2-negative cells.

Figure 2. VEGFR2 is required for SHF vascular progenitor contribution and PAA
development.

(A-F) WM-IF for VEGFR2 (magenta) and Cre-reporter (cyan), confocal imaging, and 3D

reconstruction of E10.5 **(A)** *VEGFR2^{SHF-WT}*, **(C)** *VEGFR2$^{SHF-Het}$*, and **(E)** *VEGFR2^{SHF-KO}* embryos.

Yellow brackets highlight aberrant PAAs in **C, E**. Dashed lines mark the levels of coronal optical

sections through the pharyngeal arches shown in **B, D,** and **F**. Dotted circles mark aberrant PAAs.

(G) Quantification of SHF-derived cells in each pharyngeal arch. *VEGFR2^{SHF-WT}* n = 10 arches,

VEGFR2$^{SHF-Het}$ n = 4 arches, *VEGFR2^{SHF-KO}* n = 6 arches. Each dot marks one pharyngeal arch.

Horizontal red and black lines mark means and quartiles, respectively. **(H)** Quantification of total

EC numbers in each pharyngeal arch. *VEGFR2^{SHF-WT}* n = 10 arches, *VEGFR2$^{SHF-Het}$* n = 12 arches,

VEGFR2^{SHF-KO} n = 12 arches. Each dot marks one pharyngeal arch. Horizontal red and black lines

mark means and quartiles, respectively. **(I)** The proportion of Cre-reporter-positive ECs and Cre-

reporter-negative ECs in each pharyngeal arch. Means ± SD are shown. *VEGFR2^{SHF-WT}* n = 10

arches, *VEGFR2$^{SHF-Het}$* n = 12 arches, *VEGFR2^{SHF-KO}* n = 12 arches. Data were analyzed using

two-way ANOVA with Tukey's correction for multiple testing; *$p \leq 0.05$ **$p \leq 0.01$, ***$p \leq 0.001$,

****$p < 0.0001$

Figure 3. A threshold number of SHF-derived ECs is required for proper Aortic Arch Artery
development.

(A) Percent of *VEGFR2;Isl1^{Cre}* (*VEGFR2$^{Isl1-WT}$*, *VEGFR2 $^{Isl1-Het}$*, and *VEGFR2 $^{Isl1-KO}$*) and -

VEGFR2;mef2c-AHF-Cre (*VEGFR2^{SHF-WT}*, *VEGFR2$^{SHF-Het}$*, and *VEGFR2^{SHF-KO}*) embryos without

AAA defects (white), non-lethal AAA defects (grey), or lethal AAA defects (black). n = the number

of embryos with each phenotype. **(B-D)** Transverse sections of E18.5 embryos stained with

Hematoxylin and eosin. **(B)** *VEGFR2^{SHF-WT}* embryos. **(C)** IAAB and **(D)** RERSA in *VEGFR2$^{Isl1-KO}$*

and *VEGFR2^{SHF-KO}* embryos, respectively. **(E-G)** Micro-computed tomography of E18.5

VEGFR2$^{Isl1-WT}$ **(E)** and *VEGFR2$^{Isl1-KO}$* **(F-G)** embryos. Arrows and text indicate examples of IAAB

(F) and aberrant RScA **(G)** in yellow. Dotted boxes highlight the position of a normal aortic arch.

Abbreviations: RCA – right carotid artery; LCA – left carotid artery; RScA – right subclavian

artery; LScA – left subclavian artery; BrA – brachiocephalic artery; AoA – aortic arch; PT –

pulmonary trunk; DA – ductus arteriosus; dAo – descending aorta; TR – trachea; IAA-B –

interrupted aortic arch type B; RERSA – retroesophageal right subclavian artery

Figure 4. *Tbx1* regulates endothelial compensation and rescue of PAA development.

(A-B) WM-IF for VEGFR2 (magenta) and Cre-reporter (cyan) of an E10.5 *Tbx1$^{+/-}$* embryo. **(A)** 3D

reconstruction and **(B)** coronal optical sections show the incomplete formation of the 4th PAA

(dotted circles) and that only a few non-SHF-derived ECs are present. **(C)** Quantification of ECs

in the 4th pharyngeal arches of control (*mef2c-AHF-Cre*), *Tbx1$^{+/-}$*, and *VEGFR2$^{SHF-Het}$* embryos.

Each dot marks one pharyngeal arch. Horizontal red and black lines mark means and quartiles,

respectively. **(D)** Quantification of the number of SHF-derived (Cre-reporter positive) ECs in the

4th pharyngeal arch. Error bars represent mean ± SD. **(E)** Quantification of the number of non-

SHF-derived (Cre-reporter negative) ECs in the 4th pharyngeal arch. Error bars represent mean

± SD. Control n = 14 arches, *Tbx1$^{+/-}$;mef2c-AHF-Cre* n = 6 arches, SHF-het n = 12 arches. Data

were analyzed using one-way ANOVA with Tukey's correction for multiple testing; **$p \leq 0.01$,

****$p < 0.0001$

Figure 5. Ectopic sprouts from veins in the pharyngeal region of *VEGFR2^{SHF-KO}* are absent

in *Tbx1$^{+/-}$* embryos

(A) A schematic depiction of venous and arterial vasculature in the pharyngeal region of E9.5

embryos. dAo – dorsal aorta; CV – cardinal vein; PV – pharyngeal vein. **(B-D)** 20μm-thick serial

sagittal optical sections through E9.5 *VEGFR2^{SHF-WT}* **(B)**, *VEGFR2^{SHF-KO}* **(C)**, and *Tbx1$^{+/-}$* **(D)**

embryos from the lateral to the medial side. Yellow brackets in B-D mark pharyngeal endothelial

plexus. Red stars mark absent venous sprouts in *VEGFR2^{SHF-WT}* **(B)** and *Tbx1$^{+/-}$* **(D)** embryos.

Yellow, magenta, and cyan arrows follow sprouts from the CV to the pharyngeal arch vasculature

in the *VEGFR2^{SHF-KO}* embryos **(C)**.

Figure 6. Angiogenic sprouting from veins gives rise to the compensatory endothelium that rescues PAA morphogenesis.

Dre-rox recombination was used to fate map cells of Apj-lineage in *VEGFR2^{SHF-KO}* and *VEGFR2^{SHF-WT}* embryos. Tamoxifen was injected at E9.5, and embryos were dissected at E10.5. **(A-D)** WM-IF staining detecting Dre-reporter (cyan) and VEGFR2 (magenta). 50μm-thick sagittal optical sections show *Apj^{DreERT2}* lineage labeling in the CV and intersomitic veins (yellow arrows) but not the dAo or intersomitic arteries (white arrows) of **(A)** *VEGFR2^{SHF-WT}* and **(C)** *VEGFR2^{SHF-KO}* embryos. Boxes in **B, D** are expanded in the panels on the right to show *Apj^{DreERT2}* lineage labeled ECs in the 4^{th} PAAs of *VEGFR2^{SHF-KO}* embryos **(**arrowheads, **D)** but not in control, *VEGFR2^{SHF-WT}* embryos **(B)**. **(E)** *VEGFR2^{SHF-KO}* embryos have a higher percent contribution of Apj-lineage cells in the 4^{th} PAAs. The number of Apj-lineage cells in arches was normalized by *Apj^{DreERT2}* labeling efficiency (see Methods and **Fig. S4C-D**). Each dot marks one pharyngeal arch. Horizontal red and black lines mark means and quartiles, respectively. Control n = 10 arches, mutant n = 6 arches. **(F-G)** WM-IF for COUPTF2 (magenta) and VEGFR2 (white) of E9.5 embryos. 20μm-thick sagittal optical sections: **F.** *VEGFR2^{SHF-WT}* and **G.** *VEGFR2^{SHF-KO}* embryos. Yellow arrows point to COUPTF2+ ECs in the CV. White arrows point to COUPTF2-negative ECs in 3^{rd} PAA and dAo. Arrowheads point to COUPTF2+ ECs in the 4^{th} PAA of *VEGFR2^{SHF-KO}* embryo **(G)**. **(H)** Quantification of COUPTF2+ ECs in the 4^{th} pharyngeal arches. Each dot marks one pharyngeal arch. Horizontal red and black lines mark means and quartiles, respectively. Control n = 9 arches, mutant n = 7 arches. **(I-J)** Confocal optical sections. Single-molecule RNA fluorescence *in situ* hybridization for *Apj* mRNA (magenta) combined with immunostaining for VEGFR2 (white) of E10.0 embryos. Yellow arrows point to *Apj*+ ECs in the CV. White arrows point to Apj-negative ECs in 3^{rd} PAA and dAo. **(K)** Quantification of *Apj*+ ECs in the 4^{th} pharyngeal arches. Each dot marks one pharyngeal arch. Horizontal red and black lines mark means and quartiles, respectively. Control n = 6 arches, mutant n = 7 arches. Data were analyzed using 2-tailed, unpaired Student's t-test; *p ≤ 0.05, ** p ≤ 0.01, ***p ≤ 0.001.

Figure 7. Loss of SHF-derived vascular progenitors increases *VEGFA* mRNA expression in the SHF and pharyngeal endoderm.

(A-B) Single-molecule RNA fluorescent *in situ* hybridization of E9.5 embryos detecting the expression of *VEGFA* (white) in the SHF (outlined in red) and pharyngeal endoderm (outlined in yellow). **(C)** Super plot of normalized mean fluorescence intensity of *VEGFA* mRNA in the SHF. **(D)** Super plot of normalized mean fluorescence intensity of *VEGFA* mRNA in the pharyngeal

endoderm. The mean fluorescence intensity in the SHF and pharyngeal endoderm was
normalized to the mean fluorescence intensity in the dorsal side of the embryo on the same optical
section (outlined in green). Different colors in **C, D** mark different biological samples. Light-colored
dots mark individual measurements in optical sections. Dark-colored dots mark the average of
measurements per pharyngeal arch. Red lines mark means, and black lines mark quartiles**.**
Control n = 6 arches, mutant n = 6 arches. Data were analyzed using 2-tailed, unpaired Student's
t-test; **p ≤ 0.01.

Supplemental Figures

Supplemental Figure 1. VEGFR2 is required in the *Isl1*-lineage for PAA development.

(A-F) WM-IF to detect VEGFR2 (magenta), confocal imaging, and 3D reconstruction of **(A)**
VEGFR2$^{Isl1-WT}$, **(C)** *VEGFR2$^{Isl1-Het}$*, and **(E)** *VEGFR2$^{Isl1-KO}$* embryos isolated at E10.5. Dashed lines
mark levels of coronal optical sections shown in **B, D, F**. Yellow bracket highlights aberrant PAAs
in **E**. Dotted circles mark aberrant PAAs in **D, F**. **G – I.** WM-IF on E10.5 embryos to detect
VEGFR2 (magenta) and lineage (cyan). Dotted line outlines the dorsal pericardial wall. Genotypes
are marked at the top, and somite (s) stages are included in each panel.

Supplemental Figure 2.

(A) Transverse sections of E18.5 *VEGFR2$^{SHF-Het}$* embryo showing RERSA (arrow) stained with
hematoxylin and eosin.

Supplemental Figure 3.

(A) Meta-analysis of total 4th pharyngeal arch EC numbers across five control genotypes:
VEGFR2$^{+/+}$;mef2c-AHF-Cre, *Tbx1$^{+/-}$;mef2c-AHF-Cre* (this manuscript*)*, *α5$^{f/+}$;Isl1$^{Cre/+}$*;
Fn1$^{f/+}$;Isl1$^{Cre/+}$ and wild-type (*C57BL6/J*) (analyses in Warkala et al., 2021 were performed by three
different investigators independently over course of eight years). Red lines mark means, and
black lines mark quartiles. Data were analyzed using a 2-tailed, unpaired Student's t-test; p>0.05
was considered not significant.

Supplemental Figure 4.

(A-B) WM-IF staining detecting Cre-reporter (cyan) and VEGFR2 (magenta). 50μm-thick sagittal optical sections show *ApjCreERT2* lineage labeling in the CV and intersomitic veins (yellow arrows) but not the dAo or intersomitic arteries (white arrows). Few Cre-reporter cells can be seen in the PAA endothelium. PAAs are numbered. **(C)** WM-IF staining detecting Dre-reporter (cyan), ERG (magenta), and VEGFR2 (grey). 100μm-thick sagittal optical section shows the region of the CV used to calculate Dre-recombination efficiency (outlined in yellow) in $Apj^{DreERT2}$ embryos. **(D)** Comparable Dre-recombination efficiency in $VEGFR2^{SHF-WT}$; $Apj^{DreERT2}$ and $VEGFR2^{SHF-KO}$; $Apj^{DreERT2}$ embryos at E10.5. Data were analyzed using 2-tailed, unpaired Student's t-test; p>0.05 was considered not significant.

Supplemental Tables

Supplemental Table 1. Penetrance of aberrant PAA formation. PAAs of E10.5 embryos ranging from 34-36s were analyzed via WM-IF staining for VEGFR2. Aberrant PAAs were marked as "hypoplastic" when their diameter was markedly smaller than controls or as "aplastic," indicating the absence of a PAA (e.g., **Fig. 2** and **Fig. S1**). "0" indicates PAAs comparable to controls. n = the number of individual pharyngeal arches analyzed. Percent of penetrance is reported in paratheses.

Supplemental Table 2. The average number of SHF-derived ECs in the pharyngeal arches. PAAs of E10.5 $VEGFR2^{SHF-WT}$ and $VEGFR2^{SHF-KO}$ embryos ranging from 34-36s were analyzed via WM-IF staining. The total number of SHF cells was quantified in each pharyngeal arch. Percent of VEGFR2-positive and VEGFR2-negative SHF cells was quantified. Averages ± SD are reported in the table. Statistical analyses were reported in **Fig. 2G**.

Supplemental Table 3. The average number ECs in the pharyngeal arches. PAAs of E10.5 $VEGFR2^{SHF-WT}$, $VEGFR2^{SHF-Het}$, $VEGFR2^{SHF-KO}$, and $Tbx1^{+/-}$ embryos ranging from 34-36s were analyzed via WM-IF. The total number of EC cells was quantified per pharyngeal arch. Percent of SHF-derived and non-SHF-derived ECs was quantified. Averages ± SD are reported in the table. Statistical analyses were reported in **Fig. 2H-I**.

Supplemental Table 4. Loss of VEGFR2 in the SHF is not embryonic lethal. Observed and expected values of embryos dissected between E14.5 and P0 for each genotype. n = the number of embryos observed and expected. Percent of each is reported in paratheses. Statistics were evaluated using Chi-squared test.

Supplemental Table 5. Cardiovascular defects in *VEGFR2^Isl1-KO* and *VEGFR2^SHF-KO* embryos.

Abbreviations: H&E – hematoxylin and eosin; µCT – micro-computed tomography; RScA – right

subclavian artery; IAA-B – interrupted aortic arch type B; IAA-C – interrupted aortic arch type C.

Arch artery defects lethal postnatally are labeled in red and marked by an asterisk.

Supplemental Table 6. Cardiovascular defects in *VEGFR2^Isl1-Het* and *VEGFR2^SHF-Het* embryos.

Abbreviations: H&E – hematoxylin and eosin; µCT – micro-computed tomography; RScA – right

subclavian artery.

Supplemental Table 7. Meta-analyses of PAA defects in 22Q11DS models

Supplemental Table 8. Meta-analyses of AAA defects in 22Q11DS models

References

Abrams, M.J., Basinger, T., Yuan, W., Guo, C.L., Goentoro, L., 2015. Self-repairing symmetry in jellyfish through mechanically driven reorganization. Proc Natl Acad Sci U S A 112, E3365-3373.

Aggarwal, V.S., Liao, J., Bondarev, A., Schimmang, T., Lewandoski, M., Locker, J., Shanske, A., Campione, M., Morrow, B.E., 2006. Dissection of Tbx1 and Fgf interactions in mouse models of 22q11DS suggests functional redundancy. Hum Mol Genet 15, 3219-3228.

Anderson, R.H., Bamforth, S.D., 2022. Morphogenesis of the Mammalian Aortic Arch Arteries. Front Cell Dev Biol 10, 892900.

Benjamin, E.J., Virani, S.S., Callaway, C.W., Chamberlain, A.M., Chang, A.R., Cheng, S., Chiuve, S.E., Cushman, M., Delling, F.N., Deo, R., de Ferranti, S.D., Ferguson, J.F., Fornage, M., Gillespie, C., Isasi, C.R., Jiménez, M.C., Jordan, L.C., Judd, S.E., Lackland, D., Lichtman, J.H., Lisabeth, L., Liu, S., Longenecker, C.T., Lutsey, P.L., Mackey, J.S., Matchar, D.B., Matsushita, K., Mussolino, M.E., Nasir, K., O'Flaherty, M., Palaniappan, L.P., Pandey, A., Pandey, D.K., Reeves, M.J., Ritchey, M.D., Rodriguez, C.J., Roth, G.A., Rosamond, W.D., Sampson, U.K.A.A., Satou, G.M., Shah, S.H., Spartano, N.L., Tirschwell, D.L., Tsao, C.W., Voeks, J.H., Willey, J.Z., Wilkins, J.T., Wu, J.H.Y., Alger, H.M., Wong, S.S., Muntner, P., 2018. Heart Disease and Stroke Statistics—2018 Update: A Report From the American Heart Association, Circulation, pp. e67-e492.

Calmont, A., Ivins, S., Van Bueren, K.L., Papangeli, I., Kyriakopoulou, V., Andrews, W.D., Martin, J.F., Moon, A.M., Illingworth, E.A., Basson, M.A., Scambler, P.J., 2009. Tbx1 controls cardiac neural crest cell migration during arch artery development by regulating Gbx2 expression in the pharyngeal ectoderm, Development, pp. 3173-3183.

Celoria, G., Patton, R., 1959. Congenital absence of the aortic arch. American Heart Journal 58, 407-413.

Chen, H.I., Sharma, B., Akerberg, B.N., Numi, H.J., Kivela, R., Saharinen, P., Aghajanian, H., McKay, A.S., Bogard, P.E., Chang, A.H., Jacobs, A.H., Epstein, J.A., Stankunas, K., Alitalo, K., Red-Horse, K., 2014. The sinus venosus contributes to coronary vasculature through VEGFC-stimulated angiogenesis, Development, pp. 4500-4512.

Collins-Nakai, R.L., Dick, M., Parisi-Buckley, L., Fyler, D.C., Castaneda, A.R., 1976. Interrupted

aortic arch in infancy. The Journal of Pediatrics 88, 959-962.

Goldmuntz, E., Driscol, D.A., Budarf, M.L., Zackai, E., McDonald-McGinn, D.M., Biegel, J.A.,

Emanuel, B.S., 1993. Microdeletions of chromosomal region 22q11 in patients with congenital

conotruncal cardiac defects. Journal of medical genetics 30, 807-812.

Graham, A., 2003. Development of the pharyngeal arches. Am J Med Genet A 119A, 251-256.

Guris, D.L., Duester, G., Papaioannou, V.E., Imamoto, A., 2006. Dose-dependent interaction of

Tbx1 and Crkl and locally aberrant RA signaling in a model of del22q11 syndrome. Dev Cell 10,

81-92.

Hasten, E., Morrow, B.E., 2019. Tbx1 and Foxi3 genetically interact in the pharyngeal pouch

endoderm in a mouse model for 22q11.2 deletion syndrome. PLoS Genet 15, e1008301.

Hutson, M.R., Kirby, M.L., 2007. Model systems for the study of heart development and disease

Cardiac neural crest and conotruncal malformations, Seminars in Cell & Developmental Biology,

pp. 101-110.

Jelier, R., Kruger, A., Swoger, J., Zimmermann, T., Lehner, B., 2016. Compensatory Cell

Movements Confer Robustness to Mechanical Deformation during Embryonic Development.

Cell Syst 3, 160-171.

Jerome, L.A., Papaioannou, V.E., 2001. DiGeorge syndrome phenotype in mice mutant for the

T-box gene, Tbx1, Nature Genetics, pp. 286-291.

Kau, T., Sinzig, M., Gasser, J., Lesnik, G., Rabitsch, E., Celedin, S., Eicher, W., Illiasch, H.,

Hausegger, K., 2007. Aortic Development and Anomalies, Seminars in Interventional Radiology,

pp. 141-152.

Levitt, B., Richter, J.E., 2007. Dysphagia lusoria: a comprehensive review. Dis Esophagus 20,

455-460.

Li, P., Pashmforoush, M., Sucov, H.M., 2012. Mesodermal retinoic acid signaling regulates

endothelial cell coalescence in caudal pharyngeal arch artery vasculogenesis. Dev Biol 361,

116-124.

Lindsay, E., Botta, A., Vesna, J., Carattini-Rivera, S., Cheah, Y.-C., Rosenblatt, H.M., Bradley,
A., Baldini, A., 1999. Congenital heart disease in mice deficient for the DiGeorge syndrome
region. Nature 401, 379-383.

Lindsay, E.A., Baldini, A., 2001. Recovery from arterial growth delay reduces penetrance of
cardiovascular defects in mice deleted for the DiGeorge syndrome region. Human Molecular
Genetics 10, 997-1002.

Lindsay, E.a., Vitelli, F., Su, H., Morishima, M., Huynh, T., Pramparo, T., Jurecic, V., Ogunrinu,
G., Sutherland, H.F., Scambler, P.J., Bradley, a., Baldini, a., 2001. Tbx1 haploinsufficieny in the
DiGeorge syndrome region causes aortic arch defects in mice., Nature, pp. 97-101.

McDonald-McGinn, D.M., Sullivan, K.E., 2011. Chromosome 22q11.2 deletion syndrome
(DiGeorge syndrome/velocardiofacial syndrome). Medicine (Baltimore) 90, 1-18.

McDonald-McGinn, D.M., Sullivan, K.E., Marino, B., Philip, N., Swillen, A., Vorstman, J.A.,
Zackai, E.H., Emanuel, B.S., Vermeesch, J.R., Morrow, B.E., Scambler, P.J., Bassett, A.S.,
2015. 22q11.2 deletion syndrome. Nat Rev Dis Primers 1, 15071.

Merscher, S., Funke, B., Epstein, J.A., Heyer, J., Puech, A., Lu, M.M., Xavier, R.J., Demay,
M.B., Russell, R.G., Factor, S., Tokooya, K., Jore, B.S., Lopez, M., Pandita, R.K., Lia, M.,
Carrion, D., Xu, H., Schorle, H., Kobler, J.B., Scambler, P., Wynshaw-Boris, A., Skoultchi, A.I.,
Morrow, B.E., Kucherlapati, R., 2001. TBX1 is responsible for cardiovascular defects in velo-
cardio-facial/DiGeorge syndrome, Cell, pp. 619-629.

Moerman, P., Dumoulin, M., Lauweryns, J., Van der Hauwaert, L.G., 1987. Interrupted right
aortic arch in DiGeorge syndrome., Heart, pp. 274-278.

Momma, K., 2010. Cardiovascular anomalies associated with chromosome 22q11.2 deletion
syndrome. Am J Cardiol 105, 1617-1624.

Nagelberg, D., Wang, J., Su, R., Torres-Vázquez, J., Targoff, K.L., Poss, K.D., Knaut, H., 2015.
Origin, specification, and plasticity of the great vessels of the heart, Current Biology, pp. 2099-
2110.

Paffett-Lugassy, N., Singh, R., Nevis, K.R., Guner-Ataman, B., O'Loughlin, E., Jahangiri, L.,

Harvey, R.P., Burns, C.G., Burns, C.E., 2013. Heart field origin of great vessel precursors relies

on nkx2.5-mediated vasculogenesis, Nature Cell Biology, pp. 1362-1369.

Papangeli, I., Scambler, P.J., 2013. Tbx1 Genetically Interacts With the Transforming Growth

Factor- /Bone Morphogenetic Protein Inhibitor Smad7 During Great Vessel Remodeling,

Circulation Research, pp. 90-102.

Phillips, H.M., Stothard, C.A., Shaikh Qureshi, W.M., Kousa, A.I., Briones-Leon, J.A.,

Khasawneh, R.R., O'Loughlin, C., Sanders, R., Mazzotta, S., Dodds, R., Seidel, K., Bates, T.,

Nakatomi, M., Cockell, S.J., Schneider, J.E., Mohun, T.J., Maehr, R., Kist, R., Peters, H.,

Bamforth, S.D., 2019. Pax9 is required for cardiovascular development and interacts with Tbx1

in the pharyngeal endoderm to control 4th pharyngeal arch artery morphogenesis,

Development, p. dev177618.

Randall, V., McCue, K., Roberts, C., Kyriakopoulou, V., Beddow, S., Barrett, A.N., Vitelli, F.,

Prescott, K., Shaw-Smith, C., Devriendt, K., Bosman, E., Steffes, G., Steel, K.P., Simrick, S.,

Basson, M.A., Illingworth, E., Scambler, P.J., 2009. Great vessel development requires biallelic

expression of Chd7 and Tbx1 in pharyngeal ectoderm in mice. J Clin Invest 119, 3301-3310.

Ryckebusch, L., Bertrand, N., Mesbah, K., Bajolle, F., Niederreither, K., Kelly, R.G., Zaffran, S.,

2010. Decreased levels of embryonic retinoic acid synthesis accelerate recovery from arterial

growth delay in a mouse model of DiGeorge syndrome. Circ Res 106, 686-694.

Sharma, B., Ho, L., Ford, G.H., Chen, H.I., Goldstone, A.B., Woo, Y.J., Quertermous, T.,

Reversade, B., Red-Horse, K., 2017. Alternative Progenitor Cells Compensate to Rebuild the

Coronary Vasculature in Elabela- and Apj-Deficient Hearts. Dev Cell 42, 655-666 e653.

Stoller, J.Z., Epstein, J.A., 2005. Cardiac neural crest, Seminars in Cell & Developmental

Biology, pp. 704-715.

Stothard, C.A., Mazzotta, S., Vyas, A., Schneider, J.E., Mohun, T.J., Henderson, D.J., Phillips,

H.M., Bamforth, S.D., 2020. Pax9 and Gbx2 Interact in the Pharyngeal Endoderm to Control

Cardiovascular Development. J Cardiovasc Dev Dis 7.

Taddei, I., Morishima, M., Huynh, T., Lindsay, E.A., 2001. Genetic factors are major
determinants of phenotypic variability in a mouse model of the DiGeorge:del22q11 syndromes.
PNAS 98, 11428-11431.

Tsao, C.W., Aday, A.W., Almarzooq, Z.I., Alonso, A., Beaton, A.Z., Bittencourt, M.S., Boehme,
A.K., Buxton, A.E., Carson, A.P., Commodore-Mensah, Y., Elkind, M.S.V., Evenson, K.R., Eze-
Nliam, C., Ferguson, J.F., Generoso, G., Ho, J.E., Kalani, R., Khan, S.S., Kissela, B.M.,
Knutson, K.L., Levine, D.A., Lewis, T.T., Liu, J., Loop, M.S., Ma, J., Mussolino, M.E.,
Navaneethan, S.D., Perak, A.M., Poudel, R., Rezk-Hanna, M., Roth, G.A., Schroeder, E.B.,
Shah, S.H., Thacker, E.L., VanWagner, L.B., Virani, S.S., Voecks, J.H., Wang, N.Y., Yaffe, K.,
Martin, S.S., 2022. Heart Disease and Stroke Statistics-2022 Update: A Report From the
American Heart Association. Circulation 145, e153-e639.

Van Mierop, L.H.S., Kutsche, L.M., 1986a. Cardiovascular anomalies in digeorge syndrome and
importance of neural crest as a possible pathogenetic factor, The American Journal of
Cardiology, pp. 133-137.

Van Mierop, L.H.S., Kutsche, L.M., 1986b. Cardiovascular anomalies in DiGeorge syndrome
and importance of neural crest as a possible pathogenetic factor. The American Journal of
Cardiology 58, 133-137.

Vitelli, F., Taddei, I., Morishima, M., Meyers, E., Lindsay, E.A., Baldini, A., 2002. A genetic link
between Tbx1 and fibroblast growth factor signaling. Development 129, 6.

Waddington, C.H., 1942. Canalization of development and the inheritance of the acquired
characters. Nature 150, 563-564.

Wang, X., Chen, D., Chen, K., Jubran, A., Ramirez, A., Astrof, S., 2017. Endothelium in the
pharyngeal arches 3, 4 and 6 is derived from the second heart field, Developmental Biology, pp.
108-117.

Warkala, M., Chen, D., Ramirez, A., Jubran, A., Schonning, M., Wang, X., Zhao, H., Astrof, S.,
2021. Cell-Extracellular Matrix Interactions Play Multiple Essential Roles in Aortic Arch
Development. Circ Res 128, e27-e44.

Yagi, H., Furutani, Y., Hamada, H., Sasaki, T., Asakawa, S., Minoshima, S., Ichida, F., Joo, K.,

Kimura, M., Imamura, S.-i., Kamatani, N., Momma, K., Takao, A., Nakazawa, M., Shimizu, N.,

Matsuoka, R., 2003. Role of TBX1 in human del22q11.2 syndrome, The Lancet, pp. 1366-1373.

Zhang, Z., 2006. Mesodermal expression of Tbx1 is necessary and sufficient for pharyngeal

arch and cardiac outflow tract development, Development, pp. 3587-3595.

Zhang, Z., Baldini, A., 2008. In vivo response to high-resolution variation of Tbx1 mRNA

dosage, Human Molecular Genetics, pp. 150-157.

Zhang, Z., Cerrato, F., Xu, H., Vitelli, F., Morishima, M., Vincentz, J., Furuta, Y., Ma, L., Martin,

J.F., Baldini, A., Lindsay, E., 2005. Tbx1 expression in pharyngeal epithelia is necessary for

pharyngeal arch artery development., Development (Cambridge, England), pp. 5307-5315.

Figure 1

Figure 1. Efficient ablation of VEGFR2 in the second heart field (SHF) using *mef2c-AHF-Cre* mice.

Cre-lox recombination and immunofluorescent staining were used to trace *mef2c-AHF-Cre* lineage cells (cyan) when VEGFR2 (magenta) is conditionally ablated from the SHF in embryos. **(A-E)** Optical confocal sections of E9.5 *VEGFR2^SHF-WT^* embryos show VEGFR2+ cells in the anterior **(B-C)** and posterior **(D-E)** dorsal pericardial wall outlined in yellow. **(F-I)** PAA ECs in E10.5 *VEGFR2^SHF-WT^* embryos are derived from the SHF. The box in **F** is expanded in **(G-I)**. Arrowheads point to the co-expression of VEGFR2 (magenta), ERG (orange), and Cre-reporter (cyan) in the 4th pharyngeal arch endothelium. **(J-N)** Optical sections of E9.5 *VEGFR2^SHF-KO^* embryos show the absence of VEGFR2 expression in the anterior **(K-L)** and posterior SHF dorsal pericardial wall **(M-N)**, outlined in yellow. **(O-R)** PAA ECs in E10.5 *VEGFR2^SHF-KO^* embryos are derived from a non-SHF endothelium. The box in **O** is expanded in **(P-R)**. Arrowheads point to Cre-reporter negative endothelium. Arrows point to Cre-reporter+ VEGFR2-negative cells.

Figure 2

VEGFR2 **Cre-Reporter** **DAPI**

VEGFR2^SHF-WT

A
3rd
4th
6th
36s   100µm

B
3rd
4th
6th
50µm

VEGFR2^SHF-Het

C
3rd
4th
6th
36s   100µm

D
3rd
4th
6th
50µm

VEGFR2^SHF-KO

E
3rd
4th
6th
36s   100µm

F
3rd
4th
6th
50µm

G Loss of VEGFR2 does not affect the total number of Cre-reporter+ cells in the pharyngeal arches

H Loss of VEGFR2 in the SHF does not affect the total number of ECs in the pharyngeal arches

I Loss of VEGFR2 in the SHF results in EC compensation from an alternative source

G: Number of Cre-reporter+ cells vs. 3rd PA / 4th PA / 6th PA (SHF-WT, SHF-Het, SHF-KO), ns

H: Total Number of ECs vs. 3rd PA / 4th PA / 6th PA (SHF-WT, SHF-Het, SHF-KO), ns, *, ns

I: Percent vs. 3rd PA / 4th PA / 6th PA (SHF-WT, SHF-Het, SHF-KO), **, ***, ***, ****, ***, ****

■ SHF-derived ECs
▨ non-SHF-derived ECs

Figure 2. VEGFR2 is required for SHF vascular progenitor contribution and PAA development.
(A-F) WM-IF for VEGFR2 (magenta) and Cre-reporter (cyan), confocal imaging, and 3D reconstruction of E10.5 **(A)** $VEGFR2^{SHF-WT}$, **(C)** $VEGFR2^{SHF-Het}$, and **(E)** $VEGFR2^{SHF-KO}$ embryos. Yellow brackets highlight aberrant PAAs in **C, E**. Dashed lines mark the levels of coronal optical sections through the pharyngeal arches shown in **B, D,** and **F**. Dotted circles mark aberrant PAAs. **(G)** Quantification of SHF-derived cells in each pharyngeal arch. $VEGFR2^{SHF-WT}$ n = 10 arches, $VEGFR2^{SHF-Het}$ n = 4 arches, $VEGFR2^{SHF-KO}$ n = 6 arches. Each dot marks one pharyngeal arch. Horizontal red and black lines mark means and quartiles, respectively. **(H)** Quantification of total EC numbers in each pharyngeal arch. $VEGFR2^{SHF-WT}$ n = 10 arches, $VEGFR2^{SHF-Het}$ n = 12 arches, $VEGFR2^{SHF-KO}$ n = 12 arches. Each dot marks one pharyngeal arch. Horizontal red and black lines mark means and quartiles, respectively. **(I)** The proportion of Cre-reporter-positive ECs and Cre-reporter-negative ECs in each pharyngeal arch. Means ± SD are shown. $VEGFR2^{SHF-WT}$ n = 10 arches, $VEGFR2^{SHF-Het}$ n = 12 arches, $VEGFR2^{SHF-KO}$ n = 12 arches. Data were analyzed using two-way ANOVA with Tukey's correction for multiple testing; *$p \leq 0.05$ **$p \leq 0.01$, ***$p \leq 0.001$, ****$p < 0.0001$

Figure 3

Figure 3. A threshold number of SHF-derived ECs is required for proper Aortic Arch Artery development.

(A) Percent of *VEGFR2;Isl1Cre* (*VEGFR2^Isl1-WT^, VEGFR2^Isl1-Het^,* and *VEGFR2^Isl1-KO^*) and *VEGFR2;mef2c-AHF-Cre* (*VEGFR2^SHF-WT^, VEGFR2^SHF-Het^,* and *VEGFR2^SHF-KO^*) embryos without AAA defects (white), non-lethal AAA defects (grey), or lethal AAA defects (black). n = the number of embryos with each phenotype. **(B-D)** Transverse sections of E18.5 embryos stained with Hematoxylin and eosin. **(B)** *VEGFR2^SHF-WT^* embryos. **(C)** IAA-B and **(D)** RERSA in *VEGFR2^Isl1-KO^* and *VEGFR2^SHF-KO^* embryos, respectively. **(E-G)** Micro-computed tomography of E18.5 *VEGFR2^Isl1-WT^* **(E)** and *VEGFR2^Isl1-KO^* **(F-G)** embryos. Arrows and text indicate examples of IAA-B **(F)** and aberrant RScA **(G)** in yellow. Dotted boxes highlight the position of a normal aortic arch. **Abbreviations:** RCA – right carotid artery; LCA – left carotid artery; RScA – right subclavian artery; LScA – left subclavian artery; BrA – brachiocephalic artery; AoA – aortic arch; PT – pulmonary trunk; DA – ductus arteriosus; dAo – descending aorta; TR – trachea; IAA-B – interrupted aortic arch type B; RERSA – retroesophageal right subclavian artery

Figure 4. *Tbx1* regulates endothelial compensation and rescue of PAA development.
(A-B) WM-IF for VEGFR2 (magenta) and Cre-reporter (cyan) of an E10.5 *Tbx1⁺/⁻* embryo. **(A)** 3D reconstruction and **(B)** coronal optical sections show the incomplete formation of the 4th PAA (dotted circles) and that only a few non-SHF-derived ECs are present. **(C)** Quantification of ECs in the 4th pharyngeal arches of control (*mef2c-AHF-Cre*), *Tbx1⁺/⁻*, and *VEGFR2^{SHF-Het}* embryos. Each dot marks one pharyngeal arch. Horizontal red and black lines mark means and quartiles, respectively. **(D)** Quantification of the number of SHF-derived (Cre-reporter positive) ECs in the 4th pharyngeal arch. Error bars represent mean ± SD. **(E)** Quantification of the number of non-SHF-derived (Cre-reporter negative) ECs in the 4th pharyngeal arch. Error bars represent mean ± SD. Control n = 14 arches, *Tbx1⁺/⁻* n = 6 arches, *VEGFR2^{SHF-Het}* n = 12 arches. Data were analyzed using one-way ANOVA with Tukey's correction for multiple testing; **p ≤ 0.01, ****p < 0.0001

Figure 5

Lateral → Medial

Figure 5. Ectopic sprouts from veins in the pharyngeal region of *VEGFR2^SHF-KO* are absent in *Tbx1^+/-* embryos.
(A) A schematic depiction of venous and arterial vasculature in the pharyngeal region of E9.5 embryos. dAo – dorsal aorta; CV – cardinal vein; PV – pharyngeal vein. **(B-D)** 20μm-thick serial sagittal optical sections through E9.5 *VEGFR2^SHF-WT* **(B)**, *VEGFR2^SHF-KO* **(C)**, and *Tbx1^+/-* **(D)** embryos from the lateral to the medial side. Yellow brackets in **B-D** mark pharyngeal endothelial plexus. Red stars mark absent venous sprouts in *VEGFR2^SHF-WT* **(B)** and *Tbx1^+/-* **(D)** embryos. Yellow, magenta, and cyan arrows follow sprouts from the CV to the pharyngeal arch vasculature in the VEGFR2SHF-KO embryos **(C)**.

Figure 6

Loss of VEGFR2 in the SHF leads to an increase in Dre-Reporter+ ECs in the 4th PAA

Loss of VEGFR2 in the SHF results in venous ECs in the 4th PA

Figure 6. Angiogenic sprouting from veins gives rise to the compensatory endothelium that rescues PAA morphogenesis.

Dre-rox recombination was used to fate map cells of *Apj*-lineage in *VEGFR2^{SHF-WT}* and *VEGFR2^{SHF-KO}* embryos. Tamoxifen was injected at E9.5, and embryos were dissected at E10.5. **(A-D)** WM-IF staining detecting Dre-reporter (cyan) and VEGFR2 (magenta). 50µm-thick sagittal optical sections show *Apj-DreERT2* lineage labeling in the CV and intersomitic veins (yellow arrows) but not the dAo or intersomitic arteries (white arrows) of **(A)** *VEGFR2^{SHF-WT}* and **(C)** *VEGFR2^{SHF-KO}* embryos. Boxes in **B, D** are expanded in the panels on the right to show *ApjDreERT2* lineage labeled ECs in the 4th PAAs of *VEGFR2^{SHF-KO}* embryos (arrowheads, **D**) but not in control, *VEGFR2^{SHF-WT}* embryos **(B)**. **(E)** *VEGFR2^{SHF-KO}* embryos have a higher percent contribution of *Apj*-lineage cells in the 4th PAAs. The number of *Apj*-lineage cells in arches was normalized by *ApjDreERT2* labeling efficiency (see Methods). Each dot marks one pharyngeal arch. Horizontal red and black lines mark means and quartiles, respectively. *VEGFR2^{SHF-WT}* n = 10 arches, *VEGFR2^{SHF-KO}* n = 6 arches. **(F-G)** WM-IF for COUPTF2 (magenta) and VEGFR2 (white) of E9.5 embryos. 20µm-thick sagittal optical sections: **(F)** *VEGFR2^{SHF-WT}* and **(G)** *VEGFR2^{SHF-KO}* embryos. Yellow arrows point to COUPTF2+ ECs in the CV. White arrows point to COUPTF2-negative ECs in 3rd PAA and dAo. Arrowheads point to COUPTF2+ ECs in the 4th PAA of *VEGFR2^{SHF-KO}* embryo **(G)**. **(H)** Quantification of COUPTF2+ ECs in the 4th pharyngeal arches. Each dot marks one pharyngeal arch. Horizontal red and black lines mark means and quartiles, respectively. *VEGFR2^{SHF-WT}* n = 9 arches, *VEGFR2^{SHF-KO}* n = 7 arches. **(I-J)** Confocal optical sections. Single-molecule RNA fluorescence in situ hybridization for *Apj* mRNA (magenta) combined with immunostaining for VEGFR2 (white) of E10.0 embryos. Yellow arrows point to *Apj*+ ECs in the CV. White arrows point to *Apj*-negative ECs in 3rd PAA and dAo. **(K)** Quantification of *Apj*+ ECs in the 4th pharyngeal arches. Each dot marks one pharyngeal arch. Horizontal red and black lines mark means and quartiles, respectively. *VEGFR2^{SHF-WT}* n = 6 arches, *VEGFR2^{SHF-KO}* n = 7 arches. Data were analyzed using 2-tailed, unpaired Student's t-test; *$p \leq$ 0.05, ** $p \leq$ 0.01, ***$p \leq$ 0.001.

Figure 7

Figure 7. Loss of SHF-derived vascular progenitors increases *VEGFA* mRNA expression in the SHF and pharyngeal endoderm.

(A-B) Single-molecule RNA fluorescent in situ hybridization of E9.5 embryos detecting the expression of *VEGFA* (white) in the SHF (outlined in red) and pharyngeal endoderm (outlined in yellow). **(C)** Super plot of normalized mean fluorescence intensity of *VEGFA* mRNA in the SHF. **(D)** Super plot of normalized mean fluorescence intensity of *VEGFA* mRNA in the pharyngeal endoderm. The mean fluorescence intensity in the SHF and pharyngeal endoderm was normalized to the mean fluorescence intensity in the dorsal side of the embryo on the same optical section (outlined in green). Different colors in **C, D** mark different biological samples. Light-colored dots mark individual measurements in optical sections. Dark-colored dots mark the average of measurements per pharyngeal arch. Red lines mark means, and black lines mark quartiles. *VEGFR2^{SHF-WT}* n = 6 arches, *VEGFR2^{SHF-KO}* n = 6 arches. Data were analyzed using 2-tailed, unpaired Student's t-test; **p ≤ 0.01.