Mutational signatures in wild type *Escherichia coli* strains reveal dominance of DNA polymerase errors

Sofya K. Garushyants[*,1,2], Mrudula Sane[3], Maria V. Selifanova[4], Deepa Agashe[3], Georgii A. Bazykin[1,5], Mikhail S. Gelfand[*,1,5]

[1] A.A. Kharkevich Institute for Information Transmission Problems, RAS, Moscow, Russia

[2] Present address: National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, Bethesda, MD, USA

[3] National Centre for Biological Sciences, Bengaluru, India

[4] Faculty of Bioengineering and Bioinformatics, M.V. Lomonosov Moscow State University, Moscow, Russia

[5] Skolkovo Institute of Science and Technology (Skoltech), Moscow, Russia

* Correspondence: Sofya K. Garushyants garushyants@iitp.ru; Mikhail S. Gelfand mikhail.gelfand@gmail.com

Abstract

While mutational processes operating in the *Escherichia coli* genome have been revealed by multiple laboratory experiments, the contribution of these processes to accumulation of bacterial polymorphism and evolution in natural environments is unknown. To address this question, we reconstruct signatures of distinct mutational processes from experimental data on *E. coli* hypermutators, and ask how these processes contribute to differences between naturally occurring *E. coli* strains. We show that both mutations accumulated in the course of evolution of wild type strains in nature and in the lab-grown non-mutator laboratory strains are explained predominantly by the low fidelity of DNA polymerases II and III. By contrast, contributions specific to disruption of DNA repair systems cannot be detected, suggesting that temporary accelerations of mutagenesis associated with such disruptions are unimportant for within-species evolution. These observations demonstrate that accumulation of diversity in bacterial strains in nature is predominantly associated with errors of DNA polymerases.

Introduction

Mutagenesis provides the raw material for evolution, and understanding the processes involved in it is essential for studying the evolution of bacterial genomes. The progress in this area followed the rise of genome sequencing techniques, which allowed for genome-scale comparative analyses not restricted to reporter genes [1–6]. Mutations occur in all cells at each replication cycle. However, mutation rates may differ even between laboratory strains of the same bacterial species[1]. Spontaneous mutations occur due to errors of enzymes involved in DNA replication or repair, combined with exogenous factors[7–9]. In bacteria, disruption of these processes yields hypermutator phenotypes, which are characterized by elevated mutation rates and changes in the mutational spectra[1,3,10]. Hypermutators accumulate point mutations at rates that are orders of magnitude higher than those of wildtype strains[11].

Multiple distinct hypermutator phenotypes are known. The MutS, MutL, and MutH proteins contribute to hypermutator phenotypes as parts of the DNA mismatch repair system (MMR)[4]. In wildtype laboratory strains, mutations are substantially underrepresented in transcribed regions; however, this bias disappears in MMR-defective strains, which means that MMR preferably repairs transcribed regions[1,3]. A different type of hypermutators arise due to mutations in DnaQ, DNA polymerase III ε subunit that functions as a 5'-exonuclease ensuring the accuracy of chromosome replication[12,13]. On a rich medium, mutants that carry an almost inactive DnaQ have a 10 to 1000-fold higher mutation rate than the wildtype strains[14]. Inactivation of DnaQ also leads to activation of the SOS response[15]; however, SOS-induced error-prone polymerases do not contribute to the mutation rate or spectra of

DnaQ mutants[5], indicating that the changes observed in DnaQ mutants are caused by DnaQ inactivation

itself. Furthermore, a mutator phenotype can arise from mutations in error-prone DNA polymerases IV

(UmuDC) and V (DinB), and in DNA polymerase II (PolB). Inactivation of these polymerases yields

elevated mutation rates under some conditions[16–18] but not in normally growing cells[3]. Finally,

inefficient removal of oxidized nucleotides, especially 8-oxoG, also influences mutation accumulation

rates and spectra[3]. Oxidized guanines are removed via two major pathways, MutT that hydrolyzes 8-

oxoG, or MutM or MutY that correct 8-oxoG:A mispairs[19,20].

Previous studies of mutations in bacteria mainly considered single-nucleotide substitutions regardless

of the adjacent nucleotides. Meanwhile, accounting for these nucleotides, the so-called mutational

contexts, provides a finer resolution and allows to distinguish between processes that underlie

mutagenesis. This approach has been widely applied and proved to be fruitful in cancer genomics[9,21–25].

Typically, it is assumed that individual processes affecting mutagenesis are characterized by specific

patterns of mutations, or mutational signatures. A mutational signature as used here is characterized by

the relative frequencies of six substitution subtypes: C>A (that is, the C:G pair to the A:T pair; hence

equivalent to G>T), C>G, C>T, T>A, T>C, and T>G (equivalent, respectively, to G>C, G>A, A>T,

A>G, and A>C) for each of the four possible adjacent 5' and 3' bases. This yields a total of 96 (6×4×4)

possible mutation types. In cancer genomics, relative contributions of various mutational processes are

inferred from tumor samples by decomposition of the observed mutational spectra into several standard

signatures[20]. By contrast, in bacteria, mutational signatures may be observed directly by analyzing

genomes of spontaneous mutator strains or laboratory knockouts of particular genes yielding mutation

accumulation (MA) lines. Sequencing data on MA lines in model bacterial species are now available

for strains generated in several laboratories in different set-ups, mostly for *Escherichia coli*[1,3,5,26–29].

Furthermore, data on naturally occurring mutators can be obtained from *in vitro* evolution experiments.

For example, in the *E. coli* Long-term Evolution Experiment (LTEE)[4,30], six strains acquired

spontaneous mutations that affect MMR or removal of oxidized nucleotides. Both MA lines and

evolution experiments allow for observing thousands of single nucleotide substitutions in mutator

lineages, providing sufficient resolution to characterize mutational signatures associated with various

mutational mechanisms.

In hypermutator strains, evolution primarily reflects mutational biases, with minimal contribution of

selection[30], so that mutational processes may be directly inferred from whole-genome mutational

patterns. By contrast, slower accumulation of mutations in non-mutator strains is shaped by selection,

both positive and negative[30], with most positions in bacterial genomes subject to negative selection.

Still, intergenic regions evolve in an almost neutral regime[31,32], or are at least less affected by negative

selection[33,34]. This allowed us to use intergenic regions to infer mutational processes in non-mutator

strains.

Here, we address two questions: whether the mutational signatures and mutational processes

underlying them may be inferred from experimental data on hypermutators; and how these mutational

processes contribute to accumulation of mutations in *E. coli* strains in laboratory and in nature.

Results

Meta-analysis of mutation accumulation lines in *E. coli*

To study mutations accumulated via different mutational processes, we collected all available data on

mutation accumulation in mutator strains of *E. coli* (Materials and Methods). We used sequencing data

from four main sources: (i) six mutator strains that had emerged in the LTEE experiment: Ara-1 and

Ara+6 with the MutT deficiency and Ara-2, Ara-3, Ara-4 and Ara+3 with mutations in MMR-

associated genes[4]; (ii) MA strains for all major mutational processes from the Foster laboratory[1,3,5], (iii)

mutY deficient MA strains[6]; and (iii) a set of MA strains deficient in MMR components MutH, MutL,

and MutS obtained specifically for this study (Supplementary Figure 1; Materials and Methods).

The resulting dataset included data on multiple distinct mutational processes, and for many of these

processes, data came from several sources. Specifically, strains with defective MMR systems were

collected from three independent labs, and in total yielded almost 12000 mutations (Supplementary

Figure 1). The MutT and MutY mutator strains came each from two independent labs and had

respectively nearly 9000 and 5000 mutations. We also considered strains with defective genes encoding

the epsilon subunit of DNA polymerase III (*dnaQ*), DNA polymerase II (*polB*), DNA polymerase IV

(*umuDC*), nucleotide excision repair ATPase (*uvrA*), base excision repair (BER) endonucleases III and

VIII (*nth* and *nei*, respectively), and exonucleases III and IV (*xthA* and *nfo*, respectively)[3,5]. In total, our

dataset included 40694 mutations from 17 experiments (Supplementary Figure 1).

To describe the overall mutation types characteristic of mutational processes in our data, we first

extracted single-nucleotide substitutions and combined complementary mutations, hence retaining six

types of substitutions. MA lines with similar deficiencies, such as MMR, MutT, and MutY deficient

strains obtained in different laboratories, had similar single-nucleotide mutational profiles, which

means that the observed mutational spectra were independent of experimental conditions

(Supplementary Figure 1). As shown earlier, MMR and DnaQ mutants were associated with transitions

(C>T and T>C)[3]; MutY mutants generated mostly C>A transversions; whereas MutT mutants almost

exclusively carried T>G transversions. Strains deficient in the *nth* and *nei* genes carried prevalent C>T

substitutions. MA strains deficient in *polB, umuDC, uvrA, xthA,* and *nfo* had relatively flat mutational

profiles with a higher prevalence of all six types of substitutions relative to the "wild type" laboratory

E. coli strain (Supplementary Figure 1).

Reconstruction of mutational signatures from hypermutators

Analyzing only substitutions without accounting for their contexts does not produce sufficient

resolution to distinguish, for example, between the excess of C>T mutations caused by MMR and by

errors of polymerase III. To distinguish between mutations caused by different mutational processes,

we considered three-nucleotide contexts, again collapsing complementary mutations, for a total of 96

types of mutations. We reconstructed the mutational profiles for each strain independently

(Supplementary Figure 2).

We compared the mutational profiles of each strain using cosine similarity (see Materials and

Methods), and observed that they form distinct clusters (Fig. 1a). We divided all strains into seven

groups based on similarity of their mutational profiles: MMR-deficient strains from the LTEE

experiment; DnaQ MA strains; MMR-deficient MA strains; MA strains with defective MutY, MutT,

BER endonucleases III and VIII; and PolB-deficient and similar MA strains. The clustering matched

the type of deficiency rather than the source of data. Indeed, while the spectra of MMR-deficient strains

from LTEE slightly differed from those of MA strains which had accumulated mutations over a short

period of time, the profiles in these two groups of MMR-deficient strains were still highly similar.

To reconstruct the signatures characteristic of these mutational processes, we applied three different

approaches. The direct approach was to define the mutational signature as the average mutational

profile for all lines (both MA and LTEE) with a given defect. Additionally, we used two different

techniques for *de novo* identification of signatures from a joint dataset of all lineages: non-negative

matrix factorization (NMF) and Bayesian NMF (Materials and Methods). All three methods produced

similar results (Fig 1b, Supplementary Figure 3, Supplementary Figure 4). The consistency between the

direct and the *de novo* approaches provides, to our knowledge, the first direct experimental validation

for the *de novo* approach. For further analysis, we used the signatures obtained directly from

experimental data, assuming them to be the most reliable.

To ask how adequately the obtained signatures describe mutations observed in the mutators, we

decomposed mutational profiles for each hypermutator strain into the signatures and then reconstituted

the profiles. For 18 out of 21 lineages, the reconstructed profile had >90% cosine similarity with the

original profile, and in the remaining cases, the similarity exceeded 80% (Fig 1c). The main process

was correctly identified in all experimental setups (Fig 1d).

Accumulation of mutations in non-mutator laboratory strains

To obtain mutational profiles for the strains accumulating mutations at a normal rate in laboratory

experiments, we considered non-mutator phenotypes from the LTEE (two strains) and Foster (three

strains) experiments (Materials and Methods). In total, these experiments comprised 1877 mutations

across the genome, of which 40% were C>T transitions (Supplementary Figure 5a; Supplementary

Table 1). The observed spectra differed from all profiles observed in the mutator strains, both for the

whole genome and when we only considered intergenic regions (Supplementary Figure 5a,b).

Accumulation of mutations in both LTEE mutators and MA lines was shown to be mostly driven by

mutational bias and not affected by selection[35], while in non-mutator strains, the whole genome

mutational profile differed substantially from that of intergenic regions; specifically, it comprised

considerably fewer T>C transitions (Supplementary Figure 5; Supplementary Table 1). The difference

between the mutational profiles of coding and intergenic regions arises due to stronger selection and/or

the activity of MMR in the coding regions. To control for these differences in our comparisons with

mutator strains, for further analysis of non-mutators, we considered only 375 mutations observed in

intergenic regions. This means that we were unable to study the contribution of MMR to mutation

accumulation in non-mutator strains.

Accumulation of mutations during the divergence of natural *E. coli* lineages

Finally, we considered mutational processes affecting natural populations of *E. coli*. For this, we

collected a dataset of 522 complete *E. coli* genomes, reconstructed their phylogeny based on a set of

single-copy universal genes, and selected four major clades each containing at least 40 genomes (see

Materials and Methods), for a total of 470 genomes. These clades roughly corresponded to the

established phylotypes A, B1, B2, and E (Figure 2; Supplementary Table 2). For each clade, we

reconstructed the whole-genome alignment and inferred single-nucleotide substitutions at all

phylogenetic branches with the baseml program of the PAML package [36]. To minimize the effects of

selection, we only considered substitutions in intergenic regions with more than 20000 substitution per

phylogroup (Supplementary Table 1). To additionally confirm that the reconstructed mutational

profiles are minimally affected by selection, we also separately analyzed intergenic regions between

convergent genes (Supplementary Table 1), expected to be under weaker selection because they contain

fewer regulatory sequences[33]. The reconstructed profiles were very similar between the considered

clades, as were the profiles for all intergenic regions and intergenic regions of convergent genes

(Supplementary Figure 6). In further analysis, we used the profiles built for all intergenic regions.

Laboratory non-mutator strains and natural strains have similar mutational

profiles

We find that the mutational profiles of intergenic regions of diverging natural *E. coli* lineages

are very similar to those of lab-based non-mutator strains (Figure 3a). This indicates that accumulation

of mutations in the course of evolution is driven by the same mutational processes as that observed in

non-mutator strains during laboratory MA experiments. To better understand the processes responsible

for mutation accumulation in these two settings, we decomposed the mutational profiles into a set of

six signatures of six distinct mutational processes reconstructed from the mutator data.

Both the laboratory non-mutator strains and the natural lineages are rather well described by a

combination of all six mutational processes, as indicated by the >86% cosine similarity between the

actual mutational signature and its best reconstruction from these six processes (Figure 3b,

Supplementary figure 7). The "missing" similarity in the natural lineages is very close to that in the

laboratory settings and suggests that external and internal factors that are present in the natural

environment of *E. coli* are present in laboratory conditions as well.

Consideration of contribution of each mutational process revealed that the predominant process causing

mutations is associated with the DnaQ signature, which covered 59% (57%–61%) of all explained

mutations in natural lineages and 56% (46%–65%) in laboratory non-mutators, suggesting a strong

contribution of the major replicative DNA polymerase III. The remaining variability, comprising ~45%

of explained mutations, is explained by the PolB signature, indicating the contribution of the replicative

DNA polymerase II. There was no signature of defective BER endonucleases III and VIII (Nth and

Nei), MutT, or MutY (Figure 3c). As expected, we also saw zero contribution of the MMR mutational

signature to both datasets (not shown in Figure 3d), as we considered non-coding regions, and MMR is

only active in transcribed regions.

The inferred patterns were very similar between all four *E. coli* clades, indicating that they are

characteristic of the overall *E. coli* evolution rather than lineage-specific peculiarities. The results were

stable across bootstrap replicates, and robust when we considered only intergenic regions between

convergent genes (Supplementary Figure 8). Additionally, we verified that the observed differences in

the mutational profiles of natural strains and laboratory non-mutators were not an artifact of the

ancestral state reconstruction by baseml. For that, counts for all identical mutations that had occurred at

a particular position multiple times within a clade were set to one, e.g. recurrent identical substitutions

at the same site were ignored. The resulting contributions of mutational signatures were almost

identical to the ones described above for the full dataset (Supplementary Figure 9).

Discussion

The properties of mutagenesis shape evolution, determining the load of deleterious mutations and

channeling adaptation to new conditions. While the main processes resulting in single-nucleotide

mutations are rather well understood, their relative contributions to bacterial evolution are largely

unknown. In particular, it is not clear whether experimental laboratory conditions can adequately model

mutagenesis in the complex natural environments experienced by bacterial lineages throughout their

evolution. To address this question, we combine experimentally inferred data on mutagenesis with the

mutational patterns revealed by lab MA experiments and with the reconstructed substitutions

accumulated over thousands of years in evolving *E. coli* lineages.

This approach has several limitations. First, experimental studies of mutagenesis in bacteria are in their

infancy. While tens of mutational signatures have been assigned to particular biochemical processes in

higher eukaryotes such as humans[25], the twelve mutational processes have been characterized in

bacteria, but only six single-gene knockouts have sufficient data to be accurately described. The fact

that the decomposition of the accumulated mutations into mutational signatures in our analysis (~86%

cosine similarity) is poorer than that in e.g. cancer genomics studies (>90%)[22] may be partially

explained by this deficiency. While our study reliably identifies the relative contributions of the studied

mutational processes, knowledge of new mutational processes will further clarify the picture.

Second, inference of mutagenesis from MA lines and divergence can be confounded by selection.

Bacteria have compact genomes, more than 80% of which are coding, meaning that the bulk of a

genome is subject to selection; to a weaker degree, selection also affects intergenic regions[33,34]. While

polymorphism and divergence data can be used to infer mutation patterns[37–39], the effect of selection on

the substitutions that distinguish long-term-evolved lineages is expected to be stronger than that in MA

lines (where it is very weak[35]), as these mutations have survived within populations for a longer time

and selection has had more opportunity to act. To mitigate this problem, we focused on genomic

regions that are predicted to be least affected by selection. Most of our analyses were performed on

intergenic regions, with the side effect that we were unable to study the contribution of the MMR

system that mostly acts within transcribed regions. We also considered separately the regions between

convergent genes, where the effect of selection was minimal[33]. The results were the same as for all

intergenic regions, indicating that the effect of selection was low.

Third, our divergence analysis relies on correctness of the inferred phylogeny and the ancestral state.

Systematic biases at these stages of the analysis could distort the inference of mutational spectra.

However, the phylogenetic distances between the compared strains were relatively low (Fig. 2), and an

alternative method of dealing with recurrent substitutions at a site yielded very similar results,

indicating that our results were robust in this respect.

Given these limitations, we described the mutational signatures for six determinants of spontaneous

mutagenesis in *E. coli*. Accounting for mutational contexts allowed us to quantitatively assess the

causes of spontaneous mutations in the studied bacterial strains. Decomposing the spectra of mutations

accumulated in the wild-type non-mutator strains in the laboratory, and in the natural environments

over the course of evolution, into the signatures of six mutational processes allowed us to infer the

prevalent forces responsible for mutagenesis.

We find that the mutational spectra of reconstructed substitutions that have occurred in the course of

evolution of natural *E. coli* lineages and in laboratory non-mutator *E. coli* MA strains were similar to

each other. This observation argues against the notion that *E. coli* evolution in the natural environment

involves alternating episodes of accelerated evolution in the mutator regime and slow evolution in the

non-mutator regime, suggesting instead that at the scale of evolution of different *E. coli* strains, the

contribution of atypical mutational regimes to the overall mutation accumulation is minor. This is

probably due to the high load of deleterious mutations associated with such a regime[35]. Long-term

evolution at the level of bacterial species, which involves gains and losses of repair-associated genes,

could follow a different pattern[6].

The observed spectra are similar to those associated with disruption of DnaQ and PolB. DnaQ is a core

epsilon subunit of DNA polymerase III that is required for proof-reading and is essential for accurate

DNA replication in *E. coli*[40–42]. PolB is a supplementary DNA polymerase II; while it can be induced by

the SOS response[43,44], it is also thought to be involved in DNA repair and to play an auxiliary role in

chromosomal DNA replication under normal conditions[45]. The fact that the mutational spectra

characterizing real-life evolution largely coincides with those of the enzymes involved in replication-

associated repair indicates that the accumulated mutations occur co-replicatively, and suggests that they

are associated with insufficient efficiency of these two repair enzymes during normal replication.

The near-identical contributions of PolB and DnaQ signatures to both natural *E. coli* lineages and

laboratory non-mutator *E. coli* MA strains supports the notion that the bulk of the mutations associated

with these two signatures occur co-replicatively. Indeed, if one of these components had a substantial

fraction of non-co-replicative mutations, we would expect a higher contribution of the corresponding

enzyme to the mutation spectra of natural strains, and lower, to the laboratory strains. This is because

laboratory strains have a much higher replication rate than natural strains (20 minutes vs. 15 hours per

generation[46]), and the proportion of co-replicative vs. "clock-like" mutations in the former should be

higher.

Surprisingly, no contribution of incorrect removal of 8-oxo-G, caused by ionizing radiation or

oxidative metabolism, was observed in either setting, indicating that the time spent by bacteria in

oxidized conditions is probably low.

In summary, while it has been disputed whether evolutionary experiments in monoculture bacteria

adequately represent the real evolution in nature[47], when the complexity of the real world is removed, it

appears that such experiments in *E. coli* can very well represent mutation accumulation in natural

environments. It would be interesting to expand this analysis to different evolutionary settings and

other bacterial species.

Materials and Methods

Bacterial strains

We obtained the wild-type (WT) strain of *E. coli* K-12 MG1655 from the Coli Genetic Stock Centre

(CGSC, Yale University), streaked it on LB (Luria Bertani) agar, and chose one colony at random as

the WT ancestor for subsequent experiments. We similarly obtained the mutator strains of *E. coli*

(ΔmutH, ΔmutL, ΔmutS) from the Keio collection (BW25113 strain background)[48] of gene knockouts

from the same stock centre. These gene knockouts were made by replacing open reading frames with a

Kanamycin resistance cassette, such that removing the cassette generates an in-frame deletion of the

gene. The design of gene deletion primers ensured that downstream genes were not disrupted due to

polar effects. For each strain, we moved the knockout loci from the BW25113 background into the

MG1655 (WT) background using P1-phage transduction[49]. We then removed the kanamycin resistance

marker by transforming kanamycin-resistant transductants with pCP20, a plasmid carrying the flippase

recombination gene and Amp^R resistance marker. We grew ampicillin resistant transformants at 42 ºC

in LB broth overnight to cure pCP20. We streaked out 10 µL of these cultures on LB plates. After 24

hours, we replica-plated several colonies on both kanamycin-LB agar plates and ampicillin-LB agar

plates, to screen for the loss of both kanamycin and ampicillin resistance. We PCR-sequenced the

knockout locus to confirm removal of the kanamycin cassette.

Experimental evolution under mutation accumulation (MA) and whole-genome sequencing to identify mutations

For each mutator strain, we founded 20 MA lines from a single ancestral colony (two lines per Petri
plate), incubated at 37ºC, as described earlier for wild-type MA[50]. For each line, every 24 hours we
streaked out a random colony (closest to a pre-marked spot) on a fresh LB agar plate. Every 4-5 days,
we inoculated a part of the transferred colony in LB broth at 37ºC for 2-3 hours and froze 1 mL of the
growing culture with 8% DMSO at -80 ºC. For the current study, we used stocks frozen on day 50
(~1375 generations).

For each strain, we sequenced whole genomes from the last stored colony of each lineage as follows.
We inoculated 2 µL of the frozen stock of each evolved MA line (or the ancestor) in 2 mL LB, and
allowed the cells to grow overnight at 37ºC with shaking at 200 rpm. We extracted genomic DNA
(GenElute Bacterial Genomic DNA kit, Sigma-Aldrich), quantified it (Qubit HS dsDNA assay,
Invitrogen), and pooled equal amounts of genomic DNA from each of the 20 lines for a given mutator
strain into a single tube. Thus, we prepared 3 pools of genomic DNA, one each for the mutators
Δ*mutH,* Δ*mutL,* and Δ*mutS,* each containing an equal amount of genomic DNA from each of the 20
MA lines. We then prepared paired-end libraries from the Δ*mutH,* Δ*mutL* and Δ*mutS* MA ancestors and
the three pooled genomic DNA samples using the Illumina Nextera XT DNA library preparation kit per
the manufacturer's instructions. We sequenced all libraries on the Illumina MiSeq platform using the
2x250bp paired-end V2 reaction chemistry. For the Δ*mutH,* Δ*mutL* and Δ*mutS* MA ancestors, we
obtained 1.14M, 1.49 M and 1.06 paired end reads with quality > Q30 respectively. For the pooled
evolved lines, we obtained 2.09M, 2.18M, and 2.35M paired-end reads respectively with quality >
Q30, corresponding to an average per base coverage of ~ 116x, 121x, and ~130x. For each sample, we
aligned quality-filtered reads to the NCBI reference *E. coli* K-12 MG1655 genome (RefSeq accession
ID GCA_000005845.2) using the Burrows-Wheeler short-read alignment tool. We generated pileup
files using SAMtools[51] and used VARSCAN to extract a list of base-pair substitutions and short indels
(<10bp)[52]. For the MA ancestors, we only retained mutations with >80% frequency that were
represented by at least 5 reads on both strands for further analysis. For the pooled evolved MA lines,
we retained mutations with > 5% frequency that were represented by at least 3 reads on both strands.
After removing ancestral mutations (i.e. mutations that differentiated the ancestors from the reference
E. coli genome) from the evolved lines, we identified 985, 875 and 768 SNPs in ΔmutH, ΔmutL and
ΔmutS evolved MA lines respectively.

Hypermutator data published earlier

We considered three main sources of data. (1) Sequences of six mutator strains from the LTEE
experiment: Ara-1 and Ara+6 with MutT deficiency and Ara-2, Ara-3, Ara-4, and Ara+3 with
mutations in MMR-associated genes[4]. The data for this experiment were downloaded from
http://barricklab.org/shiny/LTEE-Ecoli/. Only SNPs were selected from the obtained file. Additionally,
mutations observed at different time points for the same strain were merged to avoid repeated counting
of mutations that had occurred early in the experiment. The obtained table was converted to the vcf
format. (2) Experiments on MA strains for all major mutational processes from Patricia L. Foster's
laboratory[1,3,5]. In case of data from refs. [1,3] we obtained information about mutations from the
Supplementary Materials and converted it to the vcf format. To obtain mutations caused by inactivated
DnaQ[5], we downloaded sequencing data from SRA (SRA IDs: SRR7748246, SRR7748247,
SRR7748248, SRR7748249, SRR7748253, SRR7748306, SRR7748326, SRR7748342, SRR7748539,
SRR7748540, SRR7748541, SRR7748542, SRR7748543, SRR7748592, SRR7748593, SRR7748594,
SRR7748595, SRR7748733, SRR7748734, SRR7749018, SRR7749020, SRR7749022, SRR7749036,
SRR7749037, SRR7749062, SRR7749108). The obtained reads were trimmed with Trimmomatic-
0.39[53] with parameters LEADING:3 TRAILING:3 MINLEN:36. The trimmed paired reads were
processed as described in the original publication. The variants were called with mpileup from bcf-
tools with the minimal sequencing depth of at least 20 reads. In total, we obtained 13512 SNPs of
which four occurred more than once and were discarded, yielding 13504 mutations.

Laboratory strain data

We used non-mutator data from LTEE[4], and experiments from Patricia L. Foster's laboratory[3]. All
LTEE non-mutator strains were merged together for further analysis. Duplicated mutations were
filtered out to avoid selection effects. We considered separately all available mutations and mutations
in intergenic regions.

Reconstruction of the *E. coli* phylogeny

Complete genomes of 522 *E. coli* strains were downloaded from Genbank[54]. The complete list of
genomes is provided in Supplementary Table 1. The initial set of orthology groups (OGs) from 32
phylogenetically diverse *E. coli* and *Shigella* strains was taken from[55]. We produced alignments for
each universal group (OGs with single-copy genes present in all analyzed genomes) with ClustalW
version 2.1[56]. For each alignment, we generated a consensus sequence with the EMBOSS package ver.

6.6.0 and mapped these consensuses to the genomes with bowtie2[57]. All groups that remained universal

and single copy (690 genes) were realigned. Alignments for individual genes were concatenated and all

columns with gaps were removed. This final alignment was used to construct the phylogenetic tree with

PhyML v3.3 with the GTR substitution model[58].

Genome alignments and reconstruction of the ancestral states

Based on the *E. coli* phylogenetic tree, we selected four clades that contained at least 45 genomes. For

the genomes where phylogroup information was available (shown as colored leaves in Figure 2), we

checked that genomes within the clade belong to the same *E. coli* phylogroup.

For each of the selected clades, we constructed genome alignments with ProgressiveCactus[59]. For each

clade, we selected one representative genome that was then used to identify gene boundaries and select

intergenic regions between convergent genes (Supplementary Table 1). The output format was

converted to maf, with the representative genome serving as the reference, using the hal2maf tool from

ProgressiveCactus with parameters: --noDupes --onlySequenceNames --noAncestors --maxBlockLen

1000000. From this alignment, we extracted all variable columns in intergenic regions that (i)

contained at least five genomes and (ii) had conserved columns to their left and right. Variable columns

with adjacent variable columns or columns with gaps were discarded.

For these positions, we reconstructed the ancestral state with the baseml program from the PAML

package v4.9j[36].

Reconstruction of mutational profiles in natural strains

From the PAML results (rst), we obtained the number of substitutions in the variable column for each

site and then pooled this for all sites for each context within the clade.

We calculated the number of substitutions in all intergenic regions, and separately in intergenic regions

between convergent genes. In order to make results compatible between different clades, the number of

observed mutations in each context was normalized by the total number of mutations observed for the

clade. To control for possible incorrect reconstruction of ancestral states by baseml, in an alternative

procedure, if there were multiple identical (parallel) substitutions at the given site for the particular

clade, we counted them as just one substitution in mutation counts.

Mutational signatures

For each available experiment we calculated the contexts based on the initial strains in which the experiment was performed. Mutational signatures reconstructed from the data were calculated as the average normalized contribution of a particular substitution among all available mutator strains with the given deficiency. The reconstruction of mutational signatures was performed in R ver 4.1.1. To reconstruct signatures with the NMF approach, we used the NMF package in R. The Bayesian NMF approach was implemented with the ccfindR package. The scripts to reproduce this analysis are available on github (see Data Availability).

Deconvolution of obtained mutational profiles by mutational signatures was performed with the MutationalPatterns package[60].

Authors contributions

SKG and MSG conceived the project. MS performed the MA experiments, sequencing and variant calling from the data. SKG and MVS reconstructed mutational signatures from experimental data. SKG performed data analysis. DA, GAB and MSG supervised the research. SKG, GAB, and MSG wrote the manuscript with contributions from all authors.

Acknowledgements

The project was initiated with Arina Kolotova at the Summer School of Molecular and Theoretical Biology for high school students. We thank our colleagues, in particular Dr. Anna Kaznadzey (IITP RAS) and Dr. Olga Vakhrusheva (Skoltech), for useful discussions and suggestions. We acknowledge help and support from the Next-generation genomics facility at the National Centre for Biological Sciences (NCBS) for whole-genome sequencing of bacterial strains. This work was supported by the Russian Science Foundation (18-14-00358), the Council for Scientific and Industrial Research (CSIR-India; senior research fellowship to MS), the National Centre for Biological Sciences (NCBS-TIFR), and the Department of Atomic Energy, Government of India (Project Identification No. RTI 4006).

Data availability

All data and scripts required for the analysis of data are deposited in github: https://github.com/garushyants/Ecoli_MutationalSignatures

References

1. Lee, H., Popodi, E., Tang, H. & Foster, P. L. Rate and molecular spectrum of spontaneous mutations in the bacterium Escherichia coli as determined by whole-genome sequencing. *Proc. Natl. Acad. Sci.* **109**, E2774–E2783 (2012).

2. Foster, P. L., Hanson, A. J., Lee, H., Popodi, E. M. & Tang, H. On the Mutational Topology of the Bacterial Genome. *G3 Genes Genomes Genet.* **3**, 399–407 (2013).

3. Foster, P. L., Lee, H., Popodi, E., Townes, J. P. & Tang, H. Determinants of spontaneous mutation in the bacterium Escherichia coli as revealed by whole-genome sequencing. *Proc. Natl. Acad. Sci.* **112**, E5990–E5999 (2015).

4. Tenaillon, O. *et al.* Tempo and mode of genome evolution in a 50,000-generation experiment. *Nature* **536**, 165–170 (2016).

5. Niccum, B. A., Lee, H., MohammedIsmail, W., Tang, H. & Foster, P. L. The Spectrum of Replication Errors in the Absence of Error Correction Assayed Across the Whole Genome of Escherichia coli. *Genetics* **209**, 1043–1054 (2018).

6. Sane, M., Diwan, G. D., Bhat, B. A., Wahl, L. M. & Agashe, D. Shifts in mutation spectra enhance access to beneficial mutations. 2020.09.05.284158 Preprint at https://doi.org/10.1101/2020.09.05.284158 (2022).

7. Miller, J. H. SPONTANEOUS MUTATORS IN BACTERIA: Insights into Pathways of Mutagenesis and Repair. *Annu. Rev. Microbiol.* **50**, 625–643 (1996).

8. Nik-Zainal, S. *et al.* Landscape of somatic mutations in 560 breast cancer whole-genome sequences. *Nature* **534**, 47–54 (2016).

9. Andrianova, M. A., Bazykin, G. A., Nikolaev, S. I. & Seplyarskiy, V. B. Human mismatch repair system balances mutation rates between strands by removing more mismatches from the lagging strand. *Genome Res.* (2017) doi:10.1101/gr.219915.116.

10. Katju, V. & Bergthorsson, U. Old Trade, New Tricks: Insights into the Spontaneous Mutation

Process from the Partnering of Classical Mutation Accumulation Experiments with High-

Throughput Genomic Approaches. *Genome Biol. Evol.* **11**, 136–165 (2019).

11. Sniegowski, P. D., Gerrish, P. J. & Lenski, R. E. Evolution of high mutation rates in experimental

populations of E. coli. *Nature* **387**, 703 (1997).

12. Gentile, C. F., Yu Szi-Chieh, Serrano Sebastian Akle, Gerrish Philip J., & Sniegowski Paul D.

Competition between high- and higher-mutating strains of Escherichia coli. *Biol. Lett.* **7**, 422–424

(2011).

13. Sprouffske, K., Aguilar-Rodríguez, J., Sniegowski, P. & Wagner, A. High mutation rates limit

evolutionary adaptation in Escherichia coli. *PLOS Genet.* **14**, e1007324 (2018).

14. Cox, E. C. & Horner, D. L. Dominant mutators in Escherichia coli. *Genetics* **100**, 7–18 (1982).

15. Gautam, S., Kalidindi, R. & Humayun, M. Z. SOS induction and mutagenesis by dnaQ missense

alleles in wild type cells. *Mutat. Res. Mol. Mech. Mutagen.* **735**, 46–50 (2012).

16. Strauss, B. S., Roberts, R., Francis, L. & Pouryazdanparast, P. Role of the dinB Gene Product in

Spontaneous Mutation in Escherichia coli with an Impaired Replicative Polymerase. *J. Bacteriol.*

182, 6742–6750 (2000).

17. Bhamre, S., Gadea, B. B., Koyama, C. A., White, S. J. & Fowler, R. G. An aerobic recA-, umuC-

dependent pathway of spontaneous base-pair substitution mutagenesis in Escherichia coli. *Mutat.*

Res. Mol. Mech. Mutagen. **473**, 229–247 (2001).

18. Sanders, L. H., Rockel, A., Lu, H., Wozniak, D. J. & Sutton, M. D. Role of Pseudomonas

aeruginosa dinB-Encoded DNA Polymerase IV in Mutagenesis. *J. Bacteriol.* **188**, 8573–8585

(2006).

19. Tajiri, T., Maki, H. & Sekiguchi, M. Functional cooperation of MutT, MutM and MutY proteins in

preventing mutations caused by spontaneous oxidation of guanine nucleotide in Escherichia coli.

Mutat. Res. **336**, 257–267 (1995).

20. Fowler, R. G. *et al.* Interactions among the Escherichia coli mutT, mutM, and mutY damage

prevention pathways. *DNA Repair* **2**, 159–173 (2003).

21. Nik-Zainal, S. *et al.* Mutational Processes Molding the Genomes of 21 Breast Cancers. *Cell* **149**,

979–993 (2012).

22. Alexandrov, L. B. *et al.* Signatures of mutational processes in human cancer. *Nature* **500**, 415–421

(2013).

23. Helleday, T., Eshtad, S. & Nik-Zainal, S. Mechanisms underlying mutational signatures in human

cancers. *Nat. Rev. Genet.* **15**, 585–598 (2014).

24. Seplyarskiy, V. B., Andrianova, M. A. & Bazykin, G. A. APOBEC3A/B-induced mutagenesis is

responsible for 20% of heritable mutations in the TpCpW context. *Genome Res.* **27**, 175–184

(2017).

25. Alexandrov, L. B. *et al.* The repertoire of mutational signatures in human cancer. *Nature* **578**, 94–

101 (2020).

26. Niccum, B. A., Lee, H., MohammedIsmail, W., Tang, H. & Foster, P. L. The Symmetrical Wave

Pattern of Base-Pair Substitution Rates across the Escherichia coli Chromosome Has Multiple

Causes. *mBio* **10**, (2019).

27. Sung, W. *et al.* Asymmetric Context-Dependent Mutation Patterns Revealed through Mutation–

Accumulation Experiments. *Mol. Biol. Evol.* **32**, 1672–1683 (2015).

28. Schroeder, J. W., Hirst, W. G., Szewczyk, G. A. & Simmons, L. A. The Effect of Local Sequence

Context on Mutational Bias of Genes Encoded on the Leading and Lagging Strands. *Curr. Biol.* **26**,

692–697 (2016).

29. Schroeder, J. W., Randall, J. R., Hirst, W. G., O'Donnell, M. E. & Simmons, L. A. Mutagenic cost

of ribonucleotides in bacterial DNA. *Proc. Natl. Acad. Sci.* **114**, 11733–11738 (2017).

30. Good, B. H., McDonald, M. J., Barrick, J. E., Lenski, R. E. & Desai, M. M. The dynamics of

molecular evolution over 60,000 generations. *Nature* **551**, 45–50 (2017).

31.   Thorpe, H. A., Bayliss, S. C., Hurst, L. D. & Feil, E. J. Comparative Analyses of Selection

Operating on Nontranslated Intergenic Regions of Diverse Bacterial Species. *Genetics* **206**, 363–

376 (2017).

32.   Rocha, E. P. C. Neutral Theory, Microbial Practice: Challenges in Bacterial Population Genetics.

Mol. Biol. Evol. **35**, 1338–1347 (2018).

33.   Tsoy, O. V., Pyatnitskiy, M. A., Kazanov, M. D. & Gelfand, M. S. Evolution of transcriptional

regulation in closely related bacteria. *BMC Evol. Biol.* **12**, 200 (2012).

34.   Shelyakin, P. V., Bochkareva, O. O., Karan, A. A. & Gelfand, M. S. Micro-evolution of three

Streptococcus species: selection, antigenic variation, and horizontal gene inflow. *BMC Evol. Biol.*

19, 83 (2019).

35.   Couce, A. *et al.* Mutator genomes decay, despite sustained fitness gains, in a long-term experiment

with bacteria. *Proc. Natl. Acad. Sci.* **114**, E9026–E9035 (2017).

36.   Yang, Z. PAML 4: Phylogenetic Analysis by Maximum Likelihood. *Mol. Biol. Evol.* **24**, 1586–

1591 (2007).

37.   Seplyarskiy, V. B., Kharchenko, P., Kondrashov, A. S. & Bazykin, G. A. Heterogeneity of the

transition/transversion ratio in Drosophila and Hominidae genomes. *Mol. Biol. Evol.* **29**, 1943–

1955 (2012).

38.   Terekhanova, N. V., Seplyarskiy, V. B., Soldatov, R. A. & Bazykin, G. A. Evolution of Local

Mutation Rate and Its Determinants. *Mol. Biol. Evol.* **34**, 1100–1109 (2017).

39.   Harris, K. & Pritchard, J. K. Rapid evolution of the human mutation spectrum. *eLife* **6**, e24284

(2017).

40.   Scheuermann, R. H. & Echols, H. A separate editing exonuclease for DNA replication: the epsilon

subunit of Escherichia coli DNA polymerase III holoenzyme. *Proc. Natl. Acad. Sci. U. S. A.* **81**,

7747–7751 (1984).

41.   Dodd, T. *et al.* Polymerization and editing modes of a high-fidelity DNA polymerase are linked by

529 a well-defined path. *Nat. Commun.* **11**, 5379 (2020).

530 42. Stefan, A. *et al.* Silencing of the gene coding for the ϵ subunit of DNA polymerase III slows down

531 the growth rate of Escherichia coli populations. *FEBS Lett.* **546**, 295–299 (2003).

532 43. Rangarajan, S., Woodgate, R. & Goodman, M. F. A phenotype for enigmatic DNA polymerase II:

533 A pivotal role for pol II in replication restart in UV-irradiated Escherichia coli. *Proc. Natl. Acad.*

534 *Sci.* **96**, 9224–9229 (1999).

535 44. Whatley, Z. & Kreuzer, K. N. Mutations that Separate the Functions of the Proofreading Subunit of

536 the Escherichia coli Replicase. *G3 GenesGenomesGenetics* **5**, 1301–1311 (2015).

537 45. Banach-Orlowska, M., Fijalkowska, I. J., Schaaper, R. M. & Jonczyk, P. DNA polymerase II as a

538 fidelity factor in chromosomal DNA synthesis in Escherichia coli. *Mol. Microbiol.* **58**, 61–70

539 (2005).

540 46. Gibson, B., Wilson, D. J., Feil, E. & Eyre-Walker, A. The distribution of bacterial doubling times

541 in the wild. *Proc. R. Soc. B Biol. Sci.* **285**, 20180789 (2018).

542 47. Lenski, R. E. Experimental evolution and the dynamics of adaptation and genome evolution in

543 microbial populations. *ISME J.* **11**, 2181–2194 (2017).

544 48. Baba, T. *et al.* Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the

545 Keio collection. *Mol. Syst. Biol.* **2**, 2006.0008 (2006).

546 49. Thomason, L. C., Costantino, N. & Court, D. L. E. coli Genome Manipulation by P1 Transduction.

547 *Curr. Protoc. Mol. Biol.* **79**, 1.17.1-1.17.8 (2007).

548 50. Sane, M., Miranda, J. J. & Agashe, D. Antagonistic pleiotropy for carbon use is rare in new

549 mutations. *Evolution* **72**, 2202–2213 (2018).

550 51. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinforma. Oxf. Engl.* **25**,

551 2078–2079 (2009).

552 52. Koboldt, D. C. *et al.* VarScan: variant detection in massively parallel sequencing of individual and

553 pooled samples. *Bioinformatics* **25**, 2283–2285 (2009).

53. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence

data. *Bioinformatics* **30**, 2114–2120 (2014).

54. Benson, D. A. *et al.* GenBank. *Nucleic Acids Res.* **46**, D41–D47 (2018).

55. Gordienko, E. N., Kazanov, M. D. & Gelfand, M. S. Evolution of Pan-Genomes of Escherichia

coli, Shigella spp., and Salmonella enterica. *J. Bacteriol.* **195**, 2786–2792 (2013).

56. Larkin, M. A. *et al.* Clustal W and Clustal X version 2.0. *Bioinforma. Oxf. Engl.* **23**, 2947–2948

(2007).

57. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**, 357–

359 (2012).

58. Guindon, S. *et al.* New Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies:

Assessing the Performance of PhyML 3.0. *Syst. Biol.* **59**, 307–321 (2010).

59. Armstrong, J. *et al.* Progressive Cactus is a multiple-genome aligner for the thousand-genome era.

Nature **587**, 246–251 (2020).

60. Blokzijl, F., Janssen, R., van Boxtel, R. & Cuppen, E. MutationalPatterns: comprehensive genome-

wide analysis of mutational processes. *Genome Med.* **10**, 33 (2018).

Figure 1. Reconstruction and validation of mutational signatures from the mutator data. (a) Cosine similarity matrix between 3-letter mutational profiles for various mutator strains. (b) Signatures of mutational processes reconstructed from the mutator strains (see also Supplementary Figure 4). (c) Similarity between the original and reconstructed profiles for the mutator strains. (d) Signatures of mutational processes are correctly assigned to the mutator strains. The size of the circle represents the mean contribution of the mutational signature in the sample, and the color of the circle represents the percentage of bootstrap replicates where this signature was observed.

Figure 2. Phylogenetic tree of 522 complete *E. coli* genomes. The maximum likelihood tree was reconstructed from the concatenated alignment of universal single-copy orthologous genes (see Material and Methods). The four analyzed clades, A (yellow), B1 (green), B2 (dark green) and E (blue), are also indicated with grey arcs. Leaves for which the phylogroup is indicated in metadata are colored accordingly. Black leaves mean that no phylogroup information is available.

586
Figure 3. Contribution of mutational signatures to the evolution of intergenic regions in *E. coli* natural
and in non-mutator laboratory strains. (a) Mutational profile for non-mutator strains and clade B1.
(Profiles for all clades are shown in Supplementary Figure 6.) (b) Cosine similarity between the
original mutational profiles and the reconstructed profiles. The horizontal line is at 0.85. (c) Relative
contribution of each signature to the reconstructed profile. Each dot is the value obtained in one
bootstrap replicate. Dots are colored by the data type as in (b).
593