Early life exposure to cigarette smoke is associated with lung function changes and primes gene expression and DNA methylation changes at *Cyp1a1* upon exposure later in life

Chinonye Doris Onuzulu[1], Samantha Lee[1], Sujata Basu[2], Jeannette Comte[2], Yan Hai[1], Nikho Hizon[1], Shivam Chadha[1], Maria Shenna Fauni[1], Shana Kahnamoui[2,3], Bo Xiang[2,4], Andrew J. Halayko[2,3], Vernon W. Dolinsky[2,4], Christopher Pascoe[2,3], Meaghan J. Jones[1,2]

(1) Department of Biochemistry and Medical Genetics, University of Manitoba; (2) Children's Hospital Research Institute of Manitoba; (3) Department of Physiology and Pathophysiology; (4) Department of Pharmacology and Therapeutics, University of Manitoba

Abstract

Prenatal and early life exposure to cigarette smoke (CS) have repeatedly been shown to induce stable, long-term changes in DNA methylation (DNAm) in offspring. It has been hypothesized that these changes might be functionally related to the known outcomes of prenatal and early life CS exposure, which include impaired lung development, altered lung function and increased risk of asthma and wheeze. However, to date, few studies have examined DNAm changes induced by prenatal CS in tissues of the lung, and even fewer have attempted to examine the specific influences of prenatal versus early postnatal exposures. Here, we have established a mouse model of CS exposure which isolates the effects of prenatal and postnatal CS exposures in early life. We have used this model to measure the effects of prenatal and/or postnatal exposures on lung function and immune cell infiltration as well as DNAm and expression of *Cyp1a1*, a candidate gene previously observed to demonstrate DNAm differences upon CS exposure in humans. Our study revealed that exposure to CS prenatally and in the early postnatal period cause distinct and long-lasting differences in offspring lung function, gene expression and lung *Cyp1a1* DNAm, which are reestablished upon re-exposure to CS in

adulthood. This study creates a testable mouse model which can be used to investigate the effects of

prenatal and early postnatal CS exposures, and will contribute to the design of intervention strategies

to mediate these detrimental effects.

Introduction

The developmental origins of health and disease (DOHaD) hypothesis postulates that environmental

insults in the intrauterine and early postnatal environments program the developing fetus through

changes in cellular function and structure, and that these changes can last through adulthood[1,2].

Numerous studies have shown that factors such as maternal nutrition, infections, stress and

environmental exposure to toxins such as cigarette smoke (CS) can in fact produce long-lasting health

consequences in offspring[3,4]. CS is a common environmental toxin in humans with about 9.5% of

women of reproductive age (18-34 years) reported as active smokers in Canada in 2021[5]. Cigarette

smoke is made up of thousands of components which have detrimental effects, some of which can also

cross the placenta and affect offspring development[6–10].

Prenatal CS exposure alters offspring growth and development, lung function, and is associated with

airway hyperresponsiveness, asthma and wheeze in children[11–13]. In some cases, these outcomes persist

into adulthood, suggesting that prenatal CS 'leaves a mark' on the offspring, a process that has been

called *biological embedding*[14–16]. Epigenetic modifications have been identified as potential mechanisms

by which biological embedding takes place[16,17]. The three main components of the epigenome – DNA

methylation (DNAm), histone modifications, and non-coding RNAs[18] - are dynamic and reflect past

environmental exposures, meaning that environmental insults in early life can elicit changes in

epigenetic marks which cells would propagate over time, and thus could translate to disease phenotypes

in later life.

CS exposure *in utero* has been associated with differential DNAm across thousands of genes in exposed children, suggesting that DNAm changes might directly or indirectly link early life CS exposure with later life outcomes[19–21 21–25]. While most epigenome-wide association studies (EWAS) investigating association between maternal smoking and DNAm have been conducted in newborn umbilical cord blood[19–25], others have also reported differential DNAm in whole blood[26], placenta[27], nasal epithelia[28], buccal cells and peripheral blood granulocytes of adults[29,30]. The largest EWAS study to date which investigated effects of prenatal smoke exposure on umbilical cord blood DNAm was a meta-analysis across 13 cohorts (n = 6685) in the Pregnancy and Childhood Epigenetics Consortium (PACE)[20]. Results from the PACE study reported differential methylation at over 6000 CpGs, with some of the top sites being Aryl Hydrocarbon Receptor Repressor (*AHRR*), Cytochrome P450 1A1 (*CYP1A1*), Myosin 1G (*MYO1G*) and Growth Factor Independent 1 Transcriptional Repressor (*GFI1*). These genes have also been associated with maternal smoking in other similar EWAS studies[19,20,23,31]. They are important in detoxification, apoptosis and cell proliferation, and have been linked to the development of orofacial clefts, oncogenesis and asthma[19,20,31]. Existing studies including PACE could not investigate the effects of prenatal smoking after birth, nor could it measure DNAm in tissues that might be more proximal to CS-mediated disease, such as the lungs.

While there have been human studies outlining the long-term effects of prenatal CS exposure on offspring, one challenge of these studies is their inability to effectively account for the effects of the postnatal environment. One study which measured the long-term effects of prenatal CS on offspring lung function found no significant association between second-hand smoke exposure while pregnant and long-term offspring lung function after controlling for postnatal smoke exposure[32]. It is difficult in human studies to completely rule out the effect of postnatal smoke exposure since prenatal and postnatal exposures are typically very similar[33,34]. Therefore, there is a need to investigate the differences between DNAm signatures and health outcomes observed due to prenatal versus postnatal

CS exposure, especially since these exposures have been shown to produce mixed/confounding

effects[35–38]. Animal models provide a solution to these problems by allowing the study of the effects of

prenatal versus postnatal CS exposure, and enable research into longitudinal, cross-tissue and sex-

specific effects of early life CS exposure.

Here, we have created a mouse model to study the effects of prenatal, postnatal, and combined early

life CS on offspring DNAm and lung phenotype. Taken together, our mouse model successfully

recapitulates some of the phenotypes identified in humans exposed to prenatal CS but provided further

information on the dynamics underlying early life CS exposure. We show that early life CS exposure

alters offspring DNAm at *Cyp1a1*, lung function and gene expression in patterns specific to their early

life exposure periods. Critically, we show that the early life period is critical for creating lasting

memories, and that exposure to CS in adulthood recapitulates patterns set by early life CS exposure.

Materials and Methods

Animals

This animal experiment was approved by the Animal Research Ethics and Compliance Committee of

the University of Manitoba. Balb/C mice (Charles River Laboratories, Massachusetts, United States)

were supplied with standard laboratory chow and clean water *ad libitum* and housed, four mice of a

single sex per cage unless otherwise noted, in individually ventilated cages in a 12-hour light/dark

cycle throughout the duration of the experiment.

Development of murine model of early life CS exposure

Standard 1R6F research cigarettes (University of Kentucky, Lexington, KY) were used for this

experiment. Cigarette smoke was delivered via the use of the SCIREQ InExpose smoking robot

(SCIREQ, Montreal, QC, Canada). Female mice were left to acclimate to their environment for 1

week, and then divided into 2 groups: 16 control and 16 CS-exposed mice. Beginning at 8 weeks,

control mice were exposed to room air in a foreign cage, and CS mice were exposed to whole body CS

twice daily. Total estimated particulate matter per the certificate of analysis from the cigarette

manufacturer was 46.8 mg/cigarette, which we administered with a flow rate of 2L/min and one puff

per minute, with approximately 6 puffs per cigarette. CS mice were exposed for a total period of 9

weeks, beginning with a pre-pregnancy period of 3 weeks, after which they were mated (2 females to 1

male) with male mice. Once a vaginal plug was achieved, we separated female mice into individual

cages for the rest of the duration of the experiment. We continued all smoke and room air exposures on

CS and control mice respectively, throughout mating, pregnancy, birth, and stopped at weaning (3

weeks post-birth). At birth, we culled offspring to 6 offspring per litter, and cross-fostered half of CS

offspring with half of control offspring, matching offspring by birth date to reduce potential of

rejection. Pups were marked with a toe tattoo to identify birth group. Using our cross-fostering

strategy, we generated 4 distinct offspring groups: control with no CS exposure ("Con"), offspring

exposed to prenatal CS only ("Pre"), offspring exposed to postnatal CS only ("Post"), and offspring

that received both prenatal and postnatal CS exposures ("Full"). Throughout the experiment, offspring

were never directly exposed to CS, but were indirectly exposed *in utero*, or via breastmilk or fur from

the dams. We sacrificed and collected tissues from all offspring not cross-fostered at birth. Dams and

offspring were weighed weekly. Three weeks after birth, offspring were weaned, CS exposures

stopped, and dams underwent lung function testing followed by tissue collection. Pups were housed by

sex, 4-6 to a cage. While original plans were to assess phenotype at 8 weeks, of age, due to SARS-

CoV2-related lab lockdowns animals were maintained until 16-20 weeks of age, at which point lung

function and tissue collection were conducted.

CS re-exposure experiment

Female offspring which were not euthanized at 16-20 weeks were left undisturbed until 60 weeks of

age. Half of these female adult offspring were re-exposed for 2 hours to full body CS for 5 consecutive

days per week, for 21 days in total. The other half were not re-exposed to CS, and served as the adult

control group. At the end of CS exposure, lung function, lavage and tissue collection were also

performed.

Transthoracic echocardiography

As described previously[39], serial echocardiography was performed using a Vevo 2100 ultrasound

(Fujifilm-Visual Sonics, ON Canada) equipped with a 30 MHz transducer to assess cardiac

morphology and function in 12-14 week-old mice. Imaging by a blinded and experienced sonographer

was performed under mild anesthesia (induced with 3% isoflurane and 1.0L/min oxygen and

maintained at 1 – 1.5% isoflurane and 1.0 L/min oxygen) during echocardiography. Pulse Wave

Doppler Mode was used to measure the velocity time integral of blood flow passing through the right

ventricular outflow tract (RVOT). Vevo software was used to determine pulmonary acceleration times

(PAT), pulmonary ejection times (PET), and ventricle thickness from the RVOT for each subject.

Measurements of PAT and PET were taken from three different doppler waves for each subject and

ventricle thickness was measured from three image stills. The average of these three measurements

was then recorded.

Lung function measurement

As previously described[40,41], mice were anesthetized with sodium pentobarbital and lung function

measured using the SCREQ Flexivent small animal ventilator (SCIREQ Inc., Montreal, QC, Canada).

Total airway resistance (Rrs), Newtonian resistance (Rn), tissue resistance (G) and elastance (H) were

measured at baseline in response to nebulized saline, and changes in these parameters were also

measured in response to increasing concentrations of nebulized methacholine (3 to 50 mg/ml).

Bronchoalveolar lavage

Following lung function measurement, mouse lungs were washed twice through a tracheal cannula with 1 ml of phosphate buffer saline (PBS) each wash. The bronchoalveolar lavage fluid (BALF) was then centrifuged at 4°C at 1200 rpm for 10 minutes to obtain cell-free supernatants which were stored at -20°C for later cytokine analysis. Cell pellets were resuspended in 1 ml of PBS and total cell counts were performed using a hemocytometer. Differential cell counts were performed by placing 100ul of the resuspended pellets on glass slides using cytospin columns, and staining cells using a modified Wright-Giemsa stain (Hema 3 Stat Pack). Cell differentials were counted using a Carl Zeiss Axio Observer ZI microscope.

Blood and tissue collection and preparation

Whole blood was collected from dams at pup weaning and from offspring at 16 and 63 weeks. Blood was collected by severing the abdominal aorta, into EDTA-coated tubes, centrifuged at 4000 rpm for 15 minutes to obtain plasma. Blood cell pellets and plasma were snap-frozen in dry ice and stored at -80°C for future measurements. Left, inferior, postcaval and middle and superior lung lobes were collected separately and snap-frozen in dry ice for DNAm and gene expression analyses.

DNA/RNA isolation from lungs and blood

DNA and RNA were extracted from mouse tissues using the Invitrogen DNA and RNA isolation kits respectively. Left lungs were homogenized using the Qiagen Tissue Lyser II, followed by simultaneous DNA and RNA isolation using same Invitrogen DNA and RNA isolation kits. DNA and RNA were quantified using a NanoDrop spectrophotometer (NanoDrop Technologies, USA).

Maternal cotinine measurement

To test the concentration of nicotine in the dams, dam cotinine levels were measured in plasma 48

hours after smoking cessation at weaning of offspring, and again 13 weeks later in a subset of dams.

Plasma cotinine was measured using a commercially available ELISA kit (CalBiotech, Spring Valley,

CA), according to the manufacturer's instructions, and the concentration of cotinine in each sample

was measured at 450 nm.

Selection of mouse candidate genes to measure DNAm

To obtain candidate genes at which to measure DNAm, we selected 2 of the top differentially

methylated CpGs from the meta-analysis conducted in human cohorts[20]. The 2 chosen human CpGs

with some of the largest effect sizes were selected from genome build GRCh37/hg19: *AHRR* at

position chr5:373378 (cg05575921) and *CYP1A1* at position chr15:75019251 (cg22549041). We then

blasted the resulting human sequence against the *Mus musculus* genome assembly on NCBI. Using R

studio version 3.6, we performed a muscle[42] alignment between human and resulting mouse sequences

to detect mouse CpGs which exactly aligned with or were closest in position to the human CpGs. The

2 mouse CpGs were selected from genome build GRCm38/mm10 based on results of the alignment:

position chr13:74260517 for *Ahrr* and position chr9:57696335 for *Cyp1a1.* Importantly, the *Ahrr* gene

is not conserved between human and mice, and while we selected the closest mouse CpG, it may not

be comparable to the human position. To select a CpG position for the mouse control gene, we

selected a human CpG position which was not differentially methylated in newborn cord blood (human

genome GRCh38/hg38, *PRKAA1* cg13345558, position chr5:40796738), as reported in the PACE

study. We then identified a corresponding mouse CpG (genome build GRCm39/mm39 position

chr15:5174566) as described above.

Measurement of the effects of cross-fostering

Since we cross-fostered offspring at birth to create our model for early life CS exposure, we needed to

rule out any potential effects of the cross-fostering process on offspring. 4 female and 2 male Balb/C

mice were purchased at 6 weeks of age, separate from the mice used for the major experiment. They

were left to acclimate for 2 weeks, fed and housed as described above. At 8 weeks of age, mice were

placed together (2 females to 1 male) and mated for 3 days. At birth, we culled offspring to 6 offspring

per litter, and cross-fostered half of resulting offspring, matching offspring by birth date to reduce

potential of rejection. Pups were marked with a toe tattoo to identify birth group. Cross-fostering led to

the generation of 2 groups of offspring: non cross-fostered and cross-fostered groups. We performed

lung lavage, sacrificed and collected tissues from all offspring at 8 weeks of age.

Measurement of candidate gene DNAm

500 ng of DNA isolated from the left lungs was treated with sodium bisulfite (Zymo Research) to

generate bisulfite-converted DNA (bcDNA), following the manufacturer's protocol. Following

conversion, bcDNA was then amplified by PCR using the following conditions: 95°C for 15 min,

followed by 45 cycles of 95°C for 30 secs, 58°C for 30 secs, 72°C for 30 secs, and then followed by

72°C for 5 min. Mouse *Ahrr* and *Cyp1a1* primers were generated using the PyroMark Assay Design

software version 2.0 (Qiagen). Primer sequences can be found in Table S1. DNAm at candidate genes

was measured using the Qiagen Q48 pyrosequencer. Prior to DNAm measurement in samples, CpG

assays were validated in duplicates using mixtures of completely methylated and completely

unmethylated control DNA (0%, 25%, 50%, 75%, and 100% methylated). After validation, sample

DNAm was then measured using the optimized assay conditions.

Gene expression measurement

200ng of lung mRNA was treated with ezDNase (Thermo Fisher Scientific, Inc.) and then converted to

cDNA using Maxima cDNA synthesis Kit (Thermo Fisher Scientific, Inc.) according to the

manufacturer's protocol. Quantitative real-time RT-PCR (qPCR) was performed on the QuantStudio 3

Real-Time PCR System (Thermo Fisher Scientific, Waltham, MA), using the SYBR-Green Master

Mix (Applied Biosystems; Thermo Fisher Scientific, Inc.) to determine the relative expression levels

of *Ahrr* and *Cyp1a1* in lung tissues. *β-actin* and *Eif2a* were used as reference and normalization

controls. Samples were run in duplicates under the following cycling conditions: Holding stage, 1

cycle of 50°C for 2 min and 95°C for 2 min; cycling stage, 40 cycles of 95°C for 15 sec, 57°C for 15

sec and 72°C for 1 min; and melt curve stage, 95°C for 15 sec, 60°C for 1 min and 95°C for 15. The

primer sequences used can be found in Table S2. *Ahrr* and *Cyp1a1* mRNA levels were quantified

using the $\Delta\Delta^{Cq}$ method[43,44] and normalized against the mean *β-actin* and *Eif2a* levels in the same

sample.

Statistical analyses

Statistical analysis was done using R (version 3.6.1). Percent DNAm at each CpG was averaged across

duplicates and between-group differences were analyzed using a one-way ANOVA. Lung function and

differential cell count data were analyzed using two-way ANOVA, and multiple comparison was

performed between groups at each methacholine dose. *P* values < 0.05 were considered significant.

Adjustment for litter size effects was conducted on significant values using simple linear regression.

Figures were also produced in R, using the *ggplot2* package.

RESULTS

Development of a protocol to study the effects of early life CS exposures

We created a mouse model to study the effects of early life CS exposure on offspring, and specifically,

to isolate prenatal and postnatal CS exposures. Exposure of Balb/C dams to whole-body CS was

carried out for 9 weeks, beginning 3 weeks before mating and pregnancy, and lasting till 3 weeks after

birth (Figure 1A). To isolate the effects of prenatal and postnatal-only smoke exposures, we cross-

fostered half of the offspring born to control and smoke-exposed dams at birth, with toe tattoos to

identify cross fostered offspring (Figure 1A). All cross-fostering was carried out no more than 24

hours after birth of matched control and CS-exposed litters. Only one litter, including both cross

fostered and not cross fostered offspring was lost. All smoke-exposed dams tolerated smoke exposures

well, and our exposure paradigm did not result in loss of any dams. Since mating resulted in variable

plug/conception dates, we ensured that all analyses reported here were carried out on offspring born no

more than 48 hours apart. COVID-19 related lab lockdowns occurred shortly after weaning of the

majority of litters, resulting in changes to planned timelines.

To verify absorption of CS components, we measured cotinine levels in plasma collected from 10

dams 48 hours after smoking cessation (3 weeks after birth of pups). The remaining 12 dams' cotinine

levels were measured 13 weeks after smoking cessation (Figure 1B). Dams which were exposed to CS

had 15 times more elevated plasma cotinine levels (Mean = 1.34, SD = 0.55) 48 hours after smoking

cessation compared to control dams (Mean = 0.09, SD = 0.03) (Figure 1B, $p = 0.0068$). The relatively

low levels of cotinine observed overall in the smoke-exposed dams can be attributed to the collection

of plasma from dams 48 hours after smoking, with the reported half-life of cotinine being about 24

hours[45,46]. Cotinine levels measured 13 weeks after CS exposure showed no significant difference

between control and smoking dams (Figure 1B).

Maternal smoking can reduce litter size[47,48] and pup birth weight in mice, which might have indirect

effects on pup development[48,49]. To ensure that our DNAm and phenotypic analyses are not

confounded by such effects, we tested the effects of our smoking model on offspring birth weight and

litter size. We found a slightly larger litter size in control dams (Mean = 6.56, SD = 1.46) than smoke-

exposed ones (Mean = 5.38, SD = 1.36) (Figure 1C, T-test $p = 0.024$). Birthweight in male and female

animals was slightly but not significantly larger in CS than control pups, perhaps due to the smaller

litter size (Figure 1D). We performed sensitivity tests in downstream analyses to identify the potential

effects of litter size.

A Experimental design

B Dam cotinine levels

C Litter size

D Offspring birth weight

Figure 1: Development of a mouse model to study the effects of early life CS exposure. (A) Experimental design: Adult female mice were exposed to CS for 9 weeks, starting 3 weeks prior to mating and ending 3 weeks after birth. Half of control offspring were cross-fostered at birth with half of the CS-exposed offspring to generate 4 groups of offspring: control, prenatal CS-exposed, postnatal CS-exposed and combined prenatal and postnatal CS-exposed groups. Lungs, blood and other tissues were collected from dams and offspring at birth and at 16 weeks of age, after lung function measurement. At age 60 weeks, half of the remaining female offspring were re-exposed to CS for 3 weeks, followed by lung function and tissue collection. (B) Dam cotinine levels measured in plasma 48 hours after smoking cessation and 13 weeks after. (C) Litter size of control and smoke-exposed dams. Student t-test was used to measure differences in litter size between control and smoke-exposed dams. (D) Weight of control and CS-exposed male and female offspring at birth. Differences in birth weight were analyzed using student t-tests within sexes.

Offspring lung phenotype at 16 weeks of age, 13 weeks after cessation of smoke exposure

At baseline, only male offspring with prenatal CS exposure and female offspring with prenatal CS exposure had increased elastance (Figure 2D, $p = 0.029$) and airway resistance (Figure 2B, $p = 0.018$), respectively, compared to control offspring. When methacholine was introduced however, male offspring in the postnatal CS group showed significant decrease in responsiveness for total lung resistance, airway resistance and tissue elastance (Figures 2E, 2F and 2H), while fully exposed male offspring showed increased methacholine sensitivity for total lung resistance (Figure 2E). Similarly, full CS exposure induced decreased methacholine sensitivity for airway resistance in female offspring (Figures 2F and 2H).

When examining immune cell infiltration in the lung, female offspring exposed to postnatal only CS

had significantly elevated eosinophils (Mean = 1691, SD = 1647) compared to control females (Mean

= 475, SD = 800) (Figure 3B, p = 0.0046), while male offspring with full CS exposure had

significantly elevated lymphocytes (Mean = 3479, SD = 3064) compared to control males (Mean =

601, SD = 659) (Figure 3D, p = 0.029).

Together, our data suggested a mild phenotype induced by 9-week early life smoke exposure in

indirectly exposed offspring, as observed by persistent decline in lung function and increased

infiltration of immune cells into the lungs. Similar analyses in dams are in the Supplementary Results.

Offspring baseline lung function 13 weeks after smoke exposure

Offspring lung function 13 weeks after smoke exposure

Figure 2: Early life exposure to CS alters sensitivity to methacholine up to 13 weeks after cessation of

smoke exposure. Lung function parameters (A) Total lung resistance at baseline. (B) Airway resistance

at baseline. Female offspring exposed to prenatal CS had significantly higher Rn (Mean = 0.34, SD =

0.03) compared to female controls (Mean = 0.26, SD = 0.06). (C) Tissue resistance at baseline. (D)

Alveolar elastance at baseline. Male offspring exposed to prenatal CS had significantly higher H

(Mean = 17.50, SD = 0.63) compared to male controls (Mean = 16.40, SD = 0.91). (E) Total lung

resistance with methacholine. (F) Airway resistance with methacholine. (G) Tissue resistance with

methacholine. (H) Alveolar elastance with methacholine. Lung function was assessed by taking 90^{th}

percentile values in response to injection of saline into the lungs (baseline) and in response to

increasing doses of methacholine. Lung function data was analyzed in a sex-disaggregated manner

using a two-way ANOVA, followed by multiple comparisons between groups at each methacholine

dose where significant.

Figure 3: Early life exposure to CS increases immune cell infiltration into offspring lungs, up to 13 weeks after smoke exposure. (A) Total immune cells per mL lavage. (B) Eosinophils per mL lavage. (C) Macrophages per mL lavage. (D) Lymphocytes per mL lavage. Differential cell counts were normalized to lavage volume. Pairwise comparisons were conducted using t-tests.

Impact of CS on offspring cardiac morphometry and function

To determine whether CS exposure affected cardiac morphometry or function we used echocardiography to image the heart and examine left ventricular and right ventricular structural and functional parameters. In males, there were no effects of smoke exposure on left ventricle (LV) mass or other LV structural parameters (Table S3). In females, prenatal CS exposure reduced the intraventricular septum, the LV internal diameter and LV volume (Table S4). On the other hand, postnatal CS exposure increased LV posterior wall thickness in females. Overall, these structural

changes did not impact LV mass in females. Regarding LV functional parameters, in the males,

prenatal CS exposure significantly increased LV ejection fraction and cardiac output, though postnatal

CS and the combination of pre- and postnatal CS did not affect these parameters (Table S3). However,

in females, CS exposure did not affect LV function in any of the groups (Table S4). Next, right

ventricle (RV) morphometry and function was examined. In both males and females, RV wall

thickness was increased in prenatal CS exposed offspring and by postnatal CS as well as the

combination of prenatal and postnatal CS (Tables S3, S4). CS did not alter RV function and

hemodynamics in males (Table S3). Female RV function and hemodynamic parameters were more

sensitive to the effects of CS. Prenatal CS exposure increased the pulmonary vessel peak velocity and

peak pressure (Table S4). In addition, prenatal CS exposure followed by postnatal CS also increased

pulmonary valve peak velocity and peak pressure in females. Pulmonary acceleration time was

significantly reduced by postnatal CS and the combination of prenatal and postnatal CS in females.

DNAm and gene expression at birth and 16 weeks of age in candidate genes

Results from human studies have shown that maternal smoking causes decreased and increased DNAm

of *AHRR* and *CYP1A1* respectively, in newborn umbilical cord blood[19–21,23,26], and that these

differential methylation patterns persist in adult blood at midlife[24]. The effects of smoking on the

human lungs is less well documented, as only one study to our knowledge has reported increased

AHRR and *CYP1A1* DNAm in nasal epithelia of adult smokers[28]. Therefore, as DNAm is species and

tissue-specific, and studies investigating effects of prenatal CS exposure on the lungs is lacking, we

measured *Ahrr* and *Cyp1a1* DNAm in the lungs and blood of offspring at birth immediately after CS

exposure and at 16 weeks of age (13 weeks after cessation of smoke exposure). Since DNAm is also

associated with regulation of gene expression, we also measured *Ahrr* and *Cyp1a1* expression in dam

and offspring lungs at both timepoints.

At birth, *Ahrr* and *Cyp1a1* DNAm in the lung and blood were not significantly different between

control and smoke-exposed offspring (Figure 4A and 4C). However, both male and female offspring

born to CS-exposed dams showed lung *Cyp1a1* expression levels which were over 20 times higher

than controls (Figure 4B).

We next measured *Ahrr* and *Cyp1a1* methylation levels in lungs and blood of offspring at 16 weeks of

age to determine longitudinal effects of early life smoke exposure on offspring (Figure 4E – 4H).

There were no significant differences in *Ahrr* DNAm between control and CS-exposed groups in

offspring blood 13 weeks after cessation of smoke exposure (Supplementary figure 4B). In the lung in

male offspring only, *Ahrr* DNAm in prenatally exposed offspring was significantly higher than

offspring exposed only postnatally (Figure 4G, $p = 0.03221$), but none of the CS-exposed offspring

showed significantly altered *Ahrr* DNAm compared to control offspring. In female offspring at 16

weeks, those exposed to prenatal or postnatal CS had higher *Cyp1a1* DNAm in the lung compared to

controls (Figure 4E), while offspring with full CS exposure showed significantly decreased *Cyp1a1*

methylation compared to either postnatal (Figure 4E, $p = 0.02$) or prenatal groups (Figure 4E, $p = $

0.0049). These differences remained significant even after adjusting for possible effects of litter size

on DNAm using linear regression. The decrease in *Cyp1a1* DNAm in the lung in fully exposed female

offspring, but not in cross-fostered offspring, is similar to the direction of *Cyp1a1* DNAm in dams

(Supplementary figure 5).

We observed the same trend when we measured offspring lung *Cyp1a1* expression: increased

expression levels of *Cyp1a1* in the postnatal only and prenatal only CS groups but decreased

expression in the lungs of offspring exposed to both prenatal and postnatal CS, compared to controls,

though these differences were not statistically significant (Figure 4F). To show that the differences in

candidate gene DNAm observed in offspring was not because of changes in global DNAm, we

measured DNAm at a control gene, *Prkaa1*, which has not been associated with smoking. As expected,

Prkaa1 methylation in lungs of offspring exposed to both prenatal and postnatal CS was not

significantly different from *Prkaa1* methylation in control offspring (Supplementary Figure 6).

To rule out the potential effects of cross-fostering on DNAm, we cross fostered litters without any

exposures and measured *Cyp1a1* DNAm in lungs. We found no differences in *Cyp1a1* DNAm

between cross-fostered and non-cross fostered offspring (Supplementary figure 7).

This data suggested that early-life exposure to CS creates long lasting lung DNAm and gene

expression changes at one of our two candidate genes.

Candidate gene DNA methylation and expression in offspring lungs at birth

Candidate gene DNA methylation and expression in offspring lungs 13 weeks after smoke exposure

Figure 4: Early life exposure to CS significantly alters lung *Cyp1a1* expression at birth and *Cyp1a1*

DNAm 13 weeks after smoke cessation. (A) *Cyp1a1* DNAm in offspring lungs at birth. (B) *Cyp1a1*

expression in offspring lungs at birth was significantly increased in CS-exposed males (Mean = 19.00,

SD = 10.30) and females (Mean = 27.50, SD = 4.54) offspring, compared to control males (Mean =

1.36, SD = 0.93) and control females (Mean = 1.47, SD = 0.43). (C) *Ahrr* DNAm in offspring lungs at

birth. (D) *Ahrr* expression in offspring lungs at birth. (E) *Cyp1a1* DNAm in offspring lungs at 16

weeks of age. Prenatal and postnatal CS exposure caused an increase in female lung *Cyp1a1* DNAm,

while fully exposed offspring showed a decrease. Only prenatally exposed female offspring had

significantly elevated *Cyp1a1* DNAm (Mean = 18.80, SD = 1.42) compared to control females (Mean

= 14.10, SD = 0.26). (F) *Cyp1a1* expression in offspring lungs at 16 weeks of age. Prenatally exposed

female offspring had significantly elevated lung *Cyp1a1* expression levels (Mean = 3.28, SD = 1.44)

compared to control females (Mean = 1.48, SD = 0.47). (G) *Ahrr* DNAm in offspring lungs at 16

weeks of age. (H) *Ahrr* expression in offspring lungs at 16 weeks of age. One-way ANOVA was used

to compare DNAm values between groups ($p < 0.05$ was significant), followed by two-group

comparisons where significant.

389   Acute re-exposure to CS at 60 weeks alters lung function and re-establishes DNAm patterns

There has been some evidence in humans that maternal smoking in pregnancy induces persistent

DNAm changes until midlife, regardless of smoking habits later in life[24]. We thus sought to test what

effects re-exposure to CS in adulthood would have on the lung phenotypic and DNAm patterns set by

early life CS exposure. To achieve this, at 60 weeks of age, we re-exposed half of the offspring

remaining to CS for 3 weeks, followed by lung function, differential immune cell analyses and DNAm

measurement. At the time of re-exposure, we had a limited number of male offspring remaining, and

so the following experiments were conducted solely on female offspring. We also had a low number of

animals in the exclusively postnatal CS group, so they contribute to the baseline data only at 60 weeks

of age.

Offspring that got full CS exposure in early life, followed by re-exposure to CS at 60 weeks, had

significantly increased tissue resistance (Figure 5C) and total lung resistance (Figure 5A) levels at

baseline, while re-exposed prenatal CS offspring had increased tissue resistance compared to offspring

who had no early life CS exposure (controls), but were exposed to CS at 60 weeks (Figure 5). Immune

cell infiltration specifically by macrophages was increased only in animals with full early life CS

exposure who were also re-exposed later in life, but this increase was not statistically significant ($p =$

0.1223) (Figure 5E - 5H).

DNAm and gene expression at *Cyp1a1* showed a similar pattern, they were altered only in offspring

with full CS exposure in early life who were also re-exposed to CS (Figure 6A, $p = 0.036$). This

decrease in DNAm was not significant after adjusting for possible effects of litter size on DNAm using

linear regression, but it is unlikely that litter size continues to influence DNAm at 60 weeks of age.

Interestingly, the *Cyp1a1* DNAm patterns established upon re-exposure to CS in adulthood were

similar to those observed at 16 weeks.

Taken together, this data demonstrates that early life exposure to CS induces transient changes in lung

phenotype and *Cyp1a1* DNAm, which become re-established upon personal smoking in adulthood.

Figure 5: Acute re-exposure to CS at 60 weeks induced significant alterations in baseline lung function compared to early life control offspring which were exposed to CS in adulthood. Lung function parameters at baseline (A) total lung resistance. Offspring with full early life CS exposure and re-exposed to CS at 63 weeks had higher Rrs (Mean = 0.88, SD = 0.62) compared to early life control offspring exposed to CS at 63 weeks only (Mean = 0.35, SD = 0.11). (B) airway resistance. (C) tissue resistance. Offspring with full early life CS exposure and re-exposed to CS at 63 weeks had higher G (Mean = 6.04, SD = 3.60) compared to early life control offspring exposed to CS at 63 weeks only (Mean = 2.74, SD = 0.20). Offspring exposed to prenatal CS early in life and re-exposed to CS at 63 weeks had higher G (Mean = 3.40, SD = 0.51) compared to early life control offspring exposed to CS at 63 weeks only. (D) alveolar elastance. (E) Total immune cells per mL lavage. (F) Eosinophils per mL lavage. (G) Macrophages per mL lavage. (H) Lymphocytes per mL lavage. For lung function, 90[th] percentile values were calculated in response to saline administered into the lungs. Differential cell counts were normalized to lavage volume.

Cyp1a1 DNA methylation and expression in offspring lungs at 63 weeks

Figure 6: Re-exposure of offspring to CS at 60 weeks induced changes in lung DNAm which were

similar to those observed at 16 weeks. (A) *Cyp1a1* DNAm in offspring lungs at 63 weeks. Offspring

with full early life CS exposure and re-exposed to CS at 63 weeks had higher lung *Cyp1a1* DNAm

(Mean = 17.00, SD = 0.65) compared to early life control offspring exposed to CS at 63 weeks only

(Mean = 19.20, SD = 1.37). (B) *Cyp1a1* expression in offspring lungs at 63 weeks. Differences in

DNAm and expression were calculated using a one-way ANOVA, followed by two-group

comparisons.

DISCUSSION

The global burden of smoking still remains significant, despite numerous campaigns against its

use, and smoking in pregnancy predisposes offspring to harmful health outcomes[55–57]. CS components

have been known to pass through the placenta to the developing fetus, but it is not yet clear what

specific mechanisms link CS exposure to offspring health[27,58–60]. Past studies have shown that prenatal

exposure to CS induces differential DNAm in newborn umbilical cord blood, placenta, buccal and

nasal epithelia, and adult peripheral blood granulocytes[20,23,28,29,59]. However, previous studies in

humans have not been able to isolate the influence of postnatal exposure to smoke, as prenatal and

postnatal exposures tend to be highly correlated in humans. In addition, while the effects of maternal

smoking on peripheral tissues are well documented, less is known about the effects on the lungs,

despite significant epidemiological evidence linking maternal smoking to long term lung health,

including asthma and COPD[61–65]. Since we know that each tissue has a unique epigenetic signature[66–

74], it is essential to measure the effects of early life smoke exposure in the lungs. Therefore, we created

a mouse model to study the independent effects of prenatal and postnatal CS exposure that would also

allow us to measure lung phenotype and DNAm specifically in the lung. We used cross-fostering to

isolate prenatal and postnatal exposures rather than altering exposure to CS in the dams to avoid the

effects of nicotine withdrawal on the dams and pups[75–78]. Here, we showed that exposure to CS at

different phases early in life alters offspring lung function, DNAm and gene expression profiles, and

that these early life exposures prime the lungs such that a secondary smoke exposure in adulthood re-

establishes some of these outcomes. Tables S5 and S6 depict summaries of our findings in offspring

and dams respectively.

Cigarette smoking has been shown to cause airway hyperresponsiveness and altered lung

function in offspring[12,63,79–81]. In our model, we noted sex-specific differences in specific

manifestations of lung function in offspring even after cessation of smoke exposure for 13 weeks,

similar to those that have previously been reported in other human and mouse studies[82–85]. Many of

these studies have identified female lungs as being less sensitive to the effects of environmental toxins

such as from CS. This is hypothesized to be because important protective factors in lung development,

such as surfactant production, begin earlier in females than males[86–88]. In our model however, the sex-

specific differences we observed in lung function and immune cell infiltration were more spread out,

with female offspring exhibiting differences in certain parameters but not males, and vice versa. The

observed differences in our results compared to previous studies may be as a result of differences in

the study designs- for instance, the fact that we completely isolated CS exposures by cross fostering, as

opposed to these other studies.

In our study, lung function defects in female offspring persisted until 16 weeks of age, which is

supported by previous studies that have reported persistent effects of CS, both in human and animal

studies[63,85,94]. In one such study, maternal smoking led to a decrease in offspring lung function which

lasted till 21 years of age[65]. Similarly, another longitudinal study found persistently decreased forced

expiratory flows in 8800 children aged 8-12 years of age who were exposed to CS *in utero*[94]. We

found that the lung function alterations observed in female offspring did not last till 60 weeks, but

reappeared upon re-exposure to CS at 60 weeks of age. While we do not know when these differences

disappeared, these results suggest that early-life CS exposure alone may not be enough to induce life-

long lung function alterations that last until 60 weeks, but may prime the lungs so that once offspring

are re-exposed to CS in adulthood, lung function is altered. This idea of long term priming has been

well documented in the past[84,95,96], but few studies have extended exposures as long as those shown

here. In a similar light, one study showed that early-life secondhand smoke exposure did not alter lung

function until the administration of a postnatal allergen (*A. fumigatus* extract)[96]. Another study found

that *in utero* exposure to second-hand smoke exacerbated allergic response and lung function deficits

in offspring when exposed to ovalbumin at 23 weeks of age[82].

In addition to lung phenotype, it is important to investigate the effects of CS on DNAm, as this

could give insights into the underlying mechanisms of disease development. Cytochrome P450 (Cyp)

enzymes are hemoproteins which are involved in metabolism of drugs and other xenobiotics primarily

in the liver[97]. Its major substrates are polyaromatic hydrocarbons (PAHs) and nitrosamines (major

carcinogenic components of CS[98]), and activation by these ligands usually occurs via an aryl

hydrocarbon receptor (Ahr)-dependent pathway[99,100]. There are many subclasses of the Cyp enzymes,

but CYP1A1 is extensively studied as it is expressed at basal levels in extra-hepatic tissues[101] and is

highly inducible, leading to the speculation that it may be primarily responsible for xenobiotic

metabolism in extra-hepatic tissues[102]. When activated by Ahr, CYP1A1 directly hydroxylates or

oxidizes its bound ligand, leading to its detoxification[103–105]. CS is a potent inducer of the Ahr-Cyp

pathway both in humans and murine models[106–108], so it follows that measuring the effects of CS on

DNAm in these genes would be informative on CS-induced phenotypes.

The dynamics of *Cyp1a1* DNAm and expression over the life course in CS-exposed lungs

revealed unexpected patterns. First, we noted that maternal smoking did not alter offspring *Ahrr* or

Cyp1a1 DNAm in the lung at birth at our chosen CpGs but causes a 20-fold increase in expression of

just *Cyp1a1*. At 16 weeks, offspring exposed to only prenatal CS or only postnatal CS showed a slight

increase in *Cyp1a1* DNAm (and expression), while those exposed to both prenatal and postnatal CS

showed decreased *Cyp1a1* DNAm and expression in their left lungs compared to control offspring. We

tested the effects of cross-fostering independently and saw no difference in *Cyp1a1* DNAm, which

implies these differences are not an effect of cross-fostering itself. These opposing results could

possibly be explained by the *developmental mismatch hypothesis*, which surmises that traits that

develop in an organism in one environment may be disadvantageous in a different environment[109–111].

In other words, the mismatch between prenatal and postnatal environments in the cross-fostered

offspring would confer different characteristics from offspring with matched prenatal and postnatal

environments. The fact that the *Cyp1a1* DNAm in the prenatal CS only group and postnatal CS only

group changes in the same direction emphasizes the need for future animal models to isolate and study

them separately, as it is evident that they produce different effects and can be difficult to differentiate

in more long-term studies. The increase in *Cyp1a1* DNAm observed in the prenatal CS only group is in

fact in line with many human studies which have measured prenatal CS-induced DNAm changes in

newborn umbilical cord blood[20–23,25]. We cannot make comparisons to other mouse studies however,

as none so far has investigated effects of CS on *Cyp1a1* DNAm.

The fact that these differences in offspring *Cyp1a1* DNAm were observed 13 weeks after

smoke exposure cessation shows that similar to the lung function results, the effects of early life CS on

offspring *Cyp1a1* DNAm are long-lasting. Another mouse study similarly showed that maternal

smoking induced changes in *Cyp2a5* DNAm methylation in mouse liver which persisted till 140 days

after smoking[112]. The evidence of persistent effects of CS in humans is more abundant, as human

studies have shown that maternal smoking causes changes in the offspring epigenome which can last

till mid-life[21,23,24]. It is worth noting that our observed differences in *Cyp1a1* DNAm were not

persistent till 16 weeks in male offspring, suggesting that the effects of maternal smoking on offspring

DNAm could be sex-specific[66,113,114] .

Finally, we observed that while early life CS-induced *Cyp1a1* DNAm changes do not persist

till 60 weeks, CS-exposed animals develop impaired lung function with age. In addition, acute

exposure to CS in adulthood re-established DNAm patterns observed at 16 weeks and was associated

with an immune cell infiltration phenotype that was dependent on both early life and later life CS

exposure. The disappearance of the *Cyp1a1* DNAm patterns at 60 weeks and re-establishment by acute

smoke exposure proves that DNAm cannot be the epigenetic mechanism responsible for this response.

The next most likely candidate for a mechanism linking early life exposure with later life response is

histone modifications. One study has in fact shown that early life smoke exposure in rats causes

chromatin remodeling, marked by increased histone acetylation and altered transcription factor binding

in the lungs[115]. The DNAm and lung phenotype changes observed at this timepoint, in particular the

immune cell infiltration phenotype, are further evidence of priming. Some studies have suggested that

disappearance of detrimental phenotypes over time is an adaptation mechanism in response to

adversity[116,117], which allows affected individuals to function as close to optimal as possible, even

though they may show some altered phenotypic patterns. However, the underlying detrimental

signatures are embedded until a secondary stimulant is introduced[116,117].

Taken together, our results represent a novel model for the DOHaD hypothesis, which surmises

that early life adversities or environment can shape the offspring phenotype such that it will produce

long term consequences on an individual. While some hypotheses such as the thrifty phenotype[118] and

adaptive response[119] have been used to provide reasons why there are long-lasting health conditions

following early-life adversity, the underlying mechanisms are largely unknown[120]. Epigenetic

mechanisms like DNAm are suspected to play major roles in this phenomenon and there are many

studies which have investigated the effects of early life environment on offspring DNAm. For

example, early life smoke exposure and air pollution exposure in general has been shown to cause

conditions like asthma and cancer in adulthood, and many studies have suggested DNAm as the

underlying mechanism[20,31,121,122]. However, we have shown that in our model DNAm was not

responsible for long-term CS-associated phenotype, and future work will investigate other epigenetic

marks.

One limitation of this study is that we measured DNAm in whole mouse lungs. While this

gives us a whole picture, it does not tell what specific lung structural cell type is responsible for the

observed changes in DNAm. This is an important question, as the lungs are made up of numerous cell

types, and DNAm is cell type-specific[123–125]. It is unlikely that immune cell infiltration is the cause of

the observed DNAm changes, as lungs were lavaged to remove alveolar immune cells, and the relative

number of immune cells residing in lung tissue is very small. In addition, in a similar pilot study we

conducted prior to this experiment, we found that *Cyp1a1* DNAm in the blood of adult female control

mice is higher than *Cyp1a1* DNAm in lungs, and as a result, infiltration of immune cells from the

blood into lungs is unlikely to be the cause of the decrease in lung *Cyp1a1* DNAm in the present study

(Supplementary figure 8). A second limitation is that as our model involves administration of heavy

doses of CS and so we cannot extrapolate these results to analyze the effects of light or moderate

smoking.

In conclusion, this study has created an effective model for DOHaD to investigate the effects of

early life CS exposure, providing an essential foundation for future animal studies. Using this model,

we have shown that the effects of early life smoke exposure are embedded into the offspring,

producing phenotypes which are re-established by smoking in adulthood. Future studies are needed to

investigate these effects across the mouse epigenome and in particular, investigate the effects of early

life CS on chromatin accessibility and specific histone marks or variants, to identify the specific

mechanism responsible for the long-term persistence of response due to early life CS exposure.

Supplementary Results

Dam lung phenotype immediately after 9 weeks of smoking and 13 weeks after smoking cessation

Several studies have shown that personal smoking alters lung function parameters and induces airway hyperresponsiveness[50–52]. Therefore, we assessed the effects of CS exposure on the lungs by measuring lung function and immune cell infiltration.

We first measured the effects of personal smoking on dam lung function and immune cell infiltration, to serve as a baseline for these measurements in their offspring. We found that direct CS exposure did not alter dam lung function immediately after smoking (Supplementary figure 1), but over time induced decreased methacholine responsiveness in dam lungs, as observed by significant alterations in tissue resistance and total lung resistance values 13 weeks after smoking cessation (Supplementary figure 2). Personal smoking for 9 weeks also caused a 4-fold and 8-fold increase in eosinophils and lymphocytes respectively immediately after smoking, compared to controls (Supplementary figure 3). The immune cell infiltration differences in dams however, did not persist 13 weeks after smoking cessation (Supplementary figure 3).

Together, this data showed a mild phenotype induced by 9-week smoke exposure in dams who were directly exposed to CS, as observed by increased infiltration of immune cells into dam lungs immediately after smoking and persistent decline in lung function.

Candidate gene DNAm and expression in dams 13 weeks after smoking cessation

To determine whether DNAm changes reflect early life CS exposure in the mouse lung, we identified candidate genes to assess via pyrosequencing. We selected two CpGs with the biggest effect sizes from

a large meta-analysis of prenatal CS exposure conducted in human cohorts; *AHRR* at human position

373378 (mouse *Ahrr* position 74260517) which is not well conserved between humans and mice, and

CYP1A1 at human position 75019251 (mouse *Cyp1a1* position 57696335). In addition, we chose

position 40796738 (corresponding to the mouse gene *Prkaa1* position 5174566) as a CpG which is not

known to demonstrate DNAm changes in response to CS in human studies as a negative control[20,53,54].

We measured *Ahrr* DNAm in dam blood, since the corresponding human locus (cg05575921) is

known to be significantly altered in blood of humans exposed to CS[19–21,23]. However, we found no

significant differences in *Ahrr* DNAm in dam blood 13 weeks after smoking cessation (Supplementary

figure 4A).

On the other hand, chronic exposure to CS for 9 weeks led to a significant decrease in *Cyp1a1* DNAm

in the left lungs of exposed dams (Mean = 15.00, SD = 1.07), compared to controls (Mean = 17.60, SD

= 0.58) which lasted up to 13 weeks after smoking cessation (Supplementary figure 5, $p = 0.033$).

Similarly, direct smoke exposure caused a slight but not significant decrease in dam lung *Ahrr* DNAm

(Supplementary figure 5). *Ahrr* and *Cyp1a1* expression in the lungs of CS-exposed dams was also

slightly but not significantly elevated (Supplementary figure 5).

These results thus show that personal smoking for 9 weeks causes alterations in DNAm which persist

for 13 weeks after smoking cessation.

618   Table S1: Mouse pyrosequencing primers

Gene	Primer sequence
Ahrr	Forward: AGTTGTTGTGGTTTTTGATAAAAAGAG
	Reverse: /5BiosG/AAACACAATATACAAAAATCCAACTCAT
	Sequencing: AGGGATTTATTGGGG
Cyp1a1	Forward: TTTTGTTTGAGAGTAAAGGTTTGAATAG
	Reverse: /5BiosG/ACACCTACAAAACAACCAACTAA
	Sequencing: GTTTATATTTTTAGTGGTAGGAT
Prkaa1	Forward: AGGAAGTTTGGAGTTGATAAGAA
	Reverse: /5BiosG/CTCCCCTAAAAATAACACAACAAC
	Sequencing: ATTATATAGTAAGTTAATTGAGTTT

629   Table S2: Mouse qPCR primers

Gene	Primer sequence
Ahrr	Forward: GGACAAACAGATAGGAGCCACT
	Reverse: GCTGTTCTGTGCTCCAGTTCA
Cyp1a1	Forward: CAAGCGAAAGTGCATCGGAG
	Reverse: GAGGACCAGAAGACCGCATC
β -actin	Forward: AGAAGAGCTATGAGCTGCCTG
	Reverse: TAGAGGTCTTTACGGATGTCAACG
Eif2a	Forward: TCCCAAAAGCAGTTTGCCTTG
	Reverse: TTGTAGGTTAGGCGTCCCAG

641   Table S3: Male offspring echocardiography parameters

	Control	Prenatal Smoke	Postnatal Smoke	Prenatal and Postnatal Smoke	Prenatal Smoke	Postnatal Smoke	Interaction
Body Weight (g)	26.0 ± 0.13	25.9 ± 0.13	25.7 ± 0.06	25.7 ± 0.12	0.8872	0.4964	0.8655
LV mass (mg)	89.1 ± 0.9	89.2 ± 1.4	85.6 ± 0.8	82.6 ± 1.3	0.6957	0.1840	0.6716
LVPWd (mm)	0.75 ± 0.01	0.76 ± 0.01	0.73 ± 0.01	0.67 ± 0.01	0.4383	0.1099	0.3685
LVAWd (mm)	0.74 ± 0.01	0.75 ± 0.01	0.79 ± 0.01	0.73 ± 0.01	0.4717	0.5564	0.2913
IVSd (mm)	0.66 ± 0.01	0.69 ± 0.01	0.71 ± 0.01	0.68 ± 0.01	0.9911	0.5504	0.2704
LVIDd (mm)	4.10 ± 0.02	4.06 ± 0.02	4.02 ± 0.01	4.00 ± 0.01	0.5881	0.2374	0.9119
LV Vol (ul)	73.5 ± 0.5	69.8 ± 0.8	68.8 ± 0.6	66.6 ± 0.5	0.2022	0.0914	0.7522
EF (%)	51.8 ± 0.6	56.2 ± 0.7	53.5 ± 0.5	54.9 ± 0.4	0.0352	0.3285	0.8389
FS (%)	26.3 ± 0.4	29.2 ± 0.5	28.2 ± 0.3	29.2 ± 0.3	0.1095	0.4307	0.4225
CO (ml/min)	14.9 ± 0.3	17.0 ± 0.2	15.8 ± 0.1	16.3 ± 0.2	0.0347	0.8675	0.2108

SV (ul)	38.5 ± 0.6	40.5 ± 0.5	37.9 ± 0.4	39.5 ± 0.4	0.1241	0.6471	0.4787
IVCT (ms)	8.1 ± 0.3	9.9 ± 0.2	12.1 ± 0.3	11.3 ± 0.3	0.6817	0.0198	0.2509
IVRT (ms)	15.2 ± 0.3	13.7 ± 0.3	15.3 ± 0.2	14.2 ± 0.2	0.1118	0.7505	0.8026
E/A	2.47 ± 0.07	2.05 ± 0.05	1.88 ± 0.03	2.16 ± 0.05	0.6882	0.1546	0.0419
E'/A'	1.15 ± 0.03	1.12 ± 0.03	1.03 ± 0.02	1.00 ± 0.02	0.5589	0.1929	0.8663
E/E'	47.2 ± 0.6	42.8 ± 0.9	41.1 ± 0.5	40.3 ± 0.8	0.2331	0.0520	0.3970
MPI	0.44 ± 0.01	0.43 ± 0.01	0.51 ± 0.01	0.46 ± 0.01	0.3947	0.0944	0.4189
PAT (ms)	15.8 ± 0.2	15.7 ± 0.2	16.8 ± 0.2	15.7 ± 0.1	0.3675	0.4982	0.4769
PET (ms)	62.2 ± 0.5	60.3 ± 0.4	61.2 ± 0.3	59.6 ± 0.5	0.2517	0.5760	0.9120
PAT/PET	0.26 ± 0.01	0.26 ± 0.01	0.27 ± 0.01	0.26 ± 0.01	0.8646	0.2981	0.5046
PV Peak Velocity (mmHg)	608.0 ± 6.1	669.9 ± 7.7	637.3 ± 5.1	639.0 ± 3.0	0.1371	0.9716	0.1595

PV Peak Pressure (mmHg)	1.49 ± 0.03	1.70 ± 0.04	1.65 ± 0.03	1.64 ± 0.02	0.3104	0.6548	0.2682
RVWT (mm)	0.50 ± 0.01	0.52 ± 0.01	0.53 ± 0.00	0.57 ± 0.01	0.0005	< 0.0001	0.2881
RVOT (mm)	1.24 ± 0.01	1.19 ± 0.01	1.15 ± 0.01	1.20 ± 0.01	0.9959	0.2501	0.1421
MPAP (mmHg)	80.2 ± 0.13	80.3 ± 0.12	79.6 ± 0.11	80.3 ± 0.08	0.3678	0.4982	0.4769
HR (bpm)	390 ± 4.1	405 ± 3.8	413 ± 2.6	416 ± 3.6	0.7402	0.9087	0.2510

Abbreviations: HR-heart rate; LV- left ventricle; LVID- LV internal diameter; LVAW- LV anterior

wall; LVPW- LV posterior wall; IVS- interventricular septum; EF- ejection fraction; FS- fractional

shortening; CO- cardiac output; SV- stroke volume; IVRT- isovolumic relaxation time; E/A; ratio of

mitral valve peak velocities measured by pulsed wave doppler during the early (E) and atrial (A)

peaks; E'/A'- ratio of mitral valve peak velocities measured by tissue doppler during the early (E') and

atrial (A') peaks; E/E'- ratio of early mitral inflow velocity (E) and mitral annular early diastolic

velocity (E') measured by tissue doppler; MPI- myocardial performance index; LV Vol- end diastolic

volume; bpm- beats per minute; PAT- pulmonary acceleration time; PET- pulmonary ejection time;

RVWT- right ventricle wall thickness; RVOT- right ventricular outflow tract; MPAP- Mean

pulmonary arterial pressure

Table S4: Female offspring echocardiography parameters

	Control	Prenatal Smoke	Postnatal Smoke	Prenatal and Postnatal Smoke	Prenatal Smoke	Postnatal Smoke	Interaction
Body Weight (g)	21.4 ± 0.04	21.3 ± 0.07	21.1 ± 0.06	20.5 ± 0.07	0.2274	0.0658	0.4613
LV mass (mg)	68.1 ± 0.3	67.5 ± 0.5	69.0 ± 0.5	68.0 ± 0.6	0.7459	0.6094	0.9065
LVPWd (mm)	0.61 ± 0.002	0.63 ± 0.003	0.66 ± 0.005	0.69 ± 0.005	0.1866	0.0010	0.9604
LVAWd (mm)	0.65 ± 0.003	0.61 ± 0.003	0.65 ± 0.003	0.67 ± 0.005	0.5532	0.0733	0.1731
IVSd (mm)	0.65 ± 0.003	0.62 ± 0.003	0.67 ± 0.003	0.65 ± 0.004	0.05	0.1592	0.6054
LVIDd (mm)	3.7 ± 0.01	3.6 ± 0.01	3.7 ± 0.01	3.6 ± 0.01	0.0072	0.5422	0.3386
LV Vol (ul)	57.8 ± 0.15	55.9 ± 0.28	58.9 ± 0.34	53.9 ± 0.38	0.0112	0.7292	0.2413
EF (%)	57.1 ± 0.2	57.0 ± 0.4	55.3 ± 0.3	59.2 ± 0.3	0.1831	0.8515	0.1593
FS (%)	29.0 ± 0.1	28.7 ± 0.2	28.4 ± 0.2	30.9 ± 0.2	0.2157	0.3742	0.1142

CO (ml/min)	13.4 ± 0.07	12.5 ± 0.09	13.2 ± 0.09	13.2 ± 0.11	0.3023	0.5938	0.2920
SV (ul)	34.1 ± 0.13	32.9 ± 0.25	33.3 ± 0.24	33.2 ± 0.24	0.5342	0.8299	0.6036
IVCT (ms)	10.6 ± 0.14	10.3 ± 0.16	11.4 ± 0.15	9.4 ± 0.18	0.1162	0.9182	0.2574
IVRT (ms)	13.9 ± 0.09	14.8 ± 0.14	13.4 ± 0.10	14.0 ± 0.12	0.1287	0.1935	0.7663
E/A	2.44 ± 0.02	2.37 ± 0.03	2.38 ± 0.03	2.07 ± 0.03	0.2039	0.2141	0.2858
E'/A'	1.04 ± 0.01	1.08 ± 0.01	1.05 ± 0.01	1.11 ± 0.01	0.3278	0.6450	0.8636
E/E'	41.0 ± 0.31	39.9 ± 0.40	41.2 ± 0.33	39.5 ± 0.40	0.4263	0.9619	0.8685
MPI	0.44 ± 0.003	0.44 ± 0.004	0.45 ± 0.004	0.44 ± 0.004	0.6185	0.9261	0.9677
PAT (ms)	16.9 ± 0.10	15.9 ± 0.08	15.5 ± 0.07	15.0 ± 0.08	0.0888	0.0105	0.4307
PET (ms)	63.0 ± 0.14	64.2 ± 0.17	62.1 ± 0.22	62.0 ± 0.17	0.5380	0.0826	0.4368
PAT/PET	0.27 ± 0.002	0.25 ± 0.001	0.25 ± 0.001	0.24 ± 0.001	0.0903	0.0894	0.3316

PV Peak Vel (mm/s)	580.0 ± 2.3	615.1 ± 3.9	605.6 ± 2.7	635.6 ± 3.4	0.0294	0.1213	0.8618
PV Peak Pressure (mmHg)	1.36 ± 0.01	1.58 ± 0.02	1.53 ± 0.01	1.69 ± 0.02	0.0080	0.0494	0.7303
RVWT (mm)	0.46 ± 0.000	0.49 ± 0.002	0.51 ± 0.003	0.54 ± 0.002	0.0004	< 0.0001	0.2874
RVOT (mm)	1.08 ± 0.004	1.13 ± 0.004	1.10 ± 0.004	1.14 ± 0.005	0.0609	0.4914	0.5624
MPAP (mmHg)	79.5 ± 0.06	79.7 ± 0.07	80.1 ± 0.05	80.5 ± 0.06	0.4103	0.0265	0.7468
HR (bpm)	393 ± 1.1	389 ± 1.5	398 ± 1.6	397 ± 1.3	0.7349	0.3194	0.8034

Abbreviations: HR-heart rate; LV- left ventricle; LVID- LV internal diameter; LVAW- LV anterior wall;

LVPW- LV posterior wall; IVS- interventricular septum; EF- ejection fraction; FS- fractional

shortening; CO- cardiac output; SV- stroke volume; IVRT- isovolumic relaxation time; E/A; ratio of

mitral valve peak velocities measured by pulsed wave doppler during the early (E) and atrial (A) peaks;

E'/A'- ratio of mitral valve peak velocities measured by tissue doppler during the early (E') and atrial

(A') peaks; E/E'- ratio of early mitral inflow velocity (E) and mitral annular early diastolic velocity (E')

measured by tissue doppler; MPI- myocardial performance index; LV Vol- end diastolic volume; bpm-

beats per minute; PAT- pulmonary acceleration time; PET- pulmonary ejection time; RVWT- right

ventricle wall thickness; RVOT- right ventricular outflow tract; MPAP- Mean pulmonary arterial

pressure

Table S5: Summary of results from offspring

		Birth		16 weeks						60 weeks	
		Male	Female	Male			Female			Female	
		CS	CS	Post	Pre	Full	Post	Pre	Full	Pre	Full
Lung function at baseline	Rrs										↑
	Rn						↑				
	G									↑	↑
	H			↑							
Lung function with methacholine	Rrs			↓		↑					
	Rn			↓				↓	↓		
	G										
	H			↓							
Lung immune cell infiltration	Macrophages										
	Eosinophils						↑				
	Lymphocytes					↑					
	Total cells										
Lung DNAm	*Cyp1a1* DNAm						↑	↑*	↓*		↓
Lung gene expression	*Cyp1a1* expression	↑	↑				↑	↑			

666   At birth and 16 weeks, arrow directions are relative to control groups. At 60 weeks, arrows are relative

667   to early-life control offspring exposed to CS in adulthood only.

668   * = borderline significance.

671   Table S6: Summary of results from dams

		Immediately after 9 weeks of smoke exposure	13 weeks after smoking cessation
		CS-exposed	CS-exposed
Lung function with methacholine	Rrs		⬇
	Rn		
	G		⬇
	H		
Lung immune cell infiltration	Macrophages		
	Eosinophils	⬆	
	Lymphocytes	⬆	
	Total cells		
Lung DNAm	*Cyp1a1* DNAm		⬇
Lung gene expression	*Cyp1a1* expression		

672   Arrow directions are relative to control groups. Lung function data reported here are results from the

673   methacholine responsiveness test, not baseline.

Supplementary figure 1: Personal smoking for 9 weeks does not significantly alter lung function in

dams immediately after smoking. Lung function parameters (A) Total lung resistance at baseline. (B)

Airway resistance at baseline. (C) Tissue resistance at baseline. (D) Alveolar elastance at baseline. (E)

Total lung resistance with methacholine. (F) Airway resistance with methacholine. (G) Tissue

resistance with methacholine. (H) Alveolar elastance with methacholine. Lung function values were

measured by calculating the 90^{th} percentile in response to inhalation of saline into the lungs.

Dam baseline lung function 13 weeks after smoking cessation

Dam lung function 13 weeks after smoking cessation

Supplementary figure 2: Personal smoking affects dam lung function up to 13 weeks after smoking

cessation. Lung function parameters. (A) Total lung resistance at baseline. (B) Airway resistance at

baseline. (C) Tissue resistance at baseline. (D) Alveolar elastance at baseline. (E) Total lung resistance

with methacholine. (F) Airway resistance with methacholine. (G) Tissue resistance with methacholine.

(H) Alveolar elastance with methacholine. Lung function values were measured by calculating the 90[th]

percentile in response to inhalation of saline into the lungs.

Dam immune cell counts after 9 weeks of smoke exposure

Dam immune cell counts 13 weeks after smoking cessation

Supplementary figure 3: Personal smoking causes immediate but transient changes to immune cell

infiltration into dam lungs. Direct exposure to CS increases immune cell infiltration into the lungs

immediately after smoking, but these effects disappear 13 weeks after smoking. (A) Total immune

cells per mL lavage immediately after smoking. (B) Eosinophils per mL lavage in the CS-exposed

dams (Mean = 7528, SD = 3736) was 4 times more elevated than control dams (Mean = 1569, SD =

1029) immediately after smoking. (C) Macrophages per mL lavage immediately after smoking. (D)

Lymphocytes per mL lavage in the CS-exposed dams (Mean = 5680, SD = 2357) was over 8 times

elevated compared to control dams (Mean = 649, SD = 256) immediately after smoking. (E) Total

immune cells per mL lavage 13 weeks after smoking cessation. (F) Eosinophils per mL lavage 13

weeks after smoking cessation. (G) Macrophages per mL lavage 13 weeks after smoking cessation. (H)

Lymphocytes per mL lavage 13 weeks after smoking cessation. Differential cell counts were

normalized to lavage volume. Two-group comparisons were conducted using a student t-test.

Ahrr DNA methylation in dam and offspring blood 13 weeks after smoke exposure

717   Supplementary figure 4: Smoking did not alter *Ahrr* DNAm in dam or offspring blood 13 weeks after

718   smoke cessation. (A) *Ahrr* DNAm in dam blood 13 weeks after smoking cessation. (B) *Ahrr* DNAm in

719   offspring blood 13 weeks after cessation of smoke exposure.

Candidate gene DNA methylation and expression in dam lungs
13 weeks after smoking cessation

738   Supplementary figure 5: CS exposure caused reduced *Cyp1a1* methylation in dam lungs 13 weeks after

739   smoking cessation. (A) *Cyp1a1* DNAm in dam lungs 13 weeks after smoking cessation. (B) *Cyp1a1*

740   expression in dam lungs 13 weeks after smoking cessation. (C) *Ahrr* DNAm in dam lungs 13 weeks

741   after smoking cessation. (D) *Ahrr* expression in dam lungs 13 weeks after smoking cessation.

Prkaa1 DNA methylation in offspring lungs after 13 weeks of smoking cessation

Supplementary figure 6: Early life exposure to CS did not significantly alter offspring lung DNAm at the control gene, *Prkaa1*.

Supplementary figure 7: Cross-fostering did not significantly affect *Cyp1a1* DNAm in mouse lungs.

To rule out potential effects of cross-fostering on offspring DNAm, we measured *Cyp1a1* DNAm in

lungs of cross-fostered and non-cross fostered control Balb/C mice at 8 weeks of age. Differences in

DNAm were measured using a student t-test.

Supplementary figure 8: Pilot study - DNAm in mouse blood is higher than in the lungs. In a previous

pilot study like the present experiment, we exposed adult female BAlb/C mice to CS twice daily for 3

weeks, immediately followed by lung function and lavage as described above. We measured DNAm in

dam blood and lungs and data was analyzed as described above. Similar to dams in the present study,

we also found that exposure of adult mice to CS for 3 weeks caused a decrease in lung *Cyp1a1*

DNAm.

REFERENCES

1. Roseboom, T. J. *et al.* Effects of Prenatal Exposure to the Dutch Famine on Adult Disease in Later Life: An Overview. *Twin Research and Human Genetics* **4**, 293–298 (2001).

2. Nielsen, C. H., Larsen, A. & Nielsen, A. L. DNA methylation alterations in response to prenatal exposure of maternal cigarette smoking: A persistent epigenetic impact on health from maternal lifestyle? *Arch Toxicol* **90**, 231–245 (2016).

3. Gluckman, P. D., Hanson, M. A., Cooper, C. & Thornburg, K. L. Effect of in utero and early-life conditions on adult health and disease. *The New England journal of medicine* **359**, 61–73 (2008).

4. Mandy, M. & Nyirenda, M. Developmental Origins of Health and Disease: the relevance to developing nations. *International Health* **10**, 66–70 (2018).

5. Government of Canada, S. C. Smokers, by age group. https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1310009610 (2018).

6. Harris, J. E. Cigarette Smoke Components and Disease: Cigarette Smoke I s More Than a Triad of Tar, Nicotine, and Carbon Monoxide. 17.

7. Swan, G. E. & Lessov-Schlaggar, C. N. The Effects of Tobacco Smoke and Nicotine on Cognition and the Brain. *Neuropsychol Rev* **17**, 259–273 (2007).

8. Cheraghi, M. & Salvi, S. Environmental tobacco smoke (ETS) and respiratory health in children. *Eur J Pediatr* **168**, 897–905 (2009).

9. Ekblad, M., Korkeila, J. & Lehtonen, L. Smoking during pregnancy affects foetal brain development. *Acta Paediatrica* **104**, 12–18 (2015).

10. Rogers, J. M. Tobacco and pregnancy. *Reproductive Toxicology* **28**, 152–160 (2009).

11. CDCTobaccoFree. 2014 SGR: The Health Consequences of Smoking—50 Years of Progress. *Centers for Disease Control and Prevention* https://www.cdc.gov/tobacco/data_statistics/sgr/50th-anniversary/index.htm (2021).

12. Singh, S. P. *et al.* Prenatal Cigarette Smoke Decreases Lung cAMP and Increases Airway

Hyperresponsiveness. *Am J Respir Crit Care Med* **168**, 342–347 (2003).

13. Lawder, R., Whyte, B., Wood, R., Fischbacher, C. & Tappin, D. M. Impact of maternal smoking

on early childhood health: a retrospective cohort linked dataset analysis of 697 003 children born

in Scotland 1997–2009. *BMJ Open* **9**, (2019).

14. Hertzman, C. & Wiens, M. Child development and long-term outcomes: a population health

perspective and summary of successful interventions. *Soc Sci Med* **43**, 1083–1095 (1996).

15. Aristizabal, M. J. *et al.* Biological embedding of experience: A primer on epigenetics. *PNAS*

(2019) doi:10.1073/pnas.1820838116.

16. Demetriou, C. A. *et al.* Biological embedding of early-life exposures and disease risk in humans: a

role for DNA methylation. *Eur. J. Clin. Invest.* **45**, 303–332 (2015).

17. Cunliffe, V. T. Experience-sensitive epigenetic mechanisms, developmental plasticity, and the

biological embedding of chronic disease risk. *WIREs Systems Biology and Medicine* **7**, 53–71

(2015).

18. Kim, J. K., Samaranayake, M. & Pradhan, S. Epigenetic mechanisms in mammals. *Cell. Mol. Life*

Sci. **66**, 596 (2008).

19. Joubert, B. R. *et al.* 450K epigenome-wide scan identifies differential DNA methylation in

newborns related to maternal smoking during pregnancy. *Environmental health perspectives* **120**,

1425–1431 (2012).

20. Joubert, B. R. *et al.* DNA Methylation in Newborns and Maternal Smoking in Pregnancy:

Genome-wide Consortium Meta-analysis. *American journal of human genetics* **98**, 680–696

(2016).

21. Richmond, R. C. *et al.* Prenatal exposure to maternal smoking and offspring DNA methylation

across the lifecourse: findings from the Avon Longitudinal Study of Parents and Children

(ALSPAC). *Hum Mol Genet* **24**, 2201–2217 (2015).

22. Wiklund, P. *et al.* DNA methylation links prenatal smoking exposure to later life health outcomes

in offspring. *Clinical Epigenetics* **11**, 97 (2019).

23. Lee, K. W. K. *et al.* Prenatal Exposure to Maternal Cigarette Smoking and DNA Methylation:

Epigenome-Wide Association in a Discovery Sample of Adolescents and Replication in an

Independent Cohort at Birth through 17 Years of Age. *Environ Health Perspect* **123**, 193–199

(2015).

24. Tehranifar, P. *et al.* Maternal cigarette smoking during pregnancy and offspring DNA methylation

in midlife. *Epigenetics : official journal of the DNA Methylation Society* **13**, 129–134 (2018).

25. Richmond, R. C., Suderman, M., Langdon, R., Relton, C. L. & Davey Smith, G. DNA methylation

as a marker for prenatal smoke exposure in adults. *Int J Epidemiol* **47**, 1120–1130 (2018).

26. Rauschert, S. *et al.* Maternal Smoking During Pregnancy Induces Persistent Epigenetic Changes

Into Adolescence, Independent of Postnatal Smoke Exposure and Is Associated With

Cardiometabolic Risk. *Frontiers in Genetics* **10**, (2019).

27. Suter, M. *et al.* IN UTERO TOBACCO EXPOSURE EPIGENETICALLY MODIFIES

PLACENTAL CYP1A1 EXPRESSION. *Metabolism* **59**, 1481–1490 (2010).

28. Reddy, K. D. *et al.* Current Smoking Alters Gene Expression and DNA Methylation in the Nasal

Epithelium of Patients with Asthma. *Am J Respir Cell Mol Biol* **65**, 366–377 (2021).

29. Breton, C. V. *et al.* Prenatal tobacco smoke exposure affects global and gene-specific DNA

methylation. *Am J Respir Crit Care Med* **180**, 462–467 (2009).

30. Flom, J. D. *et al.* Prenatal smoke exposure and genomic DNA methylation in a multiethnic birth

cohort. *Cancer Epidemiol Biomarkers Prev* **20**, 2518–2523 (2011).

31. Neophytou, A. M. *et al.* In utero tobacco smoke exposure, DNA methylation, and asthma in Latino

children. *Environmental Epidemiology* **3**, e048 (2019).

32. Sherrill, D. L. *et al.* Longitudinal effects of passive smoking on pulmonary function in New

Zealand children. *Am Rev Respir Dis* **145**, 1136–1141 (1992).

33. Knopik, V. S. *et al.* Contributions of parental alcoholism, prenatal substance exposure, and genetic

transmission to child ADHD risk: a female twin study. *Psychol. Med.* **35**, 625–635 (2005).

34. Agrawal, A. *et al.* Correlates of cigarette smoking during pregnancy and its genetic and

environmental overlap with nicotine dependence. *Nicotine Tob Res* **10**, 567–578 (2008).

35. Vardavas, C. I. *et al.* The independent role of prenatal and postnatal exposure to active and passive

smoking on the development of early wheeze in children. *The European respiratory journal* **48**,

115–124 (2016).

36. Joad, J. P., Ji, C. M., Kott, K. S., Bric, J. M. & Pinkerton, K. E. In Utero and Postnatal Effects of

Sidestream Cigarette Smoke Exposure on Lung Function, Hyperresponsiveness, and

Neuroendocrine Cells in Rats. *Toxicology and Applied Pharmacology* **132**, 63–71 (1995).

37. Burke, H. *et al.* Prenatal and passive smoke exposure and incidence of asthma and wheeze:

systematic review and meta-analysis. *Pediatrics* **129**, 735–744 (2012).

38. Eskenazi, B. & Castorina, R. Association of prenatal maternal or postnatal child environmental

tobacco smoke exposure and neurodevelopmental and behavioral problems in children.

Environmental Health Perspectives **107**, 991–1000 (1999).

39. Tomczyk, M. M. *et al.* Mitochondrial Sirtuin-3 (SIRT3) Prevents Doxorubicin-Induced Dilated

Cardiomyopathy by Modulating Protein Acetylation and Oxidative Stress. *Circulation: Heart*

Failure **15**, e008547 (2022).

40. Ma, Y. *et al.* Sustained suppression of IL-13 by a vaccine attenuates airway inflammation and

remodeling in mice. *Am J Respir Cell Mol Biol* **48**, 540–549 (2013).

41. Ryu, M. H. *et al.* Chronic exposure to perfluorinated compounds: Impact on airway

hyperresponsiveness and inflammation. *Am J Physiol Lung Cell Mol Physiol* **307**, L765-774

(2014).

42. MUSCLE: multiple sequence alignment with improved accuracy and speed | IEEE Conference

Publication | IEEE Xplore. https://ieeexplore.ieee.org/abstract/document/1332560.

43. Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time

quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**, 402–408 (2001).

44. Wang, R. *et al.* Long non-coding RNA HOX transcript antisense RNA promotes expression of

14-3-3σ in non-small cell lung cancer. *Experimental and Therapeutic Medicine* **14**, 4503–4508

(2017).

45. Leong, J. W. *et al.* The Elimination Half-Life of Urinary Cotinine in Children of Tobacco-

Smoking Mothers. *Pulmonary Pharmacology & Therapeutics* **11**, 287–290 (1998).

46. Benowitz, N. L. Cotinine as a Biomarker of Environmental Tobacco Smoke Exposure.

Epidemiologic Reviews **18**, 188–204 (1996).

47. Camlin, N. J. *et al.* Maternal Smoke Exposure Impairs the Long-Term Fertility of Female

Offspring in a Murine Model1. *Biology of Reproduction* **94**, 39, 1–12 (2016).

48. Mays, C. E. Effects of Sidestream Smoke on Pregnant Mice and their Offspring. *Proceedings of*

the Indiana Academy of Science **95**, 529–536 (1985).

49. Esposito, E. R., Horn, K. H., Greene, R. M. & Pisano, M. M. An animal model of cigarette smoke-

induced in utero growth retardation. *Toxicology* **246**, 193–202 (2008).

50. Melgert, B. N. *et al.* Short-Term Smoke Exposure Attenuates Ovalbumin-Induced Airway

Inflammation in Allergic Mice. *Am J Respir Cell Mol Biol* **30**, 880–885 (2004).

51. Blacquière, M. J. *et al.* Maternal smoking during pregnancy induces airway remodelling in mice

offspring. *European Respiratory Journal* **33**, 1133–1140 (2009).

52. Lei, Y., Cao, Y.-X., Xu, C.-B. & Zhang, Y. The Raf-1 inhibitor GW5074 and dexamethasone

suppress sidestream smoke-induced airway hyperresponsiveness in mice. *Respir Res* **9**, 71 (2008).

53. Dere, E., Forgacs, A. L., Zacharewski, T. R. & Burgoon, L. D. Genome-Wide Computational

Analysis of Dioxin Response Element Location and Distribution in the Human, Mouse, and Rat

Genomes. *Chem. Res. Toxicol.* **24**, 494–504 (2011).

54. Sun, Y. V., Boverhof, D. R., Burgoon, L. D., Fielden, M. R. & Zacharewski, T. R. Comparative

analysis of dioxin response elements in human, mouse and rat genomic sequences. *Nucleic Acids*

Research **32**, 4512–4523 (2004).

55. Metzger, M. J., Halperin, A. C., Manhart, L. E. & Hawes, S. E. Association of Maternal Smoking

during Pregnancy with Infant Hospitalization and Mortality Due to Infectious Diseases. *Pediatr*

Infect Dis J **32**, e1–e7 (2013).

56. Agrawal, A. *et al.* The Effects of Maternal Smoking During Pregnancy on Offspring Outcomes.

Prev Med **50**, 13 (2010).

57. Janbazacyabar, H. *et al.* The Effects of Maternal Smoking on Pregnancy and Offspring: Possible

Role for EGF? *Frontiers in Cell and Developmental Biology* **9**, (2021).

58. Suter, M. A. & Aagaard, K. M. The impact of tobacco chemicals and nicotine on placental

development. *Prenat Diagn* **40**, 1193–1200 (2020).

59. Suter, M. *et al.* Maternal tobacco use modestly alters correlated epigenome-wide placental DNA

methylation and gene expression. *Epigenetics* **6**, 1284–1294 (2011).

60. Luck, W., Nau, H., Hansen, R. & Steldinger, R. Extent of nicotine and cotinine transfer to the

human fetus, placenta and amniotic fluid of smoking mothers. *Dev Pharmacol Ther* **8**, 384–395

(1985).

61. Beyer, D., Mitfessel, H. & Gillissen, A. Maternal smoking promotes chronic obstructive lung

disease in the offspring as adults. *Eur J Med Res* **14**, 27–31 (2009).

62. Toppila-Salmi, S. *et al.* Maternal smoking during pregnancy affects adult onset of asthma in

offspring: a follow up from birth to age 46 years. *European Respiratory Journal* **55**, (2020).

63. Gilliland, F. D., Li, Y.-F. & Peters, J. M. Effects of Maternal Smoking during Pregnancy and

Environmental Tobacco Smoke on Asthma and Wheezing in Children. *Am J Respir Crit Care*

Med **163**, 429–436 (2001).

64. Neuman, Å. *et al.* Maternal smoking in pregnancy and asthma in preschool children: a pooled

analysis of eight birth cohorts. *Am J Respir Crit Care Med* **186**, 1037–1043 (2012).

65. Hayatbakhsh, M. R. *et al.* Maternal smoking during and after pregnancy and lung function in early

adulthood: a prospective study. *Thorax* **64**, 810–814 (2009).

66. Meyer, K. F. *et al.* The fetal programming effect of prenatal smoking on Igf1r and Igf1

methylation is organ- and sex-specific. *Epigenetics* **12**, 1076–1091 (2017).

67. Lin, P.-I., Shu, H. & Mersha, T. B. Comparing DNA methylation profiles across different tissues

associated with the diagnosis of pediatric asthma. *Sci Rep* **10**, 151 (2020).

68. Song, F. *et al.* Tissue specific differentially methylated regions (TDMR): Changes in DNA

methylation during development. *Genomics* **93**, 130–139 (2009).

69. Novakovic, B. *et al.* Postnatal stability, tissue, and time specific effects of AHRR methylation

change in response to maternal smoking in pregnancy. *Epigenetics : official journal of the DNA*

Methylation Society **9**, 377–386 (2014).

70. Armstrong, D. A., Lesseur, C., Conradt, E., Lester, B. M. & Marsit, C. J. Global and gene-specific

DNA methylation across multiple tissues in early infancy: implications for children's health

research. *FASEB J* **28**, 2088–2097 (2014).

71. Zhou, J. *et al.* Tissue-specific DNA methylation is conserved across human, mouse, and rat, and

driven by primary sequence conservation. *BMC Genomics* **18**, 724 (2017).

72. Wan, J. *et al.* Characterization of tissue-specific differential DNA methylation suggests distinct

modes of positive and negative gene expression regulation. *BMC Genomics* **16**, 49 (2015).

73. Herzog, E. M. *et al.* The tissue-specific aspect of genome-wide DNA methylation in newborn and

placental tissues: implications for epigenetic epidemiologic studies. *Journal of Developmental*

Origins of Health and Disease **12**, 113–123 (2021).

74. Herzog, E. *et al.* Tissue-specific DNA methylation profiles in newborns. *Clinical epigenetics* **5**,

(2013).

75. Chellian, R. *et al.* Rodent models for nicotine withdrawal. *J Psychopharmacol* **35**, 1169–1187

(2021).

76. McLaughlin, I., Dani, J. A. & De Biasi, M. Nicotine Withdrawal. in *The Neuropharmacology of*

Nicotine Dependence (eds. Balfour, D. J. K. & Munafò, M. R.) 99–123 (Springer International

Publishing, 2015). doi:10.1007/978-3-319-13482-6_4.

77. Paolini, M. & De Biasi, M. Mechanistic insights into nicotine withdrawal. *Biochemical*

Pharmacology **82**, 996–1007 (2011).

78. Jacobsen, L. K. *et al.* Effects of smoking and smoking abstinence on cognition in adolescent

tobacco smokers. *Biological Psychiatry* **57**, 56–66 (2005).

79. Tager, I. B., Weiss, S. T., Muñoz, A., Rosner, B. & Speizer, F. E. Longitudinal Study of the

Effects of Maternal Smoking on Pulmonary Function in Children. *New England Journal of*

Medicine **309**, 699–703 (1983).

80. McEvoy, C. T. & Spindel, E. R. Pulmonary Effects of Maternal Smoking on the Fetus and Child:

Effects on Lung Development, Respiratory Morbidities, and Life Long Lung Health. *Paediatric*

Respiratory Reviews **21**, 27–33 (2017).

81. Goksör, E., Åmark, M., Alm, B., Gustafsson, P. M. & Wennergren, G. The impact of pre- and post-natal smoke exposure on future asthma and bronchial hyper-responsiveness. *Acta Paediatrica* **96**, 1030–1035 (2007).

82. Xiao, R., Noël, A., Perveen, Z. & Penn, A. L. In utero exposure to second-hand smoke activates pro-asthmatic and oncogenic miRNAs in adult asthmatic mice. *Environ Mol Mutagen* **57**, 190–199 (2016).

83. Leon Hsu, H.-H. *et al.* Prenatal Particulate Air Pollution and Asthma Onset in Urban Children. Identifying Sensitive Windows and Sex Differences. *Am J Respir Crit Care Med* **192**, 1052–1059 (2015).

84. Xiao, R. *et al.* In utero exposure to second-hand smoke aggravates the response to ovalbumin in adult mice. *Am J Respir Cell Mol Biol* **49**, 1102–1109 (2013).

85. Noël, A. *et al.* Sex-specific lung functional changes in adult mice exposed only to second-hand smoke in utero. *Respir Res* **18**, 104 (2017).

86. Nielsen, H. C. Androgen receptors influence the production of pulmonary surfactant in the testicular feminization mouse fetus. *J Clin Invest* **76**, 177–181 (1985).

87. Torday, J. S. & Nielsen, H. C. The Sex Difference in Fetal Lung Surfactant Production. *Experimental Lung Research* **12**, 1–19 (1987).

88. Carey, M. A. *et al.* The impact of sex and sex hormones on lung physiology and disease: lessons from animal studies. *American Journal of Physiology-Lung Cellular and Molecular Physiology* **293**, L272–L278 (2007).

89. Thun, M. J. *et al.* Lung Cancer Death Rates in Lifelong Nonsmokers. *JNCI: Journal of the National Cancer Institute* **98**, 691–699 (2006).

90. Carey, M. A. *et al.* It's all about sex: gender, lung development and lung disease. *Trends in Endocrinology & Metabolism* **18**, 308–313 (2007).

91. Mandhane, P. J., Greene, J. M., Cowan, J. O., Taylor, D. R. & Sears, M. R. Sex Differences in Factors Associated with Childhood- and Adolescent-Onset Wheeze. *Am J Respir Crit Care Med* **172**, 45–54 (2005).

92. Almqvist, C., Worm, M., Leynaert, B. & working group of GA2LEN WP 2.5 Gender. Impact of gender on asthma in childhood and adolescence: a GA2LEN review. *Allergy* **63**, 47–57 (2008).

93. Sennhauser, F. H. & Kühni, C. E. Prevalence of respiratory symptoms in Swiss children: Is bronchial asthma really more prevalent in boys? *Pediatric Pulmonology* **19**, 161–166 (1995).

94. Cunningham, J., Dockery, D. W. & Speizer, F. E. Maternal Smoking during Pregnancy as a Predictor of Lung Function in Children. *American Journal of Epidemiology* **139**, 1139–1152 (1994).

95. Drummond, D. *et al.* Combined Effects of in Utero and Adolescent Tobacco Smoke Exposure on Lung Function in C57Bl/6J Mice. *Environ Health Perspect* **125**, 392–399 (2017).

96. Singh, S. P. *et al.* Maternal Exposure to Secondhand Cigarette Smoke Primes the Lung for Induction of Phosphodiesterase-4D5 Isozyme and Exacerbated Th2 Responses: Rolipram Attenuates the Airway Hyperreactivity and Muscarinic Receptor Expression but Not Lung Inflammation and Atopy. *The Journal of Immunology* **183**, 2115–2121 (2009).

97. Maideen, N. M. P. Tobacco smoking and its drug interactions with comedications involving CYP and UGT enzymes and nicotine. *World Journal of Pharmacology* **8**, 14–25 (2019).

98. Anttila, S. *et al.* Methylation of Cytochrome P4501A1 Promoter in the Lung Is Associated with Tobacco Smoking. *Cancer Research* **63**, 8623–8628 (2003).

99. Hankinson, O. The aryl hydrocarbon receptor complex. *Annu Rev Pharmacol Toxicol* **35**, 307–340 (1995).

100. Kim, J. Expression of cytochromes P450 1A1 and 1B1 in human lung from smokers, non-smokers, and ex-smokers. *Toxicology and Applied Pharmacology* **199**, 210–219 (2004).

101.   Rendic, S. & Di Carlo, F. J. Human cytochrome P450 enzymes: a status report summarizing

their reactions, substrates, inducers, and inhibitors. *Drug Metab Rev* **29**, 413–580 (1997).

102.   Okey, A. B. Enzyme induction in the cytochrome P-450 system. *Pharmacology &*

Therapeutics **45**, 241–298 (1990).

103.   Ohashi, H. *et al.* The aryl hydrocarbon receptor–cytochrome P450 1A1 pathway controls lipid

accumulation and enhances the permissiveness for hepatitis C virus assembly. *J Biol Chem* **293**,

19559–19571 (2018).

104.   Rendic, S. & Guengerich, F. P. Contributions of Human Enzymes in Carcinogen Metabolism.

ACS Publications https://pubs.acs.org/doi/pdf/10.1021/tx300132k (2012) doi:10.1021/tx300132k.

105.   Nebert, D. W. *et al.* Role of the aromatic hydrocarbon receptor and [Ah] gene battery in the

oxidative stress response, cell cycle control, and apoptosis. *Biochemical Pharmacology* **59**, 65–85

(2000).

106.   Uno, S., Sakurai, K., Nebert, D. W. & Makishima, M. Protective role of cytochrome P450 1A1

(CYP1A1) against benzo[a]pyrene-induced toxicity in mouse aorta. *Toxicology* **316**, 34–42 (2014).

107.   Uno, S. *et al.* Oral Benzo[a]pyrene in Cyp1 Knockout Mouse Lines: CYP1A1 Important in

Detoxication, CYP1B1 Metabolism Required for Immune Damage Independent of Total-Body

Burden and Clearance Rate. *Mol Pharmacol* **69**, 1103–1114 (2006).

108.   Neuhold, L. A., Shirayoshi, Y., Ozato, K., Jones, J. E. & Nebert, D. W. Regulation of mouse

CYP1A1 gene expression by dioxin: requirement of two cis-acting elements during induction.

Molecular and Cellular Biology **9**, 2378–2386 (1989).

109.   Gluckman, P. D., Hanson, M. A. & Low, F. M. Evolutionary and developmental mismatches

are consequences of adaptive developmental plasticity in humans and have implications for later

disease risk. *Philosophical Transactions of the Royal Society B: Biological Sciences* **374**,

20180109 (2019).

110. Frankenhuis, W. E., Nettle, D. & McNamara, J. M. Echoes of Early Life: Recent Insights From Mathematical Modeling. *Child Development* **89**, 1504–1518 (2018).

111. Godfrey, K. M., Lillycrop, K. A., Burdge, G. C., Gluckman, P. D. & Hanson, M. A. Epigenetic Mechanisms and the Mismatch Concept of the Developmental Origins of Health and Disease. *Pediatr Res* **61**, 5–10 (2007).

112. Lkhagvadorj, K. *et al.* Prenatal smoke exposure induces persistent Cyp2a5 methylation and increases nicotine metabolism in the liver of neonatal and adult male offspring. *Epigenetics* **15**, 1370–1385 (2020).

113. Zeng, Z. *et al.* Prenatal smoke effect on mouse offspring Igf1 promoter methylation from fetal stage to adulthood is organ and sex specific. *American Journal of Physiology-Lung Cellular and Molecular Physiology* **318**, L549–L561 (2020).

114. Dehmel, S. *et al.* Intrauterine smoke exposure deregulates lung function, pulmonary transcriptomes, and in particular insulin-like growth factor (IGF)-1 in a sex-specific manner. *Sci Rep* **8**, 7547 (2018).

115. Marwick, J. A. *et al.* Cigarette Smoke Alters Chromatin Remodeling and Induces Proinflammatory Genes in Rat Lungs. *Am J Respir Cell Mol Biol* **31**, 633–642 (2004).

116. Meldrum, D. R. *et al.* Adaptive and maladaptive mechanisms of cellular priming. *Ann Surg* **226**, 587–598 (1997).

117. Kentner, A. C., Cryan, J. F. & Brummelte, S. Resilience Priming: Translational models for understanding resiliency and adaptation to early-life adversity. *Dev Psychobiol* **61**, 350–375 (2019).

118. Hales, C. N. & Barker, D. J. P. The thrifty phenotype hypothesis: Type 2 diabetes. *British Medical Bulletin* **60**, 5–20 (2001).

119.   Gluckman, P. D. & Hanson, M. A. Living with the Past: Evolution, Development, and Patterns

of Disease. *Science* **305**, 1733–1736 (2004).

120.   Hsu, C.-N. & Tain, Y.-L. Animal Models for DOHaD Research: Focus on Hypertension of

Developmental Origins. *Biomedicines* **9**, 623 (2021).

121.   Lavigne, É. *et al.* Maternal exposure to ambient air pollution and risk of early childhood

cancers: A population-based study in Ontario, Canada. *Environ Int* **100**, 139–147 (2017).

122.   Erickson, A. C. & Sbihi, H. Biological embedding, the air we breathe, and carcinogenesis. *The*

Lancet Planetary Health **2**, e149–e150 (2018).

123.   Bloushtain-Qimron, N. *et al.* Cell type-specific DNA methylation patterns in the human breast.

Proceedings of the National Academy of Sciences **105**, 14076–14081 (2008).

124.   Gervin, K. *et al.* Cell type specific DNA methylation in cord blood: a 450K-reference data set

and cell count-based validation of estimated cell type composition. *Epigenetics : official journal of*

the DNA Methylation Society 0 (2016).

125.   Novak, P. *et al.* Cell-Type Specific DNA Methylation Patterns Define Human Breast Cellular

Identity. *PLOS ONE* **7**, e52299 (2012).

1097