Functional trade-offs: exploring the temporal response of field margin plant

communities to climate change and agricultural practices

Isis Poinas[1,2*], Christine N Meynard[1], Guillaume Fried[2]

[1] CBGP, INRAE, CIRAD, Institut Agro, IRD, Univ Montpellier, Montpellier, France

[2] Anses, Laboratoire de la Santé des Végétaux, Unité Entomologie et Botanique, 755 avenue du

campus Agropolis, CS30016, 34988 Montferrier-sur-Lez Cedex, France

* Corresponding author: Isis Poinas (isispoinas@gmail.com)

14   **Number of words in abstract:** 376

15   **Number of words in main text (Introduction to Conclusion):** 4845

16   **Number of cited references:** 54

17   **Number of tables:** 1

18   **Number of figures:** 7

Abstract

Over the past decades, agricultural intensification and climate change have led to vegetation shifts. However, functional trade-offs linking traits responding to climate and farming practices are rarely analyzed, especially on large-scale empirical studies. Here we used a standardized yearly monitoring effort of agricultural field margin flora at the national scale to assess the temporal response of diversity and functional traits to variations in climate and intensity of agricultural practices. We examined temporal trends in climate (temperature, soil moisture), intensity of agricultural practices (herbicides, fertilization, margin management), plant species richness, and community-weighted means and variances of traits expected to vary both with climate and agricultural practices (e.g. seed mass, specific leaf area), across 555 sites in France between 2013 and 2021. We found clear temporal climatic trends (temperature increased while soil moisture decreased), whereas trends in agricultural practices were weak over the past decade, with only slight decreases in herbicides and margin management intensity. During the same period, functional changes in plant communities were significant, showing an increase of thermophilic species (including some Mediterranean species) with a conservative resource acquisition strategy (high stature, late and short flowering), mainly explained by climate change. The reduction in field margin management intensity resulted in a vegetation shift towards a more conservative strategy. In contrast, there was no impact from the slight temporal changes of practices conducted within the field (herbicides, fertilization). Our findings suggest that species adapted to climate change (including Mediterranean and conservative species) have temporally increased in proportion. Importantly, we identified functional trade-offs indicating that these species are also the most vulnerable to intensive agricultural practices, as they are less adapted to high levels of resources and disturbances. We put these results into the conceptual framework of Grime's CSR triangle and revealed a temporal decline of competitive and ruderal species in favor of stress-tolerant species better adapted to climate change. Choosing less intensive management can broaden the functional spectrum of agricultural plant communities, by maintaining the ability of stress-tolerant species selected by climate change to colonize habitats largely dominated by ruderals. Put together, these

46   results suggest that climate change and agricultural intensification could have synergistic negative

47   impacts on plant diversity in field margins, highlighting the need to study several biodiversity drivers

48   at the same time in anthropized landscapes.

50   **Keywords:** functional trade-offs, climate change, temporal variation, field margin, plant diversity,

51   agricultural practices, CSR strategies, Mediterranean species, phenology

Introduction

Since the 1950s, agricultural intensification has led to declining biodiversity (Emmerson et al., 2016), while climate change has caused notable changes in a wide range of taxa and habitats (Lovejoy, 2006). However, teasing out the relative importance of these two drivers on community trajectories can be quite challenging (Oliver & Morecroft, 2014). Since agricultural intensification took place in the 1950s in Europe, the main changes linked to agricultural practices (notably in terms of intensity in pesticide use and fertilization) in plant communities have likely already occurred (Lososová et al., 2004). For example, a meta-analysis considering 32 studies across Europe and covering the time period from 1939 to 2011 showed that weed species richness declined up to the 1980s, but has stabilized or even increased since then (Richner et al., 2015). Pesticide reduction plans have had so far little effect in France (Guichard et al., 2017), hindering the detection of temporal changes in biodiversity linked to changes in pesticide use. Conversely, short-term declines in species diversity due to climate change have been observed (e.g. Fonty et al., 2009), and recent temperature increases in France may impact plant communities similarly (Baude et al., 2022; Martin et al., 2019).

Temporal changes in plant communities cannot be discerned solely using taxonomic approaches due to the differing traits affected by resource availability and disturbance levels (Garnier & Navas, 2012); therefore, a functional approach provides an additional perspective to accurately understand these changes. This is particularly important in agroecosystems, where both resource (fertilization) and disturbance (herbicides, field margin management) gradients play crucial roles in shaping communities (Gaba et al., 2014; MacLaren et al., 2020). For instance, weeds with a ruderal strategy (low height and seed mass, long and early flowering, high SLA) are better adapted to agricultural disturbances, such as tillage, herbicides or management by mowing (Grime, 2006; Fried et al., 2022). At the same time, traits responding to agricultural practices can co-vary with other traits that are linked to resource acquisition, competitive ability, or climate. For example, seed mass, which is often used as a proxy for competitive ability, increases along soil fertility, temperature and solar radiation gradients (Fried et al., 2022;

Murray et al., 2004). Furthermore, correlations among different traits may represent trade-offs that

impact community adaptation (Díaz et al., 2016; Wright et al., 2004). Because of these trade-offs,

selective pressures due to one driver (e.g. climate change) may impact community properties along

the other (e.g. agricultural intensification). In this context, Grime (1977) proposed a framework called

the CSR triangle, which defines three strategies (competitiveness, stress-tolerance, and ruderality)

along two axes of variation (resource and disturbance). These strategies are correlated to multiple

traits and have proven useful to understand plant community dynamics in agroecosystems (Fried et

al., 2022). As traits responding to climate and agricultural practices may co-vary (Garnier & Navas,

2012), it can be difficult to identify the main drivers behind community changes, making this

framework potentially useful to understand such trade-offs.

To understand the complex interactions between climate change and agricultural practices, it is thus

essential to examine temporal changes in species trait distribution. For example, in French wheat

fields, ruderal species increased their frequency between the 1970s and 2000s, potentially due to their

ability to escape recurrent disturbances, such as herbicide applications (Fried et al., 2012). Inter-annual

variations in specific leaf area, leaf dry matter content and plant height are related to nitrogen supply

(Borgy et al., 2017; Gaba et al., 2014), while increased precipitations push the foliar economic spectrum

towards more acquisitive species (i.e. with higher SLA; Wheeler et al., 2023). Additionally, mean

thermal preference of plant communities, as well as their phenology, can vary over time in response

to temperature changes, even over relatively short periods (Bellard et al., 2012; Martin et al., 2019).

These temporal variations in functional traits reveal patterns that cannot be assessed solely with

space-for-time substitution.

In this study, we aimed at deciphering how inter-annual temporal variations and temporal trends in

climate (temperature, soil moisture) and agricultural practices (frequency of herbicide use, margin

management and nitrogen dose in fertilizers) in France structure species richness, trait composition

and ecological strategies of field margin plant communities. We studied the herbaceous field margin,

which represents the uncultivated vegetated area located between the cultivated strip and the

adjacent habitat. Using a standardized national monitoring effort spanning 9 years (2013-2021) in 555

agricultural field margins covering continental France, our study stands as one of the first to investigate

the temporal trends in agricultural practices and climate, and explore the response of species richness,

trait composition and ecological strategies to these trends at such extensive scales. We hypothesized

that plant traits sensitive to temperature and soil moisture will co-vary with temporal warming trends

while agricultural practices would have a comparatively weaker temporal influence on plant

communities, as we did not expect clear temporal trends in these practices. We also expected a limited

impact of agricultural practices on margin plant communities, because field margins only receive a

small amount of nitrogen and herbicides drifting from neighboring plots. Furthermore, we explored

the connection between Grime's CSR strategies, climate and farming practices. Considering that these

strategies are linked to resource and disturbance levels, we hypothesized that they would respond to

climate factors (particularly reduced soil moisture) and agricultural practices (disturbance and

resource provision through fertilization). On top of the national analyses, and because this dataset

includes the Mediterranean flora, which has been shown to respond more strongly to some

agricultural filters (Poinas et al., 2023), we included analyses separating this region from the rest of

France. We also separated vineyards from annual crops, because vineyards include very different

management practices and no crop rotation (Metay et al., 2022). Finally, we also analyzed annual plant

species separately, as they may respond more rapidly to environmental changes (Martin et al., 2019;

Fitter & Fitter, 2002).

Materials and methods

Vegetation surveys

We used vegetation data from the 500-ENI network, which is funded by the French Ministry of Agriculture (see details in Andrade et al., 2021) and monitored 555 agricultural field margins across continental France between 2013 and 2021 (with some site turnover) (**Fig. 1**). These survey sites represented three main crop types **(Appendix A, Fig. SA. 1)**: annual crops (with winter wheat or maize as the main crop production in the rotation), market gardening crops (mainly lettuce) and vineyards. The proportion of sites under organic farming was roughly 20%, but agricultural practices covered a wide range of pesticide application, fertilizers and soil management. Within each survey site, plant species were identified in ten 1 m² quadrats along the field margin (**Appendix A**, **Fig. SA.2**). Presence-absence of each species was recorded for each quadrat, which provided a frequency of occurrence from 0 to 10 in each field margin, used here as an index of relative abundance. Surveys were performed once per year at peak flowering (between the end of April and the beginning of August, depending on the region). At the national scale, this represented 4172 observations (year x site), leading to the identification of 852 taxa. Because observers changed among sites and over time (312 observers in total, each observer following on average 5 distinct sites during 4 years) and did not have the same level of expertise, we constrained our analyses to a subset of 142 focal species (Andrade et al., 2021) which are expected to be known by all the observers (and thus removing 11% of the total abundances).

Fig. 1. (A) Distribution map of the 555 field margins monitored at least one year between 2013 and

2021 in France. The black lines represent the limits of French departments. Orange: sites in

Mediterranean zone (n = 57), blue: sites in Continental zone (n = 498). The contours of the

Mediterranean zone (MZ) were derived from the Mediterranean zone and Corsica as defined in the

VégétalLocal map (Office français de la biodiversité, 2021); the rest of France will be referred to here

as Continental zone (CZ). (B) Subsets of data used in additional analyses: i) the regional scale splits the

MZ from the CZ; ii) annual crops included rotations based on wheat, maize and market gardening crops

(n = 450); perennial crops only included vineyards (n = 105); iii) annual plants (n = 61) opposed to

perennials (n = 79).

Climatic and agricultural variables

We gathered two types of explanatory variables: the first came directly from the 500-ENI network and

reflects agricultural practices assessed directly on the monitoring sites; the second one included

meteorological data from an external database (see below). Here, we chose not to include landscape

factors, as a previous study on the same dataset demonstrated that landscape variables account for a

negligible proportion of variance at the national scale, in contrast to climate (Poinas et al., 2023).

Agricultural practices were reported yearly from interviews of farmers into a standardized online

database. Data collected relate to fertilization, herbicide use and field margin management (mainly

mowing of vegetation). Daily meteorological data were extracted from the SAFRAN climate model of

Météo France, with a resolution of 8 km (Le Moigne, 2002). Meteorological data were averaged over

a one-year window prior to each floristic observation, while agricultural data were summed over the

same period (**Table 1**). We selected variables that were weakly correlated (Spearman correlation <

0.65, **Appendix B**) and have been shown to influence plant communities in previous studies (**Table 1,**

see **Appendix C** for the choice of variables).

Table 1. List of explanatory factors (blue), functional traits and ecological requirements (green) and response variables (red) with their abbreviations, units

and calculation. We have illustrated by arrows the expected link of each factor and trait to the agricultural resource (fertilization) and disturbance gradient

(herbicides and margin management), and to the climatic gradient (drought and increasing temperature). We also used arrows to illustrate the expected

direction of variation of these gradients within a year (i.e. climatic and agricultural changes according to the date of observation). Horizontal arrows indicate

contradictory findings in the literature (see **Appendix C** for the references).

Factors **Functional traits** **Response variables**	**Abbreviations**	**Units**	**Index used**	**Hypothesis of response to the temporal agricultural gradient:** **Disturbance (Di) and** **Resource (R)**	**Hypothesis of response to the temporal climatic gradient:** **Drought (Dr) and** **Temperature (T)**
Temperature	TEMP	°C	Annual mean of daily values		
Soil moisture	MOI	%	Annual mean of daily values		
Dose of nitrogen (fertilization)	N_DOSE	kg/ha	For each application: $$Dose = \frac{Quantity \times N\ of\ the\ formulation}{100}$$ Doses were summed one year before observation		
Number of herbicide treatments	HERB		Treatments were summed one year before observation		
Number of management events	MAN		An event corresponds to the occurrence of vegetation management at a given time, most often by mowing or grinding. Management events were summed one year before observation.		
Date of observation	DATE	Julian days		↘ ↘	↗ ↗

Specific leaf area	SLA	$m^2\ kg^{-1}$		↗	↘
				↗	→?
Maximum plant height	HEIGHT	m		↘	→?
				↗	→?
Seed mass	SM	g		↘	↗
				↗	↗
Flowering onset	FLOW_ON	months		↘	↗
					↗
Flowering duration	FLOW_DUR	Months		↗	↘
					↘
Ellenberg indicator for light, temperature, continentality,	EIV_L, EIV_T, EIV_K, EIV_F, EIV_R, EIV_N		High values of Ellenberg-L reflect heliophilous species (preference for light), Ellenberg-T, thermophilous species (preference for high temperatures),	EIV_N: ↗	EIV_F: ↘

moisture, pH and nutrients			Ellenberg-K, continental species (opposed to oceanic and Mediterranean species), Ellenberg-F, hygrophilous species (preference for moisture), Ellenberg-R, calcareous species (preference for calcareous soils), Ellenberg-N, nitrophilous species (preference for fertile soils) (Ellenberg, 1974)	EIV_T: ↗ ↗	
Species richness	S		Number of species	→? →?	→? →?
Trait composition	See abbreviations above for each trait preceded by "CWM_"		Community weighted means (CWM) : community trait values weighted by species abundance (Lavorel et al., 2008)		
Trait divergence	See abbreviations above for each trait preceded by "CWV_"		Community weighted variances (CWV) : community trait variability around the average value (Sonnier et al., 2010)		
PCA traits - axis 1	PCA_1		Scores of sites or observations on the 1st axis of the PCA including CWM of all traits		
PCA traits - axis 2	PCA_2		Scores of sites or observations on the 2nd axis of the PCA including CWM of all traits		
Scores of competitive, stress-tolerant and ruderal strategies	CWM_C, CWM_S, CWM_R	%	Community weighted means (CWM) : community strategy values weighted by species abundance (Lavorel et al., 2008). Species scores for each strategy are computed from the leaf area, leaf dry matter content and specific leaf area (see the algorithm of Pierce et al., 2017).	CWM_C: ↘	CWM_C: ↘

CWM_S:

CWM_S:

CWM_R:

Plant functional traits

We extracted from external databases five functional traits and six species-level indices of ecological

requirements (i.e. Ellenberg values; Ellenberg, 1974), assumed to respond to agricultural or climatic

factors (**Table 1**, **Appendix B-C**). Functional traits were missing for four species, two of which could be

imputed from an average over other species of the same genus. The remaining two species were

removed from the analysis (representing 0.01% of the total abundances among the 142 species

considered).

To characterize plant communities, we calculated species richness, community-weighted means

(CWM) and community-weighted variances (CWV) of traits and ecological requirements for

observations with at least three species (59 out of 4172 observations were excluded). The computation

was performed using the R v.4.0.0 package $_{FD}$, function *dbFD* for CWM, with the following formulas:

$\text{CWM} = \sum_{i=0}^{n} p_i \times trait_i$ **(Eq. 1)**

$\text{CWV} = \sum_{i=0}^{n} p_i \times (trait_i - CWM)^2$ **(Eq. 2)**

Where p_i is the relative abundance, $trait_i$ is the value of trait for species *i*, and *n* is the total number

of species. To correct for correlation between CWV and species richness, we used a null model

approach, shuffling the abundances in the species matrix for species of the species pool, while keeping

the species x trait matrix unchanged (Bopp et al., 2022). The species pool was defined by the site,

allowing us to investigate temporal variations. This procedure keeps trait correlations, species richness

and total abundance in a site unchanged, while dissociating abundances from trait values (Bernard-

Verdier et al., 2012). To quantify the difference between observed and null CWV, we computed effect

sizes (**Appendix D**). A positive effect size denotes a divergence in trait values within the community

(convergence for negative effect size). These effect sizes (and not the raw CWV) were used in our

analyses and referred to as CWV in the subsequent sections. We performed a normed PCA on the

CWM of traits to classify each community based on its average trait combination or ecological strategy,

which is reflected by its position on the first two axes.

Plant functional strategies

According to Grime (1988), stress (i.e. a shortage of resources such as nutrients, water and light) and

disturbance (i.e. the partial or total destruction of plant biomass) determine three main plant

strategies representing combinations of traits that are viable under conditions of low disturbances and

high resources (competitor, C), low disturbances and low resources (stress-tolerant, S) or high

disturbances and high resources (ruderal, R). Originally developed to classify individual plant species

into strategies, Grime's theory can be useful to interpret functional changes in plant communities,

especially in the context of global changes where vegetation is subject to harsher climatic conditions

(more droughts) and various levels of agricultural disturbances.

To assess these strategies, we extracted the CSR scores for 119 out of 142 focal species from Pierce et

al. (2017). CWM of CSR scores were computed by observation and were added to the PCA on the CWM

of traits as supplementary variables. They were plotted on a CSR triangle to illustrate temporal trends

in strategies.

Temporal analyses of plant communities

The general framework of analyses is presented in **Fig. 2**. First, we checked if there was a temporal

trend on the raw variables, and then we used climate and agricultural practices as predictors for the

different response variables. In all cases, we used generalized additive mixed models (GAMM) to

account for repeated measures at a site, with a Gaussian distribution in most cases (but see **Appendix**

E, **Table SE.2**), and site identity as a random effect. Observer bias was accounted for by including the

observer identity as a random term nested within sites. For each response variable (species richness,

CWM, CWV and CSR strategies) and explanatory factor (temperature, soil moisture, nitrogen dose,

herbicides and margin management), we built a first model with the year as a linear fixed effect. Then,

a second model was built for each response variable, where climate, agricultural practices and

observation date were linear explanatory factors. A first-order temporal autocorrelation structure

within sites was included (Box et al., 2015). We removed observations with missing values in climatic

and agricultural factors (1805 out of 4172 observations), and a few observations that distorted trait

distributions (**Appendix E**), resulting in varying observation numbers across models (see **Fig. 5**). We

repeated this analysis on subsets of data, including Mediterranean (MZ) vs Continental zones (CZ),

margins adjacent to annual crops vs vineyards, and annual vs perennial plant species (**Fig. 1**). For all

analyses, we chose a p-value threshold of 0.01 to focus on the effects for which our confidence level

was highest.

Fig. 2. General framework of our analyses involved two main steps. (1) Firstly, we investigated the

presence of any temporal trends in species richness, trait composition, ecological strategies, climatic

and agricultural factors, as this provides crucial insights for the subsequent analysis. Indeed, given that

we expected minimal temporal trends in agricultural practices, we did not expect significant temporal

changes in flora in response to practices. (2) Secondly, we explored how plant communities have

responded to the temporal changes in climate and agricultural practices.

Results

Temporal trends in climate, agricultural practices and plant communities

Temperatures have significantly increased by an average of 1.2°C over a decade (0.7°C in the Mediterranean Zone), while soil moisture has steadily declined (-14.1% by decade) (**Fig. 3**, and **Appendix F**). These trends differed between the Mediterranean zone (MZ) and the Continental zone (CZ), with the MZ experiencing a slower decline in soil moisture due to a high cumulative precipitation in 2019 (**Fig. 3**). Regarding agricultural practices, herbicides slightly decreased over time in vineyards (-0.9 application by decade; **Fig. 3**), with an even weaker trend in annual crops (-0.2 application by decade). Fertilization showed no significant temporal trend, except in vineyards where the cumulative dose of nitrogen has recently slightly increased (**Fig. 3**). The number of margin management events has decreased and particularly in the MZ (-0.5 by decade). Floristic surveys were conducted increasingly earlier in the season in the CZ (10.4 days earlier by decade) (**Appendix F**). Overall, there was a clear warming and drying trend in climate, but agricultural trends were weaker.

Plant species richness has slightly increased over time at the national scale (+0.1 species by decade, **Fig. 5**), even more in the MZ (+0.4 species by decade) and vineyards (+0.3 species by decade) and only for annual species (**Appendix H**). We saw an increase in the CWM of maximum height (+5.8 cm by decade), seed mass (+0.2 g by decade), flowering onset (+3.1 days by decade) and a decrease in flowering duration (-7.8 days by decade, **Fig. 5**). The requirements for light, temperature and pH have increased, while those for moisture and nitrogen have declined. CWV (i.e. computed by comparison with expected CWV in a community of same richness) have decreased for most traits indicating trait convergence, and particularly for phenological traits such as flowering onset and flowering duration (-3.6 and -2.6 days by decade respectively), while they have increased for the requirements for temperature, pH and continentality, indicating trait divergence.

Changes in functional traits were more pronounced in the MZ, particularly for the flowering onset (+8.8 days by decade) and duration (-18.9 days by decade; **Appendix F**). Conversely, changes in Ellenberg

values (environmental requirements) were only significant in the CZ and in annual crops. One

exception was the temperature (Ellenberg-T) and moisture (Ellenberg-F) requirements, which have

significantly changed in both the MZ and CZ. Interestingly, functional traits (and not environmental

requirements) showed a temporal trend mainly for annual species (**Appendix F**).

Fig. 3. Temporal changes in temperature, soil moisture, CWM of Ellenberg-F (moisture requirement),

number of herbicide applications and nitrogen dose in fertilizers. Red dots represent mean values. The

curves are from a GAM, with a smooth term on the year restricted to three effective degrees of

freedom. (A, C, E) National trend. (B, D, F) Trend by geographic area: CZ and MZ. (G, H) Trend by crop

type: annual (wheat, maize, lettuce) and perennial (vineyard). Significance of smooth terms is referred

as following: NS $p \geq 0.05$; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Functional trade-offs

The first PCA axis (named thereafter stress-tolerance axis, see **Appendix G, Fig. SG.1** for the correlation

of each CSR strategy with each axis) explained 29.9% of the variation in traits and revealed a gradient

from continental hygrophilous communities (high Ellenberg-K and F) associated with moist and

resource-rich environments (high Ellenberg-N), to Mediterranean xero-thermophilous stress-tolerant

communities (high Ellenberg-T and L, low Ellenberg-F) adapted to warm, arid and resource-poor

environments (**Fig. 4, Appendix G, Fig. SG.2**). Communities with continental species were more

nitrophilous (high Ellenberg-N), while Mediterranean communities had a higher seed mass. The second

PCA axis (named thereafter ruderal axis) explained 19.5% of the variation and contrasted stress-

tolerant/conservative communities adapted to low disturbance (low SLA, high stature, late and short

flowering) with ruderal/acquisitive communities adapted to high disturbance (high SLA, short stature,

early and long flowering).

Fig. 4. Normed PCA on CWM (first two axes) of functional traits and ecological requirements computed

by observation. The color of the dots indicates the region to which they belong and the density curve

highlights the concentration of data points in a given area. The correlations of traits to the PCA axes

are in **Appendix G**, **Fig. SG.1** and the PCA for annual and perennial species in **Fig. SG. 3**. The CWM of

strategies (in red) were added as supplementary variables. Abbreviations for CWM: CWM_SLA, specific

leaf area; CWM_HEIGHT, maximum plant height; CWM_SM, seed mass; CWM_FLOW_ON, flowering

onset; CWM_FLOW_DUR, flowering duration; CWM_EIV_L/T/K/F/R/N, requirement for

light/temperature/continentality/moisture/pH/nitrogen; CWM_C, competitive strategy; CWM_S,

stress-tolerant strategy; CWM_R, ruderal strategy.

Temporal analyses of plant communities

Climatic factors were the predominant drivers of changes in community trait composition, with high

R^2 for the temperature requirement (R^2 = 0.33) and stress-tolerance axis (see previous section, R^2 =

0.27, **Fig. 5**). Associations between each Ellenberg value and climatic factors opposed in a consistent

way Mediterranean communities to nitrophilous continental ones along the stress-tolerance axis (**Fig. 5**). Increasing temperature increased the CWM of seed mass and Ellenberg-T (requirement for temperature) and decreased the CWM of SLA. Increasing temperature led to more divergence in the requirement for temperature and moisture (compared to a community of same richness). Conversely, increasing soil moisture brought convergence in the requirement for temperature, continentality and soil pH. Increasing temperature and drought were also associated with shorter flowering duration (-2.2 days by °C and +0.15 days by % of soil moisture), and later flowering onset only for increasing temperature (+1.6 days by °C; **Fig. 5**).

Margin management was the agricultural practice with the largest impact on changes in community trait composition, with an increase in its frequency associated with more rurality (-1.2 cm in maximum height, -0.8 days in flowering onset and +2.1 days in flowering duration by management event). The date of observation also influenced changes in community trait composition, as a later observation was related to more conservative, competitive and continental communities, and to a decrease in species richness. Changes in the frequency of herbicide treatments had no significant effect, while an increasing annual nitrogen dose in fertilizers only slightly decreased the pH requirement (**Fig. 5**).

Fig. 5. Results of temporal models (GAMM) on the whole dataset, with response variables in rows and

explanatory factors in columns. The adjusted R^2, expressed as a percentage of variation, and the

number of observations are reported. Significance is indicated by colored cells, with a p-value

threshold of 0.01. Positive estimates are in red, negative estimates in blue, and the strength of the

relationship (based on the standardized estimates) is reflected by the lightness of the color (weaker

when lighter). It is important to note that the strength of the relationship can only be compared among

explanatory factors for a same response variable. Reported values are the raw estimates and can be

interpreted in the units of response and explanatory variables (e.g. an increase of 1°C in temperature

leads to an increase in 1.6 days in the CWM of flowering onset). Models with the year as explanatory

factor were run separately.

Results differed depending on the subset of data used (**Fig. 1**). In vineyards and the MZ, changes in soil

moisture did not have any influence on changes in species richness or community trait composition

(**Appendix H**) and temperature only increased the requirement for temperature (Ellenberg-T) and

decreased SLA. When margins were more frequently managed in the MZ, Mediterranean species

declined (decrease of temperature requirement and convergence towards higher values of

continentality, **Appendix H**). Increasing nitrogen dose tended to decrease the number of species in the

MZ, an effect also found on annual species. In vineyards, no agricultural effect was detected at the

national scale. Annuals were more impacted by climatic variations and seasonal effects (observation

date) than perennials, with high R^2 for temperature ($R^2 = 0.39$) and moisture ($R^2 = 0.32$) requirements.

Discussion

Our study is one of the first to provide empirical evidence that climate change is already resulting in

detectable functional changes in plant communities over a relatively short time interval of 10 years

(see also Martin et al., 2019) (**Fig. 6**). Climate change tended to favor the stress-tolerance strategy at

the expense of ruderality. These contrasting strategies highlight the functional trade-offs that prevent

field margin plants from simultaneously adapting to climate change and intensive agricultural

practices. Interestingly, reducing the frequency of margin management mimicked the impact of

climate change on community trait composition, although the trend was less pronounced. Practices

applied in the adjacent agricultural fields, including herbicide use and fertilization, had almost no effect

on changes in community trait composition.

Fig. 6. Synthesis of results based on temporal trends and temporal analyses of community response to

changes in climate and practices. Ecological strategies are based on PCA axes (Axis 1 for stress-

tolerance and Axis 2 for ruderality) and their associated traits. Explanatory factors and ecological

strategies are depicted with their temporal trend over a decade (arrows). The asterisks illustrate the

links between temporal trends in climate and practices and the resulting trends in communities,

inferred from the coefficients in **Fig. 5** (ns = non-significant). The size of the asterisks represents the

strength of the relationship, estimated from the number of impacted traits and standardized estimates

in models.

Climate as the main driver of temporal variations in field margin plant communities

Our analyses revealed a temporal shift towards more stress-tolerant and less ruderal communities,

primarily driven by climate (Díaz et al., 2016; Pakeman et al., 2009). Since meteorological variables

were extracted at a 8 km resolution, changes in soil moisture can be confidently attributed to climate

change and not to the effect of soil compaction due to cultivation. Increasing temperature and drought

favored more xerophilous and thermophilous species, with higher seed mass and lower nitrogen

requirement, thus shifting the position of communities along the stress-tolerance axis. Our results also

indicated that sites increasingly warmer and drier allowed for coexistence of a wider functional set of

species, suggesting an increased abundance of thermophilous species without any loss of cold-adapted

species. The increase in species richness over time provided additional support for this hypothesis.

The increase in temperature requirement at the community-level with rising temperatures was already

documented, but mainly by studies covering entire floras (regional or local species pool) and time

scales of several decades to a century (Salinitro et al., 2019; Tamis et al., 2005). We found that this

trend is now detectable over a short-term period of only nine years (Martin et al., 2019). Interestingly,

as in other recent studies (Duchenne et al., 2021; Martin et al., 2019), this trend was more pronounced

in northern France, while Mediterranean communities responded less to climate change (**Appendix**

H). On top of the fact that climatic trends observed in the MZ were weaker than in the CZ,

Mediterranean species are already adapted to drought and heat stress, and might be more resilient to

extinction risks (Thuiller et al., 2005). Because of their geographic position north of the Mediterranean

Sea, they might also experience some competitive release due to the lack of immigrants coming from

the south, and the northward shift of more temperate species (Duchenne et al. 2021).

Beyond the increase in temperature requirement, our models revealed additional temporal changes

related to climate change that align well with the existing literature, including a decrease in mean SLA

and an increase in mean seed mass and maximum height (Alarcón Víllora et al., 2019; Kühn et al.,

2021). These trait values (low SLA, high seed mass and height) are also known to be linked to less

intensive agriculture (Fried et al., 2012; Richner et al., 2015). In our models, we observed a similar

pattern, with less frequent margin management associated with a decrease along the ruderality axis

and an increase along the stress-tolerance axis. All of this suggests that climate change and the

evolution towards more extensive agricultural practices will select the same trait values towards more

stress-tolerant and less ruderal strategies (MacLaren et al., 2020).

Finally, temporal analyses showed additional phenological changes, suggesting that climate change

could increase the occurrence or abundance of late-flowering species, i.e. species that have high

temperature requirement to complete their life cycle (Peters et al., 2014). These phenological shifts

coincided with a decrease in trait variance, leading to trait convergence within communities. Critically,

such changes can reduce the ability of species to escape field margin management, which typically

favors species able to flower all-year-round, as expected with a ruderal strategy. As species will not be

able to advance their phenology indefinitely, this can ultimately result in species losses in the long-

term. However, farmers are likely to adapt the temporality of their practices to climate change,

mitigating some of these impacts.

Agricultural practices have a weaker impact on temporal community dynamics

Temporal variations in agricultural practices over the short-term had a weaker influence on field

margin plant communities than climatic variations (Alarcón Víllora et al., 2019; Fried et al., 2019). Field

margin management was the most impactful practice, affecting traits related to the ruderal syndrome

in a consistent way. This supports the idea that field margin management, as the only practice applied

directly in the margin, has a greater impact than herbicides and fertilization applied in the adjacent

agricultural fields, which are likely to have only collateral effects.

The lack of herbicide effect on community trait composition could arise because the number of

herbicide applications has varied little in recent years or because we have omitted some traits that

reflect herbicide tolerance (e.g. leaf cuticle thickness, hairiness). Also, reducing the intensity of

agricultural practices may not necessarily influence the trait composition of communities, because

agricultural intensification has already greatly reduced functional diversity, and highly diverse

landscapes would be required for some species to recolonize field margins.

Fertilization had minimal influence on changes in community trait composition, but reduced species

richness (Kleijn & Verbeek, 2000), an effect detected in the MZ and leading to the loss of some annual

Mediterranean species (Poinas et al., 2023). Due to functional trade-offs, we saw that nitrophilous

plant species were less thermophilous and more acidiphilous, which explains why nitrogen dose was

related to affinity for acidic soils in our models. Nitrogen dose remained constant over time, which

aligns with the weak change in global nitrophily levels in plant communities, suggesting that

eutrophication may no longer be the primary driver of changes in arable vegetation (Alignier, 2018;

Duchenne et al., 2021).

Fig. 7. Grime's CSR triangle depicting the temporal trajectory of community strategies between 2013

and 2021. To enhance clarity, we show only two levels of density curves, with each point representing

a specific site. The relative percentages of each strategy are depicted through the use of green, blue

and red colors (C = competitor, S = stress-tolerant and R = ruderal). Arrows indicate expected impacts

of climate change, disturbance, and resource levels. Temporal models (GAMM) applied to the CWM of

each strategy revealed significant decreases in ruderality (-1.14% by decade) and competitiveness (-

0.81% by decade) scores of communities, and a significant increase in stress-tolerance scores (+1.80%

by decade).

Functional trade-offs and implications for communities response to ongoing global changes

Our findings revealed that resource level (driven by fertilization) and climate vary the position of

communities along the stress-tolerance axis (see **Appendix I** for more results on the effect of

fertilization), while disturbance level (driven by field margin management) and climate vary the

position of communities along the ruderal axis (**Fig. 7**). This supports the view that functional trade-

offs are not only evident on a global scale as found by Wright et al. (2004) and Díaz et al. (2016), but

also occur within a narrower functional range (such as plants colonizing agricultural field margins). As

a result, agricultural intensification and climate change act in opposite ways on the trait composition

of field margin plant communities. Climate change favors species that are adapted to high

temperatures and drought, but not to intensive agriculture. It tends to expand the functional range for

traits related to stress-tolerance within communities (divergence), but reduces the functional range

for traits associated to ruderality (convergence). Conversely, agricultural disturbances select species

more sensitive to current climatic trends, without any particular trend in trait variance.

These trade-offs emphasize the need to consider the existing interactions between climate and

agricultural practices when predicting future community trajectories (Garnier et al., 2019; Pakeman et

al., 2009). Here, we acknowledge the difficulty in quantifying these interactions, particularly given the

limited changes observed in practices over time. However, the effects of practices were more

perceptible with analyses focused on spatial effects (**Appendix I** and see also Poinas et al., 2023),

allowing us to imagine main trends in community trajectories according to several scenarios (**Fig. 7**).

Accelerating climate change coupled with an agricultural abandonment and more extensive practices

in Europe (Miller et al., 2022; Peeters et al., 2021) will likely result in an increase in xero-thermophilous

and conservative species. However, a large part of these species are habitat specialists (e.g.

Mediterranean species as found in Munoz et al. (2017); Fried, Chauvel, et al., 2009) and have a high

affinity for calcareous soils, which will probably limit their expansion towards the CZ to restricted

calcareous areas, such as the Paris Basin. Areas where they are unable to colonize might suffer a

decrease in species richness, and this scenario could be worsen if current levels of agricultural

intensification are maintained or increased. Mediterranean species expanding in the northern half of

France could face severe agricultural intensification that would likely limit their expansion, while at the

same time ruderal species would become less frequent mostly because of drought. This highlights the

need to consider the conjunction of climate change and intensive agriculture when making future

predictions.

Conclusion

Our study highlights climate as the primary factor affecting field margin plant communities in France,

with increasing temperatures and decreasing soil moisture fostering Mediterranean, stress-tolerant

and conservative species, while negatively affecting ruderal species. In comparison, agricultural

practices had a limited effect on changes in species richness and trait composition, with field margin

management having the greatest impact. Our study suggests that the species selected by climate

change are poorly adapted to intensive farming, while the pool of species currently able to colonize

field margins is restricted to a limited functional range adapted to agricultural practices. The

persistence of intensive agricultural practices and accelerating climate change could thus have critical

consequences for the conservation of floristic diversity in agroecosystems. However, it is important to

consider the potential of adaptation of species, through intraspecific trait variability and phenotypic

plasticity (known to be particularly high in ruderal species, Baker, 1974), as it may enhance their

resilience to changing conditions. Our findings suggest a reduction in ruderality and an increase in

stress-tolerance according to Grime's strategies. Bopp (2023) highlighted a similar increase of stress-

tolerance in weeds in response to climate change, but did not observe a corresponding decrease in

ruderality. Further investigations are thus necessary to assess the generalizability of these results

across different habitats, including communities with broader or narrower functional niche, such as

weeds. Long-term monitoring programs are necessary to address other important research questions,

such as the potential time-lag in flora's response to environmental changes, the non-linearity in

temporal trends and the interactive effects between climate and agricultural changes. Finally, the

findings presented in this study call for rethinking our current agricultural model, urging us to prioritize

the development of agricultural practices that foster the creation of favorable microclimates while

minimizing local intensification. Promising approaches, such as agroforestry, hold the potential to align

agricultural production with biodiversity conservation goals by providing refuge habitats and

microclimate regulation.

Acknowledgements

The 500-ENI network is developed by the French Ministry of Agriculture under the Ecophyto

framework with funding from the French Biodiversity Agency (Office Français de la Biodiversité). We

would like to thank everyone that has collected data in the field, the farmers who provided information

on their practices, and everyone involved in the coordination of the 500-ENI data network.

Funding: This work was supported by an INRAE-ANSES thesis fellowship, an Ecophyto II+ project: GTP

500 ENI (OFB-21-1642), and the AgriBiodiv ANR-21-CE32-006-01.

Conflict of interest disclosure

All authors of this preprint declare that they have no financial conflict of interest with the content of

this article.

Appendix A-H. Supplementary data

Supplementary data associated with this article can be found, in the online version, at XXXXX.

References

Alarcón Víllora, R., Hernández Plaza, E., Navarrete, L., Sánchez, M. J., & Sánchez, A. M. (2019). Climate and tillage system drive weed communities' functional diversity in a Mediterranean cereal-legume rotation. *Agriculture, Ecosystems & Environment*, *283*, 106574. https://doi.org/10.1016/j.agee.2019.106574

Alignier, A. (2018). Two decades of change in a field margin vegetation metacommunity as a result of field margin structure and management practice changes. *Agriculture, Ecosystems & Environment*, *251*, 1–10. https://doi.org/10.1016/j.agee.2017.09.013

Andrade, C., Villers, A., Balent, G., Bar-Hen, A., Chadoeuf, J., Cylly, D., Cluzeau, D., Fried, G., Guillocheau, S., Pillon, O., Porcher, E., Tressou, J., Yamada, O., Lenne, N., Jullien, J., & Monestiez, P. (2021). A real-world implementation of a nationwide, long-term monitoring program to assess the impact of agrochemicals and agricultural practices on biodiversity. *Ecology and Evolution*, *11*(9), 3771–3793. https://doi.org/10.1002/ece3.6459

Baker, H. G. (1974). The Evolution of Weeds. *Annual Review of Ecology and Systematics*, *5*(1), 1–24. https://doi.org/10.1146/annurev.es.05.110174.000245

Baude, M., Colin, A., Calipel, C., Duvernoy, J., & Foussard, A. (2022). *Chiffres clés du climat France, Europe et Monde—Édition 2022* [Rapport ministériel]. Service des données et études statistiques - Ministère de la Transition écologique. https://www.statistiques.developpement-durable.gouv.fr/edition-numerique/chiffres-cles-du-climat-2022/pdf/chiffres-cles-du-climat-2022-integral.pdf

Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W., & Courchamp, F. (2012). Impacts of climate change on the future of biodiversity. *Ecology Letters*, *15*(4), 365–377. https://doi.org/10.1111/j.1461-0248.2011.01736.x

Bernard-Verdier, M., Navas, M.-L., Vellend, M., Violle, C., Fayolle, A., & Garnier, E. (2012). Community assembly along a soil depth gradient: Contrasting patterns of plant trait convergence and

divergence in a Mediterranean rangeland. *Journal of Ecology*, *100*(6), 1422–1433. https://doi.org/10.1111/1365-2745.12003

Bopp, M. (2023). *Dynamique spatio-temporelle des communautés d'espèces adventices en réponse aux pratiques de gestion des sols dans les vignobles: Une approche fonctionnelle* [Doctoral thesis, Montpellier, SupAgro]. https://www.theses.fr/s225831

Bopp, M.-C., Kazakou, E., Metay, A., & Fried, G. (2022). Relative importance of region, seasonality and weed management practice effects on the functional structure of weed communities in French vineyards. *Agriculture, Ecosystems & Environment*, *330*, 107892. https://doi.org/10.1016/j.agee.2022.107892

Borgy, B., Violle, C., Choler, P., Denelle, P., Munoz, F., Kattge, J., Lavorel, S., Loranger, J., Amiaud, B., Bahn, M., van Bodegom, P. M., Brisse, H., Debarros, G., Diquelou, S., Gachet, S., Jolivet, C., Lemauviel-Lavenant, S., Mikolajczak, A., Olivier, J., … Garnier, E. (2017). Plant community structure and nitrogen inputs modulate the climate signal on leaf traits. *Global Ecology and Biogeography*, *26*(10), 1138–1152. https://doi.org/10.1111/geb.12623

Box, G. E. P., Jenkins, G. M., Reinsel, G. C., & Ljung, G. M. (2015). *Time Series Analysis: Forecasting and Control*. John Wiley & Sons.

Díaz, S., Kattge, J., Cornelissen, J. H. C., Wright, I. J., Lavorel, S., Dray, S., Reu, B., Kleyer, M., Wirth, C., Colin Prentice, I., Garnier, E., Bönisch, G., Westoby, M., Poorter, H., Reich, P. B., Moles, A. T., Dickie, J., Gillison, A. N., Zanne, A. E., … Gorné, L. D. (2016). The global spectrum of plant form and function. *Nature*, *529*(7585). https://doi.org/10.1038/nature16489

Duchenne, F., Martin, G., & Porcher, E. (2021). European plants lagging behind climate change pay a climatic debt in the North, but are favoured in the South. *Ecology Letters*, *24*(6), 1178–1186. https://doi.org/10.1111/ele.13730

Ellenberg, H. (1974). Indicator values of vascular plants in central Europe. *Scripta Geobotanica*, *9*, 97.

Emmerson, M., Morales, M. B., Oñate, J. J., Batáry, P., Berendse, F., Liira, J., Aavik, T., Guerrero, I., Bommarco, R., Eggers, S., Pärt, T., Tscharntke, T., Weisser, W., Clement, L., & Bengtsson, J.

(2016). How Agricultural Intensification Affects Biodiversity and Ecosystem Services. In
Advances in Ecological Research (Vol. 55, pp. 43–97). Elsevier.
https://linkinghub.elsevier.com/retrieve/pii/S0065250416300204

Fitter, A. H., & Fitter, R. S. R. (2002). Rapid Changes in Flowering Time in British Plants. *Science*,
296(5573), 1689–1691. https://doi.org/10.1126/science.1071617

Fonty, E., Sarthou, C., Larpin, D., & Ponge, J.-F. (2009). A 10-year decrease in plant species richness on
a neotropical inselberg: Detrimental effects of global warming? *Global Change Biology*, *15*(10),
2360–2374. https://doi.org/10.1111/j.1365-2486.2009.01923.x

Fried, G., Blanchet, C., Cazenave, L., Bopp, M.-C., Kazakou, E., Metay, A., Christen, M., Alard, D., &
Cordeau, S. (2022). Consistent response of weeds according to Grime's CSR strategies along
disturbance and resource gradients in Bordeaux vineyards. *Weed Research*, *62*(5), 347–359.
https://doi.org/10.1111/wre.12549

Fried, G., Chauvel, B., & Reboud, X. (2009). A functional analysis of large-scale temporal shifts from
1970 to 2000 in weed assemblages of sunflower crops in France. *Journal of Vegetation Science*,
20(1), 49–58. https://doi.org/10.1111/j.1654-1103.2009.05284.x

Fried, G., Cordeau, S., Metay, A., & Kazakou, E. (2019). Relative importance of environmental factors
and farming practices in shaping weed communities structure and composition in French
vineyards. *Agriculture, Ecosystems & Environment*, *275*, 1–13.
https://doi.org/10.1016/j.agee.2019.01.006

Fried, G., Kazakou, E., & Gaba, S. (2012). Trajectories of weed communities explained by traits
associated with species' response to management practices. *Agriculture, Ecosystems &
Environment*, *158*, 147–155. https://doi.org/10.1016/j.agee.2012.06.005

Gaba, S., Fried, G., Kazakou, E., Chauvel, B., & Navas, M.-L. (2014). Agroecological weed control using
a functional approach: A review of cropping systems diversity. *Agronomy for Sustainable
Development*, *34*(1), 103–119. https://doi.org/10.1007/s13593-013-0166-5

Garnier, E., & Navas, M.-L. (2012). A trait-based approach to comparative functional plant ecology:

Concepts, methods and applications for agroecology. A review. *Agronomy for Sustainable*

Development, *32*(2), 365–399. https://doi.org/10.1007/s13593-011-0036-y

Garnier, E., Vile, D., Roumet, C., Lavorel, S., Grigulis, K., Navas, M.-L., & Lloret, F. (2019). Inter- and

intra-specific trait shifts among sites differing in drought conditions at the north western edge

of   the   Mediterranean   Region.   *Flora*,   *254*,   147–160.

https://doi.org/10.1016/j.flora.2018.07.009

Grime, J. P. (1977). Evidence for the existence of three primary strategies in plants and its relevance to

ecological and evolutionary theory. *The American Naturalist*, *111*(982), 1169–1194.

Grime, J. P. (1988). The C-S-R model of primary plant strategies—Origins, implications and tests. In L.

D. Gottlieb & S. K. Jain, *Plant Evolutionary Biology* (pp. 371–393). Springer Netherlands.

https://doi.org/10.1007/978-94-009-1207-6_14

Grime, J. P. (2006). Trait convergence and trait divergence in herbaceous plant communities:

Mechanisms and consequences. *Journal of Vegetation Science*, *17*(2), 255–260.

https://doi.org/10.1111/j.1654-1103.2006.tb02444.x

Guichard, L., Dedieu, F., Jeuffroy, M.-H., Meynard, J.-M., Reau, R., & Savini, I. (2017). Le plan Ecophyto

de réduction d'usage des pesticides en France: Décryptage d'un échec et raisons d'espérer.

Cahiers Agricultures, *26*(1). https://doi.org/10.1051/cagri/2017004

Kleijn, D., & Verbeek, M. (2000). Factors affecting the species composition of arable field boundary

vegetation. *Journal of Applied Ecology*, *37*(2), 256–266. https://doi.org/10.1046/j.1365-

2664.2000.00486.x

Kühn, N., Tovar, C., Carretero, J., Vandvik, V., Enquist, B. J., & Willis, K. J. (2021). Globally important

plant functional traits for coping with climate change. *Frontiers of Biogeography*, *13*(4).

https://doi.org/10.21425/F5FBG53774

Lavorel, S., Grigulis, K., McIntyre, S., Williams, N. S. G., Garden, D., Dorrough, J., Berman, S., Quétier,

F., Thébault, A., & Bonis, A. (2008). Assessing functional diversity in the field – methodology

matters! *Functional Ecology*, *22*(1), 134–147. https://doi.org/10.1111/j.1365-
2435.2007.01339.x

Le Moigne, P. (2002). *Description de l'analyse des champs de surface sur la France par le système*
SAFRAN (pp. 1–30) [Note de centre GMME]. Météo-France.

Lososová, Z., Chytrý, M., Cimalová, S., Kropáč, Z., Otýpková, Z., Pyšek, P., & Tichý, L. (2004). Weed
vegetation of arable land in Central Europe: Gradients of diversity and species composition.
Journal of Vegetation Science, *15*(3), 415–422. https://doi.org/10.1111/j.1654-
1103.2004.tb02279.x

Lovejoy, T. E. (2006). *Climate Change and Biodiversity*. Yale University Press.

MacLaren, C., Storkey, J., Menegat, A., Metcalfe, H., & Dehnen-Schmutz, K. (2020). An ecological future
for weed science to sustain crop production and the environment. A review. *Agronomy for*
Sustainable Development, *40*(4), 24. https://doi.org/10.1007/s13593-020-00631-6

Martin, G., Devictor, V., Motard, E., Machon, N., & Porcher, E. (2019). Short-term climate-induced
change in French plant communities. *Biology Letters*, *15*(7), 20190280.
https://doi.org/10.1098/rsbl.2019.0280

Metay, A., Garcia, L., Kazakou, E., & Fried, G. (2022). Effet des couverts sur la flore en viticulture.
Agronomie, Environnement & Sociétés, *12*(1). https://hal-montpellier-supagro.archives-
ouvertes.fr/hal-03776173

Miller, D., Legras, S., Barnes, A., Cazacu, M., Gava, O., Helin, J., Irvine, K., Kantelhardt, J., Landert, J.,
Latruffe, L., Mayer, A., Niedermayr, A., Povellato, A., Schaller, L., Schwarz, G., Smith, P., Vanni,
F., Védrine, L., Viaggi, D., … Vlahos, G. (2022). Creating Conditions for Harnessing the Potential
of Transitions to Agroecology in Europe and Requirements for Policy. *EuroChoices*, *21*(3), 72–
79. https://doi.org/10.1111/1746-692X.12374

Munoz, F., Fried, G., Armengot, L., Bourgeois, B., Bretagnolle, V., Chadoeuf, J., Mahaut, L., Plumejeaud,
C., Storkey, J., Violle, C., & Gaba, S. (2017). *Database of weeds in cultivation fields of France*

and UK, with ecological and biogeographical information [dataset]. Zenodo. https://doi.org/10.5281/zenodo.1112342

Murray, B. R., Brown, A. H. D., Dickman, C. R., & Crowther, M. S. (2004). Geographical gradients in seed mass in relation to climate. *Journal of Biogeography*, *31*(3), 379–388. https://doi.org/10.1046/j.0305-0270.2003.00993.x

Office français de la biodiversité. (2021). *Carte des régions d'origine—Référentiel technique de la marque Végétal local—Version 2021*. www.vegetal-local.fr

Oliver, T. H., & Morecroft, M. D. (2014). Interactions between climate change and land use change on biodiversity: Attribution problems, risks, and opportunities. *WIREs Climate Change*, *5*(3), 317–335. https://doi.org/10.1002/wcc.271

Pakeman, R. J., Lepš, J., Kleyer, M., Lavorel, S., Garnier, E., & Consortium, the V. (2009). Relative climatic, edaphic and management controls of plant functional trait signatures. *Journal of Vegetation Science*, *20*(1), 148–159. https://doi.org/10.1111/j.1654-1103.2009.05548.x

Peeters, A., Lefebvre, O., & Balogh, L. (2020). A Green Deal for implementing agroecological systems: Reforming the Common Agricultural Policy of the European Union. *Journal of Sustainable and Organic Agricultural Systems*, *70*, 83–93. https://doi.org/10.3220/LBF1610123299000

Peters, K., Breitsameter, L., & Gerowitt, B. (2014). Impact of climate change on weeds in agriculture: A review. *Agronomy for Sustainable Development*, *34*(4), 707–721. https://doi.org/10.1007/s13593-014-0245-2

Pierce, S., Negreiros, D., Cerabolini, B. E. L., Kattge, J., Díaz, S., Kleyer, M., Shipley, B., Wright, S. J., Soudzilovskaia, N. A., Onipchenko, V. G., van Bodegom, P. M., Frenette-Dussault, C., Weiher, E., Pinho, B. X., Cornelissen, J. H. C., Grime, J. P., Thompson, K., Hunt, R., Wilson, P. J., … Tampucci, D. (2017). A global method for calculating plant CSR ecological strategies applied across biomes world-wide. *Functional Ecology*, *31*(2), 444–457. https://doi.org/10.1111/1365-2435.12722

Poinas, I., Fried, G., Henckel, L., & Meynard, C. N. (2023). Agricultural drivers of field margin plant

communities are scale-dependent. *Basic and Applied Ecology*, *72*, 55–63.

https://doi.org/10.1016/j.baae.2023.08.003

Richner, N., Holderegger, R., Linder, H. P., & Walter, T. (2015). Reviewing change in the arable flora of

Europe: A meta-analysis. *Weed Research*, *55*(1), 1–13. https://doi.org/10.1111/wre.12123

Salinitro, M., Alessandrini, A., Zappi, A., & Tassoni, A. (2019). Impact of climate change and urban

development on the flora of a southern European city: Analysis of biodiversity change over a

120-year period. *Scientific Reports*, *9*(1). https://doi.org/10.1038/s41598-019-46005-1

Sonnier, G., Shipley, B., & Navas, M.-L. (2010). Quantifying relationships between traits and explicitly

measured gradients of stress and disturbance in early successional plant communities. *Journal*

of Vegetation Science, *21*(6), 1014–1024. https://doi.org/10.1111/j.1654-1103.2010.01210.x

Tamis, W. L. M., Zelfde, M. V., Der Meijden, R. V., & De Haes, H. A. U. (2005). Changes in Vascular Plant

Biodiversity in the Netherlands in the 20th Century Explained by their Climatic and other

Environmental Characteristics. *Climatic Change*, *72*(1), 37–56.

https://doi.org/10.1007/s10584-005-5287-7

Thuiller, W., Lavorel, S., Araújo, M. B., Sykes, M. T., & Prentice, I. C. (2005). Climate change threats to

plant diversity in Europe. *Proceedings of the National Academy of Sciences*, *102*(23), 8245–

8250. https://doi.org/10.1073/pnas.0409902102

Wheeler, G. R., Brassil, C. E., & Knops, J. M. H. (2023). Functional traits' annual variation exceeds

nitrogen-driven variation in grassland plant species. *Ecology*, *104*(2), e3886.

https://doi.org/10.1002/ecy.3886

Wright, I. J., Reich, P. B., Westoby, M., Ackerly, D. D., Baruch, Z., Bongers, F., Cavender-Bares, J., Chapin,

T., Cornelissen, J. H. C., Diemer, M., Flexas, J., Garnier, E., Groom, P. K., Gulias, J., Hikosaka, K.,

Lamont, B. B., Lee, T., Lee, W., Lusk, C., … Villar, R. (2004). The worldwide leaf economics

spectrum. *Nature*, *428*(6985). https://doi.org/10.1038/nature02403