Title:

Delayed formation of neural representations of space in aged mice

Authors:

Kelsey D. McDermott, M. Agustina Frechou, Jake T. Jordan, Sunaina S. Martin, J. Tiago

Gonçalves*

*Corresponding author. Email: tiago.goncalves@einsteinmed.edu

Affiliations:

Dominick P. Purpura Department of Neuroscience and Gottesmann Institute for Stem Cell Biology

and Regenerative Medicine, Albert Einstein College of Medicine, Bronx, NY

Teaser:

Representations of space in the aged mouse hippocampus are initially disrupted in novel
environments, but improve once environments become familiar.

Abstract:

Aging is associated with cognitive deficits, with spatial memory being very susceptible to

decline. The hippocampal dentate gyrus (DG) is important for processing spatial information in

the brain and is particularly vulnerable to aging, yet its sparse activity has led to difficulties in

assessing changes in this area. Using *in vivo* two-photon calcium imaging, we compared DG

neuronal activity and representations of space in young and aged mice walking on an unfamiliar

treadmill. We found that calcium activity was significantly higher and less tuned to location in

aged mice, resulting in decreased spatial information encoded in the DG. However, with

repeated exposure to the same treadmill, both spatial tuning and information levels in aged mice

became similar to young mice, while activity remained elevated. Our results show that spatial

27 representations of novel environments are impaired in the aged hippocampus and gradually

28 improve with increased familiarity. Moreover, while the aged DG is hyperexcitable, this does not

29 disrupt neural representations of familiar environments.

31
Main Text:

Introduction

As the world's population ages, it becomes more important to understand the cognitive effects of

aging and to develop therapeutic strategies for the aging brain. While healthy aging has milder

cognitive effects than neurodegenerative states, several aspects of cognitive function exhibit

decline (*1*). Memory deteriorates with age, with spatial memory being particularly susceptible to

deficits in older humans and animals (*2–4*). Older adults show orientation deficits (*5, 6*), although

they tend to perform better in familiar locations or tasks than in novel ones (*7–9*). This aging-

related decay is thought to be caused by vulnerabilities in the hippocampal circuits that mediate

spatial memory (*10, 11*). The dentate gyrus (DG) is one of the hippocampal subregions that are

most vulnerable to aging (*12–14*), and the inputs from the entorhinal cortex to DG are a main area

of dysfunction in both Alzheimer's Disease (AD) patients and AD mouse models (*15*). During

aging, there are reduced synaptic contacts from both the medial and lateral entorhinal cortex (*16*)

onto DG neurons, as well as reduced synaptic plasticity of these inputs (*17*). However, the is very

limited data about how DG activity changes with aging, primarily because neuronal activity in this

region tends to be too sparse for *in vivo* electrophysiological recordings (*18*).

Here we investigated whether aging is associated with impaired DG activity and spatial

representations, which could account for the spatial memory deficits seen in aged individuals.

Since the DG is upstream of other hippocampal areas, changes in this region could help elucidate

the causes of aging-related changes in neuronal activity (*19–22*) and neural representations of

space (*19*, *23*, *24*) found in some areas of the hippocampus but not in others. Using *in vivo* two-

photon microscopy we imaged calcium activity in a large population of DG neurons of young and

aged mice as the animals walked head-fixed on a treadmill to which they had never been exposed.

We found that aged mice have increased DG single-cell calcium activity and disrupted neural

representations of space upon their first exposure to the treadmill setting. However, further

imaging sessions on subsequent days showed that spatial representations become similar to

those of young mice as the animals familiarize themselves with the treadmill, whereas single-cell

activity remains elevated. Our results highlight the importance of novelty and familiarity to spatial

encoding in aged animals.

Results

To record neuronal activity in DG excitatory neurons, we injected young (3-4 month old) and aged

(21-26 month old) C57Bl6 mice with an AAV encoding the calcium indicator jRGECO1a (*25*) under

the CaMKII promoter (Fig. 1A). We then implanted the mice with a hippocampal imaging window

above the DG (*26*, *27*) to enable *in vivo* two-photon calcium imaging in this region (Fig. 1A,B).

After waiting 3-4 weeks to allow for viral expression and recovery from surgery, we imaged mice

on four consecutive days. During each imaging session the mice ran head-fixed along a treadmill

containing tactile cues with four distinct segments (Fig. 1A,C) that was manually moved (Fig. S1).

We recorded 10 minute videos of calcium activity and identified the regions corresponding to

active cells using Suite2p software (Fig. 1D). Previous studies found that hippocampal activity in

area CA3 becomes elevated during aging (*19*, *28*). We determined single-cell calcium activity by

measuring the area under the curve of transients and normalizing to the distance run by each

mouse (*29*) (Fig. 1E-G). We found that the aged group had higher activity levels than the young

group (Fig. 1E, p=0.00463, nested bootstrap), which aligns with previous reports of hyperactivity.

We also estimated the fraction of active neurons during each imaging session and found no

changes between groups (Fig. S2). We next investigated whether there were differences between

the spatial representations of young and aged DG neurons. Both young and aged DG neurons

have a variety of spatial responses, including place cells that only respond to a specific location

and neurons whose activity shows low spatial preference (Fig. 2A,B). We determined the spatial

Figure 1. DG is hyperactive in aged animals. A) Experimental timeline, including AAV injection, window implantation, and two-photon imaging. B) Diagram of chronic window implant over the right hemisphere of the DG. C) Diagram of imaging setup, with side view (left) and top view (right) of mouse head-fixed to treadmill with multiple tactile zones. D) Example field of view with regions of interest of active cells shaded in color. Scale bar = 100 μm. E) Mean single cell calcium activity. F) Example calcium traces (black) and corresponding treadmill positions (red) from young mice. G) Example calcium traces (black) and corresponding treadmill positions (red) from aged mice. a.u.= arbitrary units. Young N=8 mice, n=910 cells; Aged N=8 mice, n=699 cells. **p<0.01

selectivity of neuronal activity using a previously described spatial tuning index (see methods).

DG neurons in aged mice had significantly lower tuning indices than those in young mice (Fig.

2C, p<0.0001, nested bootstrap). This difference indicates that the activity of DG neurons in aged

mice was not as place-specific as that of young DG neurons, suggesting an aging-related

degradation of the spatial code. We also computed single-cell Fisher Information (FI), as a

measure of how much spatial information was encoded by individual DG neurons. By plotting

whole-recording calcium activity raster plots for neurons in the 10th percentile of FI we verified that

neurons with high spatial information behave like 'place cells' (*30*) in both young and aged

animals, as their activity is concentrated at a single location on the treadmill (Fig. 2D,E). This

place-specific response was present across laps as neurons in both cohorts of mice largely kept

the same place response in both even and odd laps. Aged mice encoded less spatial information

than young mice as denoted by a significant decrease in FI (Fig. 2F, p=0.00005, nested

bootstrap).

Figure 2. DG representations of space are impaired in aged mice. A) Tuning vector polar plots representing a highly spatially tuned cell and B) a cell with low spatial tuning. C) Spatial tuning index. D) Raster plots of tuning vectors of young mouse neurons in 10th percentile Fisher Information, sorted by the position of activity maximum for whole recording (All Laps). Raster plots of partial recording (Even Laps and Odd Laps) keeping same sorting used for whole recording. E) Same raster plots for aged mice. C) Single cell Fisher information. Statistics done with nested bootstrap analysis. Bars represent mean +/- SEM. Young N=8 mice, n=910 cells; Aged N=8 mice, n=699 cells. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. See also supplemental figures 1 and 2.

To confirm that the changes in spatial representations in aged mice corresponded to a change in

spatial memory, mice underwent behavioral testing using an object placement paradigm. This test

takes advantage of a mouse's natural preference for novelty to assess how well animals can

remember the location of objects in space, a behavior that is thought to be hippocampus-

dependent (*31*). We allowed mice to explore an arena with visual cues and two novel and identical

objects during a training trial. We then displaced one object to another location in the arena and

again allowed the mice to explore during a test trial (Fig. S3A). Young mice had a preference for

the novel object that was significantly above chance (p= 0.0017, One sample Wilcoxon test)

whereas aged mice did not score above chance (Fig S3B-C, p=0.6355, One sample Wilcoxon

test). Additionally, significantly more aged mice than young mice failed the test (Fig. S3D,

p=0.0393, Chi squared test). This change in preference can also be seen in the exploration times

per objects, in which young mice spend more time engaged with the displaced object, while aged

mice spend about an even amount of time between objects (Fig. S3E, p=0.0205 , Two-way

ANOVA with Sidak multiple comparisons test). These data confirm earlier findings of spatial

memory deficits in aged mice, which we now show is accompanied by impaired neural

representations of space in the DG.

Older individuals are better able to distinguish places in an environment that they are familiar with

than in a novel environment (*7*, *8*), so we asked how the representations of space in the DG

changed across four consecutive days of imaging, as the treadmill belt became more familiar to

the mice. Single-cell calcium activity remained elevated in aged mice through all recording days,

even though this difference was not significant in days 2 and 3 (Fig. 3A, nested bootstrap with

Bonferroni correction). While in young mice there was a slight but significant decrease in activity

(p=0.00005) across days, in aged mice activity increased over time (p<0.0001) (Fig. 3B-C, nested

bootstrap). In contrast to activity, the differences in tuning index and FI were erased over the four

days of imaging, as young and aged groups converged. The spatial tuning index of young mice

remained approximately constant over the course of all imaging sessions (p=0.19772), whereas

the tuning of aged mice underwent an increase so that the differences between both groups were

no longer significant in days 3 and 4 (p=0.00005). (Fig. 3D-F, nested bootstrap, Bonferroni

corrected in 3D). FI rose in both young and aged groups over the four imaging days, but the

increase was more pronounced in aged animals so that the differences between young and aged

animals were no longer significant in days 3 and 4 (Fig. 3G-I, nested bootstrap, Bonferroni

corrected in 3G). These data suggest that representations of space in aged animals can be

rescued by repeated exposure to the same environment.

Figure 3. Deficits in aged spatial representations are rescued with increased familiarity. A) Mean single cell calcium activity across days. B) Single cell activity in young mice on day 1 versus day 4. C) Single cell activity in aged mice on day 1 versus day 4. D) Mean tuning index across days. E) Tuning index in young mice on day 1 versus day 4. F) Tuning index in aged mice on day 1 versus day 4. G) Mean single cell Fisher Information across days. H) Fisher Information in young mice on day 1 versus day 4. I) Fisher Information in aged mice on day 1 versus day 4. Statistics done with a nested bootstrap analysis. Bars represent mean +/- SEM. a.u.= arbitrary units. Young: day 1 N=8 , n=910; day 2 N=7, n=694; day 3 N=7, n=623; day 4 N=7, n=610; Aged: day 1 N=8, n=699; day 2 N=8, n=619; day 3 N=5, n=147; day 4 N=6, n=331. ns=not significant, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Previous studies found the stability of neural representations of space was altered in aged animals

(*32, 33*). We therefore tracked the activity of individual neurons across days to investigate the

stability of the active cell ensemble (Fig. 4A-C). We first asked whether cells that were active in

the initial imaging session were reactivated on subsequent days. Surprisingly, we found no

difference between groups, suggesting that both DG neurons in young and aged mice have

similar reactivation rates (Fig. 4D, mixed effects model with Sidak multiple comparisons test). We

went on to compare the activity in the cells that were active both on day 1, when the treadmill was

novel, and day 4, when it had become familiar. The single-cell activity of matched cells increased

across days in the young mice (p= 0.00234, nested bootstrap) but not in the old mice (p=0.15981,

nested bootstrap) (Fig. 4E-F). This may reflect a ceiling for activity since the aged mice already

had higher activity on day 1. While the tuning index did not differ in either group across days (Fig.

4G-H, p=0.07626, p=0.21341), both groups saw higher spatial information, as FI was significantly

increased in reactivated cells on day 4 (Fig. 4I-J, p=0.00015, p=0.00024), which may simply

reflect the fact that FI increases sharply for the entire population across the four days. We then

asked whether the reactivated cells had a distinct profile compared to those neurons that were

not active across days. In day 1, cells that were to be reactivated did not differ from other cells in

single cell activity levels. Whether cells were reactivated or not, their activity levels were higher in

the aged mice on day 1, which recapitulates the pattern seen in the general population (Fig. 4K).

These effects were gone by day 4, again reflecting the trend in the general population (Fig. 4L).

In the aged group, cells that were reactivated showed higher tuning levels than other cells on day

1 (Fig. 4M), but this effect was also absent on day 4 (Fig. 4N). Reactivated cells had significantly

higher FI than other neurons in both the young and aged groups on both day 1 and day 4 (Fig.

4O-P). This suggests that higher spatial information is a predictor of whether neurons will be

reactivated over several days. Overall, we did not see evidence any changes in the stability of

coding ensembles in aged animals, when compared with the young cohort. However, since

reactivated neurons initially have higher tuning and spatial information than non-reactivated cells

in aged animals, and increase their FI over time, it's possible that they contribute to the

improvement of spatial representations in aged mice improve over the four days of imaging.

In summary, we have found that, in aged animals, DG neurons are hyperexcitable and impaired

in their encoding of spatial features in novel environments. Our results highlight how the defects

in spatial representations in the DG are specific to the first introduction to a novel context and can

be rescued by increasing familiarity with a given environment.

Figure 4. Reactivated cells do not have a distinct profile in aged animals. A) Example field of view used to match cells across days. Neurons that were active during recording session are shaded in color. B) Matched regions of interest from the field of view in panel A, red and white contours represent cells active on day 1 and 4. C) Examples of matched cells on day 1 and 4. D) Reactivation rate of neurons that were active on day 1. Percentages are based on imaging day versus day 1, regardless of whether the cell was active on other days. E-F) Matched single cell calcium activity in reactivated cells in young (E) and aged (F) mice. G-H) Matched tuning index in reactivated cells in young (G) and aged (H) mice. I-J) Matched single cell Fisher Information in reactivated cells in young (I) and aged (J) mice. K-L) Single cell activity of reactivated and non-reactivated cells on day 1 (K) and day 4 (L). M-N) Tuning of reactivated and non-reactivated cells on day 1 (M) and day 4 (N). O-P) Single cell Fisher Information of reactivated and non-reactivated cells on day 1 (O) and day 4 (P). Statistics done with nested bootstrap analysis. Bars represent mean +/- SEM. a.u.= arbitrary units. Young: N=6; day 1 n=617; day 1 react. n=140; day 4 n=396; day 4 react. n=140; Aged: N=5; day 1 n=375; day 1 react. n=80; day 4 n=196; day 4 react. n=80. ns=not significant, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Discussion

Aging is associated with cognitive decline in spatial memory in healthy aged humans and rodents (*2–4*). In this study we asked whether aged mice have impaired representations of space in the DG circuits that mediate spatial memory, which could account for the memory deficits seen in aged individuals.

Several studies have found that neuronal activity is elevated in the hippocampus of aged animals specifically in hippocampal area CA3 (*19–22*). Human studies using fMRI in older and younger adults have generally confirmed the hyperexcitability findings in the DG-CA3 axis (*5, 14*), although fMRI does not have the resolution to distinguish between areas DG and CA3. This aging-related hyperexcitability phenotype is thought to contribute to the memory deficits seen in aged individuals (*34, 35*). In order to understand the causes of the aging-related hyperexcitability it would be important to record neuronal activity from the DG, as this region provides the main inputs to CA3.

We used *in vivo* two-photon calcium imaging to record activity from young and aged DG neurons. This allowed us to image a large population of cells and therefore have a better chance capturing active cells in a sparsely active population. Additionally, using an imaging-based approach allowed us to study the same group of cells across time. We have found that single-cell calcium activity is indeed increased in aged DG neurons, when compared to young controls (Fig. 1E).

We also found that during the initial exposure to the treadmill, there is a significant reduction in spatial information and tuning in aged DG neurons, as compared to their young counterparts (Fig. 2B-C). By using untrained mice for our experiments we were able to capture the very first neuronal representations of a novel environment. We found that during the course of four consecutive imaging days, spatial information and tuning in the aged mice converged with those of young mice, which leads us to conclude that aged mice require additional exposure to a novel

environment in order to form adequate neural representations of space. It can therefore be said

that the spatial encoding defects seen in aged mice are specific to novel environments. This is in

agreement with previous reports that found that spatial representations in CA1 were disrupted

when novel environmental cues were introduced during a navigation task (*33*, *36*). Interestingly,

both young and aged groups saw an increase in spatial information through the four days of

imaging, whereas spatial tuning increased in aged mice but was not significantly different in young

mice. We did not find a convergence in single-cell calcium activity levels between aged and young

groups, meaning that the hyperexcitability is no impediment to aged mice eventually forming

accurate spatial representations. There are several interesting parallels between our findings and

human data showing that aged individuals are better able to navigate familiar environments than

novel ones (*8*) leading them to avoid taking novel routes in favor of navigating familiar routes.

To verify that aged mice have impairments in spatial memory, we used an object placement test

that is not physically tasking and does not involve an aversive component. Our behavior data

confirmed that aged mice are more likely to fail at an object placement task than their younger

counterparts. This goes along with studies analyzing similar spatial memory tasks that have found

deficits in aged animals (*37*, *38*). Our results also highlight the well-described heterogeneity

present during normal aging, as about half of aged mice were still able to pass the test (*39*, *40*).

In order to better compare our data to previous studies of DG activity using IEG labelling, we

determined the percentage of active cells in our imaging fields but found no difference in the

number of active cells between young and aged animals. This may be due to differences in the

activation of ensembles based on the experimental designs. However, there are also some

caveats to our experimental data that must be kept in mind: we calculate the total number of cells

from mean projections of whole calcium imaging movies, which can introduce several biases, for

example active cells will tend to be brighter and therefore more visible. We found a correlation

between the percentage of active cells and the total number of cells in the imaging field (Fig.

S2B), as fields of view with fewer detected cells had much higher variance of the fraction of active

cells. Fields with more total cells generally had more cells that were dimmer and less active but

were still counted in the total. In contrast, in fields with few cells due to areas of lower viral

expression or obscured by blood vessels, there are likely more cells present than can be

quantified in the mean projection image. These differences in imaging fields across animals

makes it difficult to calculate exact percentages, limiting the usefulness of this approach.

Previous studies have found that increased hippocampal excitability contributes to aging-related

cognitive impairments. Our experimental design did not allow us to determine whether

hyperexcitability contributes to the deficits in spatial information seen in the novel environment

during the first day of imaging, but our results suggest that instead dysfunction in the plasticity

mechanisms that underlie spatial selectivity may be to blame. Spatial selectivity can develop very

fast in CA1 hippocampal neurons through a mechanism termed behavioral timescale synaptic

plasticity (*41*) that is mostly active in novel contexts, leading to the development and stabilization

of spatial tuning within the first few minutes of exposure to a new environment (*42*). While less is

known about the dynamics of spatial selectivity in the DG, spatially tuned cells also seem to

emerge and stabilize within the first few minutes of exposure to a novel environment, with further

refinement as the animals are re-exposed to the same environment over the following days (*43*).

This is in line with our finding that spatial information increases over several days both in young

and aged animals. Previous literature has shown that aging is associated with plasticity deficits in

the hippocampus (*17*, *44*, *45*), namely in long term potentiation and depression. Given that these

plasticity mechanisms are likely most active as mice map out new environments (*42*) we speculate

that this may explain the protracted development of spatial selectivity that we see in aged mice.

Expression of some IEGs, such as *Arc* and *Egr1,* which are thought to be regulators of neuronal

plasticity, is reduced in the DG of aged animals. Deficits in the expression of these genes following

neuronal activity could be at the origin of plasticity defects in aged animals. The increased

excitability of aged DG neurons could therefore also be seen as a compensation mechanism for

both the reduced synaptic input from the entorhinal cortex, and the reduced expression of IEGs.

Another factor potentially contributing to the delayed formation of spatial representations in the

aged DG is a reduction of adult neurogenesis. Adult-born neurons are known to enhance DG

neuronal plasticity (*46*, *47*) but their numbers fall to almost zero in aged animals (*48*).

Aging-associated memory deficits have also been associated with changes in the stability of

spatial representations as some previous studies have found that the aged hippocampal spatial

code undergoes larger changes over different sessions than in young mice (*32*), whereas others

have found that spatial representations are more rigid (*24, 33*). We did not see changes in the

reactivation of cells from one session to another between groups (Fig. 4D). This is in contrast to

data showing that more distinct dentate populations express the IEG *Arc* when aged mice are re-

exposed to the same environment (*49*), which could be due to age-related changes in DNA

methylation rather than changes in neuronal excitability (*50*). In order to further investigate the

properties of cells in the aged DG over time, we matched cells that were active in our imaging

fields across days. We did not find major differences in the reactivated cells in any measures

between young and aged groups, suggesting that ensembles of neurons follow the same basic

mechanisms in the aged DG. When comparing reactivated neurons to the remainder of the

population, we found that reactivated cells had significantly higher spatial information (Fig. 4 O,P),

indicating that this may be a marker of whether a neuron will contribute to spatial representations

over time.

To summarize, our data shows that new DG spatial representations are initially impaired and take

longer to form in aged mice. These data expand our knowledge of the network activity and spatial

representations in the aged hippocampus and suggest that aging-associated hippocampal

hyperactivity is not an impediment to the formation of rich spatial representations. The protracted

refinement of spatial representations suggests that the plasticity mechanisms responsible for the

development of spatial selectivity are impaired in aged animals and are likely to be a relevant

therapeutic target for ameliorating the memory deficits associated with aging.

Acknowledgements

We thank Dr. Carolyn Pytte for the generous gift of aged mice from her laboratory at Queens

College. We thank Dr. Maria Gulinello, the director of the behavior core facilities at Einstein, for

her training and technical advice on behavior experiments. We thank Dr. Alberto Cruz-Martín for

helpful feedback and discussions.

Funding:

The Einstein Training Program in Stem Cell Research from the Empire State Stem Cell Fund

through New York State Department of Health Contract C34874GG (KDM, MAF), Whitehall

Foundation Research Grant 2019-05-71 (JTG), National Institutes of Health NINDS

R01NS125252-01A1 (JTG)

Author contributions:

Conceptualization: KDM, JTG; Methodology: KDM, MAF, JTJ, JTG; Investigation: KDM, SSM,

JTG; Visualization: KDM, JTG; Supervision: JTG; Writing—original draft: KDM, JTG; Writing—

review & editing: KDM, JTG

Competing interests:

The authors have no competing interests.

 Data availability:

The   analysis   code   used   in   this   study   is   openly   available   at

https://github.com/GoncalvesLab/McDermott-et-al-aging. All imaging data, consisting of calcium

traces for every active cell and the position of the mouse on the treadmill, will be made openly

available on a repository site before final publication. Due to their large size, raw calcium imaging

movies will be made available upon request.

Methods

Animals – Aged mice were 21-26 months old C57Bl6J females originally from Jackson Labs and

kindly gifted by Dr. Carolyn Pytte of Queens College. Young mice were 3-4 months old C57Bl6J

females purchased from Jackson Labs. All mice were housed in standard conditions with a 14/10

hour light/dark cycle. Mice were provided food and water ad libitum. All procedures were done

during the light part of the cycle and in accordance with the Einstein Institutional Animal Care and

Use Committee (Protocol #00001197).

Viral injections – Mice were anesthetized (induction: 5%, maintenance 0.5% isoflurane in O_2

vol/vol). Following anesthetization, mice were attached to a stereotactic apparatus and the right

hemisphere of the dentate gyrus (DG) was injected with 1µl of a DJ-serotype AAV vector

encoding the red-shifted calcium indicator jRGECO1a (*25*) under the CamKII promoter at 10^{12}

GC/ml titer. Viral injections were done with a pulled glass pipette using a Nanoject III microinjector

(Drummond) at previously described coordinates (*51*).

Window implantation – Following viral injection, dexamethasone (1 mg/kg) was administered

via subcutaneous injection to minimize brain swelling. Optibond (Kerr Dental) adhesive was

applied to the skull and cured to help secure the implant once attached. A 3mm diameter

craniotomy was made over the right dorsal DG and the overlaying tissue was removed by

aspiration down to the hippocampal fissure where a custom-built cylindrical titanium implant with

a glass coverslip on the bottom was inserted over the dorsal surface of the DG (*27*). The implant

and a titanium bar for head-fixation were attached to the skull with dental cement. All mice were

given carprofen (5 mg/kg, subcutaneous) as a post-surgery analgesic.

Two-photon Imaging – *In vivo* calcium imaging movies were acquired with a custom two-photon

laser scanning microscope (based on Thorlabs Bergamo) using a femtosecond-pulsed laser

(Coherent Fidelity 2, 1070 nm) and a 16x water immersion objective (0.8 NA, Nikon). Imaging

sessions were started 3-4 weeks after surgery to allow for recovery and for optimal viral

expression. During imaging sessions, mice were head-fixed to the microscope with a titanium

headbar and the microscope stage was adjusted so that the hippocampal window was

perpendicular to the axis of the objective for optimal imaging conditions. Mice were awake and

walking on a manually rotated treadmill belt throughout imaging. The treadmill contained four

zones with different textures as previously described (*52*). Ten-minute videos were acquired at

15.253 fps with a 343.6 x 343.6 µm field of view. Treadmill data was acquired using National

Instruments analog-to-digital converter and synchronized with the imaging data using Thorsync

software. To track the same field of view, the initial location was noted on the first imaging day

using a coordinate system and by taking an image of the field. During following sessions, the

coordinates and initial field image were used to match as close as possible to the same field.

Data Analysis– The open-source Suite2p software (version 10.0) was used to register and

motion correct videos, for cell detection and spike deconvolution (*53–55*). Regions of interest

corresponding to active neurons were identified by Suite2p and further manually curated in order

to identify all active cells in the field of view. The ROIMatchPub (*56*) was used to identify and

match cells in the same field of view across imaging sessions. Matched ROIs were manually

confirmed after automated detection.

Single-cell calcium activity was determined as previously described (*29*). Briefly, ΔF/F data was

filtered with a zero phase shift, third order Butterworth lowpass filter. This filtered ΔF/F was

thresholded to 2 standard deviations(σ) above a rolling-mean baseline. Single cell activity was

determined as the cumulative sum of the thresholded trace for each cell. This was then normalized

to the total distance traveled by each mouse.

Tuning indices were using deconvolved calcium traces, which were thresholded to 2σ above the

baseline. Any point not significantly above that noise threshold was set to zero. Videos were cut

to only include periods of movement >1s in duration. The treadmill band was segmented into 100

position bins, and transients were mapped to these bins according to the location of the mice on

the treadmill in order to generate a tuning vector for each cell (*26*).The mean of the thresholded

activity at each location was calculated for every neuron and normalized to the time the mouse

spent at that position. The tuning index was defined as the modulus of this normalized tuning

vector.

Single-cell Fisher Information (FI) was calculated as previously described (*29, 57*). A bias-

corrected signal to noise ratio was computed using the unfiltered ΔF/F fluorescence data for each

individual cell, where the signal is the square of the difference of the mean activity at two locations

on the treadmill, and the noise is the average variance of the activity at each location. Position

data was segmented into 20 bins.

Behavior – Object Placement –A square-shaped arena 42x42x30 (LxWxH) was set up with visual

cues along the walls and with two novel and identical objects on the floor. Mice were individually

placed in the arena for a 4 minute training trial. Following a 45 minute retention period, one object

was displaced to another location in the arena and the mice were placed again for a 4 minute test

trial. Exploration of the objects was quantified by an experimenter and confirmed with computer

tracking.

Statistics – Nested bootstrap analysis – When pooling cell data from different mice into a single experimental group, significance testing was done using a muti-level bootstrapped approach as previously described (*29*). To assess whether differences between two experimental groups are significant, a null surrogate distribution was constructed for each mouse in each group by resampling with substitution from the pool of all imaged cells. The difference between the null distributions generated for each group was calculated and the previous steps repeated to generate 100,000 bootstrap estimates of the difference between two groups' null distributions. The empirically observed value of the difference between conditions was then compared to the null distribution values with a statistical significance level (α) set at 0.05. If the empirical group differences fall outside of the 95[th] percentile of the 100,000 bootstrap estimates of the difference between the null distributions, then it is considered to be a statistically significant difference. The p-value is the proportion of bootstrap different estimates that are larger than the empirical difference between groups (or smaller, if the difference is negative). Whenever comparing more than two groups, Bonferroni's correction for multiple comparisons was applied.

- Mixed-effects models with Geisser-Greenhouse correction for matched values were used to compare mean values by mouse across imaging days. Sidak's multiple comparisons test was used to compare groups on each imaging day.

- Nonparametric one sample Wilcoxon tests were used to determine if group medians were significantly above chance levels in behavior experiments.

- A two-sided Chi squared test with 95% confidence interval was used to compare pass/fail rates in behavior experiments.

References

1. W. Jagust, Vulnerable neural systems and the borderland of brain aging and neurodegeneration. *Neuron*. **77**, 219–234 (2013).
2. C. A. Barnes, Aging and the physiology of spatial memory. *Neurobiol Aging*. **9**, 563–568 (1988).
3. C. Techentin, D. Voyer, S. D. Voyer, Spatial abilities and aging: a meta-analysis. *Exp Aging Res*. **40**, 395–425 (2014).
4. K. Konishi, S. Mckenzie, N. Etchamendy, S. Roy, V. D. Bohbot, Hippocampus-dependent spatial learning is associated with higher global cognition among healthy older adults. *Neuropsychologia*. **106**, 310–321 (2017).
5. N. Diersch, J. P. Valdes-Herrera, C. Tempelmann, T. Wolbers, Increased Hippocampal Excitability and Altered Learning Dynamics Mediate Cognitive Mapping Deficits in Human Aging. *J. Neurosci*. **41**, 3204–3221 (2021).
6. S. D. Moffat, Aging and Spatial Navigation: What Do We Know and Where Do We Go? *Neuropsychol Rev*. **19**, 478–489 (2009).
7. K. C. Kirasic, Spatial cognition and behavior in young and elderly adults: implications for learning new environments. *Psychol Aging*. **6**, 10–18 (1991).
8. V. Muffato, M. Della Giustina, C. Meneghetti, R. De Beni, Age-related differences in pointing accuracy in familiar and unfamiliar environments. *Cogn Process*. **16 Suppl 1**, 313–317 (2015).
9. N. A. Merriman, J. Ondřej, E. Roudaia, C. O'Sullivan, F. N. Newell, Familiar environments enhance object and spatial memory in both younger and older adults. *Exp Brain Res*. **234**, 1555–1574 (2016).
10. W. B. Scoville, B. Milner, Loss of recent memory after bilateral hippocampal lesions. *J. Neurol. Neurosurg. Psychiatr*. **20**, 11–21 (1957).
11. R. G. M. Morris, P. Garrud, J. N. P. Rawlins, J. O'Keefe, Place navigation impaired in rats with hippocampal lesions. *Nature*. **297**, 681–683 (1982).
12. S. A. Small, M. K. Chawla, M. Buonocore, P. R. Rapp, C. A. Barnes, Imaging correlates of brain function in monkeys and rats isolates a hippocampal subregion differentially vulnerable to aging. *Proc Natl Acad Sci U S A*. **101**, 7181–7186 (2004).
13. G. W. Small, What we need to know about age related memory loss. *BMJ*. **324**, 1502–1505 (2002).
14. M. A. Yassa, A. T. Mattfeld, S. M. Stark, C. E. L. Stark, Age-related memory deficits linked to circuit-specific disruptions in the hippocampus. *PNAS*. **108**, 8873–8878 (2011).
15. H. Moreno, W. E. Wu, T. Lee, A. Brickman, R. Mayeux, T. R. Brown, S. A. Small, Imaging the Abeta-related neurotoxicity of Alzheimer disease. *Arch Neurol*. **64**, 1467–1477 (2007).
16. S. N. Burke, C. A. Barnes, Senescent synapses and hippocampal circuit dynamics. *Trends in Neurosciences*. **33**, 153–161 (2010).
17. D. J. Froc, B. Eadie, A. M. Li, K. Wodtke, M. Tse, B. R. Christie, Reduced Synaptic Plasticity in the Lateral Perforant Path Input to the Dentate Gyrus of Aged C57BL/6 Mice. *Journal of Neurophysiology*. **90**, 32–38 (2003).
18. D. GoodSmith, X. Chen, C. Wang, S. H. Kim, H. Song, A. Burgalossi, K. M. Christian, J. J. Knierim, Spatial Representations of Granule Cells and Mossy Cells of the Dentate Gyrus. *Neuron*. **93**, 677-690.e5 (2017).
19. I. A. Wilson, S. Ikonen, M. Gallagher, H. Eichenbaum, H. Tanila, Age-associated alterations of hippocampal place cells are subregion specific. *J Neurosci*. **25**, 6877–6886 (2005).
20. Y. H. El-Hayek, C. Wu, H. Ye, J. Wang, P. L. Carlen, L. Zhang, Hippocampal excitability is increased in aged mice. *Experimental Neurology*. **247**, 710–719 (2013).
21. D. Simkin, S. Hattori, N. Ybarra, T. F. Musial, E. W. Buss, H. Richter, M. M. Oh, D. A. Nicholson, J. F. Disterhoft, Aging-Related Hyperexcitability in CA3 Pyramidal Neurons Is

Mediated by Enhanced A-Type K+ Channel Function and Expression. *J Neurosci*. **35**,
13206–13218 (2015).

22. H. Lee, Z. Wang, S. L. Zeger, M. Gallagher, J. J. Knierim, Heterogeneity of Age-Related
Neural Hyperactivity along the CA3 Transverse Axis. *J. Neurosci*. **41**, 663–673 (2021).

23. C. A. Barnes, B. L. McNaughton, J. O'Keefe, Loss of place specificity in hippocampal
complex spike cells of senescent rat. *Neurobiology of Aging*. **4**, 113–119 (1983).

24. H. Tanila, M. Shapiro, M. Gallagher, H. Eichenbaum, Brain Aging: Changes in the Nature of
Information Coding by the Hippocampus. *J. Neurosci*. **17**, 5155–5166 (1997).

25. H. Dana, B. Mohar, Y. Sun, S. Narayan, A. Gordus, J. P. Hasseman, G. Tsegaye, G. T.
Holt, A. Hu, D. Walpita, R. Patel, J. J. Macklin, C. I. Bargmann, M. B. Ahrens, E. R.
Schreiter, V. Jayaraman, L. L. Looger, K. Svoboda, D. S. Kim, Sensitive red protein calcium
indicators for imaging neural activity. *eLife*. **5**, e12727 (2016).

26. N. B. Danielson, P. Kaifosh, J. D. Zaremba, M. Lovett-Barron, J. Tsai, C. A. Denny, E. M.
Balough, A. R. Goldberg, L. J. Drew, R. Hen, A. Losonczy, M. A. Kheirbek, Distinct
Contribution of Adult-Born Hippocampal Granule Cells to Context Encoding. *Neuron*. **90**,
101–112 (2016).

27. J. T. Gonçalves, C. W. Bloyd, M. Shtrahman, S. T. Johnston, S. T. Schafer, S. L. Parylak, T.
Tran, T. Chang, F. H. Gage, In vivo imaging of dendritic pruning in dentate granule cells.
Nat Neurosci. **19**, 788–791 (2016).

28. M. A. Yassa, S. M. Stark, A. Bakker, M. S. Albert, M. Gallagher, C. E. L. Stark, High-
resolution structural and functional MRI of hippocampal CA3 and dentate gyrus in patients
with amnestic Mild Cognitive Impairment. *NeuroImage*. **51**, 1242–1252 (2010).

29. M. A. Frechou, S. S. Martin, K. D. McDermott, Ş. Gökhan, W. A. Tomé, R. Coen-Cagli, J. T.
Gonçalves, Adult neurogenesis improves spatial information encoding in the mouse
hippocampus (2022), p. 2022.11.30.518622, , doi:10.1101/2022.11.30.518622.

30. J. O'Keefe, J. Dostrovsky, The hippocampus as a spatial map. Preliminary evidence from
unit activity in the freely-moving rat. *Brain Research*. **34**, 171–175 (1971).

31. J. Haettig, Y. Sun, M. A. Wood, X. Xu, Cell-type specific inactivation of hippocampal CA1
disrupts location-dependent object recognition in the mouse. *Learn Mem*. **20**, 139–146
(2013).

32. C. A. Barnes, M. S. Suster, J. Shen, B. L. McNaughton, Multistability of cognitive maps in
the hippocampus of old rats. *Nature*. **388**, 272–275 (1997).

33. I. A. Wilson, S. Ikonen, I. Gureviciene, R. W. McMahan, M. Gallagher, H. Eichenbaum, H.
Tanila, Cognitive aging and the hippocampus: how old rats represent new environments. *J
Neurosci*. **24**, 3870–3878 (2004).

34. J. Jiménez-Balado, T. S. Eich, GABAergic dysfunction, neural network hyperactivity and
memory impairments in human aging and Alzheimer's disease. *Seminars in Cell &
Developmental Biology*. **116**, 146–159 (2021).

35. R. P. Haberman, A. Branch, M. Gallagher, Targeting Neural Hyperactivity as a Treatment to
Stem Progression of Late-Onset Alzheimer's Disease. *Neurotherapeutics*. **14**, 662–676
(2017).

36. H. Tanila, P. Sipilä, M. Shapiro, H. Eichenbaum, Brain Aging: Impaired Coding of Novel
Environmental Cues. *J Neurosci*. **17**, 5167–5174 (1997).

37. M. E. Wimmer, P. J. Hernandez, J. Blackwell, T. Abel, Aging impairs hippocampus-
dependent long-term memory for object location in mice. *Neurobiol Aging*. **33**, 2220–2224
(2012).

38. S. Yanai, T. Tago, J. Toyohara, T. Arasaki, S. Endo, Reversal of spatial memory impairment
by phosphodiesterase 3 inhibitor cilostazol is associated with reduced neuroinflammation
and increased cerebral glucose uptake in aged male mice. *Frontiers in Pharmacology*. **13**
(2022) (available at https://www.frontiersin.org/articles/10.3389/fphar.2022.1031637).

39. A. M. Hamieh, E. Camperos, A. M. Hernier, V. Castagné, C57BL/6 mice as a preclinical
model to study age-related cognitive deficits: Executive functions impairment and inter-
individual differences. *Brain Res.* **1751**, 147173 (2021).
40. A. J. Gower, Y. Lamberty, The aged mouse as a model of cognitive decline with special
emphasis on studies in NMRI mice. *Behav Brain Res.* **57**, 163–173 (1993).
41. K. C. Bittner, A. D. Milstein, C. Grienberger, S. Romani, J. C. Magee, Behavioral time scale
synaptic plasticity underlies CA1 place fields. *Science.* **357**, 1033–1036 (2017).
42. J. B. Priestley, J. C. Bowler, S. V. Rolotti, S. Fusi, A. Losonczy, Signatures of rapid plasticity
in hippocampal CA1 representations during novel experiences. *Neuron.* **110**, 1978-1992.e6
(2022).
43. S. Kim, D. Jung, S. Royer, Place cell maps slowly develop via competitive learning and
conjunctive coding in the dentate gyrus. *Nat Commun.* **11**, 4550 (2020).
44. S. N. Burke, C. A. Barnes, Neural plasticity in the ageing brain. *Nature Reviews*
Neuroscience. **7**, 30–40 (2006).
45. C. M. Norris, D. L. Korol, T. C. Foster, Increased susceptibility to induction of long-term
depression and long-term potentiation reversal during aging. *J Neurosci.* **16**, 5382–5392
(1996).
46. J. S. Snyder, N. Kee, J. M. Wojtowicz, Effects of Adult Neurogenesis on Synaptic Plasticity
in the Rat Dentate Gyrus. *Journal of Neurophysiology.* **85**, 2423–2431 (2001).
47. F. Massa, M. Koehl, T. Wiesner, N. Grosjean, J.-M. Revest, P.-V. Piazza, D. N. Abrous, S.
H. R. Oliet, Conditional reduction of adult neurogenesis impairs bidirectional hippocampal
synaptic plasticity. *Proceedings of the National Academy of Sciences.* **108**, 6644–6649
(2011).
48. H. G. Kuhn, H. Dickinson-Anson, F. H. Gage, Neurogenesis in the dentate gyrus of the adult
rat: age-related decrease of neuronal progenitor proliferation. *J. Neurosci.* **16**, 2027–2033
(1996).
49. D. F. Marrone, E. Satvat, M. J. Shaner, P. F. Worley, C. A. Barnes, Attenuated long-term
Arc expression in the aged fascia dentata. *Neurobiology of Aging.* **33**, 979–990 (2012).
50. M. R. Penner, T. L. Roth, M. K. Chawla, L. T. Hoang, E. D. Roth, F. D. Lubin, J. D. Sweatt,
P. F. Worley, C. A. Barnes, Age-related changes in Arc transcription and DNA methylation
within the hippocampus. *Neurobiology of Aging.* **32**, 2198–2210 (2011).
51. C. Zhao, E. M. Teng, R. G. Summers, G. Ming, F. H. Gage, Distinct Morphological Stages of
Dentate Granule Neuron Maturation in the Adult Mouse Hippocampus. *J. Neurosci.* **26**, 3–
11 (2006).
52. J. T. Jordan, K. D. McDermott, M. A. Frechou, M. Shtrahman, J. T. Gonçalves, Treadmill-
based task for assessing spatial memory in head-fixed mice. *STAR Protoc.* **2**, 100770
(2021).
53. M. Pachitariu, C. Stringer, M. Dipoppa, S. Schröder, L. F. Rossi, H. Dalgleish, M. Carandini,
K. D. Harris, Suite2p: beyond 10,000 neurons with standard two-photon microscopy.
bioRxiv, 061507 (2017).
54. M. Pachitariu, C. Stringer, K. D. Harris, Robustness of Spike Deconvolution for Neuronal
Calcium Imaging. *J. Neurosci.* **38**, 7976–7985 (2018).
55. J. Friedrich, P. Zhou, L. Paninski, Fast online deconvolution of calcium imaging data. *PLoS*
Comput Biol. **13**, e1005423 (2017).
56. ransona, ransona/ROIMatchPub (2023), (available at
https://github.com/ransona/ROIMatchPub).
57. I. Kanitscheider, R. Coen-Cagli, A. Kohn, A. Pouget, Measuring Fisher information
accurately in correlated neural populations. *PLoS Comput Biol.* **11**, e1004218 (2015).

533
534

Figure legends:

Figure 1. DG is hyperactive in aged animals. A) Experimental timeline, including AAV injection, window implantation, and two-photon imaging. B) Diagram of chronic window implant over the right hemisphere of the DG. C) Diagram of imaging setup, with side view (left) and top view (right) of mouse head-fixed to treadmill with multiple tactile zones. D) Example field of view with regions of interest of active cells shaded in color. Scale bar = 100 μm. E) Mean single cell calcium activity. F) Example calcium traces (black) and corresponding treadmill positions (red) from young mice. G) Example calcium traces (black) and corresponding treadmill positions (red) from aged mice. a.u.= arbitrary units. Young N=8 mice, n=910 cells; Aged N=8 mice, n=699 cells. **p<0.01

Figure 2. DG representations of space are impaired in aged mice. A) Tuning vector polar plots representing a highly spatially tuned cell and B) a cell with low spatial tuning. C) Spatial tuning index. D) Raster plots of tuning vectors of young mouse neurons in 10th percentile Fisher Information, sorted by the position of activity maximum for whole recording (All Laps). Raster plots of partial recording (Even Laps and Odd Laps) keep same sorting used for whole recording. E) Same raster plots for aged mice. C) Single cell Fisher information. Statistics done with nested bootstrap analysis. Bars represent mean +/- SEM. a.u.= arbitrary units. Young N=8 mice, n=910 cells; Aged N=8 mice, n=699 cells. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. See also supplemental figures 1 and 2.

Figure 3. Deficits in aged spatial representations are rescued with increased familiarity. A) Mean single cell calcium activity across days. B) Single cell activity in young mice on day 1 versus day 4. C) Single cell activity in aged mice on day 1 versus day 4. D) Mean tuning index across days. E) Tuning index in young mice on day 1 versus day 4. F) Tuning index in aged mice on day 1 versus day 4. G) Mean single cell Fisher Information across days. H) Fisher Information in young mice on day 1 versus day 4. I) Fisher Information in aged mice on day 1 versus day 4. Statistics done with a nested bootstrap analysis. Bars represent mean +/- SEM. a.u.= arbitrary units. Young:

day 1 N=8 , n=910; day 2 N=7, n=694; day 3 N=7, n=623; day 4 N=7, n=610; Aged: day 1 N=8, n=699; day 2 N=8, n=619; day 3 N=5, n=147; day 4 N=6, n=331. ns=not significant, $*p<0.05$, $**p<0.01$, $***p<0.001$, $****p<0.0001$.

Figure 4. Reactivated cells do not have a distinct profile in aged animals. A) Example field of view used to match cells across days. Neurons that were active during recording session are shaded in color. B) Matched regions of interest from the field of view in panel A, red and white contours represent cells active on day 1 and 4. C) Examples of matched cells on day 1 and 4. D) Reactivation rate of neurons that were active on day 1. Percentages are based on imaging day versus day 1, regardless of whether the cell was active on other days. E-F) Matched single cell calcium activity in reactivated cells in young (E) and aged (F) mice. G-H) Matched tuning index in reactivated cells in young (G) and aged (H) mice. I-J) Matched single cell Fisher Information in reactivated cells in young (I) and aged (J) mice. K-L) Single cell activity of reactivated and non-reactivated cells on day 1 (K) and day 4 (L). M-N) Tuning of reactivated and non-reactivated cells on day 1 (M) and day 4 (N). O-P) Single cell Fisher Information of reactivated and non-reactivated cells on day 1 (O) and day 4 (P). Statistics done with nested bootstrap analysis. Bars represent mean +/- SEM. a.u.= arbitrary units. Young: N=6; day 1 n=617; day 1 react. n=140; day 4 n=396; day 4 react. n=140; Aged: N=5; day 1 n=375; day 1 react. n=80; day 4 n=196; day 4 react. n=80. ns=not significant, $*p<0.05$, $**p<0.01$, $***p<0.001$, $****p<0.0001$.

Figure S1. Velocity and laps run during two photon imaging. A) Average velocity of mice walking along treadmill belt. B) Average number of laps run per imaging session. Statistics done with mixed effects model. Each point represent an individual animal. Bars represent mean +/- SEM. ns=not significant.

Figure S2. No difference in active cell number between young and aged mice. A) Percentage of cells active out of the total cells in a field of view, by mouse. Statistics done with mixed effects model. Bars represent mean +/- SEM. ns=not significant. B) Correlation of percentage active cells versus the total number of cells pooled from all imaging days. Data are shown by mouse.

Figure S3. Aged mice show a deficit in a spatial memory task. A) Schematic of the object placement behavioral paradigm. B) Novelty preference of mice measured by percentage of total time spent exploring the novel object. C) Discrimination index calculated by dividing the difference in exploration time between objects by the total exploration time [(novel-familiar)/(novel+familiar)]. D) Pass/fail rate of task, where passing was defined as at least a 53% preference for novel object. E) Exploration time dedicated to each object. Statistics in B and C were done with one sample Wilcoxon tests, D with Chi squared test and E with mixed effects model. Bars represent mean +/- SEM. Young N=13, Aged N=13. ns=not significant, *p<0.05, **p<0.01.

Figure 1

Figure 2

Figure 3

Figure 4