Combining experiments and simulations to examine the temperature-dependent behaviour of a disordered protein

Francesco Pesce and Kresten Lindorff-Larsen*

Structural Biology and NMR Laboratory, The Linderstrøm-Lang Centre for Protein Science, Department of Biology, University of Copenhagen, Copenhagen, Denmark

E-mail: lindorff@bio.ku.dk

Abstract

Intrinsically disordered proteins are a class of proteins that lack stable folded conformations and instead adopt a range of conformations that determine their biochemical functions. The temperature-dependent behaviour of such disordered proteins is complex and can vary depending on the specific protein and environment. Here, we have used molecular dynamics simulations and previously published experimental data to investigate the temperature-dependent behaviour of Histatin 5, a 24-residue-long polypeptide. We examined the hypothesis that Histatin 5 undergoes a loss of polyproline II structure with increasing temperature, leading to more compact conformations. We found that the conformational ensembles generated by the simulations generally agree with small-angle X-ray scattering data for Histatin 5, but show some discrepancies with the hydrodynamic radius as probed by pulsed-field gradient nuclear magnetic resonance spectroscopy, and with the secondary structure information derived from circular dichroism. We attempted to reconcile these differences by reweighting the conformational ensembles against the scattering and NMR data. By doing so, we were

in part able to capture the temperature-dependent behaviour of Histatin 5 and to link the observed decrease in hydrodynamic radius with increasing temperature to a loss of polyproline II structure. We were, however, unable to achieve agreement with both the scattering and NMR data within experimental errors. We discuss different possibilities for this outcome including inaccuracies in the force field, differences in conditions of the NMR and scattering experiments, and issues related to the calculation of the hydrodynamic radius from conformational ensembles. Our study highlights the importance of integrating multiple types of experimental data when modelling conformational ensembles of disordered proteins and how environmental factors such as the temperature influence them.

Introduction

Intrinsically disordered proteins (IDPs) are a class of proteins characterized by the lack of stable folded conformations. Instead, IDPs may adopt a wide range of conformations, which help determine their biochemical functions.[1,2] This ensemble of conformations, although disordered, may show sequence-dependent populations of specific regions of the Ramachandran map,[3] and the formation of transient secondary structures is common.[4,5] Therefore, they are better described as statistical coils.[6] Biophysical measurements make it possible to get insights into the propensity of IDPs to form (transient) secondary structures, although the interpretation of ensemble-averaged data is often ambiguous.[7] Alternatively, molecular dynamics (MD) simulations enable direct access to the population of conformations of IDPs at atomistic resolution, but its utility may be limited by difficulties in sampling and by the accuracy of the force fields used.[8–12]

The complexity of the problem increases even more when considering that the behaviour of IDPs is often highly dependent on the environment.[13] One factor that may affect the behaviour of IDPs is the temperature.[14–19] The temperature-dependence of the conformational properties of IDPs may, however, be complex, as they depend on the combined effects

of many different physical effects, and may vary depending on the specific protein and the specific environment in which it is found.[20,21] Other than influencing the single chain behaviour, temperature is also known to influence how IDPs interact and their tendency to form high-order entities.[22,23] Therefore, understanding how temperature affects the single-chain behaviour, may provide new insights into how aggregates and biomolecular condensates can form as a consequence of particular single-chain behaviours.[24]

Experimentally, the temperature-dependent behaviour of the local structure of IDPs has previously been studied using for example circular dichroism (CD) spectroscopy,[4,5,25–27] where changes in the spectra, although difficult to interpret at atomistic resolution, have been suggested to be due to formation of α-helices and/or loss of polyproline II (PPII) structures with increasing temperature.[15,19,28,29] This change in propensity to form certain secondary structure elements has also been linked to changes in the overall dimension of IDPs.[5,14,15,19,30] For example, it has been suggested that the loss of PPII with increasing temperature is responsible for the conformational ensemble of certain IDPs to get more compact.[15,19] Even though this might sound counter intuitive, it could be explained by the fact that the PPII structure is a particularly extended secondary structure element, though other mechanisms are also likely to be important.[14,20,21]

We here examine the case study of Histatin 5 (hereafter Hst5). This 24 residue-long polypeptide has previously been characterized by a multitude of experiments.[19,31–36] CD spectra recorded at different temperatures between 283 K and 323 K suggested that a loss of PPII structure takes place with increasing temperature. Increasing the temperature also results in a decrease of the hydrodynamic radius (R_h) measured by pulsed-field gradient (PFG) NMR, while the radius of gyration (R_g) extracted from small-angle X-ray scattering (SAXS) data appears not to be particularly sensitive to temperature. Previous attempts of computational characterization of the conformational ensembles of Hst5 at different temperatures did not completely succeed in explaining the experimental results,[19] but it was suggested that the drop in R_h with increasing temperature might be due to the temperature-dependent

change in the propensity of forming PPII structures.

To provide a structural explanation of the experimental data, we extensively sample the conformational ensemble of Hst5 at different temperatures by running MD simulations enhanced with metadynamics.[37] We then compare the conformational ensembles to the available temperature-dependent experimental data, but do not observe substantial temperature-dependent changes in compaction and propensity to form PPII structures. We refine the simulations against SAXS and R_h from PFG NMR using the Bayesian/Maximum-entropy reweighting approach,[38–40] and find that when we optimize the agreement with the R_h we also observe a temperature-dependent change in the amount of PPII structure in qualitative agreement with the CD data. On the other hand, when we combine the simulations with the SAXS data we do not observe any temperature-dependent change in compaction and PPII structure content. When we refine the ensembles simultaneously against the SAXS and NMR data, we could not achieve the same level of agreement as when optimizing against one of them at the time. We discuss this finding in light of earlier results[41] to shed light on this observation.

Results and Discussion

We sampled the conformational ensembles of Hst5 at 283 K, 293 K, 310 K and 323 K with MD simulations using the all-atom Amber ff99SB-disp (a99SB-disp) force field,[11] as it has previously been shown to be reliable in capturing the overall chain dimension of IDPs[11] and in reproducing NMR residual dipolar couplings of a peptide with high a PPII structure propensity.[42] In order to improve the sampling of the overall dimensions and secondary structure preferences of Hst5, we used Parallel Bias Metadynamics[43] (PBMetaD; see Methods) simulations with two semi-independent replicas biasing the same free-energy surface (FES).[44] We ran these simulations until the sampling of the collective variables (CVs) that we biased with PBMetaD converged within a reasonable error (*i.e.* an average error on the

estimated free-energy surface (FES) of the R_g lower than 1 kJ/mol; Fig. S1). We achieved this in all cases in a simulation time comprised between 2 and 2.5 μs per replica. We also verify that, along the simulations, both R_g and the backbone dihedrals diffuse in the CV space and do not get stuck in a specific state, supporting that the extent of biasing used in the simulations is sufficient for the system to escape local minima (Fig. S2 and S3). Furthermore, because of the well-tempered biasing[45] (see Methods), it would be ideal for the amount of bias dispensed in the system, which decreases over time, to reach a quasi-static regime, to ensure a proper convergence of the simulation. This happens when all accessible regions of the FES have been explored. In our simulations, we observe that overall the bias deposition become quasi-static, with the exception of few frames, whose R_g is outside the interval in which the bias is active (Fig. S4). Therefore this few frames are not taken into account when calculating averages and errors (see Methods).

We then calculated SAXS intensities, R_g, R_h, and per-residue secondary structure propensities, and compared these to the experimental values. We find that all simulations—apart from that at 310 K—provide a χ_r^2 between the calculated and experimental SAXS data close to 1 (Fig. 1a), which, given the use of normalized error bars for SAXS intensities (see Methods), shows that the ensembles accurately fit the SAXS experiments. As previously reported,[19] we find that the R_g values derived from the experimental SAXS profile (with the Guinier approximation, using the ATSAS package[46]) show the same average dimension for Hst5 in solution at the different temperatures (Fig. 1b). Similarly, the conformational ensembles sampled by MD, whose average R_g values are all comparable, but consistently smaller than those from the Guinier approximation. This observation is in line with the expectation that SAXS-derived R_g carries the contribution of the protein hydration layer,[47] while from the conformational ensembles the R_g is calculated on the coordinates of the protein atoms only.[48] For this reason we only rely on the χ_r^2 to SAXS data to determine whether a certain ensemble is accurate in terms of chain dimensions as described by SAXS data, since in calculating the SAXS we take into account both the protein hydration layer

and the excluded volume effect (see Methods).

In contrast, the calculated R_h does not agree well with the experimental values determined by PFG NMR at any of the temperatures (Fig. 1c). Instead, we find that the calculated R_h values are consistently smaller than the experimental values. We have previously shown that it is possible to build conformational ensembles for Hst5 that are simultaneously in good agreement with both SAXS and experimental R_h values.[41] Therefore, it is possible that the force field used is good at reproducing the protein dimension encoded in SAXS data, but not the R_h. For the ensembles at 283 K and 293 K we observe a good fit to SAXS data but a less good agreement with the experimental R_h values, which might be due to the fact that SAXS and PFG NMR probe different aspects of protein compaction.[49] Also, we note that the calculation of the R_h from conformational ensembles of IDPs might be affected by inaccuracies of forward models. A recent study has highlighted how the R_h calculated with the Kirkwood-Riseman equation is generally in agreement with experiments.[41] Nonetheless, this only works under the assumption that the R_h of the reference compound (most commonly dioxane[50]) used to derive the R_h from the diffusion data obtained by NMR is correctly estimated, and further rests on the assumption of treating R_h as an ensemble- and time-averaged quantity.[51]

The CD spectra of Hst5 show profiles often taken to represent PPII structure. The decrease in intensity of the PPII signature with increasing temperature suggests that this secondary structure element is less populated with increasing temperature.[19] Comparing CD spectra to simulations is difficult as previous attempts at calculating CD spectra from conformational ensembles of disordered peptides have shown large discrepancies between the experimental and calculated spectra.[42,52,53] Also, we are not aware of approaches to deconvolute CD spectra of disordered proteins that takes into account PPII-like signals and which can accurately extract the average content of PPII structure. For these reasons, we instead perform a qualitative comparison of our ensembles to the CD data of Hst5 by calculating the average content of PPII structure in our ensembles and examine its temperature depen-

dence. We find that the ensembles do not show any substantial differences in PPII structure at the different temperatures, with changes of at most 0.6% (Fig. 1d). All together these observations suggest that the force field underestimates the R_h from PFG NMR and does not show changes in compaction and propensity to form PPII structure with temperature.

Figure 1: Average structural features calculated from MD simulations (red squares) at four different temperatures are compared to their experimental counterparts (black circles, when applicable). (a) χ^2 to SAXS data (grey horizontal line indicates $\chi_r^2=1$, that denotes ensembles that correctly fit the experiments; see Methods), (b) R_g, (c) R_h, (d) PPII structure content. As discussed in the main text, the deviation in R_g (panel b), despite good agreement with the SAXS intensities (panel a), may likely in part be attributed to the contribution of the solvation layer to the R_g derived from Guinier fitting.[48]

To test the hypothesis that a decrease in R_h comes with a lowered propensity to form PPII structure, we optimized the agreement between the ensembles and data that encode information about the protein dimension at different temperatures. We first used the iBME[54] approach to reweight the ensembles against SAXS data. With very mild reweighting (Fig. S5), we achieved a good agreement with SAXS data for all simulations (Fig. 2a,b), but without any major change in R_h (Fig. 2c) and PPII structure propensity (Fig. 2d), and only the ensemble simulated at 310 K getting slightly more compact as measured by R_g (Fig. S6). We

note that the larger χ_r^2 before reweighting for the simulation at 310 K (and to a minor extent also that at 323 K) is likely due to a higher signal-to-noise ratio in the experimental SAXS profile at these temperatures compared to those at 283 K and 293 K (Fig. 2a). This difference arises from the use of more concentrated samples in the SAXS measurements at 310 K and 323 K.[19] This, and the fact that we do not observe major changes in the ensembles upon reweighting against SAXS, suggest that the ensembles from PBMetaD fit the experimental SAXS data well at all temperatures and differences in χ_r^2 are due to different magnitudes of the error bars among the experiments at different temperatures. As already suggested by the R_g values extracted from the experimental SAXS profiles, the SAXS data do not suggest any change in average chain dimension with temperature. Even though we do not observe any effect of the temperature on SAXS, this does not mean that the temperature does not influence the average dimension of Hst5 in solution, as SAXS and R_h may probe different aspects of protein compaction.[41,49] The fact that the simulation at 310 K gets more compact (as probed by R_g) upon reweighting, but the R_h does not change substantially, reflects that the measurements are in fact (to a certain extent) decoupled.

We also reweighted the simulations against the R_h from PFG NMR independently from the SAXS data. With the level of rewrighting that we applied, the reweighted R_h values from the simulations at 293 K, 310 K and 323 K come within the error bars associated to the experimental values, while the reweighted R_h of the simulations at 283 K remains below the experimental value (Fig 3a). The sampling at 283 K is more restricted to compact conformations and lacks expanded conformations whose population could be increased by reweighting (Fig. S7). Indeed, by reweighting the simulation at 323 K against the experimental R_h recorded at the lower temperatures, in all cases we achieve a better agreement with the experiments (Fig. S8), because there are more expanded conformations that can be up-weighted to fit the experiments. Remarkably, when improving the agreement with the experimental R_h, we also observe an overall increase in the amount of PPII structure, that follows the same trend suggested by CD data (Fig. 3b). Observing the relationship between conformational

Figure 2: Optimization of the simulations against SAXS data. (a) SAXS profiles of prior (red) and reweighted (blue) ensembles compared with the experiment (grey). (b) χ_r^2 calculated from the profiles in panel a. (c) Ensemble-averaged R_h calculated with either the prior or SAXS-optimized weights. (d) Ensemble-averaged per-residue PPII structure population calculated with either the prior or SAXS-optimized weights.

R_h and number of residues in PPII conformation, we see that, while conformations with only few residues involved in a PPII structure can be either very compact or very expanded, conformations with many residues in a PPII structure are all expanded (Fig. 3c). Indeed, by reweighting the ensembles against the experimental R_h values (that likely increases the weight of conformations with larger R_h), we also observe that conformations with five to ten residues involved in a PPII structure are up-weighted, while conformations with no or little of this secondary structure are down-weighted (Fig. 3d). The down-side arising from reweighting the simulations of Hst5 against R_h is that the agreement with SAXS data decreases (Fig. 3e), that also translates in the ensemble-averaged R_g becoming larger (Fig. S6).

As discussed above, SAXS and R_h might probe different aspects of the compaction of

Figure 3: Optimization of the simulations against R_h from PFG NMR measurements. We show the (a) R_h, (b) average PPII structure propensity and (e) χ_r^2 to SAXS data for the expectation value from the experiment (black, when applicable), the ensemble average from the simulations (red) and the simulations reweighted against the experimental R_h (green). (c) Density plot showing the number of residues involved in a PPII structure as a function of the conformational R_h, together with their relative frequency. (d) Average difference between R_h-optimized weights and prior weights, for frames with a certain number of residues involved in forming a PPII structure.

protein conformations; therefore only optimizing on one of these can lead to ensembles that lack information about the other. We therefore optimize the ensembles simultaneously against SAXS and R_h (see Methods). We find that the result of the combined reweighting is sub-optimal for both SAXS and R_h, if compared to the level of agreement that is obtained when reweighting against only one of these experiments. The χ_r^2 to SAXS ranges between 2 and 4 (Fig. 4a), while the reweighted R_h values are in all cases below the confidence interval of the experimentally derived R_h values (Fig. 4b). Nonetheless, the temperature-dependent trend of the R_h is conserved (Fig. 4b) and so is that of the PPII structure content

(Fig. 4c). The observation that the reweighting appears to compromise between satisfying the SAXS and PFG NMR data, and that it is possible to generate an ensemble for Hst5 that more closely satisfy simultaneously SAXS and R_{h} with the coarse-grained force field CALVADOS,[41,55] suggest that the two force fields sample different conformational ensembles for Hst5.

Figure 4: Optimization of the simulations against SAXS and R_{h} simultaneously. We show the (a) χ^2_r to SAXS data (b) R_{h} and (c) average PPII structure propensity from the experiment (black, when applicable), the ensemble average from the simulations (red) and the simulations reweighted against both SAXS and R_{h} (cyan).

We therefore compared the MD simulation performed at 293 K with a99SB-disp with an ensemble that we previously generated using CALVADOS at 293 K.[41] We find that these two ensembles are highly similar in terms of the R_{g} distribution, but differ substantially in terms of R_{h} (Fig. 5a). If we look at the conformational space covered by the two force fields on a two-dimensional FES defined by R_{g} and R_{h}, it is evident that the free-energy minimum in CALVADOS envelopes the experimental R_{h}, while the free-energy minimum in a99SB-disp has poor overlap with the experimental R_{h}. As a consequence, no conformations are available upon reweighting in a99SB-disp that can satisfy the experimental R_{h} while keeping still the average R_{g} of the ensemble (which leads to a good agreement to SAXS). Indeed, if we look at how the FES for a99SB-disp changes upon reweighting, we observe that, to better fit the experimental R_{h}, the minimum moves along the diagonal of the FES (Fig. 5b), determining a shift in R_{g} and a consequent loss of agreement with SAXS (Fig. 4c).

One aspect that complicates the interpretation of our results, is that there is some uncertainty related to the calculation of the R_h from conformational ensembles of IDPs. Specifically, we previously showed that for short peptides it makes a difference whether the coordinates used in calculating the R_h with the Kirkwood-Riseman equation are those of the C_α atoms or those of the centers of mass (COM) of the residues.[41] If using these last set of coordinates to calculate the R_h, we find that the free-energy minimum for a99SB-disp moves very close to the experimental R_h value (Fig. S9). Currently, it is unclear which of the two approaches is best to describe the R_h of a conformational ensemble of a disordered peptide, and the problem only makes a substantial different for short peptides such as Hst5.[41]

Figure 5: Comparison of the ensembles of Hst5 simulated with a99SB-disp (solid blue lines and blue-to-red FES) and CALVADOS (dashed yellow lines) at 293K. We show the distributions of R_g and R_h both (a) before and (b) after reweighting the a99SB-disp ensemble against SAXS and R_h from PFG NMR. For both simulations with a99SB-disp and CALVADOS we calculate R_h with the Kirkwood-Riseman equation using the C_α coordinates.

Conclusions

Integrative modelling makes it possible to provide atomistic models of IDPs by combining information from experiments probing ensemble-averaged structural features with conforma-

tional models generated for example by MD simulations. Taken alone, many experiments only provide information about a specific structural feature, *e.g.* the average chain dimension, while being mostly insensitive to other properties of the conformational ensemble. Therefore, integrating simultaneously multiple sources of independent experimental information can provide a more realistic determination of the conformational ensemble of an IDP.[6,49,56–58]

We have here focused on Hst5 as a case study, in an attempt to explain the previously observed trend showing the R_h from PFG NMR decreasing with temperature, together with a drop of the PPII structure signal from CD spectra[19] and a relatively temperature-independent R_g from SAXS. We initially faced a complicated picture, where the simulations were mostly in good agreement with SAXS data, but could not reproduce the R_h from PFG NMR and the loss of PPII structure from CD with increasing temperature. We therefore applied a reweighting approach to nudge the prior ensembles in order to achieve a better agreement with the experimental data. We were not able to achieve agreement with the SAXS and R_h data within experimental errors, but pushing the agreement towards the R_h provided a qualitative reproduction of the picture shown by CD spectra. We therefore investigated the reason for the apparent conflict between agreeing with the SAXS and R_h by comparing the simulations with a previously produced coarse-grained ensemble from CALVADOS that showed good agreement with both SAXS and R_h.[41] We found that the two ensembles are very similar in terms of their R_g values, but populate different parts of the R_h space. By consequence, there are no conformations in the a99SB-disp ensemble that can be selected upon reweighting to simultaneously satisfy both SAXS and R_h.

In light of our observations, the first conclusion is that the atomistic force field that we employed does not appear to capture the temperature-dependence shown by the experimental data. This force field, as many others of its kind, has been tested to reproduce mainly R_g extrapolated from SAXS data, and indeed it does a good job in reproducing that experimental information, which does not show any temperature dependence. When it comes to R_h, it seems that the force field is affected by an overall underestimation of the R_h, since

the experimental values are all larger than those calculated from the ensembles. We note, however, that the observed disagreement between the simulations and the R_h values may also be due to inaccuracies in the forward model for R_h including subtle effects of how to apply the Kirkwood-Riseman equation. For short peptides such as Hst5, the choice of the set of coordinates used in the equation can influence the result. If indeed we calculate the Kirkwood-Riseman equation on the COM of the residues instead of C_α, we find a better agreement of the a99SB-disp ensembles with the measurements from PFG NMR. In order to understand these issues better, we would likely need a series of small peptides or proteins with high-quality SAXS and PFG NMR experiments.[41] Furthermore, differences in the experimental conditions used in the SAXS and PFG NMR experiments[19] can potentially lead to different behaviours of the ensembles in solution. In particular, we note the different salt concentrations (150 mM NaCl for the SAXS sample and no salt for the NMR sample) that can affect the extent of interchain interactions.[34] Furthermore, given that 7 out of the 24 residues of Hst5 are histidines, the difference in pH between the SAXS buffer (7.5) and the NMR (7) can affect the protonation state of these residues and consequently affect the ensemble of conformations in solution.[33]

The possibility to calculate accurately CD spectra from conformational ensembles would also aid in understanding what is the correct way of calculating the R_h. Given the observed correlation between R_h and PPII structure content, a quantitative comparison to CD spectra could provide direct information about the possibility that the force field underestimates the PPII structure content of Hst5, together with its R_h. In absence of such a model, we decided to rely on a qualitative analysis of the secondary structures. Based on the backbone dihedral angles, we can calculate the per-residue secondary structure for each frame of our simulations. Doing so, we did not find any temperature dependence on the amount of PPII structure in our simulations, but we could qualitatively reproduce the picture from CD spectra when reweighting the ensembles against the R_h from experiments.

In summary, we have here shown an example of the importance of using multiple and

diverse experiments in order to have accurate and detailed insights into the conformational ensembles of IDPs. Even with such experiments in hand, we have also shown that it is important to have reliable models to calculate experimental observables from computational models. The development of these models is nonetheless impaired by the fact that it is far from trivial to generate accurate conformational ensembles of IDPs that can be used to train these models.[59] In the case that we examined, we have used multiple experiments to show the structural basis of the temperature-dependent behaviour of Hst5. By generating conformational ensembles that reproduce the R_h from PFG NMR measurements, we support the hypothesis that there is a link between how temperature affects the amount of PPII structures and compaction.[15,19,60,61] Some issues remain, that are related to whether the force field employed here underestimates the R_h of Hst5 or not. Future development of forward models, especially for CD, can help in better characterizing the content of PPII structures in IDPs. Furthermore, the study of more IDPs (ideally bigger, although much more difficult to sample efficiently by all-atom MD simulations) that show similar features would be useful to examine whether the behaviour observed for Hst5 occurs in other IDPs.

Methods

Molecular Dynamics Simulations

We performed MD simulations of Hst5 in solution with GROMACS 2018.6.[62] We used the Amber99SB-disp force field[11] with the TIP4P-D water model.[63] The histidine residues in the peptide were modelled as their Nϵ tautomers. We solvated the peptide in a dodecahedric box (with image distance equal to 10 nm) adding Na$^+$ and Cl$^-$ ions to neutralize the system and reach an ionic strength of 150 mM (same as used in the experiments). We ran a two-steps energy minimization, using first the steepest-descent algorithm with energy tolerance set at 200 kJ mol^{-1} nm^{-1} and than the conjugate-gradient algorithm with energy tolerance set at 100 kJ mol^{-1} nm^{-1}. Then, we equilibrated the system at the target temperature

and pressure (1 bar) by running first a 200 ps MD simulation in NVT with position restraints on the protein atoms with the v-rescale thermostat,[64] then a 1 ns MD simulation in NPT adding the Berendsen barostat.[65] For production simulations, we used the Parrinello-Rahman barostat.[66] We used a 2 fs time step in all simulations and constrained all bonds involving hydrogen atoms with the LINCS algorithm.[67]

To accelerate the sampling of the conformational space of Hst5, we employed Parallel Bias Metadynamics (PBMetaD)[43] as enhanced sampling scheme. In PBMetaD a set of user-defined one-dimensional collective variables (CVs) is subject to history-dependent bias potentials that discourage the system from visiting regions of the CV space already explored. Biasing multiple one-dimensional CVs removes the limitation on the number of CVs that affects the classical metadynamics,[37] where a single multidimensional space defined by the CVs is biased. Additionally, as in well-tempered metadynamics,[45] the extent of biasing decreases when the same region of a CV space is visited multiple times, so to have a smoother convergence than with classical metadynamics, and the free-energy surface (FES) as a function of the CVs can be estimated within a well-defined error.[68] In our PBMetaD setup we used 47 CVs, consisting in all 23 ϕ and all 23 ψ dihedral angles and the radius of gyration of the C_α atoms (R_{g,C_α}). We employ R_{g,C_α} as CV to bias because it is cheaper to compute on-the-fly, but in all further analyses we calculate the R_g for all the atoms. Gaussian bias was deposited every 400 ps. We set the initial height of the Gaussian bias to 0.3 kJ mol^{-1}. This height decreased during the simulations by means of the well-tempered feature, according to a bias factor of 24. We used an adaptive algorithm to define the width of the Gaussian bias potentials,[69] where, at each bias deposition, the width is based on the space covered by the CV in the last 2 ps. To enhance the sampling efficiency even further we used a double walker approach,[44] where two simulations run in parallel sharing the deposited bias. For the CV representing the R_{g,C_α}, we set an interval in the CV space between 0.6 and 2.5 nm outside which the biasing force of the PBMetaD does not act.[70] We did so because we observed a 'FES wall climbing effect': since the FES as a function of the R_{g,C_α} is shaped as a single

basin, as bias accumulates, new areas in the CV space, corresponding to more compact and more extended high-energy conformations, gets available (Fig. S9). Since these areas are newly explored, bias is deposited with the initial height, while for the rest of the CV space the extent of biasing is already very low. The advantage of the well-tempered feature in PB-MetaD is that when all the accessible conformations in CV space are explored, the biasing potential decreases at the point of becoming quasi-static and the simulation converges to a well definite FES. But if new areas of the CV space become continuously accessible and a large bias is deposited, the convergence is impaired. Setting the interval in the FES of the R_{g,C_α}, prevents the system from climbing the FES walls corresponding to high-energy conformations that are not relevant. Despite the use of the interval in the FES of the R_g, we still observe few frames depositing a large amount of bias outside the interval (Fig. S4). This, however, is not a problem since our calculations are not based on those few frames that explore regions of the FES outside the interval range. To enable PBMetaD in GROMACS we used PLUMED 2.5.[71]

At the end of a simulation, we recovered the total bias (B) for each conformation x explored and used it to calculate the Boltzmann weights (ω) associated to each conformation according to $\omega_x = \exp(\frac{B_s}{k_b T})$, where k_b is the Boltzmann constant and T is the temperature. The FES for a CV is calculated from the weighted histogram (H) of the CV as $FES(CV) = -k_b T \ln H(CV)$. The error on the FES and ensemble averaged values is calculated by block averaging.[68,72] The package used for these calculations is available at `https://github.com/fpesceKU/BLOCKING`.

Calculation of experimental observables

We compare the ensembles from our simulations to SAXS, PFG NMR and CD data. We calculate SAXS data with Pepsi-SAXS,[73] following the protocol we previously described:[54] the intensity of the forward scattering and constant background are fit to the experimental SAXS profile by least square fitting; the values for the density of the hydration layer and the

excluded volume are scanned on a two-dimensional grid and the combination of those that best fit the data upon reweighting is selected (Table S1).

We calculate the hydrodynamic radius (R_h) using the Kirkwood-Riseman equation[41,74,75] using either the C_α coordinates or those of the center of mass of each residue.

We calculate the per-residue secondary structure with DSSP-PPII,[76] where if two or more consecutive residues that are not assigned to any secondary structure by the DSSP algorithm[77] have backbone dihedrals correspondent to those typical of PPII structure, those residues are assigned as PPII structure. For each residue we then report the propensity for PPII structure formation as a weighted sum of the frames where a residue forms PPII structures, using either weights from PBMetaD or BME (see below).

Ensemble reweighting

We use the Bayesian/Maximum entropy (BME) approach[39] to reweight simulations in order to optimize their agreement with experimental data. This is achieved by minimizing the functional $\mathcal{L}$:[38,40]

$$\mathcal{L}(\omega_1 \cdots \omega_n) = \frac{m}{2} \chi_r^2(\omega_1 \cdots \omega_n) - \theta S_{rel}(\omega_1 \cdots \omega_n) \tag{1}$$

where m is the number of experimental data points, $(\omega_1 \cdots \omega_n)$ are the weights for each simulation frame, χ_r^2 measures the agreement between the calculated and experimental observable. The relative entropy, $S_{rel} = -\sum_{i=1}^{N} \omega_i \log\left(\frac{\omega_i}{\omega_i^0}\right)$, indicates how much the optimized weights (*i.e.* the posterior distribution, ω) diverge from the initial weights (*i.e.* the prior distribution, ω^0), θ is a free-parameter that balances the extent of minimizing the χ_r^2 vs. maximizing S_{rel}. By taking the exponential of S_{rel} ($\phi_{eff} = \exp(S_{rel})$), it is possible to gain insights into the extent of the reweighting. Generally, a low ϕ_{eff} is not desirable, as it means that a low number of frames from the simulation contributes to the calculated weighted averages. Therefore different values for θ are explored in the reweighting procedure, and that corresponding to the χ_r^2 reaching plateau with the highest ϕ_{eff} is chosen to generate the

reweighted ensemble.[39] This helps to ensure a posterior distribution with a good agreement with the experimental data while retaining as much information as possible from the prior distribution. The prior distribution, ω^0 is estimated as the weights from PBMetaD.

To reweight simulations against R_h values from PFG NMR we use the standard implementation of BME. For SAXS data we use the iterative BME (iBME).[54] iBME couples ensemble reweighting with least square fitting of scale factor (s) and constant background (cst) to the calculated SAXS profile. Therefore here the χ_r^2 in the $\mathcal{L}$ functional takes the following shape:

$$\chi_\mathrm{r}^2(\omega_1 \cdots \omega_n, s, cst) = \frac{1}{m} \sum_q^m \frac{[(s \sum_j^n \omega_j I_q(x_j) + cst) - I_q^{\mathrm{EXP}}]^2}{\sigma_{\mathrm{BIFT},q}^2} \tag{2}$$

where $I_q(x_j)$ is the calculated SAXS intensity at scattering angle q for the frame x_j, I_q^{EXP} is the experimental SAXS intensity at scattering angle q. $\sigma_{\mathrm{BIFT},q}$ is the error of the experimental intensity at scattering angle q corrected following the procedure described in.[78]

In reweighting simultaneously against R_h and SAXS we also use the standard BME, using calculated SAXS profiles with scale factor and offset optimized from previous iBME runs. Since the BME functional depends on the χ^2 between calculated and experimental data points, an issue would be that the combined reweighting would tend to optimize more towards the experimental data set with more data points (in this case, the SAXS). To avoid an unbalanced reweighting, we virtually expand the number of data points for the R_h to the same number of data points available in the SAXS profile.

Data and code availability

MD simulations are available at https://doi.org/10.17894/UCPH.4D0D9A09-B40B-4A3A-93EF-D6C249F49A90. PLUMED inputs to enable PBMetaD are available on PLUMED-NEST[79] at https://www.plumed-nest.org/eggs/23/008/. Scripts used to analyze the simulations are available at `https://github.com/KULL-Centre/_2023_Pesce_Hst5`.

Acknowledgement

We thank Marie Skepö and Birthe B. Kragelund for sharing the SAXS, NMR and CD data and for discussions on disordered proteins, and Simone Orioli for discussions about the simulations. This research was funded by the Lundbeck Foundation BRAINSTRUC initiative in structural biology (R155-2015-2666, lundbeckfonden.com). We acknowledge access to computational resources from the ROBUST Resource for Biomolecular Simulations (supported by the Novo Nordisk Foundation grant no. NF18OC0032608) and from the Biocomputing Core Facility at the Department of Biology, University of Copenhagen.

References

(1) Dyson, H. J.; Wright, P. E. Intrinsically unstructured proteins and their functions. *Nature reviews Molecular cell biology* **2005**, *6*, 197–208.

(2) Wright, P. E.; Dyson, H. J. Intrinsically disordered proteins in cellular signalling and regulation. *Nature reviews Molecular cell biology* **2015**, *16*, 18–29.

(3) Bernadó, P.; Blanchard, L.; Timmins, P.; Marion, D.; Ruigrok, R. W.; Blackledge, M. A structural model for unfolded proteins from residual dipolar couplings and small-angle x-ray scattering. *Proceedings of the National Academy of Sciences* **2005**, *102*, 17002–17007.

(4) Uversky, V. N.; Li, J.; Fink, A. L. Evidence for a partially folded intermediate in α-synuclein fibril formation. *Journal of Biological Chemistry* **2001**, *276*, 10737–10744.

(5) Jarvet, J.; Damberg, P.; Danielsson, J.; Johansson, I.; Eriksson, L.; Gräslund, A. A left-handed 3_1 helical conformation in the Alzheimer Aβ (12–28) peptide. *Febs Letters* **2003**, *555*, 371–374.

(6) Jensen, M. R.; Zweckstetter, M.; Huang, J.-r.; Blackledge, M. Exploring free-energy

landscapes of intrinsically disordered proteins at atomic resolution using NMR spectroscopy. *Chemical reviews* **2014**, *114*, 6632–6660.

(7) Schramm, A.; Bignon, C.; Brocca, S.; Grandori, R.; Santambrogio, C.; Longhi, S. An arsenal of methods for the experimental characterization of intrinsically disordered proteins–How to choose and combine them? *Archives of biochemistry and biophysics* **2019**, *676*, 108055.

(8) Palazzesi, F.; Prakash, M. K.; Bonomi, M.; Barducci, A. Accuracy of current all-atom force-fields in modeling protein disordered states. *Journal of chemical theory and computation* **2015**, *11*, 2–7.

(9) Rauscher, S.; Gapsys, V.; Gajda, M. J.; Zweckstetter, M.; De Groot, B. L.; Grubmüller, H. Structural ensembles of intrinsically disordered proteins depend strongly on force field: a comparison to experiment. *Journal of chemical theory and computation* **2015**, *11*, 5513–5524.

(10) Henriques, J.; Cragnell, C.; Skepo, M. Molecular dynamics simulations of intrinsically disordered proteins: force field evaluation and comparison with experiment. *Journal of chemical theory and computation* **2015**, *11*, 3420–3431.

(11) Robustelli, P.; Piana, S.; Shaw, D. E. Developing a molecular dynamics force field for both folded and disordered protein states. *Proceedings of the National Academy of Sciences* **2018**, *115*, E4758–E4766.

(12) Wang, W. Recent advances in atomic molecular dynamics simulation of intrinsically disordered proteins. *Physical Chemistry Chemical Physics* **2021**, *23*, 777–784.

(13) Uversky, V. N. Intrinsically disordered proteins and their environment: effects of strong denaturants, temperature, pH, counter ions, membranes, binding partners, osmolytes, and macromolecular crowding. *The protein journal* **2009**, *28*, 305–325.

(14) Nettels, D.; Müller-Späth, S.; Küster, F.; Hofmann, H.; Haenni, D.; Rüegger, S.; Reymond, L.; Hoffmann, A.; Kubelka, J.; Heinz, B., et al. Single-molecule spectroscopy of the temperature-induced collapse of unfolded proteins. *Proceedings of the National Academy of Sciences* **2009**, *106*, 20740–20745.

(15) Kjaergaard, M.; Nørholm, A.-B.; Hendus-Altenburger, R.; Pedersen, S. F.; Poulsen, F. M.; Kragelund, B. B. Temperature-dependent structural changes in intrinsically disordered proteins: Formation of α–helices or loss of polyproline II? *Protein Science* **2010**, *19*, 1555–1564.

(16) Ciasca, G.; Campi, G.; Battisti, A.; Rea, G.; Rodio, M.; Papi, M.; Pernot, P.; Tenenbaum, A.; Bianconi, A. Continuous thermal collapse of the intrinsically disordered protein tau is driven by its entropic flexible domain. *Langmuir* **2012**, *28*, 13405–13410.

(17) Wuttke, R.; Hofmann, H.; Nettels, D.; Borgia, M. B.; Mittal, J.; Best, R. B.; Schuler, B. Temperature-dependent solvation modulates the dimensions of disordered proteins. *Proceedings of the National Academy of Sciences* **2014**, *111*, 5213–5218.

(18) Zerze, G. H.; Best, R. B.; Mittal, J. Sequence-and temperature-dependent properties of unfolded and disordered proteins from atomistic simulations. *The Journal of Physical Chemistry B* **2015**, *119*, 14622–14630.

(19) Jephthah, S.; Staby, L.; Kragelund, B.; Skepo, M. Temperature dependence of intrinsically disordered proteins in simulations: What are we missing? *Journal of chemical theory and computation* **2019**, *15*, 2672–2683.

(20) Holehouse, A. S.; Pappu, R. V. Collapse transitions of proteins and the interplay among backbone, sidechain, and solvent interactions. *Annual Review of biophysics* **2018**, *47*, 19–39.

(21) Best, R. B. Emerging consensus on the collapse of unfolded and intrinsically disordered proteins in water. *Current opinion in structural biology* **2020**, *60*, 27–38.

(22) Martin, E. W.; Mittag, T. Relationship of sequence and phase separation in protein low-complexity regions. *Biochemistry* **2018**, *57*, 2478–2487.

(23) Dignon, G. L.; Zheng, W.; Kim, Y. C.; Mittal, J. Temperature-controlled liquid–liquid phase separation of disordered proteins. *ACS central science* **2019**, *5*, 821–830.

(24) Mompeán, M.; Oroz, J.; Laurents, D. V. Do polyproline II helix associations modulate biomolecular condensates? *FEBS Open bio* **2021**, *11*, 2390–2399.

(25) Gast, K.; Zirwer, D.; Damaschun, G. Are there temperature-dependent structural transitions in the" intrinsically unstructured" protein prothymosin α? *European Biophysics Journal* **2003**, *31*, 586–594.

(26) Dawson, R.; Müller, L.; Dehner, A.; Klein, C.; Kessler, H.; Buchner, J. The N-terminal domain of p53 is natively unfolded. *Journal of molecular biology* **2003**, *332*, 1131–1141.

(27) Malm, J.; Jonsson, M.; Frohm, B.; Linse, S. Structural properties of semenogelin I. *The FEBS journal* **2007**, *274*, 4503–4510.

(28) Sánchez-Puig, N.; Veprintsev, D. B.; Fersht, A. R. Human full-length Securin is a natively unfolded protein. *Protein Science* **2005**, *14*, 1410–1418.

(29) Jeganathan, S.; Von Bergen, M.; Mandelkow, E.-M.; Mandelkow, E. The natively unfolded character of tau and its aggregation to Alzheimer-like paired helical filaments. *Biochemistry* **2008**, *47*, 10526–10539.

(30) Gast, K.; Zirwer, D.; Damaschun, H.; Hahn, U.; Müller-Frohne, M.; Wirth, M.; Damaschun, G. Ribonuclease T1 has different dimensions in the thermally and chemically denatured states: a dynamic light scattering study. *FEBS letters* **1997**, *403*, 245–248.

(31) Brewer, D.; Hunter, H.; Lajoie, G. NMR studies of the antimicrobial salivary peptides histatin 3 and histatin 5 in aqueous and nonaqueous solutions. *Biochemistry and cell biology* **1998**, *76*, 247–256.

(32) Raj, P. A.; Marcus, E.; Sukumaran, D. K. Structure of human salivary histatin 5 in aqueous and nonaqueous solutions. *Biopolymers: Original Research on Biomolecules* **1998**, *45*, 51–67.

(33) Kurut, A.; Henriques, J.; Forsman, J.; Skepö, M.; Lund, M. Role of histidine for charge regulation of unstructured peptides at interfaces and in bulk. *Proteins: Structure, Function, and Bioinformatics* **2014**, *82*, 657–667.

(34) Cragnell, C.; Durand, D.; Cabane, B.; Skepö, M. Coarse-grained modeling of the intrinsically disordered protein Histatin 5 in solution: Monte Carlo simulations in combination with SAXS. *Proteins: Structure, Function, and Bioinformatics* **2016**, *84*, 777–791.

(35) Jephthah, S.; Henriques, J.; Cragnell, C.; Puri, S.; Edgerton, M.; Skepö, M. Structural characterization of histatin 5–spermidine conjugates: A combined experimental and theoretical study. *Journal of chemical information and modeling* **2017**, *57*, 1330–1341.

(36) McCaslin, T. G.; Pagba, C. V.; Yohannan, J.; Barry, B. A. Specific metallo-protein interactions and antimicrobial activity in Histatin-5, an intrinsically disordered salivary peptide. *Scientific reports* **2019**, *9*, 1–14.

(37) Laio, A.; Parrinello, M. Escaping free-energy minima. *Proceedings of the National Academy of Sciences* **2002**, *99*, 12562–12566.

(38) Hummer, G.; Köfinger, J. Bayesian ensemble refinement by replica simulations and reweighting. *The Journal of Chemical Physics* **2015**, *143*, 243150.

(39) Bottaro, S.; Bengtsen, T.; Lindorff-Larsen, K. In *Structural Bioinformatics: Methods and Protocols*; Gáspári, Z., Ed.; Springer US: New York, NY, 2020; pp 219–240.

(40) Orioli, S.; Larsen, A. H.; Bottaro, S.; Lindorff-Larsen, K. In *Computational Approaches for Understanding Dynamical Systems: Protein Folding and Assembly*; Strodel, B.,

Barz, B., Eds.; Progress in Molecular Biology and Translational Science; Academic Press, 2020; Vol. 170; pp 123–176.

(41) Pesce, F.; Newcombe, E. A.; Seiffert, P.; Tranchant, E. E.; Olsen, J. G.; Kragelund, B. B.; Lindorff-Larsen, K. Assessment of models for calculating the hydrodynamic radius of intrinsically disordered proteins. *bioRxiv* **2022**,

(42) Jephthah, S.; Pesce, F.; Lindorff-Larsen, K.; Skepo, M. Force field effects in simulations of flexible peptides with varying Polyproline II propensity. *Journal of chemical theory and computation* **2021**, *17*, 6634–6646.

(43) Pfaendtner, J.; Bonomi, M. Efficient Sampling of High-Dimensional Free-Energy Landscapes with Parallel Bias Metadynamics. *Journal of Chemical Theory and Computation* **2015**, *11*, 5062–5067, PMID: 26574304.

(44) Raiteri, P.; Laio, A.; Gervasio, F. L.; Micheletti, C.; Parrinello, M. Efficient Reconstruction of Complex Free Energy Landscapes by Multiple Walkers Metadynamics. *The Journal of Physical Chemistry B* **2006**, *110*, 3533–3539, PMID: 16494409.

(45) Barducci, A.; Bussi, G.; Parrinello, M. Well-Tempered Metadynamics: A Smoothly Converging and Tunable Free-Energy Method. *Phys. Rev. Lett.* **2008**, *100*, 020603.

(46) Manalastas-Cantos, K.; Konarev, P. V.; Hajizadeh, N. R.; Kikhney, A. G.; Petoukhov, M. V.; Molodenskiy, D. S.; Panjkovich, A.; Mertens, H. D.; Gruzinov, A.; Borges, C., et al. ATSAS 3.0: expanded functionality and new tools for small-angle scattering data analysis. *Journal of Applied Crystallography* **2021**, *54*, 343–355.

(47) Svergun, D.; Richard, S.; Koch, M.; Sayers, Z.; Kuprin, S.; Zaccai, G. Protein hydration in solution: experimental observation by x-ray and neutron scattering. *Proceedings of the National Academy of Sciences* **1998**, *95*, 2267–2272.

(48) Henriques, J.; Arleth, L.; Lindorff-Larsen, K.; Skepö, M. On the calculation of SAXS profiles of folded and intrinsically disordered proteins from computer simulations. *Journal of molecular biology* **2018**, *430*, 2521–2539.

(49) Choy, W.-Y.; Mulder, F. A.; Crowhurst, K. A.; Muhandiram, D.; Millett, I. S.; Doniach, S.; Forman-Kay, J. D.; Kay, L. E. Distribution of molecular size within an unfolded state ensemble using small-angle X-ray scattering and pulse field gradient NMR techniques. *Journal of molecular biology* **2002**, *316*, 101–112.

(50) Wilkins, D. K.; Grimshaw, S. B.; Receveur, V.; Dobson, C. M.; Jones, J. A.; Smith, L. J. Hydrodynamic Radii of Native and Denatured Proteins Measured by Pulse Field Gradient NMR Techniques. *Biochemistry* **1999**, *38*, 16424–16431, PMID: 10600103.

(51) Yamamoto, E.; Akimoto, T.; Mitsutake, A.; Metzler, R. Universal relation between instantaneous diffusivity and radius of gyration of proteins in aqueous solution. *Physical review letters* **2021**, *126*, 128101.

(52) Fagerberg, E.; Månsson, L. K.; Lenton, S.; Skepö, M. The effects of chain length on the structural properties of intrinsically disordered proteins in concentrated solutions. *The Journal of Physical Chemistry B* **2020**, *124*, 11843–11853.

(53) Gopal, S. M.; Wingbermühle, S.; Schnatwinkel, J.; Juber, S.; Herrmann, C.; Schäfer, L. V. Conformational preferences of an intrinsically disordered protein domain: A case study for modern force fields. *The Journal of Physical Chemistry B* **2020**, *125*, 24–35.

(54) Pesce, F.; Lindorff-Larsen, K. Refining conformational ensembles of flexible proteins against small-angle x-ray scattering data. *Biophysical Journal* **2021**, *120*, 5124–5135.

(55) Tesei, G.; Schulze, T. K.; Crehuet, R.; Lindorff-Larsen, K. Accurate model of liquid–liquid phase behavior of intrinsically disordered proteins from optimization of

single-chain properties. *Proceedings of the National Academy of Sciences* **2021**, *118*, e2111696118.

(56) Fuertes, G.; Banterle, N.; Ruff, K. M.; Chowdhury, A.; Mercadante, D.; Koehler, C.; Kachala, M.; Estrada Girona, G.; Milles, S.; Mishra, A., et al. Decoupling of size and shape fluctuations in heteropolymeric sequences reconciles discrepancies in SAXS vs. FRET measurements. *Proceedings of the National Academy of Sciences* **2017**, *114*, E6342–E6351.

(57) Ahmed, M. C.; Crehuet, R.; Lindorff-Larsen, K. *Intrinsically Disordered Proteins*; Springer, 2020; pp 429–445.

(58) Gomes, G.-N. W.; Krzeminski, M.; Namini, A.; Martin, E. W.; Mittag, T.; Head-Gordon, T.; Forman-Kay, J. D.; Gradinaru, C. C. Conformational ensembles of an intrinsically disordered protein consistent with NMR, SAXS, and single-molecule FRET. *Journal of the American Chemical Society* **2020**, *142*, 15697–15710.

(59) Lindorff-Larsen, K.; Kragelund, B. B. On the potential of machine learning to examine the relationship between sequence, structure, dynamics and function of intrinsically disordered proteins. *Journal of Molecular Biology* **2021**, *433*, 167196.

(60) Shi, Z.; Olson, C. A.; Rose, G. D.; Baldwin, R. L.; Kallenbach, N. R. Polyproline II structure in a sequence of seven alanine residues. *Proceedings of the National Academy of Sciences* **2002**, *99*, 9190–9195.

(61) Shi, Z.; Chen, K.; Liu, Z.; Kallenbach, N. R. Conformation of the backbone in unfolded proteins. *Chemical reviews* **2006**, *106*, 1877–1897.

(62) Abraham, M. J.; Murtola, T.; Schulz, R.; Páll, S.; Smith, J. C.; Hess, B.; Lindahl, E. GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX* **2015**, *1-2*, 19–25.

(63) Piana, S.; Donchev, A. G.; Robustelli, P.; Shaw, D. E. Water Dispersion Interactions Strongly Influence Simulated Structural Properties of Disordered Protein States. *The Journal of Physical Chemistry B* **2015**, *119*, 5113–5123, PMID: 25764013.

(64) Bussi, G.; Donadio, D.; Parrinello, M. Canonical sampling through velocity rescaling. *The Journal of Chemical Physics* **2007**, *126*, 014101.

(65) Berendsen, H. J. C.; Postma, J. P. M.; van Gunsteren, W. F.; DiNola, A.; Haak, J. R. Molecular dynamics with coupling to an external bath. *The Journal of Chemical Physics* **1984**, *81*, 3684–3690.

(66) Parrinello, M.; Rahman, A. Polymorphic transitions in single crystals: A new molecular dynamics method. *Journal of Applied Physics* **1981**, *52*, 7182–7190.

(67) Hess, B.; Bekker, H.; Berendsen, H. J. C.; Fraaije, J. G. E. M. LINCS: A linear constraint solver for molecular simulations. *Journal of Computational Chemistry* **1997**, *18*, 1463–1472.

(68) Bussi, G.; Tribello, G. A. In *Biomolecular Simulations: Methods and Protocols*; Bonomi, M., Camilloni, C., Eds.; Springer New York: New York, NY, 2019; pp 529–578.

(69) Branduardi, D.; Bussi, G.; Parrinello, M. Metadynamics with Adaptive Gaussians. *Journal of Chemical Theory and Computation* **2012**, *8*, 2247–2254, PMID: 26588957.

(70) Baftizadeh, F.; Cossio, P.; Pietrucci, F.; Laio, A. Protein Folding and Ligand-Enzyme Binding from Bias-Exchange Metadynamics Simulations. *Current Physical Chemistry* **2012**, *2*, 79–91.

(71) Tribello, G. A.; Bonomi, M.; Branduardi, D.; Camilloni, C.; Bussi, G. PLUMED 2: New feathers for an old bird. *Computer Physics Communications* **2014**, *185*, 604–613.

(72) Flyvbjerg, H.; Petersen, H. G. Error estimates on averages of correlated data. *The Journal of Chemical Physics* **1989**, *91*, 461–466.

(73) Grudinin, S.; Garkavenko, M.; Kazennov, A. *Pepsi-SAXS*: an adaptive method for rapid and accurate computation of small-angle X-ray scattering profiles. *Acta Crystallographica Section D* **2017**, *73*, 449–464.

(74) Kirkwood, J. G.; Riseman, J. The intrinsic viscosities and diffusion constants of flexible macromolecules in solution. *The Journal of Chemical Physics* **1948**, *16*, 565–573.

(75) Clisby, N.; Dünweg, B. High-precision estimate of the hydrodynamic radius for self-avoiding walks. *Physical Review E* **2016**, *94*, 052102.

(76) Mansiaux, Y.; Joseph, A. P.; Gelly, J.-C.; de Brevern, A. G. Assignment of PolyProline II conformation and analysis of sequence–structure relationship. *PloS one* **2011**, *6*, e18401.

(77) Kabsch, W.; Sander, C. Dictionary of protein secondary structure: pattern recognition of hydrogen-bonded and geometrical features. *Biopolymers: Original Research on Biomolecules* **1983**, *22*, 2577–2637.

(78) Larsen, A. H.; Pedersen, M. C. Experimental noise in small-angle scattering can be assessed using the Bayesian indirect Fourier transformation. *Journal of Applied Crystallography* **2021**, *54*, 1281–1289.

(79) Promoting transparency and reproducibility in enhanced molecular simulations. *Nature methods* **2019**, *16*, 670–673.