A phylogeny-informed characterization of global tetrapod traits addresses data gaps and biases

Mario R. Moura[1,2,3,4*], Karoline Ceron[1], Jhonny J. M. Guedes[5], Yanina V. Sica[2,3], Alessandro Catenazzi[6], Fernanda P. Werneck[7], Luís Felipe Toledo[1], and Walter Jetz[2,3*]

1. Departamento de Biologia Animal, Universidade Estadual de Campinas, Campinas, SP 13083-862, Brazil
2. Department of Ecology and Evolutionary Biology, Yale University, New Haven CT, 06511, USA
3. Center for Biodiversity and Global Change, Yale University, New Haven CT, 06511, USA
4. Departamento de Biociências, Universidade Federal da Paraíba, Areia, PB 58397-000, Brazil
5. Programa de Pós-Graduação em Ecologia e Evolução, Departamento de Ecologia, Universidade Federal de Goiás, Goiânia, GO 74690-900, Brazil
6. Department of Biological Sciences, Florida International University, Miami FL 33199, USA
7. Programa de Coleções Científicas Biológicas, Coordenação de Biodiversidade, Instituto Nacional de Pesquisas da Amazônia, Manaus, AM 69067-375, Brazil
*Corresponding author: Mario R. Moura (mariormoura@gmail.com), Walter Jetz (walter.jetz@yale.edu)

Abstract

Tetrapods (amphibian, reptiles, birds and mammals) have become a central model system for global ecology, conservation, and biodiversity science at large. But continuing data gaps, limited data standardisation, and ongoing dynamics of taxonomic nomenclature put constraints on robust integrative research in this group and potentially causes biased inference. Here we combine and harmonise taxonomic, spatial, phylogenetic, and attribute data with phylogeny-based imputation to provide a comprehensive assessment of data gaps and biases across taxa and geographic space. We find that co-missingness in attribute values increased with taxon-level completeness and richness. Prediction of missing attribute values using extensive performance and uncertainty assessment revealed substantial changes to macroecological patterns. These results highlight both the specific biases incurred by non-random data missingness and an important avenue to address them. We provide a comprehensive data resource that includes predictions and address 75 attributes, including body size, activity time, micro- and macrohabitat, geography, environmental preferences, human influence and threat status for all 33,281 tetrapod species covered in recent fully-sampled phylogenies. While there is an obvious need for further data collection and ongoing updates, the presented phylogeny-associated and -informed resource of tetrapod species attributes can support a better representation of undersampled species and their attributes in ecology and conservation.

Introduction

Over the past two decades biodiversity science has seen a dramatic growth in impactful large-scale ecological and conservation research enabled by near-global study systems such as terrestrial vertebrates, or tetrapod (amphibians, reptiles, birds, and mammals). These efforts are usually reliant on datasets spanning wide temporal, spatial, and taxonomic scales [1] that ideally are fully harmonized and well curated. Although terrestrial vertebrates are considered well-known animal groups relative to invertebrates and marine taxa [2,3], significant data gaps remain [4,5]. Consequently, trait-based research on biodiversity conservation, ecology and evolution is often performed with incomplete data.

Because this data missingness is usually highly non-random, i.e. overall trait space is represented unevenly [4], it can have pervasive consequences inference in trait-based research

[6,7] and conservation applications [6,8]. Indeed, available trait data are recognized to be biased towards certain taxa, regions, and traits [4], and often there is co-missingness, i.e. missing values for a given variable are associated with missingness in another [9]. Such congruent or aggregated (as opposed to segregated) patterns in trait data missingness can arise from societal and research preferences towards either charismatic species [3] or taxonomic groups that are easily sampled [6,10], whereas segregated patterns may reveal traits difficult to sample or even underlooked for a given taxonomic group. Documenting which groups show aggregated or segregated data missingness helps to inform strategies of data acquisition and reduce existing biases in trait databases.

While sampled trait data remain limited for many tetrapod species attributes, new methods can help to minimise these data gaps and improve our understanding of biodiversity. Past practices have included the removal of species with missing data or replacement of missing values by observed averages, but these strategies inevitably contribute to reduce statistical power and increase bias [7,9,11]. More recently, the growth in large-scale phylogenetic frameworks have boosted the development of phylogeny-based methods to impute missing values [12–15]. The phylogenetic information can be further combined with available data on other correlated traits to increase the quality of imputed values [16,17]. Among tetrapods, recent large-scale applications of imputation methods include, for example, the use of either phylogenetic regression methods or machine-learning techniques to predict missing values in trait data for amphibians [18,19], reptiles [19,20], birds [19,21], and mammals [19,21,22], as well as to inform threat statuses for data deficient and non-assessed species [20,23].

Here we leverage these methodological innovations and combine them with additional data mobilization to provide a comprehensive evaluation and prediction of key ecological attributes of all extant 33,281 tetrapod species covered in recent fully-sampled phylogenies, including 7,238 amphibians [24], 384 chelonians and crocodilians [20], 9,755 squamates and tuatara [25], 9,993 birds [26], and 5,911 mammals [27]. Specifically, we assessed 75 standardised species-level attributes combining a range of sources on taxonomy, body size, activity time, microhabitat, macrohabitat, geography, environmental preferences, human influence, and threat status. As not all species placements in the available phylogenies are yet supported with genetic data, representing one important source of uncertainty in their placement, we also evaluated phylogenetic data gap, i.e. the remaining Darwinian shortfall [28]. We taxonomically harmonised data and fill data gaps with phylogeny-based imputation methods [12–15] for which we verified the performance and remaining uncertainty through extensive validation procedures.

We use this combined dataset to assess the geographic, taxonomic, and trait-related biases imposed by existing data gaps and how their model-based closure supports improved information and biological inference. Because of a biodiversity knowledge paradox – the high biodiversity over tropics [29] but better sampling in temperate regions [4,5,30] – we expect larger unsampled fractions of the tetrapod trait space for species occurring in tropical than temperate regions. Similarly, the high research capacity (i.e., infrastructure and expertise availability) towards birds and mammals relative to amphibians and reptiles [2,3] contribute to the uneven sampling of trait space [4]. We expect larger biases in trait data among historically less sampled taxa. Finally, species' biology and sampling methodologies are known to affect species detection and collection rates in the field [31–33]. For instance, species detectability can be lower for small- than large-bodied species, and similarly so for nocturnal relative to diurnal species [34–36], whereas sampling methodologies often favour the collection and research of terrestrial and semi-terrestrial species [37–41]. We thus expect some convergent missingness across the tetrapod trait space, with undersampled species being typically small-sized, nocturnal, and fossorial or arboreal.

Methods

We reviewed available databases on global tetrapod groups and used the latest phylogeny-
based methods to create the most comprehensive tetrapod attribute dataset to date. Our
database includes various fields that track data sources and imputation techniques, which we
collectively refer to as 'attributes', along with trait data. Our procedures can be summarised in
five general steps: (i) data acquisition, (ii) taxonomic harmonisation, (iii) outlier verification,
(iv) taxonomic imputation, and (v) phylogenetic imputation. Details on the validation of our
predictions and imputation performance are available in the Supplementary Information.
116

Data Acquisition

We compiled 75 species-level attributes on taxonomy, body size, activity time, microhabitat,
macrohabitat, threat status, geography, environmental preferences, and associated data sources
for each tetrapod species. We gathered information available in several global, continental,
and regional databases, and complemented the existing data by inspecting the published
(articles, book chapters, field guides, and monographs) and grey literature (e.g., technical
reports, government documents, monographs, theses), besides incorporating unpublished data
gathered by us. To minimise the uneven representation of ecological attributes across clades,
we identified genera and families whose species did not have data on body size, activity time,
or microhabitat. We then used species belonging to such genera and families to feed
additional online searches through academic platforms (Google Scholar and Web of Science)
and included complementary attribute data whenever possible. We did not use the imputations
available at the genus level in the data sources we examined.
130

Briefly, taxonomic data were represented by taxonomic ranks, authority name, and year of
description. Body size data consisted of information on body length (mm) and body mass (g).
Activity time informed if the species was diurnal and/or nocturnal. Cathemeral or crepuscular
species were considered as both diurnal and nocturnal. Microhabitat informed five categories
of habitat use commonly reported in field guides and related literatures: fossorial, terrestrial,
aquatic, arboreal, and aerial. Microhabitat data for birds were adapted from the EltonTraits
database [42], which described the estimated relative usage for seven types of foraging
stratum. We reduce these seven categories to four by: summing the relative usage of species
foraging below water surface or on water surface in the aquatic microhabitat; summing the
relative usage of species foraging on the ground and below 2m in understory as terrestrial
microhabitat; summing the relative usage of species foraging 2m upward in canopy and just
above canopy as arboreal microhabitat. Species with aerial microhabitat were kept as
informed in EltonTraits database [42], and no fossorial bird was reported in the later source.
We then made binary the relative usage of aquatic, terrestrial, arboreal, and aerial microhabitat
using the threshold of 30% to consider a species as typical of a given microhabitat type.
146

Some attributes in the Tetrapod Traits database were derived from expert-based range maps
for amphibians [18,43,44], reptiles [20,29,43], mammals [43–45], and birds [26,44]. We
matched the authoritative expert range maps for each of the tetrapod groups with the
corresponding phylogenies and edited species ranges to ensure they represented the species
concept adopted in the corresponding phylogeny. Overall, the verification procedure for
species range maps can be summarised in eight scenarios: i) no changes, where species range
maps matched directly with binomials in the phylogeny, ii) synonyms, where species range
maps were direct synonyms to binomials in the phylogeny, requiring therefore only an
updated name, iii) no range, where extinct and domesticated species in the phylogeny did not
need a range map, iv) split, where species range maps needed to be clipped from a parent
species, or when parent species needed to have part of their range removed, vi) lumps, where
species range maps needed to be lumped back with another species range, vii) new species,
where there was no available range map and ranges were drawn from other literature sources
or a minimum convex polygon with a buffer needed to be drawn around type localities, and
viii) *de Novo* species whose ranges needed to be removed because they were not in the
phylogeny.
163

We measured range size as the number of 110×110 km equal-area grid cells intersected by each species. This spatial resolution was selected because it minimises the presence of errors related to the use of expert ranges maps [46–48]. Presence of a species in a grid cell was recorded if any part of the species distribution polygon overlapped with the grid cell. We then computed spatially-derived attributes as the within-range average value of continuous variables (e.g., annual mean temperature, proportion of urban area) weighted by species range occupancy in the cell, or as the proportion of the species range occupied by levels of categorical variables (e.g., proportion of the species range overlapped by each biogeographical realm). All within-range attributes were derived from spatial layers available online [49–52] and individually described in the Results and Discussion section. We did not derive range maps for species with uncertain geographic distribution, which refers to those with reported occurrences only at the level of administrative units (e.g., country, state) without precise location information.

Species macrohabitat data followed the IUCN Habitat Classification scheme v. 3.1 [43]. Briefly, the IUCN scheme describes 17 major habitat categories in which species can occur, besides an 18[th] category for species with unknown major habitat. We used the *rredlist* package [53] to obtain macrohabitat data directly from IUCN for 31,464 species (94.5% of all species herein considered). Our macrohabitat variables correspond only to the first level of IUCN Habitat Classification scheme. Other 157 species had macrohabitat data extracted from pertinent literature. We also used the *rredlist* package [53] to obtain non-DD assessed status for 28,885 tetrapod species based on IUCN red list v. 2021-3 [43]. For other 286 species, we used non-DD assessed statuses matching those informed in previous works using the same taxonomy of fully-sampled trees [18,20,25,54]. Data deficient species totalled 3,476 species. We did not find an assessed status for 646 species. Overall, the data acquisition involved the inspection of more than 2,600 references (Table 1). Full description of attributes and extra details on all sources consulted for each species are provided in the Results and Discussion section.

Taxonomic Harmonisation
The taxonomy of the Tetrapod Traits database followed the respective taxonomies of the recent, fully-sampled phylogenies for each of the major tetrapod groups [20,24–27]. The amphibian phylogeny taxonomy [24] followed the 19 February 2014 edition of AmphibiaWeb (http://amphibiaweb.org), with 7,238 species. The phylogeny for chelonians [20] followed the 8[th] ed. Of the Turtles of the World checklist [55], with the addition of *Aldabrachelys abrupta* and *Al. grandidieri*, and the synonymisation of *Amyda ornata with Am. cartilaginea*, adding up to 357 chelonian species, whereas for crocodilians [20], the taxonomy followed [56], complemented by the revalidation of *Crocodylus suchus* [57], *Mecistops leptorhynchus* [58], and recognition of three *Osteolaemus* species [59,60], resulting in 27 species. The taxonomy of the squamate and tuatara phylogeny [25] followed the Reptile Database update of March 2015 (http://www.reptile-database.org) with 9,755 species. For brevity, we hereafter refer to species in the later phylogeny as squamates, although we recognise that the tuatara, *Sphenodon punctatus,* is not a squamate. The taxonomy of the bird phylogeny [26] followed the Handbook of the Birds of the World [61], including 9,993 species. Finally, the taxonomy of the mammal phylogeny [27] followed IUCN [62] with modifications, and included 5,911 species.

To maximise the data usage from previous compilation efforts, and to ensure coherence among species names and the multiple data sources, we built lists with synonymies and respective valid names based on multiple taxonomic databases [43,63,64], and extracted the unique synonyms within each taxonomic database. By unique synonym, we refer to a binomial (scientific name valid or not) applied to only one valid name. We then performed a taxonomic reconciliation based on four steps:

1. ***Direct match with data sources:*** scientific names between the Tetrapod Traits (33,281 species) and each potential data source were directly paired. However, species-level attributes of close-related species may show identical values if extracted from earlier sources that treated different species as synonymies. To minimise the inclusion of duplicated values among the species-level attributes, we flagged each taxonomic match between the Tetrapod Traits database and external data sources to use each data entry only once.

2. ***Direct match with data source synonymies:*** species without a direct match after the step 1 had their taxonomy updated in the data sources using the list of unique synonymies, and a new matching operation allowed the extraction and flagging of additional data whenever possible. At the end of this procedure, all species had their scientific names restored to the taxonomy originally informed.

3. ***Direct match with Tetrapod Traits synonymies:*** some data sources may follow more recent taxonomies than those inherited from the fully-sampled phylogenies [20,24–27]. For species without a direct match after the step 2, we updated their taxonomy in the Tetrapod Traits using the list of unique synonymies and repeated the procedures of data extraction and flagging. At the end of this procedure, all species had their scientific names restored to the taxonomy originally informed.

4. ***Manual verification:*** species without a direct match after the step 3 had their scientific names individually searched by hand in specialised taxonomic databases (amphibians [63], reptiles [64], birds [65], and mammals [66]) for potential spelling errors and/or additional synonymies not included among our synonym lists. Whenever possible, we updated the taxonomy applied to data sources and then repeated the data extraction and flagging procedure. Species without data after step-4 were considered as missing data.

Outlier Verification

We implemented two approaches to detect potential inconsistencies in continuous attribute data before applying phylogeny-based methods to impute missing values.

1. ***Interquartile Range Criterion:*** species body length and body mass were $\log_{10}$ transformed and then flagged as a suspicious outlier if their value were outside the interval defined between $[q_{0.25} - 1.5 \times IQR]$ and $[q_{0.75} + 1.5 \times IQR]$, where $q_{0.25}$ and $q_{0.75}$ are respectively the first and third quartiles, and IQR is the interquartile range $[q_{0.75} - q_{0.25}]$.

2. ***Deviation from allometric relationship:*** we used interactive scatterplots to flag species with unusual deviations from the expected allometric relationship between body length and mass. We inspected allometric relationships separately for species within each Class, Order, and Suborder.

Species flagged by any of the above-mentioned approaches had their body length and/or body mass checked for validity and corrected if necessary, or even purged from the database if not confirmed by a reliable source.

Taxonomic Imputation

The global scope of the present database inevitably comes with gaps that are hard to fill, such as the scarcity of natural history data for species known from a few specimens or only the holotype [67,68]. Previous studies have tackled this challenge by using imputed values at genus-level or close-relatives [42] to reduce data missingness. Although these earlier strategies of 'taxonomic imputation' might reduce variability in attribute values, they are useful to fill gaps in highly conserved attributes, and can ultimately help increase the performance of phylogeny-based imputation methods applied in concert with correlated attribute data [16,17].

We used taxonomic imputations for two particular cases of missingness in microhabitat data: chiropterans with missing data were considered as 'aerial' (113 species), and dolphins and whales were considered as 'aquatic' (5 species). For the remaining tetrapod species, we computed the per-genus proportion of species in each type of activity time (diurnal or nocturnal), microhabitat (fossorial, terrestrial, aquatic, arboreal, aerial), and macrohabitat (17 binary variables informing the IUCN Habitat Classification scheme). If a type of activity time, microhabitat, or macrohabitat appeared in at least 70% of species in the genus, we then assumed such ecological attribute as also present among species with missing data in the respective genus. We did not use taxonomic imputation for continuous attributes. Our goal was to reduce missingness in activity time, microhabitat, and macrohabitat for groups with a well-known ecology (observed data available for at least 70% of species) before running phylogeny-based imputation methods (Table 1).

Phylogenetic Imputation

To minimise missing values in our database we used two imputation techniques proven as robust in recent methods comparisons [12,13], namely *Rphylopars* [14,15] and *missForest* [13]. The *Rphylopars* method uses the phylogenetic tree directly to impute missing values based on the phylogenetic covariances and correlated evolution between numeric attributes, whereas *missForest* is based on a random forest algorithm and allow mixed-type data.

Because of the large dimensionality of data in our analysis, we only performed imputations for body length, body mass, microhabitat, and activity time. All spatially-derived attributes showed more than 99% of data completeness (Table 1). We did not perform phylogeny-based imputations for species lacking range maps and, consequently, spatially-derived attributes. We $\log_{10}$-transformed the continuous variables (body length and body mass) to reduce skewness, and used binary variables to represent each type of microhabitat (fossorial, terrestrial, aquatic, arboreal, aerial) and activity time (diurnal and nocturnal). Note that microhabitat and activity time types are non-mutually exclusive. To account for uncertainty in fully-sampled phylogenies, we used 100 trees of each tetrapod group [20,24–27] to compute imputations. We ran *Rphylopars* using Brownian motion (BM) model of trait evolution and extracted the median value across the 100 trees for each imputed attribute per species. Because *Rphylopars* only allow continuous attributes [15], the imputed values for binary variables could assume values outside the range between 0–1. Thus, for each tetrapod group and binary attribute, we binarised imputed types of microhabitat and activity time using the 0.5 threshold.

The *missForest* algorithm does not include phylogenetic tree directly in computations [13]. To account for phylogenetic information, we selected the first 10 eigenvectors from the phylogenetic covariance matrix of each fully-sampled tree and used them in concert with the observed attribute data in the imputation routine. For each tetrapod group and across the subset of 100 trees, the selected phylogenetic eigenvectors explained on average 96.8% of the phylogenetic variation across the 100 trees of each tetrapod group (97.8% for amphibians, 99.4% for chelonians and crocodilians, 97.1% for squamates, 92.4% for birds, and 97.6% for mammals). For continuous attributes (body length and body mass), we extracted the median imputed value per species across the 100 trees. Imputed binary attributes (types of microhabitat and activity time) had their frequency computed across the 100 trees, with frequencies later made binary using the 0.5 threshold.

For birds only, we complemented the observed attribute data with 10 morphometric traits ($\log_{10}$-transformed) made recently available through the AVONET database [69], and then repeated the imputation routine using the subset of 100 fully-sampled bird trees. Given the high dimensionality of these computations, they became computationally intractable under the *Rphylopars* approach and were only performed through *missForest*. The model parameters used in *missForest* analysis for all tetrapod groups included: 500 trees to grow each decision forest (argument *ntree*), the square-root of total number of predictors as the number of

variables randomly sampled at each split (default of the argument *mtry*), and a maximum of
10 iterations for *missForest* reach the stopping criterion (default of the argument *maxiter*).

We selected the imputed values returned by the imputation approach (*Rphylopars* or
missForest) identified as most suitable (see Supplementary Information, S1 Text) for each
combination of tetrapod group (amphibians, chelonians and crocodilians, squamates, birds,
and mammals) and attribute data (body length, body mass, microhabitat, and activity time).
Additional fields clearly labelled all imputations included in the Tetrapod Traits database.
Computations were performed in R v. 4.1.0 using the packages *Rphylopars* [15], *ape* [70],
gmodels [71], and *missForest* [13].

Table 1. Summary of major groups of attributes, species counts, and sources in Tetrapod
Traits database. 'Imputed' counts indicate the number of species with taxonomic (Taxon.) or
phylogenetic (Phylog.) imputations computed. References used for data acquisition are shown
for each attribute and tetrapod group.

Attribute group	Number of species			Number of sources	Major Sources per Tetrapod group				
	Obs. Data	Taxon. Imputed	Phylog. Imputed		Amphibians	Turtles and Crocs	Squamates	Birds	Mammals
Body length	24,472	0	8,809	1559	[72–76]*	[20]*	[77]*	[78–80]*	[22,78]*
Body mass	26,613	0	6,668	460	[72,73,81–84]*	[20]*	[77]*	[42,69,80]*	[22,42,78,85–88]*
Activity time	24,512	588	8,181	347	[72,89–93]*	[94]*	[95–106]*	[42]*	[22,42,82,83,87]*
Microhabitat	31,470	1,125	686	688	[72,89,107]*	[20]*	[95–98,100–112]*	[42,107]*	[22,42,107]*
Macrohabitat	31,621	1,440	0	140	[43,44]*	[43,44]*	[43,44]*	[43,44]*	[43,44]*
Geography	33,125	0	0	7	[18,43,44]	[20,43,44]	[29]	[26,44]	[17,19,22–26]
Environment	33,125	0	0	3	[49,50]	[49,50]	[49,50]	[49,50]	[49,50]
Human Influence	33,125	0	0	3	[51,52]	[51,52]	[51,52]	[51,52]	[51,52]
Threat Status	32,647	0	0	6	[18,43,113]	[20,43,114]	[43,115]	[43,54]	[43,116]

*For brevity, the references listed include only sources contributing with at least 30 species.
All species-specific references are included in the raw files of the Tetrapod Traits database.

Patterns in Data Co-Missingness

We assessed co-occurrence patterns in data missingness across species using the
'checkerboard score' (C-score; [117]), which is less prone to type II errors than other co-
occurrence metrics computed under a null model approach [118]. The C-score was based on a
binary presence-absence matrix (PAM) of species (columns) by missing attributes (rows), the
latter represented by five binary variables informing the missingness of observed values in
body length, body mass, activity time, microhabitat, and threat status. Missing data in threat
status was represented by DD or non-assessed species. We computed the C-score using
individual PAMs for each genus and family with at least 2 species, and each pairwise
combination of attribute variable, following recommendations by [119].

To verify whether an observed C-score differed from the value expected by chance, we built a
null distribution of C-score values through a randomization algorithm that kept attribute
completeness (rows sums) fixed while considering species as equiprobable regarding their
chances of showing missing attribute values. Null distributions were built for each individual
PAM using 10K iterations with a burn-in of 500. We then computed the standardised effect-
size (SES) of C-score and associated p-values and identified the pairwise attribute
combinations with aggregated (SES C-score < 0 and p < 0.05) or segregated (SES C-score > 0
and p < 0.05) co-occurrence pattern of data missingness per taxa. Computations were
performed using the *EcoSimR* [120] package in R.

Patterns and Biases in Imputed Traits

If certain parts of the multivariate attribute space are missing, the inclusion of imputed values
may fill such 'invisible fractions' and change emergent attribute properties per taxon or

geographical assemblage (e.g., average value, geometric mean, proportion or prevalence of a certain category). But changes in attribute property per taxon can also arise from the inclusion of biased imputed values [16], since imputation errors can increase either with the percentage of missing data or when multiple attributes are imputed simultaneously [12,16,17]. To verify how the inclusion of imputed values affected attribute patterns per taxa, we calculated the per-genus and per-family relative change in attribute property after adding imputed values.

For continuous attributes (body length and mass), the attribute property was represented by the geometric mean attribute value. For binary attributes, we scored types of microhabitat and activity time into continuous metrics of verticality (0 = strictly fossorial, 0.25 = fossorial and terrestrial, 0.5 = terrestrial or aquatic, 0.75 = terrestrial and arboreal, and 1 = strictly arboreal or aerial) and nocturnality (0 = strictly diurnal, 0.5 = cathemeral or crepuscular, 1 = strictly nocturnal). We then extracted the mean verticality and nocturnality per taxon and geographical assemblage. Relative change in attribute values was represented by the ratio of attribute property with imputations (i.e., mean computed using observed + imputed data) to attribute property without imputations (mean computed using observed data only):

$$Relative\ Change = 1 - (Attribute\ Property_{obs+imp} / Attribute\ Property_{obs})$$

Positive values of relative change indicate the proportional increase in attribute property after inclusion of imputed values, whereas negative values show the opposite. We used the R function *tree.merger* of the *RRphylo* package [121] to build a tetrapod super tree by combining the global phylogenies available for amphibians [24], chelonians and crocodilians [20], squamates [25], birds [26], and mammals [27]. The full tetrapod tree topology was only used for visualization purposes. In the presence of severe data limitations (e.g., very low completeness, highly uneven sampled data, and lack of inter-correlated attributes), the relative changes in attribute properties could arise due to estimation errors instead of representing unsampled fractions of the attribute space. We assess whether the relative increase or decrease of attribute properties differed with respect to the number of simultaneously imputed attributes through Kruskal-Wallis tests.

Results and Discussion
Data Completeness
We focus our assessment of data gaps on the attributes with considerable missingness, body length, body mass, activity time, microhabitat, and threat status. Across these five focal attributes, we observed complete species-level data for 40.7% of tetrapod species (n = 13,540), 31.5% of genera (n = 1,609), and 19.7% of families (n = 102). Seven tetrapod genera showed no data for these five attributes, all of them were monotypic rodents (*Calassomys*, *Halmaheramys*, *Musseromys*, *Paucidentomys*, and *Waiomys*) or bats (*Dryadonycteris* and *Hsunycteris*). No family showed zero completeness for these five attributes combined. Lowest completeness levels were among attributes related to body length (73.5%), activity time (73.6%), body mass (80%), and microhabitat (94.5%, Table S1-S2). While assessed threat status was available for 98% of tetrapod species, this number decrease to 87.6% if DD species were treated as missing data (Fig. 1).

At the level of geographic assemblages, attribute completeness showed large variation for body length, with Neotropical and Afrotropical assemblages presenting the lowest completeness (Fig. 2A). The low spatial variability in attribute completeness for body mass, activity time, microhabitat, and assessed threat status (Fig. 2B–E) indicated that species with missing data are mostly narrow-ranged [4], which can limit the influence of data missingness in assemblage-level patterns. Of the 6,893 species missing at least two attributes (body length, body mass, activity time, microhabitat, and threat status), 77% occurred in 10 or less 110 × 110 km equal-area grid cells. If we restrict missingness to three or more attributes, then 87% of species filtered occurred in ≤10 grid cells. The completeness of phylogenetic data in the

fully-sampled trees of tetrapods [20,24–27] also showed high variation in spatial coverage (Fig. 2F), with Afrotropical and Indomalayan realms emerging with the highest Darwinian shortfall [28]. Overall, our findings help illustrate that geographical patterns in data coverage can be strongly influenced by data for wide-ranged species.

Data Co-Missingness

We found non-random co-occurrence patterns in data missingness for 20% of the pairwise attribute combinations at genus-level and 43% at family-level. Almost all non-random co-occurrence patterns showed aggregation of missing values, that is, data missingness was shared among multiple attributes for a same species, particularly among squamates and amphibians (Fig. 3A). In inspecting the co-missingness between threat status and other attributes, we found leaf-litter amphibians (*Craugastor*, *Eleutherodactylus*, and *Pristimantis*, all within the superfamily Brachycephaloidea) standing out among tetrapod genera with highest aggregation in missing values (Fig. 3B). These findings reinforce the view of amphibians and squamates as the least researched tetrapod classes [2,3], besides indicating the uneven sampling coverage of attribute space within certain clades.

Highly aggregated patterns of data missingness were more common among species-rich clades with lower proportion of missing data (Figs 3 and S5). That is, co-missingness was strongest in species-rich and well-sampled taxa, particularly in those holding high proportion of species known only from their type locality or holotype [67,122]. About 86% of the genera and 74% of families within the top 50 taxa in co-missingness (Figs 3 and S5) include 'lost taxa', that is, species that have not been reliably observed in >50 years and yet are not declared extinct [68]. While the rediscovery of such rarely observed species is not uncommon [122], their absence in attribute databases will not be easily filled through field observations [123], which could impair biogeographical research and conservation practice.

Biases uncovered by imputation-based gap filling

We found that the gap filling provided by the imputations resulted in substantial changes in aggregate attribute properties and the insights they would support. These changes occurred in both directions and were strongest in continuous attributes (body length and body mass). About 65% of tetrapod genera and 78% of tetrapod families include at least one species with phylogeny-based imputations. After the inclusion of imputed values, the average attribute value per genus decreased in 5.9% of attribute-genus combinations, and increased in 5%. At the family-level, 14% of attribute-family combinations showed increased in the average attributes after the inclusion of imputed attributes whereas 20% decreased (Fig. 4).

Changes in attribute properties at the level of geographic assemblages revealed substantial changes in the average body size in tropical regions (Fig. 5A-F). In contrast, such differences are largely absent in the temperate regions, reflecting the dedicated mobilization and resulting large availability of biodiversity data in developed countries [2,4,124], particularly throughout North America, Europe, and Asia. Average nocturnality of tetrapod assemblages tended to strongly increase after inclusion of imputed values, highlighting a global knowledge bias from lack of observed attribute data (Fig. 5H-I), particularly for squamates and to a lesser extent, amphibians (Figs S6-S11). We attribute this to both the low detectability of nocturnal species [125,126] and the general lack of data for tropical regions where nocturnal species predominate in the tropics [127]. Changes in average verticality were less pronounced across tetrapod assemblages (Fig. 5K-L), which is somewhat expected given the high completeness of microhabitat data. Even so, fossorial or arboreal species were responsible for 53% of imputed microhabitat whereas they represented about 44% of non-imputed microhabitat categories.

Although imputation biases can increase in the absence of additional correlated attributes [16], we found no relationships between the number of imputed attributes and changes in average attributes after imputation-based gap-filing (Fig. S12). That is, changes in average attributes

478 were not higher in genera and families including species with multiple missing attributes.
479 Moreover, changes in average attributes were either mostly unrelated with co-missingness
480 (Fig. S13) or occurred in opposite directions, with low co-missingness leading to greater
481 changes.
482
Broadly, we found that the magnitude of change in average attributes from imputation-based
gap-filling was associated with both completeness and richness (Figs S14-S16). Those taxa
and assemblages with minor completeness and with least species saw the strongest changes in
average attribute value. Lower attribute completeness creates more opportunities for variation
in attribute space after imputation-based gap-filing, whereas a high species richness can dilute
the importance of imputed values in the sample.
489

Tetrapod Traits – a comprehensive, imputation informed attributes database
We make the full phylogenetically coherent database, Tetrapod Traits, publicly available to
support a range of research applications in ecology, evolution, and conservation and to help
minimize the impacts of biased data in this system. Briefly, the Tetrapod Traits database
includes 75 fields registering observed and imputed species-level attributes followed by
respective sources across 33,281 tetrapod species. Attributes in this database show at least
98% of completeness after including imputed values and are organized in 9 major groups
(Table 1, see Table S1 for the complete description of each attribute):
498

1. **Taxonomy –** includes 8 attributes that inform scientific names and respective higher-
    level taxonomic ranks, authority name, and year of species description. Field names:
    Scientific.Name, Genus, Family, Suborder, Order, Class, Authority, and
    YearOfDescription.

2. **Tree –** includes 2 attributes that indicate which fully-sampled phylogeny contains the
    species, besides informing if the species phylogenetic relationship was imputed or not
    in the phylogeny. Field names: TreeTaxon, TreeImputed.

3. **Body size –** includes 7 attributes that inform length, mass, and data sources on species
    sizes, and details on the imputation of species length or mass. Field names:
    BodyLength_mm, LengthMeasure, ImputedLength, SourceBodyLength,
    BodyMass_g, ImputedMass, SourceBodyMass.

4. **Activity time –** includes 5 attributes that inform period of activity (e.g., diurnal,
    fossorial) as dummy variables, data sources, details on the imputation of species
    activity time, and a diurnality score. Field names: Diu, Noc, ImputedActTime,
    SourceActTime, Nocturnality.

5. **Microhabitat –** includes 8 attributes inform habitat use (e.g., fossorial, terrestrial,
    aquatic, arboreal, aerial) as dummy variables, data sources, details on the imputation
    of microhabitat, and a verticality score. Field names: Fos, Ter, Aqu, Arb, Aer,
    ImputedHabitat, SourceHabitat, Verticality.

6. **Macrohabitat –** includes 19 attributes that inform major habitat types according to
    the IUCN classification, the sum of major habitats, data source, and details on the
    imputation of macrohabitat. Field names: MajorHabitat_1 to MajorHabitat_10,
    MajorHabitat_12 to MajorHabitat_17, MajorHabitatSum, ImputedMajorHabitat,
    SourceMajorHabitat.

7. **Geography –** includes 12 attributes that inform species range size, average latitude
    and longitude, and the proportion of species range within each biogeographical realm.
    Field names: RangeSize, Longitude, Latitude, Afrotropic, Australasia, IndoMalay,
    Neartic, Neotropic, Oceania, Paleartic, Antartic.

8. **Environment –** includes 5 attributes that inform the average environmental condition within the species geographic range. Field names: AnnuMeanTemp, AnnuPrecip, TempSeasonality, PrecipSeasonality, Elevation.

9. **Human Influence –** includes 10 attributes that inform the within-range average or proportion of human-related activities, assessed statuses and sources. Field names: ETA50K, HumanDensity, PropAnthroHyde, PropUrbanArea, PropCroplandArea, PropPastureArea, PropRangelandArea, PropGrazingArea, AssessedStatus, SourceStatus.

Conclusion

The growth in mobilized attribute data over the last decade has enabled significant progress and increased geographic and taxonomic generality in comparative biology [82,128,129], macroevolution [130], macroecology [131], and conservation biology [132–134]. But research outcomes and their interpretation often remained constrained by non-randomly missing data [6,8,17], and lack of standardisation across data dimensions – trait, phylogenetic, spatial – and data sources [135]. In concert with taxonomic updates [136,137], these issues have imposed additional limits on multi-taxon investigations and global biodiversity synthesis [138–140]. Here, after documenting remaining global data gaps for central traits, we were able to use imputation to fill the gaps and illustrate the knowledge biases these missing attribute data had imposed. By using phylogeny-based imputation methods in concert with inter-related ecological attributes, we were able reveal the hitherto invisible part of the attributes space. Our findings confirmed the predominance of attributes co-missingness, and provide a comprehensive assessment of data gaps and biases across tetrapod groups.

The attributes predictions for species missing data brought to light the key correlates and identified the significant consequences these gaps can have for inference. For instance, larger body size has been related to both high species detectability [34,38,125] and discovery probability [141–143], larger research effort and public interest [40,41,144,145], increased availability of assessed status [146,147], and better data sampling. Despite the research difficulties that a large size may impose – e.g., high species sensitivity to disturbances [148,149], lower abundance [150,151], uneasy collection or transport – the presence of multiple data biases towards small-sized species confirm the struggle of biodiversity scientists to know well the smallest beings in nature. In summary, different knowledge shortfalls [139] can share the similar determinants.

Biodiversity knowledge shortfalls were also clearly related to activity time and microhabitat. Nocturnal species often show lower detectability [125], are typically less researched [40] and underassessed with respect to threat status [146], which helps explain why diurnal species are more likely to have activity time data. In addition, the high detectability of terrestrial than arboreal or fossorial species with standard sampling methods [38,152] can lead to less biased occurrence data [153] and more research effort [30,40]. Although the species foraging stratum may constrain certain aspects of our biodiversity knowledge, we find only a limited influence of verticality on microhabitat data availability.

Attribute imputations can be limited and themselves be biased [16,17]. For tetrapods and this study, these concerns were minimized by an already substantial initial data coverage and a gap-oriented mobilization of additional data. In only 276 tetrapod species (0.8%) all four focal attributes (body length, body mass, activity time, and microhabitat) had to be imputed. No family had the four focal attributes missing in all species, and only eight genera (0.15% of total genera) were missing the four focal attributes. Two important diagnostics offered strong support for the robustness of the attribute predictions. First, there was no association between the number of imputed attributes and changes in average attributes after imputation-based gap-filling (Fig. S12). Second, there was no positive association between co-missingness in

attribute data and changes in average attribute values (Fig. S13). We encourage researchers to carefully consider how assumptions and uncertainty of any imputed data might affect downstream use and interpretation, and the information we developed here is no exception. For many tetrapod research cases, however, we expect the inclusion of species with imputed data to be preferrable to the wholesale exclusion of the data-limited species.

We demonstrate that even for tetrapod species, a central model system in biodiversity science, our ecological knowledge attribute data are lacking consistently across certain types of traits, taxa, and regions. Through careful imputation we uncover that these gaps have brought about attribute distributions that have strong geographical and taxonomic biases. We expect the approach and likely the same general insights regarding biodiversity knowledge shortfalls to transcend taxa and systems. The illustrated consequence of biased data can percolate through trait-based metrics and comparative analyses, and ultimately have the potential to mislead research on functional diversity [8,19,154], extinction risk [6,21,155], and ecogeography [156–158]. Here we have provided a new phylogeny-based attribute databased that can help minimise the impact of biased data and support new research avenues in tetrapod conservation, ecology and evolution.

Data Availability

The Tetrapod Traits database and the R-code used for data validation will be made available on vertlife.org and GitHub upon acceptance for publication.

Acknowledgements

We genuinely thank the many authors of the numerous sources consulted during the preparation of this database. This work was only possible because of their valuable efforts. To São Paulo Research Foundation (FAPESP) for grants in support of MRM (#2021/11840-6 and #2022/12231-6), LFT (#2016/25358-3) and KC (#2020/12558-0). To Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for the fellowship to JJMG. To Conselho Nacional de Desenvolvimento Científico - CNPq for research grants in support of FPW (#311504/2020-5) and LFT (#302834/2020-6). WJ also acknowledges support from National Science Foundation (NSF) grant DEB-1441737 and NASA grants 80NSSC17K0282 and 80NSSC18K0435.

Author Contributions

Conceptualization: Mario R. Moura and Walter Jetz

Data Curation: Mario R. Moura, Karoline Ceron, Jhonny J. M. Guedes, Yanina V. Sica, Alessandro Catenazzi, Fernanda P. Werneck, Luís Felipe Toledo

Formal Analysis: Mario R. Moura

Funding Acquisition: Walter Jetz

Investigation: Mario R. Moura

Methodology: Mario R. Moura

Software: Mario R. Moura

Supervision: Walter Jetz

Visualization: Mario R. Moura and Walter Jetz

Writing – original draft preparation: Mario R. Moura

Writing – review & editing: Mario R. Moura, Karoline Ceron, Jhonny J. M. Guedes, Yanina V. Sica, Alessandro Catenazzi, Fernanda P. Werneck, Luís Felipe Toledo, and Walter Jetz

Competing interests

The authors declare no competing financial interests.

References

1. Beck J, Ballesteros-Mejia L, Buchmann CM, Dengler J, Fritz SA, Gruber B, et al. What's on the horizon for macroecology? Ecography (Cop). 2012;35: 673–683. doi:10.1111/j.1600-0587.2012.07364.x

2. Titley MA, Snaddon JL, Turner EC. Scientific research on animal biodiversity is systematically biased towards vertebrates and temperate regions. Schierwater B, editor. PLoS One. 2017;12: e0189577. doi:10.1371/journal.pone.0189577

3. Troudet J, Grandcolas P, Blin A, Vignes-Lebbe R, Legendre F. Taxonomic bias in biodiversity data and societal preferences. Sci Rep. 2017;7: 9132. doi:10.1038/s41598-017-09084-6

4. Etard A, Morrill S, Newbold T. Global gaps in trait data for terrestrial vertebrates. Sheard C, editor. Glob Ecol Biogeogr. 2020;29: 2143–2158. doi:10.1111/geb.13184

5. Oliver RY, Meyer C, Ranipeta A, Winner K, Jetz W. Global and national trends, gaps, and opportunities in documenting and monitoring species distributions. Moritz C, editor. PLOS Biol. 2021;19: e3001336. doi:10.1371/journal.pbio.3001336

6. González-Suárez M, Lucas PM, Revilla E. Biases in comparative analyses of extinction risk: mind the gap. J Anim Ecol. 2012;81: 1211–1222. doi:10.1111/j.1365-2656.2012.01999.x

7. Kim SW, Blomberg SP, Pandolfi JM. Transcending data gaps: a framework to reduce inferential errors in ecological analyses. Ecol Lett. 2018;21: 1200–1210. doi:10.1111/ele.13089

8. Pakeman RJ. Functional trait metrics are sensitive to the completeness of the species' trait data? Oksanen J, editor. Methods Ecol Evol. 2014;5: 9–15. doi:10.1111/2041-210X.12136

9. Nakagawa S, Freckleton RP. Missing inaction: the dangers of ignoring missing data. Trends Ecol Evol. 2008;23: 592–596. doi:10.1016/j.tree.2008.06.014

10. Sandel B, Gutiérrez AG, Reich PB, Schrodt F, Dickie J, Kattge J. Estimating the missing species bias in plant trait measurements. J Veg Sci. 2015;26: 828–838. doi:10.1111/jvs.12292

11. Taugourdeau S, Villerd J, Plantureux S, Huguenin□Elie O, Amiaud B. Filling the gap in functional trait databases: use of ecological hypotheses to replace missing data. Ecol Evol. 2014;4: 944–958. doi:10.1002/ece3.989

12. Penone C, Davidson AD, Shoemaker KT, Di Marco M, Rondinini C, Brooks TM, et al. Imputation of missing data in life-history trait datasets: which approach performs the best? Freckleton R, editor. Methods Ecol Evol. 2014;5: 961–970. doi:10.1111/2041-210X.12232

13. Stekhoven DJ, Buhlmann P. MissForest--non-parametric missing value imputation for mixed-type data. Bioinformatics. 2012;28: 112–118. doi:10.1093/bioinformatics/btr597

14. Bruggeman J, Heringa J, Brandt BW. PhyloPars: estimation of missing parameter values using phylogeny. Nucleic Acids Res. 2009;37: W179–W184. doi:10.1093/nar/gkp370

15. Goolsby EW, Bruggeman J, Ané C. Rphylopars□: fast multivariate phylogenetic comparative methods for missing data and within-species variation. Fitzjohn R, editor. Methods Ecol Evol. 2017;8: 22–27. doi:10.1111/2041-210X.12612

16. Jardim L, Bini LM, Diniz-Filho JAF, Villalobos F. A Cautionary Note on Phylogenetic Signal Estimation from Imputed Databases. Evol Biol. 2021;48: 246–258.

doi:10.1007/s11692-021-09534-0

17.   Johnson TF, Isaac NJB, Paviolo A, González␣Suárez M. Handling missing values in
trait data. Schrodt F, editor. Glob Ecol Biogeogr. 2021;30: 51–62.
doi:10.1111/geb.13185

18.   González-del-Pliego P, Freckleton RP, Edwards DP, Koo MS, Scheffers BR, Pyron
RA, et al. Phylogenetic and Trait-Based Prediction of Extinction Risk for Data-
Deficient Amphibians. Curr Biol. 2019;29: 1557-1563.e3.
doi:10.1016/j.cub.2019.04.005

19.   Carmona CP, Tamme R, Pärtel M, de Bello F, Brosse S, Capdevila P, et al. Erosion of
global functional diversity across the tree of life. Sci Adv. 2021;7.
doi:10.1126/sciadv.abf2675

20.   Colston TJ, Kulkarni P, Jetz W, Pyron RA. Phylogenetic and spatial distribution of
evolutionary diversification, isolation, and threat in turtles and crocodilians (non-avian
archosauromorphs). BMC Evol Biol. 2020;20: 81. doi:10.1186/s12862-020-01642-3

21.   Cooke RSC, Eigenbrod F, Bates AE. Projected losses of global mammal and bird
ecological strategies. Nat Commun. 2019;10: 2279. doi:10.1038/s41467-019-10284-z

22.   Soria CD, Pacifici M, Di Marco M, Stephen SM, Rondinini C. COMBINE: a
coalesced mammal database of intrinsic and extrinsic traits. Ecology. 2021;102:
e03344. doi:10.1002/ECY.3344/SUPPINFO

23.   Caetano GHO, Chapple DG, Grenyer R, Raz T, Rosenblatt J, Tingley R, et al.
Automated assessment reveals that the extinction risk of reptiles is widely
underestimated across space and phylogeny. Jordano P, editor. PLOS Biol. 2022;20:
e3001544. doi:10.1371/journal.pbio.3001544

24.   Jetz W, Pyron RA. The interplay of past diversification and evolutionary isolation with
present imperilment across the amphibian tree of life. Nat Ecol Evol. 2018;2: 850–858.
doi:10.1038/s41559-018-0515-5

25.   Tonini JFR, Beard KH, Ferreira RB, Jetz W, Pyron RA. Fully-sampled phylogenies of
squamates reveal evolutionary patterns in threat status. Biol Conserv. 2016;204: 23–
31. doi:10.1016/j.biocon.2016.03.039

26.   Jetz W, Thomas GH, Joy JB, Hartmann K, Mooers AO. The global diversity of birds
in space and time. Nature. 2012;491: 444–448. doi:10.1038/nature11631

27.   Upham NS, Esselstyn JA, Jetz W. Inferring the mammal tree: Species-level sets of
phylogenies for questions in ecology, evolution, and conservation. Tanentzap AJ,
editor. PLOS Biol. 2019;17: e3000494. doi:10.1371/journal.pbio.3000494

28.   Diniz-Filho JAF, Loyola RD, Raia P, Mooers AO, Bini LM. Darwinian shortfalls in
biodiversity conservation. Trends Ecol Evol. 2013;28: 689–695.
doi:10.1016/j.tree.2013.09.003

29.   Roll U, Feldman A, Novosolov M, Allison A, Bauer AM, Bernard R, et al. The global
distribution of tetrapods reveals a need for targeted reptile conservation. Nat Ecol
Evol. 2017;1: 1677–1682. doi:10.1038/s41559-017-0332-2

30.   Guedes JJM, Moura MR, Diniz␣Filho JAF. Species out of sight: elucidating the
determinants of research effort in global reptiles. Ecography (Cop). 2023; Early View.
doi:10.1111/ecog.06491

31.   Kéry M, Schmidt BR. Imperfect detection and its consequences for monitoring for
conservation. Community Ecol. 2008;9: 207–216. doi:10.1556/ComEc.9.2008.2.10

32.   Mazerolle MJ, Bailey LL, Kendall WL, Royle JA, Converse SJ, Nichols JD. Making
great leaps forward: Accounting for detectability in herpetological field studies. J
Herpetol. 2007;41: 672–689. doi:10.1670/07-061.1

33.   Jarzyna MA, Jetz W. Detecting the Multiple Facets of Biodiversity. Trends Ecol Evol.
2016;31: 527–538. doi:10.1016/j.tree.2016.04.002

34.   Anderson AS, Marques TA, Shoo LP, Williams SE. Detectability in Audio-Visual
Surveys of Tropical Rainforest Birds: The Influence of Species, Weather and Habitat
Characteristics. Ambrosini R, editor. PLoS One. 2015;10: e0128464.
doi:10.1371/journal.pone.0128464

35.   Einoder LD, Southwell DM, Lahoz-Monfort JJ, Gillespie GR, Fisher A, Wintle BA.

Occupancy and detectability modelling of vertebrates in northern Australia using multiple sampling methods. Crowther MS, editor. PLoS One. 2018;13: e0203304. doi:10.1371/journal.pone.0203304

36. Richardson E, Nimmo DG, Avitabile S, Tworkowski L, Watson SJ, Welbourne D, et al. Camera traps and pitfalls: an evaluation of two methods for surveying reptiles in a semiarid ecosystem. Wildl Res. 2017;44: 637. doi:10.1071/WR16048

37. Ribeiro-Júnior MA, Gardner TA, Ávila-Pires TC. Evaluating the Effectiveness of Herpetofaunal Sampling Techniques across a Gradient of Habitat Change in a Tropical Forest Landscape. J Herpetol. 2008;42: 733. doi:10.1670/07-097R3.1

38. Chutipong W, Lynam A, Steinmetz R, Savini T, Gale G. Sampling mammalian carnivores in western Thailand: Issues of rarity and detectability. Raffles Bull Zool. 2014;62: 521–535.

39. Moura MR, Costa HC, Peixoto MA, Carvalho ALG, Santana DJ, Vasconcelos HL. Geographical and socioeconomic determinants of species discovery trends in a biodiversity hotspot. Biol Conserv. 2018;220: 237–244. doi:10.1016/j.biocon.2018.01.024

40. Brooke ZM, Bielby J, Nambiar K, Carbone C. Correlates of Research Effort in Carnivores: Body Size, Range Size and Diet Matter. PLoS One. 2014;9: e93195. doi:10.1371/journal.pone.0093195

41. Guedes JJM, Moura MR, Alexandre F. Diniz□Filho J. Species out of sight: elucidating the determinants of research effort in global reptiles. Ecography (Cop). 2023;2023: e06491. doi:10.1111/ecog.06491

42. Wilman H, Belmaker J, Simpson J, de la Rosa C, Rivadeneira MM, Jetz W. EltonTraits 1.0: Species-level foraging attributes of the world's birds and mammals. Ecology. 2014;95: 2027–2027. doi:10.1890/13-1917.1

43. IUCN - International Union for Conservation of Nature. IUCN Red List of Threatened Species. In: Version 2021.3. 2021 p. www.iucnredlist.org.

44. Jetz W, McPherson JM, Guralnick RP. Integrating biodiversity distribution knowledge: toward a global map of life. Trends Ecol Evol. 2012;27: 151–159.

45. Marsh CJ, Sica Y V., Burgin CJ, Dorman WA, Anderson RC, Toro Mijares I, et al. Expert range maps of global mammal distributions harmonised to three taxonomic authorities. J Biogeogr. 2022; 1–14. doi:10.1111/jbi.14330

46. Hurlbert AH, Jetz W. Species richness, hotspots, and the scale dependence of range maps in ecology and conservation. Proc Natl Acad Sci. 2007;104: 13384–13389.

47. Hurlbert AH, White EP. Disparity between range map-and survey-based analyses of species richness: patterns, processes and implications. Ecol Lett. 2005;8: 319–327.

48. Jetz W, Sekercioglu CH, Watson JEM. Ecological correlates and conservation implications of overestimating species geographic ranges. Conserv Biol. 2008;22: 110–119. doi:10.1111/j.1523-1739.2007.00847.x

49. Karger DN, Conrad O, Böhner J, Kawohl T, Kreft H, Soria-Auza RW, et al. Climatologies at high resolution for the earth's land surface areas. Sci Data. 2017;4: 170122. doi:10.1038/sdata.2017.122

50. Amatulli G, Domisch S, Tuanmu M-N, Parmentier B, Ranipeta A, Malczyk J, et al. A suite of global, cross-scale topographic variables for environmental and biodiversity modeling. Sci Data. 2018;5: 180040. doi:10.1038/sdata.2018.40

51. Klein Goldewijk K, Beusen A, Doelman J, Stehfest E. Anthropogenic land use estimates for the Holocene-HYDE 3.2. Earth Syst Sci Data. 2017;9: 927–953. doi:10.5194/essd-9-927-2017

52. Nelson A. Estimated travel time to the nearest city of 50,000 or more people in year 2000. Ispra: Global Environment Monitoring Unit - Joint Research Centre of the European Commission; 2008. Available: http://forobs.jrc.ec.europa.eu/products/gam/

53. Scott Chamberlain. rredlist: "IUCN" Red List Client. 2020. p. R package version 0.7.0. Available: https://cran.r-project.org/package=rredlist

54. Jetz W, Thomas GH, Joy JB, Redding DW, Hartmann K, Mooers AO. Global Distribution and Conservation of Evolutionary Distinctness in Birds. Curr Biol.

806   2014;24: 919–930. doi:10.1016/j.cub.2014.03.011

807   55.   Anders G J Rhodin, Iverson JB, Van-Dijk PP, Buhlmann KA, Pritchard PCH,
808   Mittermeier RA. Turtles of the world: Annotated checklist and atlas of taxonomy,
809   synonymy, distribution, and conservation status (8th Ed.). New York: Chelonian
810   Research   Foundation;   2017.   Available:
811   https://books.google.com/books/about/Turtles_of_the_World_Annotated_Checklist.ht
812   ml?id=9PSOswEACAAJ

813   56.   Oaks JR. A time-calibrated species tree of Crocodylia reveals a recent radiation of the
814   true crocodiles. Evolution (N Y). 2011;65: 3285–3297. doi:10.1111/j.1558-
815   5646.2011.01373.x

816   57.   Meredith RW, Hekkala ER, Amato G, Gatesy J. A phylogenetic hypothesis for
817   *Crocodylus* (Crocodylia) based on mitochondrial DNA: Evidence for a trans-Atlantic
818   voyage from Africa to the New World. Mol Phylogenet Evol. 2011;60: 183–191.
819   doi:10.1016/j.ympev.2011.03.026

820   58.   Shirley MH, Carr AN, Nestler JH, Vliet KA, Brochu CA. Systematic revision of the
821   living African Slender-snouted Crocodiles (Mecistops Gray, 1844). Zootaxa.
822   2018;4504: 151. doi:10.11646/zootaxa.4504.2.1

823   59.   Franke FA, Schmidt F, Borgwardt C, Bernhard D, Bleidorn C, Engelmann W-E, et al.
824   Genetic differentiation of the African dwarf crocodile *Osteolaemus tetraspis* Cope,
825   1861 (Crocodylia: Crocodylidae) and consequences for European zoos. Org Divers
826   Evol. 2013;13: 255–266. doi:10.1007/s13127-012-0107-1

827   60.   Shirley MH, Villanova VL, Vliet KA, Austin JD. Genetic barcoding facilitates captive
828   and wild management of three cryptic African crocodile species complexes. Anim
829   Conserv. 2015;18: 322–330. doi:10.1111/acv.12176

830   61.   del Hoyo J, Elliott A, Sargatal J, Christie DA. Handbook of the Birds of the World.
831   Barcelona: Lynx Edicions;

832   62.   IUCN - International Union for Conservation of Nature. IUCN Red List of Threatened
833   Species. In: Version 2015.1. 2015 p. www.iucnredlist.org.

834   63.   Frost DR. Amphibian Species of the World: an Online Reference. Version 6.1. In:
835   American Museums of Natural History, New York, USA [Internet]. 2021 [cited 20 Jan
836   2022]. Available: https://amphibiansoftheworld.amnh.org/

837   64.   Uetz P, Freed P, Hosek J. The Reptile Database. 2021 [cited 20 Jan 2022]. Available:
838   http://www.reptile-database.org

839   65.   Lepage D, Vaidya G, Guralnick R. Avibase – a database system for managing and
840   organizing   taxonomic   concepts.   Zookeys.   2014;420:   117–135.
841   doi:10.3897/zookeys.420.7089

842   66.   Burgin CJ, Colella JP, Kahn PL, Upham NS. How many species of mammals are
843   there? J Mammal. 2018;99: 1–14. doi:10.1093/jmammal/gyx147

844   67.   Meiri S, Bauer AM, Allison A, Castro-Herrera F, Chirio L, Colli G, et al. Extinct,
845   obscure or imaginary: The lizard species with the smallest ranges. Stevens G, editor.
846   Divers Distrib. 2018;24: 262–273. doi:10.1111/ddi.12678

847   68.   Martin TE, Bennett GC, Fairbairn A, Mooers AO. 'Lost' taxa and their conservation
848   implications. Anim Conserv. 2022;Early View: 1–11. doi:10.1111/acv.12788

849   69.   Tobias JA, Sheard C, Pigot AL, Devenish AJM, Yang J, Sayol F, et al. AVONET:
850   morphological, ecological and geographical data for all birds. Ecol Lett. 2022;25: 581–
851   597. doi:10.1111/ELE.13898

852   70.   Paradis E, Claude J, Strimmer K. APE: Analyses of Phylogenetics and Evolution in R
853   language. Bioinformatics. 2004;20: 289–290. doi:10.1093/bioinformatics/btg412

854   71.   Warnes GR, Bolker B, Lumley T, Johnson RC. gmodels: Various R Programming
855   Tools for Model Fitting. 2018. p. R package version 2.18.1. Available: https://cran.r-
856   project.org/package=gmodels

857   72.   Oliveira BF, São-Pedro VA, Santos-Barrera G, Penone C, Costa GC. AmphiBIO, a
858   global database for amphibian ecological traits. Sci Data. 2017;4: 170123.
859   doi:10.1038/sdata.2017.123

860   73.   Zamudio KR, Bell RC, Nali RC, Haddad CFB, Prado CPA. Polyandry, Predation, and

the Evolution of Frog Reproductive Modes. Am Nat. 2016;188: S41–S61. doi:10.1086/687547

74.   Manamendra-Arachchi K, Pethiyagoda R. The Sri Lankan shrub-frogs of the genus Philautus Gistel, 1848 (Ranidae: Rhacophorinae), with description of 27 new species. Raffles Bull Zool. 2005;12: 163–303.

75.   Liang F, Shuqin H, Changyuan Y, Yongzhao H. Fauna Sinica: Amphibia（Vol. 3）. Beijing: Science Press; 2009.

76.   Channing A, Ro□del M-O. Field guide to the frogs & other amphibians of Africa. Cape Town: Struik Nature; 2019.

77.   Feldman A, Sabath N, Pyron RA, Mayrose I, Meiri S. Body sizes and diversification rates of lizards, snakes, amphisbaenians and the tuatara. Glob Ecol Biogeogr. 2016;25: 187–197. doi:10.1111/geb.12398

78.   Myhrvold NP, Baldridge E, Chan B, Sivam D, Freeman DL, Ernest SKM. An amniote life history database to perform comparative analyses with birds, mammals, and reptiles. Dryad Digit Repos. 2014;96: 3109. doi:10.5061/dryad.t6m96

79.   Meyer de Schauensee R. The birds of China. Washington, DC: Smithsonian Institution Press; 1984.

80.   Hutchins M, Jackson JA, Bock WJ, Olendorf D. Grzimek's Animal Life Encyclopedia, v. 8-11: Birds. 2nd editio. Farmington Hills: Gale Group; 2003.

81.   Santini L, Benítez-López A, Ficetola GF, Huijbregts MAJ. Length–mass allometries in amphibians. Integr Zool. 2018;13: 36–45. doi:10.1111/1749-4877.12268

82.   Stark G, Pincheira-Donoso D, Meiri S. No evidence for the 'rate-of-living' theory across the tetrapod tree of life. Glob Ecol Biogeogr. 2020;29: 857–884. doi:10.1111/GEB.13069

83.   Stark G, Pincheira-Donoso D, Meiri S. CORRIGENDUM: No evidence for the 'rate□of□living' theory across the tetrapod tree of life. Glob Ecol Biogeogr. 2021;30: 2514–2514. doi:10.1111/geb.13400

84.   Jacobsen NHG. A Herpetological survey of the Transvaal. University of Natal. 1989. Available: https://researchspace.ukzn.ac.za/xmlui/handle/10413/8461?show=full

85.   Faurby S, Davis M, Pedersen RØ, Schowanek SD, Antonelli1 A, Svenning J. PHYLACINE 1.2: The Phylogenetic Atlas of Mammal Macroecology. Ecology. 2018;99: 2626–2626. doi:10.1002/ecy.2443

86.   Faurby S, Svenning J-C. Resurrection of the Island Rule: Human-Driven Extinctions Have Obscured a Basic Evolutionary Pattern. Am Nat. 2016;187: 812–820. doi:10.1086/686268

87.   Jones KE, Bielby J, Cardillo M, Fritz SA, O'Dell J, Orme CDL, et al. PanTHERIA: a species-level database of life history, ecology, and geography of extant and recently extinct mammals. Michener WK, editor. Ecology. 2009;90: 2648–2648. doi:10.1890/08-1494.1

88.   Kingdon J, Happold D, Butynski T, Hoffmann M, Happold M, Kalina J. Mammals of Africa v. 1-6. 1st editio. London: Bloomsbury Publishing; 2013.

89.   Haddad CFB, Toledo LF, Prado CPA, Loebmann D, Gasparini JL, Sazima I. Guide to the amphibians of the Atlantic Forest: diversity and biology. São Paulo: Anolis Book; 2013.

90.   Stark G, Meiri S. Cold and dark captivity: Drivers of amphibian longevity. Glob Ecol Biogeogr. 2018;27: 1384–1397. doi:10.1111/geb.12804

91.   Stark G, Meiri S. CORRIGENDUM: Cold and dark captivity: Drivers of amphibian longevity. Glob Ecol Biogeogr. 2021;30: 2513–2513. doi:10.1111/geb.13399

92.   Thomas KN, Gower DJ, Bell RC, Fujita MK, Schott RK, Streicher JW. Eye size and investment in frogs and toads correlate with adult habitat, activity pattern and breeding ecology. Proc R Soc B Biol Sci. 2020;287: 20201393. doi:10.1098/rspb.2020.1393

93.   Neves MO, Cabral H, Pedrozo M, Ferreira VL, Moura MR, Santana DJ. Dataset of occurrences and ecological traits of amphibians from Upper Paraguay River Basin, central South America. Nat Conserv. 2020;41: 71–89. doi:10.3897/natureconservation.41.54265

94.   Lovich JE, Gibbons W. Turtles of the World: A guide to every family. Princeton:
Princeton University Press; 2021.
95.   Meiri S. Traits of lizards of the world: Variation around a successful evolutionary
design. Glob Ecol Biogeogr. 2018;27: 1168–1172. doi:10.1111/geb.12773
96.   Harrington SM, de Haan JM, Shapiro L, Ruane S. Habits and characteristics of
arboreal snakes worldwide: arboreality constrains body size but does not affect lineage
diversification. Biol J Linn Soc. 2018;125: 61–71. doi:10.1093/biolinnean/bly097
97.   O'Shea M. The Book of Snakes: A life-size guide to six hundred species from around
the world. Chicago: University of Chicago Press; 2018.
98.   Pietersen D, Verburgt L, Davies J. Snakes and other reptiles of Zambia and Malawi.
Cape Town: Struik Nature; 2021.
99.   Scharf I, Feldman A, Novosolov M, Pincheira-Donoso D, Das I, Böhm M, et al. Late
bloomers and baby boomers: ecological drivers of longevity in squamates and the
tuatara. Glob Ecol Biogeogr. 2015;24: 396–405. doi:10.1111/geb.12244
100. Feldman A, Meiri S. Australian snakes do not follow Bergmann's rule. Evol Biol.
2014;41: 327–335. doi:10.1007/s11692-014-9271-x
101. Guedes TB, Nogueira C, Marques OAV. Diversity, natural history, and geographic
distribution of snakes in the Caatinga, Northeastern Brazil. Zootaxa. 2014;3863: 1–93.
doi:10.11646/zootaxa.3863.1.1
102. Fraga R, Lima AP, Prudente ALC, Magnusson WE. Guide to the Snakes of the
Manaus region - Central Amazonia. Manaus: Editora INPA; 2013.
103. Marques OA V, Eterovic A, Sazima I. Snakes of the Brazilian Atlantic Forest: An
illustrated field guide for the Serra do Mar range. Ribeirão Preto: Holos; 2004.
104. Marques OA V, Eterovic A, Nogueira CC, Sazima I. Serpentes do Cerrado: Guia
ilustrado. Ribeirão Preto: Holos; 2015.
105. Chippaux J-P, Jackson K. Snakes of Central and Western Africa. Baltimore Maryland:
Johns Hopkins University Press; 2019.
106. Vogel G. Venomous snakes of Asia. Frankfurt: Edition Chimaira; 2006.
107. Munstermann MJ, Heim NA, McCauley DJ, Payne JL, Upham NS, Wang SC, et al. A
global ecological signal of extinction risk in terrestrial vertebrates. Conserv Biol. 2021;
1–13. doi:10.1111/cobi.13852
108. Bars-Closel M, Kohlsdorf T, Moen DS, Wiens JJ. Diversification rates are more
strongly related to microhabitat than climate in squamate reptiles (lizards and snakes).
Evolution (N Y). 2017;71: 2243–2261. doi:10.1111/evo.13305
109. Branch B. Photographic guide to snakes, other reptiles and amphibians of East Africa.
Cape Town: Random House Struik; 2014.
110. Branch B. A photographic guide to snakes and other reptiles of Southern Africa. Struik
Nature; 2002.
111. Figueroa A. Phylogenetic Relationships and Evolution of Snakes. University of New
Orleans. 2016. Available: https://scholarworks.uno.edu/td/2222/
112. Campbell JA, Lamar WW. The venomous reptiles of the Western Hemisphere - 2
volumes. Ithaca: Comstock Pub. Associates; 2004.
113. IUCN - International Union for Conservation of Nature. IUCN Red List of Threatened
Species. In: Version 2016.3. 2016 p. www.iucnredlist.org.
114. IUCN - International Union for Conservation of Nature. IUCN Red List of Threatened
Species. In: Version 2018.2. 2018 p. www.iucnredlist.org.
115. IUCN - International Union for Conservation of Nature. IUCN Red List of Threatened
Species. In: Version 2019.3. 2020 p. www.iucnredlist.org.
116. IUCN - International Union for Conservation of Nature. IUCN Red List of Threatened
Species. In: Version 2019.2. 2019 p. www.iucnredlist.org.
117. Stone L, Roberts A. The checkerboard score and species distributions. Oecologia.
1990;85: 74–79. doi:10.1007/BF00317345
118. Gotelli NJ. Null model analysis of species co-occurrence patterns. Ecology. 2000;81:
2606–2621. doi:10.1890/0012-9658(2000)081[2606:NMAOSC]2.0.CO;2
119. Ulrich W, Gotelli NJ. Pattern detection in null model analysis. Oikos. 2013;122: 2–18.

971   doi:10.1111/j.1600-0706.2012.20325.x

120. Gotelli NJ, Hart E, Ellison A. Null Model Analysis for Ecological Data. R package version 0.1.0. 2015. pp. 1–53.

121. Castiglione S, Tesone G, Piccolo M, Melchionna M, Mondanaro A, Serio C, et al. A new method for testing evolutionary rate variation and shifts in phenotypic evolution. Cooper N, editor. Methods Ecol Evol. 2018;9: 974–983. doi:10.1111/2041-210X.12954

122. Scheffers BR, Yong DL, Harris JBC, Giam X, Sodhi NS. The World's Rediscovered Species: Back from the Brink? Gratwicke B, editor. PLoS One. 2011;6: e22531. doi:10.1371/journal.pone.0022531

123. Ladle R, Jepson P, Malhado ACM, Jennings S, Barua M. The causes and biogeographical significance of species' rediscovery. Front Biogeogr. 2012;3: 111–118. doi:10.21425/F5FBG12432

124. Meyer C, Kreft H, Guralnick R, Jetz W. Global priorities for an effective information basis of biodiversity distributions. Nat Commun. 2015;6: 8221. doi:10.1038/ncomms9221

125. Einoder LD, Southwell DM, Lahoz-Monfort JJ, Gillespie GR, Fisher A, Wintle BA. Occupancy and detectability modelling of vertebrates in northern Australia using multiple sampling methods. Crowther MS, editor. PLoS One. 2018;13: e0203304. doi:10.1371/journal.pone.0203304

126. Richardson E, Nimmo DG, Avitabile S, Tworkowski L, Watson SJ, Welbourne D, et al. Camera traps and pitfalls: an evaluation of two methods for surveying reptiles in a semiarid ecosystem. Wildl Res. 2017;44: 637. doi:10.1071/WR16048

127. Vidan E, Roll U, Bauer A, Grismer L, Guo P, Maza E, et al. The Eurasian hot nightlife: Environmental forces associated with nocturnality in lizards. Glob Ecol Biogeogr. 2017;26: 1316–1325. doi:10.1111/geb.12643

128. Atwood TB, Valentine SA, Hammill E, McCauley DJ, Madin EMP, Beard KH, et al. Herbivores at the highest risk of extinction among mammals, birds, and reptiles. Sci Adv. 2020;6. doi:10.1126/sciadv.abb8458

129. Anderson SR, Wiens JJ. Out of the dark: 350 million years of conservatism and evolution in diel activity patterns in vertebrates. Evolution (N Y). 2017; 1–16. doi:10.1111/evo.13284

130. Pigot AL, Sheard C, Miller ET, Bregman TP, Freeman BG, Roll U, et al. Macroevolutionary convergence connects morphological form to ecological function in birds. Nat Ecol Evol. 2020;4: 230–239. doi:10.1038/s41559-019-1070-4

131. Violle C, Reich PB, Pacala SW, Enquist BJ, Kattge J. The emergence and promise of functional biogeography. Proc Natl Acad Sci. 2014;111: 13690–13696. doi:10.1073/pnas.1415442111

132. Cardillo M, Mace GM, Gittleman JL, Jones KE, Bielby J, Purvis A. The predictability of extinction: biological and external correlates of decline in mammals. Proc R Soc B Biol Sci. 2008;275: 1441–1448. doi:10.1098/rspb.2008.0179

133. Pollock LJ, Thuiller W, Jetz W. Large conservation gains possible for global biodiversity facets. Nature. 2017;546: 141–144. doi:10.1038/nature22368

134. Lee TM, Jetz W. Unravelling the structure of species extinction risk for predictive conservation science. Proc R Soc B Biol Sci. 2011;278: 1329–1338. doi:10.1098/rspb.2010.1877

135. Schneider FD, Fichtmueller D, Gossner MM, Güntsch A, Jochum M, König‐Ries B, et al. Towards an ecological trait‐data standard. Orme D, editor. Methods Ecol Evol. 2019;10: 2006–2019. doi:10.1111/2041-210X.13288

136. Bremer K, Bremer B, Karis P, Källersjö M. Time for change in taxonomy. Nature. 1990;343: 202–202. doi:10.1038/343202a0

137. Garnett ST, Christidis L. Taxonomy anarchy hampers conservation. Nature. 2017;546: 25–27. doi:10.1038/546025a

138. Wüest RO, Zimmermann NE, Zurell D, Alexander JM, Fritz SA, Hof C, et al. Macroecology in the age of Big Data – Where to go from here? J Biogeogr. 2019;

1026   jbi.13633. doi:10.1111/jbi.13633

139. Hortal J, de Bello F, Diniz-Filho JAF, Lewinsohn TM, Lobo JM, Ladle RJ. Seven Shortfalls that Beset Large-Scale Knowledge of Biodiversity. Annu Rev Ecol Evol Syst. 2015;46: annurev-ecolsys-112414-054400. doi:10.1146/annurev-ecolsys-112414-054400

140. Heberling JM, Miller JT, Noesgaard D, Weingart SB, Schigel D. Data integration enables global biodiversity synthesis. Proc Natl Acad Sci. 2021;118. doi:10.1073/pnas.2018093118

141. Moura MR, Jetz W. Shortfalls and opportunities in terrestrial vertebrate species discovery. Nat Ecol Evol. 2021;5: 631–639. doi:10.1038/s41559-021-01411-5

142. Diniz-Filho JAF, Bastos RP, Rangel TFLVB, Bini LM, Carvalho P, Silva RJ. Macroecological correlates and spatial patterns of anuran description dates in the Brazilian Cerrado. Glob Ecol Biogeogr. 2005;14: 469–477. doi:10.1111/j.1466-822x.2005.00165.x

143. Meiri S. Small, rare and trendy: traits and biogeography of lizards described in the 21st century. J Zool. 2016;299: 251–261. doi:10.1111/jzo.12356

144. Roll U, Mittermeier JC, Diaz GI, Novosolov M, Feldman A, Itescu Y, et al. Using Wikipedia page views to explore the cultural importance of global reptiles. Biol Conserv. 2016;204: 42–50. doi:10.1016/j.biocon.2016.03.037

145. Tyler EHM, Somerfield PJ, Berghe E Vanden, Bremner J, Jackson E, Langmead O, et al. Extensive gaps and biases in our knowledge of a well-known fauna: implications for integrating biological traits into macroecology. Glob Ecol Biogeogr. 2012;21: 922–934. doi:10.1111/j.1466-8238.2011.00726.x

146. Meiri S, Chapple DG. Biases in the current knowledge of threat status in lizards, and bridging the 'assessment gap.' Biol Conserv. 2016;204: 6–15. doi:10.1016/j.biocon.2016.03.009

147. Miqueleiz I, Bohm M, Ariño AH, Miranda R. Assessment gaps and biases in knowledge of conservation status of fishes. Aquat Conserv Mar Freshw Ecosyst. 2020;30: 225–236. doi:10.1002/aqc.3282

148. Blumstein DT. Developing an evolutionary ecology of fear: how life history and natural history traits affect disturbance tolerance in birds. Anim Behav. 2006;71: 389–399. doi:10.1016/j.anbehav.2005.05.010

149. Henle K, Davies KF, Kleyer M, Margules C, Settele J. Predictors of Species Sensitivity to Fragmentation. Biodivers Conserv. 2004;13: 207–251. doi:10.1023/B:BIOC.0000004319.91643.9e

150. Robinson JG, Redford KH. Body Size, Diet, and Population Density of Neotropical Forest Mammals. Am Nat. 1986;128: 665–680. doi:10.1086/284596

151. White EP, Ernest SKM, Kerkhoff AJ, Enquist BJ. Relationships between body size and abundance in ecology. Trends Ecol Evol. 2007;22: 323–330. doi:10.1016/j.tree.2007.03.007

152. Ribeiro-Júnior MA, Gardner TA, Ávila-Pires TC. Evaluating the Effectiveness of Herpetofaunal Sampling Techniques across a Gradient of Habitat Change in a Tropical Forest Landscape. J Herpetol. 2008;42: 733. doi:10.1670/07-097R3.1

153. Meyer C, Jetz W, Guralnick RP, Fritz SA, Kreft H. Range geometry and socio-economics dominate species-level biases in occurrence information. Glob Ecol Biogeogr. 2016. doi:10.1111/geb.12483

154. Cooke RSC, Bates AE, Eigenbrod F. Global trade-offs of functional redundancy and functional dispersion for birds and mammals. Schrodt F, editor. Glob Ecol Biogeogr. 2019;28: 484–495. doi:10.1111/geb.12869

155. Vilela B, Villalobos F, Rodríguez MÁ, Terribile LC. Body Size, Extinction Risk and Knowledge Bias in New World Snakes. Lambertucci SA, editor. PLoS One. 2014;9: e113429. doi:10.1371/journal.pone.0113429

156. Vidan E, Novosolov M, Bauer AM, Herrera FC, Chirio L, Campos CN, et al. The global biogeography of lizard functional groups. J Biogeogr. 2019;46: 2147–2158. doi:10.1111/jbi.13667

157. Oliveira BF, Scheffers BR. Vertical stratification influences global patterns of biodiversity. Ecography (Cop). 2019;42: 249–249. doi:10.1111/ecog.03636

158. Gaston KJ, Chown SL, Evans KL. Ecogeographical rules: elements of a synthesis. J Biogeogr. 2008;35: 483–500. doi:10.1111/j.1365-2699.2007.01772.x

Figure Legends

Figure 1. Attribute data coverage (completeness) of tetrapod species by family. Colour rings show the proportion of species with observed attributes in each tetrapod family for the five attributes with lowest completeness. Phylogenetic relationships shown are simple the family-level topology of tetrapod families.

Figure 2. Average data completeness across tetrapod assemblages. Proportion of species with observed values for: (A) body length, (B) body mass, (D) activity time, (E) microhabitat, (G) assessed threat status, (H) phylogenetic data, (J) attribute set, i.e., the average pattern for maps depicted in ABDE, and (K) the complete database, i.e., average pattern for maps depicted in ABDEGH. Maps show grid cell assemblages of 110×110 km size in an equal-area projection, and latitudinal plots C, F, I and L illustrate the average of these cells value across per latitude. See Fig. S4 for spatial patterns of attribute completeness for each tetrapod group. Colour ramps followed Jenks natural breaks classification.

Figure 3. The top 50 tetrapod genera with most pronounced patterns of shared data missingness. For each genus, the aggregation metric equals the median value of the standardised effect-size (SES) of the C-score metric computed across (A) all pairwise attribute combinations; or (B) attribute pairs mandatorily involving threat status. Gray numbers of the left side of each panel indicate the per-genus species richness. See Fig. S5 for aggregation patterns computed at the family-level.

Figure 4. Changes in family-level attributes average after imputation-based gap-filling. For continuous data (BL, body length and BM, body mass), relative change in attribute property quantify the proportional decrease or increase in the geometric mean attribute value per-genus. For binary attributes (types of activity time and microhabitat), relative change in attribute property measured the decrease or increase in mean species nocturnality (Noct) and verticality (Vert) score. The inset donut chart denotes the proportion of genera (inner donut) and families (outer donut) by category of attribute change. See Fig. S11 for individual charts of each tetrapod group.

Figure 5. Changes in average attribute per tetrapod assemblage after imputation-based gap-filing. For each grid cell, maps show the average species attribute value and respective relative change in attribute value after inclusion of imputed values for (A─C) body length, (D─F) body mass, (G─I) nocturnality, (J─L) verticality. Body length and body mass were $\log_{10}$ transformed before computations. Gray cells indicate assemblages without species with imputed values. See Figs S6-S10 for patterns of attribute change for each tetrapod group.