A phylogeny-informed characterisation of global tetrapod traits addresses data gaps and biases

Mario R. Moura[1,2,3,4*], Karoline Ceron[1], Jhonny J. M. Guedes[5], Rosana Chen-Zhao[1], Yanina V. Sica[3,4], Julie Hart[4,6], Wendy Dorman[4,7], Julia M. Portmann[3,4], Pamela Gonzalez-del-Pliego[4,8], Ajay Ranipeta[4], Alessandro Catenazzi[9], Fernanda P. Werneck[10], Luís Felipe Toledo[1], Nathan S. Upham[11], João F. R. Tonini[12], Timothy J. Colston[13], Robert Guralnick[14], Rauri C. K. Bowie[15], R. Alexander Pyron[16], and Walter Jetz[3,4*]

1. Departamento de Biologia Animal, Universidade Estadual de Campinas, Campinas, SP 13083-862, Brazil
2. Departamento de Biociências, Universidade Federal da Paraíba, Areia, PB 58397-000, Brazil
3. Department of Ecology and Evolutionary Biology, Yale University, New Haven CT, 06511, USA
4. Center for Biodiversity and Global Change, Yale University, New Haven CT, 06511, USA
5. Programa de Pós-Graduação em Ecologia e Evolução, Departamento de Ecologia, Universidade Federal de Goiás, Goiânia, GO 74690-900, Brazil
6. New York Natural Heritage Program, State University of New York College of Environmental Science and Forestry, Albany, NY 12233, USA
7. Department of Natural Resources and Environmental Sciences (NRES), University of Illinois Urbana-Champaign, Urbana, IL 61801, USA
8. Rui Nabeiro Biodiversity Chair, MED Institute, Universidade de Évora, 7000 Évora, Portugal
9. Department of Biological Sciences, Florida International University, Miami FL 33199, USA
10. Programa de Coleções Científicas Biológicas, Coordenação de Biodiversidade, Instituto Nacional de Pesquisas da Amazônia, Manaus, AM 69067-375, Brazil
11. School of Life Sciences, Arizona State University, Tempe, AZ 85284, USA
12. Department of Biology, University of Richmond, Richmond VA, 23173, USA
13. Biology Department, University of Puerto Rico at Mayagüez, Mayagüez, Puerto Rico, 00681
14. Florida Museum of Natural History, University of Florida, Gainesville, FL 32611, USA
15. Museum of Vertebrate Zoology and Department of Integrative Biology, University of California, Berkeley, Berkeley CA 9470, USA
16. Department of Biological Sciences, The George Washington University, Washington DC, 20052, USA

*Corresponding authors: Mario R. Moura (mariormoura@gmail.com), Walter Jetz (walter.jetz@yale.edu)

Abstract

Tetrapods (amphibian, reptiles, birds and mammals) are model systems for global biodiversity science, but continuing data gaps, limited data standardisation, and ongoing flux in taxonomic nomenclature constrain integrative research on this group and potentially cause biased inference. We combined and harmonised taxonomic, spatial, phylogenetic, and attribute data with phylogeny-based multiple imputation to provide a comprehensive data resource (TetrapodTraits 1.0.0) that includes values, predictions, and sources for body size, activity time, micro- and macrohabitat, ecosystem, threat status, biogeography, insularity, environmental preferences and human influence, for all 33,281 tetrapod species covered in recent fully sampled phylogenies. We assess gaps and biases across taxa and space, finding that shared data missing in attribute values increased with taxon-level completeness and richness across clades. Prediction of missing attribute values using multiple imputation revealed substantial changes in estimated macroecological patterns. These results highlight

biases incurred by non-random missingness and strategies to best address them. While there is an obvious need for further data collection and updates, our phylogeny- -informed database of tetrapod traits can support a more comprehensive representation of tetrapod species and their attributes in ecology, evolution, and conservation research.

Introduction

Over the past two decades biodiversity science has seen a dramatic growth in large-scale research in ecology, evolution, and conservation biology, enabled by near-global coverage for study systems such as terrestrial vertebrates, or Tetrapoda (amphibians, reptiles, birds, and mammals). These efforts usually rely on datasets spanning wide temporal, spatial, and taxonomic scales [1] that ideally are fully harmonised and well curated. Despite terrestrial vertebrates being a relatively well-known animal group when compared to invertebrates and plants [2,3], notable gaps persist across various attributes, including fundamental aspects of species natural history [4,5]. Consequently, trait-based research on biodiversity is often hampered by spatially and phylogenetically incomplete datasets [6–8].

Understanding causes of missingness is instrumental to advancing biodiversity data coverage. For any species attribute, there are observed and unobserved entries, each with a probability of being missing [9]. When all entries, whether observed or unobserved, share the same likelihood of being missing, data are said to be missing completely at random (MCAR). If missingness affects only observed entries, the data is termed missing at random (MAR). For example, a depleted digital scale battery makes weighing subsequent specimens impossible in the field, resulting in MAR data. Data is considered missing not at random (MNAR) when missingness is tied to unobserved entries, indicating a link to the missing values themselves. These three missing mechanisms – MCAR, MAR, MNAR – can lead to different configurations of the invisible fraction of the trait space [9–11].

While some missing mechanisms (e.g., MCAR, MAR) primarily affect a single attribute, the underlying cause behind an attribute MNAR can influence multiple variables, resulting in co-missingness or shared gaps. For instance, a species might lack information on multiple ecological aspects due to being known from only a few specimens with no details on where, when, and how they were found [12]. Similar circumstances apply to rare species or those collected solely through passive sampling techniques (e.g., pitfall traps), leaving ecological data unobserved. Indeed, bias is recognised in the availability of trait data for certain taxa, regions, and traits [4], and missing values for a given variable may be associated with incompleteness in others. Such congruent or aggregated (as opposed to segregated) patterns in trait missingness can arise from societal and research preferences for charismatic species [3], easily sampled taxonomic groups, or accessible geographical regions [6,13]. Conversely, segregated patterns may reveal traits and taxa that are challenging to sample or under-represented.

Despite the continued limitation of sampled data for many attributes in tetrapods, new methods can help to minimise these gaps and improve our understanding of biodiversity. Past practices have included the removal of species with missing data or the replacement of missing values by observed averages, but these strategies may ultimately reduce statistical power and increase bias [7,10,14]. More recently, growth in large-scale phylogenetic analyses has boosted the development of methods to increase the accuracy of imputing missing values [15–18]. Among tetrapods, recent large-scale applications of imputation methods include the use of phylogenetic regression methods and machine-learning techniques to predict missing values in trait data for amphibians [19], reptiles [19,20], birds [19,21], and mammals [19,21,22], as well as to inform threat statuses for data deficient and non-assessed species [20,23,24].

We leveraged a fast and automated multiple imputation technique with additional data mobilisation to provide a comprehensive database and assessment of key ecological attributes

of all extant 33,281 tetrapod species covered in recent fully-sampled phylogenies, including 7,238 amphibians [25], 384 chelonians and crocodilians [20], 9,755 squamates and tuatara [26], 9,993 birds [27], and 5,911 mammals [28]. Our assessment covered standardised species-level attributes for taxonomy, body size, activity time, microhabitat, macrohabitat, ecosystem, threat status, biogeography, insularity, environmental preferences, and human influence. Since not all species have genetic data (representing an important source of the remaining uncertainty about their placements in available phylogenies), we also evaluated completeness in genetic sequences [29]. We pinpointed taxa exhibiting pronounced shared missingness in natural history data to inform new strategies for data acquisition and mitigate biases in trait databases. To enhance database consistency, we taxonomically harmonised data sources and filled gaps using a phylogeny-based multiple imputation method [30–32] for which we verified the performance and associated uncertainty.

We use this gap-filled database to assess the geographic, taxonomic, and trait-related biases and evaluate how their model-based closure supports improved information and biological inference. Due to the biodiversity knowledge paradox – high biodiversity in the tropics [33] but better taxonomic sampling in temperate regions [4,5,34] – we expect larger unsampled fractions in the tetrapod trait space for tropical species. Similarly, the high research capacity (i.e., infrastructure and expertise availability) dedicated to birds and mammals relative to amphibians and reptiles [2,3] contributes to the uneven sampling of trait space [4], likely producing larger biases among historically undersampled taxa. Finally, species biology and sampling methodologies are known to affect detection and collection rates in the field [35–37]. For instance, detectability is typically lower for small- than large-bodied species, and similarly so for nocturnal relative to diurnal taxa [38–40], whereas sampling methodologies often favour the collection and research of species living on the surface compared to fossorial or arboreal groups [34,41–44]. We thus anticipate convergent missingness across trait space in tetrapods, and expect undersampled species to typically be small, nocturnal, and fossorial or arboreal.

Methods

We curated and assembled available databases for global tetrapod groups and used the latest phylogeny-based methods to create the most comprehensive tetrapod attribute dataset to date. While the TetrapodTraits database also covers a wide range of attributes derived from species range maps (see Table S1, Supporting Information), our focus regarding the imputations primarily centred on natural history traits, specifically: body length, body mass, activity time, and microhabitat. Our procedures can be summarised in five general steps: (i) data acquisition, (ii) taxonomic harmonisation, (iii) outlier verification, (iv) taxonomic imputation, and (v) phylogenetic multiple imputation.

Data Acquisition

We compiled species-level attributes regarding taxonomy, body size, activity time, microhabitat, macrohabitat, ecosystem, threat status, biogeography, insularity, environmental preferences, and associated data sources for each tetrapod species (Table S1). We gathered information from several global, continental, and regional databases, and complemented the existing data from published (articles, book chapters, field guides, and monographs) and grey literature (e.g., technical reports, government documents, monographs, theses). We also incorporated unpublished data gathered during fieldwork performed by some of us. To minimise the uneven representation of ecological attributes across clades, we initially identified genera and families whose species did not have available data on body size, activity time, or microhabitat. We then used species belonging to these genera and families to carry out additional online searches on academic platforms (Google Scholar and Web of Science) and included complementary attribute data whenever possible. To improve the chances of finding relevant natural history data [45,46], we conducted these searches using natural history terms in English (e.g., activity time, microhabitat, body size, length, mass, weight),

Portuguese (e.g., tempo de atividade, micro-habitat, tamanho de corpo, comprimento, massa, peso), and Spanish (e.g., tiempo de actividad, microhabitat, tamaño del cuerpo, longitud, masa, peso) along with the respective species scientific name or unique synonyms (see Taxonomic Harmonisation section). When sources were available in other languages, we employed translation tools for inspection (e.g., Google Translate). In our examination of the data sources, we did not use trait values provided solely at the genus level (e.g., mean value per genus).

Briefly, taxonomic data were represented by higher-level taxonomic ranks (Class, Order, Family), scientific name (same spelling as used in recent fully-sampled phylogenies), authority name, and year of description. Three broad natural history traits – body size, activity time, and microhabitat – have been compiled and harmonised across different tetrapod groups. Body size data consisted of information on body length (mm) and body mass (g). Activity time encompassed whether the species was diurnal and/or nocturnal. Cathemeral or crepuscular species were considered as both diurnal and nocturnal. Microhabitat included five categories of habitat use commonly reported in field guides and related literature: fossorial, terrestrial, aquatic, arboreal, and aerial.

Microhabitat categories are not mutually exclusive, meaning that a species can be present in more than one category to represent intermediate microhabitats, such as semifossorial (which involves both fossorial and terrestrial categories) or semiarboreal (which combines terrestrial and arboreal). Exceptionally for birds, we adapted microhabitat data from the EltonTraits database [47], which describes the estimated relative usage for seven types of foraging stratum. To make our definition of microhabitat similar across tetrapod groups, we reduced these seven categories to four by: summing the relative usage of species foraging below the water surface or on the water surface in the aquatic microhabitat; summing the relative usage of species foraging on the ground and below 2m in understorey as terrestrial; summing the relative usage of species foraging 2m upward in the canopy and just above canopy as arboreal. Species with aerial microhabitat were kept as defined in EltonTraits database [47], and no fossorial bird was reported in the later source. We then made binary the relative usage of aquatic, terrestrial, arboreal, and aerial microhabitat using a threshold of 30% to consider a species as typical of a given microhabitat type. In a departure from previous mammal databases that treated fossorial and terrestrial species collectively as 'terrestrial', we have reviewed microhabitat data to consider fossorial life-style separately from terrestrial [22,47].

Macrohabitat data followed the IUCN Habitat Classification scheme v. 3.1 [48]. This scheme describes 17 major habitat categories in which species can occur, and an 18th category for species with unknown major habitat (not included here). We also gathered data on species' major ecosystem (terrestrial, freshwater, and marine). For both macrohabitat and ecosystem, we initially used the *rredlist* package [49] to obtain macrohabitat for 31,740 species, and ecosystem for 32,442 species. Our macrohabitat variables correspond only to the first level of IUCN Habitat Classification scheme. For an additional 1,322 species, we extracted macrohabitat data from relevant literature, bringing the coverage to 33,019 species (99.2% of all species considered). Ecosystem data was extracted from the literature for another 837 species, encompassing 33,279 species and accounting for >99,9% of the total number of species.

We used the *rredlist* package [49] to obtain non-DD assessed status for 29,181 tetrapod species based on IUCN red list v. 2023-1 [50]. For 392 species not available via *rredlist*, we used non-DD assessed statuses matching those described in previous assessments and included in works using the same taxonomy of fully-sampled trees [20,24,26,51]. Data deficient species totalled 2,924 species. We did not find an assessed status for 555 species. To enhance the usability of TetrapodTraits, we also provide the respective IUCN binomials for 32,207 species based on IUCN 2023-1 [50].

To compute spatially-based attributes in TetrapodTraits, we derived expert-based range maps
for amphibians [24,48,52], reptiles [20,33,48], mammals [48,52–54], and birds [27,52]. We
matched the authoritative expert range maps for each of the tetrapod groups with the
corresponding phylogenies and edited species ranges to ensure that they represented the
species concept adopted in the corresponding phylogeny. Overall, our verification procedure
of the species range maps can be summarised under 10 scenarios: (i) no changes, where
species range maps matched directly with binomials in the phylogeny, (ii) synonyms, where
species range maps were direct synonyms to binomials in the phylogeny, thus requiring only
an updated name, (iii) split, where species range maps needed to be clipped from a parent
species, or when parent species needed to have part of their range removed, (iv) lumps, where
species range maps needed to be combined with those of other species, (v) new species-1,
where no range map was previously available, so we derived ranges based on recent literature,
(vi) new species-2, in the absence of any published map, we drew 10 km radius buffer around
point occurrence data (including the species type locality), (vii) new species-3, in the absence
of point occurrence data, we drew a polygon around nearby geographical features reported in
the literature (e.g. boundaries of a municipality or protected area). We used two additional
scenarios for extinct species [54] by referencing the natural ranges of either (viii) extant or
(ix) extinct species that coexisted with fossil records of the target extinct species. The last
scenario refers to (x) domesticated species, which were represented by their natural ranges
before domestication. *Homo sapiens* had its range map represented by the overlapping of all
range maps. We did not derive range maps for 11 species (1 amphibian, 2 bats, and 8
squamates) because information on their occurrence was either vaguely defined (e.g.,
continental land mass or very large administrative unit) or completely absent.

Species expert-based maps were used to compute different attributes related to species range
and biogeography. We extracted the latitude and longitude centroids of each range map.
Range size was measured as the number of 110×110 km equal-area grid cells intersected by
each species, a spatial resolution that minimises the presence of errors related to the use of
expert ranges maps [55–57]. We recorded the presence of a species in a grid cell if any part of
the species distribution polygon overlapped with the grid cell. We then computed the
proportion of the species range overlapped by each biogeographical realm.

To define if a species was insular endemic or not, we used the literature available [58,59]. We
further completed insularity data by registering species whose range maps intersected with
minor (< 2 km²) and major islands worldwide. Island vector data was sourced from Natural
Earth (www.naturalearthdata.com, [60]) databases, v. 4.1.0 and v. 5.1.1 for minor and major
islands, respectively. Species missing range maps were assumed as non-insular based on the
collection of type specimens within major continental land masses (e.g., South Asia, South
America, West Africa, Madagascar).

Finally, spatially-based attributes associated with environment or human influence were
computed as the within-range average value of continuous variables (e.g., annual mean
temperature, proportion of urban area) [61–64] weighted by species range occupancy in the
cell. All within-range attributes were derived from spatial layers available online [61–64] and
are individually described in the Results and Discussion section (see also Table S1).

Taxonomic Harmonisation
The taxonomy of the TetrapodTraits database follows the respective taxonomies of the recent,
fully-sampled phylogenies for each of the major tetrapod groups [20,25–28]. The amphibian
phylogeny taxonomy [25] follows the 19 February 2014 edition of AmphibiaWeb
(http://amphibiaweb.org), with 7,238 species. The phylogeny for chelonians [20] follows the
Turtles of the World (8th ed.) checklist [65], with the addition of *Aldabrachelys abrupta* and
Al. grandidieri, and the synonymisation of *Amyda ornata with Am. cartilaginea*, adding to
357 chelonian species. For crocodilians [20], the taxonomy followed [66], complemented by
the revalidation of *Crocodylus suchus* [67] and *Mecistops leptorhynchus* [68], and the

recognition of three *Osteolaemus* species [69,70], resulting in 27 species. The taxonomy of the squamate and tuatara phylogeny [26] follows the Reptile Database update of March 2015 (http://www.reptile-database.org) with 9,755 species. For brevity, we refer hereafter to species in the latter phylogeny as squamates, although we recognise that the tuatara, *Sphenodon punctatus,* is not a squamate. The taxonomy of the bird phylogeny [27] followed the Handbook of the Birds of the World [71], including 9,993 species. Finally, the taxonomy of the mammal phylogeny [28] follows the IUCN [72] with modifications resulting in a net addition of 398 species, bringing the total to 5,911 species.

To maximise data usage from previous compilation efforts, and to ensure coherence among species names and the multiple data sources, we built lists of synonyms and valid names based on multiple taxonomic databases [48,73,74], and extracted the unique synonyms in each taxonomic database. By unique synonym, we refer to a binomial (scientific name valid or not) applied to only one valid name. We then performed taxonomic reconciliation based on four steps:

1. ***Direct match with data sources:*** We directly paired the names of each of the 33,281 species in TetrapodTraits with the potential source of the data. Species-level attributes of closely related species could appear as identical values if the attributes had been extracted from sources in which different species were treated as synonyms. We minimised the inclusion of duplicated values by flagging each taxonomic match between TetrapodTraits and external data sources to ensure that each data entry was made only once.
2. ***Direct match with data source synonyms:*** For the species we were unable to directly match with the data source in step 1, we updated the taxonomy using the list of unique synonyms, and then performed a new matching operation which allowed us to extract and flag additional data whenever possible.
3. ***Direct match with TetrapodTraits synonymies:*** Some data sources may follow more recent taxonomies than those inherited from the fully-sampled phylogenies [20,25–28]. For species without a direct match after step 2, we updated their taxonomy in the TetrapodTraits using the list of unique synonyms and repeated the data extraction and flagging procedures.
4. ***Manual verification:*** For species without a direct match after step 3, we manually searched the specialised taxonomic databases (amphibians [73], reptiles [74], birds [75], and mammals [76]) for potential spelling errors and/or additional synonyms not yet included among our synonym lists. Whenever possible, we updated the taxonomy applied to data sources and then repeated the data extraction and flagging procedure. Species without data after the completion of step-4 were classified as missing data.

Outlier Verification

We implemented two approaches to detect potential inconsistencies in continuous attribute data, body length and body mass, before applying phylogeny-based methods to impute missing values.

1. ***Interquartile Range Criterion:*** Species body length and body mass were $\log_{10}$ transformed and then flagged as outliers if their value were outside the interval defined between $[q_{0.25} - 1.5 \times IQR]$ and $[q_{0.75} + 1.5 \times IQR]$, where $q_{0.25}$ and $q_{0.75}$ are respectively the first and third quartiles, and IQR is the interquartile range $[q_{0.25} - q_{0.75}]$.
2. ***Deviation from allometric relationship:*** Although allometric escape is a phenomenon observed in nature, we used interactive scatterplots to flag species with unusual deviations from the expected allometric relationship between body length and mass. We inspected allometric relationships separately for species within each Class, Order, and Suborder.

For species flagged in steps one or two above, we checked body length and/or body mass for validity and corrected these values where necessary. Data entries that could not be confirmed using a reliable source were purged from the database.

Taxonomic Imputation

The global scope of the present database inevitably includes some gaps that are hard to fill, e.g., natural history data for species known only from the holotype or a few specimens [12,77]. Previous studies have addressed this challenge and reduced data missingness by using values imputed at the level of genus or from close-relatives [47]. Although these earlier strategies of 'taxonomic imputation' might artificially reduce variability in attribute values, they are useful for filling gaps in highly conserved attributes, and can ultimately help increase the performance of phylogeny-based imputation methods applied in concert with correlated attribute data [78,79].

We used taxonomic imputations for two cases of missing data in microhabitat: chiropterans, who were considered 'aerial' (66 species), and dolphins and whales who were considered 'aquatic' (5 species). For the remaining tetrapod species, we computed the per-genus proportion of species in each type of activity time (diurnal or nocturnal), microhabitat (fossorial, terrestrial, aquatic, arboreal, aerial), macrohabitat (17 binary variables informing the IUCN Habitat Classification scheme), and ecosystem (terrestrial, freshwater, marine). If a type of activity time, microhabitat, macrohabitat, or ecosystem appeared in at least 70% of species in the genus, we assumed this ecological attribute was also present among species with missing data in the respective genus. Our goal was to reduce missing values in activity time, microhabitat, and macrohabitat for groups with well-known ecologies (observed data available for at least 70% of species) before running phylogeny-based imputation methods. We did not use taxonomic imputation for continuous attributes.

Phylogenetic Multiple Imputation

To minimise missing values and capture their uncertainty, we applied the *mixgb* method [30], a recently developed approach that combines the tree-based algorithm *XGBoost* [80] with predictive mean matching (PMM) [81], a multiple imputation technique. *XGBoost* captures interactions and nonlinear relations among variables, while PMM, alongside subsampling, addresses variability associated with missing data. PMM assigns imputed values to each missing entry based on a group of k donors whose predicted values are the most similar among the observed entries. One donor is then randomly selected, and its observed value is used for imputation [81,82]. The process is repeated m times to produce multiple imputations. When PMM uses a single donor without subsampling, imputations are expected to be identical.

The *XGBoost* does not directly include a phylogenetic tree into its computations. To account for phylogenetic information, we used the phylogenetic covariance matrix of each fully-sampled tree [20,25–28] to derive a set of phylogenetic filters (eigenvectors). We determined the number of phylogenetic filters to retain using the broken stick rule [83]. The selection of phylogenetic filters was performed separately for each tetrapod group and across the subset of 100 trees.

To assess the reliability of imputed data, we initially filtered a subset of species with complete data within each tetrapod group. Then, we randomly partitioned these subsets into 10 folds for cross-validation. In each iteration, one fold was excluded from the training process and used as testing data in subsequent modelling. Continuous variables (body length and body mass) were log_{10}-transformed to reduce skewness, while binary variables represented types of microhabitat (fossorial, terrestrial, aquatic, arboreal, aerial) and activity time (diurnal and nocturnal). Note that microhabitat and activity time types are non-mutually exclusive. For birds only, we complemented the observed attribute data with 10 morphometric traits (log_{10}-transformed) made recently available through the AVONET database [84].

XGBoost places a central emphasis on the tuning of hyperparameters, covering aspects such as learning rates, tree topology, subsampling, weighting, and regularisation [80,85]. Our tuning procedure began with an initial grid search, exploring 1K parameter combinations uniformly draw from specified ranges for five key hyperparameters: learning rate (η) from 0.01 to 0.3, maximum tree depth from 3 to 12, subsample portion of training data from 0.7 to 1, minimum child weight from 0.5 to 1.5, and number of boosting iterations (*nrounds*) from 30 to 1000. For continuous traits, our goal was to minimise the Normalised Root Mean Square Error (NRMSE) in *XGBoost* regression models using the 'reg:squarederror' objective, while for binary traits, we sought to reduce misclassification error in models using the 'binary:logistic' objective. We refined the hyperparameter selection by reassessing model performance with an additional 1K parameter combinations uniformly drawn from the parameter ranges defined by the top 5% of models. The tuning procedure was performed separately for each combination of response variable and tetrapod group. Other *XGBoost* parameters were kept at their default values.

Following hyperparameter tuning, we trained *mixgb* models under the 10-fold cross-validation approach. Each *mixgb* model considered the predictive mean matching with 10 donors, and provided 10 imputations for each missing entry. This process yielded a total of 10K imputed values for each missing entry across the subset of 100 phylogenetic trees (= 10 imputed values $\times$ 10 validation folds $\times$ 100 phylogenies). In each iteration, we assessed the reliability of imputations using four distinct validation metrics:

1. ***Pearson correlation:*** computed between imputed and observed values for continuous attributes (body length and body mass).
2. ***Regression slope:*** computed between imputed and observed values, where slope values > 1 indicate overestimation, and a slope < 1 indicates the underestimation for continuous attributes (body length and body mass).
3. ***Normalised Root Mean Square Error (NRMSE):*** computed for continuous attributes (body length and body mass), with lower values indicating higher accuracy [86].
4. ***Accuracy:*** measured the proportion of correctly classified entries, computed for binary categories representing microhabitat (fossorial, terrestrial, aquatic, arboreal, aerial) and activity time (diurnal and nocturnal).

Computations were carried out in R version 4.2.3 using the *mixgb* [30], *ape* [87], *gmodels* [88], *hydroGOF* [89], and *stats* [90] packages. The availability of the 10K imputed values per missing entry per species are reported under Data Availability.

We recognise that this section does not constitute a comprehensive investigation into the impacts of proportion of missing data and the degree of shared missingness on imputation performance. Delving into this aspect is beyond the scope of the present study and is an area for future research. For further details on mechanisms of data missingness, see [91]. Instead, our goal is to illustrate the utility of multiple imputation to uncover directional bias in natural history data.

Patterns of Shared Missing Data

We assessed co-occurrence patterns in missing data across species using the 'checkerboard score' (C-score; [92]), which is less prone to type II errors than other co-occurrence metrics computed under a null model approach [93]. The C-score was based on a binary presence-absence matrix (PAM) of species (columns) and missing attributes (rows). These attributes were represented by five binary variables informing the absence of observed values in body length, body mass, activity time, microhabitat, and threat status. Missing data in threat status were represented by Data Deficient (DD) or non-assessed species. We computed the C-score using individual PAMs for each genus and family with at least two species, and each pairwise combination of attribute variables, following recommendations by [94].

To verify whether an observed C-score differed from the value expected by chance, we built a null distribution of C-score values using a randomisation algorithm in which attribute completeness (rows sums) remained fixed while the probability of showing missing attribute values was considered equal for all species. Null distributions were built for each individual PAM using 10K iterations with a burn-in of 500. We then computed the standardised effect-size (SES) of the C-score and associated p-values and identified the pairwise attribute combinations with an aggregated (SES C-score < 0 and p < 0.05) or segregated (SES C-score > 0 and p < 0.05) co-occurrence pattern of missing data per taxa. Computations were performed using the *EcoSimR* [95] package in R.

Patterns and Biases in Imputed Traits

If certain parts of the multivariate attribute space are missing, the inclusion of imputed values may fill such 'invisible fractions' and change emergent attribute properties per taxon or geographical assemblage (e.g., average value, geometric mean, proportion or prevalence of a certain category). But changes in attribute property per taxon can also arise from the inclusion of biased imputed values [78], since imputation errors can increase either with the percentage of missing data or when multiple attributes are imputed simultaneously [15,78,79]. To verify how the inclusion of imputed values affected attribute patterns per taxon, we calculated the per-genus and per-family relative change in attribute property after adding imputed values.

Initially, we extracted the median (for continuous traits, body length and body mass) or the mean (for binary traits, categories of activity time and microhabitat) value per missing attribute across 10K imputations. We then computed attribute properties per-genus, -family, and -assemblage using (i) observed data only, and (ii) the combined dataset of observed and imputed data. For continuous attributes, we used the geometric mean due to its lower sensitivity to outliers and fluctuations in the median position value. For binary attributes, we computed the mean based on two scores derived from microhabitat and activity time categories: verticality (0 = strictly fossorial, 0.25 = fossorial and terrestrial, 0.5 = terrestrial or aquatic, 0.75 = terrestrial and arboreal, and 1 = strictly arboreal or aerial) and nocturnality (0 = strictly diurnal, 0.5 = cathemeral or crepuscular, 1 = strictly nocturnal). Relative change in attribute property was expressed as the ratio of attribute property obtained with the combined dataset to attribute property with the observed data only. We subtracted the relative change from 1 to centre it around zero:

$$Relative\ Change = 1 - (Attribute\ Property_{obs+imp}/Attribute\ Property_{obs})$$

Positive values of relative change indicate the proportional increase in attribute property after inclusion of imputed values, whereas negative values show the opposite. We used the R function *tree.merger* of the *RRphylo* package [96] to build a tetrapod super tree by combining the available global phylogenies for amphibians [25], chelonians and crocodilians [20], squamates [26], birds [27], and mammals [28]. The full tetrapod tree topology was only used for visualisation purposes. In the presence of severe data limitations (e.g., very low completeness, highly uneven sampled data, and lack of inter-correlated attributes), the relative changes in attribute properties could arise due to estimation errors instead of representing unsampled fractions of the attribute space. We therefore used Kruskal-Wallis tests to assess whether the relative increase or decrease of attribute properties differed with respect to the number of simultaneously imputed attributes.

Results and Discussion

TetrapodTraits offers insights into species-level attributes related to taxonomy, body size, activity time, microhabitat, macrohabitat, ecosystem, geography, environmental preferences, and threat status of tetrapod species (Table S1). Compilation efforts of previous works have significantly benefited TetrapodTraits, with these major data sources contributing to 68% of the natural history data entries. Additional information, whether compiled or revised from

original sources (e.g., field guides, taxonomic revisions, and species descriptions), accounted for 14.2% of data entries, and imputed values represented 18.1% of the natural history information (Fig. 1). In total, data acquisition involved the scanning of more than 3,400 references, including previously published databases. Imputation performance demonstrates high overall effectiveness across all tetrapods (Figs S1-S3), particularly for body mass (average Pearson correlation coefficient, r = 0.950, range = 0.692–0.973) and body length (r = 0.933, range = 0.585–0.976). Average accuracy is also high for activity time (0.880, range = 0.625–0.886) and microhabitat (0.897, range = 0.742–0.891). The number of observed, taxonomically imputed, or phylogenetically imputed data entries per tetrapod group are reported in Table S2, Supporting Information.

Data Completeness

Our assessment of data gaps focused on five attributes with considerable missing data: body length, body mass, activity time, microhabitat, and threat status. Across these five attributes, we observed complete species-level data for 43% of tetrapod species (n = 14,321), 36.9% of genera (n = 1,883), and 22.4% of families (n = 116). No genus or family showed zero completeness for these five attributes combined. The lowest completeness levels were among attributes related to body length (75.6%), activity time (77.1%), body mass (79.7%), and microhabitat (95.3%, Table S2). While assessed threat status was available for 98.3% of tetrapod species, this number decreased to 89.5% when DD species were treated as missing data (Fig. 2).

Across geographic assemblages, missingness of body length showed large variation, with Neotropical and Afrotropical species presenting the lowest completeness (Fig. 3A). The low spatial variability in attribute completeness for body mass, activity time, microhabitat, and assessed threat status (Fig. 3B–E) reveals that species with missing data are mostly narrow-ranged [4], which can limit the influence of these gaps in assemblage-level patterns. Of the 6,233 species missing at least two of the five attributes (body length, body mass, activity time, microhabitat, and threat status), 77.3% occurred in 10 or fewer 110 × 110 km equal-area grid cells. If we restrict missing data to three or more attributes, then 87% of species filtered occur in ≤10 grid cells. The completeness of genetic data in the fully-sampled trees of tetrapods [20,25–28] also showed high variation in spatial coverage (Fig. 3F), with the Afrotropical and Indomalayan realms emerging with the highest shortfall in genetic data [29]. Overall, our findings help illustrate that geographical patterns in data coverage can be strongly influenced by data for wide-ranged species.

Patterns of Shared Missing Data

We found non-random co-occurrence patterns in data gaps for 15.5% of the pairwise attribute combinations at the genus-level and for 37% at the family-level. Almost all non-random co-occurrence patterns showed aggregation of missing values, that is, missing data was shared among multiple attributes for the same species, particularly among squamates and amphibians (Fig. 4A). This supports the perspective that amphibians and squamates are the least researched tetrapod classes [2,3,34]. In our examination of shared gaps between threat status and other attributes, we found higher co-missingness in several tetrapod genera representative of leaf litter-dwelling (*Craugastor*, *Eleutherodactylus*, *Lerista*, and *Pristimantis*), fossorial and semifossorial (*Atractus*, *Crocidura*, and *Tantilla*), and arboreal or scansorial (*Cyrtodactylus*, *Dendroica*, and *Sphaerodactylus*) species. These findings indicate the uneven sampling coverage of attribute space within certain clades.

Aggregated data gaps were more common among species-rich clades with a lower proportion of missing data (Figs 4 and S9). That is, shared gaps were strongest in species-rich and well-sampled taxa, particularly in those holding many species known only from their type locality or holotype [77,97]. About 80% of the genera and 66% of families within the top 50 taxa in shared gaps include 'lost taxa', species that have not been reliably observed in >50 years but are not yet declared extinct [12]. While the rediscovery of 'lost' species is somewhat common

[97], their absence in attribute databases will not be easily filled through field observations [98]. The data gaps resulting from these poorly-known species often prevent their inclusion in trait-based research, potentially harming biogeographical research and conservation practice.

Biases Uncovered by Imputation-Based Gap Filling

We found that the use of imputations to fill gaps resulted in substantial changes in aggregate attribute properties and the insights they could support. These changes occurred in both directions (increasing and decreasing values) and were strongest in continuous attributes (body length and body mass). After the inclusion of imputed values, the average attribute value per genus decreased in 6.1% of the attribute-genus combinations and increased in 5.4% of these combinations. At the family-level, after the inclusion of imputed attributes, 16.3% of the attribute-family combinations showed an increase in the average values whereas 19.4% decreased (Figs 5 and S15).

Changes in attribute properties at the level of geographic assemblages revealed substantial alterations in average body size in tropical regions (Fig. 6A-F). Such differences are mostly absent in more temperate regions, reflecting the dedicated mobilisation and the resulting large availability of natural history data throughout North America, Europe, and Asia [2,4,99]. Average nocturnality of tetrapod assemblages tended to increase strongly after the inclusion of imputed values, pointing to a global knowledge bias from lack of observed attribute data (Fig. 6H-I), particularly for squamates and to a lesser extent, amphibians (Figs S10-S14). We attribute this to both the low detectability of nocturnal species [39,40] and to the general lack of data for tropical regions where nocturnal species predominate [100]. Changes in average verticality were less pronounced across tetrapod assemblages (Fig. 6K-L), which is somewhat expected given the relatively high completeness of microhabitat data.

Although imputation biases can increase in the absence of additional correlated attributes [78], we found no relationship between the number of imputed attributes and changes in attribute averages after imputation-based gap-filling (Fig. S16). That is, changes in average attributes were not higher in genera and families that included species with multiple missing attributes. Moreover, changes in average attributes were either mostly unrelated to shared gaps (Fig. S17) or occurred in opposite directions, with low shared gaps leading to greater changes.

Broadly, we found that the magnitude of change in average attributes from imputation-based gap-filling was associated with both completeness and richness (Figs S18-S20). Those taxa and assemblages with great incompleteness and with fewest species saw the most pronounced changes in average attribute values. Lower attribute completeness creates more opportunities for variation in attribute space after imputation-based gap-filling, whereas low species richness can enhance the importance of imputed values in the sample. Future research replacing imputed values with empirical data will reveal the accuracy of these estimates over time.

TetrapodTraits – a comprehensive, imputation informed attributes database

The full phylogenetically coherent database we developed, TetrapodTraits, is being made publicly available to support a range of research applications in ecology, evolution, and conservation and to help minimise the impacts of biased data in this model system. The database includes 24 species-level attributes linked to their respective sources across 33,281 tetrapod species. Entries across this database show at least 98% of completeness after the inclusion of imputed values. Specific fields clearly label data sources and imputations in the TetrapodTraits (Table S1), while additional tables record the 10K values per missing entry per species.

1. **Taxonomy –** includes 8 attributes that inform scientific names and respective higher-level taxonomic ranks, authority name, and year of species description. Field names:

Scientific.Name, Genus, Family, Suborder, Order, Class, Authority, and YearOfDescription.

2. **Phylogenetic tree –** includes 2 attributes that notify which fully-sampled phylogeny contains the species, along with whether the species placement was imputed or not in the phylogeny. Field names: TreeTaxon, TreeImputed.

3. **Body size –** includes 7 attributes that inform length, mass, and data sources on species sizes, and details on the imputation of species length or mass. Field names: BodyLength_mm, LengthMeasure, ImputedLength, SourceBodyLength, BodyMass_g, ImputedMass, SourceBodyMass.

4. **Activity time –** includes 5 attributes that describe period of activity (e.g., diurnal, fossorial) as dummy (binary) variables, data sources, details on the imputation of species activity time, and a nocturnality score. Field names: Diu, Noc, ImputedActTime, SourceActTime, Nocturnality.

5. **Microhabitat –** includes 8 attributes covering habitat use (e.g., fossorial, terrestrial, aquatic, arboreal, aerial) as dummy (binary) variables, data sources, details on the imputation of microhabitat, and a verticality score. Field names: Fos, Ter, Aqu, Arb, Aer, ImputedHabitat, SourceHabitat, Verticality.

6. **Macrohabitat –** includes 19 attributes that reflect major habitat types according to the IUCN classification, the sum of major habitats, data source, and details on the imputation of macrohabitat. Field names: MajorHabitat_1 to MajorHabitat_10, MajorHabitat_12 to MajorHabitat_17, MajorHabitatSum, ImputedMajorHabitat, SourceMajorHabitat. MajorHabitat_11, representing the marine deep ocean floor (unoccupied by any species in our database), is not included here.

7. **Ecosystem –** includes 6 attributes covering species ecosystem (e.g., terrestrial, freshwater, marine) as dummy (binary) variables, the sum of ecosystem types, data sources, and details on the imputation of ecosystem. Field names: EcoTer, EcoFresh, EcoMar, EcosystemSum, ImputedEcosystem, SourceEcosystem.

8. **Threat status –** includes 3 attributes that inform the assessed threat statuses according to IUCN red list and related literature. Field names: IUCN_Binomial, AssessedStatus, SourceStatus.

9. **RangeSize** – the number of 110×110 km grid cells covered by the species range map.

10. **Latitude** – coordinate centroid of the species range map.

11. **Longitude** – coordinate centroid of the species range map.

12. **Biogeography –** includes 8 attributes that present the proportion of species range within each biogeographical realm. Field names: Afrotropic, Australasia, IndoMalay, Nearctic, Neotropic, Oceania, Palearctic, Antarctic [101].

13. **Insularity –** includes 2 attributes that notify if a species is insular endemic (binary, 1 = yes, 0 = no), followed by the respective data source. Field names: Insularity, SourceInsularity.

14. **AnnuMeanTemp –** Average within-range annual mean temperature (Celsius degree). Data derived from CHELSA v. 1.2 [62].

15. **AnnuPrecip** – Average within-range annual precipitation (mm). Data derived from CHELSA v. 1.2 [62].

16. **TempSeasonality** – Average within-range temperature seasonality (Standard deviation $\times$ 100). Data derived from CHELSA v. 1.2 [62].

17. **PrecipSeasonality** – Average within-range precipitation seasonality (Coefficient of Variation). Data derived from CHELSA v. 1.2 [62].

18. **Elevation** – Average within-range elevation (metres). Data derived from topographic layers in EarthEnv [61].

19. **ETA50K** – Average within-range estimated time to travel to cities with a population >50K in the year 2015. Data from [64].

20. **HumanDensity** – Average within-range human population density in 2017. Data derived from HYDE v. 3.2 [63].

21. **PropUrbanArea** – Proportion of species range map covered by built-up area, such as towns, cities, etc. at year 2017 [63].

22. **PropCroplandArea** – Proportion of species range map covered by cropland area, identical to FAO's category 'Arable land and permanent crops' at year 2017 [63].

23. **PropPastureArea** – Proportion of species range map covered by cropland, defined as Grazing land with an aridity index > 0.5, assumed to be more intensively managed (converted in climate models) at year 2017 [63].

24. **PropRangelandArea** – Proportion of species range map covered by rangeland, defined as Grazing land with an aridity index < 0.5, assumed to be less or not managed (not converted in climate models) at year 2017 [63].

Conclusion

The growth in mobilised attribute data over the last several decades has enabled significant progress and increased geographic and taxonomic generality for global studies in comparative biology [102–104], macroevolution [105], macroecology [106], and conservation [107–109]. Despite the richness of available data, research outcomes and their interpretation often remained constrained by non-randomly missing data [6,8,79], and by the lack of standardisation across data dimensions (trait, phylogenetic, spatial) and data sources [110]. We show a clear latitudinal bias in the sampling of genetic data, with limited availability of genetic samples towards the Equator. In concert with taxonomic updates [111,112], these issues have imposed significant limits on multi-taxon investigations and global biodiversity synthesis [113–115]. We documented remaining global data gaps for key traits and then used imputation to fill these gaps, revealing the biases in our knowledge due to the missingness of attribute data. By using phylogeny-based imputation methods in concert with inter-related ecological attributes, we uncovered the hitherto invisible part of attribute space. Our findings confirmed the predominance of attributes that share missing data and provide a comprehensive assessment of gaps and biases across tetrapod groups.

Attributes predictions for species with missing data brought to light key correlates and allowed us to identify the significant consequences these gaps can have for inference. For instance, larger body size has been related to both high species detectability [38,39,42] and discovery probability [116–118], larger research effort and public interest [34,44,119,120], increased availability of assessed status [121,122], and better data sampling. Despite the

difficulties associated with researching large-sized species – e.g., high sensitivity to disturbances [123,124], lower abundance [125,126], challenges with collection or transport – the presence of multiple data biases towards small-sized species confirm the struggle of biodiversity scientists to accurately characterise the smallest organisms and the lower tail of size distributions in nature.

Biodiversity knowledge shortfalls are also clearly related to activity time and microhabitat. Nocturnal species often show lower detectability [39], and typically are less researched [44] and underassessed with respect to threat status [121], which helps explain why diurnal species are more likely to have activity time data. In addition, the higher detectability of terrestrial than arboreal or fossorial species via standard sampling methods [41,42] can lead to less biased occurrence data [127] and more research effort [34,44]. Although foraging stratum may constrain certain aspects of our biodiversity knowledge, verticality has a limited influence on the availability of microhabitat data.

Attribute imputations can be limited and biased [78,79]. For tetrapods generally and this study specifically, these concerns were minimised by the already substantial taxon-specific datasets and by the gap-oriented mobilisation of additional data. We had to impute all four focal attributes (body length, body mass, activity time, and microhabitat) in only 186 tetrapod species (<0.25%). No family or genus had the four focal attributes missing in all species. Two important diagnostics offered strong support for the robustness of our attribute predictions. First, there was no association between the number of imputed attributes and changes in average attributes after imputation-based gap-filling (Fig. S16). Second, there was no positive association between shared gaps in attribute data and changes in average attribute values (Fig. S17). We encourage researchers to consider how assumptions and uncertainty of any imputed data might affect downstream use and interpretation. However, for many tetrapod research cases, including species with imputed data is preferable to wholesale exclusion of data-limited species.

We have demonstrated that even for Tetrapoda, a central model system in biodiversity science, our ecological knowledge attribute data are lacking consistently across certain types of traits, taxa, and regions. Through careful imputation, we discovered that these gaps have led to attribute distributions with strong geographical and taxonomic biases. We expect that our approach and likely our general insights regarding biodiversity knowledge shortfalls will transcend taxa and systems. The consequences of biased data can percolate through trait-based metrics and comparative analyses, and may ultimately mislead research on functional diversity [8,19,128], extinction risk [6,21,129], and ecogeography [130–132]. The new phylogeny-based attribute database we have constructed can help minimise the impact of biased data and support new avenues of research in tetrapod conservation, ecology, and evolution.

Supporting Information
The Supporting Information includes Tables S1-S2 and Figures S1-S20.

Code Availability
The TetrapodTraits database and R-code used for data validation will be made available on vertlife.org and GitHub upon acceptance for publication.

Acknowledgements
We deeply thank the many authors of the numerous sources consulted during the preparation of this database. This work was possible only because of their valuable efforts. We gratefully acknowledge support from the following sources: São Paulo Research Foundation (FAPESP) for grants supporting MRM (#2021/11840-6 and #2022/12231-6), LFT (#2016/25358-3), KC (#2020/12558-0), and RZC (#2022/15247-0); Coordenação de Aperfeiçoamento de Pessoal de

Nível Superior (CAPES) for the fellowship to JJMG; Conselho Nacional de Desenvolvimento Científico - CNPq for research grants in support of FPW (#311504/2020-5) and LFT (#302834/2020-6); U.S. National Science Foundation (NSF) for grants supporting RAP (DEB-1441719), RCKB (DEB-1441652), and WJ (DEB-1441737 and DEB-1441719). WJ also acknowledges support from NASA grants 80NSSC17K0282 and 80NSSC18K0435.

Author Contributions

Conceptualisation: Mario R. Moura and Walter Jetz

Data Curation: Mario R. Moura, Karoline Ceron, Jhonny J. M. Guedes, Rosana Chen-Zhao, Yanina V. Sica, Julie Hart, Wendy Dorman, Julia M. Portmann, Pamela Gonzalez-del-Pliego, Ajay Ranipeta, Alessandro Catenazzi, Fernanda P. Werneck, Luís Felipe Toledo, Nathan S. Upham, Timothy J. Colston

Formal Analysis: Mario R. Moura

Funding Acquisition: R. Alexander Pyron, Rauri C.K. Bowie, Walter Jetz

Investigation: Mario R. Moura

Methodology: Mario R. Moura

Software: Mario R. Moura

Supervision: Walter Jetz

Visualisation: Mario R. Moura and Walter Jetz

Writing – original draft preparation: Mario R. Moura

Writing – review & editing: all authors

Competing interests

The authors declare no competing financial interests.

References

1. Beck J, Ballesteros-Mejia L, Buchmann CM, Dengler J, Fritz SA, Gruber B, et al. What's on the horizon for macroecology? Ecography. 2012;35: 673–683. doi:10.1111/j.1600-0587.2012.07364.x
2. Titley MA, Snaddon JL, Turner EC. Scientific research on animal biodiversity is systematically biased towards vertebrates and temperate regions. Schierwater B, editor. PLoS One. 2017;12: e0189577. doi:10.1371/journal.pone.0189577
3. Troudet J, Grandcolas P, Blin A, Vignes-Lebbe R, Legendre F. Taxonomic bias in biodiversity data and societal preferences. Sci Rep. 2017;7: 9132. doi:10.1038/s41598-017-09084-6
4. Etard A, Morrill S, Newbold T. Global gaps in trait data for terrestrial vertebrates. Sheard C, editor. Glob Ecol Biogeogr. 2020;29: 2143–2158. doi:10.1111/geb.13184
5. Oliver RY, Meyer C, Ranipeta A, Winner K, Jetz W. Global and national trends, gaps, and opportunities in documenting and monitoring species distributions. Moritz C, editor. PLOS Biol. 2021;19: e3001336. doi:10.1371/journal.pbio.3001336
6. González-Suárez M, Lucas PM, Revilla E. Biases in comparative analyses of extinction risk: mind the gap. J Anim Ecol. 2012;81: 1211–1222. doi:10.1111/j.1365-2656.2012.01999.x
7. Kim SW, Blomberg SP, Pandolfi JM. Transcending data gaps: a framework to reduce

inferential errors in ecological analyses. Ecol Lett. 2018;21: 1200–1210.
doi:10.1111/ele.13089

8.   Pakeman RJ. Functional trait metrics are sensitive to the completeness of the species'
trait data? Oksanen J, editor. Methods Ecol Evol. 2014;5: 9–15. doi:10.1111/2041-
210X.12136

9.   Rubin DB. Inference and Missing Data. Biometrika. 1976;63: 581.
doi:10.2307/2335739

10.   Nakagawa S, Freckleton RP. Missing inaction: the dangers of ignoring missing data.
Trends Ecol Evol. 2008;23: 592–596. doi:10.1016/j.tree.2008.06.014

11.   Nakagawa S, Freckleton RP. Model averaging, missing data and multiple imputation: a
case study for behavioural ecology. Behav Ecol Sociobiol. 2011;65: 103–116.
doi:10.1007/s00265-010-1044-7

12.   Martin TE, Bennett GC, Fairbairn A, Mooers AO. 'Lost' taxa and their conservation
implications. Anim Conserv. 2023;26: 14–24. doi:10.1111/acv.12788

13.   Sandel B, Gutiérrez AG, Reich PB, Schrodt F, Dickie J, Kattge J. Estimating the
missing species bias in plant trait measurements. J Veg Sci. 2015;26: 828–838.
doi:10.1111/jvs.12292

14.   Taugourdeau S, Villerd J, Plantureux S, Huguenin☐Elie O, Amiaud B. Filling the gap
in functional trait databases: use of ecological hypotheses to replace missing data. Ecol
Evol. 2014;4: 944–958. doi:10.1002/ece3.989

15.   Penone C, Davidson AD, Shoemaker KT, Di Marco M, Rondinini C, Brooks TM, et al.
Imputation of missing data in life-history trait datasets: which approach performs the
best? Methods Ecol Evol. 2014;5: 961–970. doi:10.1111/2041-210X.12232

16.   Stekhoven DJ, Buhlmann P. MissForest--non-parametric missing value imputation for
mixed-type data. Bioinformatics. 2012;28: 112–118.
doi:10.1093/bioinformatics/btr597

17.   Bruggeman J, Heringa J, Brandt BW. PhyloPars: estimation of missing parameter
values using phylogeny. Nucleic Acids Res. 2009;37: W179–W184.
doi:10.1093/nar/gkp370

18.   Goolsby EW, Bruggeman J, Ané C. Rphylopars: fast multivariate phylogenetic
comparative methods for missing data and within☐species variation. Fitzjohn R,
editor. Methods Ecol Evol. 2017;8: 22–27. doi:10.1111/2041-210X.12612

19.   Carmona CP, Tamme R, Pärtel M, de Bello F, Brosse S, Capdevila P, et al. Erosion of
global functional diversity across the tree of life. Sci Adv. 2021;7: eabf2675.
doi:10.1126/sciadv.abf2675

20.   Colston TJ, Kulkarni P, Jetz W, Pyron RA. Phylogenetic and spatial distribution of
evolutionary diversification, isolation, and threat in turtles and crocodilians (non-avian
archosauromorphs). BMC Evol Biol. 2020;20: 81. doi:10.1186/s12862-020-01642-3

21.   Cooke RSC, Eigenbrod F, Bates AE. Projected losses of global mammal and bird
ecological strategies. Nat Commun. 2019;10: 2279. doi:10.1038/s41467-019-10284-z

22.   Soria CD, Pacifici M, Di Marco M, Stephen SM, Rondinini C. COMBINE: a
coalesced mammal database of intrinsic and extrinsic traits. Ecology. 2021;102:
e03344. doi:10.1002/ECY.3344/SUPPINFO

23.   Caetano GH de O, Chapple DG, Grenyer R, Raz T, Rosenblatt J, Tingley R, et al.
Automated assessment reveals that the extinction risk of reptiles is widely
underestimated across space and phylogeny. PLOS Biol. 2022;20: e3001544.
doi:10.1371/journal.pbio.3001544

24.   González-del-Pliego P, Freckleton RP, Edwards DP, Koo MS, Scheffers BR, Pyron
RA, et al. Phylogenetic and Trait-Based Prediction of Extinction Risk for Data-
Deficient Amphibians. Curr Biol. 2019;29: 1557-1563.e3.
doi:10.1016/j.cub.2019.04.005

25.   Jetz W, Pyron RA. The interplay of past diversification and evolutionary isolation with
present imperilment across the amphibian tree of life. Nat Ecol Evol. 2018;2: 850–858.
doi:10.1038/s41559-018-0515-5

26.   Tonini JFR, Beard KH, Ferreira RB, Jetz W, Pyron RA. Fully-sampled phylogenies of

squamates reveal evolutionary patterns in threat status. Biol Conserv. 2016;204: 23–
31. doi:10.1016/j.biocon.2016.03.039

27.   Jetz W, Thomas GH, Joy JB, Hartmann K, Mooers AO. The global diversity of birds
in space and time. Nature. 2012;491: 444–448. doi:10.1038/nature11631

28.   Upham NS, Esselstyn JA, Jetz W. Inferring the mammal tree: Species-level sets of
phylogenies for questions in ecology, evolution, and conservation. PLOS Biol.
2019;17: e3000494. doi:10.1371/journal.pbio.3000494

29.   Diniz-Filho JAF, Loyola RD, Raia P, Mooers AO, Bini LM. Darwinian shortfalls in
biodiversity conservation. Trends Ecol Evol. 2013;28: 689–695.
doi:10.1016/j.tree.2013.09.003

30.   Deng Y, Lumley T. Multiple Imputation Through XGBoost. J Comput Graph Stat.
2023; 1–19. doi:10.1080/10618600.2023.2252501

31.   Buuren S van, Groothuis-Oudshoorn K. mice: Multivariate Imputation by Chained
Equations in R. J Stat Softw. 2011;45. doi:10.18637/jss.v045.i03

32.   Rässler S, Rubin DB, Zell ER. Imputation. WIREs Comput Stat. 2013;5: 20–29.
doi:10.1002/wics.1240

33.   Roll U, Feldman A, Novosolov M, Allison A, Bauer AM, Bernard R, et al. The global
distribution of tetrapods reveals a need for targeted reptile conservation. Nat Ecol
Evol. 2017;1: 1677–1682. doi:10.1038/s41559-017-0332-2

34.   Guedes JJM, Moura MR, Alexandre F. Diniz☐Filho J. Species out of sight: elucidating
the determinants of research effort in global reptiles. Ecography. 2023;2023: e06491.
doi:10.1111/ecog.06491

35.   Kéry M, Schmidt BR. Imperfect detection and its consequences for monitoring for
conservation. Community Ecol. 2008;9: 207–216. doi:10.1556/ComEc.9.2008.2.10

36.   Mazerolle MJ, Bailey LL, Kendall WL, Royle JA, Converse SJ, Nichols JD. Making
great leaps forward: Accounting for detectability in herpetological field studies. J
Herpetol. 2007;41: 672–689. doi:10.1670/07-061.1

37.   Jarzyna MA, Jetz W. Detecting the Multiple Facets of Biodiversity. Trends Ecol Evol.
2016;31: 527–538. doi:10.1016/j.tree.2016.04.002

38.   Anderson AS, Marques TA, Shoo LP, Williams SE. Detectability in Audio-Visual
Surveys of Tropical Rainforest Birds: The Influence of Species, Weather and Habitat
Characteristics. PLoS One. 2015;10: e0128464. doi:10.1371/journal.pone.0128464

39.   Einoder LD, Southwell DM, Lahoz-Monfort JJ, Gillespie GR, Fisher A, Wintle BA.
Occupancy and detectability modelling of vertebrates in northern Australia using
multiple sampling methods. PLoS One. 2018;13: e0203304.
doi:10.1371/journal.pone.0203304

40.   Richardson E, Nimmo DG, Avitabile S, Tworkowski L, Watson SJ, Welbourne D, et
al. Camera traps and pitfalls: an evaluation of two methods for surveying reptiles in a
semiarid ecosystem. Wildl Res. 2017;44: 637. doi:10.1071/WR16048

41.   Ribeiro-Júnior MA, Gardner TA, Ávila-Pires TC. Evaluating the Effectiveness of
Herpetofaunal Sampling Techniques across a Gradient of Habitat Change in a Tropical
Forest Landscape. J Herpetol. 2008;42: 733. doi:10.1670/07-097R3.1

42.   Chutipong W, Lynam A, Steinmetz R, Savini T, Gale G. Sampling mammalian
carnivores in western Thailand: Issues of rarity and detectability. Raffles Bull Zool.
2014;62: 521–535.

43.   Moura MR, Costa HC, Peixoto MA, Carvalho ALG, Santana DJ, Vasconcelos HL.
Geographical and socioeconomic determinants of species discovery trends in a
biodiversity hotspot. Biol Conserv. 2018;220: 237–244.
doi:10.1016/j.biocon.2018.01.024

44.   Brooke ZM, Bielby J, Nambiar K, Carbone C. Correlates of Research Effort in
Carnivores: Body Size, Range Size and Diet Matter. PLoS One. 2014;9: e93195.
doi:10.1371/journal.pone.0093195

45.   Amano T, Berdejo-Espinola V, Christie AP, Willott K, Akasaka M, Báldi A, et al.
Tapping into non-English-language science for the conservation of global biodiversity.
PLOS Biol. 2021;19: e3001296. doi:10.1371/journal.pbio.3001296

46. Zenni RD, Barlow J, Pettorelli N, Stephens P, Rader R, Siqueira T, et al. Multi‑lingual literature searches are needed to unveil global knowledge. J Appl Ecol. 2023;60: 380–383. doi:10.1111/1365-2664.14370

47. Wilman H, Belmaker J, Simpson J, de la Rosa C, Rivadeneira MM, Jetz W. EltonTraits 1.0: Species-level foraging attributes of the world's birds and mammals. Ecology. 2014;95: 2027–2027. doi:10.1890/13-1917.1

48. IUCN - International Union for Conservation of Nature. IUCN Red List of Threatened Species. In: Version 2021.3. 2021 p. www.iucnredlist.org.

49. Scott Chamberlain. rredlist: "IUCN" Red List Client. 2020. p. R package version 0.7.0. Available: https://cran.r-project.org/package=rredlist

50. IUCN - International Union for Conservation of Nature. IUCN Red List of Threatened Species. In: Version 2023.1. 2024 p. www.iucnredlist.org.

51. Jetz W, Thomas GH, Joy JB, Redding DW, Hartmann K, Mooers AO. Global Distribution and Conservation of Evolutionary Distinctness in Birds. Curr Biol. 2014;24: 919–930. doi:10.1016/j.cub.2014.03.011

52. Jetz W, McPherson JM, Guralnick RP. Integrating biodiversity distribution knowledge: toward a global map of life. Trends Ecol Evol. 2012;27: 151–159.

53. Marsh CJ, Sica Y V., Burgin CJ, Dorman WA, Anderson RC, Toro Mijares I, et al. Expert range maps of global mammal distributions harmonised to three taxonomic authorities. J Biogeogr. 2022; 1–14. doi:10.1111/jbi.14330

54. Faurby S, Davis M, Pedersen RØ, Schowanek SD, Antonelli1 A, Svenning J. PHYLACINE 1.2: The Phylogenetic Atlas of Mammal Macroecology. Ecology. 2018;99: 2626–2626. doi:10.1002/ecy.2443

55. Hurlbert AH, Jetz W. Species richness, hotspots, and the scale dependence of range maps in ecology and conservation. Proc Natl Acad Sci. 2007;104: 13384–13389.

56. Hurlbert AH, White EP. Disparity between range map-and survey-based analyses of species richness: patterns, processes and implications. Ecol Lett. 2005;8: 319–327.

57. Jetz W, Sekercioglu CH, Watson JEM. Ecological correlates and conservation implications of overestimating species geographic ranges. Conserv Biol. 2008;22: 110–119. doi:10.1111/j.1523-1739.2007.00847.x

58. Marino C, Leclerc C, Bellard C. Profiling insular vertebrates prone to biological invasions: What makes them vulnerable? Glob Chang Biol. 2022;28: 1077–1090. doi:10.1111/gcb.15941

59. Meiri S, Avila L, Bauer AM, Chapple DG, Das I, Doan TM, et al. The global diversity and distribution of lizard clutch sizes. McGill B, editor. Glob Ecol Biogeogr. 2020;29: 1515–1530. doi:10.1111/geb.13124

60. Kelso NV, Patterson T. Introducing natural earth data-naturalearthdata. com. Geogr Tech. 2010;5: 25.

61. Amatulli G, Domisch S, Tuanmu M-N, Parmentier B, Ranipeta A, Malczyk J, et al. A suite of global, cross-scale topographic variables for environmental and biodiversity modeling. Sci Data. 2018;5: 180040. doi:10.1038/sdata.2018.40

62. Karger DN, Conrad O, Böhner J, Kawohl T, Kreft H, Soria-Auza RW, et al. Climatologies at high resolution for the earth's land surface areas. Sci Data. 2017;4: 170122. doi:10.1038/sdata.2017.122

63. Klein Goldewijk K, Beusen A, Doelman J, Stehfest E. Anthropogenic land use estimates for the Holocene-HYDE 3.2. Earth Syst Sci Data. 2017;9: 927–953. doi:10.5194/essd-9-927-2017

64. Nelson A. Estimated travel time to the nearest city of 50,000 or more people in year 2000. Ispra: Global Environment Monitoring Unit - Joint Research Centre of the European Commission; 2008. Available: http://forobs.jrc.ec.europa.eu/products/gam/

65. Anders G J Rhodin, Iverson JB, Van-Dijk PP, Buhlmann KA, Pritchard PCH, Mittermeier RA. Turtles of the world: Annotated checklist and atlas of taxonomy, synonymy, distribution, and conservation status (8th Ed.). New York: Chelonian Research Foundation; 2017. Available: https://books.google.com/books/about/Turtles_of_the_World_Annotated_Checklist.ht

ml?id=9PSOswEACAAJ

66.   Oaks JR. A time-calibrated species tree of Crocodylia reveals a recent radiation of the
true crocodiles. Evolution. 2011;65: 3285–3297. doi:10.1111/j.1558-
5646.2011.01373.x

67.   Meredith RW, Hekkala ER, Amato G, Gatesy J. A phylogenetic hypothesis for
Crocodylus (Crocodylia) based on mitochondrial DNA: Evidence for a trans-Atlantic
voyage from Africa to the New World. Mol Phylogenet Evol. 2011;60: 183–191.
doi:10.1016/j.ympev.2011.03.026

68.   Shirley MH, Carr AN, Nestler JH, Vliet KA, Brochu CA. Systematic revision of the
living African Slender-snouted Crocodiles (Mecistops Gray, 1844). Zootaxa.
2018;4504: 151. doi:10.11646/zootaxa.4504.2.1

69.   Franke FA, Schmidt F, Borgwardt C, Bernhard D, Bleidorn C, Engelmann W-E, et al.
Genetic differentiation of the African dwarf crocodile *Osteolaemus tetraspis* Cope,
1861 (Crocodylia: Crocodylidae) and consequences for European zoos. Org Divers
Evol. 2013;13: 255–266. doi:10.1007/s13127-012-0107-1

70.   Shirley MH, Villanova VL, Vliet KA, Austin JD. Genetic barcoding facilitates captive
and wild management of three cryptic African crocodile species complexes. Anim
Conserv. 2015;18: 322–330. doi:10.1111/acv.12176

71.   del Hoyo J, Elliott A, Sargatal J, Christie DA. Handbook of the Birds of the World.
Barcelona: Lynx Edicions; 1992.

72.   IUCN - International Union for Conservation of Nature. IUCN Red List of Threatened
Species. In: Version 2015.1. 2015 p. www.iucnredlist.org.

73.   Frost DR. Amphibian Species of the World: an Online Reference. Version 6.1. In:
American Museums of Natural History, New York, USA [Internet]. 2021 [cited 20 Jan
2022]. Available: https://amphibiansoftheworld.amnh.org/

74.   Uetz P, Freed P, Hosek J. The Reptile Database. 2021 [cited 20 Jan 2022]. Available:
http://www.reptile-database.org

75.   Lepage D, Vaidya G, Guralnick R. Avibase – a database system for managing and
organizing taxonomic concepts. Zookeys. 2014;420: 117–135.
doi:10.3897/zookeys.420.7089

76.   Burgin CJ, Colella JP, Kahn PL, Upham NS. How many species of mammals are
there? J Mammal. 2018;99: 1–14. doi:10.1093/jmammal/gyx147

77.   Meiri S, Bauer AM, Allison A, Castro-Herrera F, Chirio L, Colli G, et al. Extinct,
obscure or imaginary: The lizard species with the smallest ranges. Stevens G, editor.
Divers Distrib. 2018;24: 262–273. doi:10.1111/ddi.12678

78.   Jardim L, Bini LM, Diniz-Filho JAF, Villalobos F. A Cautionary Note on Phylogenetic
Signal Estimation from Imputed Databases. Evol Biol. 2021;48: 246–258.
doi:10.1007/s11692-021-09534-0

79.   Johnson TF, Isaac NJB, Paviolo A, González‑Suárez M. Handling missing values in
trait data. Glob Ecol Biogeogr. 2021;30: 51–62. doi:10.1111/geb.13185

80.   Chen T, Guestrin C. XGBoost: A scalable tree boosting system. Proceedings of the
22nd ACM SIGKDD International Conference on Knowledge Discovery and Data
Mining. New York, NY, USA: ACM; 2016. pp. 785–794.
doi:10.1145/2939672.2939785

81.   Little RJA. Missing-Data Adjustments in Large Surveys. J Bus Econ Stat. 1988;6: 287.
doi:10.2307/1391878

82.   Rubin DB. Statistical Matching Using File Concatenation with Adjusted Weights and
Multiple Imputations. J Bus Econ Stat. 1986;4: 87. doi:10.2307/1391390

83.   Jackson DA. Stopping Rules in Principal Components Analysis: A Comparison of
Heuristical and Statistical Approaches. Ecology. 1993;74: 2204–2214.
doi:10.2307/1939574

84.   Tobias JA, Sheard C, Pigot AL, Devenish AJM, Yang J, Sayol F, et al. AVONET:
morphological, ecological and geographical data for all birds. Ecol Lett. 2022;25: 581–
597. doi:10.1111/ELE.13898

85.   Probst P, Bischl B, Boulesteix A-L. Tunability: Importance of Hyperparameters of

Machine Learning Algorithms. J Mach Learn Res. 2019;20: 1–32.

86. Barnston AG. Correspondence among the Correlation, RMSE, and Heidke Forecast Verification Measures; Refinement of the Heidke Score. Weather Forecast. 1992;7: 699–709. doi:10.1175/1520-0434(1992)007<0699:CATCRA>2.0.CO;2

87. Paradis E, Claude J, Strimmer K. APE: Analyses of Phylogenetics and Evolution in R language. Bioinformatics. 2004;20: 289–290. doi:10.1093/bioinformatics/btg412

88. Warnes GR, Bolker B, Lumley T, Johnson RC. gmodels: Various R Programming Tools for Model Fitting. 2018. p. R package version 2.18.1. Available: https://cran.r-project.org/package=gmodels

89. Zambrano-Bigiarini M. hydroGOF: Goodness-of-Fit Functions for Comparison of Simulated and Observed Hydrological Time Series. 2020. p. v. 0.4-0. doi:10.5281/zenodo.3707013

90. R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: The R Foundation for Statistical Computing Platform; 2022. p. v. 4.2.0.

91. Little RJA, Rubin DB. Statistical Analysis with Missing Data. Hokoben: John Wiley & Sons, Inc.; 2020.

92. Stone L, Roberts A. The checkerboard score and species distributions. Oecologia. 1990;85: 74–79. doi:10.1007/BF00317345

93. Gotelli NJ. Null model analysis of species co-occurrence patterns. Ecology. 2000;81: 2606–2621. doi:10.1890/0012-9658(2000)081[2606:NMAOSC]2.0.CO;2

94. Ulrich W, Gotelli NJ. Pattern detection in null model analysis. Oikos. 2013;122: 2–18. doi:10.1111/j.1600-0706.2012.20325.x

95. Gotelli NJ, Hart E, Ellison A. Null Model Analysis for Ecological Data. R package version 0.1.0. 2015. pp. 1–53.

96. Castiglione S, Tesone G, Piccolo M, Melchionna M, Mondanaro A, Serio C, et al. A new method for testing evolutionary rate variation and shifts in phenotypic evolution. Cooper N, editor. Methods Ecol Evol. 2018;9: 974–983. doi:10.1111/2041-210X.12954

97. Scheffers BR, Yong DL, Harris JBC, Giam X, Sodhi NS. The World's Rediscovered Species: Back from the Brink? Gratwicke B, editor. PLoS One. 2011;6: e22531. doi:10.1371/journal.pone.0022531

98. Ladle R, Jepson P, Malhado ACM, Jennings S, Barua M. The causes and biogeographical significance of species' rediscovery. Front Biogeogr. 2012;3: 111–118. doi:10.21425/F5FBG12432

99. Meyer C, Kreft H, Guralnick R, Jetz W. Global priorities for an effective information basis of biodiversity distributions. Nat Commun. 2015;6: 8221. doi:10.1038/ncomms9221

100. Vidan E, Roll U, Bauer A, Grismer L, Guo P, Maza E, et al. The Eurasian hot nightlife: Environmental forces associated with nocturnality in lizards. Glob Ecol Biogeogr. 2017;26: 1316–1325. doi:10.1111/geb.12643

101. Dinerstein E, Olson D, Joshi A, Vynne C, Burgess ND, Wikramanayake E, et al. An ecoregion-based approach to protecting half the terrestrial realm. Bioscience. 2017;67: 534–545. doi:10.1093/biosci/bix014

102. Stark G, Pincheira-Donoso D, Meiri S. No evidence for the 'rate-of-living' theory across the tetrapod tree of life. Glob Ecol Biogeogr. 2020;29: 857–884. doi:10.1111/GEB.13069

103. Atwood TB, Valentine SA, Hammill E, McCauley DJ, Madin EMP, Beard KH, et al. Herbivores at the highest risk of extinction among mammals, birds, and reptiles. Sci Adv. 2020;6. doi:10.1126/sciadv.abb8458

104. Anderson SR, Wiens JJ. Out of the dark: 350 million years of conservatism and evolution in diel activity patterns in vertebrates. Evolution. 2017; 1–16. doi:10.1111/evo.13284

105. Pigot AL, Sheard C, Miller ET, Bregman TP, Freeman BG, Roll U, et al. Macroevolutionary convergence connects morphological form to ecological function in birds. Nat Ecol Evol. 2020;4: 230–239. doi:10.1038/s41559-019-1070-4

106. Violle C, Reich PB, Pacala SW, Enquist BJ, Kattge J. The emergence and promise of functional biogeography. Proc Natl Acad Sci. 2014;111: 13690–13696. doi:10.1073/pnas.1415442111

107. Cardillo M, Mace GM, Gittleman JL, Jones KE, Bielby J, Purvis A. The predictability of extinction: biological and external correlates of decline in mammals. Proc R Soc B Biol Sci. 2008;275: 1441–1448. doi:10.1098/rspb.2008.0179

108. Pollock LJ, Thuiller W, Jetz W. Large conservation gains possible for global biodiversity facets. Nature. 2017;546: 141–144. doi:10.1038/nature22368

109. Lee TM, Jetz W. Unravelling the structure of species extinction risk for predictive conservation science. Proc R Soc B Biol Sci. 2011;278: 1329–1338. doi:10.1098/rspb.2010.1877

110. Schneider FD, Fichtmueller D, Gossner MM, Güntsch A, Jochum M, König‐Ries B, et al. Towards an ecological trait‐data standard. Methods Ecol Evol. 2019;10: 2006–2019. doi:10.1111/2041-210X.13288

111. Bremer K, Bremer B, Karis P, Källersjö M. Time for change in taxonomy. Nature. 1990;343: 202–202. doi:10.1038/343202a0

112. Garnett ST, Christidis L. Taxonomy anarchy hampers conservation. Nature. 2017;546: 25–27. doi:10.1038/546025a

113. Wüest RO, Zimmermann NE, Zurell D, Alexander JM, Fritz SA, Hof C, et al. Macroecology in the age of Big Data – Where to go from here? J Biogeogr. 2019; jbi.13633. doi:10.1111/jbi.13633

114. Hortal J, de Bello F, Diniz-Filho JAF, Lewinsohn TM, Lobo JM, Ladle RJ. Seven Shortfalls that Beset Large-Scale Knowledge of Biodiversity. Annu Rev Ecol Evol Syst. 2015;46: 523–549. doi:10.1146/annurev-ecolsys-112414-054400

115. Heberling JM, Miller JT, Noesgaard D, Weingart SB, Schigel D. Data integration enables global biodiversity synthesis. Proc Natl Acad Sci. 2021;118. doi:10.1073/pnas.2018093118

116. Moura MR, Jetz W. Shortfalls and opportunities in terrestrial vertebrate species discovery. Nat Ecol Evol. 2021;5: 631–639. doi:10.1038/s41559-021-01411-5

117. Diniz-Filho JAF, Bastos RP, Rangel TFLVB, Bini LM, Carvalho P, Silva RJ. Macroecological correlates and spatial patterns of anuran description dates in the Brazilian Cerrado. Glob Ecol Biogeogr. 2005;14: 469–477. doi:10.1111/j.1466-822x.2005.00165.x

118. Meiri S. Small, rare and trendy: traits and biogeography of lizards described in the 21st century. J Zool. 2016;299: 251–261. doi:10.1111/jzo.12356

119. Roll U, Mittermeier JC, Diaz GI, Novosolov M, Feldman A, Itescu Y, et al. Using Wikipedia page views to explore the cultural importance of global reptiles. Biol Conserv. 2016;204: 42–50. doi:10.1016/j.biocon.2016.03.037

120. Tyler EHM, Somerfield PJ, Berghe E Vanden, Bremner J, Jackson E, Langmead O, et al. Extensive gaps and biases in our knowledge of a well-known fauna: implications for integrating biological traits into macroecology. Glob Ecol Biogeogr. 2012;21: 922–934. doi:10.1111/j.1466-8238.2011.00726.x

121. Meiri S, Chapple DG. Biases in the current knowledge of threat status in lizards, and bridging the 'assessment gap.' Biol Conserv. 2016;204: 6–15. doi:10.1016/j.biocon.2016.03.009

122. Miqueleiz I, Bohm M, Ariño AH, Miranda R. Assessment gaps and biases in knowledge of conservation status of fishes. Aquat Conserv Mar Freshw Ecosyst. 2020;30: 225–236. doi:10.1002/aqc.3282

123. Blumstein DT. Developing an evolutionary ecology of fear: how life history and natural history traits affect disturbance tolerance in birds. Anim Behav. 2006;71: 389–399. doi:10.1016/j.anbehav.2005.05.010

124. Henle K, Davies KF, Kleyer M, Margules C, Settele J. Predictors of Species Sensitivity to Fragmentation. Biodivers Conserv. 2004;13: 207–251. doi:10.1023/B:BIOC.0000004319.91643.9e

125. Robinson JG, Redford KH. Body Size, Diet, and Population Density of Neotropical

Forest Mammals. Am Nat. 1986;128: 665–680. doi:10.1086/284596

126.   White EP, Ernest SKM, Kerkhoff AJ, Enquist BJ. Relationships between body size
and abundance in ecology. Trends Ecol Evol. 2007;22: 323–330.
doi:10.1016/j.tree.2007.03.007

127.   Meyer C, Jetz W, Guralnick RP, Fritz SA, Kreft H. Range geometry and socio-
economics dominate species-level biases in occurrence information. Glob Ecol
Biogeogr. 2016. doi:10.1111/geb.12483

128.   Cooke RSC, Bates AE, Eigenbrod F. Global trade-offs of functional redundancy and
functional dispersion for birds and mammals. Glob Ecol Biogeogr. 2019;28: 484–495.
doi:10.1111/geb.12869

129.   Vilela B, Villalobos F, Rodríguez MÁ, Terribile LC. Body Size, Extinction Risk and
Knowledge Bias in New World Snakes. PLoS One. 2014;9: e113429.
doi:10.1371/journal.pone.0113429

130.   Vidan E, Novosolov M, Bauer AM, Herrera FC, Chirio L, Campos CN, et al. The
global biogeography of lizard functional groups. J Biogeogr. 2019;46: 2147–2158.
doi:10.1111/jbi.13667

131.   Oliveira BF, Scheffers BR. Vertical stratification influences global patterns of
biodiversity. Ecography. 2019;42: 249–249. doi:10.1111/ecog.03636

132.   Gaston KJ, Chown SL, Evans KL. Ecogeographical rules: elements of a synthesis. J
Biogeogr. 2008;35: 483–500. doi:10.1111/j.1365-2699.2007.01772.x

1170
1171

Figure Legends

1173

Figure 1. Relative contribution of major data sources to natural history information in TetrapodTraits. For improved readability, only sources containing data entries for at least 500 species are shown. Boxes labelled 'Other sources' encompass combined information not included within major data sources, often compiled from field guides and taxonomic-specific literature for tetrapod species.

1179

Figure 2. Attribute data coverage (completeness) of tetrapod species by family. Colour rings show the proportion of species with observed attributes in each tetrapod family for the five attributes with the lowest completeness. Phylogenetic relationships shown are simply the family-level topology of tetrapod families. Darker colours indicate higher completeness.

1184

Figure 3. Average data completeness across tetrapod assemblages. Proportion of species with observed values for: (A) body length, (B) body mass, (D) activity time, (E) microhabitat, (G) assessed threat status, (H) phylogenetic data, (J) attribute set, i.e., the average pattern for maps depicted in ABDE, and (K) the complete database, i.e., average pattern for maps depicted in ABDEGH. Maps show grid cell assemblages of 110×110 km size in an equal-area projection. Latitudinal plots (C, F, I, L) show the average values of these cells across latitudes. Colour ramps followed Jenks' natural breaks classification. See Figs S4-S8 for spatial patterns of attribute completeness for each tetrapod group.

1193

Figure 4. The top 50 tetrapod genera with the most pronounced patterns of shared data gaps. For each genus, the aggregation metric equals the median value of the standardised effect-size (SES) of the C-score metric computed across (A) all pairwise attribute combinations among the five focal traits; and (B) all attribute pairs involving threat status. Grey numbers on the left side of each panel indicate the species richness per genus. See Fig. S9 for aggregation patterns computed at the family-level.

1200

Figure 5. Changes in family-level attributes values after imputation-based gap-filling. For continuous attributes (BL, body length; BM, body mass), relative change in attribute property quantifies the proportional change in the geometric mean attribute per genus. For

binary attributes (types of activity time and microhabitat), relative change in attribute property measures the change in mean species nocturnality (Noct) and verticality (Vert) scores. The inset donut chart denotes the proportion of genera (inner donut) and families (outer donut) by category of attribute change. The y-axis of histograms is square-rooted to improve readability.

Figure 6. Changes in average attribute per tetrapod assemblage after imputation-based gap-filling. For each grid cell, maps show the average species attribute value and respective relative change in attribute value after inclusion of imputed values for (A-C) body length, (D-F) body mass, (G-I) nocturnality, and (J-L) verticality. Body length and body mass were $\log_{10}$ transformed before computations. Grey cells indicate assemblages without species with imputed values. See Figs S10-S14 for patterns of attribute change for each tetrapod group.

A

Genus	N
Amphisbaena	106
Pteropus	64
Varanus	77
Typhlops	49
Oligosoma	41
Mus	39
Dipsas	36
Peromyscus	58
Atelopus	97
Chalcides	31
Eutropis	31
Dendroica	30
Naja	29
Buteo	27
Gerbillus	48
Cyrtopodion	28
Indotyphlops	24
Plestiodon	47
Microgale	22
Lycodon	51
Pristimantis	460
Rattus	66
Atractus	138
Leptodactylus	77
Pipistrellus	49
Callulops	20
Acanthodactylus	42
Anolis	395
Akodon	44
Liolaemus	245
Hebius	40
Sphaerodactylus	103
Hipposideros	73
Cyrtodactylus	196
Hemidactylus	132
Epictia	27
Hyperolius	144
Sceloporus	97
Eleutherodactylus	187
Litoria	162
Aspidoscelis	40
Crocidura	192
Caprimulgus	58
Parvoscincus	24
Lerista	94
Sphenomorphus	117
Tantilla	62
Emoia	78
Agama	45
Plethodon	55

Legend: ■ Amphibians ■ Squamates □ Birds □ Mammals

Inset top: $R = 0.379$, $p < 2.2e{-16}$ — SES C-score vs Average data missingness
Inset bottom: $R = -0.541$, $p < 2.2e{-16}$ — SES C-score vs $\log_{10}$ Per genus richness

Aggregation in missing values (SES C−score)

B

Genus	N
Oligosoma	41
Dendroica	30
Naja	29
Buteo	27
Pristimantis	460
Liolaemus	245
Sphaerodactylus	103
Cyrtodactylus	196
Atractus	138
Lerista	94
Anolis	395
Hyperolius	144
Erythrolamprus	48
Hemidactylus	132
Emoia	78
Scinax	111
Caprimulgus	58
Leptodactylus	77
Agama	45
Rhacophorus	84
Litoria	162
Proceratophrys	36
Hipposideros	73
Rana	100
Calotes	25
Phrynobatrachus	88
Myotis	114
Tantilla	62
Carlia	44
Crocidura	192
Eremias	35
Philautus	52
Dipodomys	21
Hypsiboas	91
Scelotes	24
Crotalus	39
Rhinolophus	87
Ameiva	30
Ameerega	31
Hebius	40
Japalura	28
Fejervarya	42
Vipera	22
Sphenomorphus	117
Platymantis	72
Reithrodontomys	22
Cnemaspis	117
Gehyra	41
Uperoleia	27
Oligoryzomys	20

Avg. Data Missingness 0.0 0.2 0.4

Inset top: $R = 0.471$, $p = 4.01e{-11}$ — SES C-score vs Average data missingness
Inset bottom: $R = -0.564$, $p = 3.68e{-16}$ — SES C-score vs $\log_{10}$ Per genus richness

Aggregation in missing values (SES C−score)