Tuning the stator subunit of the flagellar motor with coiled-coil engineering.

Pietro Ridone[1], Daniel L. Winter[1], Matthew A. B. Baker[1]

1.   School of Biotechnology and Biomolecular Science, UNSW Sydney, Australia.

Abstract:

Many bacteria swim driven by an extracellular filament rotated by the bacterial flagellar
motor. This motor is powered by the stator complex, $MotA_5MotB_2$, a heterodimeric complex
which forms an ion channel which couples energy from the ion motive force to torque
generation. Recent structural work revealed that stator complex consists of a ring of five
MotA subunits which rotate around a central dimer of MotB subunits. Transmembrane (TM)
domains TM3 and TM4 from MotA combine with the single TM domain from MotB to form
two separate ion channels within this complex. Much is known about the ion binding site
and ion specificity; however, to date, no modelling has been undertaken to explore the
MotB-MotB dimer stability and the role of MotB conformational dynamics during rotation.
Here, we modelled the central MotB dimer using coiled-coil engineering and modelling
principles and calculated free energies to identify stable states in the operating cycle of the
stator. We found 3 stable coiled-coil states with dimer interface angles of 28°, 56° and 64°.
We tested the effect of strategic mutagenesis on the comparative energy of the states and
correlated motility with a specific hierarchy of stability between the three states. In general,
our results indicate agreement with existing models describing a 36° rotation step of the
MotA pentameric ring during the power stroke and provide an energetic basis for the
coordinated rotation of the central MotB dimer based on coiled-coil modelling.

Introduction:

The bacterial flagellar motor (BFM) is a complex nanoscale engine that converts ion flux into the rotation of a flagellum to propel bacteria through their surroundings[1,2]. A crucial component of this complex is the stator subunit which is responsible for the torque-generating step of the motor[3,4]. In *E. coli,* the stator subunit is in fact composed of two transmembrane (TM) proteins (MotA and MotB) which assemble around the basal rotor of the flagellar motor and form a proton channel at their interface[5]. The inward flux of ions across the complex is efficiently coupled to a mechanical stimulus applied onto the rotor protein FliG which ultimately drives the rotation of a single flagellum[6].

The structure of the full $MotA_5MotB_2$ complex was solved in 2020[7,8], revealing that the stator complex likely acted as a rotary motor itself. Further recent structural work has identified the basis for ion selectivity and many catalytic residues involved in the process[9]. The current consensus model is that a transmembrane pentameric ring of MotA subunits rotates around a central dimer stalk of MotB TM domains[9,10]. The model details the conformational waypoints through which the MotA subunit passes during the gating cycle of the stator, but our understanding of the dynamics, in general, is limited, and in particularly for the MotB subunit central stalk.

The central MotB dimer is important because it harbours the catalytic centre. It has been proposed to rotate in place between protonated and non-protonated states[10]. Previous work suggested that the MotB TM domains exchange their interface contacts during gating and may pivot along their helical axis[11], in a similar fashion to the proposed mechanism for the homologous TolR protein[12]. The dimerization of MotB at the TM domain was shown to be essential for motility as modifications to TM residues that restrict or alter the conformational space of the dimer (e.g., cysteine crosslinking and tryptophan-scanning mutagenesis) generally diminished motility[11, 12].

The MotB stalk appears to assemble as a dimeric parallel coiled-coil in recently published structures[7,8]. Coiled-coils are well-characterized structural motifs where alpha-helices interact to form dimers or higher order oligomeric states[13]. A repeating seven residues-long "heptad" motif with alternating hydrophobic (*h*) and polar (*p*) residues forming the general pattern *hpphppp* along the alpha-helices results in a core interface where the hydrophobic residues engage in knob-into-hole (KIH) interactions. The sequence-structure relationship of coiled-coils is well understood, and design principles have been established to create synthetic coiled-coils. Dimers are a specific example: it has been shown that isoleucine and leucine in positions *a* and *d* of a repeating heptad *abcdefg* (corresponding to the pattern *hpphppp*) greatly favor dimerization over other oligomeric states, whereas isoleucine at both positions favors trimers[14]. Moreover, the structure of coiled-coils can be parametrically modelled, which greatly accelerates the calculation of their physical properties, such as interaction strengths[15]. Parametric modelling entails the use of very few parameters (the radius, interface angle, and pitch of a coiled-coil) to describe the entire backbone structure. Parametric modelling, in combination with the BUDE (Bristol University Docking Engine)[16] forcefield, has been successfully applied to predict the structures of natural and synthetic coiled-coils[17,18].

In this study, we investigated the dimerization interface of MotB using parametric modelling
to represent the MotB dimer as a dynamic coiled-coil. This simplification of the MotB
structure enables the rapid calculation of the total energy of the system to discriminate
between stable and unstable conformations. From these results, we rationally designed
stator sequence variants to be tested *in vivo* and to better understand how MotB dimer
stability may relate to overall conformational dynamics of the stator complex during its
gating cycle.

In this study, we investigated the dimerization interface of MotB using parametric modelling
to represent the MotB dimer as a dynamic coiled-coil. This simplification of the MotB
structure enables the rapid calculation of the total energy of the system to discriminate
between stable and unstable conformations. From these results, we rationally designed
stator sequence variants to be tested *in vivo* and to better understand how MotB dimer
stability may relate to the conformational dynamics of the stator complex during its gating
cycle. By testing the *in vivo* motility MotB variants predicted to differ in stable and unstable
conformational states, we aimed to identify which coiled-coil conformations were
correlated with motility, and to inform the structural changes undertaken by *Ec*MotB and its
extant homologs during their gating cycle.

Results

We modelled the *Ec*MotB TM domain as a parallel coiled-coil based on the most recent
structures of MotB from *B. subtilis* (PDBID: 6YSL), *C. sporogenes* (PDBID: 6YSF) and *C. jejuni*
(PDBID: 6YKM)[7,8] (Fig. 1). A set of ISAMBARD models of the 20-residue long *Ec*MotB peptide
IAYA**D**FMTAMMAFFLVMWLI (corresponding to residues 28-47 of WT *Ec*MotB; the catalytic
residue D32 is indicated in bold) with varying interface angles were aligned to the crystal
structures to formally identify the "register" of the MotB coiled-coil, that is, what positions
in the coiled-coil heptad each residue occupies. Based on RMSD calculations and the
orientation of the catalytic aspartate (D32), we established that the 20-residue long MotB
dimer stalk is best described as a coiled-coil starting at position *e*. In other words, residues
A31, M34, M38, F41, and W45 occupy positions *a* and *d* along the coiled-coil, forming the
core interface. These results agree with previous reports where mutation of these residues
to cysteine led to crosslinking between the two MotB subunits, indicating proximity[19].
Further analysis using CCBuilder of the peptide IAYADFMTAMMAFFLVMWLI as a parallel
dimeric coiled-coil starting at position *e* indicated that the five core residues could form KIH
interactions.

In light of these results, we decided to model MotB as a dimeric coiled to inform our *in vivo*
mutagenesis experiments. To this end, rather than simply considering the most stable
conformation calculated by ISAMBARD, we considered the conformational space of MotB by
generating coiled-coil structural models with varying interface angles and radii and then, for
each combination of the two parameter values, calculating the total energy of the system
using the BUDE forcefield (hereafter, BUDE). This *in silico* conformational screen revealed
the potential existence of wells (more stable configurations) and peaks (unstable
configurations) across the energy landscape of the WT *Ec*MotB (Fig. 2A). Interestingly, the

deepest stability well (#4, Fig. 2B) did not correspond to the configuration adopted in the
cryo-EM structures (#1, Fig. 2B).

Next, we considered strategic point mutation of coiled-coils to investigate if the stability of
the MotB dimer would change after replacing the key *a* and *d* residues forming the core
interface. We considered the design rules of synthetic coiled-coils and hypothesised that
alanine mutations would systematically weaken the coiled-coil interaction by disrupting the
hydrophobic KIH interactions, whereas isoleucine and leucine mutations would lead to more
stable KIH interactions and thus a more stable coiled-coil (when isoleucine and leucine
occupy positions *a* and *d*) or to steric clashes (when isoleucine occupies both positions *a* and
d). Therefore, core residues were systematically mutated *in silico* to alanine, isoleucine, or
leucine at positions *a* and *d* in over 200 permutations (see Method and Supplementary
Table 1). In addition, we also considered the M38G mutation that appeared to disrupt KIH
interactions in CCBuilder analyses. In total, a library of 221 sequence variants was generated
in silico.

To test the above hypothesis *in silico*, we repeated the conformational analysis using
ISAMBARD for each MotB variant modelled as a coiled-coil and compared the energy
landscape of each sequence variant to the WT sequence. Different mutations resulted in a
shift of the energy wells towards different interface angles and radii values and, in some
extreme cases, to the absence of stability wells. The energy peaks corresponding to
potential inaccessible conformations also differed. Altogether, the results indicate that the
selected mutations in the core of the MotB dimerization interface could affect its accessible
conformational space and, presumably, affect cell motility. Further to this analysis, the
three-dimensional (3D) BUDE energy landscapes were collapsed into two-dimensional (2D)
energy profiles by averaging the energy values for radii between 4 to 5 Å (Fig. 2C). This
range of radii values in BUDE encompassed the radii observed in the reference structures
(4.7 to 4.9 Å) and contained most of the energy wells and peaks of interest to compare
sequence variants. In these dimensionally reduced datasets, we subsequently considered
the interface angle parameter as the main variable affecting the energy of the coiled-coil
system.

We next sought to select a subset of mutants to verify whether the differences in calculated
energy profiles for MotB sequence variants correlated with BFM activity. Comparison of the
2D BUDE energy profiles suggested that single mutations could be sufficient to alter the
accessible conformational space of MotB, and therefore affect cell motility (Fig. 2C, 2D). We
selected 9 single mutants (SI Fig. 1) and an additional 10 double mutants and 4 triple
mutants (SI Fig. 2) that showed varied energy profiles for *in vivo* experimentation. We
restricted our *in vivo* testing to variants displaying alanine and/or isoleucine replacements.

We recombinantly expressed the selected sequence variants of MotB in an *E. coli* Δ*motAB*
background strain and assessed the motility of bacteria on swim plates (Fig. 3A) and at a
single cell level (Fig. 3B). Some of the single point mutations affected the size of swim rings
(M34A, M34I, M38A, F41A) whereas all other mutations, including the selected double and
triple mutants, abolished motility (Supplementary Fig. 3). Amongst the motile mutants, the
first two had a comparable (M34A) or markedly positive (M34I) effect on motility, while

M38A and F41A decreased the size of the swim rings (Fig.3A). Motility was confirmed at the
single cell level when these four variants were tested in tunnel slides. Tethered cell assay
measurement of cell speed revealed a significant reduction in cell speed for all the motile
variants including M34I (p < 0.001, T test) (Fig. 3B). The M34I variant was also found to have
a significant decrease in clockwise (CW) bias (p < 0.05) when compared to M34A indicating
an increased propensity for counterclockwise (CCW) motor runs over tumbles
(Supplementary Fig. 4), which may explain the increased size of swim rings for this variant in
spite of the reduced motility. Interestingly, M34I was able to swim in higher density agar
(0.4%) better than WT or any other variant, suggesting a change in torque generation in the
mutant stator (SI Fig. 5).

A comparison of the mean 2D BUDE profiles of all motile and non-motile variants suggested
that the stability of the coiled-coil around the interface angle of 28°, where M34 plays a key
role in KIH interaction, was crucial for function (Fig. 3C). The influence on the total energy
from mutation at that site, adjacent to the catalytic aspartate residue (D32), appears to be
correlated with a motile phenotype. When comparing the 2D BUDE profiles of each MotB
variant tested *in vivo*, we noticed a similarity in the shapes of the BUDE profiles of the
motile variants, whereas the profiles of the non-motile variants diverged substantially (Fig.
3D). Aligning the profiles with respect to their peak values (BUDE^{max}) revealed additional
differences between motile and non-motile profiles than the strength of interaction at the
28° configuration alone. The BUDE gap between the minimum value of each profile and the
peak corresponding to the 44° configuration appeared to be similar across the motile
variants.

Further comparison of the 2D profiles of all point mutants suggested a pattern of specific
interface angles, or configurations, associated with motility (SI Fig. 6). All the *Ec*MotB
variants found to be motile had the total energies associated with the 66° (cross), 28°
(square), 56° (circle) and 44° (triangle) configurations in ascending order of magnitude. The
exceptions to this pattern were the single mutants A31I, F41I and M38G, which featured
destabilized 44°, 28°, and 28° configurations, respectively, and the double mutants A31I
M34I and A31I M38A which featured a destabilized 44° configuration. This suggested that
the relative stability of the 28°, 56° and 64° configurations play a role in determining the
function of MotB and that this pattern may reflect the order of configurations undertaken
by WT MotB during its gating.

We captured the features associated with motility variants by defining two metrics, *f1* and
f2 (Fig. 2E), where *f1* = BUDE$^{56°}$ - BUDE$^{28°}$ and *f2* = BUDE^{min} – BUDE^{max} (Fig. 3E). The first
metric, *f1*, is the BUDE energy gap at an interface angle of 28°, associated with motile
variants (Fig 3C), and the reference interface angle of 56°, predicted to form a KIH
interaction with the conserved residue F41 (Fig. 2B, 'circle' config). Residue F41 appeared to
have the more consistent value across the profiles of all variants (SI Fig. 7). With the *f2*
metric we measured the energy gap between the minimum and maximum BUDE values
across the profile, or the energy barrier separating the energy wells. Separating all variants
on the basis of these two metrics revealed a cluster of optimal *f1* and *f2* values for the
motile variants of *E. coli* MotB (Fig. 3F). These results suggest that a combination of

energetic parameters contained within the 2D profiles could be descriptive of functional
EcMotB variants and that the relative stability of each configuration needs to be maintained
to operate the E. coli stator.

We also searched for other metrics that could cluster and separate motile and non-motile
sequences to facilitate the prediction of motile variants of EcMotB from their 2D profile
models. We defined $f3 = [(BUDE^{56°} - BUDE^{28°}) - (\Delta BUDE^{56°} - \Delta BUDE^{28°})]$ and observed that this
metric was correlated with motility only when yielding a positive value, where $\Delta BUDE$ is the
difference in BUDE energy between the mutant and WT MotB at the given interface angle.
Additionally, we defined $f4 = (BUDE^{min} - BUDE^{max} - BUDE^{max}) = (BUDE^{min} - 2*BUDE^{max})$. This
considered not only the range (the difference between min and max) but also the maximum
value as an indicator of the magnitude of the BUDE value (allowing, e.g., to distinguish
between two samples with range of 200, but with a $BUDE^{max}$ of -500 or -400 respectively).
This quantity, $f4$, was found to correlate with motility when positive. We observed that both
$f3$ and $f4$ were positive only in the functional variants of E. coli MotB studied here
(Supplementary Fig. 8).

Finally, we explored a larger set of homologous MotB sequences to determine whether our
observations were applicable to other extant motile bacteria. We generated a 91 member
phylogeny of MotB using RAxML[20] across diverse clades (Fig. 4A, SI Fig. 9, 10) and carried
out the same BUDE analysis on 70 representative MotB sequences across the phylogeny
(Supplementary Fig. 9A, 10) to produce a 2D BUDE energy profile for each of the 20
residues-long TM domain modelled as a coiled-coil (SI Fig. 9B). To compare the energy
profiles to each other and to the phylogenetic clustering, we translated each BUDE profile
relative to its $BUDE^{max}$ value (Fig. 4B). This analysis of the energy profiles distinguished 7
distinct groups (Fig. 4C) that matched the phylogenetic clades (Fig. 4A), and each displaying
one or more conserved amino acid residues amongst the representative positions a and d of
their heptad repeats, evidenced by their consensus logos (Fig. 4C). These groups clustered
around specific values of the $f1$ and $f2$ metrics (Fig. 4D). Out of these 70 sequences, 66 were
found to have $f1 > 0$, while only a subset had $f2 > 0$ (SI Fig. 9C). Clusters were separated by
the identity of their residue 38 (position a) and shared the conserved D32 and F41 (d)
residues. Positions 31 (a), 34 (d) and 45 (a) also displayed unique residues in specific clusters
(Fig. 4A, 4C).

Discussion

In this study we constructed several MotB stator variants based on *in silico* designs following coiled-coil design principles and measured their capacity to drive swimming *in vivo* in *E. coli*. Because the residues targeted by mutagenesis were at the hydrophobic interface of the coiled-coil structure, we assumed that the effects of amino acid replacement would be confined to MotB and its dimerization and thus restricted our analyses to symmetric configurations of the MotB dimer. In other words, we assumed that movement (i.e. rotation) of one MotB subunit was always accompanied by the same movement of the second subunit. We sought to examine how motility of the overall stator complex was disrupted when assembly of the coiled-coil via knobs-in-holes (KIH) was altered. Altogether, we focused on variations of the interface angle between the two MotB alpha-helices, which correspond to the rotation of the alpha-helices relative to each other. This type of rotation is known as a register shift, as the interface angle is related to the coiled-coil register, and register shifts have been shown elsewhere to underpin biological processes such as dynein recruitment[21,22] and the activity of proteasomal ATPase[23].

We tested a mutant with the intent to disrupt motility (M38G), by disrupting the KIH interactions between the coils. A gain-of-function variant (M34I), designed to have a stronger coiled-coil interaction, demonstrated that coiled-coil stability could be linked to improved motility, but we also observed that "tightening" the coiled-coil at the wrong knob could disrupt function entirely. For example, mutating the central knob towards a new type of KIH interaction (M38I mutant) was predicted to stabilize only the 64° conformation. This mutation yielded a non-motile variant. In contrast, energy calculations indicated the M34I mutation could increase dimer stability for all analysed coiled coil configurations, including the 64° confirmation, and yet this mutation yielded a motile variant. The reason for improved swimming of this variant on swim plates remains unclear because the motor performance appeared to be reduced compared to WT in single-cell assays (both in tethered and free-swimming assays). In single-cell assays, M34I was indistinguishable from the M34A variant, while M34A displayed WT phenotype on swim plates in contrast to M34I.

Swimming of M34I, WT, M34A, M38A and F41A correlated with shared properties of their BUDE energy profiles. Although a bulk comparison of our motile and non-motile variants showed a difference in their BUDE magnitude at a specific configuration on the profile (the 28° configuration) we sought to identify other features in the BUDE profile to use for classifying motile and non-motile classes. We defined two metrics, *f1* and *f2,* to differentiate between the two classes and the effect of each of point mutant on motility. In this framework, M38A maintains the energy profile shape of the WT; M38I reduces $BUDE^{max}$ and abolishes function; F41A destabilizes the 56° configuration, maintaining a positive value for *f1* and therefore a functional stator, while F41I stabilizes it and abolishes function by lowering *f1* below zero; W45A increases $BUDE^{max}$ beyond a functional level and W45I destabilizes the 28°configurations, thus lowering *f1* to a negative value. A31I reduces $BUDE^{max}$ compared to WT and abolishes function.

Using *f1* as an indicator of functionality in a stator implies that the interaction is dominated by the knobs at position 34 and the conserved residue F41. We thus speculate that a

transition between these two configurations is essential for the function of the stator. Previous work involving chemical cross-linking between MotB coiled-coil residues suggested that the TM structure is flexible and able to exchange the residues at the CC interface during a rotation of ~36°[19]. In line with those considerations, we found that BUDE profile features at all four configurations (28°, 44°, 56° and 64° configurations in EcMotB) were predictive of stator functionality, that is, motility on swim plates.

The f1 and f2 parameters were useful also in describing and clustering extant MotBs. Indeed, it may be that these clusters represent families of MotB homologs that are related in their functional characteristics. Notably, V. alginolyticus PomB and E. coli MotB, of which functional chimeras exist[24], belong to the same cluster (green, Fig. 4). These two stators are known to differ in their power source (Na^+ and H^+ respectively) and differ in the identity of their homologous residue 45 (L45 for Na^+, W45 for H^+) while showing similar 2D profiles. Another notable cluster is the one shared by B. subtilis, C. sporogenes and C. jejuni (black, Fig. 3). These three species are sources of the most detailed cryo-EM structures of the $MotA_5B_2$ stator complex to date. It is possible that these complexes were more stable and thus easier to purify and resolve due to stronger MotB coil interaction than the E. coli or V. alginolyticus homologs.

We noted a linear relationship between the hydrophobicity of residue 38 with the $BUDE^{max}$ magnitude. MotB clusters displaying F, L or M at position 38 had a proportionally high $BUDE^{max}$ peak in their profiles, with F38 being the highest and M38 the lowest (SI Fig. 9B). The height of this peak could be related to the presence of stable intermediate conformations available during the transition between the two most stable configurations at 28° and 64°.

Elsewhere we have extensively examined large phylogenies of MotB[25]. Here we calculated a small phylogeny that was representative of wider flagellar diversity to compare how energetics clustered across a model of evolutionary history. Our MotB classes obtained from clustering similar energetic profiles closely overlap clades from our phylogeny obtained using traditional tree-building approach (Fig. 3A). This suggests that conservation of sequence identity is closely linked to conservation of functional traits, and indeed it is notable that BUDE free energy profiles calculated across only the 20 transmembrane residues so naturally fell into clusters that matched those generated by sequence similarity from phylogenetic consideration across the full length MotB. Together, these results show that cluster-based comparison of BUDE free energies can recapitulate phylogenetic clades and shows that there is evolutionary evidence for conservation of distinct stable states. This clustering based on MotB-MotB dimer stability could be further examined in greater detail, for example, with regard to species that have specific high-torque operating environments (e.g., Spirochaetes sp.). These species typically have large rotors to generate larger torques[5,26–28], but there could be signatures of conservation of increased stator stability elsewhere that may correlate with high-torque or viscous operating conditions.

Based on the model proposed by Rieu et al.[10], we describe a model of mechanochemical gating that includes the conformational changes in the MotB TM domain studied here (Fig.5). Our model assumes that the rotation of the MotA pentamer can occur only if both

catalytic aspartates in MotB are protonated. We propose that a single protonated aspartate,
engaging MotA with MotB, couples the synchronized rotation of both $MotA_5$ and $MotB_2$ by
36° clockwise, and is then released into the cytoplasm after the entry of a second proton
binding onto the other disengaged aspartate. Our model also assumes that the binding of a
proton to the disengaged aspartate initiates the power stroke, which then occurs as both
aspartates are protonated as postulated by Santiveri et al.[7,10] The newly protonated site
triggers a conformational change at the MotA-MotB interface that promotes the
engagement of the protonated aspartate with MotA at the second catalytic aspartate, the
cytoplasmic pore exit site, as the complex rotates 36° clockwise.

In the Oster et al.[4] model, a 'kink' occurs in TM3 of MotA upon H^+ binding to the aspartate
and when MotB engages with TM3/4 of MotA this results in a swivelling motion of the TM3
of MotA that is the basis of the power stroke mechanism. Our data suggests that MotB
might be coupled with the movement of MotA during its swivelling motion (a CW rotation)
and switch between at least two of the stable configurations studied here, the 28° and 64°
ones. Our model takes into account the 5:2 stoichiometry detected in crystal structures and
considers only one catalytic site at the time to be participating in a power stroke. The
concerted CW rotation of the MotB TMs during the power stroke facilitates the alternating
engagement of MotA subunits separated by $2/5^{th}$ of a revolution, while maintaining MotB
always in a parallel coiled coil state and without the need for vertical displacement of the
MotB coils during the proton release stage. The activation of individual MotA subunits
during the power stroke is reminiscent of the model proposed by Xing et al. but involving
the alternating translocation of protons across the stator[29].

Whilst we cannot determine the exact motions of the MotB TM dimer during the switch
between its two most stable configurations, we suspect that the relative stability of the
intermediate configurations (like the 56° one) is important for the completion of a 36°
rotation of the interface angle of MotB. It is also possible for other coiled-coil parameters
(pitch and radius) to change during this rotation which may stabilise the coiled coil at
configurations considered inaccessible (like the 44° configuration). Mutations that further
stabilise certain configurations, for example a more hydrophobic KIH at position 38 of the
EcMotB, would increase the height of the $BUDE^{max}$ peak (Fig.4). This might interfere with the
establishment of these intermediate conformations which need to be crossed during the
rotation from the 28° to the 64° configuration.

The function of the flagellar stator complex clearly depends not only on the pore lining
residues between MotA and MotB and torque-generating interactions between MotA and
FliG[9,30]. Our overall conclusion is that it also depends on the degree of interaction between
the two TM domains of MotB that assemble as a coiled-coil at the core of the stator and the
conformational states the dimer can access. The residues responsible for knob-in-hole
interactions at the dimerization interface of E. coli MotB, and possibly other extant
homologs, appear to tolerate onlyl those mutations which do not shift their coiled-coil
energetics too far from wild-type. Consideration of these energetics will improve our ability
to engineer mutant MotB subunits, chimeric stators, or synthetic combinations of multiple
flagellar motor proteins.

Figure 1. Modelling the transmembrane domain of MotB as a dimeric coiled-coil.

A) The stator subunit of the bacterial flagellar motor (BFM) of *E. coli* (MotAB) drives the rotation of the large rotor complex by converting the proton motive force into mechanical torque. The stator unit MotA$_5$MotB$_2$ acts as an ion channel and each MotA has 4 transmembrane (TM) domains while MotB has a single TM domain and exists as a dimer in the centre of the complex. B) The crystal structure of *B. subtilis* MotA$_5$B$_2$[PDB 6YSL] (left) was used as the starting point for *E. coli* MotB TM modelling. The MotB dimer is shown in pink. C) A CCBuilder model of the homologous sequence from *E. coli* was generated (blue) and aligned to the reference *Bs*MotB dimer (purple). A coiled-coil model in the *e* register was selected based on the orientation of the catalytic aspartates D32 (*E.coli*) and D24 (*B. subtilis*) (middle, dotted circles in the side and bottom views of the TM domain). Residues A31, M34, M38, F41 and W45 were found to be at the coiled-coil interface (right), corresponding to positions *a* and *d* of the heptad repeat able to participate in knob-in-kole (KIH) interactions (labelled in red on the atomic model and heptad diagram underneath). An alignment of the two homologous amino acid sequences and the selected register is shown on the bottom-right, with *E.coli* residues at the *a* and *d* positions highlighted in red and numbered.

Figure 2. Characterization of the MotB TM coiled-coil interaction energy. (A) Free energy
(BUDE) between MotB dimer coils represented as heatmap indicating BUDE value vs
interface angle and radius. Four configurations are highlighted by symbols. (B) Atomic
models of the MotB coiled-coil for four configurations (28˚, 44˚, 56˚ and 64˚, identified by
the square, triangle, circle and cross symbols respectively). The main amino acid residue
responsible for KIH interactions are labelled in red. (C) 2D BUDE profile calculated from
mean across ten radii values from (A). (D) predicted BUDE energy changes at each of the
four configurations (symbols) after amino acid replacement at positions *a* and *d* of the MotB
coil with amino acids alanine, (E) ΔBUDE (BUDE^mut – BUDE^wt) for each of four configurations.
The BUDE value calculated at interface angles 28°, 44° 56° and 64° is calculated as the
average of the two neighbouring data points above the black horizontal bars. Substitutions
are indicated by colour (Ala, black) or isoleucine (Ile, red).

Figure 3. Testing coiled-coil variants in _E.coli_ and parametrization of 2D BUDE profiles.

A) Swim plate assay of _E. coli_ ΔMotAB + pDB108 expressing WT MotA and either WT (top) or

mutant MotB's on LB agar 0.3% supplied with 0.1 % arabinose and Chloramphenicol after 24

hr incubation at 30'C. See supplementary figure 3 for all variants tested in this study. B)

Single _E. coli_ cell swimming speed measured in motility buffer. The 10 fastest cells for each

mutant are plotted here. Bars indicate Mean speed (µm/s) + SD. C) Average 2D BUDE plots

of motile (green) vs non-motile (black) MotB variants. The KIH present in MotB in the 28°

configuration is highlighted in red on the atomic model in the dashed box. Bars indicate

Mean + SD. D) Individual 2D BUDE plots of motile (green) vs non-motile (black) MotB

variants, translated according to their respective $BUDE^{max}$ values. E) The *f1* and *f2*

parameters visualized on the 2D BUDE profile of *E. coli* WT MotB. F) Individual variants

plotted on a graph (X = *f1* and Y = *f2*). The cluster of motile variants is highlighted in green.

Figure 4. BUDE analysis of extant MotB sequences.

A) Phylogeny of extant, full-length MotBs. Clades are labelled by colored boxes
corresponding to the clusters highligted in B. A representative member of each clade, or
representative bacterial phylum, are shown next to each box. The conserved amino acids
indicative of each clade are shown next to each box and numbered according to *E.coli*
residues. A detailed phylogeny is provided in SI Figure 10. B) 2D BUDE plots of extant WT

MotB variants, translated by their $BUDE^{max}$ values. C) Consensus logos for each plot in B.
Each logo is inscribed in a box matching the colour of its respective plots in A. Knob-in-hole
residues at positions homologous to *Ec*MotB 31, 34, 38, 41 and 45 are highlighted in vertical
grey boxes. D) Clustered populations of extant MotB stators plotted on a graph (X = *f1* and Y
= *f2*). The formulas used to calculate *f1* and *f2* for each homolog are provided below the X
axis.

Figure 5. Rotary model showing the role of MotB coiled-coil conformations in the mechanochemical cycle of the stator complex.

A) Section of the *Bs*MotAB [PDB 6YSL] structure highlighting the central MotB dimer (blue) and transmembrane helices TM3 and TM4 of MotA (grey) interfacing the MotB dimer. A sphere model (red) indicates the position and orientation of the catalytic aspartate in MotB. A cartoon representation of the crystal structure and legend are shown on the right. The dashed lines below separate the stator cross-section in ten identical 36° sectors. B) MotB TM BUDE fluctuations between three configurations during the gating cycle of the stator. The red arrows indicate binding or unbinding of a proton at the left or right aspartate (H- or

-H, respectively). A yellow star indicates the power stroke and 36° clockwise rotation of the
stator. Configuration symbols and representative KIH interactions are also shown on the
right of the model. C) Six stages of the mechanochemical cycle model described in B. A
proton is transferred into the cytoplasm through an exit pore immediately after a power
stroke (1). A change in the MotB TM configuration follows, which rotates 36°
counterclockwise from the 28° (1) to the 64° configuration (3), via the 56° configuration (2).
Once in the 64° configuration, the aspartate on the left subunit engages MotA at the TM3-4
interface in proximity of the cytoplasmic exit pore (3). A proton then enters from the
opposite periplasmic entry pore to bind the unprotonated aspartate causing a
conformational change of the coiled-coil and the coordinated rotation of the MotB dimer
and MotA pentamer by 36° clockwise (a power stroke). This rotation restores the coiled-coil
to the 28° configuration and the engaged ion is released into the cytoplasm. The cycle
repeats at the previously disengaged aspartate in (4), (5) and (6) eventually resulting in a
new power stroke and the return to stage (1).

Materials and Methods

Structural modelling of coiled-coils.

The *B. subtilis* MotA$_5$B$_2$ structure (PDB:6YSL)[8] was used as a reference to model the
transmembrane domain of *E. coli* MotB as coiled-coil. We used ISAMBARD to create a
specification whereby a parallel dimeric coiled-coil backbone structure could be generated
from three parameters: the interface angle, the radius, and the pitch. The coiled-coil
backbone structure for the amino acid sequence IAYADFMTAMMAFFLVMWLI
(corresponding to the *E. coli* MotB residues 28-47) was optimized using the genetic
algorithm evolutionary optimizer from ISAMBARD and SCWRL 4.0 for the optimization of
side chain orientations to minimize the total energy of the system as calculated by the BUDE
force field. Optimizations were performed using seven different initial interface angle values
which corresponded to the seven possible registers of a coiled-coil. The seven optimized
coiled-coil models were aligned to the resolved structure of the MotB dimer stalk in PDB
structure 6YSL. The alignment revealed that the best initial interface angle that resulted in a
good structural match after optimization to the MotB stalk was 51.4°, which corresponds to
position *e* for the first residue in the amino acid sequence. The optimized parameter values
were 4.54 Å radius, 232.03 Å pitch, and 74.85° interaction angle.

Calculation of the energy landscapes of the MotB coiled-coils

We used in-house Python scripts to generate a list of 221 sequence variants of the amino
acid sequence IAY**A**DF**M**TAM**M**AFFLVM**W**LI where residues A31, M34, M38, F41, W45 were
systematically mutated to alanine in every permutation; separately the sequence was
mutated with isoleucine (for residues A31, M38, and W45), asparagine (for residue M38),
and leucine (for residues M34 and F41) in every permutation. These mutations follow
coiled-coil design principles to weaken or strengthen dimeric coiled-coils. Finally, additional
MotB mutants were added to the list (SI Table 1). For each sequence variant, we used
ISAMBARD to generate coiled-coil backbone structural models with radii values ranging
from 4 to 6 Å (in 0.1 Å increments) and interface angle values ranging from 10 to 86° (in 4°
increments). The pitch was kept constant to a value of to 150 Å. These ranges of values
encompass the optimized parameters that best matched the reference structures. The
sidechain orientations were optimised using SCWRL 4.0 as part of the ISAMBARD script. For
each conformational variant, the total energy of the system was calculated using the BUDE
forcefield. The three-dimensional data sets (BUDE total energy as a function of interface
angle and radius) were plotted as heatmaps using in-house Python scripts and Matplotlib
library. Two-dimensional datasets were generated by integrating the total energy values for
a subset of the radii values (4 to 5 Å) which contained the stability wells and instability peaks
of interest. The integrated total energy values were plotted as a function of the interface
angles. In some analyses, we focused on four regions in the three-dimensional dataset for
comparison of the sequence variants (centred on list combinations of radii and interface
angles) and a representative energy value for each of the four regions was calculated by
averaging two neighbouring points from the two-dimensional dataset (see Fig. 2C). For the
identification of knob-into-hole interactions, MotB was modelled as a dimeric coiled-coil

starting at register *e* using CCBuilder 2.0, which evaluates the presence of knob-into-holes
using SOCKET[31].

E. coli strains, plasmids, and culture media

E. coli strain RP437 Δ*motAB* was prepared using the No-Scar method[32]. MotA and MotB
were expressed from plasmid pDB108 (Cm$^+$) after induction with 0.1 mM Arabinose. FliC-
sticky *E. coli* RP437 cells (Δ*motAB* Δ*cheY* Δ*pilA fliC*st)[33] were used to perform the tethered
cell assay experiments. Liquid cell culturing was done using LB broth (85 mM NaCl, 0.5%
yeast extract, and 1% Bacto tryptone). Point mutations in *motB* on the pDB108 plasmid
were generated using the QuikChange™ technique and verified by Sanger sequencing. A list
of primers used is provided in Supplementary Table 2. Cells were cultured on agar plates
composed of LB broth and 1% Bacto agar (BD Biosciences, U.S.A.). Swim plate cultures were
performed on the same substrates adjusted for agar content (0.3% Bacto agar).

Single cell swimming assay and tethered cell assay.

Free-swimming cells were grown overnight in TB buffer (85 mM NaCl, 1% Bacto tryptone) at
30°C to OD$_{600}$ of ~0.5 then washed three times in 1 ml MB buffer (85 mM NaCl and 10 mM
KPi, pH=7.0) before resuspension in 500 µl of MB buffer and imaging in a tunnel slide. The
tethered cell assay was performed as previously described[34]. Briefly, 1 ml of cells [optical
density at 600 nm (OD600) = 0.5] grown in TB buffer was sheared by passing the cell
suspension through a 26-gauge syringe needle 30 times. These cells were then washed
three times in 1 ml MB and lastly resuspended in 500 µl of the same buffer. Then, 20 µL of
suspension was loaded into a tunnel slide prefilled with MB buffer and incubated for 10 min
at room temperature. The unbound cells were then removed from the tunnel slide by
washing with a total of 200 µl of MB buffer (~10 times the tunnel slide volume) and then
loaded onto a phase contrast microscope for video recording. Time-lapse videos were
recorded at 40x magnification on a phase contrast microscope (Nikon). Time-lapse videos
were collected using a camera (Chameleon3 CM3, Point Grey Research) recording 20-s-long
videos at 20 frames/s. Experiments involving single-cell tracking were recorded at 60
frames/s. A custom LabView software[35] was used as previously reported to estimate specific
rotational parameters of the tethered cells such as rotation frequency (speed), clockwise
and counterclockwise bias, switching frequency, and speed of swimming cells.

REFERENCES

1. Nakamura, S., and Minamino, T. (2019). Flagella-driven motility of bacteria. Biomolecules *9*, 279. 10.3390/biom9070279.

2. Mondino, S., San Martin, F., and Buschiazzo, A. (2022). 3D cryo-EM imaging of bacterial flagella: Novel structural and mechanistic insights into cell motility. Journal of Biological Chemistry *298*, 102105. 10.1016/j.jbc.2022.102105.

3. Nirody, J.A., Sun, Y.-R., and Lo, C.-J. (2017). The biophysicist's guide to the bacterial flagellar motor. Advances in Physics: X *2*, 324–343. 10.1080/23746149.2017.1289120.

4. Mandadapu, K.K., Nirody, J.A., Berry, R.M., and Oster, G. (2015). Mechanics of torque generation in the bacterial flagellar motor. Proc Natl Acad Sci USA *112*, E4381–E4389. 10.1073/pnas.1501734112.

5. Guo, S., and Liu, J. (2022). The Bacterial Flagellar Motor: Insights Into Torque Generation, Rotational Switching, and Mechanosensing. Frontiers in Microbiology *13*.

6. Hu, H., Santiveri, M., Wadhwa, N., Berg, H.C., Erhardt, M., and Taylor, N.M.I. (2022). Structural basis of torque generation in the bi-directional bacterial flagellar motor. Trends in Biochemical Sciences *47*, 160–172. 10.1016/j.tibs.2021.06.005.

7. Santiveri, M., Roa-Eguiara, A., Kühne, C., Wadhwa, N., Hu, H., Berg, H.C., Erhardt, M., and Taylor, N.M.I. (2020). Structure and Function of Stator Units of the Bacterial Flagellar Motor. Cell *183*, 244-257.e16. 10.1016/j.cell.2020.08.016.

8. Deme, J.C., Johnson, S., Vickery, O., Muellbauer, A., Monkhouse, H., Griffiths, T., James, R.H., Berks, B.C., Coulton, J.W., Stansfeld, P.J., et al. (2020). Structures of the stator complex that drives rotation of the bacterial flagellum. Nature Microbiology, 1–12. 10.1038/s41564-020-0788-8.

9. Hu, H., Popp, P.F., Santiveri, M., Roa-Eguiara, A., Yan, Y., Liu, Z., Wadhwa, N., Wang, Y., Erhardt, M., and Taylor, N.M.I. (2022). Mechanisms of ion selectivity and rotor coupling in the bacterial flagellar sodium-driven stator unit. 2022.11.25.517900. 10.1101/2022.11.25.517900.

10. Rieu, M., Krutyholowa, R., Taylor, N.M.I., and Berry, R.M. (2022). A new class of biological ion-driven rotary molecular motors with 5:2 symmetry. Frontiers in Microbiology *13*.

11. Sharp, L.L., Zhou, J., and Blair, D.F. (1995). Tryptophan-scanning mutagenesis of MotB, an integral membrane protein essential for flagellar rotation in Escherichia coli. Biochemistry *34*, 9166–9171. 10.1021/bi00028a028.

12. Zhang, X.Y.-Z., Goemaere, E.L., Seddiki, N., Célia, H., Gavioli, M., Cascales, E., and Lloubes, R. (2011). Mapping the interactions between *Escherichia coli* TolQ transmembrane segments. J. Biol. Chem. *286*, 11756–11764. 10.1074/jbc.M110.192773.

536 13. Lupas, A.N., and Bassler, J. (2017). Coiled Coils – A Model System for the 21st Century.
537   Trends in Biochemical Sciences *42*, 130–140. 10.1016/j.tibs.2016.10.007.

538 14. Fletcher, J.M., Boyle, A.L., Bruning, M., Bartlett, G.J., Vincent, T.L., Zaccai, N.R.,
539   Armstrong, C.T., Bromley, E.H.C., Booth, P.J., Brady, R.L., et al. (2012). A Basis Set of *de*
540   *Novo* Coiled-Coil Peptide Oligomers for Rational Protein Design and Synthetic Biology.
541   ACS Synth. Biol. *1*, 240–250. 10.1021/sb300028q.

542 15. Crick, F.H.C. (1953). The Fourier transform of a coiled-coil. Acta Cryst *6*, 685–689.
543   10.1107/S0365110X53001952.

544 16. McIntosh-Smith, S., Price, J., Sessions, R.B., and Ibarra, A.A. (2015). High performance *in*
545   *silico* virtual drug screening on many-core processors. The International Journal of High
546   Performance Computing Applications *29*, 119–134. 10.1177/1094342014528252.

547 17. Wood, C.W., Bruning, M., Ibarra, A.Á., Bartlett, G.J., Thomson, A.R., Sessions, R.B.,
548   Brady, R.L., and Woolfson, D.N. (2014). CCBuilder: an interactive web-based tool for
549   building, designing and assessing coiled-coil protein assemblies. Bioinformatics *30*,
550   3029–3035. 10.1093/bioinformatics/btu502.

551 18. Wood, C.W., and Woolfson, D.N. (2018). CCBuilder 2.0: Powerful and accessible coiled-
552   coil modeling: Powerful and Accessible Coiled-Coil Modeling. Protein Science *27*, 103–
553   111. 10.1002/pro.3279.

554 19. Braun, T.F., and Blair, D.F. (2001). Targeted disulfide cross-linking of the MotB protein of
555   Escherichia coli: evidence for two H(+) channels in the stator Complex. Biochemistry *40*,
556   13051–13059. 10.1021/bi011264g.

557 20. Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-
558   analysis of large phylogenies. Bioinformatics *30*, 1312–1313.
559   10.1093/bioinformatics/btu033.

560 21. Choi, J., Park, H., and Seok, C. (2011). How does a registry change in dynein's coiled-coil
561   stalk drive binding of dynein to microtubules. Biochemistry *50*, 7629–7636.
562   10.1021/bi200834k.

563 22. Noell, C.R., Loh, J.Y., Debler, E.W., Loftus, K.M., Cui, H., Russ, B.B., Zhang, K., Goyal, P.,
564   and Solmaz, S.R. (2019). Role of Coiled-Coil Registry Shifts in the Activation of Human
565   Bicaudal D2 for Dynein Recruitment upon Cargo Binding. J. Phys. Chem. Lett. *10*, 4362–
566   4367. 10.1021/acs.jpclett.9b01865.

567 23. Snoberger, A., Brettrager, E.J., and Smith, D.M. (2018). Conformational switching in the
568   coiled-coil domains of a proteasomal ATPase regulates substrate processing. Nat
569   Commun *9*, 2374. 10.1038/s41467-018-04731-6.

570 24. Nishino, Y., Onoue, Y., Kojima, S., and Homma, M. (2015). Functional chimeras of
571   flagellar stator proteins between E. coli MotB and Vibrio PomB at the periplasmic region
572   in Vibrio or E. coli. Microbiologyopen *4*, 323–331. 10.1002/mbo3.240.

25. Islam, M.I., Lin, A., Lai, Y.-W., Matzke, N.J., and Baker, M.A.B. (2020). Ancestral Sequence
Reconstructions of MotB Are Proton-Motile and Require MotA for Motility. Front.
Microbiol. *11*. 10.3389/fmicb.2020.625837.

26. Beeby, M., Ferreira, J.L., Tripp, P., Albers, S.-V., and Mitchell, D.R. (2020). Propulsive
nanomachines: the convergent evolution of archaella, flagella and cilia. FEMS
Microbiology Reviews *44*, 253–304. 10.1093/femsre/fuaa006.

27. Beeby, M., Ribardo, D.A., Brennan, C.A., Ruby, E.G., Jensen, G.J., Hendrixson, D.R., and
Hultgren, S.J. (2016). Diverse high-torque bacterial flagellar motors assemble wider
stator rings using a conserved protein scaffold. Proceedings of the National Academy of
Sciences of the United States of America *113*, E1917–E1926. 10.1073/pnas.1518952113.

28. Carroll, B.L., and Liu, J. (2020). Structural Conservation and Adaptation of the Bacterial
Flagella Motor. Biomolecules *10*, 1492. 10.3390/biom10111492.

29. Xing, J., Bai, F., Berry, R., and Oster, G. (2006). Torque–speed relationship of the
bacterial flagellar motor. Proceedings of the National Academy of Sciences of the United
States of America *103*, 1260–1265. 10.1073/pnas.0507959103.

30. Islam, M.I., Ridone, P., Lin, A., Michie, K.A., Matzke, N.J., Hochberg, G., and Baker, M.A.
(2022). Ancestral reconstruction of the MotA stator subunit reveals that conserved
residues far from the pore are required to drive flagellar motility. 2022.10.17.512626.
10.1101/2022.10.17.512626.

31. Walshaw, J., and Woolfson, D.N. (2001). SOCKET: a program for identifying and
analysing coiled-coil motifs within protein structures11Edited by J. Thornton. Journal of
Molecular Biology *307*, 1427–1450. 10.1006/jmbi.2001.4545.

32. Reisch, C.R., and Prather, K.L.J. (2015). The no-SCAR (Scarless Cas9 Assisted
Recombineering) system for genome editing in Escherichia coli. Sci Rep *5*, 15096.
10.1038/srep15096.

33. Sowa, Y., Rowe, A.D., Leake, M.C., Yakushi, T., Homma, M., Ishijima, A., and Berry, R.M.
(2005). Direct observation of steps in rotation of the bacterial flagellar motor. Nature
437, 916–919. 10.1038/nature04003.

34. Islam, M.I., Bae, J.H., Ishida, T., Ridone, P., Lin, J., Kelso, M.J., Sowa, Y., Buckley, B.J., and
Baker, M. a. B. (2021). Novel Amiloride Derivatives That Inhibit Bacterial Motility across
Multiple Strains and Stator Types. J Bacteriol *203*, e0036721. 10.1128/JB.00367-21.

35. Ishida, T., Ito, R., Clark, J., Matzke, N.J., Sowa, Y., and Baker, M.A.B. (2019). Sodium-
powered stators of the bacterial flagellar motor can generate torque in the presence of
phenamil with mutations near the peptidoglycan-binding region. Molecular
Microbiology *111*, 1689–1699. https://doi.org/10.1111/mmi.14246.

608