Title page

Movements and habitat selection of a marsupial carnivore in a modified landscape

Authors

Evie M Jones[1,3], Amelia J Koch[1,2], James M Pay[1], Dydee F Mann[2], Menna E Jones[1], Rodrigo

K Hamede[1]

[1]School of Natural Sciences, University of Tasmania, Private Bag 55, Hobart, TAS, 7001,

Australia.

[2]Forest Practices Authority, 30 Patrick St, Hobart, TAS, 7001, Australia.

[3]Corresponding author: evie.jones@utas.edu.au

Abstract

Landscape modification is a major threat to carnivores worldwide, but modified landscapes

can also provide important habitat for these species, as protected areas alone are insufficient.

Understanding how carnivores move through modified landscapes, such as production

forests, can inform management strategies to improve the value of these landscapes to

carnivores. Little is known about habitat selection by marsupial carnivores in production

forests, where they occupy a similar ecological niche to their more well-studied eutherian

counterparts. We used GPS tracking, Hidden Markov Models, and Manly's selection ratios to

identify the habitat selected in 3 movement states indicating resting/feeding, foraging, and

travel by the largest marsupial carnivore, the Tasmanian devil (*Sarcophilus harrisii*), in a

timber plantation-dominated landscape. Devils did not show preferences for any of native

forest, native grassland, and plantation in any movement state. Within plantations, devils

preferred a plantation age of 4–7 years (selection ratio [*wi*] = 1.52) and slightly avoided older

plantations (8–13 years *wi* = 0.74, 14+ years *wi* = 0.65). Devils preferred roads (state 1: *wi* =

2.71, state 2: *wi* = 2.48, state 3: *wi* = 2.97) and plantation edges (state 1: *wi* = 2.38, state 2: *wi*

= 2.24, state 3: *wi* = 2.78) in all movement states, and moved faster on roads (0.06 [95% CI

0.04 to 0.09] m.sec^{-1}) and edges (0.08 [0.06 to 0.10] m.sec^{-1}) than away from them, indicating

they use them for foraging and travel. No measured habitat variables influenced devil home

range size. To support devils in plantation landscapes, we recommend maintaining a

heterogeneous landscape of different plantation ages and native remnants and reducing the

risk of vehicle collisions by minimizing forestry traffic at night. Tasmanian devils share

similar adaptable traits to generalist eutherian carnivore species in their use of modified

landscapes. Plantations can provide valuable habitat for this and other threatened predator

species.

Keywords

plantation, Tasmanian devil, forestry, GPS telemetry, roads, edges, foraging, production

forest

Background

With the ongoing global loss of untouched forests (1), there is increasing focus on finding

strategies to preserve and also enhance biodiversity in human-modified landscapes (2).

Modified habitats such as production forests can make an important contribution to

maintaining forest ecosystems, as protected areas alone are insufficient (3). For instance,

replacing native forests with timber plantations (henceforth 'plantations') negatively affects

forest biodiversity (4), but plantations often contain higher biodiversity than other land uses

such as agriculture, and can decrease harvesting pressure on native forest (2). Conservation

strategies in production forests should be informed by ecological research to maximize

effectiveness and cost-efficiency (5).

Carnivores are especially sensitive to modified landscapes (6) and knowledge of their habitat

use within these landscapes can inform practices to sustain their populations. This is

particularly important as carnivores have key ecological effects such as limiting populations

of prey and smaller predators (7), effects that are dampened in modified landscapes (8).

Improving carnivore conservation in anthropogenic landscapes such as production forests can

therefore benefit entire ecosystems in these landscapes. Carnivore species can differ

markedly in both their habitat selection within plantations (9,10) and their responses to

forestry operations (11). While the use of production forests by eutherian carnivores has been

well studied (12), little is known about the use of these landscapes by marsupial carnivores,

which occupy a similar ecological niche and face comparable threats from habitat loss (13).

As well as displaying broad preferences among habitats, carnivores may also show finer scale

differences in the habitats used for different behaviors, such as travel, foraging and denning

(14). Understanding carnivore movements within these landscapes can help elucidate these

behaviors and highlight important habitat characteristics (15). Animals often change their

movements in response to altered forest structure (16). Habitat features such as windrows

(linear piles of logging debris) (17) and corridors (18) can facilitate animal movement across

production forest landscapes, and plantations can connect patches of native forest (19).

Anthropogenic linear features in production forest landscapes, such as roads and edges

between natural and modified vegetation, can be used by carnivores for travel and hunting

(20,21) but also expose them to threats such as vehicle collisions (22). Investigating how

carnivores move through production forestry landscapes, and the behaviors influencing

habitat selection, can inform management strategies to promote important habitat features in

the landscape.

With plantation forest cover in Australia increasing, fauna conservation is an important

consideration in plantation management (23). This is particularly relevant in Tasmania where

plantations comprise nearly 10% of total forest cover (24), and thus have significant potential

to supplement native forest habitats and contribute to maintaining ecological function of the

broader Tasmanian forest ecosystems if managed with conservation in mind.

Tasmania is home to the world's largest marsupial carnivore, the Tasmanian devil. Devils are

endangered due to a rare transmissible cancer, devil facial tumor disease (DFTD) (25), that

has caused a steep population decline of over 80% since 1996 (26), leaving them vulnerable

to other threats. Devils play multiple important roles in Tasmanian ecosystems: as

morphologically specialized scavengers, they influence carrion persistence in the landscape

(27), and as top predators, they help maintain trophic function and suppress feral cats (*Felis*

catus) (28). Finding ways to protect devils in human-modified landscapes is a priority to

maintain ecosystem function. Previous research investigated devil use of anthropogenic

pastoral landscapes, finding that devils foraged along the pasture-forest interface and used

pastures mainly for travel (29). Another study found devil abundance increases with

plantation extent in production forest landscapes (30). Movements and habitat selection by

devils in plantations have not been studied.

Our study provides the first analysis of marsupial carnivore habitat selection and movements

within plantations, complementing similar research on eutherian carnivores, and enhancing

our understanding of carnivore use of production forests. We used GPS tracking to explore

the use of a plantation landscape by devils, with the intent to provide recommendations for

plantation managers to sustain devil populations. We focused mainly on adult female devils,

a key demographic to protect in critically diminished populations, particularly when they are

raising young (April to January) (31,32). We aimed to answer 3 questions:

1. At a plantation landscape scale, how do female devils select among habitat types for

different behaviors, and how do different habitats influence devil home range sizes?

2. At a plantation-specific scale, how do female devils select among plantation ages for

different behaviors?

3. How do devils move through plantation landscapes?

Methods

Study area

This study was undertaken in Surrey Hills, a eucalypt plantation landscape in northwest

Tasmania (-41°S, 145°E) (Figure 1). The site is embedded in an intensively managed forestry

region containing a mosaic of *Eucalyptus nitens* plantation stands of varying ages and

remnant native forest and grassland patches. It has an average annual rainfall of 2,163 mm

and mean maximum temperature of 12.3°C (33). The plantations have mostly been

established since the mid-1970s on previously native vegetation land, and the remaining

native vegetation primarily comprises highland *Poa* spp. grasslands, *Eucalyptus delegatensis*

grassy forests and cool-temperate rainforest dominated by *Nothofagus cunninghamii* (34).

DFTD has been present in the landscape for about 6 years, causing steep population declines

of devils (26).

Figure 1: Study site within Tasmania (inset), overlaid with the combined home ranges of all devils in
the study (95% utilization distribution from Biased Random Bridges).

GPS telemetry

We captured adult devils (> 2 years old) in custom-built traps (made from 80 cm by 30 cm

PVC pipe) and fitted them with GPS radio-collars (Q4000ER GPS, Telemetry Solutions,

Concord USA) between April-October 2021. Collars weighed 140 g, less than 3% of the

body weight of collared individuals, and were fitted with corrosive bolts that degrade over

time and eventually allow the collar to drop off (35). As devils are primarily nocturnal (36),

collars were set to take 15 minute fixes every second night from approximately sunset until

sunrise (17:30 to 07:00 April and August, 17:00 to 07:30 May to July, 18:00 to 06:00

September, 18:30 to 05:30 October) and were turned off during the day to conserve batteries.

We set 'GPS additional time' to 30 seconds, which forces the GPS unit to keep searching for

satellites even after it acquires a fix, improving fix accuracy. Collars were also equipped with

a VHF unit to assist with relocation.

GPS data screening

To account for errors in animal locations acquired by the GPS, fixes with less than 3

satellites, horizontal dilution of precision (HDOP) > 7 (as in Andersen et al. [29]), or

unrealistic step lengths (> 4000 m) were removed from the analysis. Lower values of HDOP

indicate higher location precision. Because the successful fix rate was low, we tested for bias

in the success rate of GPS fixes due to vegetation type or plantation age. First, we left 10

collars stationary overnight in each of 3 vegetation types – native forest remnant, 1–5 year

old plantation, and 10–15 year old plantation – and compared percentages of successful fixes

among habitats. Second, we used straight-line interpolation to approximate the locations of

failed fixes ('approx' function in R package 'stats') and compared the percentages of points

in native forest, plantation and native grassland between real and interpolated points.

Interpolated points were not used for analysis.

To see how many failed GPS fixes could be attributed to animals resting in a den, ambient

temperatures recorded by the collars were plotted and compared between successful and

failed fixes. Temperatures > 25°C appeared to indicate animals were in a den. Collar

temperatures were unlikely to reach 25°C while animals were active since fixes were

programmed only at night, and devils were collared over winter and spring in a relatively

cold climate (annual mean max temp 12.3°C [33]).

Home ranges

We calculated devil home ranges using biased random bridges (BRBs) (37), which quantify

an animal's utilization distribution (UD) based on movement paths rather than just the

density of individual fixes. We identified 95% UDs for each individual devil. Home ranges

were also estimated using 100% minimum convex polygons (MCPs) (38) for comparison

with the broader literature. This is a commonly reported method of home range estimation,

though it is prone to overestimating the areas actually used by animals (39,40). We did not

use kernel density estimates (KDEs) (41), another widely used method, as the limited data

and decentralized spread of GPS points for many individuals made this method unsuitable for

our study. Analyses were carried out using the 'adehabitatHR' package in the R programming

environment (42).

We constructed maps of the study site using vegetation data from the TasVeg 3.0 layer (43)

and plantation features provided by the landowners, Forico Pty Ltd. Vegetation classes were

grouped into 3 habitat types: plantation, native forest, and native grassland (including

moorland and scrub). Plantation ages were grouped into 5 classes: < 1, 1–3, 4–7, 8–13 and

14+ years, age classes used by the forest industry due to different fuel loads. To examine if

habitat variables influenced devil home range size (95% UDs), linear models were

constructed using home range size as a response variable and proportions of different habitat

types and plantation ages within each devil's UD as predictors. These variables were included

in separate models as proportions of different habitats/plantation ages would be correlated

with one another, and plantation ages would be correlated with the percentage of plantation.

Sex was also used as a predictor variable, both alone and in combination with other variables.

We selected final candidate models using Akaike's Information Criterion values adjusted for

small sample size (AICc) (44) according to the methods of Richards (45), where models with

ΔAICc > 6 were excluded, as were more complex models where simpler, nested models had

smaller AICc values. To identify how much of the devils' home ranges were affected by

recent and active logging, and infer how this may influence their behavior, we calculated the

percentage of each devil's home range that was logged during the collaring period (April to

October 2021) and in the 12 months preceding it.

Movement states

To identify different movement states of female devils that may influence their habitat

selection, we classified the GPS fixes for each animal into 3 different movement states using

Hidden Markov Models (HMMs) (46). This method classifies movement paths according to

differences in turning angle and step length between successive locations, and assumes that

these characteristics are driven by underlying behaviors. Movement states can therefore

approximate behavioral states to some extent. We used past HMM results from devils to

guide our choice of initial parameters for each state (47) (see Supplementary Table 1,

Additional File 1): state 1, stationary (e.g., resting and/or feeding at carcass); state 2,

characterized by short step lengths and variable turning angles indicating tortuous movement

(e.g., foraging); and state 3, characterized by long step lengths and relatively constant turning

angles indicating directional movement (e.g., travel). Analyses were carried out using the

'moveHMM' package in R (48).

Sequential GPS tracking data are often temporally autocorrelated which can violate

assumptions for statistical inference (49). To account for this, once movement states were

assigned to all GPS fixes, we used variograms (function 'variogram' in R package 'ctmm'

[50]) for individual devils in each state to find a time threshold at which variance stabilized

and each location could be considered independent. We subsampled the GPS data by the time

threshold specific to each movement state (function 'track_resample' in R package 'amt';

[51]) and used the subsampled data for habitat selection analysis. We did not have sufficient

data to analyze selection of plantation ages by different movement states, so for plantation-

specific habitat selection analyses, we repeated this process and subsampled the data by the

time threshold for all combined states.

Habitat selection and animal movements

For landscape-scale habitat selection analyses and for the plantation-specific analyses, we

used the same 'habitat type' categories and 'plantation age' classes, respectively, as above.

To investigate how animals may use plantation edges (edges between plantation and other

habitat types, excluding roads) and roads for movement and foraging, we created 3 buffer

zones around plantation edges and roads: < 20 m (on roads/edges, as in Andersen et al. [29]),

20–50 m (near roads/edges) and > 50 m (away from roads/edges). We calculated the

percentage of each habitat type, plantation age class, and road/edge buffer in each devil's

95% MCP (to reduce undue influence of outlying points), buffered by 5%, to represent

available habitat. We used MCPs to define available habitat as we needed to measure all

potentially available habitat rather than just the areas used by the animal. For each successful

GPS point within the buffered MCPs we identified the habitat type, plantation age class (if

applicable), and their distance to plantation edges and roads, split into the above buffers.

These calculations were performed in QGIS 3.14$^{®}$.

We used Manly's selection ratios to investigate habitat selection by female devils in each

HMM movement state, a method that tests for differences in used vs. available habitat (52).

As there were insufficient male devils to examine sex differences in habitat selection, the 3

male devils were excluded from this analysis. We used a design III analysis, which

recognizes that the available habitat differs among individuals (52). The selection ratio is the

proportional use divided by proportional availability of each habitat attribute class, with a

selection ratio (*wi*) > 1 indicating that a habitat type was used proportionally more than

expected based on availability and < 1 proportionally less. The percentages of habitat

attributes within buffered MCPs represented available habitat, while the locations of GPS

points within the buffered MCPs represented used habitat. Univariate models were run for

each movement state with the predictors of habitat type, plantation age, and distance to roads

and plantation edges. Analyses were undertaken using the 'adehabitatHS' package in R (53).

Velocity

To support movement state analyses and help clarify devil movements in different habitats,

we calculated the velocity of each devil at each GPS point using the difference in time and

distance between 2 consecutive points using the 'sp' package in R (54). GPS points following

a failed fix were ignored. The distance from each GPS point to the nearest road and

plantation edge was calculated in QGIS 3.14$^{®}$ and classified into 'on road/edge' (< 20 m),

'near road/edge' (20–50 m) and 'away from road/edge' (> 50 m). We tested for correlation

between the values for plantation edges and roads using a polychoric correlation test

(function 'polychor' in package 'polycor' in R) and found low correlation (0.36), so these

variables could be included in the same model. To investigate if devils move at different

speeds in different habitats, we constructed linear mixed effects models using 'velocity' as a

response variable and distance to road, distance to edge, habitat type, and plantation age as

predictors. Animal ID was included as a random effect. We ran 2 sets of models: 'whole-of-

landscape' models, including all successful GPS fixes and all predictors except plantation

age; and 'plantation-only' models, including only GPS fixes inside plantations, and all

predictors except habitat type (to investigate the influence of plantation age). Final candidate

models were selected using the same process as for the home range models (44), using AIC

for whole-of-landscape models and AICc for plantation-only models.

Results

We deployed GPS radio-collars on 15 adult devils (12 females, 3 males) between April and

October 2021. One collar was unable to be retrieved at the end of the study, though some data

were downloaded when the animal was recaptured during the study. A total of 7406

successful GPS fixes remained after removal of GPS errors, a success rate of 23% (successful

vs programmed fixes). After accounting for temperatures recorded by the collar that indicated

the animal was in a den (> 25 °C), the fix success rate increased to 52%, indicating much of

the missing data was due to devils being in underground burrows where the collars could not

detect satellites. Data were retrieved from between 24 and 164 days per animal (mean 77 ±

49 SD), with 152 to 1452 successful fixes per individual (494 ± 384) (see Supplementary

Table 2, Additional File 1). A total of 7066 points were used for analysis, following removal

of points outside each animal's buffered 95% MCP.

The stationary collar test had a fix success rate of 99% in plantation age 1–5 years, 97.2% in

age 10–15 years, and 93.6% in native forest, similar to the difference in fix success rate

among habitats in other studies (29). Comparison of the proportion of real vs. interpolated

fixes that fell in each habitat category showed little difference between habitats (native forest

45.7% and 45.8% respectively, plantation 43.8% and 45.5%, grassland 10.4% and 8.8 %),

indicating that any difference in fix success rate among habitats did not introduce a

significant bias, though there was considerable variation among individuals (see

Supplementary Table 3, Additional File 1).

The best performing HMM differentiated the 3 movement states as: state 1, representing little

or no movement, with a mean step length of 16.37 ± 13.09 m (± SD) and a wide distribution

of turning angles centered on 3.13 rad (concentration 0.39); state 2, with intermediate step

lengths of mean 177.84 ± 148.25 m and a wide distribution of turning angles centered on 0.03

rad (concentration 0.52); and state 3, with longer steps of mean 527.03 ± 228.58 and a narrow

distribution of turning angles centered on 0.03 rad (concentration 1.66), suggesting faster

directional movement (see Supplementary Figure 1, Additional File 1). GPS data variance

stabilized and temporal autocorrelation decreased around 3 hours for state 1 and combined

states, and 4 hours for states 2 and 3. Data were subsampled by these state-specific time

thresholds for habitat selection analysis, leaving 338 observations for state 1, 469 for state 2,

712 for state 3, and 859 for combined states. All HMM results can be found in

Supplementary Table 4, Additional File 1.

Landscape-scale habitat selection and home ranges

At a landscape scale, behavior-specific habitat selection ratios for female devils showed that

devils did not select among habitat types (plantation, native forest, and native

grassland/scrub) in any movement state (state 1: $p = 0.48$, state 2: $p = 0.69$, state 3: $p = 0.56$)

(Figure 2a).

Home range sizes (95% UDs from BRBs) were a mean of 1849 (416 SE) ha for female devils

and 3343 (1388 SE) ha for males. Core use areas (50% UDs) were 318 (66 SE) ha for

females and 467 (102 SE) ha for males. 100% MCPs were a mean of 3481 (1475 SE) ha for

females and 6326 (4401 SE) ha for males. UD and MCP estimations for individual devils can

be found in Supplementary Table 2, Additional File 1. A single model remained in the final

candidate set for predictors of devil home range size (95% UDs) and included only sex

(**Error! Reference source not found.**). Home ranges of males were 1494 (95% CI 566 to

2421) ha larger than females. The average recently logged area within each devil's home

range was 3.46% (0–9.15%, see Supplementary Table 1, Additional File 1). Male home range

results should be treated with caution as there were only 3 male devils in the study.

Home range size (ha)				
Fixed effects			Model description	
Intercept	Sex		df	AICc
1849 [1433 to 2264] (female)	1494 [566 to 2421] (male)		3	234.6

Velocity (m.sec^{-1}) (whole-of-landscape)

Fixed effects			Random effects		Model description	
Intercept (m)	Distance to road (m)	Distance to plantation edge (m)	Intercept (animal ID)	Residual	df	AIC
0.41 [0.38 to 0.44] (< 20)	-0.09 [-0.12 to -0.06] (20–50) -0.06 [-0.09 to -0.04] (> 50)	-0.07 [-0.09 to 0.05] (20–50) -0.08 [-0.10 to -0.06] (> 50)	0.002 (0.04 SD)	0.09 (0.30 SD)	5	2511.3

Table 1: Final candidate models with parameter estimates for predicting devil home range size and
velocity in plantations (95% CIs in square brackets []).

Plantation-specific habitat selection

Within plantations, habitat selection ratios for female devils indicated devils used plantations

of different ages non-randomly ($p < 0.001$). Female devils selected for age 4–7 years ($wi =$

1.52 [95% CI 0.98–2.05]), slightly avoiding older age classes 8–13 ($wi = 0.74$ [0.28–1.20])

and 14+ years ($wi = 0.65$ [0.16-1.15]) and neither selecting nor avoiding young age classes <

4 years (Figure 2b).

Devil movements in plantation landscapes

Female devils selected habitats non-randomly in all movement states regarding distance to

roads (state 1: $p = 0.1$, state 2: $p = 0.01$, state 3: $p < 0.001$) and plantation edges (state 1: $p <$

0.001, state 2: $p < 0.001$, state 3: $p < 0.001$) (Figure 2c). Female devils used areas < 20 m

from plantation edges (state 1: $wi = 2.38$ [95% CI 1.66–3.10], state 2: $wi = 2.24$ [1.63–2.85],

state 3: $wi = 2.78$ [2.09–3.46]) and roads (state 1: $wi = 2.71$ [1.38–4.05], state 2: $wi = 2.48$

[1.59–3.37], state 3: $wi = 2.97$ [2.10–3.82]) more than twice as often as expected for all

states. To a lesser extent, they selected for areas 20–50 m from plantation edges ($wi = 1.47$

[95% CI 1.11–1.82]) and roads ($wi = 1.69$ [1.19–2.20]) for state 2 only. They avoided areas >

50 m from edges (state 1: $wi = 0.68$ [95% CI 0.47–0.89], state 2: $wi = 0.64$ [0.49–0.79], state

3: $wi = 0.62$ [0.49–0.75]) and roads (state 1: $wi = 0.87$ [0.79–0.95], state 2: $wi = 0.87$ [0.81–

0.92], state 3: $wi = 0.89$ [0.83–0.96]) in all states, particularly edges.

Figure 2: Habitat selection by adult female devils regarding habitat type (a), plantation age (b) and
distance to roads/edges (c) in 3 movement states. Movement states were not considered for plantation
ages due to limited data. Error bars show 95% CIs. A selection ratio of 1 indicates attributes were
selected in proportion to availability; 0.5 indicates they were used half as often as expected; and 2
indicates they were used twice as often as expected, based on availability within buffered 95% MCPs.

A single model remained in the final whole-of-landscape candidate set predicting devil

velocity and this model included both distance to roads and plantation edges (**Error!**

Reference source not found.). Devils moved faster on roads and edges (< 20 m) than away

from them (Figure 3). In the plantation-only models, only distance to plantation edges

remained in the final candidate list (see Supplementary Table 5, Additional File 1). As

plantation-only models used a subset of the whole-of-landscape data and the results were

similar, we consider whole-of-landscape results more relevant for interpretation.

Figure 3: Predicted velocity of devils with distance to roads and plantation edges using whole-of-
landscape data. Error bars represent 95% CIs.

Discussion

Our results demonstrate the adaptability of devils to human-modified landscapes,

highlighting the need to manage these landscapes as valuable devil habitat. Devils did not

show preferences among different habitat types at a landscape scale, though at a finer scale

they showed selection for specific plantation ages, and they preferentially used plantation

edges and roads to hunt and to move through the landscape.

Landscape-scale habitat selection and home ranges

At a landscape scale, female devils did not show selection among plantation, native forest

remnants, and native grassland in any movement state, and no habitat attributes considered

affected devil home range sizes. This demonstrates the adaptability of devils to human-

modified landscapes, consistent with Jones et al. (30)'s finding that devil abundance slightly

increased with plantation extent in production forest landscapes. Devils are generalist

carnivores (29), and generalist species usually respond more positively to human-modified

landscapes than specialists (55). Other generalist carnivores such as raccoon dogs (56) even

use plantations to a greater extent than native forest. Plantations likely provide abundant prey

for devils. Medium-sized herbivores, wallabies and pademelons (Family Macropodidae;

Macropus rufogriseus and *Thylogale billardierii*), and brushtail possums (*Trichosurus*

vulpecula), preferred devil prey (57), are common enough in Tasmanian plantations that they

are regularly culled to protect seedlings (24). Culling along with roadkill from forestry

vehicles would provide abundant carrion for this specialized scavenger (58). There appear to

be sufficient resources available in these landscapes for an opportunistic predator and

facultative scavenger to exploit.

Notably, home ranges of devils in this study were substantially larger than those in Andersen

et al. (59) (100% MCPs: females 2072 [277 SD] ha, males 3105 [309 SD]), which was

conducted in an area that was free of DFTD at the time of the study and contains a mix of

conservation and agricultural land. This is surprising as female devils decreased their home

range sizes in another landscape that suffered significant population decline due to DFTD

(60), likely due to less competition for resources. In the present study, we would therefore

expect smaller home range sizes than in Andersen et al. (59) due to low devil density in our

study landscape caused by high DFTD mortality (26). As home range size tends to increase

with lower quality habitat (61), this result may indicate that the plantation-dominated

landscape in our study area was less ideal devil habitat than the more protected landscape in

Andersen et al. (59). Perhaps protected landscapes provide more spatially and temporally

constant and reliable resources (e.g., prey and den sites) than a production forest landscape

that is always changing due to harvesting. For instance, black bears (*Ursus americanus*) were

found to have larger home ranges in plantation dominated than in unmanaged forest

landscapes (62).

Plantation-specific habitat selection

Within plantations, female devils selected for plantation aged 4–7 years and avoided older

plantations. Younger plantations may contain a higher density of wallabies and brushtail

possums which browse in plantations (63). Some other carnivores, such as American marten

(*Martes americana*) (64), similarly prefer younger post-logging native forest, likely due to

prey availability. Additionally, near-surface vegetation cover in hardwood plantation, e.g.,

grass and shrubs, increase with plantation age in Tasmania (K. Parkins, pers. comm.), which

may create a dense understory layer that restricts movement for devils in older plantations.

While many carnivores prefer dense understory in plantations (11,65), this is often to avoid

other predators or stalk prey. Devils are pounce-pursuit predators, capable of fast but short

pursuit but dependent on some degree of ambush to get sufficiently close (58). They prefer

forested rather than open environments and use forest edges to intercept herbivore prey as

they move between daytime refuges and open areas to feed at night (29,66). The habitat

structure offered by young plantations 4–7 years old may provide sufficient cover to conceal

them from prey but not enough to significantly obstruct movement and potentially impede

hunting and prey capture. Very young plantations may be too open for devils to be able to

effectively ambush prey which could see them coming and move out of range.

It is interesting that female devils did not appear to avoid recently logged plantation (< 1

year), although this result was based on just 4 individuals. Considering female devils likewise

did not avoid plantations 1–3 years old and actively selected for 4–7 years old, and that

plantation forestry landscapes have high prey abundance, logging is unlikely to have a long-

term negative effect on devil use of the landscape, as long as it is on a schedule that maintains

plantation age heterogeneity in the landscape. Furthermore, as we found female devils

avoided older plantation, logging removes habitat that is already apparently nonpreferred

(coupes are logged at ~12–15 years old in this landscape). Other factors likely to influence

the effect of logging on devils include the quality of remaining habitat in the surrounding area

and home range (e.g., plantation ages and native forest remnants) (23); the timing of logging

i.e., if logging occurs during the maternal denning season and could disrupt denning

behavior; and the overall area and proportion of home range harvested (67). We could not

directly investigate how devils responded to active logging, however, the proportion of their

individual home ranges that was logged during the study and in the preceding year was small

(mean 3.5%, range 0–9%), so even short-term displacement due to logging may not have had

a major effect on devils in this study. Extrapolation of these results to other years and

landscapes should be treated with caution due to variations in logging area size and intensity.

The lack of an apparent response to logging by devils is particularly surprising as the logging

method in the landscape is clearfelling, which involves removing all trees from the harvest

footprint and typically has a more negative effect on biodiversity than other logging methods

(68). Some carnivores can respond positively to clearfelling in plantations, however, perhaps

due to increased prey availability (69).

Devil movements in plantation landscapes

Devils preferentially used roads and plantation edges to move through the landscape in all

movement states. This is consistent with Andersen et al. (29)'s discovery that devils use

anthropogenic linear features for movement and foraging. Devils moved faster on roads and

edges, indicating they use them for travel. Edges of plantation coupes are cleared of trees to

allow vehicle access, potentially providing less obstructed movement paths for devils. Other

carnivores such as wolves similarly use linear features such as roads and seismic lines for

travel, allowing them to move further and faster (70). Our results indicate devils also forage

along roads and edges. As well as scavenging roadkill along roads (29), devils may use

plantation edges to intercept macropod prey moving between the denser shelter of native

forest where they rest during the day to browse in more open plantations at night (63), as was

found for the forest-pasture interface in Andersen et al. (29). Other carnivores likewise select

for edges in production forest landscapes (14,71), which can increase hunting success (20).

Female devils selected for areas close to (20–50 m), but not on, roads and edges for state 2

only. State 2 represents shorter, more tortuous movements which may indicate foraging,

contrasted with longer, straighter step lengths for state 3 which suggests travel, and stationary

behavior in state 1, i.e. resting or feeding at a carcass. Selection of areas 20–50 m from

roads/edges in state 2 may be explained by devils searching for food near roads/edges but in

cover to intercept prey, combined with a higher chance of picking up scents of prey or

carcasses that are close to roads/edges while travelling along them.

Management implications

As plantations appear to be valuable habitat for devils, we recommend implementing

strategies to maintain habitat suitability for devils in plantation landscapes. Preserving a

heterogeneous landscape with a mosaic of different plantation ages – along with patches of

native vegetation remnants (23) – is likely to benefit devils due to their broad use of different

habitat types and their preferences among plantation ages. The use of plantation landscapes

by devils exposes them to potential risks from human use of the landscape such as timber

harvesting operations, vehicle collisions, and lead poisoning from consuming culled browser

carcasses (72). Our finding that devils preferentially use roads make them particularly

vulnerable to vehicle collisions, a major cause of devil mortality (22). Logging in some

coupes occurs during 24 hours in this landscape. Since measures targeted at changing driver

behavior such as implementing speed limits and increasing driver awareness have limited

effectiveness (22), the most effective way to mitigate this risk to devils may be to minimize

forestry traffic at night. Our study suggests that the risk to devils from harvesting operations

is likely to be minimal, as devils slightly avoided the plantation age class most likely to be

harvested (14+ years), and the age class they preferred (4–7 years) is unlikely to be harvested.

One caveat is that we were unable to investigate denning behavior of devils in this study, so

the risk of damage to devil dens during logging operations – particularly maternal dens,

which may harbor young devils unable to escape – remains unknown.

Conclusions

Our study reinforces the high adaptability of devils to human-modified landscapes (29),

similar to other generalist carnivores (12). Not only did devils show no selection between

plantation and native habitat remnants, they took active advantage of modified habitat

features – edges and roads – for foraging opportunities and expedited travel. Similar

behavior has been observed before in devils (29) as well as other carnivores (70). Habitat

selection by carnivores in production forests is highly variable among species, such as

selection among harvested forest ages: some carnivores, like devils, prefer young-mid stage

regrowth (73), while others select for mature forest (74). Identifying species-specific habitat

selection in human-modified landscapes is therefore important. While it is essential to

recognize that carnivores as a group are particularly vulnerable to human impacts (6), it is

also important to understand that many generalist carnivores are adaptable to modified

landscapes, providing impetus to manage these landscapes as valuable habitat for these

species. Production forests are an excellent example – they are often incorrectly viewed as of

little value to conservation, exposing them to mismanagement and habitat degradation (2).

Recognizing that production forests and other modified landscapes provide important habitat

to devils and other carnivores is key to maintaining populations of these predators beyond

protected areas.

List of abbreviations

Devil facial tumor disease (DFTD)

Supporting Information

Additional information is available online in the Supporting Information section at the end of

the online article. The authors are solely responsible for the content and functionality of

these materials. Queries (other than absence of the material) should be directed to the

corresponding author.

Declarations

Ethics approval

This work was carried out under a permit from the University of Tasmania's Animal Ethics

Committee (permit number 23211).

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the

corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by a matched funding scheme through Forest and Wood Products

Australia Limited, with funding provided by Forico Pty Ltd, SFM, Sustainable Timbers

Tasmania, Timberlands, Norske Skog, and Private Forests Tasmania. It was also supported

by the Holsworth Wildlife Research Endowment [grant number H0027981], the Dr Eric

Guiler Tasmanian Devil Research Grant, and a Forest Practices Authority student research

grant. These funding bodies were not involved in the design of the study or the collection,

analysis, and interpretation of data or in writing the manuscript.

Authors' contributions

EJ was a major contributor in designing the study, acquiring funding, conducting fieldwork

for data collection, analysing the data and writing the manuscript. AK, DM, MJ and RH were

major contributors to study design. RH assisted with fieldwork issues. JP and MJ were major

contributors to data analysis. All authors contributed to writing the manuscript.

Acknowledgements

We thank Forico Pty Ltd for providing accommodation, data and access to their land, and T.

Patterson for assistance with data analysis. We are grateful to the many volunteers who

helped to collect data and samples.

References

1.   Curtis PG, Slay CM, Harris NL, Tyukavina A, Hansen MC. Classifying drivers of

498   global forest loss. Science. 2018 Sep 14;361(6407):1108–11.

499   2.   Brockerhoff EG, Jactel H, Parrotta JA, Quine CP, Sayer J. Plantation forests and

500   biodiversity: oxymoron or opportunity? Biodivers Conserv. 2008 May 1;17(5):925–

501   51.

502   3.   Hayes T, Ostrom E. Conserving the world's forests: Are protected areas the only

503   way? Ind L Rev. 2005;38(3):595–618.

4.   Kanowski J, Catterall CP, Wardell-Johnson GW. Consequences of broadscale timber

plantations for biodiversity in cleared rainforest landscapes of tropical and subtropical

Australia. Forest Ecology and Management. 2005 Apr 5;208(1):359–72.

5.   Hartley MJ. Rationale and methods for conserving biodiversity in plantation forests.

Forest Ecology and Management. 2002 Jan 1;155(1):81–95.

6.   Ripple WJ, Estes JA, Beschta RL, Wilmers CC, Ritchie EG, Hebblewhite M, et al.

Status and ecological effects of the world□s largest carnivores. Science. 2014 Jan

10;343(6167):1241484.

7.   Ritchie EG, Johnson CN. Predator interactions, mesopredator release and biodiversity

conservation. Ecology Letters. 2009;12(9):982–98.

8.   Kuijper DPJ, Sahlén E, Elmhagen B, Chamaillé-Jammes S, Sand H, Lone K, et al.

Paws without claws? Ecological effects of large carnivores in anthropogenic

landscapes. Proc Biol Sci [Internet]. 2016 Oct 26 [cited 2020 Jan 4];283. Available

from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5095381/

9.   Lantschner MV, Rusch V, Hayes JP. Habitat use by carnivores at different spatial

scales in a plantation forest landscape in Patagonia, Argentina. Forest Ecology and

Management. 2012 Apr 1;269:271–8.

10.   Moreira-Arce D, Vergara PM, Boutin S, Carrasco G, Briones R, Soto GE, et al.

Mesocarnivores respond to fine-grain habitat structure in a mosaic landscape

comprised by commercial forest plantations in southern Chile. Forest Ecology and

Management. 2016 Jun 1;369:135–43.

11.   Escudero-Páez SP, Botero-Delgadillo E, Estades CF. Effect of plantation clearcutting

on carnivore presence in industrial forest landscapes in south-central Chile.

Mammalia. 2018;83(2):115–24.

12.   Ferreira AS, Peres CA, Bogoni JA, Cassano CR. Use of agroecosystem matrix

529   habitats by mammalian carnivores (Carnivora): a global☐scale analysis. Mammal

530   Review. 2018 Oct;48(4):312–27.

531   13.   Jones M, Archer M, Dickman C. Convergence in ecomorphology and guild structure

532   among marsupial and placental carnivores. In: Predators with Pouches: The Biology

533   of Carnivorous Marsupials. Melbourne Australia: CSIRO Publishing; 2003.

534   14.   Houle M, Fortin D, Dussault C, Courtois R, Ouellet JP. Cumulative effects of forestry

535   on habitat use by gray wolf (*Canis lupus*) in the boreal forest. Landscape Ecol. 2010

536   Mar 1;25(3):419–33.

537   15.   Fuller AK, Harrison DJ. Movement paths reveal scale-dependent habitat decisions by

538   Canada lynx. Journal of Mammalogy. 2010;91(5):1269–79.

539   16.   Moriarty KM, Epps CW, Zielinski WJ. Forest thinning changes movement patterns

540   and habitat use by Pacific marten. The Journal of Wildlife Management.

541   2016;80(4):621–33.

542   17.   Sullivan TP, Sullivan DS. Responses of red-backed voles (*Myodes gapperi*) to

543   windrows of woody debris along forest–clearcut edges. Wildl Res. 2014 Oct

544   27;41(3):212–21.

545   18.   MacDonald MA. The role of corridors in biodiversity conservation in production

546   forest landscapes: a literature review. Tasforests. 2003;14:41–52.

547   19.   Lyra-Jorge MC, Ciocheti G, Pivello VR. Carnivore mammals in a fragmented

548   landscape in northeast of São Paulo State, Brazil. Biodivers Conserv. 2008 Mar

549   26;17(7):1573–80.

550   20.   Bojarska K, Kwiatkowska M, Skórka P, Gula R, Theuerkauf J, Okarma H.

551   Anthropogenic environmental traps: Where do wolves kill their prey in a commercial

552   forest? Forest Ecology and Management. 2017 Aug;397:117–25.

21. Gurarie E, Suutarinen J, Kojola I, Ovaskainen O. Summer movements, predation and

habitat use of wolves in human modified boreal forests. Oecologia. 2011 Apr

1;165(4):891–903.

22. Jones ME. Road upgrade, road mortality and remedial measures: impacts on a

population of eastern quolls and Tasmanian devils. Wildl Res. 2000;27(3):289–96.

23. Lindenmayer DB, Hobbs RJ. Fauna conservation in Australian plantation forests – a

review. Biological Conservation. 2004 Sep 1;119(2):151–68.

24. FPA. State of the forests Tasmania 2017 [Internet]. Hobart, Tasmania: Forest

Practices Authority; 2017 [cited 2020 May 6]. Available from:

https://www.fpa.tas.gov.au/__data/assets/pdf_file/0005/163418/State_of_the_Forests_

Report_2017_-_erratum_Feb_2018.pdf

25. Hawkins CE, McCallum H, Mooney N, Jones M, Holdsworth M. *Sarcophilus*

harrisii. The IUCN Red List of Threatened Species [Internet]. e.T40540A10331066.

2008 [cited 2020 Mar 20]. Available from:

https://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T40540A10331066.en.

26. Cunningham CX, Comte S, McCallum H, Hamilton DG, Hamede R, Storfer A, et al.

Quantifying 25 years of disease-caused declines in Tasmanian devil populations: host

density drives spatial pathogen spread. Ecology Letters. 2021;24(5):958–69.

27. Cunningham CX, Johnson CN, Barmuta LA, Hollings T, Woehler EJ, Jones ME. Top

carnivore decline has cascading effects on scavengers and carrion persistence. Proc R

Soc B. 2018 Dec 5;285(1892):20181582.

28. Cunningham CX, Johnson CN, Jones ME. A native apex predator limits an invasive

mesopredator and protects native prey: Tasmanian devils protecting bandicoots from

cats. Ecology Letters. 2020;23(4):711–21.

29.   Andersen GE, Johnson CN, Barmuta LA, Jones ME. Use of anthropogenic linear

features by two medium-sized carnivores in reserved and agricultural landscapes. Sci

Rep. 2017 Dec;7(1):11624.

30.   Jones EM, Koch AJ, Mann DF, Hamede RK, Jones ME. What drives the abundance

of marsupial carnivores in production forest landscapes? Forest Ecology and

Management. 2023 Feb 1;529:120745.

31.   Bryant SL. Seasonal breeding in the eastern quoll *Dasyurus viverrinus* (Marsupialia:

Dasyuridae) [PhD thesis]. [Hobart]: University of Tasmania; 1988.

32.   Pemberton D. Social organisation and behaviour of the Tasmanian devil, *Sarcophilus

harrisii* [Internet] [PhD thesis]. [Hobart]: University of Tasmania; 1990 [cited 2021

Nov 27]. Available from: https://eprints.utas.edu.au/11406/1/front.pdf

33.   Bureau of Meteorology. Climate Data Online [Internet]. 2022 [cited 2022 Oct 10].

Available from: http://www.bom.gov.au/climate/data/

34.   Onfray R. Fires, Farms and Forests: a human history of Surrey Hills, north-west

Tasmania. Tasmania: Forty South Publishing Bookshop; 2021.

35.   Thalmann S. Evaluation of a degradable time-release mechanism for telemetry

collars. Aust Mammalogy. 2013;35(2):241–4.

36.   Jones ME, Grigg GC, Beard LA. Body temperatures and activity patterns of

Tasmanian devils (<>Sarcophilus harrisii</i>) and eastern quolls (*Dasyurus

viverrinus*) through a subalpine winter. Physiological Zoology. 1997;70(1):53–60.

37.   Benhamou S. Dynamic Approach to Space and Habitat Use Based on Biased Random

Bridges. PLOS ONE. 2011 Jan 26;6(1):e14592.

38.   Mohr CO. Table of equivalent populations of North American small mammals. The

American Midland Naturalist. 1947;37(1):223–49.

39.   Burgman MA, Fox JC. Bias in species range estimates from minimum convex

polygons: implications for conservation and options for improved planning. Animal

Conservation forum. 2003 Feb;6(1):19–28.

40.   Hemson G, Johnson P, South A, Kenward R, Ripley R, McDonald D. Are kernels the

mustard? Data from Global Positioning System (GPS) collars suggests problems for

kernel home-range analyses with least-squares cross-validation. Journal of Animal

Ecology. 2005;74(3):455–63.

41.   Worton BJ. Kernel Methods for Estimating the Utilization Distribution in Home-

Range Studies. Ecology. 1989;70(1):164–8.

42.   Calenge C. Home Range Estimation in R: the adehabitatHR Package. 2019;61.

43.   DPIPWE. TASVEG 3.0 [Internet]. LISTdata. 2017 [cited 2022 Jan 5]. Available

from: https://www.thelist.tas.gov.au/app/content/data/geo-meta-data-

record?detailRecordUID=6e3d672d-cdc1-40f3-8ee4-b45c5de68b68

44.   Burnham KP, Anderson DR. Model selection and multimodel inference: a practical

information-theoretic approach [Internet]. USA: Springer New York; 2002 [cited

2022 May 26]. Available from: http://sutlib2.sut.ac.th/sut_contents/H79182.pdf

45.   Richards SA. Dealing with overdispersed count data in applied ecology. Journal of

Applied Ecology. 2008;45(1):218–27.

46.   Patterson TA, Thomas L, Wilcox C, Ovaskainen O, Matthiopoulos J. State–space

models of individual animal movement. Trends in Ecology & Evolution. 2008 Feb

1;23(2):87–94.

47.   Comte S. The spatial epidemiology of the Devil Facial Tumour Disease in long-

diseased populations of its unique host, the Tasmanian devil [Internet] [PhD thesis].

[Hobart]: University of Tasmania; 2019 [cited 2022 May 9]. Available from:

https://eprints.utas.edu.au/31698/1/Comte_whole_thesis_ex_pub_mat.pdf

48.   Michelot T, Langrock R, Patterson T. moveHMM: An R package for the analysis of

627   animal movement data. 2019.

628   49.   Boyce MS, Pitt J, Northrup JM, Morehouse AT, Knopff KH, Cristescu B, et al.

629   Temporal autocorrelation functions for movement rates from global positioning

630   system radiotelemetry data. Philosophical Transactions of the Royal Society B:

631   Biological Sciences. 2010 Jul 27;365(1550):2213–9.

632   50.   Calabrese JM, Fleming CH, Gurarie E. ctmm: an R package for analyzing animal

633   relocation data as a continuous-time stochastic process. Methods in Ecology and

634   Evolution. 2016;7(9):1124–32.

635   51.   Signer J, Fieberg J, Avgar T. Animal movement tools (amt): R package for managing

636   tracking data and conducting habitat selection analyses. Ecology and Evolution.

637   2019;9(2):880–90.

638   52.   Manly BF, McDonald L, Thomas DL, McDonald TL, Erickson WP. Resource

639   Selection by Animals: Statistical Design and Analysis for Field Studies. Springer

640   Science & Business Media; 2007. 233 p.

641   53.   Calenge C. Exploratory Analysis of the Habitat Selection by the Wildlife in R: the

642   adehabitatHS Package. 2011;60.

643   54.   Pebesma E, Bivand R, Rowlingson B, Gomez-Rubio V. sp: Classes and Methods for

644   Spatial data [Internet]. 2022 [cited 2022 Oct 6]. Available from: https://cran.r-

645   project.org/web/packages/sp/sp.pdf

646   55.   Dotta G, Verdade LM. Trophic categories in a mammal assemblage: diversity in an

647   agricultural landscape. Biota Neotropica. 2007;7(2):287–92.

648   56.   Hwang HS, Son SH, Kang H, Rhim SJ. Ecological factors influencing the winter

649   abundance of mammals in temperate forest. fozo. 2014 Dec;63(4):296–300.

650   57.   Andersen GE, Johnson CN, Barmuta LA, Jones ME. Dietary partitioning of

651   Australia's two marsupial hypercarnivores, the Tasmanian devil and the spotted-tailed

652   quoll, across their shared distributional range. Crowther MS, editor. PLoS ONE. 2017

653   Nov 27;12(11):e0188529.

654   58.   Jones M, Archer M, Dickman C. Predators with pouches: the biology of carnivorous

655   marsupials. CSIRO Publishing; 2003. 508 p.

656   59.   Andersen GE, Johnson CN, Jones ME. Space use and temporal partitioning of

657   sympatric Tasmanian devils and spotted-tailed quolls. Austral Ecology.

658   2020;45(3):355–65.

659   60.   Comte S, Carver S, Hamede R, Jones M. Changes in spatial organization following an

660   acute epizootic: Tasmanian devils and their transmissible cancer. Global Ecology and

661   Conservation. 2020 Jun 1;22:e00993.

662   61.   Kittle AM, Anderson M, Avgar T, Baker JA, Brown GS, Hagens J, et al. Wolves

663   adapt territory size, not pack size to local habitat quality. Journal of Animal Ecology.

664   2015;84(5):1177–86.

665   62.   Jones MD, Pelton MR. Female American black bear use of managed forest and

666   agricultural lands in coastal North Carolina. Ursus. 2003;14(2):188–97.

667   63.   Le Mar K, Mcarthur C. Comparison of habitat selection by two sympatric macropods,

668   *Thylogale billardierii* and *Macropus rufogriseus rufogriseus*, in a patchy eucalypt-

669   forestry environment. Austral Ecol. 2005 Sep;30(6):674–83.

670   64.   Fuller AK, Harrison DJ. Influence of partial timber harvesting on American martens

671   in north-central Maine. The Journal of Wildlife Management. 2005;69(2):710–22.

672   65.   Caryl FM, Quine CP, Park KJ. Martens in the matrix: the importance of nonforested

673   habitats for forest carnivores in fragmented landscapes. Journal of Mammalogy.

674   2012;93(2):464–74.

66.   Jones ME, Barmuta LA. Niche differentiation among sympatric Australian dasyurid

carnivores. Journal of Mammalogy. 2000;81(2):14.

67.   Leonard TD, Taylor PD, Warkentin IG. Landscape Structure and Spatial Scale Affect

Space use by Songbirds in Naturally Patchy and Harvested Boreal Forests. The

Condor. 2008 Aug 1;110(3):467–81.

68.   Chaudhary A, Burivalova Z, Koh LP, Hellweg S. Impact of forest management on

species richness: Global meta-analysis and economic trade-offs. Sci Rep. 2016

Jul;6(1):23954.

69.   Eom TK, Hwang HS, Lee JK, Bae HK, Park CR, Lim JH, et al. Effects of forestry

practices on habitat variables and mammal abundance in a Japanese larch plantation.

wbio [Internet]. 2019 Dec [cited 2020 Jun 18];2020(1). Available from:

https://bioone.org/journals/Wildlife-Biology/volume-2020/issue-1/wlb.00566/Effects-

of-forestry-practices-on-habitat-variables-and-mammal-

abundance/10.2981/wlb.00566.full

70.   Dickie M, Serrouya R, McNay RS, Boutin S. Faster and farther: wolf movement on

linear features and implications for hunting behaviour. Journal of Applied Ecology.

2017;54(1):253–63.

71.   Simons□Legaard EM, Harrison DJ, Krohn WB, Vashon JH. Canada lynx occurrence

and forest management in the Acadian Forest. The Journal of Wildlife Management.

2013;77(3):567–78.

72.   Hivert LG, Clarke JR, Peck SJ, Lawrence C, Brown WE, Huxtable SJ, et al. High

blood lead concentrations in captive Tasmanian devils (*Sarcophilus harrisii*): a threat

to the conservation of the species? Aust Vet J. 2018 Nov;96(11):442–9.

73.   Boisjoly D, Ouellet JP, Courtois R. Coyote habitat selection and management

implications for the Gaspésie caribou. The Journal of Wildlife Management.

2010;74(1):3–11.

74.   Buskirk SW, Yiqing M, Li X, Zhaowen J. Winter habitat ecology of sables (*Martes*

zibellina) in relation to forest management in China. Ecological Applications.

1996;6(1):318–25.