Article title: Intra and interspecific diversity in a tropical plant clade alter herbivory and

ecosystem resilience

Authors[*]:

Ari J. Grele[1], Tara J. Massad[2], Kathryn A. Uckele[1], Lee A. Dyer[1,3], Yasmine Antonini[4], Laura

Braga[4], Matthew L. Forister[1,3], Lidia Sulca Garro[5], Massuo J. Kato[6], Humberto G. Lopez[1], Andre

R. Nascimento[7], Thomas L. Parchman[1,8], Wilmer R. Simbaña[9], Angela M. Smilanich[1], John O.

Stireman[10], Eric J. Tepe[11], Thomas R. Walla[12], Lora A. Richards[1,3]

Affiliations

[1]Program in Ecology, Evolution, and Conservation Biology, Department of Biology, University

of Nevada, Reno, NV 89557, USA.

[2]Department of Scientific Services, Gorongosa National Park, Sofala, Mozambique.

[3]Hitchcock Center for Chemical Ecology, University of Nevada, Reno, NV 89557, USA.

[4]Lab. de Biodiversidade, Departamento de Biodiversidade, Evolução e Meio Ambiente. Instituto

de Ciências Exatas e Biológicas. Universidade Federal de Ouro Preto (UFOP). Ouro Preto, MG,

Brazil

[5]Departamento de Entomología, Museo de Historia Natural, Universidad Nacional Mayor de San

Marcos, Av. Arenales 1256 Jesús María, Lima 14, Peru.

[6]Department of Fundamental Chemistry, Institute of Chemistry, University of São Paulo, São

Paulo, Brazil.

[7]Department of Ecology, Universidade Federal de Goiás, Goiânia, GO, Brazil.

[8]Department of Biology, University of Nevada, Reno, NV 89557, USA

[9]Yanayacu Biological Station, Cosanga, Napo Province, Ecuador.

[10]Department of Biological Sciences, Wright State University, Dayton, OH 45435, USA.

[11]Department of Biological Sciences, University of Cincinnati, Cincinnati, OH 45221, USA.

[12]Department of Biology, Mesa State College, Grand Junction, Colorado 81501, USA

*Emails for authors are as follows: **A.J.G.:** arijg21@gmail.com, **T.J.M.:** tmassad77@gmail.com,

K.A.U: katuckele@gmail.com, **L.A.D.:** ldyer@unr.edu, **Y.A.**: antonini.y@gmail.com, **L.B**.:

lblepidoptera@gmail.com, **M.L.F.:** forister@gmail.com, **L.S.G.:** lidia.sulca@gmail.com,

M.J.K.: massuojorge@gmail.com, **H.G.L.:** humgar@hotmail.com, **A.R.N.:**

andre110479@gmail.com, **T.L.P.:** tparchman@unr.edu, **W.R.S.:** rosendocat@yahoo.com,

A.M.S.: smilanich@gmail.com, **J.O.S.:** john.stireman@wright.edu, **E.J.T.:**

tepeej@ucmail.uc.edu, **T.R.W.:** metawalla@gmail.com, **L.A.R.:** lorar@unr.edu

Correspondence and requests for materials should be addressed to Lora Richards,

lorar@unr.edu. Dept of Biology, UNR, MS:314, 1664 N. Virginia St. Reno, NV 89523

Data accessibility statement:

Data and code used for all analyses and figures can be accessed via the Dryad repository at

https://doi.org/10.5061/dryad.8w9ghx3rf

Abstract

46   Declines in biodiversity generated by anthropogenic stressors at both species and

47 population levels can alter emergent processes instrumental to ecosystem function and resilience.

48 As such, understanding the role of biodiversity in ecosystem function and its response to climate

49 perturbation is increasingly important, especially in tropical systems where responses to changes

50 in biodiversity are less predictable and more challenging to assess experimentally. Using large

51 scale transplant experiments conducted at five neotropical sites, we documented the impacts of

52 changes in intraspecific and interspecific plant richness in the genus *Piper* on insect herbivory,

53 insect richness, and ecosystem resilience to perturbations in water availability. We found that

54 reductions of both intraspecific and interspecific *Piper* diversity had measurable and site specific

55 effects on herbivory, herbivorous insect richness, and plant mortality. The responses of these

56 ecosystem-relevant processes to reduced intraspecific *Piper* richness were often similar in

57 magnitude to the effects of reduced interspecific richness. Increased water availability reduced

58 herbivory by 4.2% overall, and the response of herbivorous insect richness and herbivory to

59 water availability were altered by both intra and interspecific richness in a site dependent

60 manner. Our results underscore the role of intraspecific and interspecific richness as foundations

61 of ecosystem function and the importance of community and location specific contingencies in

62 controlling function in complex tropical systems.

64 **Keywords:** Biodiversity, Ecosystem function, Herbivores, *Piper*, Plant-Insect interactions,

65 Water availability

Introduction

As climate change and anthropogenic activity alter ecosystems at unprecedented rates, it has become critical to understand the consequences of biodiversity loss on ecosystem processes and the maintenance of ecosystem processes through species interactions. A complex mix of anthropogenic forces are eroding multiple dimensions of global biological diversity, including plant intraspecific, interspecific, and functional diversity (1,2). Plant diversity affects herbivore abundance and diversity, thereby influencing biomass allocation and energy fluxes between trophic levels (3). Because losses of plant diversity can destabilize the flow of resources to higher trophic levels (3,4), understanding the connection between biodiversity and trophic interactions is necessary to predict the consequences of the loss of primary producer biodiversity on ecosystem traits such as resilience to environmental perturbation (5,6). While climate change is associated with increased absolute precipitation in some regions and decreased precipitation in others, IPCC models predict mid-century increases in the frequency of extreme precipitation events in Central and South America (7–9), a phenomenon already observable in many ecosystems (10). Measuring how precipitation change will affect the relationship between biodiversity and ecosystem function is therefore increasingly important, particularly in diverse tropical systems.

Very few multi-site, manipulative diversity experiments have been reported from tropical areas compared to temperate environments (11), limiting our knowledge of the role of biodiversity in ecosystem function in the most species-rich regions of the planet (12). In the context of several established hypotheses (Table 1) we investigate how multiple dimensions of plant diversity affect ecosystem processes at five neotropical sites and explore how diversity

modulates how ecosystems respond to changes in water availability at three of those sites. We

focus on ecosystem responses that represent changes in energy fluxes between trophic levels as

measured by herbivory, herbivore diversity and plant mortality. By altering plant uptake of

nutrients and plant defense production, abnormal levels of precipitation can alter herbivore

pressure, affecting the movement of resources into higher trophic levels (13). As such, extreme

dry and wet periods of climate are expected to strongly perturb plant-insect interactions and

thereby alter ecosystem function (13–15). The insurance hypothesis suggests that greater

biodiversity can act to stabilize ecosystems and improve their resilience to environmental change

(4,16). While greater interspecific plant richness is expected to lead to increased diversity in

higher trophic levels due to the accumulation of specialist herbivores, field studies have

demonstrated both positive and neutral effects of interspecific plant richness on ecosystem

resilience (17,18). Despite traditional views that interspecific richness has a greater impact on

ecosystem processes than intraspecific diversity (19), recent research suggests that plant intra-

and interspecific richness can have similar effects on ecosystem productivity and consumer

abundance (20,21). As changes to intraspecific richness can alter the diversity of resources

available to herbivores (22), it is necessary to consider both inter and intraspecific diversity when

investigating the effects of biodiversity loss on ecosystem processes.

We conducted common garden experiments at five sites across Central and South

America to test the insurance hypothesis by quantifying 1) the relative strength of intra- and

interspecific plant richness in driving ecosystem function and 2), the effects of increased water

availability on ecosystem function. Using 33 species in the genus *Piper* (Piperaceae) as a model

system (23), we manipulated intra- and interspecific plant richness in Costa Rica, Ecuador, Peru,

and two sites in Brazil. We additionally manipulated water availability in Costa Rica, Ecuador,

and Peru (Figs. 1 & S1 A). We predicted that reduced *Piper* diversity would lead to reduced

diversity of higher trophic levels, that water addition would lead to altered herbivore pressure,

and that lower *Piper* diversity would be associated with more extreme changes in herbivory and

plant mortality in response to water addition. Finally, we predicted that changes in intraspecific

and interspecific plant richness would affect ecosystem processes, including herbivory, herbivore

diversity, and plant mortality, with similar magnitudes.

Methods

Study sites and focal plant genus

We conducted a large-scale transplant experiment replicated across five sites spanning

42° latitude in the Neotropics (Fig. 1 & Table S1) encompassing a range in annual precipitation

from 1271 mm to 4495 mm (Table S1). At each site, we studied herbivory on planted individuals

in the genus *Piper* (Piperaceae) in response to experimental treatments. Study sites included

lowland equatorial humid forest at La Selva Biological Station, Costa Rica; high elevation

equatorial humid forest in Yanayacu Biological Station, Ecuador; high elevation equatorial

humid forest at El Fundo Génova, Peru; lowland seasonally dry gallery forest in the cerrado

within the phytogeographic domain of the Atlantic Forest in Mogi-Guaçu Biological Reserve,

Brazil; and lowland seasonally semideciduous forest in the transition between the Atlantic Forest

and the cerrado phytogeographic domains in Uaimii State Forest, Brazil. Climate classifications

follow the Köppen-Geiger climate model (Table S1; 22).

Multiple species of *Piper* are found at all sites, ranging from 11 species in Mogi-Guaçu to

50 species in Costa Rica (25). *Piper* is an ideal genus for large-scale comparative studies, as it is

found across the Neotropics and subtropics, from ~10° N to about ~32° S. In addition to being

widespread, *Piper* is abundant and diverse across its range, encompassing ~1000 species in the

Neotropics (26). *Piper* has been the subject of detailed studies of herbivory, and its herbivore

fauna has been surveyed across its range (27–38). Members of the genus host both specialist

caterpillars and beetles as well as generalist caterpillars and Orthopterans (23,27–30,32). Leaf

damage patterns produced by the different classes of herbivores are well documented and allow

for the determination of unique taxa of herbivores (32).

Experimental design

Executing experiments across the Americas presents challenges, including nuanced

variations in methodologies at each site. Nonetheless, the advantages of this expansive and

consistent approach provides a greater understanding in the role of biodiversity in ecosystem

function than examining isolated single-site studies. Here we describe the experimental design

applied across sites, see Figures 1 & S1, and Tables S1 & S2 for site specific details.

At each study site, a factorial experiment was implemented to test the effects of plant

interspecific and intraspecific richness on herbivory, variation in herbivory, and insect richness.

Experimental plants were propagated from cuttings of naturally occurring *Piper* plants (typically

with three nodes and zero to one leaves). At each site a subset of naturally occurring *Piper*

species was selected to act as a species pool for each experimental plot (Table S2). These species

were selected to constitute a breadth of genetic and functional diversity representative of the

Piper community present at each site. In all sites except Uaimii, experimental plots measured 4

m in diameter and contained 12 *Piper* individuals, planted either in pots with locally derived soil

(for experiments manipulating water availability) or directly into the ground (for experiments

without water additions in Mogi-Guaçu and Uaimii; Table S1) and cultivated without fertilizer or

irrigation. Interspecific richness treatment levels consisted of single species monoculture plots,

two species plots, and high richness plots with the maximum number of species available to

produce cuttings (Table S1). The minimum number of *Piper* species in a high richness plot was

three in Uaimii, and the maximum was 12 in Costa Rica and Ecuador. In plots with more than

one species, species were randomly sampled from the species pool for that site.

An intraspecific richness treatment was crossed with the interspecific richness treatment.

Intraspecific richness was manipulated by clones taken from a single mother plant (low richness)

or individual cuttings all taken from unique mother plants (high richness). In the high

intraspecific richness treatment, cuttings were from different *Piper* individuals growing at least

10 m apart, to eliminate the likelihood of a shared root system thus representing genetically

unique individuals. In Costa Rica and Ecuador, the high interspecific richness plots only

included individuals from unique mother plants because all cuttings were taken from different

species.

A water addition treatment was crossed with the inter- and intraspecific richness

treatments in Costa Rica, Ecuador, and Peru in order to examine how IPCC predicted regional

mid-century increases in precipitation in these regions (9) may affect herbivory and trophic

interactions. Plants were cultivated in five to six liter pots with drainage holes. Each potted plant

under the water addition treatment was watered with 2 L of water twice per month in Costa Rica

and Ecuador. In Peru, 2 L of water were added to plants every three months. Water was rapidly

applied as a flooding event to completely saturate the soil. Plant cultivation periods lasted in

excess of one year at all sites, and as such water was added in both the wet season and dry

season. Experimental plots were randomly located in the study sites; in Mogi-Guaçu and Uaimii

they were organized in three replicate blocks. There was a minimum of 20 m between plots in

blocks and 100 m between blocks. Randomly located plots were separated by a minimum of 50

m. Experimental periods lasted between 1.4 and 2.8 years, depending on the site (Table S1), and

all measurements of herbivory were conducted on leaves that were initiated during these periods.

Cuttings were replaced if they died during the first three months of the experiment. One of the

authors was present at all of the sites for multiple visits to ensure as much standardization of

treatment applications as possible.

Piper mortality resulted in a reduction in species richness in many plots and the loss of

some treatment combinations, notably the loss of all *Piper* in unwatered, low intraspecific

richness plots in Peru (Fig. S1B). Plots originally planted as high interspecific richness

treatments had a final richness of between 1 and 12 species in Costa Rica, 9 and 11 species in

Ecuador, 1 and 3 species in Peru, 2 and 4 species in Mogi, and only 2 species in Uaimii.

To determine the effects of site variation in natural levels of precipitation on the outcomes of the

water addition treatment, the absolute level of precipitation and precipitation anomalies relative

to climate normals were collected for each month of the experimental periods in Costa Rica,

Ecuador, and Peru (Fig. S2). Climate data for Costa Rica were obtained from La Selva

Biological Station. Data for Ecuador and Peru were obtained using TerraClimate (39), and an

interpolation error in the precipitation for Peru in February 2016, was corrected using data from

the National Service of Meteorology and Hydrology of Peru (SENAMHI).

Measures of herbivory and insect richness

Plots were open to naturally occurring herbivores, and herbivory was recorded by taking

photographs of all the experimental leaves at the end of the experimental period. Additional

photos were taken every three months in Uaimii and Mogi-Guaçu, and in the first five months of

the experiment in Costa Rica. These data were used to measure herbivory and to determine the

types of herbivores feeding on leaves based on patterns of damage. When herbivores were

observed on plants, they were photographed but were left to continue feeding so as to not

interfere with the experiment. Herbivory was quantified by eye for each type of herbivore on

each leaf by a single parataxonomist with extensive experience measuring herbivory on *Piper*

following established protocols (32). The amount of leaf area consumed was measured in

relation to the total leaf area by visually dividing the leaf into equal sized segments to determine

the percent area missing. This was measured as a continuous value to the greatest possible

accuracy, typically 1-5% of the total leaf area. Insect herbivores were identified to the lowest

taxonomic level possible based on their damage patterns (genus for specialist Lepidoptera,

family for generalist Lepidoptera, family for Coleoptera, order for Orthoptera). Direct

observations of herbivores were rare, so only damage patterns were used in analyses. As insect

damage patterns on leaves are tightly correlated with insect richness in tropical forests (40), the

different types of damage recorded were used as proxy for the richness of above ground insect

herbivores on *Piper*. Hereafter, the term "insect richness" refers to the richness of insect

herbivore damage patterns on plants.

Data analyses

The percentage of leaf tissue consumed by each insect taxon (based on damage patterns)

on individual leaves was summed to determine the total percentage of herbivory on each leaf,

and the variance in herbivory was calculated as the variance in leaf damage within each plant in

a plot. Due to high mortality rates across sites, *Piper* interspecific richness in each plot was the

final number of *Piper* species present in each plot at the end of the experiment (rather than the

number of species planted) and was analyzed as a continuous covariate rather than a categorical

treatment. Because most plant deaths occurred early in the experiments, final *Piper* interspecific

richness more accurately reflects the local plant richness experienced by herbivores. Interspecific

diversity for each plot was analyzed as the proportion of the interspecific diversity present in the

species pools at each site. This enabled easier comparisons between intra and interspecific

richness, as intraspecific richness was only quantified as low and high, where high treatments

represent the maximum intraspecific richness available at each site. The effect of intra and

interspecific diversity on each response variable is reported as the change in response as diversity

changes from the lowest possible value at each site to the highest.

The effects of intraspecific richness, interspecific richness, and water addition on percent

leaf area consumed, the percentage of damaged leaves, variance in herbivory, and insect richness

were analyzed across and within sites using hierarchical Bayesian models (HBM). This

framework acts as the Bayesian equivalent of a random-effects model where site is a random

effect, allowing for generalized parameter estimates across sites. Analyses testing *Piper*

intraspecific richness and interspecific richness were conducted using data from all sites; the

effects of water addition and its interactions with richness were conducted using data from

Ecuador, Costa Rica, and Peru. Due to high mortality in Peru, interactions between the water

addition and intraspecific richness treatments could not be modeled for measures of herbivory or

insect richness, and were only tested for survival.

We acknowledge that treatment level combinations were not the same across the different

sites, but experimental designs of this nature are encompassed within the framework of random-

effects models, where different levels of random factors, such as site or year, consist of treatment

levels that are unique to that site or year. This type of experimental design goes back to the

origins of mixed models (41,42), and the lack of interactions between fixed and random effects

increases generality in these models (43). Even at a single site, manipulated variables in

ecological experiments do not even consist of the same manipulations across all of the units of

replication, as they suffer from problems such as multiple versions of treatments, interference,

and noncompliance (44)

Bayesian structural equation models (BSEMs) were constructed for each site using all

treatments as exogenous variables and insect richness and herbivory as endogenous variables.

Three path models were developed under the assumptions that 1) intraspecific and interspecific

richness may influence herbivory both directly and indirectly by modulating insect richness 2)

water addition may influence both herbivory and insect richness and 3) insect richness may

influence herbivory directly (Table 1). Models I-III were tested in Ecuador, Costa Rica and Peru.

while a model without the water addition variable was used to analyze data from Mogi-Guaçu

and Uaimii. Additional models incorporating interactions between intraspecific richness,

interspecific richness, and water availability were tested in Costa Rica, Ecuador, and Peru.

Models were run at the leaf level for herbivory and insect richness, and at the plant level

for the percentage of leaves with damage, variance of herbivory, and *Piper* survival. For all

HBMs and BSEMs, model convergence was estimated visually using traceplots and an $\hat{R}$

discriminatory threshold of 1.1 (45). Model fit was determined via posterior predictive checks

(PPC) using the sum of squares of the residuals as the discriminatory function (46). A PPC near

0.5 indicates a high model fit, while values near one or zero indicate poor fit. BSEM models

were further compared using the Deviance Information Criterion (DIC).

All Bayesian models were written in JAGS via the jagsUI package in R (47) using

Markov chain Monte Carlo (MCMC) sampling with weakly-informative priors. Residuals were

modeled as normally distributed based on PPC comparisons between models. Models using

gamma distributions for herbivory and binomial distributions for damage presence and mortality

were found to consistently underestimate the magnitude of variance in the data based on PPC.

For the majority of models, MCMC runs were conducted for 10,000 iterations using the first

1,000 iterations as a burn-in phase to generate posterior distributions of parameter estimates for

each response variable. HBMs modeling interactions required 20,000 iterations with the first

5,000 as burn-ins for all models to consistently converge. Mean parameter estimates and 95%

credible intervals (CIs) were calculated for all responses. 95% CIs which do not cross the y-axis

are typically associated with less than a 2.5% type S error rate (48). A posterior probability of

direction (PD) was calculated based on the percentage of the posterior distribution responding in

the same direction as the median response. A PD of 95%, for example, indicates that the same

direction of response (e.g. a positive or negative response) was observed in 95% of iterations,

regardless of the magnitude of the response (49). PDs less than 95%, indicate lower confidence

that a relationship exists, but can still be interpreted as the probability that an effect exists.

To analyze the effects of the experimental treatments on plant survival, survivorship

curves were constructed for all sites. Analyses of *Piper* survival were based on the initially

planted interspecific richness treatment of each plot because mortality occurred early in the

experiment. The effects of water addition, intra- and interspecific richness and species identity

on *Piper* survival were analyzed using Cox proportional hazard models. All data were analyzed

using R software v4.1.0 (50).

Results

Considerable variation in herbivory and plant mortality was observed among study sites.

Percent herbivory was lowest at Mogi where only 9% of leaf tissue was consumed by herbivores

compared to 22% of leaf tissue consumed in Uaimii. *Piper* mortality was highest in Peru (89%),

likely due to El Niño related drought, and was lowest in Ecuador (27%; Fig. S3A). We identified

herbivore damage from 13 insect taxa at the five study sites, as well as damage by leaf miners of

unknown orders. Damage from a total of 10 taxa were observed in Costa Rica, Ecuador, and

Peru, 9 taxa were observed in Mogi, and 8 taxa were observed in Uaimii (Fig. S3 B). The

majority of taxa in Peru, Mogi, and Uaimii were generalist herbivores, while 60% of taxa in

Costa Rica and 58% of taxa in Ecuador were *Piper* specialists. The proportion of leaf tissue

consumed by specialists was only greater than generalist damage in Costa Rica.

Our experiments revealed pronounced heterogeneity in ecosystem responses to water

availability and *Piper* diversity between sites (Figs. S4 & S5). Posterior predictive checks

(PPCs) for all hierarchical Bayesian models (HBMs), and for models I, and III were within 0.03

of 0.5, indicating models fit well. Model III (Table 1) was selected as the most parsimonious

causal model for BSEMs in Costa Rica (PPC = 0.499), Ecuador (PPC = 0.499), and Peru (PPC =

0.498). Fit was high for models in Mogi (PPC = 0.5) and Uaimii (PPC = 0.497). Across all sites

where the water addition treatment was applied, percent herbivory was 4.2±3.6% (mean ± 95%

CI) lower in plots that received additional water (PD = 98.7%; Fig. 2 A-B). Greater *Piper*

interspecific richness was associated with a 15±18.6% increase in the richness of insect

herbivores (PD = 95.0%; Fig. 2E) and an indirect increase in herbivory was mediated by insect

richness (Fig. 3). Insect richness was associated with an 8.8±2.8% increase in herbivory per

insect taxon present (PD = 100%) and a 6.7±6.9% increase in the percentage of leaves with any

damage (PP = 97%) (Figs. 2A, B). Intra- and interspecific richness affected herbivory, although

effects varied in strength and direction across sites (Fig. 3, Table S3). Intraspecific richness had

similar or greater effects on plant mortality and insect richness than interspecific richness.

However, intra- and interspecific richness often had contrasting directions of effect on insect

richness and measures of herbivore pressure (Figs. 3, S4B-C & S5B-C). For example, in Costa

Rica insect richness was 6.0 ± 5.9% (PD = 93.7%) lower in plots with high interspecific plant

richness, while high intraspecific richness increased insect richness by 7.2 ± 6.9% (PD = 98.1%).

In contrast, in Ecuador high interspecific richness was associated with a 43.6 ±6.4% (PD =

100%) increase in insect richness, and high intraspecific richness decreased insect richness by

13.6 ±5.2% (PD = 100%; Fig. S5B-C).

The effects of water addition were altered by *Piper* intra- and interspecific richness at all

sites (Fig. 4). Water availability reduced herbivory in Costa Rica (4.7 ±2.5%, PD = 100%) and

Peru (5.1 ±3.7%, PD = 100%), but this effect was only present in Ecuador when intraspecific

richness was high (Fig. S6). Across sites, water addition had negligible or negative effects on

insect richness at low interspecific richness, but this pattern was reduced or reversed when

interspecific richness was high. Insect richness increased by 20.8 ±18.9% more in

interspecifically diverse plots water compared to unwatered plots in Costa Rica (PD = 99%).

Insect richness increased by 15.2 ±16.8% more in rich, watered plots in Ecuador (PD = 96%),

and 29.3 ±31.5% more in Peru (PD = 98%). Additionally, water addition had a negligible effect

on insect richness in Ecuador when intraspecific richness was low, but increased insect richness

by 12.1% when intraspecific richness was high (PD = 99%; Fig. 4B).

Water addition had a negative effect on *Piper* survival in Costa Rica when intraspecific

richness was low, but improved survival by 12.1 ±8.5% in high intraspecific richness plots (PD =

99%; Fig. 4C). Interactions with water in Peru may have been influenced by an El Niño related

drought which resulted in high *Piper* mortality, while the typically wetter sites in Costa Rica and

Ecuador experienced greater precipitation than average (Fig. S2).

Analysis of plant survival

Water addition did not affect *Piper* survival in Costa Rica ($z = -1.2$, $P > 0.2$), or in

Ecuador ($z = 0.2$, $P > 0.8$), but survival was reduced by 48% in plots without water in Peru ($z =$

3.2, $P = 0.001$). Intraspecific richness increased survival by 38% in Costa Rica ($z = -3.3$, $P <$

0.001) and by 32% in Ecuador ($z = -4.9$, $P < 0.001$), but had no effect on survival in Peru ($z = -$

1.55, $P > 0.1$), Mogi ($z = -0.22$, $P > 0.8$), or Uaimii ($z = 0.18$, $P > 0.8$). Interspecific richness had

no effect on survival in Costa Rica ($z = -1.5$, $P > 0.1$), Ecuador ($z = -0.56$, $P > 0.5$), Peru ($z = -$

0.49, $P > 0.6$), or Uaimii ($z = 0.21$, $P > 0.8$). In Mogi, survival was reduced by 13% in plots with

higher interspecific richness (z = -2.16, P = 0.031). There was an interaction between

intraspecific richness and water addition in Costa Rica and Peru. Survival in Costa Rica

increased in response to water in high intraspecific richness plots, and decreased in response to

water in low intraspecific richness plots (z = 3.8, P < 0.001), while the opposite pattern was

observed in Peru (z = -2.4, P = 0.02; Fig. S7). Although plant die-offs cause a loss in richness in

some plots, plant species identity was not related to survival with the exception of *P. peltatum*,

which had the lowest survival rate of any species planted in Costa Rica. Further statistical results

are available in the supplement.

Discussion

Our results demonstrate two key patterns. First, the strength of effects of intraspecific

richness on higher trophic levels is comparable to that of interspecific richness, supporting our

predictions and corroborating recent studies demonstrating the importance of intraspecific

richness (20,21,51). However, the direction of the effect of intraspecific richness on herbivory,

insect richness, and plant survival can vary dramatically from the direction of effect of

interspecific richness, in contrast to our predictions. Second, we found that perturbations in water

availability can have complex effects on herbivores and plant survival, and that these effects can

be modulated by plant diversity in a context sensitive manner. While our prediction that water

availability would influence herbivory across sites was supported, our results suggest that

biodiversity loss and climatic perturbations may have dramatically different effects on ecosystem

function at local scales, which may diminish our ability to predict how local communities will

change as anthropogenic stressors increase.

As we did not directly measure plant stress or nutrition, it is difficult to determine the

exact mechanism through which water addition reduced herbivory. The presence of an El Niño

weather pattern during the course of the Peru experiment may have led the water addition

treatment to relieve plants from drought stress, while water addition in Costa Rica and Ecuador

may have added to water stress in treated plants as these sites received an above average level of

rainfall during the course of the experiment. Despite this, water addition consistently suppressed

herbivory in Costa Rica, Ecuador, and Peru under natural levels of *Piper* diversity, suggesting

that predicted increases in precipitation in the next century (52) will dramatically alter the flow

of resources from primary producers. Although we were only able to record the richness of insect

herbivory patterns, this measure is indicative of the functional diversity of insect herbivores on

Piper and changes to this value represent changes in interactions between *Piper* and higher

trophic levels (32,40). As such, the additional reduction in effects of water addition on insect

richness when *Piper* richness was low suggest that biodiversity loss in tropical systems will alter

the ability of higher trophic levels to respond to environmental perturbations.

While our prediction that increased *Piper* interspecific richness would lead to increased

insect diversity was met in the majority of sites, interspecific richness was associated with

decreased insect diversity in Costa Rica and Peru. As herbivore taxa can be differently affected

by manipulations of diversity (53), variation in the direction of the effect of intra- and

interspecific richness may be due in part to variation in the composition of insect communities

and herbivore pressure measured across sites. Changes in neighborhood effects when small

numbers of plant species dominate a community can lead either to the reduction or increase of

herbivore pressure, dependent both on the nature of plant species lost and herbivore species

present in a given community (54). As such, local variation in community composition has the

potential to greatly alter the effects of both climate change and biodiversity loss on ecosystem

function.

Although experimental methods varied somewhat between study sites, this cannot fully

explain the level of heterogeneity observed in ecosystem response. For example, the methods

employed in Costa Rica and Ecuador were nearly identical, and yet the directions of effect of

intra- and interspecific diversity on insect richness were reversed in these sites (Fig. S5 B-C).

There was considerable variation in both biotic (Fig. S3 B) and abiotic factors (Table S1) across

sites, which may have contributed to the heterogeneity observed in ecosystem response to *Piper*

diversity. Regardless of how biodiversity loss effects ecosystem function at large scales,

variation in abiotic and biotic factors at locals scales can alter these effects, reducing our ability

to predict how anthropogenic activity will alter ecosystem function. This is especially relevant in

tropical systems, which have been the subject of far fewer studies of ecosystem function than

temperate ecosystems. As such, our knowledge of local effects on the relationship between

biodiversity and ecosystem function remains limited in these systems.

A long-standing question in ecology has been the extent to which ecosystem function

increases with biodiversity and if this relationship plateaus at a level past which ecological

redundancy predominates. Recent results from less complex temperate grassland systems suggest

that these ecosystems can be described by a mostly linear relationship between richness and

function, where even rare species make unique contributions to ecosystem function (55). In these

systems, high contingency can be expected, where ecosystem-level effects will depend on most

of the interacting species. In contrast, we might expect that diverse, tropical communities could

be characterized by greater ecological redundancy and thus be subject to less contingency

(56,57). Despite these expectations, our results demonstrate heterogeneity in ecosystem response

to changes in both intraspecific and interspecific richness in five tropical sites, suggesting that

complexity in these systems may not reduce the contingency effects of biodiversity loss.

Understanding the impact of biodiversity loss in tropical forests is fundamental to our ability to

conserve those systems, and our findings highlight the importance of approaching the study of

changes in ecosystem function as context sensitive responses in complex ecosystems.

Acknowledgments

The authors would like to thank Hacienda San Isidro, Ecuador for access to their forest reserve

for experiments, the Meteorological Service from Peru (SENAMHI) for access to their weather

data, and IEF, Minas Gerais, for access to Uaimii Forest Reserve during experiments. The

collecting permit in Peru was issued by Peruvian Ministry of Agriculture (RDG N 288-2015-

SERFOR-DGGSPFFS).

Funding sources

Funding was supplied through NSF grants DEB-1442103 to LAD, LAR, TLP, AMS, and DEB-

1442075 to EJT

Author contributions:

L.A.R., L.A.D., A.M.S., T.L.P., E.J.T., J.O.S., M.J.K., and T.J.M. developed the original idea for

the research and secured funding. L.A.S., T.J.M, L.A.D, E.J.T., and H.G.L. collected Peru data.

H.G.L. and L.A.D. collected Costa Rica data. W.R.S., T.R.W., and L.A.D. collected Ecuador

data. T.J.M and A.R.N collected Mogi data. Y.A., L.B., T.J.M, and A.R.N. collected Uaimii

data. W.R.S measured herbivory for all sites. A.J.G. analyzed the data and wrote the first draft of

the manuscript. All authors contributed to analysis ideas and to writing revisions.

Competing interests: The authors declare no competing interests

460   References

1. Morris RJ. Anthropogenic impacts on tropical forest biodiversity: a network structure and ecosystem functioning perspective. Philos Trans R Soc B Biol Sci. 2010 Nov 27;365(1558):3709–18.

2. Oliver TH, Morecroft MD. Interactions between climate change and land use change on biodiversity: attribution problems, risks, and opportunities. WIREs Clim Change. 2014;5(3):317–35.

3. Ebeling A, Rzanny M, Lange M, Eisenhauer N, Hertzog LR, Meyer ST, et al. Plant diversity induces shifts in the functional structure and diversity across trophic levels. Oikos. 2018;127(2):208–19.

4. Naeem S, Li S. Biodiversity enhances ecosystem reliability. Nature. 1997 Dec;390(6659):507–9.

5. Elmqvist T, Folke C, Nyström M, Peterson G, Bengtsson J, Walker B, et al. Response diversity, ecosystem change, and resilience. Front Ecol Environ. 2003;1(9):488–94.

6. Oliver TH, Heard MS, Isaac NJB, Roy DB, Procter D, Eigenbrod F, et al. Biodiversity and Resilience of Ecosystem Functions. Trends Ecol Evol. 2015 Nov 1;30(11):673–84.

7. Easterling DR. Climate Extremes: Observations, Modeling, and Impacts. Science. 2000 Sep 22;289(5487):2068–74.

8. O'Gorman PA, Schneider T. The physical basis for increases in precipitation extremes in simulations of 21st-century climate change. Proc Natl Acad Sci. 2009 Sep 1;106(35):14773–7.

9. Field CB, Barros VR, Mastrandrea MD, Mach KJ, Abdrabo MAK, Adger WN, et al. Summary for Policy Makers: Working group 11 contribution to the fifth assessment report of the Intergovernmental panel on Climate Change [Internet]. Cambridge University Press; 2014 [cited 2023 Feb 10]. (Climate Change 2014: Impacts, Adaptation and Vulnerability - Contributions of the Working Group II to the Fifth Assessment Report). Available from: http://www.mendeley.com/research/summary-policy-makers-52

10. Fischer EM, Knutti R. Observed heavy precipitation increase confirms theory and early models. Nat Clim Change. 2016 Nov;6(11):986–91.

11. Clarke DA, York PH, Rasheed MA, Northfield TD. Does Biodiversity–Ecosystem Function Literature Neglect Tropical Ecosystems? Trends Ecol Evol. 2017 May 1;32(5):320–3.

12. Gentry AH. Tropical forest biodiversity: distributional patterns and their conservational significance. Oikos. 1992;19–28.

13. White TCR. A hypothesis to explain outbreaks of looper caterpillars, with special reference to populations of Selidosema suavis in a plantation of Pinus radiata in New Zealand. Oecologia. 1974 Dec 1;16(4):279–301.

14. Côté IM, Darling ES. Rethinking Ecosystem Resilience in the Face of Climate Change. PLOS Biol. 2010 Jul 27;8(7):e1000438.

15. Koricheva J, Larsson S, Haukioja E. Insect performance on experimentally stressed woody plants: A meta-analysis. Annu Rev Entomol. 1998;43:195.

16. Yachi S, Loreau M. Biodiversity and ecosystem productivity in a fluctuating environment: The insurance hypothesis. Proc Natl Acad Sci. 1999 Feb 16;96(4):1463–8.

17. Klaus VH, Hölzel N, Prati D, Schmitt B, Schöning I, Schrumpf M, et al. Plant diversity moderates drought stress in grasslands: Implications from a large real-world study on 13C natural abundances. Sci Total Environ. 2016 Oct 1;566–567:215–22.

18. Lanta V, Doležal J, Zemková L, Lepš J. Communities of different plant diversity respond similarly to drought stress: experimental evidence from field non-weeded and greenhouse conditions. Naturwissenschaften. 2012;99(6):473–82.

19. Hughes AR, Inouye BD, Johnson MTJ, Underwood N, Vellend M. Ecological consequences of genetic diversity. Ecol Lett. 2008;11(6):609–23.

20. Raffard A, Santoul F, Cucherousset J, Blanchet S. The community and ecosystem consequences of intraspecific diversity: a meta-analysis. Biol Rev. 2019;94(2):648–61.

21. Koricheva J, Hayes D. The relative importance of plant intraspecific diversity in structuring arthropod communities: A meta-analysis. Funct Ecol. 2018;32(7):1704–17.

22. Crutsinger GM, Collins MD, Fordyce JA, Gompert Z, Nice CC, Sanders NJ. Plant Genotypic Diversity Predicts Community Structure and Governs an Ecosystem Process. Science. 2006 Aug 18;313(5789):966–8.

23. Dyer LA, Palmer ADN, editors. Piper: A Model Genus for Studies of Phytochemistry, Ecology, and Evolution [Internet]. Boston, MA: Springer US; 2004 [cited 2021 Feb 26]. Available from: http://link.springer.com/10.1007/978-0-387-30599-8

24. Kottek M, Grieser J, Beck C, Rudolf B, Rubel F. World map of the Köppen-Geiger climate classification updated. 2006;

25. Salazar D, Jaramillo MA, Marquis RJ. Chemical similarity and local community assembly in the species rich tropical genus *Piper*. Ecology. 2016 Nov;97(11):3176–83.

26. G. Davidse, C. Ulloa Ulloa, H. M. Hernández Macías & S. Knapp (Eds.), editor. Piperaceae. In: Flora Mesoamericana. St. Louis: Missouri Botanical Garden Press; 2020. p. i–xix; 1–590.

27. Dyer LA, Letourneau DK. Trophic cascades in a complex terrestrial community. Proc Natl Acad Sci. 1999;96(9):5072–6.

28. Dyer LA, Letourneau D. Top-down and bottom-up diversity cascades in detrital vs. living food webs. Ecol Lett. 2003;6(1):60–8.

29. Dyer LA, Letourneau DK, Dodson CD, Tobler MA, Stireman III JO, Hsu A. Ecological causes and consequences of variation in defensive chemistry of a Neotropical shrub. Ecology. 2004;85(10):2795–803.

30. Letourneau DK, Dyer LA, Vega C G. Indirect effects of a top predator on a rain forest understory plant community. Ecology. 2004;85(8):2144–52.

31. Connahs H, Rodríguez-Castañeda G, Walters T, Walla T, Dyer L, Miller J. Geographic variation in host-specificity and parasitoid pressure of an herbivore (Geometridae) associated with the tropical genus Piper (Piperaceae). J Insect Sci. 2009;9(1).

32. Dyer LA, Letourneau DK, Chavarria GV, Amoretti DS. Herbivores on a dominant understory shrub increase local plant diversity in rain forest communities. Ecology. 2010;91(12):3707–18.

33. Bodner F, Strutzenberger P, Brehm G, Fiedler K. Species richness and host specificity among caterpillar ensembles on shrubs in the Andes of southern Ecuador. Neotrop Entomol. 2012;41(5):375–85.

34. Abarca M, Boege K, Zaldívar-Riverón A, Giray T. Shelter-building behavior and natural history of two pyralid caterpillars feeding on Piper stipulaceum. J Insect Sci. 2014;14(1).

35. Glassmire AE, Jeffrey CS, Forister ML, Parchman TL, Nice CC, Jahner JP, et al. Intraspecific phytochemical variation shapes community and population structure for specialist caterpillars. New Phytol. 2016;212(1):208–19.

36. Slinn HL, Richards LA, Dyer LA, Hurtado PJ, Smilanich AM. Across multiple species, phytochemical diversity and herbivore diet breadth have cascading effects on herbivore immunity and parasitism in a tropical model system. Front Plant Sci. 2018;9:656.

37. Cosmo LG, Nascimento AR, Cogni R, Freitas AV. Temporal distribution in a tri-trophic system associated with Piper amalago L. in a tropical seasonal forest. Arthropod-Plant Interact. 2019;13(4):647–52.

38. Campos-Moreno DF, Dyer LA, Salcido D, Massad TJ, Pérez-Lachaud G, Tepe EJ, et al. Importance of interaction rewiring in determining spatial and temporal turnover of tritrophic (Piper-caterpillar-parasitoid) metanetworks in the Yucatán Península, México. Biotropica. 2021;

39. Abatzoglou JT, Dobrowski SZ, Parks SA, Hegewisch KC. TerraClimate, a high-resolution

global dataset of monthly climate and climatic water balance from 1958–2015. Sci Data. 2018 Jan 9;5(1):170191.

40. Carvalho MR, Wilf P, Barrios H, Windsor DM, Currano ED, Labandeira CC, et al. Insect Leaf-Chewing Damage Tracks Herbivore Richness in Modern and Ancient Forests. PLOS ONE. 2014 May 2;9(5):e94950.

41. Fisher RA. XV.—The correlation between relatives on the supposition of Mendelian inheritance. Earth Environ Sci Trans R Soc Edinb. 1919;52(2):399–433.

42. Henderson CR, Kempthorne O, Searle SR, von Krosigk CM. The Estimation of Environmental and Genetic Trends from Records Subject to Culling. Biometrics. 1959;15(2):192–218.

43. Abelson R. P.(1995). Statistics as principled argument. Hillsdale, NJ: Lawrence Erlbaum;

44. Kimmel K, Dee LE, Avolio ML, Ferraro PJ. Causal assumptions and causal inference in ecological experiments. Trends Ecol Evol. 2021 Dec 1;36(12):1141–52.

45. Gelman A, Rubin DB. Inference from Iterative Simulation Using Multiple Sequences. Stat Sci. 1992 Nov;7(4):457–72.

46. Gelman A, Meng XL, Stern H. Posterior predictive assessment of model fitness via realized discrepancies. Stat Sin. 1996;733–60.

47. Kellner K, Kellner MK, SystemRequirements J. Package 'jagsUI.' 2021;

48. Gelman A, Tuerlinckx F. Type S error rates for classical and Bayesian single and multiple comparison procedures. Comput Stat. 2000 Sep 1;15(3):373–90.

49. Makowski D, Ben-Shachar MS, Chen SHA, Lüdecke D. Indices of Effect Existence and Significance in the Bayesian Framework. Front Psychol. 2019;10:2767.

50. R. Core Team. R: A language and environment for statistical computing. 2013;

51. Cook-Patton SC, McArt SH, Parachnowitsch AL, Thaler JS, Agrawal AA. A direct comparison of the consequences of plant genotypic and species diversity on communities and ecosystem function. Ecology. 2011 Apr;92(4):915–23.

52. Field CB, Barros VR, Intergovernmental Panel on Climate Change, editors. In: Climate change 2014: impacts, adaptation, and vulnerability: Working Group II contribution to the fifth assessment report of the Intergovernmental Panel on Climate Change. New York, NY: Cambridge University Press; 2014.

53. Agrawal AA, Lau JA, Hambäck PA. Community heterogeneity and the evolution of interactions between plants and insect herbivores. Q Rev Biol. 2006 Dec;81(4):349–76.

54. Barbosa P, Hines J, Kaplan I, Martinson H, Szczepaniec A, Szendrei Z. Associational Resistance and Associational Susceptibility: Having Right or Wrong Neighbors. Annu Rev Ecol Evol Syst. 2009 Dec;40(1):1–20.

55. Isbell F, Calcagno V, Hector A, Connolly J, Harpole WS, Reich PB, et al. High plant diversity is needed to maintain ecosystem services. Nature. 2011 Sep;477(7363):199–202.

56. Naeem S. Species Redundancy and Ecosystem Reliability. Conserv Biol. 1998;12(1):39–45.

57. Rosenfeld JS. Functional redundancy in ecology and conservation. Oikos. 2002;98(1):156–62.

58. Root RB. Organization of a Plant-Arthropod Association in Simple and Diverse Habitats: The Fauna of Collards (Brassica Oleracea). Ecol Monogr. 1973;43(1):95–124.

59. Gely C, Laurance SGW, Stork NE. How do herbivorous insects respond to drought stress in trees? Biol Rev. 2020;95(2):434–48.

60. Lenhart PA, Eubanks MD, Behmer ST. Water stress in grasslands: dynamic responses of plants and insect herbivores. Oikos. 2015;124(3):381–90.

 Figures and tables

474 **Table 1.** *Path models and explanatory hypotheses.* Black arrows indicate causal paths between

475 variables

Hypothesis	Causal paths	Sites tested
Bottom-up diversity: Host plant diversity impacts insect richness through multiple mechanisms, such as reducing host density, masking or amplifying host signals, or altering the proportion of specialist herbivores (53,54,58)	All models; paths A, C	All sites
Neighborhood effects: Plant diversity directly affects herbivory through mechanisms which do not alter herbivore diversity (53,54)	All models; paths B, D	All sites
Water affects herbivore diversity: Changes in water availability induce changes to plant nutrition and defenses which can benefit or harm different herbivore taxa, leading to changes in herbivore diversity (59,60)	Models I, II; path G	Costa Rica, Ecuador, Peru
Water affects herbivory: Water addition directly affects plant physiology, altering both the nutritive quality of plant tissue and the ability of plants to combat herbivores and leading to changes in herbivory (13)	Models I, III; path F	Costa Rica, Ecuador, Peru

Figure 1. *Treatments of intraspecific richness, interspecific richness, and water addition used in each of the five study sites*. White tiles represent treatment combinations which were not tested in a given site

Figure 2. *Hierarchical Bayesian model parameter estimates for the effects of water availability,*

as well as intraspecific and interspecific Piper *richness on herbivory, herbivorous insect*

richness, and Piper *survivorship across sites*

Violins represent the cross-site posterior parameter distribution for each relationship in site level

hierarchical Bayesian models. Black lines represent the median posterior estimate and white bars

represent 95% credible intervals. Percentages above violins indicate the probability of an effect

being positive or negative (as indicated by a negative probability) in response to an increase of

the independent variable. Distributions for water addition compare watered and control plots;

distributions for interspecific richness compare *Piper* species richness standardized as the

proportion of the maximum richness used at a site; distributions for intraspecific richness

compare low and high intraspecific richness treatments; distributions for insect richness compare

responses per insect taxon present on an individual leaf

Figure 3. *Direct and indirect effects of plant diversity and water availability on insect*

herbivores at five study sites. Bayesian structural equation models comparing effects of different

drivers of herbivorous insect richness and herbivory at five sites. Standardized path coefficients

are means of the posterior distribution for the effects estimated at each causal path. Positive

relationships are indicated in blue with triangular heads, and negative relationships are indicated

in red with circular heads. Black arrows indicate path coefficients of zero magnitude. Dot plots

545   summarize the standardized mean of the posterior distribution for each causal path with 95% and

546   80% credible intervals. Asterisks indicate causal paths where the probability of an effect being

547   positive or negative is >95%

Figure 4. *Interactions between intraspecific or interspecific richness and water availability on*

herbivory, herbivorous insect richness, and Piper *survival*

Bars indicate mean response and standard error of the mean. Percentages above each sub-figure

indicate the probability that the two slopes are different as calculated using site level hierarchical

589   Bayesian models. Due to high mortality in Peru, interactions between water and intraspecific

590   richness could not be compared for any responses except mortality. Single and two species

591   richness plots have been combined for visualization purposes only

Supplementary figures and tables

Table S1. *Study site characteristics and experimental details*

	La Selva Biological Station	Yanayacu Biological Station	El Fundo Génova	Mogi-Guaçu Biological Reserve	Uaimii State Forest
Location	N9.9400, W84.0433	S0.6008, W77.8903	S11.0948, W75.3516	S32.2510, W47.1558	S30.2957, W43.5731
Mean elevation (m a.s.l.)	35	2124	1133	627	800
Mean annual precipitation (mm)	4495	2900	1767	1271	1600
Mean annual T (°C)	25	16	23	26	19
Date planted	2015-03-01	2015-07-01	2015-10-01	2017-02-01	2018-02-20
Date harvested	2017-02-01	2016-12-01	2017-07-01	2019-12-01	2019-12-20
Number of species in high richness plots	12	12	6	4	3
Planted in pots / ground	Pots	Pots	Pots	Ground	Ground
Water addition treatment applied	Yes	Yes	Yes	No	No
Number of *Piper* planted	360	360	360	432	144
Number of *Piper* surviving till the end of the experiment	200	259	56	182	123

*Climate data from La Selva Biological station are for the experimental period (2015-2018) and were provided by the Organization for Tropical Studies. Data from El Fundo Génova are based on nearby San Ramón (https://en.climate-data.org/south-america/peru/junin/san-ramon-28556/). Data from Ecuador were provided by Yanayacu Biological Station. Data for Mogi-Guaçu Biological Reserve are from January 2017-December 2019 and are from the Centro Integrado de Informações Agrometeorológicas of São Paulo. Data for Uaimii State Forest are from the Plano de Manejo FLOE Uaimii, Instituto Estadual de Florestas of Minas Gerais

626   **Table S2.** Species of *Piper* used at each study location

Study location				
La Selva Biological Station	Yanayacu Biological Station	El Fundo Génova	Mogi-Guaçu Biological Reserve	Uaimii State Forest
arboreum	grande	arboreum	arboreum	lepturum
peltatum	peludo	reticulatum	crassinervium	corcovadense
biolleyi	ecuadorense	glabribaccum	richardifolium	vicosanum
sancti-felicis	perareolatum	chanchamayanum	miquelianum	
reticulatum	baezanum	armatum		
nudifolium	escabrosa	lechlerianum		
multiplinervium	kelleii			
decurrens	hispidum			
pseudobumbratum	schupii			
imperiale	pubinervulum			
garagaranum	pequenia			
urostachyum				

641   **Table S3.** Mean parameter estimates and probability of direction (PD) for the effects of increases

642   in intraspecific diversity, interspecific richness, water availability and insect richness on

643   measures of herbivory, plant mortality, and insect richness

Site	Predictor variable	Response variable	Mean parameter estimate	PD
All	Intraspecific richness	Percent herbivory	-0.87	63.59%
All	Intraspecific richness	Presence of damage	-3.26	71.60%
All	Intraspecific richness	Variance in herbivory	-9.97	57.31%
All	Intraspecific richness	Insect richness	-0.02	59.41%
All	Intraspecific richness	Plant survival	6.50	92.48%
All	Insect richness	Percent herbivory	8.78	100.00%
All	Insect richness	Presence of damage	6.63	96.73%
All	Insect richness	Variance in herbivory	17.86	71.13%
All	Insect richness	Plant survival	3.35	66.21%
All	Interspecific richness	Percent herbivory	0.02	50.34%
All	Interspecific richness	Presence of damage	7.40	83.01%
All	Interspecific richness	Variance in herbivory	-52.43	73.46%
All	Interspecific richness	Insect richness	0.20	95.00%
All	Interspecific richness	Plant survival	10.66	80.22%
All	Water availability	Percent herbivory	-4.19	98.73%
All	Water availability	Presence of damage	-6.33	89.46%
All	Water availability	Variance in herbivory	-118.10	91.35%
All	Water availability	Insect richness	-0.07	64.52%
All	Water availability	Plant survival	5.15	79.29%
Costa Rica	Intraspecific richness	Percent herbivory	-0.10	52.68%
Costa Rica	Intraspecific richness	Presence of damage	2.86	79.84%
Costa Rica	Intraspecific richness	Variance in herbivory	37.95	72.01%
Costa Rica	Intraspecific richness	Insect richness	0.10	98.07%
Costa Rica	Intraspecific richness	Plant survival	3.72	80.91%
Costa Rica	Insect richness	Percent herbivory	7.53	100.00%
Costa Rica	Insect richness	Presence of damage	5.60	100.00%
Costa Rica	Insect richness	Variance in herbivory	-6.74	65.69%
Costa Rica	Insect richness	Plant survival	3.60	66.25%
Costa Rica	Interspecific richness	Percent herbivory	-2.43	91.53%
Costa Rica	Interspecific richness	Presence of damage	-4.89	89.55%
Costa Rica	Interspecific richness	Variance in herbivory	-89.88	83.97%
Costa Rica	Interspecific richness	Insect richness	-0.08	93.43%
Costa Rica	Interspecific richness	Plant survival	18.69	99.61%
Costa Rica	Water availability	Percent herbivory	-4.76	99.99%
Costa Rica	Water availability	Presence of damage	-7.96	99.90%
Costa Rica	Water availability	Variance in herbivory	-151.05	98.04%

Costa Rica	Water availability	Insect richness	-0.02	62.33%
Costa Rica	Water availability	Plant survival	12.09	99.74%
Ecuador	Intraspecific richness	Percent herbivory	-3.50	99.87%
Ecuador	Intraspecific richness	Presence of damage	-8.10	99.95%
Ecuador	Intraspecific richness	Variance in herbivory	-80.24	92.45%
Ecuador	Intraspecific richness	Insect richness	-0.18	100.00%
Ecuador	Intraspecific richness	Plant survival	20.46	100.00%
Ecuador	Insect richness	Percent herbivory	8.68	100.00%
Ecuador	Insect richness	Presence of damage	8.23	100.00%
Ecuador	Insect richness	Variance in herbivory	22.77	97.99%
Ecuador	Insect richness	Plant survival	7.46	81.29%
Ecuador	Interspecific richness	Percent herbivory	-1.74	89.56%
Ecuador	Interspecific richness	Presence of damage	8.46	99.73%
Ecuador	Interspecific richness	Variance in herbivory	16.01	57.91%
Ecuador	Interspecific richness	Insect richness	0.58	100.00%
Ecuador	Interspecific richness	Plant survival	7.78	85.86%
Ecuador	Water availability	Percent herbivory	-2.67	99.52%
Ecuador	Water availability	Presence of damage	-5.72	99.78%
Ecuador	Water availability	Variance in herbivory	-28.52	68.94%
Ecuador	Water availability	Insect richness	0.04	85.57%
Ecuador	Water availability	Plant survival	5.51	90.57%
Mogi	Intraspecific richness	Percent herbivory	-2.15	98.59%
Mogi	Intraspecific richness	Presence of damage	-4.29	97.74%
Mogi	Intraspecific richness	Variance in herbivory	-24.33	67.12%
Mogi	Intraspecific richness	Insect richness	0.02	75.41%
Mogi	Intraspecific richness	Plant survival	1.57	66.65%
Mogi	Insect richness	Percent herbivory	11.17	100.00%
Mogi	Insect richness	Presence of damage	12.22	100.00%
Mogi	Insect richness	Variance in herbivory	80.94	99.99%
Mogi	Insect richness	Plant survival	1.95	58.03%
Mogi	Interspecific richness	Percent herbivory	2.02	90.42%
Mogi	Interspecific richness	Presence of damage	6.39	97.05%
Mogi	Interspecific richness	Variance in herbivory	-116.52	90.64%
Mogi	Interspecific richness	Insect richness	0.04	80.94%
Mogi	Interspecific richness	Plant survival	17.14	99.05%
Peru	Intraspecific richness	Percent herbivory	-0.09	51.56%
Peru	Intraspecific richness	Presence of damage	-9.11	94.39%
Peru	Intraspecific richness	Variance in herbivory	-58.63	73.32%
Peru	Intraspecific richness	Insect richness	-0.03	63.89%
Peru	Intraspecific richness	Plant survival	2.84	74.08%
Peru	Insect richness	Percent herbivory	8.09	100.00%
Peru	Insect richness	Presence of damage	4.38	99.51%
Peru	Insect richness	Variance in herbivory	30.07	87.18%
Peru	Insect richness	Plant survival	-2.31	61.10%

Peru	Interspecific richness	Percent herbivory	0.14	50.60%
Peru	Interspecific richness	Presence of damage	16.49	97.23%
Peru	Interspecific richness	Variance in herbivory	-69.46	70.67%
Peru	Interspecific richness	Insect richness	0.16	91.04%
Peru	Interspecific richness	Plant survival	5.70	77.33%
Peru	Water availability	Percent herbivory	-5.15	99.75%
Peru	Water availability	Presence of damage	-5.27	89.79%
Peru	Water availability	Variance in herbivory	-173.87	94.77%
Peru	Water availability	Insect richness	-0.22	99.69%
Peru	Water availability	Plant survival	-2.10	69.02%
Uaimii	Intraspecific richness	Percent herbivory	1.51	71.49%
Uaimii	Intraspecific richness	Presence of damage	2.21	64.77%
Uaimii	Intraspecific richness	Variance in herbivory	74.93	79.39%
Uaimii	Intraspecific richness	Insect richness	-0.01	55.41%
Uaimii	Intraspecific richness	Plant survival	3.92	77.51%
Uaimii	Insect richness	Percent herbivory	8.41	100.00%
Uaimii	Insect richness	Presence of damage	2.83	85.39%
Uaimii	Insect richness	Variance in herbivory	-36.56	83.52%
Uaimii	Insect richness	Plant survival	5.98	75.15%
Uaimii	Interspecific richness	Percent herbivory	2.23	70.86%
Uaimii	Interspecific richness	Presence of damage	10.61	87.21%
Uaimii	Interspecific richness	Variance in herbivory	-3.20	52.59%
Uaimii	Interspecific richness	Insect richness	0.27	98.14%
Uaimii	Interspecific richness	Plant survival	3.53	62.35%

Figure S1. *Treatments and number of plots used across sites*

A) Treatments of intraspecific richness, interspecific richness, and water addition at the

beginning of the experimental period at each of the five study sites, and B) final treatments at the

end of the experimental period in each site. White numerals indicate number of plots used.

Changes in *Piper* species richness and the loss of some treatment combinations was due to *Piper*

mortality over the course of the experimental period

Figure S2. *Precipitation levels at study sites where the water addition treatment was applied*

A) Monthly precipitation over the experimental period at Costa Rica (CR), Ecuador (EC), and

Peru (PE). B) difference in monthly precipitation from climate normals at the three sites across

the experimental periods for those sites. Bars indicate median values, black points indicate mean

values. C) Cumulative precipitation over the course of the experimental period. Dotted lines

indicate average cumulative precipitation, shaded regions indicate 95%, 80% and 50% quantiles.

Climate normals are based on monthly precipitation for a period from 1958 to 1998

Figure S3. *Overall herbivory, plant mortality, and insect richness at five study sites*

A) Percent herbivory, percentage of leaves with any damage, herbivorous insect richness, and

percent mortality of *Piper* in Costa Rica (CR), Ecuador (EC), Peru (PE), Uaimii (UI), and Mogi-

Guaçu (MG), across treatments. B) Proportion of herbivorous insect taxa observed at each site

measured by feeding damage patterns. Colored regions indicate the percentage of damage

observations contributed by each taxon. C) Proportion of herbivory attributed to each taxon at

five study sites. Colored regions represent the percentage of each leaf consumed by each taxon at

each site. White tiles represent sites where no leaf damage by that insect taxon was observed

Figure S4. *HBM parameter estimates of percent herbivory, percentage of leaves with damage,*

variance in herbivory, and percent Piper *survival against levels of A) water addition, B)* Piper

intraspecific richness, C) Piper *interspecific richness, and D) insect richness at each site.*

Violins represent the posterior parameter distribution for each relationship across sites and within

sites in Costa Rica (CR), Ecuador (EC), Peru (PE), Uaimii (UI), and Mogi-Guaçu (MG). Black

lines represent the median posterior estimate while white bars represent 95% credibility

intervals. Percentages above violins indicate the probability of an effect being entirely positive or

entirely negative in response to an increase of the manipulated variable. Distributions for water

addition compare unwatered control and watered plots; distributions for interspecific richness

compare *Piper* species richness and are standardized as a percentage of the high richness

treatment level; distributions for intraspecific richness compare low and high intraspecific

richness plots; distributions for insect richness compare responses per insect taxon present at the

leaf level

Figure S5. *HBM posterior parameter estimates of insect richness compared to levels of A) water*

addition, B) Piper *intraspecific richness, and C)* Piper *interspecific richness*

Violins represent the posterior parameter distribution for each relationship across sites and within

sites at Costa Rica (CR), Ecuador (EC), Peru (PE), Uaimii (UI), and Mogi-Guaçu (MG). Black

lines represent the median posterior estimate while white bars represent 95% credibility

intervals. Percentages above violins indicate the probability of an effect being entirely positive or

entirely negative in response to an increase of the manipulated variable. Distributions for water

addition compare the unwatered control and watered plots; distributions for interspecific richness

compare *Piper* species richness and are standardized as a percentage of the high richness

treatment level; distributions for intraspecific richness compare low and high intraspecific

richness plots

Figure S6. *Bayesian structural equation models for drivers of insect richness, herbivory and*

Piper survival at three sites, including interactions between intraspecific and interspecific

richness, and water addition. Path coefficients indicate the standardized mean of the posterior

distribution for each causal path. Positive relationships are indicated in blue with triangular

heads, and negative relationships are indicated in red with circular heads. Dot plots indicate the

standardized mean of the posterior distribution for each causal path in A with 95% and 80%

credibility intervals. Asterisks indicate causal paths where the probability of an effect being

entirely positive or entirely negative is > 95%

Figure S7. *Percent Piper survival over time in five sites, compared to levels of intraspecific*

richness and water addition

50