1 **Title**
2 **The commensal protist *Tritrichomonas musculus* exhibits a dynamic life cycle that induces B cell-**
3 **modulated remodeling of the gut microbiota**
4
5 **Authors**

6 Ana Popovic[1,2], Eric Yixiao Cao[3], Joanna Han[3], Nirvana Nursimulu[1,4], Eliza V.C. Alves-Ferreira[5], Andrea
7 Kennard[5], Noor Alsmadi[3], Michael E. Grigg[5], Arthur Mortha[3], John Parkinson[1,2,6],*
8
Affiliations
[1]Program in Molecular Medicine, Hospital for Sick Children, Toronto, Ontario, Canada
[2]Department of Biochemistry, University of Toronto, Toronto, Ontario, Canada
[3]Department of Immunology, University of Toronto, Toronto, Ontario, Canada
[4]Department of Computer Science, University of Toronto, Toronto, Ontario, Canada
[5]Molecular Parasitology Section, Laboratory of Parasitic Diseases, NIAID, National Institutes of Health,
Bethesda, Maryland, USA
[6]Department of Molecular Genetics, University of Toronto, Toronto, Ontario, Canada
17
18 **Corresponding authors**
19 *To whom correspondence should be addressed:
20 john.parkinson@utoronto.ca
21
22

23 Keywords: Eukaryotic microbiota; Parasites; Tritrichomonas; Microbiome; Gut pathogens
24
25

Summary

Commensal protists and gut bacterial communities exhibit complex relationships, mediated at least in part through host immunity. To improve our understanding of this tripartite interplay, we investigated community and functional dynamics between the murine protist *Tritrichomonas musculus* (*T. mu*) and intestinal bacteria in healthy and B cell-deficient mice. We identified dramatic, *T. mu*-dependent remodeling of resident bacterial composition and metabolism in parallel to increases in *T. mu* metabolism and the expression of putative virulence factors, responses accelerated in the absence of B cells. Upregulation of iron acquisition and tryptophan consumption revealed competition between bacteria and the protist. Single cell transcriptomics identified distinct *T. mu* life stages, providing evidence for active growth, sexual replication and the formation of pseudocysts. Unique cell states were validated *in situ*, and appeared modulated by constituent bacteria, suggesting their role in protist development. Our results reveal complex microbial dynamics during the establishment of a commensal protist.

Introduction

Intestinal microbiota are key regulators of host health and the development of disease[1,2]. The majority of studies currently address roles of bacteria and fungi, however emerging data supports an appreciable presence of protists in the gut[3-5]. These include species considered commensals, diarrheagenic parasites and pathobionts that exert influential roles on resident bacterial communities. Conversely, bacterial communities have been shown to impact the ability of protists to successfully colonize[6,7]. In addition to direct interactions such as competition for resources, protist-bacterial cross-talk appears largely mediated through the modulation of host immune pathways[8,9]. Currently little is known concerning the molecular basis which drive these complex protist:host:bacterial interactions.

Among protists capable of both asymptomatic gut colonization and virulence are members of the trichomonads, microbes prevalent in the human gut with links to colitis, irritable bowel syndrome and colon cancer[5,10-13]. In mice, trichomonads *Tritrichomonas musculus* (*T. mu*) and *Tritrichomonas muris* establish chronic asymptomatic infections, which result in elevated baseline intestinal immune activation, dependent on increases in IFN-gamma producing Th1 and interleukin-17 producing Th17 cells. Although sustained inflammation from prolonged *T. mu* colonization can increase susceptibility to T-cell-driven colitis, it also confers protection to mice against enteric infections with *Salmonella*[9,14-16]. The stimulation of intestinal tuft cells by *Tritrichomonas*-produced succinate furthermore leads to a Th2-based immune response, triggering structural remodelling of the intestinal epithelium, and ultimately restricts colonization by parasitic helminths[14,19,20]. Colonization by *Tritrichomonas* spp. has also been shown to shift commensal bacterial diversity and the abundance of *Helicobacter* and Lactobacillaceae taxa[17,18]. Similarly, bacteria can modulate *T. mu* engraftment. For example, *Bifidobacterium* spp. together with inulin supplements were shown to influence *T. mu* colonization in the context of a high fat diet, suggesting complex microbial interactions[17]. These interactions between protist and bacteria may also be driven through immunoglobulins (Ig) A and M, antibodies secreted by B cells and elevated in response to *T. mu* colonization[9,14]. Although both bacteria

and protists are known to impact B cell antibody responses, it is not well understood what role B cells play

in modulating the physiology of a complex community comprised of members from diverse microbial

kingdoms[9,21,22].

Here, we begin to unravel the tripartite interactions between *T. mu*, resident gut bacteria and the host through

the application of 16S rRNA sequence surveys and transcriptomics to microbial communities in healthy

(WT) and B cell-deficient (*muMT*[-/-]) mice. We show that introduction of *T. mu* induces dramatic

diversification of gut bacterial populations in both host genotypes during the first 28 days of colonization.

Shifts in bacterial growth and metabolism, in response to protist colonization, track with upregulation of *T.*

mu core metabolic processes and the expression of putative virulence factors, changes which are accelerated

in the absence of B cells. To dissect the activities of *T. mu* in the mouse intestinal tract at high resolution,

we performed single cell transcriptomics. This exceptional dataset reveals actively growing and replicating

trophozoites, and evidence of *T. mu* sexual replication and a pseudocyst stage. Using fluorescent RNA

labelling of uniquely expressed genes, we validate these protist cell states *in situ*. Finally, we describe

potential cross-kingdom competition for resources between protist and resident bacteria, specifically in the

acquisition and utilization iron and key amino acids tryptophan and arginine. Our data demonstrate the co-

adaptation of resident bacteria and a new protist commensal, which together reshape the gut environment

in a manner dependent on the host immune response.

Results

***T. mu* colonization drives diversification of the intestinal microbiome**

Colonization with *T. mu* induces profound changes in the mouse intestinal immune landscape, including

increased levels of host IgA, an abundant antibody known to maintain gut microbial homeostasis[9,22]. To

explore the downstream impact on resident gut microbiota, and how this may be influenced by antibody-

producing B cells, we measured microbial composition and gene expression during the first 28 days of *T.*

mu colonization in WT and B cell-deficient (*muMt*[-/-]) C57Bl/6 mice, through 16S and metatranscriptomics

sequencing (Figure 1A). Protists expanded rapidly in mouse caeca, in both the presence and absence of B

cells, reaching 54 million and 68 million cells respectively (Figure 1B). Host IgA levels in WT mice tracked

T. mu growth, and were significantly higher relative to naïve mice at both days 2 and 28 ($p < 0.05$)

confirming previous reports (Figure 1C)[23].

The introduction of *T. mu* dramatically altered gut bacterial composition in both WT and *muMt*[-/-] mice

(Figure 2A). Sequencing of fecal 16S revealed 470 bacterial taxa (OTUs with ≥5 reads) in mouse fecal

samples, with median 176 OTUs per sample (Figure 2A and Tables S1 and S2). Naïve mice were dominated

by Bacteroidetes and Firmicutes, including members of Muribaculaceae, Bacteroidaceae, Lactobacillaceae

and Lachnospiraceae (Figure 2A). Upon colonization with *T. mu*, bacterial richness (observed OTUs)

increased, with a delayed increase in *muMt*[-/-] mice, before receding to near day 0 levels (LME, day 7 vs day

0, $\beta = 94.7$, 95% CI [49.0, 140.5], $p < 0.001$; day 14 vs day 0 in *muMt*[-/-], $\beta = 66.4$, 95% CI [1.8, 131.1], $p <$

0.05; two-way ANOVA, day 28 colonized vs naïve, *ns*) (Figure 2B). Increases in evenness (Shannon

diversity), however, persisted to day 28 across both host genotypes, reflecting expansions in multiple taxa

from Firmicutes, Bacteroidetes, Actinobacteria and Proteobacteria clades (LME, day 2 vs day 0 x host

genotype, $\beta = -0.52$, 95% CI [-1.02, -0.01], $p < 0.05$; day 7 vs day 0 x host genotype, $\beta = -1.00$, 95% CI [-

1.50, -0.49], $p < 0.001$; days 2 to 28 vs day 0, $p < 0.001$; two-way ANOVA, day 28 colonized vs naïve,

$F(1,12) = 7.06$, $p < 0.05$) (Figures 2A and 2B). A spike in evenness during the first week in WT hosts is the

result of rapid initial expansions of multiple taxa compared to a single Lachnospiraceae taxon in *muMt*[-/-]

mice (Figure 2A). While bacterial communities in WT and *muMt*[-/-] mice differed prior to infection, their

compositions changed progressively after day 2 of protist colonization and trended toward a convergent

community profile (Adonis, time x host, $F(5,55) = 6.2$, $p < 0.001$; pre-infected vs day 2, *ns*) (Figure 2C).

That compositions in day 28 control mice remained most similar to those at day 0, prior to *T. mu*

colonization, underlines a protist-driven shift (Adonis, day 28 ctl vs day 0, protist x host genotype, $F_{(1,15)}$

= 5.7, $p < 0.05$; *ns* in WT mice). We identified 55 taxa that changed significantly as a result of *T. mu* ($p <$

0.05), clustered into six distinct dynamics patterns (Figure 2D). The majority of bacteria increased in

abundance in both WT and *muMt*$^{-/-}$ hosts: 26 taxa continued expanding until the termination of the

experiment (including *Bifidobacterium*, *Parabacteroides*, *Alistipes* and two lactobacilli) and 10 taxa peaked

at day 21 after which their abundances fell (e.g. the Proteobacteria *Desulfovibrio* and *Helicobacter*). Taxa

whose abundances decreased as a result of *T. mu* colonization (11 taxa) included *Blautia* and several other

clostridia, taxa from the Muribaculaceae family and *Enterorhabdus*. Five taxa exhibited discordant patterns

between host genotypes, including *Parasutterella*, *Bacteroides*, and a species of Lachnospiraceae which

experienced temporary reductions in WT mice but expanded significantly in *muMt*$^{-/-}$ mice; and two species

of Muribaculaceae which displayed the opposite response. We identified 82 taxa with host genotype-

dependent patterns, independent of the protist, however as WT and *muMt*$^{-/-}$ mice were not co-housed, we

cannot rule out caging effects. Changes in bacterial abundances were validated through qPCR using clade-

specific primers for four groups: Firmicutes, Bacteroidetes, Actinobacteria and Gammaproteobacteria (day

21 vs day 0, $p < 0.05$, except Firmicutes in *muMT*$^{-/-}$, $p = 0.057$, and Bacteroidetes in WT and *muMT*$^{-/-}$, $p =$

0.057) (Figure 2E).

***T. mu* induces changes in bacterial metabolism**

To evaluate how *T. mu*-induced shifts in gut bacterial composition translate to changes in community

function, we performed bulk RNA sequencing (metatranscriptomics) of mouse caecal contents at days 2

and 28 of the experiment, timepoints reflecting early and chronic stages of infection (Figure 3A).

Sequencing produced a total of 2.4 billion paired reads across 32 samples (median 76.8 million) (Table S3).

After filtering for low quality, host and bacterial rRNA, 835 million reads were identified as bacterial (741

million in naïve and 94 million in colonized mice), 157 million as *T. mu* and 349 million reads remained

unannotated (293 million in naïve and 56 million in colonized mice) (Figure S1 and Table S3). The majority

of reads mapping to bacterial transcripts belonged to Firmicutes taxa (597 million), followed by Bacteroidetes (75.4 million) and Proteobacteria (21.5 million) (Figure 3A). Colonization with *T. mu* associated with blooms in Bacteroidetes and particularly Proteobacteria activity. Bacteroidetes transcripts increased in proportion from 14.9% to 26.6% in WT colonized mice and from 12.4% to 24.2% in *muMT*[-/-] colonized mice, whereas naïve mice displayed relatively modest increases (9.1% to 11.5% in WT; 5.9% to 7.6% in *muMT*[-/-]) (Figure 3A). Changes in proteobacterial activity were more striking, varying between 1% and 3.4% in naïve mice, but increasing from 4% to 15.6% of reads in WT mice by day 28, and already accounting for 13.7% of reads by day 2 in *muMT*[-/-] mice, suggesting accelerated increases and a possible role for B cells in regulating these taxa. The majority of these reads belonged the *Helicobacteraceae* clade (9.1% WT and 6.2% *muMT*[-/-] at day 28), bacteria previously associated with enteric inflammation and exacerbated disease in T cell-driven colitis [9,14,24].

The presence of *T. mu* RNA, rRNA in particular, substantially reduced transcriptional information from bacteria in colonized mice, from a median of 22.1 million reads per naïve mouse to 2.1 million (Table S3). Protozoan rRNA is not captured by commercial rRNA depletion kits used in sequencing library preparation. To account for the variation in sequencing depth, we evaluated gene expression in data rarefied to 600,000 bacterial reads per sample. Principal component analysis of bacterial gene expression re-capitulated patterns observed in 16S compositional data (Figures 2C and 3B). Bacterial expression differed between host genotypes, and progressively changed after introduction of *T. mu*. The protist induced overall upregulation of bacterial genes, with strong responses in *muMt*[-/-] mice already by day 2 (Figure 3C). Interestingly, host genotype-related differences decreased over time in both colonized and naïve mice (Figure 3C).

The analysis of bacterial metabolic pathways revealed an association between *T. mu* colonization and increased bacterial glycolysis, carbohydrate metabolism, peptidoglycan, polysaccharide and O-Antigen

sugar biosynthesis, and the biosynthesis and metabolism of numerous amino acids (Figure 3D). Amino acid

biosynthesis and metabolism, in particular, showed earlier upregulation in the absence of B cells. At day 2,

we detected metabolism primarily in Firmicutes taxa, but noted shifts toward Bacteroidetes and

Proteobacteria by day 28 (Figures S2 and S3). The expansion of Proteobacteria (facultative anaerobes) and

increased cycling through succinyl-CoA synthetase (SucC) in the TCA cycle, an enzyme suppressed under

anaerobic conditions[25], may suggest oxygen exposure, characteristic of gut inflammation and dysbiosis

(Figure S2). Increasing metabolism, and particularly the expression of enzymes required for biosynthesis

of surface lipopolysaccharides (LPS) and O-antigen sugars, the variable polysaccharide coating the outer

LPS layer, suggests that the establishment of a chronic infection by *T. mu* may induce bacterial growth in

the caecum, and specifically the growth of particular taxa.

To further explore this notion, we probed for changes in the expression of bacterial cell division and cell

wall biosynthetic machinery. We first identified homologues to 44 proteins involved in these processes, as

previously identified in the model *E. coli* system, many of which are conserved across bacteria and form a

network of functionally associated protein modules[26] (Figure 3E). Homologues to 40 of these proteins were

detected in the experiment, and 27 were present in the rarefied dataset (represented by 667 unique genes).

Patterns of gene expression were broadly consistent with those described in other pathways—genes were

upregulated during *T. mu* colonization, with earlier responses in B cell-deficient mice (12 bacterial genes

upregulated in *muMT*$^{-/-}$ mice at day 2, compared to three in WT mice), and a shift from predominantly

Firmicutes gene expression at day 2 to Bacteroidetes (primarily *Bacteroides* and *Parabacteroides*) and

Proteobacteria (primarily *Helicobacter*) at day 28, suggesting an environment favouring gram-negative

taxa. At day 2, gene *ftsZ*, the initiator of bacterial cell division, was upregulated nearly 160 and 1600-fold

in colonized WT and *muMT*$^{-/-}$ mice, respectively. At this stage, several additional genes were upregulated

in *muMT*$^{-/-}$ mice, including *ftsK* (~95-fold) which functions in chromosome segregation, *mrdA* and *mrdB*

(40 and 114-fold) which have roles in bacterial cell shape and elongation, and peptidoglycan biogenesis

genes *murCDE* (84 to 206-fold). By day 28, these genes were upregulated in bacteria from both groups of

mice, along with cell division protein *ftsA* (112 and 274-fold). Other key cell division genes (e.g., ftsBLNQ)

were not detected in the rarefied data. Genes active in protein secretion (*secA*, *yjjK*, *prlA*) followed similar

patterns, and the porin *ompA*, which is known to modulate infection and host immunity in gram-negative

bacteria, was upregulated at day 28 (62 and 247-fold in WT and *muMT*$^{-/-}$ mice). A number of genes showed

host genotype-specific expression at day 28, including peptidoglycan biosynthesis genes *murG* and *mraY*

(upregulated in the WT host) and nutrient stress response gene *relA* (upregulated in *muMt*$^{-/-}$ microbiota).

Collectively, these data demonstrate substantial changes in bacterial functions in response to the

establishment of a new protist resident.

Profiling of *T. mu* gene expression during colonization required assembly of a draft metagenome

Concurrent with changes in bacterial gene expression during *T. mu* colonization, we tracked the activities

of the protist itself in the mouse gut. To allow mapping and interpretation of gene expression in the protist,

a draft metagenome assembly of purified *T. mu* was produced using one million PacBio sequences (median

length 11,924 bp), polished with 8.6 million Illumina MiSeq reads. The assembly consisted of 1,780 contigs

with a total length of 225 Mbp. Sequences were AT-rich (71.4%) with repetitive regions across 52% of the

length (see Tables S4 and S5 for metagenome statistics). A total of 26,723 genes were predicted (mean

length 1,878 bp), of which 18,405 were identified as protein coding. We also identified two distinct rRNA

sequences, inclusive of the 18S, 5.8S, 28S and internal transcribed spacers (ITS), with 97% sequence

identity, suggesting the presence of one additional tritrichomonad organism (Figure 4A). Phylogenetic

comparison of the ITS sequences with other parabasalids confirmed that both belong to the

Tritrichomonadidae family (Figure 4B). The major sequence (81% of all reads) was synonymous with *T.*

mu, whereas the minor sequence (19% of reads) belonged to a novel species distinct from all parabasalids

catalogued in the GenBank database, with 85.4% identity at the ITS locus to *T. muris*.

Functional annotation of protist genes identified 1,932 unique Pfam domains, 999 unique Gene Ontology (GO) terms and 471 unique enzymes (EC numbers). Among the top 20 most frequent Pfam domains in *T. mu* is an interesting BspA-type Leucine rich repeat region (PF13306), identified in 317 protist genes and present also in large numbers in *T. vaginalis*, *E. histolytica* and *E. dispar*[27,28]. Seven of these genes were closely related to *T. vaginalis* TvBspA625, a cell surface protein with known antigenic properties (Tables S6 and S7)[27]. In bacteria, these domains were reported in virulence factors that mediate interactions with the host extracellular matrix and promote invasion at mucosal sites[29,30]. We also identified adhesins, lectins and cysteine proteases, other protein families implicated in facilitating host interactions and parasite infection (Tables S6 and S7). Such proteins localize to the cell surface or are secreted, and may represent key factors in *T. mu*'s ability to establish a chronic colonization[27,31-33]. We furthermore report homologues to enzymes of the arginine dihydrolase pathway, a pathway linked to immune escape in *G. duodenalis* through the depletion of local L-arginine, ultimately inhibiting host production of toxic nitrous oxide[34-36]. Enzymes in this pathway (arginine deiminase (ADI), TMU_00005093 and TMU_00022804; ornithine carbamoyltransferase (OCT), TMU_00009104; and carbamate kinase (CK), TMU_00010220 and TMU_00019703) have also been identified in *T. vaginalis* and *T. foetus*[37], and may contribute to chronicity of colonization. Future research is set to validate the role of these enzymes in host interactions and permanent colonization.

T. mu gene expression changes during colonization

The assembly of the *T. mu* metagenome allowed us to map 157 million metatranscriptomic reads to the protist. Of these, 20% mapped uniquely across 17,212 genes (between 247,166 and 3,810,677 reads per sample), and were used to explore *T. mu* activity during colonization of a healthy and B cell-deficient gut. Gene expression changed over time in both hosts (666 DE genes in WT and 627 in *muMt*[-/-] mice, $p < 0.05$), and the presence of host B cells appeared to have a greater impact later in the infection, at day 28 (Figures 4C and 4D). Pathway analysis showed upregulation of *T. mu* carbohydrate metabolism, including galactose

(ec00052) and starch and sucrose (ec00500) during colonization, increased degradation of glycans

including N-glycans and/or gangliosides (ec00511), as well as upregulation of the tricarboxylic acid cycle

(TCA; ec00020) and glycolysis (ec00010), consistent with growth (Figure 4E). The expression of 41 *T. mu*

genes changed in a B cell-dependent manner, including four BspA-like proteins (with predicted

extracellular immunogenic regions), which increase in expression from day 2 to 28 in WT mice (Figure 4F,

Table S8). Interestingly, in *muMT*$^{-/-}$ mice, these genes showed either no change or decreasing expression

over time (TMU_00005602 and TMU_00010916; $p < 0.05$). Other genes with potential cytoadherence and

virulence roles (BspA-like, adhesins, lectins and cysteine proteases) exhibited a diverse array of expression

changes in response to colonization time and/or the presence of B cells (Figures 4G and S4). For example,

homologues of adhesins AP33 and AP120 (TMU_00013231 and TMU_00017025) and TvBspA625

(TMU_00015121), genes with documented roles in *T. vaginalis* infection, were upregulated at day 28, with

intriguing increases already at day 2 in *muMT*$^{-/-}$ mice (Figure 4G; day 28 vs day 2 in WT, $p < 0.05$ and $p <$

0.01; *muMT*$^{-/-}$, *ns*; WT vs *muMT*$^{-/-}$ at day 2, *ns*). Conversely two of three identified cysteine proteases

(TMU_00012588 and TMU_00013765) were downregulated at day 28 (Figure S4B). These data

demonstrate that dynamic metabolic and accessory functions are required by the protist as it adapts to the

host environment.

256 Single cell profiling of *T. mu* reveals distinct life cycle stages

Since trichomonads display a complex life cycle featuring a variety of cell states (e.g. trophozoite,

amoeboid, pseudocyst) we applied single cell RNA sequencing (scRNA-Seq) to explore whether the

dramatic changes in protist gene expression, defined in the previous section, reflects a shift in the

distribution of cell states. Here, we were also interested in seeing how the absence of the bacterial

microbiome might impact the colonization and life cycle propagation of *T. mu*, and therefore performed

scRNA-Seq on protists isolated from both conventionalized and germ-free (GF) mice. GF mice are also

known to possess an immature B cell compartment with low levels of IgA prior to colonization[38].

Sequencing produced 232 and 224 million reads in GF and conventionalized mice, respectively, with 70%

and 76% uniquely mapped to the protist genome assembly (Table S9). Gene expression was analysed in

3,000 *T. mu* cells per host, with medians of 3,776 and 3,140 transcripts and 823 and 589 genes per cell. We

identified 15 distinct protist subpopulations, subsequently assigned to three 'superclusters' (Figure 5A).

Clusters 1 through 4 (designated supercluster 'A') and clusters 10 and 11 were notably enriched in protists

isolated from the GF mouse, while clusters 5 through 7 (within supercluster 'B') and 15 were represented

primarily by protists from the conventionalized mouse (Figure 5B).

Functional enrichment of GO terms in genes upregulated in cell clusters suggested the presence of actively

metabolizing cells, those at various stages of cell cycle, and protists potentially undergoing pseudocyst

formation (Figure 5C). Clusters 3 and 5-7, for example, were enriched in carbohydrate metabolism (7 GO

terms), including galactose and monosaccharide metabolism (GO:0006012; GO:0005996). Cytoskeleton

remodeling and actin polymerization/depolymerization (19 GO terms) were upregulated in clusters 1-3 and

6-8, and biosynthetic processes (14 GO terms), including peptide and macromolecule biosynthesis

(GO:0043043; GO:0009059) were enriched in clusters 1-4 and cluster 8, as well as in clusters 12-14

(supercluster 'C'). We therefore hypothesize that clusters 1-4 ('A') and 5-11 ('B') may represent actively

feeding and growing *T. mu* trophozoites. Sphingolipid metabolism (GO:0006665), which contributes to

regulation of *Giardia* encystation[39], was enriched only in cluster 8. Interestingly oxidative stress response

(GO:0006979) was upregulated in the neighbouring clusters 12 and 13, and the metabolism and

biosynthesis of biogenic amines (GO:0042401, GO:0006576) were upregulated in cluster 14. As biogenic

amines increase in *Entamoeba invadens* during encystation[40], we suggest that the oxygen stress response

may represent stress-induced activation of the encystation transcriptional program in clusters 12 and 13,

and that cluster 14 may represent a pseudocyst form of *T. mu*, previously identified in *T. foetus* and *T.*

vaginalis[41,42]. Cluster 14, incidentally, has the lowest transcriptional complexity, defined by the ratio of

detected genes per transcripts, and the fewest detected genes (Figure 5D).

Enrichment of chromosome and DNA reorganization GO terms in clusters 1, 5 and 11 (eight GO terms), and DNA replication and expression of cell cycle checkpoint genes (10 GO terms) in cluster 10, suggests these protists may be undergoing cell cycle. We note that cluster 15, which shares some of these functions, also features increased DNA recombination (GO:0006310) and may be undergoing sexual or parasexual replication. To lend support to these findings, we carried out a comprehensive analysis of G1/S (*hisH3*, *hisH4*, *tk*) and G2 (*pcna*, *cycA* and *cycB*) cell cycle marker genes, as well as those specific to and/or required for meiosis (Tables S6 and S7)[43,44]. Functional enrichment scores were calculated based on the average expression of each gene set relative to randomly chosen control genes. Scores for expression of G1/S marker genes were highest in Clusters 1 and 11, and cluster 10 scored highest for the G2 phase (G1/S cluster 1 or cluster 11 vs remaining clusters, $p < 0.001$; cluster 1 vs cluster 11, $p < 0.01$; G2 cluster 10 vs remaining clusters, $p < 0.001$; Figures 5E and 5F). We confirmed increased expressions of histones H3 and H4 (highest during G1/S) and thymidine kinase (*tk*; expressed during S phase) in clusters 1 and 11 (Figures S5A and S5B). Genes which achieve highest expression during G2 phase (*pcna* and cyclins *cycA* and *cycB*), and minichromosome maintenance (*mcm*) genes, expressed during G1/S and G2 phases, were upregulated in cluster 10.

To probe for evidence of meiosis, we searched for 26 essential proteins—previously identified in *T. vaginalis*[44] and involved in recombination, and synaptonemal and meiotic sister chromatid cohesion complexes—in the *T. mu* proteome. Homologues to 24 of these proteins were present, including all eight with meiosis-specific activity: Dmc1, Spo11-1, Hop1, Hop2A, Hop2B, Mnd1, Msh4, Msh5 and Mer3 (Table S6). Spo11-1 mapped to two highly similar *T. mu* proteins, however the expression of only one was detected in our experiment. As above, we calculated an enrichment score for meiosis-specific gene expression across all clusters. Cluster 15, representing a small group of 60 cells (52 from the conventionalized and eight from the GF mouse), scored highest ($p < 0.001$; Figures 5E and 5F). In addition

to meiosis-specific genes *dmc1*, *hop2A* and *mnd1*, required genes *rad1*, *mre11*, *smc2*, *smc3* and *smc5* were

also significantly upregulated in cluster 15 ($p < 0.001$; Figure S5C), consistent with protists undergoing

sexual or parasexual replication. *Smc2* and *smc5* were upregulated also in cluster 10, supporting DNA

replication ($p < 0.001$). Given the differences in cluster distributions between conventionalized and GF

mice (Figure 5B), including the proportion of cells undergoing meiosis, our results suggest that the

constituent microbiome plays a role in modulating the protist's life cycle.

Protist subpopulations express distinct sets of virulence-related genes

Since the distribution of protist life cycle stages are altered in the absence of an intact constituent

microbiome, we were interested in how these differences might impact the ability of *T. mu* to colonize. We

therefore examined the expression of genes suspected to facilitate colonization (BspA-like proteins,

adhesins, lectins and cysteine proteases; Figure 5G). All four groups of genes were expressed in *T. mu*

isolated from both GF and conventionalized hosts, and their expression patterns were broadly defined by

the three major superclusters (A, B and C). Of the 317 identified BspA-like genes, we detected 176 in the

scRNA-Seq data. Remarkably, the expression of many of these genes was restricted to specific

subpopulations and, with the exception of clusters 12-14 (putative encysting stages), they differed between

protists from the GF and conventionalized mice (Table S10). A number of genes, for example, was enriched

in clusters 1-4 across both hosts (e.g. TMU_00019122, TMU_00026100), while the exclusive expression

of others in the conventionalized mouse (e.g. TMU_00025647, TMU_00025644) led to separation of these

clusters based on the presence of a constituent microbiome. Gene expression patterns in the two hosts were

similar in each of clusters 12-14, however a distinct shift in cluster 14 may reflect the terminal stage of

encystation. Notably, cells associated with cluster 15 (the putative sexual stage) exhibited distinct profiles

of BspA-like genes between the GF and conventionalized hosts, supporting the idea that the constituent

microbiome impacts multiple stages of the protist life cycle.

Eight of the nine *T. mu* adhesin homologues, which in *T. vaginalis* mediate binding to host epithelial cells

(AP33, AP51, AP65 and AP120)[31], were detected and notably depleted in clusters 4, 13 and 14 (Figure 5G).

The latter is perhaps indicative of the switch to pseudocyst forms that may facilitate transmission. Lectin

homologues, likewise displayed distinct patterns of expression across clusters, with similar depletion in

clusters 12-14, as well as in the putative sexual stage, cluster 15. Interestingly, cells from clusters 7-10

expressed more lectins in the GF mouse relative to their conventionalized counterparts. Similar to the other

groups of putative virulence factors, cysteine proteases exhibited distinct expression profiles across clusters,

with clear differences between the GF and conventionalized mice and limited expression in cluster 14. Of

the four cysteine proteases expressed in putative pseudocysts, two are members of the C54 peptidase family

with similarity to yeast ATG4 which is involved in autophagy. C54 cysteine proteinases, along with other

autophagy related proteins, have in *T. vaginalis* been implicated in cell starvation and differentiation[45,46].

This dynamic arsenal of putative virulence factors appears to have lifecycle stage-specific roles in *T. mu*,

that are further modulated by the presence of a constituent microbiota. Their distinct patterns of expression

further support the transition to a pseudocyst form in clusters 12-14, while clusters 1-4 and 5-11 represent

two distinct, actively growing forms of the protist, perhaps the trophozoite and amoeboid states.

Distinct *T. mu* cell stages are detected *in vivo*

We validated the presence of distinct *T. mu* subpopulations in the intestinal lumen through *in situ*

fluorescent labelling of protozoan mRNAs that were uniquely expressed or distinctively upregulated across

specific clusters (Figure 6A). Microscopic imaging of caecal sections from WT mice revealed protists as

large, nucleated cells in the gut lumen, with no apparent adherence to the host epithelium (Figure 6B).

Transcript TMU_00005724, which was expressed in a majority of cells of supercluster 'B', was labelled in

31% of caecal protists (Figures 6A and 6C). TMU_00016742, expressed in a subset of 'B' (clusters 5, 6,

and 9), was detected in 9% of cells also positive for TMU_00005724, and was uniquely expressed in another

19%. Fluorescent *in situ* hybridization flow cytometry (FiSH-Flow) revealed expression of

TMU_00005724 across all protists freshly isolated from caeca (Figure 6D), with reproducible detection of

cluster-specific transcripts TMU_00016742 (clusters 5, 6 and 9), TMU_00009244 (cluster 12) and

TMU_00001185 (cluster 14) and in 2.6%, 0.08% and 0.19% of cells, respectively (Figure 6E). Greater

proportions of protists expressed TMU_00016742 in *muMT*$^{-/-}$ mice ($p < 0.05$) and trended towards more

TMU_00001185-positive but fewer TMU_00009244-expressing cells (*ns*), suggesting possible B cell

influence on the *T. mu* transcriptional program (Figure 6E). Since TMU_00001185 is uniquely expressed

in cells of cluster 14, those we hypothesize to represent *T. mu* pseudocysts, and pseudocyst formation occurs

during egress from the gastrointestinal tract, we tracked TMU_00001185 expression in protists freshly

isolated from the caecum and colon, and for up to 3 days of *in vitro* culturing (Figure 6F). Caecal isolates

contained the fewest TMU_00001185-expressing cells, consistent with their identity as pseudocysts

undergoing egress through the colon. In line with our speculations, proportions of TMU_00001185-

expressing cells increased during *in vitro* culturing. Since chitin is a critical component of cysts or

pseudocysts in other trichomonads and protozoan parasites, we measured levels of chitin in freshly isolated

or cultured *T. mu* by staining with fluorophore-labelled wheat-germ agglutinin (WGA). WGA fluorescence

charted a pattern similar to that of TMU_00001185: lowest detection in fresh caecal isolates, with a steady

increase during *in vitro* culturing (Figures 6G and 6H)[42,47,48]. Studies are ongoing to characterize the

pseudocyst stage at the ultrastructural level. These *in situ* results validate the presence of transcriptionally

distinct groups of protists, and provide an effective means to track *T. mu* population dynamics and

functional adaptations during colonization.

***T. mu* and gut bacteria compete for nutrients**

We surmise that the establishment of a chronic *T. mu* infection depends on its ability to compete with

resident bacteria, as well as the host, for nutrients in the intestinal tract. Recently we showed a link between

iron supplementation (an essential and limiting micronutrient required for microbial growth), protist

colonization and microbiome structure[5]. Since iron is essential for trichomonad growth, and pathogenesis

in urogenital tract infection[49,50], we hypothesized that its consumption by intestinal *T. mu* would exert

pressure on bacterial iron acquisition and storage systems. Our metatranscriptomic data confirmed

increasing *T. mu* iron consumption through the upregulation of the pyruvate:ferredoxin oxidoreductase

(TMU_00017025, $p < 0.01$) and predicted ferredoxin-1 (TMU_00007489, *ns*), key iron-sulfur cluster

containing proteins required for trichomonad energy production (Figure 7A)[51,52]. The enzymes were

expressed across protist populations, with reductions in cluster 4, and in clusters 13 and 14, which aligns

with reduced energy requirements in the pseudocyst stage (Figure 7B). The upregulation instead of

ferredoxins TMU_00018447 and TMU_00005891 in clusters 4, and particularly 12 and 14 may be linked

to encystation-related stress responses. Similar expression of these genes in *T. mu* from conventionalized

and GF mice may suggest that the protist is agnostic in terms of iron acquisition to the resident microbiome.

In response to the increasing demand for iron from the protist, bacteria showed upregulated iron acquisition

and storage systems (Figures 7C and 7D). Gene expression in *T. mu*-colonized mice was enriched for

siderophore transport genes at day 28, predominantly in *Helicobacter*, *Bacteroides* and *Parabacteroides* (p

< 0.001), and for genes with identified siderophore transport *potential* already at day 2 (WT, $p < 0.05$;

muMt[-/-], $p < 0.001$; Figures 7C and S5C). Numerous siderophore synthesis genes were also upregulated at

days 2 and 28 (Figure 7D), in addition to iron gene regulation (WT, $p < 0.01$; *muMt*[-/-], $p < 0.05$; primarily

Firmicutes), iron storage (WT, *ns*; *muMt*[-/-], $p < 0.05$; *Mucispirillum schaedleri* and unidentified bacteria)

and magnetosome formation (WT, $p < 0.05$; *muMt*[-/-], $p < 0.01$; primarily *Bacteroides*) (Figures 7C and

S5C). Host B cell status appeared to have little impact on bacterial iron-related machinery in the context of

the protist (Figure S6B), which may suggest a more direct competition between resident microbiota and the

new colonizer for an increasingly limited nutrient.

Another nutrient at the interface of gut microbiota and host is the essential amino acid tryptophan.

Tryptophan plays a central role in the formation of bioactive metabolites related to host physiology e.g.

including indole, tryptamine, kynuneurine and serotonin. The gut microbiome has emerged as a potent

driver for modulating its metabolism[53], and we therefore examined how *T. mu* and resident bacteria might

compete for tryptophan and route its conversion. In *T. mu*, a putative tryptophanase (EC 4.1.99.1;

TMU_0006921), which converts tryptophan to indole, was significantly upregulated at day 28 relative to

day 2 (WT, $p < 0.05$; *muMt$^{-/-}$*, $p < 0.01$; Figure 7E). Enzymes associated with alternative pathways, in the

conversion of tryptophan to kynurenate (EC 3.5.1.9; TMU_00003049) or to tryptamine (EC 4.1.1.28;

TMU_00000947), were expressed at relatively low levels and exhibited no significant differences over

time, suggesting that *T. mu* preferentially drives production of indole (Figure 7E). This metabolite, also

previously shown to be an end product of tryptophan metabolism in *T. foetus* and *T. vaginalis*, is known to

impact host immune response, mucosal homeostasis and barrier function, and serves in interspecies

signaling among bacteria[54-56]. Select bacteria, in the presence of *T. mu*, appear to upregulate their own

tryptophanases (Figure 7F). Six of 19 bacterial tryptophanases were differentially expressed in *T. mu*

colonized-mice: the two most highly expressed increased in expression between days 2 and 28, and mapped

to *Bacteroides* and *Prevotella* (CDE07436; colonized vs naïve at day 2 in WT, $p < 0.05$, and *muMT$^{-/-}$*, $p <$

0.01; at day 28 in both hosts, $p < 0.0001$), and to *Oscillibacter* (R9MHE9; colonized vs naïve at day 28 in

both hosts, $p < 0.001$; WT vs *muMt$^{-/-}$* at day 28,; day 28 vs day 2 in WT colonized, $p < 0.01$). Four identified

arylformamidases (EC 3.5.1.9) showed no significant *T. mu*-dependent differences, and two other enzymes

with possible activity in the kynurenine pathway (tryptophan 2,3-dioxygenase, EC 1.13.11.11; kynurenine

aminotransferase, EC 2.6.1.7) were absent after data rarefaction. This may suggest that tryptophan

metabolism is further driven toward the production of indole by gut bacteria in the presence of the protist.

In line with these observations, our single cell data showed upregulation of *T. mu* tryptophanase in the

presence of a constituent microbiota, and reduced expression during transition to the pseudocyst form in

clusters 13 and 14 (Figure 7E), consistent with reduced dependence on host mucosal homeostasis.

The consumption of arginine in the gut, an important source of energy for both trichomonads and bacteria,

also has an immunomodulatory role by limiting host production of antimicrobial NO[36,57]. Enzymes of the

associated arginine dihydrolase pathway—arginine deiminase (ADI; EC 3.5.3.6), ornithine

transcarbamylase (OTC; EC 2.1.3.3) and carbamate kinase (CK; EC 2.7.2.2)—are encoded by both the

protist and bacteria. We explored competition for the nutrient by tracking the expression of these enzymes

among *T. mu* and the resident microbiota. None of the enzymes expressed by *T. mu* were significantly

associated with either colonization time or B cell status, suggesting continual arginine consumption (Figure

S4D). Interestingly, the two isoforms of ADI, the only enzyme specific to this pathway, were selectively

upregulated in protist subpopulations, either clusters 1-4 (TMU_00005093) or clusters 5-11 and 15

(TMU_00022804) (Figure 7G). The same pattern was observed for the two isoforms of CK

(TMU_00010220 associated with clusters 1-3 and TMU_00019703 with clusters 6-11). The single OTC

showed greater expression in clusters 5-15, and consistently so in conventionalized mice (GF vs

conventionalized, clusters 5-14, $p < 0.001$; cluster 15, $p < 0.05$), while the constituent microbiota showed

no significant effects on ADI and CK genes. The presence of *T. mu*, however, may impact arginine

metabolism in select bacteria. Numerous enzymes with potential activity in the arginine dihydrolase

pathway were detected in bacteria (104 ADI, 740 OTC and 396 CK). In the rarefied data used to evaluate

gene expression, a single ADI (belonging to a member of the Lactobacillales order) was detected in *T. mu*-

colonized mice only, with highest expression at day 28 (Figure 7H). Of the two most highly expressed OTC

and CK genes each, one OTC (belonging to *Helicobacter*) increased in expression in both WT and *muMT*-

/- colonized mice, whereas one CK (from an unclassified bacterium) increased in WT colonized mice only

(day 28, $p < 0.05$). These data show complex changes in microbial arginine consumption during protist

colonization.

The dynamics of the acquisition and metabolism machinery for these three key gut nutrients suggests

competitive (iron) and even cooperative relationships (tryptophan) between the protist and the resident

microbiota, and provide a high resolution view of complex tripartite relationships in the gut ecosystem.

Discussion

Gut colonization with *T. mu* and its close relative *T. muris* is known to stimulate the mouse immune

system[9,14]. Such immune changes induced by other parasitic protozoa were shown to mediate antagonistic

and mutualistic interactions with intestinal bacteria, suggesting the intriguing possibility that the resident

microbiome may mediate protist colonization or virulence, and vice versa[7,8,58]. Here, we profiled the

activities of *T. mu* during a new colonization event, and report its ability to reshape both resident bacterial

composition and gene expression in healthy hosts, and more rapidly in those with a compromised immune-

mediated ability to regulate the microbiota. Protist-induced shifts in bacterial composition have previously

been documented for other protozoa, including *G. duodenalis*, *T. foetus* and *T. gondii,* as well as in a recent

study of *T. mu*[8,17,18,59]. Some of the compositional changes that we noted also occurred in the study by Wei

et al., including a reduction of *Lactobacillaceae* and increase in Bacteroidetes and *Parabacteroides*

abundances[17]. However, while the previous study found reduced bacterial diversity and *Helicobacter*

abundance after colonization, we observed the opposite in both healthy and B cell-deficient mice (possibly

due to distinct mouse lines, animal facilities and variations in *Tritrichomonas* strains), and indeed increased

Helicobacter RNA expression in the presence of *T. mu*. Increased abundance of *Helicobacter* has also been

reported after *T. foetus* colonization of cats[18].

Increased bacterial metabolism and peptidoglycan biogenesis, and particularly the shift from Firmicutes to

gram negative taxa (Bacteroidetes and Proteobacteria) at day 28, is a particularly interesting consequence

of colonization, and may signal environmental pressure favouring the growth of particular taxa. As gram

negative bacteria harbor surface LPS capable of boosting immunity, their increasing abundance and activity

may contribute to the intestinal immune activation by *T. mu*[9,60]. The earlier occurrence of these changes (at

day 2) in B cell-deficient mice suggests a failure to control commensal bacteria, protist expansion or both.

Absence of B cells is known to result in aberrant control of gut bacterial communities, and we recently

showed altered IgA reactivity against commensal bacteria in the presence of *T. mu*[22,61,62]. We speculate this

reactivity might serve to limit adverse responses by the resident bacteria as the protist attempts to colonize.

The reactivity of IgA against trichomonad cell surface proteins has also previously been demonstrated in

T. vaginalis, *T. foetus* and *Trichomonas tenax*[63]. Here, we show earlier upregulation of several putative

surface proteins in *T. mu* in the absence of B cells and IgA, including TMU_00015121, a homologue of the

antigenic *T. vaginalis* BspA625 protein previously detected in clinical isolates, and homologues of *T.*

vaginalis AP33 and AP120 adhesins[27]. These may signal changes in protist immunogenicity modulated by

B cells, or a more rapid expansion of TMU_00015121-expressing *T. mu* in their absence. This would be

consistent with the central role of antibodies (including IgA) in parasite clearance. While IgA-deficient

mice are unable to clear *Giardia* infections, for example, there is no evidence that antibodies against *T. mu*

will facilitate protozoan clearance from the intestine[64].

We noted distinct patterns of expression among *T. mu* subpopulations of BspA-like proteins, adhesins,

lectins and cysteine proteases, protein families implicated in host cell binding or parasite virulence[27,31-33].

This may suggest a change in host adhesive properties or antigenicity as the protist adopts different cell

states during its life cycle. Related trichomonads *T. vaginalis* and *T. foetus* are known to exist in one of

three forms—actively growing trophozoites, host-adherent amoeboid cells or environmentally resistant

pseudocysts[41,42,65-67]. The capture of actively metabolizing *T. mu* undergoing cell cycle in the scRNA-Seq

data, and cells with stress-activated pathways neighbouring those expressing relatively fewer genes (with

little to no expression of adhesins) suggests we have captured at least the trophozoite and pseudocyst states.

The upregulation of biogenic amine biosynthesis in the putative pseudocyst is in line with the increased

levels of biogenic amines observed, for example, during *Entamoeba invadens* encystation[40]. Although

adhesins were depleted in these *T. mu* cells, the binding of *T. foetus* cysts to vaginal epithelia was previously

demonstrated through electron microscopy, a possible survival strategy against expulsion[68]. If *T. mu*

pseudocyst forms are capable of cytoadherence, the distinct set of expressed BspA-like genes may

contribute to such a strategy. To gain access to host epithelia, particularly a trophozoite or amoeboid *T. mu*

would need to penetrate the intestinal mucous layer. *T. vaginalis* utilizes cysteine proteases to degrade host

mucins along the vaginal wall[69]. In addition to the large repertoire of cysteine proteases in the *T. mu* genome,

enzymes of the N-glycan degradation pathway, encoded by both *T. mu* and *T. vaginalis*, are also capable of

degrading O-glycans, the dominant glycans found in mucins[70].

Since encystation occurs prior to exit from the intestinal tract, or under conditions of stress, we expected

few pseudocysts in the caecum. In fresh caecal isolates, we indeed detected transcripts of the pseudocyst

marker gene in fewer than 1% of protists. This proportion increased during incubation *in vitro*, accompanied

by a reduction in motile cells, lending support to cluster 14 as pseudocysts. A similar transcript-labelling

approach could be used to capture cells in the amoeboid state. Exposure to host epithelia and cell surface

glycoproteins in other trichomonads triggers a transformation of trophozoites to host-adherent amoeboid

cells[65,66]. Identification of uniquely expressed transcripts in such cells of *T. mu* could help to track the

appearance of amoeboid cells, particularly in contexts of induced colitis and models of colorectal cancer

which are exacerbated by *T. mu* coloinzation[9]. One possibility is that we have already captured the

amoeboid state in our scRNA-Seq data, in supercluster 'A'. The majority of protists in these populations

were recovered from the GF mouse, which are known to have a more penetrable intestinal mucus[71]. These

cells express different suites of adhesins, cysteine proteases and enzymes involved in the degradation of

arginine relative to supercluster 'B', which we suggest represents the trophozoite form. Our attempts to

capture the cells of state 'A' in mice with conventional microbiomes using fluorescent labelling have so far

failed. As microscopic analysis both in our study and that of Wei *et al.* found no evidence of host adherence,

it is possible that the more accessible host epithelium in GF mice promotes transformation to an amoeboid

state[14]. This could have important implications for protist colonizations in a diseased or inflamed gut where

the mucus layer is compromised.

One intriguing finding was the expression of meiosis-specific genes in a small subpopulation of *T. mu* cells,

suggesting the presence of sexual or parasexual replication. Sexual recombination has previously been

hypothesized in *T. vaginalis*, based on the presence and expression of core meiotic genes in its genome (27

of 29)[44,72]. Synapsed chromosomes observed in small subpopulations of cultured *T. vaginalis* and linkage

disequilibrium in its genome, suggesting the non-random segregation of genetic alleles, have also been

presented as evidence of a sexual cycle[72-74]. Meiotic genes have similarly been identified in *G. duodenalis*,

known to undergo parasexual recombination[75]. The expression of meiosis-specific gene *hop2B* in other *T.*

mu clusters suggests this gene may have alternative roles in cellular processes, seen also in the ciliate

Tetrahymenoa thermophila which requires *hop2B* for vegetative growth[76]. While our data suggests that the

presence of a microbiome may promote meiosis (as noted by the higher proportion of meiotic cells in the

conventionalized relative to GF mice), we nonetheless note that despite extensive efforts to purify *T. mu*

from mouse caeca prior to infection of new mice, we detected bacterial DNA in the GF mouse. Unlike the

GF mouse, however, the conventionalized mouse was exposed to commensal bacteria for four weeks prior

to the introduction of *T. mu*. This would have allowed ample time for microbiome-induced changes in

intestinal morphology and immune cell differentiation, which reach a new homeostasis within eight to 30

days[77]. Consequently, the protist would have encountered two very different intestinal environments.

The upregulation of bacterial iron acquisition machinery, consequent to *T. mu* colonization, could be a

competitive response to nutrient depletion by *T. mu*, and may function as a complementary means to

maintain control over protist growth. Iron is an essential micronutrient. To limit the growth and virulence

of invading pathogens, the host restricts access to luminal iron through sequestration by proteins such as

lactoferrin[49,50,78]. Parasitic trichomonads require particularly high quantities to sustain their growth. In *T.*

vaginalis, enzymes of the hydrogenosome, an energy-producing organelle resembling mitochondria,

achieve peak activity at over 200 μM iron, whereas at 100 μM, growth is restricted and the protist

upregulates iron sequestration[50]. Low iron conditions have also been shown to hinder trichomonad adhesion

to host epithelia[79]. Iron uptake by resident bacteria has the potential, therefore, to further restrict protist

growth and virulence, an effect which may be lost in a context of surplus iron provided for example by

nutritional supplements. Previous studies have associated iron supplements with increased growth of

enteropathogens, diarrhea and intestinal inflammation in infants[5]. In a cohort of young children from

Pakistan supplemented with iron and vitamins, we recently showed increased carriage of numerous

intestinal protozoa and fungi, with possible carry-over effects to bacterial composition[5]. In the present

study, we noted blooms of siderophore transport in *Helicobacter* and iron storage in *Mucispirillum*

schaedleri, a pathobiont causally associated with Crohn's-like colitis in immunodeficient mice[80].

Interestingly, changes in iron levels have also been shown to alter expression of *T. vaginalis* BspA-like

genes *in vitro*, potentially influencing protist virulence[27]. The question of how iron supplementation or

restriction influences the balance and/or virulence of gut protozoa and bacteria remains an important

question. In addition to iron, we also noted a potential for competition over the catabolism of tryptophan,

with the expression of a putative tryptophanase by the protist. Recently there has been much interest in the

production of neuroactive kynurenines from tryptophan by the gut microbiome, and their potential role in

the gut brain access[81,82]. Through shifting the production of indole at the expense of alternative products,

tryptamine and kynurenine, our findings suggest that colonization by the protist may similarly modulate

the gut-brain access.

We note also several limitations in the metatranscriptomic study. Although we used littermate controls to

compare the microbiota of protist-colonized and naïve mice, host genotype-dependent differences may have

been confounded by caging effects. Differences in bacterial composition and gene expression diminished

over time in *T. mu* colonized mice, possibly a due in part to transfer of bacteria adherent to the protist. In

the microbial RNA-Seq data, we were limited in our ability to predict bacterial gene function and taxonomy

by the current available reference microbial data. We note that the large number of unclassified RNA

sequences may belong to taxa identified by 16S rRNA data but for which genome related information is

sparse, or to non-coding regulatory RNAs not examined in this study. Metatranscriptomics also revealed

the unexpected presence of murine astrovirus in mouse caeca, represented by 5.5 million reads and

assembled into full-length genomes belonging to at least two viral strains. As this virus is capable of altering

the host intestinal barrier, its presence represents a potential confounding factor in delineating protist-

specific effects on the microbiome[83]. It nevertheless showcases the potential of this method to identify

additional agents contributing to inflammatory disease. We note that we found no association between virus

detection, protozoan colonization and host genotype in this study.

Through a series of sequencing studies, we showed that introduction of a new protist leads to significant

remodeling of resident microbiota in the gut. As these changes are accelerated in the absence of B cells, we

suggest that antibodies moderate the expansion of the protist during the early days of colonization. We

furthermore demonstrated powerful means to track protist activities and cell states through combined

sequence-based and *in situ* methodologies, which will contribute substantially to our ability to decode the

strategies employed by gut parasites during colonization.

Author contributions

J.P., A.M., M.E.G., and A.P. conceived and designed the study. A.P. and A.K. isolated nucleic material for

high throughput sequencing, and A.P. performed sequence data analyses. E.Y.C. performed mouse

colonization experiments, performed flow cytometry and microscopic analyses. N.N. performed protozoan

enzyme annotation. N.A. and J.H. participated in sample collection and *in vitro* culturing experiments.

E.C.A.F. generated the phylogenetic tree. A.P., J.P., A.M. and M.E.G. wrote the manuscript. All authors

reviewed and/or edited the paper.

Acknowledgements

This work was funded by the Canadian Institutes for Health Research (MRT-168043) to J.P., M.E.G. and A.M; the Natural Sciences and Engineering Research Council (RGPIN-2019-06852) to J.P.; the Intramural Research Program of the National Institute of Allergy and Infectious Diseases at the National Institutes of Health to M.E.G.; a Restracomp scholarship administered by the Research Training Centre (Hospital for Sick Children) and a graduate scholarship from the Government of Ontario to A.P. A.M. is supported by the Canadian Foundation for Innovation John R. Evans Leaders Fund, the Canadian Institutes for Health Research (PJT-388337, PJT-480765) and a Natural Sciences and Engineering Research Council (RGPIN-2019-04521). A.M. is the Tier 2 Canadian Research Chair in Mucosal Immunology and supported by the Tier 2 CRC-CIHR program (CRC-2021-00511). Computing resources were provided by the SciNet High Performance Computing (HPC) Consortium; SciNet is funded by the Canada Foundation for Innovation under the auspices of Compute Canada, the Government of Ontario, Ontario Research Fund - Research Excellence, and the University of Toronto.

Declaration of interests

The authors declare no competing interests.

     Main figure titles and legends

Figure 1. Study design to test the impact of _T. mu_ colonization on resident microbiota.

(A) Schematic of microbial interactions explored in the present study using 16S sequencing,

metatranscriptomics (RNA-Seq) and/or single-cell transcriptomics (scRNA-Seq). (B) Experimental design.

Groups of four WT and _muMT_$^{-/-}$ mice were colonized with _T. mu_. Fecal pellets and caecal samples were

collected at indicated timepoints for 16S and metatranscriptomic profiling. (C) Expansion of _T. mu_ cells in

WT mouse caeca and (D) mucosal IgA levels over 28 days of colonization.

Figure 2. _T. mu_ colonization drives bacterial diversification.

(A) Relative abundance of bacterial composition as determined by 16S amplicon sequencing of mouse fecal

pellets. Plots represent profiles from four WT and _muMt_$^{-/-}$ mice sampled over a four-week colonization time

course, as well as two groups of uninfected control mice sampled at the end of the experiment. Colours

represent genus-level abundances within indicated phylum colour groups. (B) Bacterial richness (observed

OTUs) and evenness (Shannon diversity) over the course of _T. mu_ colonization. (C) Principal coordinate

analysis showing compositional differences among samples, as determined Bray-Curtis dissimilarities. (D)

Patterns of bacterial changes during _T. mu_ colonization in WT (grey) and _muMt_$^{-/-}$ (red) mice. Significantly

differentially abundant OTUs were clustered into groups based on patterns of abundance changes. Pie charts

show taxonomies of OTUs included in each cluster. (E) Phylum-level abundance determined through qPCR

using clade-specific primers, at before (day 0) or 21 days after colonization with _T. mu_ in WT _(top)_ and

muMt$^{-/-}$ mice _(bottom)_. Abundances were normalized to Ct values obtained with universal 16S rRNA

probes. Error bars represent SEM. Mann-Whitney U test for significance was performed, and significance

is indicated as follows: *$p < 0.05$, ** < 0.01, _ns_ not significant.

Figure 3. Bacterial gene expression is altered in the presence of the protist.

(A) Taxonomic profiles of putative bacterial mRNA reads. (B) Principal component analysis of bacterial gene expression. Each circle denotes a sample isolated from a single mouse, coloured by host genotype, presence of *T. mu* and timepoint, as shown. (C) Numbers of significantly upregulated or downregulated bacterial genes between protist-colonized and naïve mice, or due to host genotype (*muMt*[-/-] vs WT), grouped as indicated (n = 3 or 4). (D) Bacterial metabolic pathway enrichment associated with *T. mu* colonization. Enrichment was determined by the overrepresentation of EC terms in differentially expressed genes of KEGG-defined pathways. Total ECs detected in the experiment and in the rarefied gene expression matrix used in the analysis are indicated in the bar charts to the right. Significantly enriched pathways are represented as coloured dots: grey indicates *ns*, and the gradient from yellow to red represents decreasing *p* values, beginning from *p*<0.05. Sizes of dots represent odds ratios. Significance was calculated using Fisher's exact tests. Asterisks indicate infinite odds ratios. (E) Interaction network of cell division, cell wall biogenesis and secretory genes, based on a previously determined network of *E. coli* proteins[26]. Coloured nodes represent protein homologues identified in the metatranscriptomic data, and edges are previously described protein-protein or functional interactions. Genes significantly up- and downregulated are denoted with blue and red borders, respectively. Node sizes correlate with log2-fold changes between *T. mu*-colonized and naïve mice. Colours within node pie charts represent proportions of gene expression assigned to various taxa, as shown: yellows represent Proteobacteria, greens are Bacteroidetes, pinks and purples are Firmicutes, and black denotes unclassified bacteria. Colours encircling groups of proteins represent functional modules as previously described[26].

Figure 4. Protist gene expression changes during colonization of WT and *muMt*[-/-] mice.

(A) Graphical depiction of two trichomonad rRNA sequences identified in the metagenome assembly. The major sequence is identical to *T. mu* and maps to 81% of reads, and the minor sequence maps to 19% of

reads. The sequences were aligned using MUSCLE. Mismatches in the minor sequence to *T. mu* are indicated in black. (B) Phylogeny of the two distinct ribosomal ITS sequences recovered from the trichomonad metagenome. An unrooted tree was constructed from aligned ITS regions using the Maximum-likelihood method. The major sequence refers to *T. mu*, showing 99.4% identity to the reference, and the minor sequence shares 85.4% identity with *T. muris*. Scale = number of SNPs per site. (C) Principal component analysis of protist gene expression patterns over time, in WT or *muMt$^{-/-}$* mouse caeca. (D) Numbers of differentially expressed genes associated with colonization time (day 28 vs day 2) or host genotype (*muMT$^{-/-}$* vs WT). (E) Metabolic pathway enrichment in *T. mu* during colonization. The dotplot represents enrichment of EC terms in KEGG-defined pathways, within the set of genes differentially expressed over time or based on B cell status. Coloured dots represent those significantly enriched: the gradient from yellow to red marks decreasing *p* values, beginning from *p*<0.05. Grey dots are *ns*. Significance was evaluated using Fisher's exact tests, and dot sizes correlate with odds ratios. Total ECs detected in the experiment are shown in the bar charts to the right. (F) Normalized expression of BspA-like genes showing interaction with host B cell status and (G) differentially expressed homologous to genes implicated in *T. vaginalis* virulence. Differences were evaluated using the Wilcoxon rank-sum test, and adjusted for multiple testing using the Benjamini-Hochberg approach. *$p < 0.05$, ** < 0.01, *** < 0.001, *ns*, non-significant.

Figure 5. Single cell transcriptional profiling of caecal *T. mu* populations.

(A) Protist subpopulations isolated from colonized GF and conventionalized mouse caeca. Each point represents one of 6000 protists cells, clustered by gene expression profiles using UMAP dimensionality reduction, and coloured by cluster designation (3000 cells from each of two hosts). Groups of clusters, designated 'superclusters' A, B or C, are indicated. (B) Numbers of cells assigned to each cluster by host are shown. Total cluster sizes are shown on the right. (C) Heatmap of GO term enrichment within *T. mu* clusters. Rows represent significantly enriched GO terms in one or more clusters (columns). Cells are

coloured and clustered by (log-transformed) numbers of upregulated proteins per GO term. (D)

Transcriptional complexity of protist clusters, defined as numbers of genes per transcript. Boxplots indicate

medians and interquartile ranges. (E) Violin plots and (F) feature plots depicting G1/S, G2 and meiotic

scores in cells across each *T. mu* cluster. Scores were determined based on the average expression of pre-

defined gene sets relative to randomly chosen control genes. Significance was tested using the two-sided

Wilcoxon-rank sum test, $*p<0.05$, $**<0.01$, $***<0.001$. (G) Heatmap showing expression of (*top* to *bottom*)

BspA-like proteins, putative adhesin proteins, lectins and Cysteine proteases across protist clusters. Colours

to the right of Cysteine proteases indicate the protein family (Pfam).

Figure 6. *T. mu* subpopulations are detected *in situ*.

(A) Feature plots showing the expression of select cluster-specific genes. (B) Immunofluorescence

micrograph of mouse cecal tissue after colonization with *T. mu* for 21 days. Nuclear staining (blue) shows

host and protozoan DNA, and white dotted lines denote the mucus layer. (scale bar = 10μm). (C) RNAscope

immunofluorescence of *T. mu* transcripts TMU_00005724 (green) and TMU_00016742 (red) on sections

of cecal tissue. Nuclear DNA is stained blue. Panels on either side (scale bars = 10μm), and centre images

(scale bars = 1μm). Frequencies of protists expressing cluster-specific genes are shown on the right. (D)

FISH-Flow analysis of freshly isolated *T. mu* at 21 days post colonization. Histogram shows control

unstained cells (grey) and cells stained with fluorescent probes for TMU_00005724 (purple) (E) FISH-

Flow analysis of *T. mu* hybridized with fluorescent RNA probes for the TMU_00005724 transcript,

multiplexed with probes for TMU_00016742, TMU_00009244 or TMU_00001185 transcripts. Contour

plots indicate fluorescence within isolated protists from a representative experiment from WT mice.

Percentages of cells positive for hybridized probes are indicated within gates. Adjacent bar plots show

frequencies of *T. mu* positive for transcripts 28 days post colonization in WT (grey) and *muMt*$^{-/-}$ (red) mice

(n=4 per group). Significance was tested using the Wilcoxon rank sum test. (F) Percentages of protists

freshly isolated from WT mouse caeca and colons, or cultured *in vitro* for one and three days, which were

positive for the TMU_00001185 transcript or (G) stained with WGA-FITC. Representative contour plots

of WGA-stained protists are shown (n=3 animals or culture plates per group). Significance was tested using

one-way ANOVA, and adjusted for multiple comparisons using Tukey's test. (H) Cytospins of WGA-

stained protists. Representative images are shown at 63x magnification. Error bars represent SEM. *$p <$

0.05, **$p < 0.01$, *ns* not significant.

Figure 7. *T. mu* and resident bacteria compete for resources in the gut lumen

Shown are changes in expression of bacterial and *T. mu* genes related to iron, A-D, generation of tryptophan

catabolites, E and F, and the arginine dihydrolase pathway, G and H. (A) Expression of putative *T. mu*

pyruvate:ferredoxin oxidoreductase (PFOR) and ferredoxin (Fdx) genes in metatranscriptomic data, at days

2 and 28 of the experiment. (B) PFOR and Fdx gene expression among protist subpopulations isolated from

GF and conventionalized mice. The colour intensity correlates with average expression across cells and the

sizes of the dots indicate the percentage of cells in each group with detected expression. (C) Enrichment of

iron-related gene expression in bacteria, in the context of *T. mu* colonization. Shown are numbers of DE

genes per iron-related gene family in *T. mu*-colonized versus naïve mice at days 2 and 28. Total numbers

of genes identified in each family in the metatranscriptomic data are indicated in brackets. Significance of

enrichment was tested using the hypergeometric test. (D) Median log2 fold-change of bacterial siderophore

transport and synthesis gene expression between colonized and naïve mice. Significantly DE genes are

shown as coloured points: the gradient from yellow to red marks decreasing *p* values, beginning from

p<0.05. (E) Expression of *T. mu* and (F) the most highly expressed bacterial (read count >100) enzymes

with predicted function in the production of tryptophan metabolites. Dotplots in E show gene expression

among protist subpopulations, as above. Pie charts in F represent the total numbers of tested and DE

bacterial enzymes. Blue and green dots in the graphic show enzyme activities identified in *T. mu* and/or

bacterial genes, respectively. (G) Expression of *T. mu* enzymes associated with the arginine dihydrolase

pathway among protist subpopulations isolated from GF conventionalized mice, as above. (H) Expression

of bacterial enzymes associated with the arginine dihydrolase pathway. Shown are the only ADI present in

the rarefied gene count matrix, and the two differentially expressed OTC and CK genes. Gene expression

in metatranscriptomics data was evaluated using DESeq2, and in scRNA-Seq data using Seurat. All

expression values are derived from normalized gene counts. Boxplots and violin plots show medians and

interquartile ranges. *P* values were corrected for multiple testing using the Benjamini-Hochberg approach.

*$p<0.05$, **<0.01, ***<0.001

 STAR Methods

Mice

C57Bl/6J and B6.129S2-*Ighm*[tm1Cgn]/J (*muMt*[-/-]) female mice were purchased from Jackson Laboratory (Bar

Harbor, ME, USA) and maintained in specific pathogen free (SPF) conditions at the University of Toronto

Division of Comparative Medicine animal facility. Animals were housed in a closed caging system and

provided with an irradiated chow diet (Envigo Teklad 2918), non-acidified water (reverse-osmosis and UV-

sterilized) with a 12hr light/dark cycle. Experiments were conducted at 6-8W of age, using age and sex

matched controls. Following colonization with *T. mu*, animals were housed in separated cages to avoid

contamination of non-colonized mice. Animal experiments were approved by the Local Animal Care

Committee (LACC) at the Faculty of Medicine, University of Toronto (animal use protocol 20012400 to

AM).

For the single-cell RNA sequencing experiment, 12-week-old germ-free (GF), female C57Bl/6 mice were

purchased from Taconic Biosciences (Germantown, NY, USA) and maintained in the GF facility until the

day of analysis, when they were transported to specific pathogen free (SPF) conditions. Those

conventionalized with a microbiome received a 200 μL bacterial suspension through oral gavage, derived

from two C56Bl/6J fecal pellets resuspended in 1ml sterile PBS, followed by centrifugation at 1000rpm for

3min to remove large debris. Conventionalized mice were then maintained under SPF conditions for 4

weeks prior to *T. mu* colonization.

***T. mu* colonization**

T. mu used in this study was originally isolated at the Icahn School of Medicine at the Mount Sinai animal

facility in New York[9]. Purification of *T. mu* from mice was performed as previously described[9]. The cecal

content of a colonized C57Bl/6J mouse was harvested into 20ml sterile cold PBS and filtered through a

70µm cell strainer. Cecal contents were centrifuged at 500g for 7min at 4°C. The supernatant was discarded,

and the pellet was washed twice with 40ml sterile PBS, and resuspended in 5ml 40% Percoll overlaid on

5ml of 80% Percoll. Samples were centrifuged at 1300g for 20min at 20°C with the breaks off. The

interphase containing the protist was collected, washed with sterile PBS, pelleted at 4°C, and resuspended

in 5ml sterile PBS and filtered as above. Protist cells were sorted into sterile PBS on a BD Influx Cell Sorter

(BD Biosciences, Franklin Lakes, NJ, USA) using the 100µm nozzle at 27psi at 4°C, with >99% purity.

Sorted cells were once more pelleted as above and resuspended in sterile PBS. Two million collected *T. mu*

cells were orally gavaged immediately after into each C57Bl/6J wildtype (WT) or *muMt*[-/-] mouse, or for the

scRNA-Seq experiment into 16-week-old female GF or previously conventionalized C57Bl/6 mice.

Trichomonad genome sequencing and annotation

High molecular weight genomic DNA was extracted from 100 million sorted tritrichomonad cells using the

MagAttract HMW DNA Kit (QIAGEN, Hilden, Germany). Library generation and sequencing using

PacBio Sequel technology on two SMRT cells were carried out at the McMaster University Farncombe

Metagenomics Facility (Hamilton, Canada). Reads were error corrected using CANU v1.8[84], and

subsequently assembled into contigs using Flye v.2.4.1[85]. Contig sequences were polished through

alignment with eight million 300 bp paired-end Illumina MiSeq reads (sequenced at the National Institute

of Allergy and Infectious Diseases, NIH, Bethesda, Maryland) using BWA[86] and Pilon v.1.23[87]. The

genome assembly is available at: https://github.com/ParkinsonLab/Tritrichomonas-murine-microbiome-

interactions/draft-genome-assembly

Genome annotation was carried out using the Maker v2.31 pipeline[88] as follows. Repetitive regions were

identified *de novo* and masked using RepeatModeler v.1.0.11[89] and RepeatMasker v.4.0.7[90]. Transfer RNA

genes were predicted with tRNAscan-SE v2.0.6[91]. *Ab initio* prediction of gene models was carried out using

SNAP version 2013-11-29[92] and Augustus version 3.3.1[93], with an initial two rounds of training using

SNAP followed by one using Augustus with BUSCO v.2.0.1[94]. Reference information for gene prediction

included data from related trichomonad organisms: the *T. foetus* expressed sequence tag (EST) library

(CX159216.1), *T. foetus* strain K (3AUP000179807) and *Trichomonas vaginalis* G3 (3AUP000001542)

proteomes, and available *Pentatrichomonas hominis* and *Dientamoeba fragilis* protein sequences within

the NCBI nr database (accessed August 21, 2019). Functional annotation including Gene Ontology (GO)

term and PFAM prediction was performed with InterProScan v.5.30-69.0[95] and the hmmscan algorithm,

from HmmerWeb version 2.41.2[96], with minimum 1e-05 E-value cut-offs. Enzyme Commission (EC)

numbers were assigned to proteins using the enzyme prediction program ARCHITECT[97] with minimum

0.5 confidence scores. Genes encoding adhesins, meiosis and cell cycle-related proteins were identified

based on protein-level homology with relevant *T. vaginalis* proteins[31,44] as follows. Sequences of *T.*

vaginalis proteins were retrieved from the TrichDB database[98], and submitted for sequence similarity

searches against the *T. mu* proteome using BLAST[99], with minimal scoring criteria of E-value 1e-5, 30%

sequence identity and 40% query coverage. Where proteins matched multiple candidates in *T. mu* with a

10% or greater difference in sequence identity, only the highest scoring match was retained.

For functional comparison of *T. mu* genes with other protozoa and a model organism, GO terms, PFAM

domains and EC numbers were downloaded for *T. vaginalis* G3 (release 2.0), *Giardia intestinalis*

assemblage A isolate WB (release 1.0) and *Entamoeba dispar* SAW760 (release 49) from VEuPathDB[100],

and the *Tritrichomonas foetus* (UP000179807), *Blastocystis* sp. ST1 (UP000078348) and *Saccharomyces*

cerevisiae (UP000002311) proteomes in UniProt[101] (accessed Oct 27, 2020). GO terms were summarized

into GO Slim categories using the R package GSEABase[102].

Protist phylogeny analysis

Ribosomal internal transcribed spacer (ITS) sequences were extracted from PacBio whole genome

sequence data. Parabasalid sequences downloaded from the GenBank database

(https://www.ncbi.nlm.nih.gov/genbank/) and used in the analysis of phylogeny are listed in Table S11.

Multiple sequence alignments were generated using the MUSCLE algorithm[103], and an unrooted phylogeny

was constructed using the Maximum-likelihood method with a Tamura-Nei model[104] in the MEGA-X

software[105]. Bootstrap values were generated using 1000 replicates, with only values >65% shown.

scRNA sequencing and analysis of gut-colonizing protist

One GF and one conventionalized C56Bl/6J female mouse (Taconic Biosciences, Germantown, NY, USA)

were colonized with *T. mu* through oral gavage, as described above. Four weeks post colonization, the GF

mouse was transferred to the SPF facility, where the conventionalized mouse was being housed, and protist

cells were isolated from the caecal contents as described above. Purified *T. mu* was immediately transferred

on ice to the Princess Margaret Genomics Centre (Toronto, Canada) for STAMP library preparation using

the DropSeq approach and sequencing on the NextSeq 500 platform (Illumina, San Diego, CA, USA)[106].

Reads were processed using Drop-seq Tools v.1.13[107] as per the authors' instructions and aligned to the *T.*

mu genome assembly using STAR v. 2.5.3a[108]. A total of 3,000 cells, with minimum 200 detected genes

and 500 transcripts, were selected from each mouse for gene expression analysis in Seurat v4[109,110]. Only

genes detected in a minimum of 3 cells were included. Anchor cells were identified using

FindVariableFeatures and FindIntegrationAnchors functions based on 2000 most variable genes, and used

to align the transcriptome datasets. Cells were grouped into clusters based on gene expression using graph-

based clustering with the following parameters: resolution=0.8, principal component dimensions=18;

clusters were visualized using UMAP nonlinear dimensionality reduction[111].

Differences in metabolic pathway regulation among protist cell clusters were identified as follows.

Differentially expressed genes were identified for each cluster using the FindAllMarkers function in Seurat,

with minimum *p*-value 0.05, adjusted for multiple testing using Bonferroni correction. All enzymes in the

T. mu gene set were identified by their assigned EC numbers. Differential regulation of metabolic pathways

(the set of enzymes associated with each pathway, as defined by the Kyoto Encyclopedia of Genes and

Genomes, KEGG[112]) between cell clusters was determined using the hypergeometric test, to test for

overrepresentation of pathway enzymes within the differentially expressed gene set. Only pathways having

a minimum of 5 and maximum of 500 enzymes in the *T. mu* genome assembly were considered. Functional

enrichment within differentially expressed genes was also performed using GO term overrepresentation

with the topGO package[113] and the Fisher's Exact test. The AddModuleScore function in Seurat was used

to score enrichment of custom-defined gene sets (e.g. meiosis, G1/S and G2 phase genes), and score

differences between clusters were tested for significance using the two-sided Wilcoxon-rank sum test.

Benjamini-Hochberg correction was applied for multiple testing[114]. Heatmaps of gene expression were

generated using the pheatmap 1.0.12[115] package in R and clustered using the Ward.D2 hierarchical

clustering algorithm.

16S microbiome profiling and metatranscriptomics

Groups of four WT or *muMt*[-/-] six-week-old female mice were infected with two million protists each by

oral gavage for profiling of 16S rRNA and microbial gene expression. A schematic of the experiment is

presented in Figure 1a. Protist burdens were determined in mouse caeca using a hemocytometer.

16S microbiome profiling. For 16S profiling, mouse fecal pellets were collected from individual mice and

stored at -80°C prior to *T. mu* colonization and at days 2, 7, 14, 21 and 28 post-colonization. Pellets from

two groups of uninfected control mice were collected at day 28 for comparison. Microbial DNA was

extracted using the DNeasy PowerSoil Kit (QIAGEN, Hilden, Germany) according to manufacturer

instructions, with an initial 5min bead beating step using the FastPrep-24 homogenizer (MP Biomedicals,

Santa Ana, CA, USA) at 5M/s. Generation of 16S rRNA amplicons and sequencing were carried out by

the Integrated Microbiome Resource facility at Dalhousie University (Halifax, Canada). Briefly, 16S

variable regions 6 to 8 (V6-V8) were amplified using B969F and BA1406R primers[116] and Phusion DNA

polymerase (New England Biolabs, Ipswich, MA, USA). The first amplification round was carried out

using primers fused to Illumina adapters and multiplexing barcodes, followed by 30 cycles as follows: 98°C

for 10s, 55°C for 30s, 72°C for 30s, and a final extension at 72°C for 5min. Libraries were generated and

250 bp paired-end sequencing was carried out with MiSeq v2 chemistry (Illumina, San Diego, CA, USA).

Data were quality filtered, denoised and clustered *de novo* to 97% operational taxonomic units (OTUs)

using Dada2[117] within QIIME2[118] (parameters: truncate f 280, r 240, trim 10 (f/r), min-parent-fold 5). OTU

taxonomies were assigned using QIIME2 with a custom classifier trained on V6-V8 regions extracted from

SILVA[119] v132 nr SSU sequences and clustered to 97% sequence identity.

Bacterial diversity and differential abundance. Diversity and multivariate analyses were performed with

R-packages Phyloseq 1.26.1[120] and vegan v.2.5[121]. Microbiota richness (number of OTUs) and evenness

(Shannon Diversity Index, H) values were averaged from 100 independently rarefied datasets to the

minimum read depth (19,427). To evaluate the effects of *T. mu* colonization time and host genotype on

bacterial diversity, we implemented linear mixed effects (LME) models using the package lme4[122], and

performed a two-way ANOVA on day 28 samples to compare diversities of infected and naïve mice. Beta

diversity analyses, using Bray Curtis and weighted Unifrac (wUnifrac) dissimilarity scores, were carried

out on rarefied data, using OTUs present at a minimum of 10 reads in 10% of samples, and significance

was tested using the adonis function in vegan. Differential taxon abundance was evaluated with DESeq2

v.1.22.2[123], and groups of taxa with similar changes in abundance were identified using DEGreport[124].

Microbial metatranscriptomics. Caecal contents were recovered from sacrificed *T. mu*-colonized and

uninfected mice at days 2 and 28 of the experiment and immediately frozen at -80°C. On the same day, 30-

90mg caecal samples were added to 1ml ice cold TRIzol (Invitrogen, Carlsbad, CA, USA) containing

200mg 0.1mm Zirconia beads (BioSpec Products, Bartlesville, OK, USA). Samples were homogenized

with three 2min cycles of bead-beating at 25 Hz using the TissueLyser (QIAGEN, Hilden, Germany), and

chilled on ice for 5min between cycles. Homogenized samples were incubated on ice for an additional

15min, followed by 5min at room temperature. Total RNA extraction was achieved using the

PureLink RNA Mini Kit (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's TRIzol Plus

Total Transcriptome Isolation protocol. Prokaryotic and eukaryotic rRNA was depleted using the QIAseq

FastSelect rRNA HMR and 5S/16S/23S kits (QIAGEN, Hilden, Germany). Sequencing libraries were

prepared using NEBNext Ultra II Directional RNA Library Prep Kits for Illumina (New England Biolabs,

Ipswich, MA, USA) and paired-end sequenced (150 bp) on the NovaSeq 6000 platform (Illumina, San

Diego, CA, USA) at Genome Quebec (Montréal, Canada). Low quality reads, host and vector

contamination, rRNA and tRNA reads were filtered out using the MetaPro pipeline[125]. Remaining reads

were mapped to the *T. mu* genome assembly using STAR v. 2.5.3a[108], and all unmapped reads were returned

to MetaPro for contig assembly, gene model prediction and functional and taxonomic annotation. Proteins

with iron-related functions were identified using FeGenie[126] searches of the annotated gene and protein

sequences from the metatranscriptome as well as the *T. mu* proteome. Putative cell wall biogenesis

machinery was identified by conducting sequence similarity searches of translated genes and proteins in

the metatranscriptomics data against previously described proteins[26] in the *E. coli* MG1655 proteome using

DIAMOND v0.9.22[127] with a minimum E-value 1e-10.

Differential gene expression and functional enrichment. Differential gene expression analyses were

performed using DESeq2 v.1.22.2[123]. Analysis for *T. mu* was performed on genes prefiltered for a minimum

detection of two reads in two samples. Due to differences in read depth of bacterial genes between *T. mu*-

colonized and naïve samples, gene count matrices were rarefied to 600,000 reads per sample. Differential

gene expression was calculated for 15 independent rarefactions, and median log2-fold changes and average

false discovery rate-corrected *p* values were recorded. Principal component analyses of *T. mu* and bacterial

gene expression were performed on DESeq2 variance stabilized counts. Functional enrichment was

evaluated using EC or GO term overrepresentation in differentially expressed genes, as described for single

cell data above, where enrichment of up- or downregulated enzymes in any given pathway was used as a

proxy for pathway activity.

RNA-FISH Flow Cytometry

Purified *T. mu* were divided into staining aliquots of 4 million cells and samples were washed twice with

PBS by spinning at 500g for 7min at 4°C. Cells were stained with Zombie Aqua viability dye (BioLegend,

San Diego, CA, USA) in PBS in the dark (1:1000 – 15min – 4°C), washed twice with PBS, and the

remaining steps were performed as described in the Stellaris RNA-FISH protocol for cells in suspension.

After hybridization, cells were resuspended in Wash Buffer B and analyzed on a BD LSR Fortessa X-20

cell analyzer (BD Biosciences, Franklin Lakes, NJ, USA). For detection of chitin on putative pseudocysts,

T. mu was isolated from the mouse caecum and colon at day 21 post colonization and stained with wheat

germ agglutinin (WGA)-FITC (Sigma Aldrich, St. Louis, MO, USA) in FACS buffer (PBS – 2% FBS –

5mn EDTA) in the dark. To monitor encystation of protists upon isolation from the gastrointestinal tract,

isolated protists were either stained immediately or cultured anaerobically *in vitro* for one to three days

prior to staining. Cytospins were visualized using a Zeiss AXIO Observer microscope (Carl Zeiss AG, Jena,

Germany).

RNAScope

RNAScope was performed as indicated in the RNAScope Multiplex Fluorescent Reagent Kit v2 (Advanced

Cell Diagnostics, Newark, CA, USA) protocol. Approximately 0.5cm long cecum sections were excised

from *T. mu*-colonized C57Bl/6J WT or *muMt*[-/-] mice, placed in 10% neutral buffered formalin and fixed

overnight at RT with gentle agitation. The next day, samples were gently washed twice with PBS, placed

in 70% ethanol, and embedded in paraffin and sliced to 7μm sections at the Toronto Centre for

Phenogenomics. Paraffin sections were pretreated as follows. Samples were baked at 40°C for 30min in a

HybEZ Oven (Advanced Cell Diagnostics, Newark, CA, USA), and hydrogen peroxide was applied at RT

for 10min. Antigen target retrieval was conducted at 99°C in a water bath under the 15min standard antigen

retrieval procedure. A barrier was created around sections using an ImmEdge pen (Vector Laboratories,

Burlingame, CA, USA) and allowed to dry at RT for 15min. Samples were treated with protease at 40°C

for 30min in a HybEZ Oven, and stored in sterile saline sodium citrate solution overnight (175.3g NaCl –

88.2g sodium citrate – 800ml ddH2O – pH 7). Multiplex probe amplification was conducted as per the

manufacturer's instructions: TSA Plus Fluorophores Fluorescein and Cyanine 3 were hybridized against

protist probes TMU_00005724 and TMU_00016742 respectively, and samples were visualized using a

Zeiss AXIO Observer microscope (Carl Zeiss AG, Jena, Germany).

qPCR

DNA Isolation for quantitative PCR. DNA isolation was performed from frozen fecal pellets using the

QIAQuick qPCR purification kit (QIAGEN Hilden, Germany) with modifications. Fecal weights were

measured, and pellets were thawed on ice for 10min. SDS buffer (8g SDS – 30mL H_2O – 0.22μm) and DNA

buffer A (20mM Tris pH 8 – 2mM EDTA – 200mM NaCl) were combined in a 1:1.41 ratio to prepare SDS-

A buffer. Samples were transferred to cryogenic 2ml tubes and 400μl zirconium beads, 550μl of phenol-

chloroform, and 482μl of SDS-A buffer were mixed with each sample. Samples were homogenized using

the Mini-BeadBeater (Biospec Products, Bartlesville, OK, USA) at 4000rpm for 5min, centrifuged at 1800g

for 5min, and the supernatant was transferred into a PCR purification plate containing 650μl of PM buffer.

Samples were washed twice with PE buffer, centrifuged and eluted into 100μl H_2O. DNA concentrations

were quantified using the Quant-iT kit (Thermo Fisher Scientific, Waltham, MA, USA).

RNA isolation and reverse transcription of mRNA for quantitative real time RT PCR. Two million *T. mu*

cells were combined with 500μL TRIzol (Invitrogen, Waltham, MA, USA) and incubated at RT for 5min.

100μL chloroform was added to each sample, samples were mixed and incubated at RT for 3min and

centrifuged for 15min at 12,000g at 4°C. The aqueous phase containing RNA was mixed with 250μL

isopropanol, mixed and incubated at 4°C for 10min. Samples were centrifuged for 10min at 12,000g at 4°C,

the supernatant was discarded and 500μL of 75% ethanol was added. Samples were vortexed for 10s and

centrifuged for 5min at 7,500g at 4°C. The supernatant was discarded and samples were allowed to air dry

for 10min at RT. RNA was resuspended in 50μL of RNase-free water and stored at -80°C. *T. mu* mRNA

was converted to cDNA using the High-Capacity RNA-to-cDNA kit (Applied Biosystems, Waltham, MA,

USA). In brief, 2μg purified RNA per reaction was mixed with 2X Reverse Transcription Buffer Mix, 20X

Reverse Transcription Enzyme Mix and nuclease free H_2O (up to 20μL). The thermal cycling conditions

were conducted as follows: 37°C for 60min – 95°C for 5min – infinite hold at 4°C.

Quantitative Real-Time RT-PCR. qPCR was performed following the PowerTrack SYBR Green Master

Mix kit instructions (Applied Biosystems, Waltham, MA, USA). In brief, 10ng of DNA was mixed with

2X PowerTrack Master Mix, and 400nM of forward/reverse primers in 10μL reactions. The thermal cycling

conditions were as follows: enzyme inactivation at 95°C for 2 min, 40 cycles of denaturation at 95°C for

15s and annealing/extension at 60°C for 60s. Relative expression of target genes was normalized to the *T.*

mu glyceraldehyde 3-phosphate dehydrogenase (GAPDH) or universal 16S gene for *T. mu* and bacterial

genes respectively, and calculated using the delta-delta Ct method. Primer sequences are included in Table

S12.

1008

IgA ELISA

Between 100μl and 200μl of mouse blood was collected and serum was isolated using Microvette CV 300

Z serum separation tubes (SARSTEDT, Newton, NC. USA) according to the manufacturer's protocol, and

stored at -80°C. Immunoglobulins were absorbed onto high-bound Nunc MaxiSorp ELISA plates

(BioLegend, San Diego, CA, USA) and anti-IgA ELISA was performed using anti-mouse IgA-HRP

antibody (SouthernBiotech, Birmingham, AL, USA). In brief, 50μl of anti-mouse Ig master mix

(SouthernBiotech, Birmingham, AL, USA) was diluted 1:2000 in PBS, and used to coat 96-well ELISA

plates overnight at 4°C. The following day, 200μl of 1% bovine serum albumin (Sigma-Aldrich, St. Louis,

MI, USA) in PBS was added to plates to block for 1hr at 37°C. 50μl of purified IgA (1:1000 in assay buffer)

or serum (1:100 in assay buffer) was added to 100μl assay buffer (0.1M Tris-HCl, 0.1M NaCl, 1% FBS,

0.05% Tween-20), and 6x 1 in 3 serial dilutions were prepared. Mixtures were incubated for 1 hour at 37°C

and washed 3 times with 200μl 0.1% PBST. After adding 50μl of anti-mouse IgA-HRP (1:1000 in assay

buffer) (SouthernBiotech, Birmingham, AL, USA), mixtures were incubated for 1 hour more at 37°C and

washes repeated as before. 50μl of TMB substrate (BioLegend, San Diego, CA, USA) was added, incubated

for 30s, and reactions were stopped with the addition of 50μl of H_2SO_4. Concentrations were measured by

spectrophotometry (450 nm spectral absorbance) and values were calibrated against those of the purified

IgA standard curve.

1026

1027 **Data availability**

1028 Sequence data generated in this study have been deposited to the NCBI Sequence Read Archive under the

1029 BioProject identifiers PRJNA913581 and PRJNA914770.

1030

1031   **Supplemental item titles**

1032

Table S1. 16S sequencing read and OTU counts

Table S2. 16S OTU table

Table S3. Metatranscriptomic read processing and classification

Table S4. Metagenome characteristics.

Table S5. Frequency of predicted transfer RNA genes in the metagenome assembly.

Table S6. Putative *T. mu* homologues of potential virulence and meiosis-specific genes.

Table S7. *T. mu* HMMER protein domain predictions

Table S8. Genes differentially expressed in *T. mu* during WT or B-cell deficient mouse colonization

Table S9. Single cell RNA sequencing statistics of *T. mu*

Table S10. Read counts for genes containing BspA-like domains across *T. mu* subpopulations

Table S11. Parabasalid ribosomal ITS sequences used in the analysis of phylogeny

Table S12. DNA primers used for quantitative RT-PCR

1045

References

Fan, Y. & Pedersen, O. Gut microbiota in human metabolic health and disease. *Nat Rev Microbiol* **19**, 55-71, doi:10.1038/s41579-020-0433-9 (2021).

Libertucci, J. & Young, V. B. The role of the microbiota in infectious diseases. *Nat Microbiol* **4**, 35-45, doi:10.1038/s41564-018-0278-4 (2019).

Sardinha-Silva, A., Alves-Ferreira, E. V. C. & Grigg, M. E. Intestinal immune responses to commensal and pathogenic protozoa. *Front Immunol* **13**, 963723, doi:10.3389/fimmu.2022.963723 (2022).

Scanlan, P. D. *et al.* The microbial eukaryote Blastocystis is a prevalent and diverse member of the healthy human gut microbiota. *FEMS microbiology ecology* **90**, 326-330, doi:10.1111/1574-6941.12396 (2014).

Popovic, A. *et al.* Micronutrient supplements can promote disruptive protozoan and fungal communities in the developing infant gut. *Nat Commun* **12**, 6729, doi:10.1038/s41467-021-27010-3 (2021).

Benson, A., Pifer, R., Behrendt, C. L., Hooper, L. V. & Yarovinsky, F. Gut commensal bacteria direct a protective immune response against Toxoplasma gondii. *Cell Host Microbe* **6**, 187-196, doi:10.1016/j.chom.2009.06.005 (2009).

Chappell, C. L. *et al.* Fecal Indole as a Biomarker of Susceptibility to Cryptosporidium Infection. *Infect Immun* **84**, 2299-2306, doi:10.1128/IAI.00336-16 (2016).

Raetz, M. *et al.* Parasite-induced TH1 cells and intestinal dysbiosis cooperate in IFN-gamma-dependent elimination of Paneth cells. *Nat Immunol* **14**, 136-142, doi:10.1038/ni.2508 (2013).

Chudnovskiy, A. *et al.* Host-Protozoan Interactions Protect from Mucosal Infections through Activation of the Inflammasome. *Cell* **167**, 444-456 e414, doi:10.1016/j.cell.2016.08.076 (2016).

Jokelainen, P. *et al.* Dientamoeba fragilis, a Commensal in Children in Danish Day Care Centers. *J Clin Microbiol* **55**, 1707-1713, doi:10.1128/JCM.00037-17 (2017).

Cuffari, C., Oligny, L. & Seidman, E. G. Dientamoeba fragilis masquerading as allergic colitis. *J Pediatr Gastroenterol Nutr* **26**, 16-20, doi:10.1097/00005176-199801000-00003 (1998).

Yakoob, J. *et al.* Blastocystis hominis and Dientamoeba fragilis in patients fulfilling irritable bowel syndrome criteria. *Parasitol Res* **107**, 679-684, doi:10.1007/s00436-010-1918-7 (2010).

Zhang, N. *et al.* High prevalence of Pentatrichomonas hominis infection in gastrointestinal cancer patients. *Parasit Vectors* **12**, 423, doi:10.1186/s13071-019-3684-4 (2019).

Escalante, N. K. *et al.* The common mouse protozoa Tritrichomonas muris alters mucosal T cell homeostasis and colitis susceptibility. *The Journal of experimental medicine* **213**, 2841-2850, doi:10.1084/jem.20161776 (2016).

Tuzlak, L. *et al.* Ultrastructural identification and molecular characterization of two new parabasalid species that naturally colonize laboratory mice, Tritrichomonas musculus and Tritrichomonas casperi. *BioRxiv*, doi:https://doi.org/10.1101/2023.01.20.524969 (2023).

Chiaranunt, P. *et al.* NLRP1B and NLRP3 Control the Host Response following Colonization with the Commensal Protist Tritrichomonas musculus. *J Immunol* **208**, 1782-1789, doi:10.4049/jimmunol.2100802 (2022).

Wei, Y. *et al.* Commensal Bacteria Impact a Protozoan's Integration into the Murine Gut Microbiota in a Dietary Nutrient-Dependent Manner. *Appl Environ Microbiol* **86**, doi:10.1128/AEM.00303-20 (2020).

Bierlein, M., Hedgespeth, B. A., Azcarate-Peril, M. A., Stauffer, S. H. & Gookin, J. L. Dysbiosis of fecal microbiota in cats with naturally occurring and experimentally induced Tritrichomonas foetus infection. *PLoS One* **16**, e0246957, doi:10.1371/journal.pone.0246957 (2021).

Schneider, C. *et al.* A Metabolite-Triggered Tuft Cell-ILC2 Circuit Drives Small Intestinal Remodeling. *Cell* **174**, 271-284 e214, doi:10.1016/j.cell.2018.05.014 (2018).

20   Howitt, M. R. *et al.* Tuft cells, taste-chemosensory cells, orchestrate parasite type 2 immunity in
the gut. *Science* **351**, 1329-1333, doi:10.1126/science.aaf1648 (2016).
21   Ost, K. S. *et al.* Adaptive immunity induces mutualism between commensal eukaryotes. *Nature*
596, 114-118, doi:10.1038/s41586-021-03722-w (2021).
22   Pabst, O., Cerovic, V. & Hornef, M. Secretory IgA in the Coordination of Establishment and
Maintenance of the Microbiota. *Trends Immunol* **37**, 287-296, doi:10.1016/j.it.2016.03.002
(2016).
23   Rojas, O. L. *et al.* Recirculating Intestinal IgA-Producing Cells Regulate Neuroinflammation via
IL-10. *Cell* **176**, 610-624 e618, doi:10.1016/j.cell.2018.11.035 (2019).
24   Polk, D. B. & Peek, R. M., Jr. Helicobacter pylori: gastric cancer and beyond. *Nat Rev Cancer*
10, 403-414, doi:10.1038/nrc2857 (2010).
25   Partridge, J. D., Scott, C., Tang, Y., Poole, R. K. & Green, J. Escherichia coli transcriptome
dynamics during the transition from anaerobic to aerobic conditions. *J Biol Chem* **281**, 27806-
27815, doi:10.1074/jbc.M603450200 (2006).
26   Peregrin-Alvarez, J. M., Xiong, X., Su, C. & Parkinson, J. The Modular Organization of Protein
Interactions in Escherichia coli. *PLoS Comput Biol* **5**, e1000523,
doi:10.1371/journal.pcbi.1000523 (2009).
27   Noel, C. J. *et al.* Trichomonas vaginalis vast BspA-like gene family: evidence for functional
diversity from structural organisation and transcriptomics. *BMC Genomics* **11**, 99,
doi:10.1186/1471-2164-11-99 (2010).
28   Wilson, I. W. *et al.* Genetic Diversity and Gene Family Expansions in Members of the Genus
Entamoeba. *Genome Biol Evol* **11**, 688-705, doi:10.1093/gbe/evz009 (2019).
29   Inagaki, S., Onishi, S., Kuramitsu, H. K. & Sharma, A. Porphyromonas gingivalis vesicles
enhance attachment, and the leucine-rich repeat BspA protein is required for invasion of epithelial
cells by "Tannerella forsythia". *Infect Immun* **74**, 5023-5028, doi:10.1128/IAI.00062-06 (2006).
30   Rego, S. *et al.* Structural and Functional Analysis of Cell Wall-anchored Polypeptide Adhesin
BspA in Streptococcus agalactiae. *J Biol Chem* **291**, 15985-16000, doi:10.1074/jbc.M116.726562
(2016).
31   Garcia, A. F. & Alderete, J. Characterization of the Trichomonas vaginalis surface-associated
AP65 and binding domain interacting with trichomonads and host cells. *BMC Microbiol* **7**, 116,
doi:10.1186/1471-2180-7-116 (2007).
32   Guzman-Tellez, P. *et al.* Lectins as virulence factors in Entamoeba histolytica and free-living
amoebae. *Future Microbiol* **15**, 919-936, doi:10.2217/fmb-2019-0275 (2020).
33   Siqueira-Neto, J. L. *et al.* Cysteine proteases in protozoan parasites. *PLoS Negl Trop Dis* **12**,
e0006512, doi:10.1371/journal.pntd.0006512 (2018).
34   Das, P., Lahiri, A., Lahiri, A. & Chakravortty, D. Modulation of the arginase pathway in the
context of microbial pathogenesis: a metabolic enzyme moonlighting as an immune modulator.
PLoS Pathog **6**, e1000899, doi:10.1371/journal.ppat.1000899 (2010).
35   Rath, M., Muller, I., Kropf, P., Closs, E. I. & Munder, M. Metabolism via Arginase or Nitric
Oxide Synthase: Two Competing Arginine Pathways in Macrophages. *Front Immunol* **5**, 532,
doi:10.3389/fimmu.2014.00532 (2014).
36   Stadelmann, B., Merino, M. C., Persson, L. & Svard, S. G. Arginine consumption by the
intestinal parasite Giardia intestinalis reduces proliferation of intestinal epithelial cells. *PLoS One*
7, e45325, doi:10.1371/journal.pone.0045325 (2012).
37   Yarlett, N., Lindmark, D. G., Goldberg, B., Moharrami, M. A. & Bacchi, C. J. Subcellular
localization of the enzymes of the arginine dihydrolase pathway in Trichomonas vaginalis and
Tritrichomonas foetus. *J Eukaryot Microbiol* **41**, 554-559, doi:10.1111/j.1550-
7408.1994.tb01516.x (1994).
38   Hapfelmeier, S. *et al.* Reversible microbial colonization of germ-free mice reveals the dynamics
of IgA immune responses. *Science* **328**, 1705-1709, doi:10.1126/science.1188454 (2010).

39   Hernandez, Y. *et al.* Novel role of sphingolipid synthesis genes in regulating giardial encystation.
Infect Immun **76**, 2939-2949, doi:10.1128/IAI.00116-08 (2008).
40   Jeelani, G. *et al.* Metabolic profiling of the protozoan parasite Entamoeba invadens revealed
activation of unpredicted pathway during encystation. *PLoS One* **7**, e37740,
doi:10.1371/journal.pone.0037740 (2012).
41   Pereira-Neves, A. & Benchimol, M. Tritrichomonas foetus: budding from multinucleated
pseudocysts. *Protist* **160**, 536-551, doi:10.1016/j.protis.2009.05.001 (2009).
42   Beri, D. *et al.* Demonstration and Characterization of Cyst-Like Structures in the Life Cycle of
Trichomonas vaginalis. *Front Cell Infect Microbiol* **9**, 430, doi:10.3389/fcimb.2019.00430
(2019).
43   Horlock-Roberts, K. *et al.* Drug-Free Approach To Study the Unusual Cell Cycle of Giardia
intestinalis. *mSphere* **2**, doi:10.1128/mSphere.00384-16 (2017).
44   Malik, S. B., Pightling, A. W., Stefaniak, L. M., Schurko, A. M. & Logsdon, J. M., Jr. An
expanded inventory of conserved meiotic genes provides evidence for sex in Trichomonas
vaginalis. *PLoS One* **3**, e2879, doi:10.1371/journal.pone.0002879 (2007).
45   Hirt, R. P. *et al.* Trichomonas vaginalis pathobiology new insights from the genome sequence.
Adv Parasitol **77**, 87-140, doi:10.1016/B978-0-12-391429-3.00006-X (2011).
46   Brennand, A. *et al.* Autophagy in parasitic protists: unique features and drug targets. *Mol
Biochem Parasitol* **177**, 83-99, doi:10.1016/j.molbiopara.2011.02.003 (2011).
47   Arroyo-Begovich, A., Carabez-Trejo, A. & Ruiz-Herrera, J. Identification of the structural
component in the cyst wall of Entamoeba invadens. *J Parasitol* **66**, 735-741 (1980).
48   Ward, H. D., Alroy, J., Lev, B. I., Keusch, G. T. & Pereira, M. E. Identification of chitin as a
structural component of Giardia cysts. *Infect Immun* **49**, 629-634, doi:10.1128/iai.49.3.629-
634.1985 (1985).
49   Castro, C. *et al.* Iron-modulated pseudocyst formation in Tritrichomonas foetus. *Parasitology*
143, 1034-1042, doi:10.1017/S0031182016000573 (2016).
50   Lehker, M. W. & Alderete, J. F. Iron regulates growth of Trichomonas vaginalis and the
expression of immunogenic trichomonad proteins. *Mol Microbiol* **6**, 123-132, doi:10.1111/j.1365-
2958.1992.tb00844.x (1992).
51   Vanacova, S. *et al.* Iron-induced changes in pyruvate metabolism of Tritrichomonas foetus and
involvement of iron in expression of hydrogenosomal proteins. *Microbiology (Reading)* **147**, 53-
62, doi:10.1099/00221287-147-1-53 (2001).
52   Horvathova, L. *et al.* Transcriptomic identification of iron-regulated and iron-independent gene
copies within the heavily duplicated Trichomonas vaginalis genome. *Genome Biol Evol* **4**, 1017-
1029, doi:10.1093/gbe/evs078 (2012).
53   Agus, A., Planchais, J. & Sokol, H. Gut Microbiota Regulation of Tryptophan Metabolism in
Health and Disease. *Cell Host Microbe* **23**, 716-724, doi:10.1016/j.chom.2018.05.003 (2018).
54   Lloyd, D., Lauritsen, F. R. & Degn, H. The parasitic flagellates Trichomonas vaginalis and
Tritrichomonas foetus produce indole and dimethyl disulphide: direct characterization by
membrane inlet tandem mass spectrometry. *J Gen Microbiol* **137**, 1743-1747,
doi:10.1099/00221287-137-7-1743 (1991).
55   Li, X., Zhang, B., Hu, Y. & Zhao, Y. New Insights Into Gut-Bacteria-Derived Indole and Its
Derivatives in Intestinal and Liver Diseases. *Front Pharmacol* **12**, 769501,
doi:10.3389/fphar.2021.769501 (2021).
56   Zhang, L. S. & Davies, S. S. Microbial metabolism of dietary components to bioactive
metabolites: opportunities for new therapeutic interventions. *Genome Med* **8**, 46,
doi:10.1186/s13073-016-0296-x (2016).
57   Novak, L. *et al.* Arginine deiminase pathway enzymes: evolutionary history in metamonads and
other eukaryotes. *BMC Evol Biol* **16**, 197, doi:10.1186/s12862-016-0771-4 (2016).
58   Yilmaz, B. *et al.* Gut microbiota elicits a protective immune response against malaria
transmission. *Cell* **159**, 1277-1289, doi:10.1016/j.cell.2014.10.053 (2014).

59 Barash, N. R., Maloney, J. G., Singer, S. M. & Dawson, S. C. Giardia Alters Commensal
Microbial Diversity throughout the Murine Gut. *Infect Immun* **85**, doi:10.1128/iai.00948-16
(2017).
60 Wang, J., Chen, W. D. & Wang, Y. D. The Relationship Between Gut Microbiota and
Inflammatory Diseases: The Role of Macrophages. *Front Microbiol* **11**, 1065,
doi:10.3389/fmicb.2020.01065 (2020).
61 Cao, E. *et al.* The protozoan commensal Tritrichomonas musculis is a natural adjuvant for
mucosal IgA. *BioRxiv*, doi:https://doi.org/10.1101/2022.10.08.511442 (2022).
62 Catanzaro, J. R. *et al.* IgA-deficient humans exhibit gut microbiota dysbiosis despite secretion of
compensatory IgM. *Sci Rep* **9**, 13574, doi:10.1038/s41598-019-49923-2 (2019).
63 Mundodi, V., Kucknoor, A. S., Chang, T. H. & Alderete, J. F. A novel surface protein of
Trichomonas vaginalis is regulated independently by low iron and contact with vaginal epithelial
cells. *BMC Microbiol* **6**, 6, doi:10.1186/1471-2180-6-6 (2006).
64 Langford, T. D. *et al.* Central importance of immunoglobulin A in host defense against Giardia
spp. *Infect Immun* **70**, 11-18, doi:10.1128/IAI.70.1.11-18.2002 (2002).
65 Kusdian, G., Woehle, C., Martin, W. F. & Gould, S. B. The actin-based machinery of
Trichomonas vaginalis mediates flagellate-amoeboid transition and migration across host tissue.
Cell Microbiol **15**, 1707-1721, doi:10.1111/cmi.12144 (2013).
66 Petropolis, D. B., Fernandes Rodrigues, J. C., da Rocha-Azevedo, B. & Silva-Filho, F. C. E. The
binding of Tritrichomonas foetus to immobilized laminin-1 and its role in the cytotoxicity exerted
by the parasite. *Microbiology (Reading)* **154**, 2283-2290, doi:10.1099/mic.0.2007/015941-0
(2008).
67 Dias-Lopes, G. *et al.* Morphologic study of the effect of iron on pseudocyst formation in
Trichomonas vaginalis and its interaction with human epithelial cells. *Mem Inst Oswaldo Cruz*
112, 664-673, doi:10.1590/0074-02760170032 (2017).
68 Mariante, R. M., Lopes, L. C. & Benchimol, M. Tritrichomonas foetus pseudocysts adhere to
vaginal epithelial cells in a contact-dependent manner. *Parasitol Res* **92**, 303-312,
doi:10.1007/s00436-003-1026-z (2004).
69 Lehker, M. W. & Sweeney, D. Trichomonad invasion of the mucous layer requires adhesins,
mucinases, and motility. *Sex Transm Infect* **75**, 231-238, doi:10.1136/sti.75.4.231 (1999).
70 Alsmark, C. *et al.* Patterns of prokaryotic lateral gene transfers affecting parasitic microbial
eukaryotes. *Genome Biol* **14**, R19, doi:10.1186/gb-2013-14-2-r19 (2013).
71 Johansson, M. E. *et al.* Normalization of Host Intestinal Mucus Layers Requires Long-Term
Microbial Colonization. *Cell Host Microbe* **18**, 582-592, doi:10.1016/j.chom.2015.10.007 (2015).
72 Bradic, M. *et al.* Genetic Indicators of Drug Resistance in the Highly Repetitive Genome of
Trichomonas vaginalis. *Genome Biol Evol* **9**, 1658-1672, doi:10.1093/gbe/evx110 (2017).
73 Drmota, T. & Kral, J. Karyotype of Trichomonas vaginalis. *European Journal of Protistology* **33**,
131-135 (1997).
74 Bradic, M. & Carlton, J. M. Does the common sexually transmitted parasite Trichomonas
vaginalis have sex? *PLoS Pathog* **14**, e1006831, doi:10.1371/journal.ppat.1006831 (2018).
75 Ramesh, M. A., Malik, S. B. & Logsdon, J. M., Jr. A phylogenomic inventory of meiotic genes;
evidence for sex in Giardia and an early eukaryotic origin of meiosis. *Current biology : CB* **15**,
185-191, doi:10.1016/j.cub.2005.01.003 (2005).
76 Mochizuki, K., Novatchkova, M. & Loidl, J. DNA double-strand breaks, but not crossovers, are
required for the reorganization of meiotic nuclei in Tetrahymena. *J Cell Sci* **121**, 2148-2158,
doi:10.1242/jcs.031799 (2008).
77 El Aidy, S. *et al.* Temporal and spatial interplay of microbiota and intestinal mucosa drive
establishment of immune homeostasis in conventionalized mice. *Mucosal Immunol* **5**, 567-579,
doi:10.1038/mi.2012.32 (2012).
78 Gastelum-Martinez, A. *et al.* Iron-modulated virulence factors of Entamoeba histolytica. *Future*
Microbiol **13**, 1329-1341, doi:10.2217/fmb-2018-0066 (2018).

79   Alderete, J. F., Nguyen, J., Mundodi, V. & Lehker, M. W. Heme-iron increases levels of AP65-
1248   mediated adherence by Trichomonas vaginalis. *Microb Pathog* **36**, 263-271,
1249   doi:10.1016/j.micpath.2003.12.007 (2004).

1250   80   Herp, S., Durai Raj, A. C., Salvado Silva, M., Woelfel, S. & Stecher, B. The human symbiont
1251   Mucispirillum schaedleri: causality in health and disease. *Med Microbiol Immunol* **210**, 173-179,
1252   doi:10.1007/s00430-021-00702-9 (2021).

1253   81   Chen, L. M. *et al.* Tryptophan-kynurenine metabolism: a link between the gut and brain for
1254   depression in inflammatory bowel disease. *J Neuroinflammation* **18**, 135, doi:10.1186/s12974-
1255   021-02175-2 (2021).

1256   82   Ortega, M. A. *et al.* Microbiota-gut-brain axis mechanisms in the complex network of bipolar
1257   disorders: potential clinical implications and translational opportunities. *Mol Psychiatry*,
1258   doi:10.1038/s41380-023-01964-w (2023).

1259   83   Cortez, V. *et al.* Astrovirus infects actively secreting goblet cells and alters the gut mucus barrier.
1260   *Nat Commun* **11**, 2097, doi:10.1038/s41467-020-15999-y (2020).

1261   84   Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and
1262   repeat separation. *Genome Res* **27**, 722-736, doi:10.1101/gr.215087.116 (2017).

1263   85   Kolmogorov, M., Yuan, J., Lin, Y. & Pevzner, P. A. Assembly of long, error-prone reads using
1264   repeat graphs. *Nat Biotechnol* **37**, 540-546, doi:10.1038/s41587-019-0072-8 (2019).

1265   86   Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
1266   *arXiv:1303.3997v1 [q-bio.GN]* (2013).

1267   87   Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant detection and
1268   genome assembly improvement. *PLoS One* **9**, e112963, doi:10.1371/journal.pone.0112963
1269   (2014).

1270   88   Cantarel, B. L. *et al.* MAKER: an easy-to-use annotation pipeline designed for emerging model
1271   organism genomes. *Genome Res* **18**, 188-196, doi:10.1101/gr.6743907 (2008).

1272   89   Smit, A. & Hubley, R. *RepeatModeler Open-1.0* (2008-2015).

1273   90   Smit, A., Hubley, R. & Green, P. *RepeatMasker Open-4.0.*, <http://www.repeatmasker.org>
1274   (2013-2015).

1275   91   Chan, P. P. & Lowe, T. M. tRNAscan-SE: Searching for tRNA Genes in Genomic Sequences.
1276   *Methods Mol Biol* **1962**, 1-14, doi:10.1007/978-1-4939-9173-0_1 (2019).

1277   92   Korf, I. Gene finding in novel genomes. *BMC Bioinformatics* **5**, 59, doi:10.1186/1471-2105-5-59
1278   (2004).

1279   93   Stanke, M., Schoffmann, O., Morgenstern, B. & Waack, S. Gene prediction in eukaryotes with a
1280   generalized hidden Markov model that uses hints from external sources. *BMC Bioinformatics* **7**,
1281   62, doi:10.1186/1471-2105-7-62 (2006).

1282   94   Simao, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M. BUSCO:
1283   assessing genome assembly and annotation completeness with single-copy orthologs.
1284   *Bioinformatics* **31**, 3210-3212, doi:10.1093/bioinformatics/btv351 (2015).

1285   95   Jones, P. *et al.* InterProScan 5: genome-scale protein function classification. *Bioinformatics* **30**,
1286   1236-1240, doi:10.1093/bioinformatics/btu031 (2014).

1287   96   Finn, R. D., Clements, J. & Eddy, S. R. HMMER web server: interactive sequence similarity
1288   searching. *Nucleic Acids Res* **39**, W29-37, doi:10.1093/nar/gkr367 (2011).

1289   97   Nursimulu, N., Moses, A. M. & Parkinson, J. Architect: A tool for aiding the reconstruction of
1290   high-quality metabolic models through improved enzyme annotation. *PLoS Comput Biol* **18**,
1291   e1010452, doi:10.1371/journal.pcbi.1010452 (2022).

1292   98   Aurrecoechea, C. *et al.* GiardiaDB and TrichDB: integrated genomic resources for the eukaryotic
1293   protist pathogens Giardia lamblia and Trichomonas vaginalis. *Nucleic Acids Res* **37**, D526-530,
1294   doi:10.1093/nar/gkn631 (2009).

1295   99   Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search
1296   tool. *J Mol Biol* **215**, 403-410, doi:10.1016/S0022-2836(05)80360-2 (1990).

1297 100 Amos, B. *et al.* VEuPathDB: the eukaryotic pathogen, vector and host bioinformatics resource
1298   center. *Nucleic Acids Res*, doi:10.1093/nar/gkab929 (2021).

1299 101 UniProt, C. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res* **47**, D506-D515,
1300   doi:10.1093/nar/gky1049 (2019).

1301 102 Morgan, M., Falcon S & R, G. *GSEABase: Gene set enrichment data structures and methods. R*
1302   *package version 1.52.*, <https://rdrr.io/bioc/GSEABase/man/GSEABase-package.html> (2020).

1303 103 Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high throughput.
1304   *Nucleic Acids Res* **32**, 1792-1797, doi:10.1093/nar/gkh340 (2004).

1305 104 Tamura, K. & Nei, M. Estimation of the number of nucleotide substitutions in the control region
1306   of mitochondrial DNA in humans and chimpanzees. *Mol Biol Evol* **10**, 512-526,
1307   doi:10.1093/oxfordjournals.molbev.a040023 (1993).

1308 105 Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular Evolutionary
1309   Genetics Analysis across Computing Platforms. *Mol Biol Evol* **35**, 1547-1549,
1310   doi:10.1093/molbev/msy096 (2018).

1311 106 Macosko, E. Z. *et al.* Highly Parallel Genome-wide Expression Profiling of Individual Cells
1312   Using Nanoliter Droplets. *Cell* **161**, 1202-1214, doi:10.1016/j.cell.2015.05.002 (2015).

1313 107 Nemesh, J. & Wysoker, A. *Drop-seq tools v1.13*, <https://github.com/broadinstitute/Drop-
1314   seq/releases> (

1315 108 Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21,
1316   doi:10.1093/bioinformatics/bts635 (2013).

1317 109 Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell
1318   transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol* **36**,
1319   411-420, doi:10.1038/nbt.4096 (2018).

1320 110 Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587 e3529,
1321   doi:10.1016/j.cell.2021.04.048 (2021).

1322 111 McInnes, L., Healy, J., Saul, N. & Großberger, L. UMAP: Uniform Manifold Approximation and
1323   Projection for Dimension Reduction. *Journal of Open Source Software* **3**, 861,
1324   doi:https://doi.org/10.21105/joss.00861 (2018).

1325 112 Okuda, S. *et al.* KEGG Atlas mapping for global analysis of metabolic pathways. *Nucleic Acids*
1326   *Res* **36**, W423-426, doi:10.1093/nar/gkn282 (2008).

1327 113 Alexa, A. & Rahnenfuhrer, J. *topGO: Enrichment Analysis for Gene Ontology. R package version*
1328   *2.42.0.*, <https://github.com/lyijin/topGO_pipeline> (2020).

1329 114 Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful
1330   Approach to Multiple Testing. *Journal of the Royal statistical society: series B* **57**, 289-300,
1331   doi:https://doi.org/10.1111/j.2517-6161.1995.tb02031.x (1995).

1332 115 Kolde, R. *pheatmap: Pretty Heatmaps. R package version 1.0.12.*, <https://CRAN.R-
1333   project.org/package=pheatmap> (2019).

1334 116 Comeau, A. M., Li, W. K., Tremblay, J. E., Carmack, E. C. & Lovejoy, C. Arctic Ocean
1335   microbial community structure before and after the 2007 record sea ice minimum. *PLoS One* **6**,
1336   e27492, doi:10.1371/journal.pone.0027492 (2011).

1337 117 Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon data.
1338   *Nat Methods* **13**, 581-583, doi:10.1038/nmeth.3869 (2016).

1339 118 Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data science using
1340   QIIME 2. *Nat Biotechnol* **37**, 852-857, doi:10.1038/s41587-019-0209-9 (2019).

1341 119 Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data processing and
1342   web-based tools. *Nucleic Acids Res* **41**, D590-596, doi:10.1093/nar/gks1219 (2013).

1343 120 McMurdie, P. J. & Holmes, S. phyloseq: an R package for reproducible interactive analysis and
1344   graphics of microbiome census data. *PLoS One* **8**, e61217, doi:10.1371/journal.pone.0061217
1345   (2013).

1346 121 Oksanen, J. B., F. Guillaume; Friendly, Michael; Kindt, Roeland; Legendre, Pierre; McGlinn,
1347   Dan; Minchin, Peter R. ; O'Hara, R. B.; Simpson, Gavin L.; Solymos, Peter; Henry M.; Stevens,

H.; Szoecs, Eduard; Wagner, Helene. *vegan: Community Ecology Package*, <https://CRAN.R-
project.org/package=vegan> (2017).
122   Bates, D., Maechler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using
lme4. *Journal of Statistical Software* **67**, 1-48, doi:10.18637/jss.v067.i01 (2015).
123   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**, 550, doi:10.1186/s13059-014-0550-8 (2014).
124   Pantano, L. *DEGreport: Report of DEG analysis. R package version 1.26.0*,
<http://lpantano.github.io/DEGreport/> (2020).
125   Taj, B. *et al.* MetaPro: A scalable and reproducible data processing and analysis pipeline for
metatranscriptomic investigation of microbial communities. *BioRxiv*,
doi:https://doi.org/10.1101/2021.02.23.432558 (2021).
126   Garber, A. I. *et al.* FeGenie: A Comprehensive Tool for the Identification of Iron Genes and Iron
Gene Neighborhoods in Genome and Metagenome Assemblies. *Front Microbiol* **11**, 37,
doi:10.3389/fmicb.2020.00037 (2020).
127   Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using DIAMOND. *Nat*
Methods **12**, 59-60, doi:10.1038/nmeth.3176 (2015).

1364

1365

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7