The commensal protist *Tritrichomonas musculus* exhibits a dynamic life cycle that induces extensive remodeling of the gut microbiota

Authors

Ana Popovic[1,2], Eric Yixiao Cao[3], Joanna Han[3], Nirvana Nursimulu[1,4], Eliza V.C. Alves-Ferreira[5], Kyle Burrows[3], Andrea Kennard[5], Noor Alsmadi[3], Michael E. Grigg[5], Arthur Mortha[3], John Parkinson[1,2,6,*]

Affiliations

[1]Program in Molecular Medicine, Hospital for Sick Children, Toronto, Ontario, Canada

[2]Department of Biochemistry, University of Toronto, Toronto, Ontario, Canada

[3]Department of Immunology, University of Toronto, Toronto, Ontario, Canada

[4]Department of Computer Science, University of Toronto, Toronto, Ontario, Canada

[5]Molecular Parasitology Section, Laboratory of Parasitic Diseases, NIAID, National Institutes of Health, Bethesda, Maryland, USA

[6]Department of Molecular Genetics, University of Toronto, Toronto, Ontario, Canada

Corresponding authors

*To whom correspondence should be addressed:

jparkin@sickkids.ca

Keywords: Eukaryotic microbiota; Protists; Tritrichomonas; Microbiome; Gut pathogens

Abstract

Commensal protists and gut bacterial communities exhibit complex relationships, mediated at least in part through host immunity. To improve our understanding of this tripartite interplay, we investigated community and functional dynamics between the murine protist *Tritrichomonas musculus* (*T. mu*) and intestinal bacteria in healthy and B cell-deficient mice. We identified dramatic, protist-driven remodeling of resident microbiome growth and activities, in parallel with *T. mu* functional changes, accelerated in the absence of B cells. Metatranscriptomic data revealed nutrient-based competition between bacteria and the protist. Single cell transcriptomics identified distinct *T. mu* life stages, providing new evidence for trichomonad sexual replication and the formation of pseudocysts. Unique cell states were validated *in situ* through microscopy and flow cytometry. Our results reveal complex microbial dynamics during the establishment of a commensal protist in the gut, and provide valuable datasets to drive future mechanistic studies.

Introduction

Research into host-associated microbiomes has established the critical role of gut bacteria in health and disease[1,2]. Emerging data supports an appreciable presence of gut-dwelling protists with host immunomodulatory capabilities and influence over resident bacterial communities, suggesting these microbes are essential players in organizing the gut environment[3-7]. Studies in this discipline are in their infancies however and lack understanding of both the ability of protists to colonize and persist and protist:bacterial cross-talk in the context of host immunity.

The protist *Tritrichomonas musculus* (*T. mu*) is a common resident of the murine gut, and a relative of prevalent human trichomonads with links to colitis, irritable bowel syndrome and colon cancer[5,8-12]. *T. mu* establishes chronic asymptomatic infections, resulting in elevated baseline intestinal immune activation and structural remodelling of the intestinal epithelium[6,13-15]. Protist-produced succinate furthermore stimulates a Th2-based immune response that restricts infection by enteric parasites[14-16]. Although the sustained inflammation associated with *T. mu* confers protection to mice against pathogenic bacteria, it can increase susceptibility to T-cell-driven colitis and tumorigenesis, suggesting *T. mu* behaves as a pathobiont. Evidence shows its presence influences gut bacterial composition, and that conversely *T. mu* engraftment may be modulated by bacterial taxa such as *Bifidobacterium* spp. through unknown mechanisms[17]. These microbial interactions may in part be driven through host immunoglobulins (Ig) A and M, antibodies secreted by B cells and elevated in response to *T. mu*[6,15]. Both bacteria and protists have known effects on B cell antibody responses, but it is not well understood how host B cells modulate cross-kingdom microbial interactions[6,18,19].

Here we track *T .mu* activities during the first 28 days of gut colonization, together with its interactions with resident bacteria in the context of a healthy WT host and B cell-deficient (*muMT*[-/-]) mice through 16S sequence surveys and microbial transcriptomics. We show that *T. mu* expansion is accelerated in the

absence of host B cells. Its colonization dramatically remodels bacterial composition and function,

concurrent with shifting protist metabolism and virulence factor expression - changes accelerated in the

absence of host B cells. We provide detailed transcriptomic analyses suggesting cross-kingdom competition

for key dietary nutrients. Finally, we conduct single-cell level characterization of the *T. mu* life cycle and

validate cell stages *in situ* through fluorescence-based labelling of stage-specific transcripts. Our data reveal

B cell-modulated co-adaptation between resident bacteria and a newly colonizing protist, and provide

valuable datasets to drive future mechanistic studies.

Materials and Methods

Mice

C57Bl/6J and B6.129S2-*Ighm^{tm1Cgn}*/J (*muMt^{-/-}*) female mice (Jackson Laboratory, Bar Harbor, ME, USA)

were maintained in specific pathogen free (SPF) conditions at the University of Toronto Division of

Comparative Medicine. Experiments were conducted at 6-8W of age, using age and sex matched controls.

For single-cell protist sequencing, 12W germ-free (GF) female C57Bl/6 mice (Taconic Biosciences,

Germantown, NY, USA) were housed in the GF facility until the day of analysis, when they were

transported to SPF conditions. Mice conventionalized with a microbiome received a bacterial suspension

through oral gavage derived from two C56Bl/6J fecal pellets in PBS, and were maintained under SPF

conditions for 4 weeks prior to protist colonization. Animals were housed in a closed caging system and

provided with an irradiated chow diet (Envigo Teklad 2918), non-acidified water (reverse-osmosis and UV-

sterilized) with a 12hr light/dark cycle. Following *T. mu* colonization, animals were housed in separate

cages. Animal experiments were approved by the Local Animal Care Committee (LACC) at the Faculty of

Medicine, University of Toronto (animal use protocol 20012400 to AM).

Protist colonization

Purification of *T. mu* was performed as previously described[6]. Briefly, cecal contents of a colonized

C57Bl/6J mouse were filtered through a 70µm cell strainer, washed with PBS, and protists were collected

from the interphase after density centrifugation through 40% Percoll overlaid on 80% Percoll. Cells were

sorted into PBS on a BD Influx Cell Sorter (BD Biosciences, Franklin Lakes, NJ, USA) using the 100µm

nozzle at 27psi at 4°C, with >99% purity. Mice were orally gavaged immediately after with two million *T.*

mu cells. Protists were quantified using a hemocytometer.

Genome sequencing and annotation

Genomic DNA was extracted from sorted protists using the MagAttract HMW DNA Kit (QIAGEN, Hilden,

Germany), and sequenced using PacBio Sequel technology on two SMRT cells at the McMaster University

Farncombe Metagenomics Facility (Hamilton, Canada). Reads were error corrected using CANU v1.8[20],

assembled into contigs using Flye v.2.4.1[21], and subsequently polished with 8.6 million 300 bp paired-end

Illumina MiSeq reads sequenced at the National Institute of Allergy and Infectious Diseases (Bethesda,

Maryland) using BWA[22] and Pilon v.1.23[23]. Annotation was carried out with the Maker v2.31 pipeline[24]

using SNAP v.2013-11-29[25] and Augustus v.3.3.1[26] for gene model prediction. Functional annotation was

performed with InterProScan v.5.30-69.0[27], HmmerWeb v.2.41.2[28] and Architect[29]. Genes encoding

adhesins, meiosis and cell cycle-related proteins were identified based on sequence homology with *T.*

vaginalis proteins retrieved from the TrichDB database[30-32]. Detailed methods are provided in

supplementary   information.   The   genome   assembly   is   accessible   at:

https://github.com/ParkinsonLab/Tritrichomonas-murine-microbiome-interactions/draft-genome-

assembly.

Phylogenetic analysis

Parabasalid ribosomal internal transcribed spacer (ITS) sequences were downloaded from GenBank[33]

(Table S12) and compared to ITS in the *T. mu* genome assembly. Multiple sequence alignments were

generated using MUSCLE, and an unrooted phylogeny was constructed using the Maximum-likelihood

method with a Tamura-Nei model in the MEGA-X software[34-36]. Bootstrap values were generated using

1000 replicates.

scRNA-Seq analysis

A GF and conventionalized mouse were colonized with *T. mu* through oral gavage, as described above.

Four weeks post colonization, protists were purified from caecal contents and immediately transferred on

ice to the Princess Margaret Genomics Centre (Toronto, Canada) for STAMP library preparation using

Drop-seq technology and sequenced on a NextSeq 500 (Illumina, San Diego, CA, USA)[37]. Reads were

processed using Drop-seq Tools v.1.13 and aligned to the protist genome assembly using STAR v.

2.5.3a[38,39]. Three thousand protists per mouse (minimum 200 genes, 500 transcripts) were analyzed in

Seurat v4 and the two transcriptomes were aligned using the FindIntegrationAnchors feature[40,41]. Cells were

grouped using graph-based clustering (0.8 resolution, 18 principal components) and visualized via

UMAP[42]. Differentially expressed (DE) genes were identified using the FindAllMarkers function, and

functional enrichments were determined based on overrepresentation of pathway enzymes as defined by

KEGG using the hypergeometric test, or GO terms using the topGO package and the Fisher's Exact test[43,44].

Enrichments of custom-defined gene sets (meiosis, G1/S and G2 phase genes) were scored with the

AddModuleScore function, and evaluated using two-sided Wilcoxon rank-sum tests. Benjamini-Hochberg

correction was applied for multiple testing[45]. Heatmaps were generated using pheatmap 1.0.12 and

Ward.D2 clustering[46].

16S microbiome profiling and metatranscriptomics

Groups of four WT or *muMt*[-/-] 6W female mice were infected with protists for microbial composition and

gene expression profiling. A schematic of the experiment is presented in Fig. 1b.

16S profiling. Mouse fecal pellets were collected from the same group of WT or *muMT[-/-]* mice at days 0, 2,

7, 14, 21 and 28, or uninfected control mice at day 28, and stored at -80°C. DNA was extracted using the

DNeasy PowerSoil Kit (QIAGEN, Hilden, Germany) according to manufacturer instructions, with a 5 min

bead beating step using the FastPrep-24 homogenizer (MP Biomedicals, Santa Ana, CA, USA) at 5M/s.

16S rRNA amplicons were generated using B969F and BA1406R primers and paired-end sequenced (250

bp) with MiSeq v2 chemistry (Illumina, San Diego, CA, USA) at the Integrated Microbiome Resource

facility at Dalhousie University (Halifax, Canada)[47]. Reads were pre-processed using Dada2 within

QIIME2, subsequently *de novo* clustered to 97% operational taxonomic units (OTUs), and classified with

a custom classifier trained on V6-V8 regions extracted from the SILVA v132 database[48-50]. Diversity and

multivariate analyses were performed with Phyloseq 1.26.1[51] and vegan v.2.5[52]. Average alpha diversities

were evaluated from 100 independently rarefied datasets to the minimum read depth, using linear mixed

effects (LME) models with lme4 or a two-way ANOVA[53]. Beta diversities were determined for rarefied

OTU data and tested using the adonis function in vegan. Differentially abundant taxa were identified with

DESeq2 v.1.22.2 and grouped by abundance pattern with DEGreport[54,55].

Metatranscriptomics. Caecal content from groups of four WT or *muMT[-/-]* mice was collected at day 2 or 28

of the experiment. RNA was extracted using TRIzol (Invitrogen, Carlsbad, CA, USA) and homogenized

with 0.1 mm Zirconia beads (BioSpec Products, Bartlesville, OK, USA) using a TissueLyser (QIAGEN,

Hilden, Germany), followed by the PureLink RNA Mini Kit (Invitrogen, Carlsbad, CA, USA) according to

the manufacturer TRIzol Plus Total Transcriptome Isolation protocol. rRNA was depleted using the

QIAseq FastSelect rRNA HMR and 5S/16S/23S kits (QIAGEN, Hilden, Germany), and libraries were

paired-end sequenced (150 bp) on a NovaSeq 6000 (Illumina, San Diego, CA, USA) at Genome Quebec

(Montréal, Canada). Reads were processed using the MetaPro pipeline. High quality reads were mapped to

the *T. mu* genome assembly using STAR v. 2.5.3a, and all remaining unmapped reads were returned to

MetaPro for functional and taxonomic annotation[39,56]. Proteins with iron-related functions were predicted

using FeGenie[57]. Putative cell division and cell wall biogenesis machinery was identified through sequence

similarity with previously described proteins in the *E. coli* MG1655 proteome using DIAMOND

v0.9.22[58,59]. DE was evaluated with DESeq2 v.1.22.2[54]. Due to read depth differences of bacterial sequences

in protist-colonized and uninfected mice, reduced due to protist rRNA (Table S4), bacterial gene counts

were rarefied to 600,000 reads per sample, and DE was averaged from 15 independent rarefactions.

Principal component analyses were performed on DESeq2-variance stabilized counts. Functional

enrichment was evaluated using EC or GO term overrepresentation as described above.

RNA-FISH Flow Cytometry

Four million purified protists were stained with Zombie Aqua viability dye (BioLegend, San Diego, CA,

USA) in PBS in the dark (1:1000, 15min, 4°C), washed twice with PBS, and the remaining steps were

performed as described in the Stellaris RNA-FISH protocol for cells in suspension. After hybridization,

cells were sorted on a BD LSR Fortessa X-20 cell analyzer (BD Biosciences, Franklin Lakes, NJ, USA).

To test for pseudocyst formation, protists were isolated from the mouse caecum or colon and stained with

wheat germ agglutinin (WGA)-FITC (Sigma Aldrich, St. Louis, MO, USA) in FACS buffer (PBS, 2% FBS,

5 mM EDTA) in the dark, either immediately or 1-3 days after anaerobic in vitro culturing. Cytospins were

visualized using a Zeiss AXIO Observer microscope (Carl Zeiss AG, Jena, Germany).

RNAScope

RNAScope was performed as per the RNAScope Multiplex Fluorescent Reagent Kit v2 (Advanced Cell

Diagnostics, Newark, CA, USA) protocol. Approximately 0.5 cm cecum sections were excised from *T. mu-*

colonized mice, placed in 10% neutral buffered formalin and fixed overnight at room temperature (RT)

with gentle agitation. The following day, samples were washed with PBS, placed in 70% ethanol, embedded

in paraffin and sliced to 7 μm sections at the Toronto Centre for Phenogenomics. Paraffin sections were

baked at 40°C for 30 min in a HybEZ Oven (Advanced Cell Diagnostics, Newark, CA, USA), and treated

with hydrogen peroxide at RT for 10min. Antigen target retrieval was conducted at 99°C under the 15 min

standard procedure. A barrier was created around sections using an ImmEdge pen (Vector Laboratories,

Burlingame, CA, USA) and allowed to dry for 15 min. Samples were treated with protease at 40°C for 30

min, and stored in saline sodium citrate solution overnight (175.3 g NaCl, 88.2 g sodium citrate, 800 mL

ddH$_2$O, pH 7). TSA Plus Fluorophores Fluorescein and Cyanine 3 were hybridized against protist probes

TMU_00005724 and TMU_00016742 respectively, and samples were visualized using a Zeiss AXIO

Observer microscope (Carl Zeiss AG, Jena, Germany).

Transmission Electron Microscopy

Protist pellets were prepared using standard methods for the Embed 812 resin kit (Electron Microscopy

Sciences, EMS)[60]. Briefly, samples were fixed with 4% paraformaldehyde, 1% glutaraldehyde in phosphate

buffer (PB, 0.1M, pH 7.2), followed by 1% OsO4 in PB in the dark. After washing with PB, they were

dehydrated in a 30%-100% gradient ethanol series, and infiltrated with increasing amounts of Embed 812

resin (EMS) in propylene oxide and cured in molds at 65°C. 80 nm sections were prepared with a Reichert

Ultracut E microtome (Leica) on 300 mesh copper grids (EMS), and counter stained with saturated 5%

uranyl acetate (EMS), followed by Reynold's lead citrate (EMS). Sections were imaged using a Talos

L120C transmission electron microscope (ThermoFisher Scientific, Waltham, Massachusetts). See

supplementary methods for details.

qPCR

208   qPCR of fecal DNA was performed using PowerTrack SYBR Green Master Mix (Applied Biosystems,

209   Waltham, MA, USA). In brief, 10 ng DNA was mixed with 2X PowerTrack Master Mix and 400 nM

210   primers, and treated as follows: enzyme inactivation at 95°C for 2 min, 40 cycles of denaturation at 95°C

211   for 15s and annealing/extension at 60°C for 60s. Relative abundances were normalized to the universal 16S

212   gene, and calculated using the delta-delta Ct method. Primers are provided in Table S13.

Results

***T. mu* colonization drives diversification of the intestinal microbiome**

Colonization with *T. mu* induces profound changes in the mouse gastrointestinal tract, including the immune landscape known to maintain gut microbial homeostasis[6,19]. In light of this, we characterized the impact of *T. mu* engraftment on the local bacterial community in a healthy and immune-impaired host (Fig. 1a). We tracked bacterial composition and activities in WT and B cell-deficient (*muMt*[-/-]) C57Bl/6 mice through 16S rRNA surveys of mouse fecal pellets and metatranscriptomics of caecal microbiota (Fig. 1b, Tables S1-S3) during the first 28 days of *T. mu* colonization. Protist expansion was accelerated in *muMt*[-/-] mice, reaching its maximum level within the first 7 days, while in WT mice, expansion steadily increased until day 28 (Fig. 1c).

T. mu dramatically altered gut bacterial composition in both WT and *muMt*[-/-] mice (Fig. 1d). Bacterial richness increased in the first week of colonization, with a delay in *muMt*[-/-] mice, before receding to near day 0 levels, while increased evenness persisted to day 28, reflecting expansions in multiple taxa (Fig. 1e). Communities in WT and *muMt*[-/-] mice differed prior to infection, but converged toward similar profiles over the course of the experiment (Fig. 1f). Those in day 28 control mice remained similar to day 0 naïve bacterial compositions, underlining a protist-driven shift. We observed six distinct patterns of growth dynamics in *T. mu*-modulated bacteria (Fig. 1g, Table S4). The majority changed congruently in WT and *muMt*[-/-] mice, and included taxa that increased throughout (e.g. *Bifidobacterium* and *Parabacteroides*), peaked at day 21 (e.g. *Helicobacter* and *Lachnospiraceae*), or decreased after *T. mu* colonization; five taxa exhibited discordant patterns (e.g. *Parasutterella* and *Bacteroides*). Changes in bacterial abundances were validated through qPCR (Fig. 1h).

T. mu drives changes in bacterial metabolism

238
To investigate the impact of *T. mu* on community functions, we performed bulk RNA sequencing of mouse

caecal contents at days 2 and 28 (early and chronic stages of infection) (Supplementary Fig. 1 and Table

S3). 835 million reads were identified as bacterial (741 million in naïve and 94 million in colonized mice)

and 157 million as *T. mu*. Colonization with *T. mu* associated with blooms in activities of Proteobacteria

and Bacteroidetes (Fig. 2a). The increase in proteobacterial gene expression was accelerated in *muMT*[-/-]

mice, accounting for 14% of reads by day 2 compared to 4% in colonized WT mice and 1-3% in naïve

mice, suggesting a role for B cells in regulating protist-induced bacterial gene expression. The majority of

these reads mapped to *Helicobacteraceae* (esp. *Helicobacter ganmani* and *Helicobacter rodentium*)

previously associated with enteric inflammation and exacerbated colitis[6,15,61].

Principal component analysis of bacterial gene expression re-capitulated the 16S data, with an overall

upregulation of bacterial genes in *T. mu*-colonized mice (Fig. 2b,c). Although *muMt*[-/-] mice exhibited higher

protist-dependent upregulation at day 2, B cell-associated differences converged over time (Fig. 2c).

Pathway analysis revealed upregulation of multiple metabolic pathways including glycolysis, metabolism

of amino acids, and biosynthesis of peptidoglycans, polysaccharides, O-Antigen sugars and fatty acids (Fig.

2d). Metabolic activity shifted from Firmicutes at day 2 to Bacteroidetes and Proteobacteria by day 28

(Supplementary Fig. 2, 3). The expansion of Proteobacteria, particularly in the expression of succinyl-CoA

synthetase (SucC) which is typically suppressed under anaerobic conditions, may indicate increased

exposure to oxygen characteristic of gut inflammation and dysbiosis (Supplementary Fig.2)[62].

Given the central role of these pathways in bacterial growth, we examined the expression of homologues

to 44 genes associated with bacterial cell division and cell wall biosynthetic machinery in *E. coli* (Fig. 2e)[58].

Patterns were consistent with those described—genes were upregulated during *T. mu* colonization, with

earlier responses in *muMt*[-/-] mice and a shift from predominantly Firmicutes at day 2 to Bacteroidetes and

Proteobacteria (primarily *Helicobacter*) at day 28. Day 2 bacteria from *muMt*[-/-] mice were associated with

increased chromosome segregation (e.g. *ftsK* and *ftsZ*), cell elongation (e.g. *mrdA* and *mrdB*) and

peptidoglycan biogenesis (e.g. *murCDE*). By day 28, these genes were upregulated in bacteria across both

hosts, together with genes involved in protein secretion (e.g. *secA*, *yjjK*, *prlA*) and porin *ompA*, which

modulates infection and host immunity in Gram-negative bacteria. Collectively, these data demonstrate

substantial changes in bacterial activities in response to a new protist, and a shift to an ecological niche

favouring the growth of Gram-negative taxa.

***T. mu* activity changes in a B cell-dependent manner**

To monitor *T. mu* gene expression during colonization, we generated a draft assembly of the protist genome

from one million PacBio sequences (median length 11,924 bp) and 8.6 million Illumina MiSeq reads.

Assembly statistics are available in Tables S4 and S5. Functional annotation predicts 26,723 genes, with

999 unique GO terms, 417 enzymes and 1,932 Pfam domains (Table S6). We also identified a second

distinct rRNA locus with 97% sequence identity to *T. mu*, suggesting the presence of an additional organism

at lower abundance (19% of rRNA reads) (Fig. 3a). Phylogenetic comparison confirmed this novel species

also belonged to the Tritrichomonadidae order (Fig. 3b).

We mapped 157 million metatranscriptomic reads to the protist, of which 20% mapped to coding regions

(Table S3). *T. mu* gene expression changed during colonization in both WT and *muMt*[-/-] hosts (666 and 627

DE genes, respectively), and a host B cell-modulating effect was apparent at day 28 (Fig. 3c,d). Consistent

with protist growth, *T. mu* upregulated the expression of enzymes involved in energy production (e.g. the

tricarboxylic acid cycle (TCA) and glycolysis) and carbohydrate metabolism (e.g. galactose, and starch and

sucrose) (Fig. 3e). We probed genes with potential roles in establishing colonization: adhesins, BspA-type

Leucine rich repeat regions, lectins and cysteine proteases (Tables S6-S8), protein families previously

implicated in facilitating host interactions and parasite infection[31,63-66]. Homologues to adhesins AP33 and

AP120, and TvBspA625, proteins with documented roles in *T. vaginalis* virulence and host

cytoadherence[31,63], were upregulated at day 28—already at day 2 in *muMt*[-/-] mice (Fig. 3f). The expression

of four additional BspA-like proteins with predicted extracellular immunogenic regions changed in B cell-

dependent manners, upregulated only in WT mice (TMU_00001180, TMU_00003472, TMU_00005602,

TMU_00010916; Fig. 3g, Table S9). The remaining genes showed varied changes during colonization

and/or in the presence of B cells (Supplementary Fig. 4b), suggesting dynamic metabolic and accessory

functions required as the protist adapts to the host environment. Future research is set to validate the role

of these proteins.

Single cell profiling of *T. mu* reveals distinct life cycle stages

The trichomonad life cycle features a variety of cell states (e.g. trophozoite, amoeboid, pseudocyst). To

investigate shifts in *T. mu* cell states during colonization we carried out transcriptional profiling at the single

cell level (Table S10). To further explore whether resident bacteria and B cells can influence these states,

we characterized protists from a conventionalized and GF mouse, as GF mice are known to possess an

immature B cell compartment[67]. Clustering of gene expression profiles from 6,000 protists (3,000 per host)

revealed 15 distinct cell populations, assigned to three 'superclusters' (Fig. 4a). Clusters 1 through 4

(designated supercluster A) and clusters 10 and 11 were enriched in protists from the GF mouse, while

clusters 5 through 7 and 15 were composed primarily of protists from the conventionalized mouse (Fig.

4b).

Functional enrichment of GO terms suggested the presence of actively metabolizing cells, protists at various

stages of cell cycle, and those undergoing pseudocyst formation (Fig. 4c). Clusters 1-3 and 5-8, for example,

were enriched in cytoskeleton & actin remodeling, carbohydrate metabolism and biosynthetic processes

and may represent actively feeding and growing *T. mu* trophozoites. Cluster 8 is enriched in sphingolipid

metabolism, which contributes to regulation of *Giardia* encystation[68]. Since cluster 14 is enriched in

biogenic amine metabolism, a pathway associated with encystation in *Entamoeba invadens*[69], and exhibits

lowest transcriptional complexity (Fig. 4d), we suggest these cells represents a pseudocyst form of *T.*

mu[70,71]. Furthermore, the upregulation of oxidative stress response in the neighbouring clusters 12 and 13

suggests the encystation transcriptional program may be triggered through stress-induced pathways.

Enrichment of chromosome reorganization, DNA replication and cell cycle checkpoint in clusters 1, 5, 10

and 11 identified protists undergoing cell cycle. While cluster 15 shared cell cycle terms with these clusters,

it was uniquely enriched in DNA recombination, indicative of sexual or parasexual replication. To

investigate further, we generated functional enrichment scores for the expression of *T. mu* homologs

associated with the G1/S and G2 stages of the cell cycle, and those associated with meiosis (Tables S6 and

S7)[30,72]. Scores for expression of G1/S marker genes were highest in clusters 1 and 11, and cluster 10 scored

highest for the G2 phase (Fig. 4e,f; Supplementary Fig. 5a,b). Cluster 15 scored highest for meiosis-related

markers (Fig. 4f). In addition to meiosis-specific genes *dmc1*, *hop2A* and *mnd1*, required genes *rad1*, *mre11*,

smc2, *smc3* and *smc5* were also significantly upregulated in cluster 15 (Supplementary Fig. 5c), consistent

with sexual or parasexual replication. Of the 60 cells associated with this this cluster, only 8 were from the

GF mouse. Together with other differences in cluster distributions between conventionalized and GF mice

(Fig. 4b), our results suggest that the constituent microbiome may play a role in modulating the protist life

cycle. However, we note that despite purification and antibiotic treatment of the protist, we detected

bacterial DNA in the GF mouse, limiting our conclusions on microbiome impact.

Protist populations express distinct sets of virulence genes

Since the distribution of protist cell states was altered in the GF mouse, we investigated how these changes

might impact the expression of genes implicated in colonization (BspA-like proteins, adhesins, lectins and

cysteine proteases; Fig. 4g). Many genes exhibited cluster and/or host-specificity, with profiles broadly

defined by the three major superclusters (Table S11). Notably, cells in cluster 15 (the putative sexual stage)

expressed distinct BspA-like genes in GF and conventionalized hosts, implicating a role for the microbiome

in regulating the protist life cycle, while putative encysting stages (clusters 12-14) shared similar expression

patterns in both mice. Of the nine putative adhesins, which mediate binding to host epithelia in *T.*

vaginalis[31], we detected expression for eight (Fig. 4g). These genes were absent from pseudocyst-associated

clusters 13 and 14, supporting their role in colonization. Lectins were depleted in clusters 12-14, as well as

the putative sexual stage, cluster 15. Of the four cysteine proteases expressed in putative pseudocysts, two

are members of the C54 peptidase family which have previously been implicated in cell starvation and

differentiation[73,74]. These data present a dynamic arsenal of virulence factors, exhibiting cell stage and host-

specific expression, potentially modulated by resident microbiota, which may mediate protist colonization

and transmission.

Distinct *T. mu* cell stages are detected *in vivo*

We validated the presence of distinct *T. mu* populations in the mouse intestine through *in situ* fluorescent

labelling of cluster-specific mRNAs. Microscopic imaging of caecal sections from WT mice revealed

protists as large, nucleated cells restricted to the gut lumen (Fig. 5a). Protists expressing one or more cell-

stage specific gene could be visibly identified in the cecum (e.g. TMU_00005724, specific to cells of

supercluster B, and/or TMU_00016742 expressed in a subset of B; Fig. 5b,c). Fluorescent *in situ*

hybridization flow cytometry (FiSH-Flow) revealed expression of TMU_00005724 in all caecal protists

(Fig. 5d). Consistent with cluster-specific profiles, expression of TMU_00016742 (clusters 5, 6 and 9)

exhibited higher overlap with TMU_00005724 than did TMU_00009244 (cluster 12) and TMU_00001185

(cluster 14) (Fig. 5c,e). *muMT*[-/-] mice harbored more TMU_00016742-expressing protists, suggesting B

cell influence on the *T. mu* transcriptional program (Fig. 5e). Since TMU_00001185 was unique to cells of

cluster 14 (predicted pseudocysts) and pseudocyst formation occurs during host egress, we tested for its

expression in protists freshly isolated from the caecum, colon, and for up to 3 days of *in vitro* culturing

(Fig. 5f). As expected, caecal isolates contained fewest TMU_00001185-expressing cells, and their

proportions increased during *in vitro* culturing. Supporting their identity as pseudocysts, the same pattern

was observed for presence of chitin, a critical component of cysts and pseudocysts, stained using

fluorophore-labelled wheat-germ agglutinin (WGA) (Fig. 5g, Supplementary Fig. 6)[71,75,76]. WGA-labelled

protists furthermore showed thicker cell walls, consistent with cyst-like structures (Fig. 5h). The *in situ* data

confirm the presence of transcriptionally distinct groups of protists in the cecum, and illustrate the utility

of cluster-specific genes to monitor *T. mu* life cycle dynamics during colonization.

_T. mu_ and gut bacteria compete for nutrients

As *T. mu* colonizes the gut, it must compete with resident microbiota and the host for nutrients. Since

trichomonads require high concentrations of iron for growth[77,78], we hypothesized that its consumption by

T. mu would exert pressure on bacterial iron acquisition and storage systems. Metatranscriptomic data

confirmed increasing protist iron consumption through the upregulation of ferredoxin-1 (TMU_00007489),

an enzyme required for energy production, and also adhesin TMU_00017025 with predicted

pyruvate:ferredoxin oxidoreductase activity (Fig. 6a,3f)[79,80]. Both enzymes were widely expressed across

protist cell states, but reduced in clusters 13 and 14, consistent with lower energy requirements of a

pseudocyst (Fig. 6b). The upregulation instead of ferredoxins TMU_00018447 and TMU_00005891 in

clusters 12 and 14 may be specific to encysting forms, and potentially involved in mediating stress response.

Similar expression of these genes in conventionalized and GF mice suggests *T. mu* is agnostic in terms of

iron acquisition to the resident microbiome. In response to *T. mu*, expression of bacterial iron acquisition

and storage systems increased, including siderophore synthesis and transport genes (Fig. 6c,d),

predominantly in *Helicobacter*, *Bacteroides*, *Parabacteroides* and *Mucispirillum schaedleri*

(Supplementary Fig. 7b). Host B cell status minimally impacted these systems, suggesting more direct

competition between *T. mu* and resident microbiota (Supplementary Fig. 7c).

We also tracked expression of genes associated with tryptophan and arginine metabolism, amino acids

implicated in host:microbiome interactions[81,82]. The protist tryptophanase TMU_0006921 (EC 4.1.99.1)

was upregulated at day 28 (Fig. 6e). As enzymes in competing pathways (conversion of tryptophan to

kynurenate and tryptamine) exhibited minimal expression with no temporal shift, *T. mu* appears to

preferentially drive production of indole (Fig. 6e). Remarkably, the two most highly expressed bacterial

tryptophanases were also upregulated in *T. mu* colonized-mice suggesting bacteria further promote the

indole production in the presence of the protist (Fig. 6f). Consistent with this, scRNA-Seq data showed

upregulation of *T. mu* tryptophanase in the conventionalized mouse and reduction in putative pseudocyst

clusters (13 and 14) suggesting reduced dependence on host mucosal homeostasis (Fig. 6e).

Arginine is an important source of energy for both trichomonads and bacteria. Its depletion by microbes

through the arginine dihydrolase pathway has an additional immunomodulatory role, limiting host

production of antimicrobial NO[81,82]. None of the three *T. mu* enzymes in the associated pathway: arginine

deiminase (ADI), ornithine transcarbamylase (OTC) or carbamate kinase (CK), were significantly

associated with either colonization time or host B cell status, suggesting continual arginine consumption

(Supplementary Fig. 4d). We did note selective expression across protist cell states (Fig. 6g). Conversely,

T. mu colonization was associated with increased expression of several bacterial arginine dihydrolase genes,

including a Lactobacillales ADI and *Helicobacter* OTC (Fig. 6h).

The dynamics of the acquisition and metabolism machinery for these key gut nutrients suggests competitive

as well as cooperative relationships between the protist and the resident microbiota.

Discussion

Previous work has revealed that protists mediate antagonistic and mutualistic interactions with intestinal

bacteria with consequences for the host, primarily through studies of microbial composition[6,7,15,83,84]. Here

we demonstrated interactions between *T. mu* and resident microbiota at a functional level in a healthy and

immunodeficient host, and identified genes predicted to facilitate protist colonization. We showed that,

consistent with a previous study, *T. mu* induces longitudinal shifts in bacterial composition[17]. However, our

study notes increased bacterial diversity and importantly the abundance and activity of *Helicobacter* spp.

in the presence of *T. mu*, reported also during *T. foetus* colonization[85]. The shift from Firmicutes to

Bacteroidetes and Proteobacteria in colonized mice suggests environmental pressure favouring Gram

negative taxa. As Gram negative bacteria harbor surface LPS capable of boosting immunity, their bloom

may contribute to the intestinal immune activation observed during *T. mu* engraftment[6,86]. The earlier

occurrence of these changes in B cell-deficient mice suggests a failure to control commensal bacteria that

might otherwise be mediated through *T. mu*-induced IgA reactivity[19,87,88]. We speculate this reactivity might

serve to limit adverse responses by the bacteria as the protist attempts to colonize. Similarly, the accelerated

expansion of *T. mu* and earlier upregulation of its adhesins and TMU_00015121, a homologue of the

antigenic *T. vaginalis* BspA625 protein, signals changes in protist immunogenicity and suggests that B cell

surveillance plays a critical role during early protist infection[63,89].

Within our scRNASeq data, we identified distinct protist populations representing different stages of the *T.*

mu life cycle. Cluster-specific patterns of expression of protein families implicated in host cell binding or

parasite virulence suggest the protist alters its host adhesive properties or antigenicity as it transitions

through its life cycle[31,63,65,66]. *T. vaginalis* and *T. foetus* exist in one of three forms—actively growing

trophozoites, host-adherent amoeboid cells and environmentally resistant pseudocysts[70,71,90-92]. The capture

of actively metabolizing *T. mu* in the lumen, together with cells exhibiting reduced activity (with little to

no expression of adhesins) and stress-activated pathways suggests our data captured the trophozoite and

pseudocyst states. Supporting the ability of *T. mu* to penetrate the intestinal mucous layer and gain access

to host epithelia, we identified a large repertoire of expressed cysteine proteases and enzymes of the N-

glycan degradation pathway, capable of degrading O-glycans, the dominant glycans found in mucins[93,94].

We validated the pseudocyst stage using cluster-specific and chitin markers on mouse intestinal contents and *in vitro* cultures, and TEM of purified protists. A similar transcript-labelling approach could be used to capture the amoeboid state through exposure to host epithelia[90,91]. One possibility is that we have already captured the amoeboid state in our scRNA-Seq data, in supercluster A. The majority of protists in these populations were recovered from the GF mouse, which are known to have a more penetrable intestinal mucus[95]. These cells express different suites of adhesins, cysteine proteases and enzymes involved in arginine metabolism. Their identification has clinically relevant implications in a diseased or inflamed gut where the protist may further exacerbate symptoms[6]. The discovery of meiosis-specific genes is particularly intriguing as it suggests the presence of sexual replication, with implications for host adaptation and immune evasion. Sexual recombination has previously been hypothesized in *T. vaginalis* and *G. duodenalis*[30,96-99]. While our data may suggest that the microbiome promotes meiosis (as noted by the low proportion of meiotic cells in the GF mouse), we did detect bacterial DNA in the GF mouse. Differences in meiotic cells may therefore arise from timing of microbiome exposure. Although unlike the GF mouse, the conventionalized mouse was exposed to commensal bacteria for four weeks prior to *T. mu* infection, which would have affected intestinal morphology and immune cell differentiation, additional experiments are needed to confirm the role of the microbiome on the *T. mu* life cycle[100]. To date, attempts at axenic culturing of *T. mu* have proven challenging.

To successfully colonize, *T. mu* must compete with resident bacteria for nutrients. Parasitic trichomonads are known to require high quantities of iron to sustain their growth, whereas low iron conditions hinder their adhesion to host epithelia[78,101]. To limit growth and virulence of invading pathogens, hosts restrict access to luminal iron through sequestration by proteins such as lactoferrin[77,78,102]. The upregulation of bacterial iron acquisition machinery in response to *T. mu* has the potential, therefore, to regulate protist growth and virulence, an effect which may be lost in a context of surplus iron provided for example by

nutritional supplements[5]. We also noted synergistic increased expression of genes metabolizing tryptophan

to indole in *T. mu* and bacteria. Production of indole by *T. foetus* and *T. vaginalis* was shown to impact host

immunity and barrier function, and bacterial signalling[103-105]. Recently there has been much interest in the

production of neuroactive kynurenines from tryptophan by the gut microbiome[106,107]. Through promoting

production of indole at the expense of alternative products, tryptamine and kynurenine, our findings

question whether protists similarly modulate the gut-brain access.

The ecological shift induced by *T. mu* identifies it as a keystone species of the murine gut microbiome.

Applying similar sequence-based and *in situ* methodologies will provide insight into how other gut protists

might impact gut health.

Acknowledgements

This work was funded by the Canadian Institutes of Health Research (MRT-168043) to J.P., M.E.G. and

A.M; the Natural Sciences and Engineering Research Council (RGPIN-2019-06852) to J.P.; the Intramural

Research Program of the National Institute of Allergy and Infectious Diseases at the National Institutes of

Health to M.E.G.; a Restracomp scholarship administered by the Research Training Centre (Hospital for

Sick Children) and a graduate scholarship from the Government of Ontario to A.P. A.M. is supported by

the Canadian Foundation for Innovation John R. Evans Leaders Fund, the Canadian Institutes of Health

Research (PJT-388337, PJT-480765) and a Natural Sciences and Engineering Research Council (RGPIN-

2019-04521). A.M. is the Tier 2 Canadian Research Chair in Mucosal Immunology and supported by the

Tier 2 CRC-CIHR program (CRC-2021-00511). Computing resources were provided by the SciNet High

Performance Computing (HPC) Consortium; SciNet is funded by the Canada Foundation for Innovation

under the auspices of Compute Canada, the Government of Ontario, Ontario Research Fund - Research

Excellence, and the University of Toronto. We thank the Temerty Faculty of Medicine Microscopy Imaging

Lab at the University of Toronto for TEM sample preparation and training.

Author contributions

J.P., A.M., M.E.G., and A.P. conceived and designed the study. A.P. and A.K. isolated nucleic material for

high throughput sequencing, and A.P. performed sequence data analyses. E.Y.C. performed mouse

colonization experiments, performed flow cytometry and microscopic analyses. N.N. performed protozoan

enzyme annotation. N.A. and J.H. participated in sample collection and *in vitro* culturing experiments.

E.C.A.F. generated the phylogenetic tree. A.P., J.P., A.M. and M.E.G. wrote the manuscript. All authors

reviewed and/or edited the paper.

Competing Interests

The authors declare no competing interests.

Data Availability Statement

Sequence data generated in this study have been deposited to the NCBI Sequence Read Archive under the

BioProject identifiers PRJNA913581 and PRJNA914770.

 Figure Legends

Figure 1. *T. mu* colonization drives bacterial diversification.

a, Schematic of microbial interactions explored in the present study using 16S sequencing,

metatranscriptomics (RNA-Seq) and/or single-cell transcriptomics (scRNA-Seq). **b**, Experimental design.

Groups of four WT and *muMT*$^{-/-}$ mice were colonized with *T. mu*. Fecal pellets and caecal samples were

collected at indicated timepoints for 16S and metatranscriptomic profiling. **c**, Expansion of *T. mu* cells in

WT and *muMt*$^{-/-}$ mouse caeca over 28 days of colonization (n=4 or 5 per group). Differences were tested

using a two-sided t-test. **d**, Relative abundance of bacterial taxa determined through 16S sequencing of

mouse fecal pellets from two groups of WT and *muMt*$^{-/-}$ mice sampled over 28 days of colonization, and

two groups of uninfected controls sampled on day 28 (n=4 per group). Colours represent genus-level

abundances within indicated phylum colour groups. **e**, Bacterial richness (observed OTUs) and evenness

(Shannon diversity) over the course of *T. mu* colonization. Differences between timepoints were evaluated

using LME, and between colonized and naïve mice at day 28 using two-way ANOVA. **f**, Principal

coordinate analysis of Bray-Curtis dissimilarities between samples. Significance was tested using

permutational variance in Adonis. **g**, Patterns of bacterial abundance changes during *T. mu* colonization in

WT (grey) and *muMt*$^{-/-}$ (red) mice. Pie charts show the taxonomic make up of each cluster. **h**, Phylum-level

bacterial abundance before (day 0) or 21 days after protist colonization, determined through qPCR using

clade-specific primers. Abundances were normalized to Ct values obtained with universal 16S rRNA

probes. Bars show means. Significance was evaluated using the Wilcoxon rank-sum test: *$p < 0.05$, ** <

0.01, *ns* not significant.

Figure 2. Bacterial gene expression is altered in the presence of the protist.

a, Taxonomic profiles of putative bacterial mRNA reads. **b**, Principal component analysis of bacterial gene

expression. Each circle denotes a sample isolated from a single mouse, coloured by host B cell status,

presence of *T. mu* and timepoint, as shown. **c**, Numbers of significantly upregulated or downregulated

bacterial genes between protist-colonized and naïve mice, or due to host B cell status, grouped as indicated

(n = 3 or 4). **d**, Bacterial metabolic pathway enrichment associated with *T. mu* colonization. Enrichment

was determined by the overrepresentation of EC terms in differentially expressed genes of KEGG-defined

pathways. Total ECs detected in the experiment and in the rarefied gene expression matrix used in the

analysis are indicated in the bar charts to the right. Significantly enriched pathways are represented as

coloured dots: grey indicates *ns*, and the gradient from yellow to red represents decreasing *p* values,

beginning from *p*<0.05. Sizes of dots represent odds ratios. Significance was calculated using Fisher's exact

tests. Asterisks indicate infinite odds ratios. **e**, Network of cell division, cell wall biogenesis and secretory

genes, based on a previously determined *E. coli* protein network[58]. Coloured nodes represent protein

homologues identified in the metatranscriptomic data, and edges are previously described protein-protein

or functional interactions. Nodes in grey represent undetected genes. Genes significantly up- and

downregulated are denoted with blue and red borders, respectively. Node sizes correlate with log2-fold

changes between *T. mu*-colonized and naïve mice. Colours within node pie charts represent proportions of

gene expression assigned to various taxa, as shown, and colours encircling groups of proteins depict

functional modules as previously described.

Figure 3. Protist gene expression changes during colonization of WT and *muMt*-/- mice.

a, Graphical depiction of aligned rRNA sequences identified in the protist metagenome assembly. The

dominant sequence was synonymous with *T. mu* and mapped to 81% of reads; the second distinct sequence

mapped to 19% of reads. Sequences were aligned using MUSCLE. Mismatches in the sequence of the novel

organism to *T. mu* are indicated in black. **b**, Phylogeny of the two ribosomal ITS sequences identified in

the metagenome. An unrooted tree was constructed from aligned sequences using the Maximum-likelihood

method. Scale = number of SNPs per site. Bootstrap values >65% are shown. **c**, Principal component

analysis of protist gene expression patterns at days 2 and 28, in WT or *muMt*-/- mouse caeca. **d**, Numbers

of differentially expressed genes associated with colonization time or host B cell status. **e**, Changes in protist

metabolism during colonization. The dotplot represents enrichment of *T. mu* EC terms in KEGG-defined

pathways, within sets of genes differentially expressed over time or based on host B cell status. Coloured

dots represent significantly enriched pathways: the gradient from yellow to red marks decreasing *p* values,

beginning from $p < 0.05$. Grey dots are *ns*. Significance was evaluated using Fisher's exact tests, and dot

sizes correlate with odds ratios. Bar charts to the right show total ECs detected in the experiment. **f**,

Normalized expression of BspA-like genes showing interaction between time and host B cell status, and **g**,

differentially expressed BspA-like gene and adhesins homologous to *T. vaginalis* virulence factors.

Differences were evaluated using the Wilcoxon rank-sum test, and adjusted for multiple testing using the

Benjamini-Hochberg approach. $*p < 0.05$, $** < 0.01$, $*** < 0.001$, *ns*, non-significant.

Figure 4. Single cell transcriptional profiling of caecal *T. mu* populations.

a, Protist subpopulations isolated from colonized GF and conventionalized mouse caeca. Each point

represents one of 6000 protists cells, clustered by gene expression profiles using UMAP dimensionality

reduction, and coloured by cluster designation (3000 cells from each of two hosts). Groups of clusters,

designated 'superclusters' A, B or C, are indicated. **b**, Numbers of cells assigned to each cluster by host are

shown. Total cluster sizes are shown on the right. **c**, Heatmap of GO term enrichment within *T. mu* clusters.

Rows represent significantly enriched GO terms in one or more clusters (columns). Cells are coloured and

clustered by (log-transformed) numbers of upregulated transcripts per GO term. **d**, Transcriptional

complexity of protist clusters, defined as numbers of genes per transcript. Boxplots indicate medians and

interquartile ranges. **e**, Violin plots and **f**, feature plots depicting G1/S, G2 and meiosis scores in cells across

each *T. mu* cluster. Scores were determined based on the average expression of pre-defined gene sets

relative to randomly chosen control genes. Significance was evaluated using the two-sided Wilcoxon rank-

sum test, $*p<0.05$, $**<0.01$, $***<0.001$. Asterisks above clusters indicate significantly higher score relative

to all other clusters. Lines denote specific comparisons between two clusters. **g**, Heatmap showing

expression of (*top* to *bottom*) BspA-like proteins, putative adhesin proteins, lectins and Cysteine proteases

across protist clusters. Colours to the right of Cysteine proteases indicate the protein family (Pfam).

Figure 5. *T. mu* subpopulations are detected *in situ*.

a, Immunofluorescence micrograph of mouse cecal tissue after colonization with *T. mu* for 21 days. Nuclear staining (blue) shows host and protozoan DNA, and white dotted lines denote the mucus layer. Scale bar = 10μm. **b**, RNAscope immunofluorescence of *T. mu* transcripts TMU_00005724 (green) and TMU_00016742 (red) on sections of cecal tissue. Nuclear DNA is stained blue. Left image scale bar = 10μm; right images scale bars = 1μm. **c**, Feature plots showing the expression of cluster-specific genes. **d**, FISH-Flow analysis of freshly isolated *T. mu* at 21 days post colonization. Histogram shows control unstained cells (grey) and cells stained with fluorescent probes for TMU_00005724 (purple) **e**, FISH-Flow analysis of *T. mu* isolated 28 days post colonization from WT or *muMt$^{-/-}$* mice. Numbers adjacent to gates represent the average percentage +/-standard deviation of probe-expressing *T. mu*. Data shown is representative of 4 independent experiments, and adjacent plots show results from all experiments. Significance was tested using the Wilcoxon rank-sum test. **f**, Percentages of TMU_00001185 probe-positive or **g**, WGA-FITC stained protists freshly isolated from WT mouse caeca and colons, or cultured *in vitro* for one and three days. n=3 animals or culture plates per group, from four independent experiments. Significance was tested using one-way ANOVA, and adjusted for multiple comparisons using Tukey's test. **h**, Cell wall thickness of FACS-sorted protists hybridized or not to WGA-FITC. Cells were imaged using TEM, and cell walls were measured for all cells (5-10) across 7 view fields per group. Scale bars, 100 nm. Bars are means and whiskers show interquartile ranges. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, *ns* not significant.

Figure 6. *T. mu* and resident bacteria compete for resources in the gut lumen

Shown are changes in expression of bacterial and *T. mu* genes associated with iron, **a-d**, generation of tryptophan catabolites, **e** and **f**, and the arginine dihydrolase pathway, **g** and **h**. **a**, Expression of putative *T. mu* ferredoxin (Fdx) genes in metatranscriptomic data at days 2 and 28. **b**, PFOR and Fdx gene expression among protist subpopulations isolated from GF and conventionalized mice. Colour intensity correlates with average expression across cells and dot sizes represent percentages of cells in each cluster with detected expression. **c**, Enrichment of iron-related gene expression in bacteria, in the context of *T. mu* colonization. Shown are numbers of DE genes per iron-related gene family in *T. mu*-colonized versus naïve mice at days 2 and 28. Total numbers of genes identified in each family in the metatranscriptomic data are indicated in brackets. Significance of enrichment was tested using the hypergeometric test. **d**, Median log2 fold-change of bacterial siderophore transport and synthesis gene expression between colonized and naïve mice. Significantly DE genes are shown as coloured points: the gradient from yellow to red marks decreasing *p* values, beginning from *p*<0.05. **e**, Expression of *T. mu* and **f**, the most highly expressed bacterial (read count >100) enzymes with predicted function in the production of tryptophan metabolites. Dotplots in E show gene expression among protist subpopulations, as above. Pie charts in F represent the total numbers of tested and DE bacterial enzymes. Blue and green dots in the graphic show enzyme activities identified in *T. mu* and/or bacterial genes, respectively. **g**, Expression of *T. mu* enzymes associated with the arginine dihydrolase pathway among protist subpopulations isolated from GF conventionalized mice, as above. **h**, Expression of bacterial enzymes associated with the arginine dihydrolase pathway. Shown are the only ADI present in the rarefied gene count matrix, and the two differentially expressed OTC and CK genes. Gene expression in metatranscriptomics data was evaluated using DESeq2, and in scRNA-Seq data using Seurat. All expression values are derived from normalized gene counts. Boxplots and violin plots show medians and interquartile ranges. *P* values were corrected for multiple testing using the Benjamini-Hochberg approach. *p<0.05, **<0.01, ***<0.001

References

Fan, Y. & Pedersen, O. Gut microbiota in human metabolic health and disease. *Nat Rev Microbiol* **19**, 55-71, doi:10.1038/s41579-020-0433-9 (2021).

Libertucci, J. & Young, V. B. The role of the microbiota in infectious diseases. *Nat Microbiol* **4**, 35-45, doi:10.1038/s41564-018-0278-4 (2019).

Sardinha-Silva, A., Alves-Ferreira, E. V. C. & Grigg, M. E. Intestinal immune responses to commensal and pathogenic protozoa. *Front Immunol* **13**, 963723, doi:10.3389/fimmu.2022.963723 (2022).

Scanlan, P. D. *et al.* The microbial eukaryote Blastocystis is a prevalent and diverse member of the healthy human gut microbiota. *FEMS microbiology ecology* **90**, 326-330, doi:10.1111/1574-6941.12396 (2014).

Popovic, A. *et al.* Micronutrient supplements can promote disruptive protozoan and fungal communities in the developing infant gut. *Nat Commun* **12**, 6729, doi:10.1038/s41467-021-27010-3 (2021).

Chudnovskiy, A. *et al.* Host-Protozoan Interactions Protect from Mucosal Infections through Activation of the Inflammasome. *Cell* **167**, 444-456 e414, doi:10.1016/j.cell.2016.08.076 (2016).

Raetz, M. *et al.* Parasite-induced TH1 cells and intestinal dysbiosis cooperate in IFN-gamma-dependent elimination of Paneth cells. *Nat Immunol* **14**, 136-142, doi:10.1038/ni.2508 (2013).

Jokelainen, P. *et al.* Dientamoeba fragilis, a Commensal in Children in Danish Day Care Centers. *J Clin Microbiol* **55**, 1707-1713, doi:10.1128/JCM.00037-17 (2017).

Cuffari, C., Oligny, L. & Seidman, E. G. Dientamoeba fragilis masquerading as allergic colitis. *J Pediatr Gastroenterol Nutr* **26**, 16-20, doi:10.1097/00005176-199801000-00003 (1998).

Yakoob, J. *et al.* Blastocystis hominis and Dientamoeba fragilis in patients fulfilling irritable bowel syndrome criteria. *Parasitol Res* **107**, 679-684, doi:10.1007/s00436-010-1918-7 (2010).

Zhang, N. *et al.* High prevalence of Pentatrichomonas hominis infection in gastrointestinal cancer patients. *Parasit Vectors* **12**, 423, doi:10.1186/s13071-019-3684-4 (2019).

Tuzlak, L. *et al.* Ultrastructural identification and molecular characterization of two new parabasalid species that naturally colonize laboratory mice, Tritrichomonas musculus and Tritrichomonas casperi. *BioRxiv*, doi:https://doi.org/10.1101/2023.01.20.524969 (2023).

Chiaranunt, P. *et al.* NLRP1B and NLRP3 Control the Host Response following Colonization with the Commensal Protist Tritrichomonas musculis. *J Immunol* **208**, 1782-1789, doi:10.4049/jimmunol.2100802 (2022).

Schneider, C. *et al.* A Metabolite-Triggered Tuft Cell-ILC2 Circuit Drives Small Intestinal Remodeling. *Cell* **174**, 271-284 e214, doi:10.1016/j.cell.2018.05.014 (2018).

Escalante, N. K. *et al.* The common mouse protozoa Tritrichomonas muris alters mucosal T cell homeostasis and colitis susceptibility. *The Journal of experimental medicine* **213**, 2841-2850, doi:10.1084/jem.20161776 (2016).

Howitt, M. R. *et al.* Tuft cells, taste-chemosensory cells, orchestrate parasite type 2 immunity in the gut. *Science* **351**, 1329-1333, doi:10.1126/science.aaf1648 (2016).

Wei, Y. *et al.* Commensal Bacteria Impact a Protozoan's Integration into the Murine Gut Microbiota in a Dietary Nutrient-Dependent Manner. *Appl Environ Microbiol* **86**, doi:10.1128/AEM.00303-20 (2020).

Ost, K. S. *et al.* Adaptive immunity induces mutualism between commensal eukaryotes. *Nature* **596**, 114-118, doi:10.1038/s41586-021-03722-w (2021).

Pabst, O., Cerovic, V. & Hornef, M. Secretory IgA in the Coordination of Establishment and Maintenance of the Microbiota. *Trends Immunol* **37**, 287-296, doi:10.1016/j.it.2016.03.002 (2016).

Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome Res* **27**, 722-736, doi:10.1101/gr.215087.116 (2017).

21 Kolmogorov, M., Yuan, J., Lin, Y. & Pevzner, P. A. Assembly of long, error-prone reads using
repeat graphs. *Nat Biotechnol* **37**, 540-546, doi:10.1038/s41587-019-0072-8 (2019).
22 Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv:1303.3997v1 [q-bio.GN] (2013).
23 Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant detection and
genome assembly improvement. *PLoS One* **9**, e112963, doi:10.1371/journal.pone.0112963
(2014).
24 Cantarel, B. L. *et al.* MAKER: an easy-to-use annotation pipeline designed for emerging model
organism genomes. *Genome Res* **18**, 188-196, doi:10.1101/gr.6743907 (2008).
25 Korf, I. Gene finding in novel genomes. *BMC Bioinformatics* **5**, 59, doi:10.1186/1471-2105-5-59
(2004).
26 Stanke, M., Schoffmann, O., Morgenstern, B. & Waack, S. Gene prediction in eukaryotes with a
generalized hidden Markov model that uses hints from external sources. *BMC Bioinformatics* **7**,
62, doi:10.1186/1471-2105-7-62 (2006).
27 Jones, P. *et al.* InterProScan 5: genome-scale protein function classification. *Bioinformatics* **30**,
1236-1240, doi:10.1093/bioinformatics/btu031 (2014).
28 Finn, R. D., Clements, J. & Eddy, S. R. HMMER web server: interactive sequence similarity
searching. *Nucleic Acids Res* **39**, W29-37, doi:10.1093/nar/gkr367 (2011).
29 Nursimulu, N., Moses, A. M. & Parkinson, J. Architect: A tool for aiding the reconstruction of
high-quality metabolic models through improved enzyme annotation. *PLoS Comput Biol* **18**,
e1010452, doi:10.1371/journal.pcbi.1010452 (2022).
30 Malik, S. B., Pightling, A. W., Stefaniak, L. M., Schurko, A. M. & Logsdon, J. M., Jr. An
expanded inventory of conserved meiotic genes provides evidence for sex in Trichomonas
vaginalis. *PLoS One* **3**, e2879, doi:10.1371/journal.pone.0002879 (2007).
31 Garcia, A. F. & Alderete, J. Characterization of the Trichomonas vaginalis surface-associated
AP65 and binding domain interacting with trichomonads and host cells. *BMC Microbiol* **7**, 116,
doi:10.1186/1471-2180-7-116 (2007).
32 Aurrecoechea, C. *et al.* GiardiaDB and TrichDB: integrated genomic resources for the eukaryotic
protist pathogens Giardia lamblia and Trichomonas vaginalis. *Nucleic Acids Res* **37**, D526-530,
doi:10.1093/nar/gkn631 (2009).
33 Benson, D. A. *et al.* GenBank. *Nucleic Acids Res* **42**, D32-37, doi:10.1093/nar/gkt1030 (2014).
34 Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Res **32**, 1792-1797, doi:10.1093/nar/gkh340 (2004).
35 Tamura, K. & Nei, M. Estimation of the number of nucleotide substitutions in the control region
of mitochondrial DNA in humans and chimpanzees. *Mol Biol Evol* **10**, 512-526,
doi:10.1093/oxfordjournals.molbev.a040023 (1993).
36 Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular Evolutionary
Genetics Analysis across Computing Platforms. *Mol Biol Evol* **35**, 1547-1549,
doi:10.1093/molbev/msy096 (2018).
37 Macosko, E. Z. *et al.* Highly Parallel Genome-wide Expression Profiling of Individual Cells
Using Nanoliter Droplets. *Cell* **161**, 1202-1214, doi:10.1016/j.cell.2015.05.002 (2015).
38 Nemesh, J. & Wysoker, A. *Drop-seq tools v1.13*, <https://github.com/broadinstitute/Drop-
seq/releases> (
39 Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21,
doi:10.1093/bioinformatics/bts635 (2013).
40 Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell
transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol* **36**,
411-420, doi:10.1038/nbt.4096 (2018).
41 Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587 e3529,
doi:10.1016/j.cell.2021.04.048 (2021).

42   McInnes, L., Healy, J., Saul, N. & Großberger, L. UMAP: Uniform Manifold Approximation and
Projection for Dimension Reduction. *Journal of Open Source Software* **3**, 861,
doi:https://doi.org/10.21105/joss.00861 (2018).
43   Okuda, S. *et al.* KEGG Atlas mapping for global analysis of metabolic pathways. *Nucleic Acids*
Res **36**, W423-426, doi:10.1093/nar/gkn282 (2008).
44   Alexa, A. & Rahnenfuhrer, J. *topGO: Enrichment Analysis for Gene Ontology. R package version*
2.42.0., <https://github.com/lyijin/topGO_pipeline> (2020).
45   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. *Journal of the Royal statistical society: series B* **57**, 289-300,
doi:https://doi.org/10.1111/j.2517-6161.1995.tb02031.x (1995).
46   Kolde, R. *pheatmap: Pretty Heatmaps. R package version 1.0.12.*, <https://CRAN.R-
project.org/package=pheatmap> (2019).
47   Comeau, A. M., Li, W. K., Tremblay, J. E., Carmack, E. C. & Lovejoy, C. Arctic Ocean
microbial community structure before and after the 2007 record sea ice minimum. *PLoS One* **6**,
e27492, doi:10.1371/journal.pone.0027492 (2011).
48   Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon data.
Nat Methods **13**, 581-583, doi:10.1038/nmeth.3869 (2016).
49   Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data science using
QIIME 2. *Nat Biotechnol* **37**, 852-857, doi:10.1038/s41587-019-0209-9 (2019).
50   Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data processing and
web-based tools. *Nucleic Acids Res* **41**, D590-596, doi:10.1093/nar/gks1219 (2013).
51   McMurdie, P. J. & Holmes, S. phyloseq: an R package for reproducible interactive analysis and
graphics of microbiome census data. *PLoS One* **8**, e61217, doi:10.1371/journal.pone.0061217
(2013).
52   Oksanen, J. B., F. Guillaume; Friendly, Michael; Kindt, Roeland; Legendre, Pierre; McGlinn,
Dan; Minchin, Peter R. ; O'Hara, R. B.; Simpson, Gavin L.; Solymos, Peter; Henry M.; Stevens,
H.; Szoecs, Eduard; Wagner, Helene. *vegan: Community Ecology Package*, <https://CRAN.R-
project.org/package=vegan> (2017).
53   Bates, D., Maechler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using
lme4. *Journal of Statistical Software* **67**, 1-48, doi:10.18637/jss.v067.i01 (2015).
54   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**, 550, doi:10.1186/s13059-014-0550-8 (2014).
55   Pantano, L. *DEGreport: Report of DEG analysis. R package version 1.26.0*,
<http://lpantano.github.io/DEGreport/> (2020).
56   Taj, B. *et al.* MetaPro: A scalable and reproducible data processing and analysis pipeline for
metatranscriptomic investigation of microbial communities. *BioRxiv*,
doi:https://doi.org/10.1101/2021.02.23.432558 (2021).
57   Garber, A. I. *et al.* FeGenie: A Comprehensive Tool for the Identification of Iron Genes and Iron
Gene Neighborhoods in Genome and Metagenome Assemblies. *Front Microbiol* **11**, 37,
doi:10.3389/fmicb.2020.00037 (2020).
58   Peregrin-Alvarez, J. M., Xiong, X., Su, C. & Parkinson, J. The Modular Organization of Protein
Interactions in Escherichia coli. *PLoS Comput Biol* **5**, e1000523,
doi:10.1371/journal.pcbi.1000523 (2009).
59   Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using DIAMOND. *Nat*
Methods **12**, 59-60, doi:10.1038/nmeth.3176 (2015).
60   Hayat, M. A. *Principles and Techniques of Electron Microscopy*. (Hodder Arnold H& S, 1981).
61   Polk, D. B. & Peek, R. M., Jr. Helicobacter pylori: gastric cancer and beyond. *Nat Rev Cancer*
10, 403-414, doi:10.1038/nrc2857 (2010).
62   Partridge, J. D., Scott, C., Tang, Y., Poole, R. K. & Green, J. Escherichia coli transcriptome
dynamics during the transition from anaerobic to aerobic conditions. *J Biol Chem* **281**, 27806-
27815, doi:10.1074/jbc.M603450200 (2006).

63 Noel, C. J. *et al.* Trichomonas vaginalis vast BspA-like gene family: evidence for functional
diversity from structural organisation and transcriptomics. *BMC Genomics* **11**, 99,
doi:10.1186/1471-2164-11-99 (2010).
64 Wilson, I. W. *et al.* Genetic Diversity and Gene Family Expansions in Members of the Genus
Entamoeba. *Genome Biol Evol* **11**, 688-705, doi:10.1093/gbe/evz009 (2019).
65 Guzman-Tellez, P. *et al.* Lectins as virulence factors in Entamoeba histolytica and free-living
amoebae. *Future Microbiol* **15**, 919-936, doi:10.2217/fmb-2019-0275 (2020).
66 Siqueira-Neto, J. L. *et al.* Cysteine proteases in protozoan parasites. *PLoS Negl Trop Dis* **12**,
e0006512, doi:10.1371/journal.pntd.0006512 (2018).
67 Hapfelmeier, S. *et al.* Reversible microbial colonization of germ-free mice reveals the dynamics
of IgA immune responses. *Science* **328**, 1705-1709, doi:10.1126/science.1188454 (2010).
68 Hernandez, Y. *et al.* Novel role of sphingolipid synthesis genes in regulating giardial encystation.
Infect Immun **76**, 2939-2949, doi:10.1128/IAI.00116-08 (2008).
69 Jeelani, G. *et al.* Metabolic profiling of the protozoan parasite Entamoeba invadens revealed
activation of unpredicted pathway during encystation. *PLoS One* **7**, e37740,
doi:10.1371/journal.pone.0037740 (2012).
70 Pereira-Neves, A. & Benchimol, M. Tritrichomonas foetus: budding from multinucleated
pseudocysts. *Protist* **160**, 536-551, doi:10.1016/j.protis.2009.05.001 (2009).
71 Beri, D. *et al.* Demonstration and Characterization of Cyst-Like Structures in the Life Cycle of
Trichomonas vaginalis. *Front Cell Infect Microbiol* **9**, 430, doi:10.3389/fcimb.2019.00430
(2019).
72 Horlock-Roberts, K. *et al.* Drug-Free Approach To Study the Unusual Cell Cycle of Giardia
intestinalis. *mSphere* **2**, doi:10.1128/mSphere.00384-16 (2017).
73 Hirt, R. P. *et al.* Trichomonas vaginalis pathobiology new insights from the genome sequence.
Adv Parasitol **77**, 87-140, doi:10.1016/B978-0-12-391429-3.00006-X (2011).
74 Brennand, A. *et al.* Autophagy in parasitic protists: unique features and drug targets. *Mol*
Biochem Parasitol **177**, 83-99, doi:10.1016/j.molbiopara.2011.02.003 (2011).
75 Arroyo-Begovich, A., Carabez-Trejo, A. & Ruiz-Herrera, J. Identification of the structural
component in the cyst wall of Entamoeba invadens. *J Parasitol* **66**, 735-741 (1980).
76 Ward, H. D., Alroy, J., Lev, B. I., Keusch, G. T. & Pereira, M. E. Identification of chitin as a
structural component of Giardia cysts. *Infect Immun* **49**, 629-634, doi:10.1128/iai.49.3.629-
634.1985 (1985).
77 Castro, C. *et al.* Iron-modulated pseudocyst formation in Tritrichomonas foetus. *Parasitology*
143, 1034-1042, doi:10.1017/S0031182016000573 (2016).
78 Lehker, M. W. & Alderete, J. F. Iron regulates growth of Trichomonas vaginalis and the
expression of immunogenic trichomonad proteins. *Mol Microbiol* **6**, 123-132, doi:10.1111/j.1365-
2958.1992.tb00844.x (1992).
79 Vanacova, S. *et al.* Iron-induced changes in pyruvate metabolism of Tritrichomonas foetus and
involvement of iron in expression of hydrogenosomal proteins. *Microbiology (Reading)* **147**, 53-
62, doi:10.1099/00221287-147-1-53 (2001).
80 Horvathova, L. *et al.* Transcriptomic identification of iron-regulated and iron-independent gene
copies within the heavily duplicated Trichomonas vaginalis genome. *Genome Biol Evol* **4**, 1017-
1029, doi:10.1093/gbe/evs078 (2012).
81 Novak, L. *et al.* Arginine deiminase pathway enzymes: evolutionary history in metamonads and
other eukaryotes. *BMC Evol Biol* **16**, 197, doi:10.1186/s12862-016-0771-4 (2016).
82 Stadelmann, B., Merino, M. C., Persson, L. & Svard, S. G. Arginine consumption by the
intestinal parasite Giardia intestinalis reduces proliferation of intestinal epithelial cells. *PLoS One*
7, e45325, doi:10.1371/journal.pone.0045325 (2012).
83 Chappell, C. L. *et al.* Fecal Indole as a Biomarker of Susceptibility to Cryptosporidium Infection.
Infect Immun **84**, 2299-2306, doi:10.1128/IAI.00336-16 (2016).

84 Yilmaz, B. *et al.* Gut microbiota elicits a protective immune response against malaria
transmission. *Cell* **159**, 1277-1289, doi:10.1016/j.cell.2014.10.053 (2014).
85 Bierlein, M., Hedgespeth, B. A., Azcarate-Peril, M. A., Stauffer, S. H. & Gookin, J. L. Dysbiosis
of fecal microbiota in cats with naturally occurring and experimentally induced Tritrichomonas
foetus infection. *PLoS One* **16**, e0246957, doi:10.1371/journal.pone.0246957 (2021).
86 Wang, J., Chen, W. D. & Wang, Y. D. The Relationship Between Gut Microbiota and
Inflammatory Diseases: The Role of Macrophages. *Front Microbiol* **11**, 1065,
doi:10.3389/fmicb.2020.01065 (2020).
87 Cao, E. *et al.* The protozoan commensal Tritrichomonas musculis is a natural adjuvant for
mucosal IgA. *BioRxiv*, doi:https://doi.org/10.1101/2022.10.08.511442 (2022).
88 Catanzaro, J. R. *et al.* IgA-deficient humans exhibit gut microbiota dysbiosis despite secretion of
compensatory IgM. *Sci Rep* **9**, 13574, doi:10.1038/s41598-019-49923-2 (2019).
89 Mundodi, V., Kucknoor, A. S., Chang, T. H. & Alderete, J. F. A novel surface protein of
Trichomonas vaginalis is regulated independently by low iron and contact with vaginal epithelial
cells. *BMC Microbiol* **6**, 6, doi:10.1186/1471-2180-6-6 (2006).
90 Kusdian, G., Woehle, C., Martin, W. F. & Gould, S. B. The actin-based machinery of
Trichomonas vaginalis mediates flagellate-amoeboid transition and migration across host tissue.
Cell Microbiol **15**, 1707-1721, doi:10.1111/cmi.12144 (2013).
91 Petropolis, D. B., Fernandes Rodrigues, J. C., da Rocha-Azevedo, B. & Silva-Filho, F. C. E. The
binding of Tritrichomonas foetus to immobilized laminin-1 and its role in the cytotoxicity exerted
by the parasite. *Microbiology (Reading)* **154**, 2283-2290, doi:10.1099/mic.0.2007/015941-0
(2008).
92 Dias-Lopes, G. *et al.* Morphologic study of the effect of iron on pseudocyst formation in
Trichomonas vaginalis and its interaction with human epithelial cells. *Mem Inst Oswaldo Cruz*
112, 664-673, doi:10.1590/0074-02760170032 (2017).
93 Lehker, M. W. & Sweeney, D. Trichomonad invasion of the mucous layer requires adhesins,
mucinases, and motility. *Sex Transm Infect* **75**, 231-238, doi:10.1136/sti.75.4.231 (1999).
94 Alsmark, C. *et al.* Patterns of prokaryotic lateral gene transfers affecting parasitic microbial
eukaryotes. *Genome Biol* **14**, R19, doi:10.1186/gb-2013-14-2-r19 (2013).
95 Johansson, M. E. *et al.* Normalization of Host Intestinal Mucus Layers Requires Long-Term
Microbial Colonization. *Cell Host Microbe* **18**, 582-592, doi:10.1016/j.chom.2015.10.007 (2015).
96 Bradic, M. *et al.* Genetic Indicators of Drug Resistance in the Highly Repetitive Genome of
Trichomonas vaginalis. *Genome Biol Evol* **9**, 1658-1672, doi:10.1093/gbe/evx110 (2017).
97 Drmota, T. & Kral, J. Karyotype of Trichomonas vaginalis. *European Journal of Protistology* **33**,
131-135 (1997).
98 Bradic, M. & Carlton, J. M. Does the common sexually transmitted parasite Trichomonas
vaginalis have sex? *PLoS Pathog* **14**, e1006831, doi:10.1371/journal.ppat.1006831 (2018).
99 Ramesh, M. A., Malik, S. B. & Logsdon, J. M., Jr. A phylogenomic inventory of meiotic genes;
evidence for sex in Giardia and an early eukaryotic origin of meiosis. *Current biology : CB* **15**,
185-191, doi:10.1016/j.cub.2005.01.003 (2005).
100 El Aidy, S. *et al.* Temporal and spatial interplay of microbiota and intestinal mucosa drive
establishment of immune homeostasis in conventionalized mice. *Mucosal Immunol* **5**, 567-579,
doi:10.1038/mi.2012.32 (2012).
101 Alderete, J. F., Nguyen, J., Mundodi, V. & Lehker, M. W. Heme-iron increases levels of AP65-
mediated adherence by Trichomonas vaginalis. *Microb Pathog* **36**, 263-271,
doi:10.1016/j.micpath.2003.12.007 (2004).
102 Gastelum-Martinez, A. *et al.* Iron-modulated virulence factors of Entamoeba histolytica. *Future*
Microbiol **13**, 1329-1341, doi:10.2217/fmb-2018-0066 (2018).
103 Lloyd, D., Lauritsen, F. R. & Degn, H. The parasitic flagellates Trichomonas vaginalis and
Tritrichomonas foetus produce indole and dimethyl disulphide: direct characterization by

membrane inlet tandem mass spectrometry. *J Gen Microbiol* **137**, 1743-1747,
doi:10.1099/00221287-137-7-1743 (1991).
104   Li, X., Zhang, B., Hu, Y. & Zhao, Y. New Insights Into Gut-Bacteria-Derived Indole and Its
Derivatives in Intestinal and Liver Diseases. *Front Pharmacol* **12**, 769501,
doi:10.3389/fphar.2021.769501 (2021).
105   Zhang, L. S. & Davies, S. S. Microbial metabolism of dietary components to bioactive
metabolites: opportunities for new therapeutic interventions. *Genome Med* **8**, 46,
doi:10.1186/s13073-016-0296-x (2016).
106   Chen, L. M. *et al.* Tryptophan-kynurenine metabolism: a link between the gut and brain for
depression in inflammatory bowel disease. *J Neuroinflammation* **18**, 135, doi:10.1186/s12974-
021-02175-2 (2021).
107   Ortega, M. A. *et al.* Microbiota-gut-brain axis mechanisms in the complex network of bipolar
disorders: potential clinical implications and translational opportunities. *Mol Psychiatry*,
doi:10.1038/s41380-023-01964-w (2023).

893
894

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6