Climate Change Factors Interact to Shift Functional Microbial Composition in Peatlands

Christopher L. Kilner[1,✉], Alyssa A. Carrell[2], Daniel J. Wieczynski[1], Samantha Votzke[1], Andrea Yammine[1], Katrina DeWitt[1], Dale A. Pelletier[2], David J. Weston[2], and Jean P. Gibert[1]

[1]Department of Biology, Duke University, Durham, NC 27708
[2]Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, TN, 37830

Microbes play a major role in the global carbon cycle that fuels climate change. But how microbes may in turn respond to climate change remains poorly understood. Here, we collect data from a long-term whole-ecosystem warming experiment at a boreal peatland to address how temperature and CO_2 jointly influence protist communities — i.e., abundant and diverse, but poorly understood, microbial Eukaryotes. Protists influence ecosystem function directly through photosynthesis and respiration, and indirectly through predation on decomposers (bacteria, archaea, and fungi). Using a combination of high-throughput fluid imaging and 18S amplicon sequencing, we report climate-induced, community-wide shifts in protist community functional composition (e.g., size, shape, and metabolism) that could alter the overall function of peatland ecosystems. We also show that these responses to warming and elevated CO_2 are the result of taxonomic turnover. Surprisingly, our results clearly show strong interactive effects between temperature and CO_2, such that the effects of warming on functional composition are generally reversed by elevated CO_2. These findings show how the interactive effects of warming and rising CO_2 could alter the structure and function of peatland microbial food webs — a fragile ecosystem that stores 25% of terrestrial carbon and is increasingly threatened by human exploitation.

climate change | microbial diversity | community ecology | functional traits
Correspondence: *kilnerc@tcd.ie*

Introduction

Greenhouse gas emissions set the pace of climate change (1) and most ecosystems on Earth influence this process by serving as carbon (C) sources or sinks (2). Overall, terrestrial ecosystems mitigate $\approx$30 % of anthropogenic CO_2 emissions (3, 4). The predominant terrestrial carbon sink — peatlands — store 100% more carbon than all forests while covering less than 3% of the globe (5–8). *Sphagnum* mosses dominate peatlands (9) and are responsible for most C sequestration in the form of recalcitrant peat (10). They harbor diverse communities of single-celled (*i.e.,* bacteria, archaea, protists) and multi-celled (*i.e.,* fungi, metazoa) organisms that influence C cycling directly (11, 12), through respiration of $\approx$100 Pg $\cdot$ C $\cdot$ yr^{-1} in soils alone (13, 14), and indirectly, by supplying nitrogen needed for *Sphagnum* moss growth (15–17), converting methane into CO_2 necessary for photosynthesis (18–20), providing stress tolerance to the host mosses (21, 22), and assisting in host-pathogen defense (23).

How ecological interactions with other members of the moss-associated community modulate the effects of these microbes on peatland C cycling (24–26), is poorly understood. Indeed, predation by metazoans plays a prominent role in peatland C cycling (27), so it is likely to also matter within microbial communities. Unicellular Eukaryotes collectively known as protists store 2$\times$ more C globally than all animals combined (28, 29), contribute to C cycling through mixotrophic metabolism (30), and also serve as one of the principal biological controls on microbial respiration and photosynthesis through predation and competition (31–33). Protists also directly influence plant growth and health through their effects on the rhizosphere (31, 34, 35) and phyllosphere (36, 37) — thus indirectly modulating plant growth through their effects on beneficial microorganisms (38, 39).

But how protist communities and associated function may respond to climate change in peatlands, is not known. Yet, simple protist functional traits — *i.e.*, characteristics controlling how organisms respond to the environment or affect ecosystem properties (40) — like cell size, cell shape, and cellular contents, can be used to predict warming effects on protist communities and ecosystem-level function because of underlying links to metabolism and respiration rates(41). By quantifying changes in functional traits of these protist communities, in addition to changes in abundances, biomass and composition, we can infer changes in their ecological function and their impacts in peatland C cycling. For instance, metabolic demand and total respiration rates increase with increasing temperature, favoring smaller organisms (42, 43) with a shape (e.g., long and slim) that maximizes energy intake and minimizes energetic costs (44, 45). Increased metabolic demands can impact resource acquisition strategies (i.e., increased heterotrophy over autotrophy) (46–49), resulting in measurable changes in cellular optical properties — e.g., red/green color ratio or the heterogeneity of gray-scale values, which have been linked to cellular contents and cell viability (50, 51). Consequently, changes in functional composition could alter net C fluxes and other elements within peatlands (52–55).

Here, we address how peatland protist communities change in novel climates by quantifying shifts in abundance, biomass, taxonomic composition, and functional trait composition, in long-term whole-ecosystem temperature and CO_2 manipulations of living moss and superficial peat (Fig. 1A) at the SPRUCE long-term whole-ecosystem warming experimental site (56). We quantified protist abundance, biomass,

Fig. 1. (A) Experimental approach and design; samples were collected in a 2-factorial mesocosm design in boreal peatlands with variation in temperature treatment (+0, +2.25, +4.5, +6.75, +9℃) and CO_2 (+0 PPM, +450 PPM) concentrations. Protist samples were collected from peat within each treatment and processed *ex situ*. (B) After incubation (for 24 hours in climate controlled growth chambers, Carolina protist liquid media, 22℃ and 65% humidity), samples were analyzed via fluid imaging to quantify abundances across treatments and physical and optical functional traits. (C) Functional composition data hypothesized on first-principles of metabolic and ecological theory. We anticipate decreases in size (e.g., volume), shape (e.g., aspect ratio), and cellular contents (e.g., sigma intensity) under warming, with an increase in red/green ratio (e.g., photosynthesis). Under elevated CO_2 we expect larger cell volumes, more roundedness, higher cellular activity, and lower red/green ratios.

and community functional traits through fluid imaging and used 18S rRNA gene sequencing to infer mechanisms of change in observed protist community functional trait composition (*i.e.*, taxonomic shifts or phenotypic plasticity). We test the hypotheses that increasing temperatures lead to: (1) a decrease in protists' body size — i.e., the temperature-size rule (42, 44), (2) an increase in surface-to-volume ratio (aspect ratio), as protists optimize shape to increase energy uptake (surface) and reduce metabolic costs (volume) (57), (3) an increase in cellular optical heterogeneity (or cellular "contents"), as temperature-induced increased metabolic rates requires more cellular machinery (58), and (4) increased heterotrophy (*i.e.*, increased red/green ratios) to keep up with metabolic demands. Additionally, we expect elevated CO_2 to result in an increase in total community biomass (59, 60) as more CO_2 fuels photosynthesis at the base of the food web.

This increase in energy availability at the base of the microbial food web should also result in (Fig. 1C): 1) an increase

in protist volume in response to more energy availability, 2) an increase in aspect-ratio as increased available energy releases cells from shape metabolic constraints, 3) an increase in sigma-intensity, due to increased production of cellular machinery to process available energy, and, 4) an increase in autotrophy (i.e., decrease in the red/green ratio of cells) relative to heterotrophy (42, 61, 62). We show that warming and CO_2 have significant, negative interactive effects, on the taxonomic and functional composition of peatland protist communities, which has important consequences for C cycling in these systems.

Materials and Methods

A. Study Site and Field Sampling. Our study site is situated within the S1-Bog of the Marcell Experimental Forest (47° 30.4760′ N, 93° 27.1620′ W; 418 m above mean sea level), MN, USA, and sits on the southern edge of the temperate boreal forest range in North America, with a sub-humid continental climate. The S1-Bog — above a water

table with marginal groundwater influence — consists of an ombrotrophic peatland dominated by a forest canopy of black spruce (*Picea mariana*) and tamarack (*Larix laricinia*). Below the canopy, the site contains a mixture of ericaceous scrubs, herbaceous sedges and grasses, and a bryophyte layer dominated by *Sphagnum* spp. mosses. Mosses are distributed in hollows (*S. angustifolium* & *S. fallax*) and in drier hummocks (*S. divinum* previously called *S. magellanicum*). Other bryophytes include feather mosses (*Pleurozium* spp.) and haircap mosses (*Polytrichum* spp.). For further description of the site and below-ground peat profile, see (63) and (64).

We collected superficial peat (1-5 cms) from whole-ecosystem experimental manipulations of black spruce (*Picea mariana*) dominated peatlands at the SPRUCE experiment. The *P. mariana*–*Sphagnum* spp. raised bog experimental ecosystem contains open-top chamber systems with a factorial-based design of ambient +0 ppm and +450 ppm CO_2 treatments across ambient (+0, +2.25, +4.5, +6.75, +9°C soil (initiated June 2014) and air (initiated June 2015) warming treatments (Fig. 1; (63)). Peat and living plant samples collected from two sub-plot locations across ten 12 meter diameter chambers were placed in individual sterile bags and over-night shipped to Durham, NC, for imaging of protist communities while additional samples were flash-frozen and over-night shipped to Oak Ridge, TN, for DNA extraction and 18S amplicon sequencing. Upon reception of samples for imaging, we removed the material from sterile bags and transferred the samples to Carolina protist media (Liquid Protozoan Nutrient Concentrate; 13-2350) through glass funnels. We collected the discharge from the funnels in 250 ml glass jars and kept the cultures at 22°C and 65% humidity for 24 hours before fluid imaging.

B. Density and Biomass. Protist — operationally defined in this study as all non-fungal unicellular Eukaryotes — densities were estimated as simple counts ($\hat{E}$) of the number of imaged particles per volume analyzed. This estimate accounts for the FlowCam efficiency, which screens roughly 75% of the sampled volume. Biomass was estimated by using efficiency-corrected measurements of total protist volume (see supplementary materials).

C. Trait Measurements. We quantified all functional traits (Table S1) through fluid imaging with a FlowCAM (Yokagawa Fluid Imaging, USA) of 6 mL water samples from incubated jars at $10\times$ magnification. We manually curated the ensuing cell images through the Visual Spreadsheet software program (Yokagawa Fluid Imaging, USA), by discarding all non-protist images and retaining only high quality protist images for trait quantification (n = 211,039, all protist images, regardless of quality, were kept for abundance estimations). To test our hypotheses about size, shape, cellular contents, and energy acquisition, we selected four traits (volume, aspect ratio, sigma intensity, red/green ratio) representing these functional composition categories from a larger set of 37 physical and optical functional traits of possible proxies by

eliminating highly correlated and co-linear traits (see supplementary materials; Table S1, Fig. S1A).

D. Analyses. We tested the effects of temperature and CO_2 levels on the traits of interest with generalized linear models (GLMs) as well as the abundance and biomass of the protist communities with generalized additive models (GAMs). To test hypotheses 1-4, we first clustered individual cellular observations into size classes based upon the natural log of protists' geodesic length (e.g., longest axis) using Gaussian finite mixture modelling with the the R package *mclust* (65). We chose length as it was less correlated with the traits of interest and would reduce co-linearity in our models between the size class cluster and the functional traits (Fig. S2). We selected an optimal set of 5 size classes with a Gaussian mixture model and Bayesian Information Criterion (BIC) (Fig. S3). We then ran GLMs for each functional trait against size class (factor), temperature (continuous), and carbon dioxide level (factor). Associated Pearson correlation coefficients and p-values were obtained via the *ggpubr* R package (66) and reported for each size class (Fig. S4). Additionally, we examined changes across environments in the estimated abundance and biomass (grams) of the community per size class with GAMs (k = 5) through the R package *mgcv* (67), reporting associated Pearson correlations and p-values for each size class (Fig. S5).

E. Amplicon Sequencing. To understand the possible mechanisms behind observed changes in abundance, biomass and functional traits, we extracted genomic DNA from our samples with the DNeasy 96 Plant Kit (Qiagen) and amplified the V4 region of the 18S rRNA gene following (68). Samples were multiplexed, pooled and sequenced on an Illumina MiSeq instrument with paired end kit (2 x 251) at Oak Ridge National Laboratory. We processed protist sequences with QIIME 2 v 2021.2 platform. Paired sequences were demultiplexed with the plugin demux and quality filtered (denoised, dereplicated, chimera filtered and pair-end merged) and processed in Sequence Variants (SVs) with the dada2 plugin (69). Taxonomy was assigned using the Pr2 (18S) database (70) and sequences assigned as "bacteria", Gregarinomorphea, Bacillariophyta, or Embryophyceae were removed (as these are not considered true protists phyla). We used sequence variants to calculate alpha diversity as counts and the Shannon diversity index, and calculated beta diversity as a Bray-Curtis distance with the phyloseq package (71).

Results

Experimental manipulations of temperature and CO_2 levels induced a considerable (80-fold) change in total observed protist abundance (F = 2.2554; p = 0.02244, Fig. S5A) and biomass (F = 2.2504; p = 0.02273, Fig. S5B) across treatments (Fig. 2A), as well as a shift in the relative dominance of size classes (Fig. 2B). Across all body size classes, protist abundance and biomass increased with temperature under ambient CO_2 (Fig. 2A). This trend reverted under elevated CO_2, with major declines in biomass and abundance

Fig. 2. Estimated protist abundances. (A) Estimated total abundance data on the natural log scale for size classes across temperatures under ambient and elevated CO_2 treatments. Data indicates that abundances increase across size classes with temperature under ambient CO_2 levels but this trend is reversed under elevated CO2 levels. (B) Relative abundance of size classes by CO_2 treatment calculated from total estimated abundance (A) across temperatures. Results show that different size classes respond differentially to environmental treatments.

across size classes (Fig. 2A). The proportion of abundance within size classes remained relatively consistent with increasing temperature under ambient CO_2 conditions; however, under elevated CO_2 conditions, the relative proportion of larger protists increased, at the expense of the two smallest size classes (Fig. 2B).

A three-way interaction between temperature, CO_2 concentration, and size class drove significant changes in the functional trait composition of protist communities (Fig. 3): size (F = 4,007; $p \ll 2.2x10^{-16}$); shape (F = 13,204; $p \ll 2.2 \times 10^{-16}$); cellular contents (F = 789.05; p $\ll 2.2x10^{-16}$); and energy acquisition (F = 657.15; p $\ll 2.2x10^{-16}$). The magnitude of change was driven by the CO_2 treatment, with elevated CO_2 rates of change outpacing that of ambient CO_2 by up to a factor of 25 (Fig. 3). Under ambient CO_2, protists got smaller (Figs. 3A, S6), less round (Figs. 3B, S7), showed more cellular contents (Figs. 3C, S8), and became redder (*i.e.*, more heterotrophic) at lower levels of warming but greener (*i.e.*, more autotrophic) at higher levels of warming (Fig. 3D, Fig. S9). Conversely, protists under elevated CO_2 got larger (Fig. 3A, S6), rounder (Fig. 3B, S7), had fewer cellular contents (Fig. 3C, S8), and got redder (Fig. 3D, S9). Thus, with the exception of the two smallest size classes for red/green ratio, protists displayed inverse patterns for all traits in response to increasing temperature under ambient and elevated CO_2 levels.

We also find that some of these functional responses are at least in part mediated by protist size, as protist volumes' rate of change with warming was strongly size-dependent (Fig. S10A, (0.014 < p < 0.066), c.f. (43)): larger size classes responded more strongly to temperature than smaller ones. While the rates of change in all traits display interesting non-

linear size-dependency, we found weaker support for allometry in shape, cellular contents, and energy acquisition (Fig. S10B-D).

Importantly, observed changes in community functional trait values could be the results of plastic change across all or most protists in the community, or the result of compositional shifts that led to communities whose functional traits differed from the original ones. Amplicon sequencing results show that important taxonomic shifts (Fig. 5A) underlie the observed changes in functional traits (Fig. 4). Indeed, increasing temperature leads to a decrease in the diversity of microbial communities (Fig. 5B,C) — a finding consistent with multiple studies across biomes that find climate change decreases diversity of species across scales (72–74). Curiously, temperature interacts with CO_2 such that the effects of CO_2 treatment decrease at warmer temperatures, leading to a near convergence in the taxonomic compositional similarity between ambient and elevated CO_2 as Bray-Curtis dissimilarity decreases rapidly (Fig. 5D).

Discussion

Understanding how moss-associated microbial communities change with temperature and CO_2 concentrations is vital for predicting how peatland ecosystems will respond to climate change (7, 75, 76) and alter the global C cycle (77–79). Here, we find strong evidence of changes in the abundance, biomass, functional traits, and composition of important but poorly understood members of peatland microbial communities that are driven by an interaction between warming and CO_2 concentration (Figs. 2-4). We show that peatland protist communities are sensitive to climate change, and under ambient CO_2 match our expectations based upon

    Kilner *et al.* | Climate Change Factors Interact to Shift Functional Microbial Composition in Peatlands

Fig. 3. Rate of change across temperatures (β) for functional traits by protist size class and CO_2 treatments (lollipops). Slopes were calculated from GLMs. Mean slope across size classes displayed as dashed line. Positive slope values indicate: (A, Volume) increasing size; (B, Aspect Ratio) higher symmetry and roundedness (shape); (C, Sigma Intensity) higher optical heterogeneity; and (D, Red/Green ratio) increasing redness and/or decreasing greenness.

metabolic theory — getting smaller, less round, more active, and more heterotrophic. However, these trends are reversed by elevated CO_2 concentrations (Fig. 3), and that the magnitude and direction of these changes are driven by nonlinear size-dependence (Fig. S10). Last, while protist communities converge in taxonomic composition between ambient and elevated CO2 levels as temperature rises, they diverge in functional trait composition, suggesting important but yet unknown effects on peatland function with environmental change.

Protist reductions in volume (Figs. 3A, S6) and roundedness (Fig. 3B, Fig. S7) observed at SPRUCE under ambient CO_2 and elevated temperatures can be easily predicted from metabolic theory. This is important because it suggests that we may be able to anticipate complex temperature and CO_2 responses in natural and intricate microbial communities based on first principles. For example, temperature selects for smaller organisms and/or organisms with larger surface-to-volume ratio, who can meet higher metabolic demands while conserving resources as metabolism costs increase at higher temperatures (80). This negative relationship between size and temperature —the "temperature-size rule" (42, 44)—is well documented for both single-celled (42) and multi-celled organisms (81–83), but was so far thought to be

a plastic response.

We show that the temperature-size rule can result from taxonomic shifts that selects for smaller organisms (Fig 3). Similarly, shape shifts (e.g., a decrease in roundedness with temperature) likely results from selection to conserve energy by simultaneously reducing volume (which increases metabolic costs), and increasing surface (which increases resource acquisition) (84). Changes in protist shape and size have multiple ecological consequences, including population dynamics and ecological interactions within communities (33, 41, 85), as well as total respiration, which decreases with volume

(86). Shifts in size and shape can occur quite rapidly and can be as dramatic as 50% or more (61, 87). Likely, species that cannot rapidly respond to temperatures are thus replaced by those that can, leading to high species turnover (Fig. 4).

Based on lack of evidence in the literature, we initially assumed that warming-induced patterns on non-aquatic protists in cell shape and contents would not differ between CO_2 treatments. However, shocking trend reversals between CO_2 treatments (Figs. 3, 4, S6-S9) suggest that joint temperature and CO_2 impacts may be much more complex than previously thought. Despite our observations, the current ecolog-

Fig. 4. Change in (A) protist order composition and (B, C) species diversity of protist community across temperatures and CO_2 treatments. (A) Protists' order composition shifts both with temperature increase and within CO_2 treatment, with a decrease in (B) total species composition and (C) species diversity at higher temperatures. (D) Species composition becomes more homogeneous—lower Bray-Curtis distance—between CO_2 treatments at higher temperatures, indicating an interactive effect (pearson's correlation = -0.72).

Fig. 5. Combined normalized change in biomass and resource acquisition across quadrant (I-IV) outcomes. Under elevated CO_2, the increase in respiration is offset by a loss in biomass of these organisms (IV). Given normalization of difference, slopes $\leq$ -1 indicate a net reduction in heterotrophy. Under ambient CO_2 conditions, an increase in net photosynthesis occurs for the two largest size classes (II), as both biomass and green reflectance increase. The net carbon dynamics in these plots may be offset by the normalized relative increase in biomass of the more heterotrophic smaller size classes (I).

ical paradigm posits that increased volume (Fig. 3A), decreased surface-to-volume ratio (Fig. 3B) and decreased cellular contents (Fig. 3C) should be disfavored under higher temperatures (88), but this only happens under ambient CO_2 conditions (Fig. 3). While interactive effects of CO_2 and temperature have also been observed in aquatic phytoplankton systems (59, 60, 89), these systems displayed a shift in magnitude of traits, but not a change in the sign of the relationships, as observed here, showing the importance of addressing temperature impacts on microbial organisms in the light of simultaneous CO_2 increases.

These trend reversal may be due to CO_2-induced pH changes of the peat and/or the stoichiometry of diffusion under higher temperatures and elevated CO_2. Experiments on ocean-dwelling phytoplankton have found that the interaction between warming temperatures and elevated CO_2 can induce a more rapid decrease in community biomass and chlorophyll-a concentration (90) and that elevated CO_2 may lessen the demand for a higher surface-to-volume ratio as diffusion becomes easier across the cellular membrane at higher concentrations of dissolved CO_2. These changes may also be due to the increased efficiency of heterotrophic resource acquisition compared to photosynthesis (91); the shift towards heterotrophic resource acquisition (Fig. 3D) under elevated CO_2 conditions may lessen the need for energy conservation. The observed trait trends may also be an indirect effect of elevated CO_2, as increased carbon dioxide levels may alter *Sphagnum* functions as well as nutrient cycling and mineralization (92)—especially of N (93). Elevated CO_2 increases the exudation of soluble organic compounds (94)—especially sugars—that contribute to the structure of the microbial community, often increasing the abundance of bacteria and heterotrophic protists (95). With higher abundances of bacteria, protists may obtain more energy through predation. Alternatively, elevated CO_2 may induce the *Sphagnum* and other plants in the peatland system to release more nitrogen either

directly through increases in N-releasing enzymes or through symbiosis with mycorrhizal fungi (96, 97).

Interestingly, the trend reversal in traits, biomass, and abundance induced by elevated CO_2 are size-dependent(Fig. S10). We find strong support for a super-linear scaling relationship between body size and volume that indicates protist populations might undergo density-dependent (population) and density-independent (individual phenotypic) responses to climate change (41, 61, 87, 98, 99). Our results indicate a significant relationship between proportional change of traits and size—larger protists' size (Fig. S10A) and shape (Fig. S10B) change more than smaller protists per unit volume (Figs. S10A, S10B) while cellular contents follow a sub-linear size-dependency (Fig. S10C). Because smaller protist phenotypes shift much less and these smaller individuals dominate the relative abundance of the system (Figs. 2B, S3B), there may be embedded resistance to phenotypic change in these communities overall. This trend portends potential eco-evolutionary responses of protists' under environmental stressors as phenotypic traits and plasticity may drive the success and survival of individuals responding to changing metabolic demands (100, 101).

Conclusion

Our findings suggest important ways in which climate change may be influencing C cycling on peatland ecosystems. Given protists' various energy acquisition strategies, changes in the balance of autotrophy versus heterotrophy can have cascading effects on the biological carbon cycle (102–105). Due to their distribution predominantly within the mid- and high-latitudes of the Northern hemisphere, peatlands are especially sensitive to climate change (106–109). The increase in the red/green ratio of 70% of size classes in our study (Figs. 6, S9) may be alarming; a decrease in the red/green ratio is associated with decreased autotrophy and, therefore, car-

bon uptake (110). However, when we normalize the β coefficents of our models, the overall increase in heterotrophy is outweighed by a decrease in biomass of heterotrophic organisms and an increase in biomass of putatively photosynthetic organisms, likely indicating a net reduction in heterotrophy (Figs. 5, S9). These tantalizing data-driven hypotheses remain untested, but are suggestive of important ways in which environmental change both influences and is influenced by rapid microbial responses in functional trait composition in globally important—and highly threatened—ecosystems like peatlands.

AUTHOR CONTRIBUTIONS

CLK and JPG conceptualized the study. AC, DAP and DJW collected SPRUCE samples. AAC prepared and provided amplicon sequencing data. JPG, SV, and AY cultured specimens and collected protists traits with FlowCam. CLK performed all analyses with support from DJW, AC, and JPG. CLK wrote the first draft and all authors contributed substantially to subsequent versions.

FUNDING

This work was supported by a US Department of Energy, Office of Science, Office of Biological and Environmental Research, Genomic Science Program Grant to J.P.G., under Award DE-SC0020362.

ACKNOWLEDGEMENTS

We would like to thank Dawn M. Klingeman from Oak Ridge National Laboratory for running the 18S rRNA samples on the sequencer.

Bibliography

1. Will Steffen, Johan Rockström, Katherine Richardson, Timothy M Lenton, Carl Folke, Diana Liverman, Colin P Summerhayes, Anthony D Barnosky, Sarah E Cornell, Michel Crucifix, et al. Trajectories of the earth system in the anthropocene. *Proceedings of the National Academy of Sciences*, 115(33):8252–8259, 2018.

2. Benjamin D Stocker, Raphael Roth, Fortunat Joos, Renato Spahni, Marco Steinacher, Soenke Zaehle, Lex Bouwman, Iain Colin Prentice, et al. Multiple greenhouse-gas feedbacks from the land biosphere under future climate change scenarios. *Nature Climate Change*, 3(7):666–672, 2013.

3. Josep G Canadell, Corinne Le Quéré, Michael R Raupach, Christopher B Field, Erik T Buitenhuis, Philippe Ciais, Thomas J Conway, Nathan P Gillett, RA Houghton, and Gregg Marland. Contributions to accelerating atmospheric co2 growth from economic activity, carbon intensity, and efficiency of natural sinks. *Proceedings of the national academy of sciences*, 104(47):18866–18870, 2007.

4. J.G. Canadell, P.M.S. Monteiro, M.H. Costa, L. Cotrim da Cunha, P.M. Cox, A.V. Eliseev, S. Henson, M. Ishii, S. Jaccard, C. Koven, A. Lohila, P.K. Patra, S. Piao, J. Rogelj, S. Syampungani, S. Zaehle, and K. Zickfeld. *Global Carbon and other Biogeochemical Cycles and Feedbacks*, page 673–816. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2021. doi: 10.1017/9781009157896.007.

5. Florian Humpenöder, Kristine Karstens, Hermann Lotze-Campen, Jens Leifeld, Lorenzo Menichetti, Alexandra Barthelmes, and Alexander Popp. Peatland protection and restoration are key for climate change mitigation. *Environmental Research Letters*, 15(10):104093, 2020.

6. Jiren Xu, Paul J Morris, Junguo Liu, and Joseph Holden. Peatmap: Refining estimates of global peatland distribution based on a meta-analysis. *Catena*, 160:134–140, 2018.

7. J Loisel, Angela V Gallego-Sala, and Z Yu. Global-scale pattern of peatland sphagnum growth driven by photosynthetically active radiation and growing season length. *Biogeosciences*, 9(7):2737–2746, 2012.

8. Zicheng Yu, Julie Loisel, Daniel P Brosseau, David W Beilman, and Stephanie J Hunt. Global peatland dynamics since the last glacial maximum. *Geophysical research letters*, 37(13), 2010.

9. Merritt R Turetsky, Benjamin Bond-Lamberty, E Euskirchen, Julie Talbot, Steve Frolking, A David McGuire, and E-S Tuittila. The resilience and functional role of moss in boreal and arctic ecosystems. *New Phytologist*, 196(1):49–67, 2012.

10. Aljosja Hooijer, Susan Page, JG Canadell, Marcel Silvius, Jaap Kwadijk, H Wösten, and Jyrki Jauhiainen. Current and future co 2 emissions from drained peatlands in southeast asia. *Biogeosciences*, 7(5):1505–1514, 2010.

11. D Gilbert, Christian Amblard, Gilles Bourdier, and A-J Francez. The microbial loop at the surface of a peatland: structure, function, and impact of nutrient input. *Microbial ecology*, 35(1):83–93, 1998.

12. Daniel Gilbert and Edward AD Mitchell. Microbial diversity in sphagnum peatlands. *Developments in earth surface processes*, 9:287–318, 2006.

13. Christos Gougoulias, Joanna M Clark, and Liz J Shaw. The role of soil microbes in the global carbon cycle: tracking the below-ground microbial processing of plant-derived carbon for manipulating carbon dynamics in agricultural systems. *Journal of the Science of Food and Agriculture*, 94(12):2362–2371, 2014.

14. Susan Trumbore. Carbon respired by terrestrial ecosystems–recent progress and challenges. *Global Change Biology*, 12(2):141–153, 2006.

15. Andreas Berg, Åsa Danielsson, and Bo H Svensson. Transfer of fixed-n from n2-fixing cyanobacteria associated with the moss sphagnum riparium results in enhanced growth of the moss. *Plant and soil*, 362(1):271–278, 2013.

16. Melanie A Vile, R Kelman Wieder, Tatjana Živković, Kimberli D Scott, Dale H Vitt, Jeremy A Hartsock, Christine L Iosue, James C Quinn, Meaghan Petix, Hope M Fillingim, et al. N2-fixation by methanotrophs sustains carbon and nitrogen accumulation in pristine peatlands. *Biogeochemistry*, 121(2):317–328, 2014.

17. Kathrin Rousk, Davey L Jones, and Thomas H DeLuca. Moss-cyanobacteria associations as biogenic sources of nitrogen in boreal forest ecosystems. *Frontiers in Microbiology*, 4:150, 2013.

18. Nardy Kip, Wenjing Ouyang, Julia van Winden, Ashna Raghoebarsing, Laura van Niftrik, Arjan Pol, Yao Pan, Levente Bodrossy, Elly G van Donselaar, Gert-Jan Reichart, et al. Detection, isolation, and characterization of acidophilic methanotrophs from sphagnum mosses. *Applied and environmental microbiology*, 77(16):5643–5654, 2011.

19. Nardy Kip, Christian Fritz, ES Langelaan, Yao Pan, Lev Bodrossy, Veronica Pancotto, MSM Jetten, AJP Smolders, and HJM Op den Camp. Methanotrophic activity and diversity in different sphagnum magellanicum dominated habitats in the southernmost peat bogs of patagonia. *Biogeosciences*, 9(1):47–55, 2012.

20. Max Reumer, Monika Harnisz, Hyo Jung Lee, Andreas Reim, Oliver Grunert, Anuliina Putkinen, Hannu Fritze, Paul LE Bodelier, and Adrian Ho. Impact of peat mining and restoration on methane turnover potential and methane-cycling microorganisms in a northern bog. *Applied and environmental microbiology*, 84(3):e02218–17, 2018.

21. AV Shcherbakov, AV Bragina, E Yu Kuzmina, Christian Berg, AN Muntyan, NM Makarova, NV Malfanova, Massimiliano Cardinale, Gabriele Berg, VK Chebotar, et al. Endophytic bacteria of sphagnum mosses as promising objects of agricultural microbiology. *Microbiology*, 82(3):306–315, 2013.

22. Alyssa A Carrell, Travis J Lawrence, Kristine Grace M Cabugao, Dana L Carper, Dale A Pelletier, Jun Hyung Lee, Sara S Jawdy, Jane Grimwood, Jeremy Schmutz, Paul J Hanson, et al. Habitat-adapted microbial communities mediate sphagnum peatmoss resilience to warming. *New Phytologist*, 234(6):2111–2125, 2022.

23. Katja Opelt, Vladimir Chobot, Franz Hadacek, Susan Schönmann, Leo Eberl, and Gabriele Berg. Investigations of the structure and function of bacterial communities associated with sphagnum mosses. *Environmental microbiology*, 9(11):2795–2809, 2007.

24. Verity G Salmon, Deanne J Brice, Scott Bridgham, Joanne Childs, Jake Graham, Natalie A Griffiths, Kirsten Hofmockel, Colleen M Iversen, Terri M Jicha, Randy K Kolka, et al. Nitrogen and phosphorus cycling in an ombrotrophic peatland: a benchmark for assessing change. *Plant and Soil*, 466(1):649–674, 2021.

25. Alyssa A Carrell, Max Kolton, Jennifer B Glass, Dale A Pelletier, Melissa J Warren, Joel E Kostka, Colleen M Iversen, Paul J Hanson, and David J Weston. Experimental warming alters the community composition, diversity, and n2 fixation activity of peat moss (sphagnum fallax) microbiomes. *Global Change Biology*, 25(9):2993–3004, 2019.

26. Alyssa A Carrell, Dušan Veličković, Travis J Lawrence, Benjamin P Bowen, Katherine B Louie, Dana L Carper, Rosalie K Chu, Hugh D Mitchell, Galya Orr, Lye Meng Markillie, et al. Novel metabolic interactions and environmental conditions mediate the boreal peatmoss-cyanobacteria mutualism. *The ISME journal*, 16(4):1074–1085, 2022.

27. Kevin H Wyatt, Kevin S McCann, Allison R Rober, and Merritt R Turetsky. Trophic interactions regulate peatland carbon cycling. *Ecology Letters*, 24(4):781–790, 2021.

28. Yinon M Bar-On, Rob Phillips, and Ron Milo. The biomass distribution on earth. *Proceedings of the National Academy of Sciences*, 115(25):6506–6511, 2018.

29. Ben Bond-Lamberty and Allison Thomson. Temperature-associated increases in the global soil respiration record. *Nature*, 464(7288):579–582, 2010.

30. Vincent EJ Jassey, Constant Signarbieux, Stephan Hättenschwiler, Luca Bragazza, Alexandre Buttler, Frédéric Delarue, Bertrand Fournier, Daniel Gilbert, Fatima Laggoun-Défarge, Enrique Lara, et al. An unexpected role for mixotrophs in the response of peatland carbon cycling to climate warming. *Scientific reports*, 5(1):1–10, 2015.

31. Zhilei Gao, Ida Karlsson, Stefan Geisen, George Kowalchuk, and Alexandre Jousset. Protists: puppet masters of the rhizosphere microbiome. *Trends in Plant Science*, 24(2):165–176, 2019.

32. Madhav P Thakur and Stefan Geisen. Trophic regulations of the soil microbiome. *Trends in microbiology*, 27(9):771–780, 2019.

33. Jennifer D Rocca, Andrea Yammine, Marie Simonin, and Jean P Gibert. Protist predation influences the temperature response of bacterial communities. *Frontiers in microbiology*, 13, 2022.

34. Wu Xiong, Yuqi Song, Keming Yang, Yian Gu, Zhong Wei, George A Kowalchuk, Yangchun Xu, Alexandre Jousset, Qirong Shen, and Stefan Geisen. Rhizosphere protists are key determinants of plant health. *Microbiome*, 8(1):1–9, 2020.

35. Javier A Ceja-Navarro, Yuan Wang, Daliang Ning, Abelardo Arellano, Leila Ramanculova, Mengting Maggie Yuan, Alyssa Byer, Kelly D Craven, Malay C Saha, Eoin L Brodie, et al. Protist diversity and community complexity in the rhizosphere of switchgrass are dynamic as plants develop. *Microbiome*, 9(1):1–18, 2021.

36. Daniel Gómez-Pérez, Monja Schmid, Vasvi Chaudhry, Ana Velic, Boris Maček, Ariane Kemen, and Eric Kemen. Proteins released into the plant apoplast by the obligate parasitic protist albugo selectively repress phyllosphere-associated bacteria. *bioRxiv*, page 11, 2022. doi: 10.1101/2022.05.16.492175.

37. Iqra Bashir, Aadil Farooq War, Iflah Rafiq, Zafar A Reshi, Irfan Rashid, and Yogesh S Shouche. Phyllosphere microbiome: Diversity and functions. *Microbiological Research*, 254:126888, 2022.

38. Stefan Geisen, Enrique Lara, Edward AD Mitchell, Eckhard Völcker, and Valentyna Krashevska. Soil protist life matters! *Soil Organisms*, 92(3):189–196, 2020.

39. Sai Guo, Wu Xiong, Xinnan Hang, Zhilei Gao, Zixuan Jiao, Hongjun Liu, Yani Mo, Nan Zhang, George A Kowalchuk, Rong Li, et al. Protists as main indicators and determinants of plant performance. *Microbiome*, 9(1):1–11, 2021.

40. Cyrille Violle, Marie-Laure Navas, Denis Vile, Elena Kazakou, Claire Fortunel, Irène Hummel, and Eric Garnier. Let the concept of trait be functional! *Oikos*, 116(5):882–892, 2007.

41. Daniel J Wieczynski, Pranav Singla, Adrian Doan, Alexandra Singleton, Ze-Yi Han,

Samantha Votzke, Andrea Yammine, and Jean P Gibert. Linking species traits and demography to explain complex temperature responses across levels of organization. *Proceedings of the National Academy of Sciences*, 118(42), 2021.

42. David Atkinson, Benjamin J Ciotti, and David JS Montagnes. Protists decrease in size linearly with temperature: ca. 2.5% c⁻1. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 270(1533):2605–2611, 2003.

43. David JS Montagnes, Susan A Kimmance, and David Wilson. Effects of global and local temperature changes on free living, aquatic protists. In *Conference Proceedings of the 14th International Conference on Comparative Physiology, Climate Changes: Effects on Plants, Animals, and Humans (eds Bolis CL, Keines R, Elia M et al.). Available on CD ROM (from the senior author)*, pages 1–13, 2002.

44. D Atkinson. Temperature and organism size: a biological law for ectotherms? *Adv Ecol Res*, 25:1–58, 1994.

45. A Clarke and KPP Fraser. Why does metabolism scale with temperature? *Functional ecology*, 18(2):243–251, 2004.

46. K. E. Fussmann, B. Rosenbaum, U. Brose, and B.C. Rall. Interactive effects of shifting body size and feeding adaptation drive interaction strengths of protist predators under warming. *bioRxiv*, pages 1–34, 2017. doi: 10.1101/101675.

47. Susanne Menden-Deuer, Caitlyn Lawrence, and Gayantonia Franzè. Herbivorous protist growth and grazing rates at in situ and artificially elevated temperatures during an arctic phytoplankton spring bloom. *PeerJ*, 6:e5264, 2018.

48. Gabriel Yvon-Durocher, J Iwan Jones, Mark Trimmer, Guy Woodward, and Jose M Montoya. Warming alters the metabolic balance of ecosystems. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365(1549):2117–2126, 2010.

49. Susanne Wilken, Jef Huisman, Suzanne Naus-Wiezer, and Ellen Van Donk. Mixotrophic organisms become more heterotrophic with rising temperature. *Ecology letters*, 16(2):225–233, 2013.

50. M Matsubayashi, H Ando, I Kimata, H Nakagawa, M Furuya, H Tani, and K Sasai. Morphological changes and viability of cryptosporidium parvum sporozoites after excystation in cell-free culture media. *Parasitology*, 137(13):1861–1866, 2010.

51. AS Sediq, R Klem, MR Nejadnik, P Meij, and Wim Jiskoot. Label-free, flow-imaging methods for determination of cell concentration and viability. *Pharmaceutical research*, 35(8):1–10, 2018.

52. Ulrich Brose, Jennifer A Dunne, Jose M Montoya, Owen L Petchey, Florian D Schneider, and Ute Jacob. Climate change in size-structured ecosystems, 2012.

53. Jean P Gibert. Temperature directly and indirectly influences food web structure. *Scientific reports*, 9(1):5312, 2019.

54. Mary I O'Connor, Michael F Piehler, Dina M Leech, Andrea Anton, and John F Bruno. Warming and resource availability shift food web structure and metabolism. *PLoS biology*, 7(8):e1000178, 2009.

55. Daniel J Wieczynski, Kristin M Yoshimura, Elizabeth R Denison, Stefan Geisen, Jennifer M DeBruyn, A Jonathan Shaw, David J Weston, Dale A Pelletier, Steven W Wilhelm, and Jean P Gibert. Viral infections likely mediate microbial controls on ecosystem responses to global warming. *FEMS Microbiology Ecology*, page fiad016, 2023.

56. Paul J Hanson, Jeffery S Riggs, W Robert Nettles, Jana R Phillips, Misha B Krassovski, Leslie A Hook, Lianhong Gu, Andrew D Richardson, Donald M Aubrecht, Daniel M Ricciuto, et al. Attaining whole-ecosystem warming using air and deep-soil heating methods with an elevated co 2 atmosphere. *Biogeosciences*, 14(4):861–883, 2017.

57. David Atkinson, Simon A Morley, and Roger N Hughes. From cells to colonies: at what levels of body organization does the 'temperature-size rule' apply? *Evolution & development*, 8(2):202–214, 2006.

58. HyeonSeok Shin, Seong-Joo Hong, Chan Yoo, Mi Han, Hookeun Lee, Hyung-Kyoon Choi, Suhyung Cho, Choul-Gyun Lee, and Byung-Kwan Cho. Genome-wide transcriptome analysis revealed organelle specific responses to temperature variations in algae. *Scientific reports*, 6(1):1–11, 2016.

59. Ulrich Sommer, Carolin Paul, and Maria Moustaka-Gouni. Warming and ocean acidification effects on phytoplankton—from species shifts to size shifts within species in a mesocosm experiment. *PloS one*, 10(5):e0125239, 2015.

60. Ulrich Sommer, Kalista H Peter, Savvas Genitsaris, and Maria Moustaka-Gouni. Do marine phytoplankton follow b ergmann's rule sensu lato? *Biological Reviews*, 92(2):1011–1026, 2017.

61. Jean P Gibert, Ze-Yi Han, Daniel J Wieczynski, Samantha Votzke, and Andrea Yammine. Feedbacks between size and density determine rapid eco-phenotypic dynamics. *Functional Ecology*, 36(7):1668–1680, 2022.

62. Y Nakano, K Miyatake, H Okuno, K Hamazaki, S Takenaka, N Honami, M Kiyota, I Aiga, and J Kondo. Growth of photosynthetic algae euglena in high co2 conditions and its photosynthetic characteristics. In *International Symposium on Plant Production in Closed Ecosystems 440*, pages 49–54, 1996.

63. PJ Hanson, AL Gill, X Xu, JR Phillips, DJ Weston, RK Kolka, JS Riggs, and LA Hook. Intermediate-scale community-level flux of co2 and ch4 in a minnesota peatland: putting the spruce project in a global context. *Biogeochemistry*, 129(3):255–272, 2016.

64. Malak M Tfaily, William T Cooper, Joel E Kostka, Patrick R Chanton, Christopher W Schadt, Paul J Hanson, Colleen M Iversen, and Jeffrey P Chanton. Organic matter transformation in the peat column at marcell experimental forest: humification and vertical stratification. *Journal of Geophysical Research: Biogeosciences*, 119(4):661–675, 2014.

65. Chris Fraley and Adrian E Raftery. Model-based clustering, discriminant analysis, and density estimation. *Journal of the American statistical Association*, 97(458):611–631, 2002.

66. Alboukadel Kassambara and Maintainer Alboukadel Kassambara. Package 'ggpubr'. *R package version 0.1*, 6, 2020.

67. Simon Wood and Maintainer Simon Wood. Package 'mgcv'. *R package version*, 1(29):729, 2015.

68. Luisa W Hugerth, Emilie EL Muller, Yue OO Hu, Laura AM Lebrun, Hugo Roume, Daniel Lundin, Paul Wilmes, and Anders F Andersson. Systematic design of 18s rrna gene primers for determining eukaryotic diversity in microbial consortia. *PloS one*, 9(4):e95567, 2014.

69. Benjamin J Callahan, Paul J McMurdie, Michael J Rosen, Andrew W Han, Amy Jo A Johnson, and Susan P Holmes. Dada2: High-resolution sample inference from illumina amplicon data. *Nature methods*, 13(7):581–583, 2016.

70. Laure Guillou, Dipankar Bachar, Stéphane Audic, David Bass, Cédric Berney, Lucie Bittner, Christophe Boutte, Gaétan Burgaud, Colomban de Vargas, Johan Decelle, et al. The protist ribosomal reference database (pr2): a catalog of unicellular eukaryote small subunit rrna sequences with curated taxonomy. *Nucleic acids research*, 41(D1):D597–D604, 2012.

71. Paul J McMurdie and Susan Holmes. phyloseq: an r package for reproducible interactive analysis and graphics of microbiome census data. *PloS one*, 8(4):e61217, 2013.

72. Roberto Danovaro, Antonio Dell'Anno, and Antonio Pusceddu. Biodiversity response to climate change in a warm deep sea. *Ecology Letters*, 7(9):821–828, 2004.

73. Luke O Frishkoff, Daniel S Karp, Leithen K M'Gonigle, Chase D Mendenhall, Jim Zook, Claire Kremen, Elizabeth A Hadly, and Gretchen C Daily. Loss of avian phylogenetic diversity in neotropical agricultural systems. *science*, 345(6202):1343–1346, 2014.

74. Zhenghu Zhou, Chuankuan Wang, and Yiqi Luo. Meta-analysis of the impacts of global change factors on soil microbial diversity and functionality. *Nature communications*, 11(1):1–10, 2020.

75. Håkan Rydin, Urban Gunnarsson, and Sebastian Sundberg. The role of sphagnum in peatland development and persistence. In *Boreal peatland ecosystems*, pages 47–65. Springer, 2006.

76. JM Waddington, M Strack, and MJ Greenwood. Toward restoring the net carbon sink function of degraded peatlands: Short-term response in co2 exchange to ecosystem-scale restoration. *Journal of Geophysical Research: Biogeosciences*, 115(G1), 2010.

77. Richard D Bardgett, Chris Freeman, and Nicholas J Ostle. Microbial contributions to climate change through carbon cycle feedbacks. *The ISME journal*, 2(8):805–814, 2008.

78. Jizhong Zhou, Kai Xue, Jianping Xie, YE Deng, Liyou Wu, Xiaoli Cheng, Shenfeng Fei, Shiping Deng, Zhili He, Joy D Van Nostrand, et al. Microbial mediation of carbon-cycle feedbacks to climate warming. *Nature Climate Change*, 2(2):106–110, 2012.

79. Thomas P Smith, Thomas JH Thomas, Bernardo García-Carreras, Sofía Sal, Gabriel Yvon-Durocher, Thomas Bell, and Samrāt Pawar. Community-level respiration of prokaryotic microbes may rise with global warming. *Nature communications*, 10(1):1–11, 2019.

80. James F Gillooly, James H Brown, Geoffrey B West, Van M Savage, and Eric L Charnov. Effects of size and temperature on metabolic rate. *science*, 293(5538):2248–2251, 2001.

81. Janet L Gardner, Anne Peters, Michael R Kearney, Leo Joseph, and Robert Heinsohn. Declining body size: a third universal response to warming? *Trends in ecology & evolution*, 26(6):285–291, 2011.

82. Jean P Gibert and John P DeLong. Temperature alters food web body-size structure. *Biology letters*, 10(8):20140473, 2014.

83. Wilco CEP Verberk, David Atkinson, K Natan Hoefnagel, Andrew G Hirst, Curtis R Horne, and Henk Siepel. Shrinking body sizes in response to warming: explanations for the temperature–size rule with special emphasis on the role of oxygen. *Biological Reviews*, 96(1):247–268, 2021.

84. John P DeLong, Chad E Brassil, Emma K Erickson, Valery E Forbes, Etsuko N Moriyama, and Wayne R Riekhof. Dynamic thermal reaction norms and body size oscillations challenge explanations of the temperature–size rule. *Evolutionary Ecology Research*, 18(3):293–303, 2017.

85. Jean P Gibert, Rachel L Allen, Ron J Hruska III, and John P DeLong. The ecological consequences of environmentally induced phenotypic changes. *Ecology Letters*, 20(8):997–1003, 2017.

86. John P DeLong, Jordan G Okie, Melanie E Moses, Richard M Sibly, and James H Brown. Shifts in metabolic scaling, production, and efficiency across major evolutionary transitions of life. *Proceedings of the National Academy of Sciences*, 107(29):12941–12945, 2010.

87. Jean P Gibert, Daniel J Wieczynski, Ze-Yi Han, and Andrea Yammine. Rapid eco-phenotypic feedback and the temperature response of biomass dynamics. *Ecology and Evolution*, 13(1):e9685, 2023.

88. Joel G Kingsolver and Raymond B Huey. Size, temperature, and fitness: three rules. *Evolutionary Ecology Research*, 10(2):251–268, 2008.

89. Lennart T Bach, Santiago Alvarez-Fernandez, Thomas Hornick, Annegret Stuhr, and Ulf Riebesell. Simulated ocean acidification reveals winners and losers in coastal phytoplankton. *PloS one*, 12(11):e0188198, 2017.

90. Alexandra Coello-Camba, Susana Agustí, Johnna Holding, Jesús M Arrieta, and Carlos M Duarte. Interactive effect of temperature and co2 increase in arctic phytoplankton. *Frontiers in Marine Science*, 1:49, 2014.

91. Andrew Clarke. *Principles of thermal ecology: Temperature, energy and life*. Oxford University Press, 2017.

92. Allison R Rober, Kevin S McCann, Merritt R Turetsky, and Kevin H Wyatt. Cascading effects of predators on algal size structure. *Journal of Phycology*, 58(2):308–317, 2022.

93. Edward B Rastetter, Göran I Ågren, and Gaius R Shaver. Responses of n-limited ecosystems to increased co2: A balanced-nutrition, coupled-element-cycles model. *Ecological Applications*, 7(2):444–460, 1997.

94. Petra MA Fransson and Emma M Johansson. Elevated co2 and nitrogen influence exudation of soluble organic compounds by ectomycorrhizal root systems. *FEMS Microbiology Ecology*, 71(2):186–196, 2010.

95. Edward AD Mitchell, Daniel Gilbert, Alexandre Buttler, Christian Amblard, Philippe Grosvernier, and J-M Gobat. Structure of microbial communities in sphagnum peatlands and effect of atmospheric carbon dioxide enrichment. *Microbial ecology*, 46(2):187–199, 2003.

96. Ina C Meier, Seth G Pritchard, Edward R Brzostek, M Luke McCormack, and Richard P Phillips. The rhizosphere and hyphosphere differ in their impacts on carbon and nitrogen cycling in forests exposed to elevated co 2. *New Phytologist*, 205(3):1164–1174, 2015.

97. Shauna M Uselman, Robert G Qualls, and Richard B Thomas. A test of a potential short cut in the nitrogen cycle: The role of exudation of symbiotically fixed nitrogen from the roots of a n-fixing tree and the effects of increased atmospheric co2 and temperature. *Plant and Soil*, 210(1):21–32, 1999.

98. Takehito Yoshida, Laura E Jones, Stephen P Ellner, Gregor F Fussmann, and Nelson G Hairston. Rapid evolution drives ecological dynamics in a predator–prey system. *Nature*,

424(6946):303–306, 2003.

99. Ze-Yi Han, Daniel J Wieczynski, Andrea Yammine, and Jean P Gibert. Temperature and nutrients drive eco-phenotypic dynamics in a microbial food web. *Proceedings of the Royal Society B*, 290(1992):20222263, 2023.

100. John P DeLong and Jean P Gibert. Gillespie eco-evolutionary models (gem s) reveal the role of heritable trait variation in eco-evolutionary dynamics. *Ecology and evolution*, 6(4):935–945, 2016.

101. Thomas M Luhring and John P DeLong. Trophic cascades alter eco-evolutionary dynamics and body size evolution. *Proceedings of the Royal Society B*, 287(1938):20200526, 2020.

102. Aditee Mitra, Kevin J Flynn, Joann M Burkholder, Terrje Berge, Albert Calbet, John A Raven, Edna Granéli, Pat M Glibert, Per Juel Hansen, Diane K Stoecker, et al. The role of mixotrophic protists in the biological carbon pump. *Biogeosciences*, 11(4):995–1005, 2014.

103. Zhenfeng Liu, Victoria Campbell, Karla B Heidelberg, and David A Caron. Gene expression characterizes different nutritional strategies among three mixotrophic protists. *FEMS Microbiology Ecology*, 92(7):fiw106, 2016.

104. Marc-André Selosse, Marie Charpin, and Fabrice Not. Mixotrophy everywhere on land and in water: the grand écart hypothesis. *Ecology Letters*, 20(2):246–263, 2017.

105. Daniel Wieczynski, Holly Moeller, and Jean-Philippe Gibert. Mixotrophs generate carbon tipping points under warming, 2022.

106. J Limpens, F Berendse, Christian Blodau, JG Canadell, C Freeman, J Holden, N Roulet, Håkan Rydin, and Gabriela Schaepman-Strub. Peatlands and the carbon cycle: from local processes to global implications–a synthesis. *Biogeosciences*, 5(5):1475–1491, 2008.

107. Steve Frolking, Julie Talbot, Miriam C Jones, Claire C Treat, J Boone Kauffman, Eeva-Stiina Tuittila, and Nigel Roulet. Peatlands in the earth's 21st century climate system. *Environmental Reviews*, 19(NA):371–396, 2011.

108. Eville Gorham, Clarence Lehman, Arthur Dyke, Dicky Clymo, and Joannes Janssens. Long-term carbon sequestration in north american peatlands. *Quaternary Science Reviews*, 58:77–82, 2012.

109. Qinglong You, Ziyi Cai, Nick Pepin, Deliang Chen, Bodo Ahrens, Zhihong Jiang, Fangying Wu, Shichang Kang, Ruonan Zhang, Tonghua Wu, et al. Warming amplification over the arctic pole and third pole: Trends, mechanisms and consequences. *Earth-Science Reviews*, 217:103625, 2021.

110. Coral J Fung Shek. *Dynamic Imaging Particle Technology for Quantitative Morphological Analysis and Cell Counting*. PhD thesis, Johns Hopkins University, 2015.