1

BMP signaling in the *C. elegans* immune response to bacterial pathogen regulates

anti-microbial peptide expression through cell non-autonomous activity

Emma Jo Ciccarelli [1,2], Moshe Bendelstein [1], Cathy Savage-Dunn [1,2] *

[1] Department of Biology, Queens College, CUNY, Flushing NY

[2] PhD Program in Biology, The Graduate Center, CUNY, New York NY

* Corresponding author

E-mail: cathy.savagedunn@qc.cuny.edu

Short title: Tissue specificity in *C. elegans* host-pathogen response

Abstract

Host response to pathogens recruits multiple tissues in part through conserved cell signaling pathways. In *C. elegans*, the bone morphogenetic protein (BMP) like DBL-1 signaling pathway has a role in the response to infection in addition to other roles in development as well as several post-developmental functions. In the regulation of body size and fat storage, the DBL-1 pathway has been shown to act through cell autonomous and non-autonomous signaling in the epidermis (hypodermis). We have now elucidated the tissues that respond to DBL-1 signaling in the response to bacterial pathogens. The receptors and Smad signal transducers for DBL-1 are expressed in the pharynx, intestine, and epidermis. We demonstrate that expression of receptor-regulated Smad (R-Smad) gene *sma-3* in either the epidermis or pharynx is sufficient to improve the impaired survival phenotype of *sma-3* mutants and that expression of *sma-3* in the intestine has no effect when exposing worms to bacterial infection of the intestine. We also show by qRT-PCR that two antimicrobial peptide (AMP) genes – *abf-2* and *cnc-2* – are regulated by DBL-1 signaling through R-Smad SMA-3. We show that this AMP regulation is similarly cell non-autonomous, as expression of *sma-3* in either the epidermis or the pharynx is sufficient to induce both *abf-2* and *cnc-2* expression in animals exposed to bacterial infection. These results demonstrate that in response to bacterial pathogen, the DBL-1 pathway regulates AMP expression through R-Smad SMA-3 in a cell non-autonomous manner to confer immunity against infection.

Author Summary

Animals exposed to infection mount a defense through immune activation. Innate immune responses are regulated by conserved cell signaling. One conserved signaling pathway involved in the *C. elegans* immune response is the BMP-like DBL-1 pathway. Here we demonstrate that cell non-autonomous signaling through DBL-1 mediator SMA-3 plays a significant role in the response to bacterial intestinal infection. We also identify two antimicrobial peptides regulated by this signaling mechanism in response to bacterial infection. Our work provides insight into the way in which the BMP-like signaling triggers a systemic response to regulate immunity.

Introduction

Animals encounter a diverse range of microorganisms in their environments, many of which can pose a risk for infection. This exposure has resulted in the development of immune networks to protect the host from pathogenic threats. Innate immunity provides an immediate response to infection and functions through highly conserved mechanisms. In humans, innate immunity involves physical barriers such as the skin, recruitment of phagocytic cells, and the release of antimicrobial peptides. Expression of these antimicrobial peptides is regulated by a number of signaling pathways and these mechanisms are conserved across metazoan species [1-9]. The soil-dwelling nematode *Caenorhabditis elegans* naturally encounters many pathogens, thus necessitating a functioning immune response system to provide adequate protection. The *C. elegans* lacks an antibody-based acquired immunity, so response is limited to innate immunity including the release of antimicrobial peptides, however the signaling pathways responsible for their regulation are highly conserved, even in mammals. For this reason, *C. elegans* has served as

a useful organism in which to study many of the signaling pathways that are involved in

mammalian innate immunity and allows for the study of how they regulate an antimicrobial

response [3-9].

One pathway that has been shown to be involved in the *C. elegans* immune response is the BMP-

like DBL-1 signaling pathway [10]. BMPs are members of the large conserved Transforming

Growth Factor beta (TGF-β) family of secreted peptide growth factors [11]. Members of the

TGF-β family play critical roles in adaptive and innate immunity [12]. In *C. elegans,* the DBL-1

pathway is initiated through the neuronally expressed ligand DBL-1 [13 and 14]). The receptors

DAF-4 and SMA-6 along with the signaling mediators SMA-2, SMA-3, and SMA-4 are

endogenously expressed in the epidermis (also known as hypodermis in *C. elegans*), the

intestine, and the pharynx [10, 14-19]. DBL-1 signaling plays a significant role in body size

regulation, male tail development, mesodermal patterning, and lipid accumulation [14-26] in

addition to its role in the *C. elegans* immune response [10]. This pathway has been well studied

in the *C. elegans* response to the gram-negative bacterium *Serratia marcescens* [10]. Our interest

in this study was to determine the way in which DBL-1 signaling plays a broader role in the

immune response against bacterial pathogens and the mechanisms through which this specific

function is carried out.

Here we show that DBL-1 signaling through the receptor mediated R-Smad SMA-3 is required

for protection against bacterial pathogen in *C. elegans*. We demonstrate that expression of *sma-3*

in pharyngeal muscle or in epidermis is sufficient to restore survival of *sma-3* deficient animals.

We additionally demonstrate that DBL-1 signaling regulates expression of antimicrobial peptides

ABF-2 (antibacterial factor related) [27] and CNC-2 (Caenacin) [28] and that this regulation is

through R-Smad SMA-3. We further show that SMA-3 regulates ABF-2 and CNC-2 expression

in a cell non-autonomous manner, with *sma-3* expression in either epidermal or pharyngeal tissue

proving sufficient to induce ABF-2 and CNC-2 expression following pathogen exposure. We

also demonstrate that *sma-3* mutants suffer a reduced pharyngeal pumping rate that could

increase sensitivity to pathogens and that expression of *sma-3* in pharyngeal muscle restores the

pumping rate to control levels.

Results

DBL-1 Pathway R-Smad SMA-3 is required for survival against bacterial pathogen

It has previously been shown that DBL-1 signaling activity is required for *C. elegans* survival against infection by the gram-negative bacterium *S. marcescens* [10]. It has also been shown that in response to fungal infection with *D. coniospora*, DBL-1 signals through a non-canonical signaling cascade to provide protection for the *C. elegans* host [28]. To determine whether the components of the DBL-1 pathway are required for the response to other bacterial pathogens, we analyzed the survival phenotypes of R-Smad mutant *sma-3*. Animals were transferred from *E. coli* food to pathogenic bacteria at the fourth larval (L4) stage and survival on pathogen was monitored through adulthood until dead. When we compare the survival of *sma-3* mutant animals, we see that they display a significantly reduced survival phenotype, as compared to control animals when exposed to *S. marcescens* infection, thus indicating that SMA-3 is involved in the response to *Serratia* infection (Fig 1B). We similarly observed that mutating *sma-3* results in reduced survival when infected with the gram-negative bacterium *P. luminescens* (Fig 1C). We conclude that SMA-3 plays a significant role in *C. elegans* survival against different bacterial pathogens.

Fig 1. DBL-1 Pathway R-Smad SMA-3 is required for survival against bacterial pathogen

A. BMP signaling in *C. elegans* through the DBL-1 pathway. B. Survival of *sma-3(wk30)* on *S. marcescens.* N values: Control (99), *sma-3* (55). C. Survival of *sma-3(wk30)* on *P. luminescens*. N values: Control (67), *sma-3* (63). Survival analyses were repeated. Statistical analysis done using Log-rank (Mantel-Cox) Test. *** P ≤ 0.001. **** P < 0.0001.

Expressing *sma-3* in pharyngeal muscle rescues survival of *sma-3* mutants

As previously noted, DBL-1 pathway Smads – SMA-2, SMA-3, and SMA-4 – are endogenously expressed in the epidermis, the intestine, and the pharynx [10 and 14-19]. *sma-3* mutants have a small body size phenotype compared to control animals [14, 16-26]. Previous work has shown that expression of *sma-3* in the epidermis alone is sufficient to rescue the body size phenotype in a *sma-3* mutant background [17]. Additionally, DBL-1 signaling is cell non-autonomous in its effect on regulation of intestinal lipid stores, as expression of *sma-3* in the epidermis is sufficient to rescue the low-fat phenotype of DBL-1 pathway mutants [26]. From this information we wanted to see in which tissue(s) SMA-3 activity would be required to provide protection against bacterial infection. We used transgenic *C. elegans* strains with a *sma-3* mutant background to conduct these experiments. These strains have *sma-3* expressed under tissue-specific promoters (epidermal expression using *vha-7p*; intestinal expression using *vha-6p*; pharyngeal muscle expression using *myo-2p*; control using endogenous *sma-3p* promoter), allowing for expression of *sma-3* in a tissue-specific manner. Upon exposure to *S. marcescens* we found that expression of *sma-3* in either epidermal or pharyngeal tissue resulted in rescue of the *sma-3* mutant reduced survival phenotype. Pharynx-specific expression of *sma-3* provided the most significant rescue of the *sma-3* phenotype (Fig 2A). We repeated this experiment using *P. luminescens* bacteria as our pathogen and similarly found that pharyngeal expression of *sma-3* alone was sufficient to rescue the *sma-3* mutant survival phenotype (Fig 2B). Interestingly, on this pathogen, intestinal expression of *sma-3* resulted in a worse survival outcome compared to our *sma-3* mutants. From these results we conclude that the pharynx, and to an extent the epidermis, plays a significant role for DBL-1 pathway activity in response to bacterial infection in *C. elegans*.

Figure 2. Expressing *sma-3* in pharyngeal muscle rescues survival of *sma-3* mutants

A. Survival analysis of tissue specific *sma-3* expressing strains on *S. marcescens*. Tissue specific

strains have *sma-3* expressed under regulation of tissue specific promotors in a *sma-3(wk30)*

mutant background.

N values: Control (47), *sma-3* (45) Epidermis (41), Intestine (58), Pharynx (41). B. Survival of

tissue specific *sma-3* expressing strains on *P. luminescens.* N values: Control (102), *sma-3* (82)

Epidermis (63), Intestine (134), Pharynx (131). Survival analyses were repeated. Statistical

analysis done using Log-rank (Mantel-Cox) Test. ns P > 0.05. * P ≤ 0.05. ** P ≤ 0.01. *** P ≤

0.001. **** P < 0.0001. Black asterisks show significance relative to control; magenta shows

significance relative to *sma-3* mutant.

DBL-1 regulates expression of AMPs *abf-2* and *cnc-2* through SMA-3

We next wanted to see if the protective nature of DBL-1 signaling against bacterial infection is due to induction of antimicrobial defense genes. We began by testing a panel of 12 immune response genes via qRT-PCR to determine whether DBL-1 signaling regulates their expression when animals are exposed to infection (Supplemental Figure S1). L4 animals were transferred from control plates to pathogen and collected as Day 1 adults following 24-hour exposure to pathogen. From this analysis we found two genes that demonstrated the expression pattern we expected if their regulation was mediated through DBL-1. Both *abf-2* and *cnc-2* were significantly upregulated in control animals upon 24-hour exposure to bacterial pathogen. Neither of these genes had a significant increase in expression in *dbl-1* mutant animals under the same infection conditions (Fig 3A).

ABF-2 is an antibacterial factor related protein that has been shown to be expressed in the pharynx of *C. elegans* and is involved in the response against bacterial infection [26]. *cnc-2* encodes a Caenacin protein previously shown to be involved in the *C. elegans* immune response to fungal infection, where it is regulated in epidermal tissue [28]. These results suggest that both *abf-2* and *cnc-2* are regulated by DBL-1 signaling during the response to bacterial pathogen. We then used qRT-PCR to test whether *sma-3* mutants demonstrate any change in *abf-2* levels when exposed to bacterial pathogen. Following 24-hour exposure to *P. luminescens* bacteria, *sma-3* mutants demonstrated no increase in *abf-2* levels as compared to control conditions. Additionally, *sma-3* mutants expressed overall lower levels of *abf-2* than control animals even in control conditions (Fig 3B). From these data we conclude that DBL-1 signaling does regulate

expression of known immune response genes and that loss of the signaling effector SMA-3

prevents induction of *abf-2* expression. This observation indicates that SMA-3 plays a role in the

regulation of antimicrobial peptide ABF-2. When we use qRT-PCR to assess the expression

pattern of *cnc-2* in *sma-3* mutants we find a similar pattern for *cnc-2* expression. Control animals

have a significant increase in *cnc-2* expression under infection conditions while *sma-3* mutants

demonstrate no significant change in *cnc-2* expression when exposed to bacterial pathogen (Fig

3B).

As previously noted, earlier work has shown that *cnc-2* is involved in the *C. elegans* response to

fungal infection [28]. We therefore followed up these experiments with a survival analysis of a

cnc-2 mutant on *P. luminescens* to ensure that *cnc-2* plays a role in bacterial infection as well as

fungal infection. Our results demonstrate that *cnc-2* loss of function mutant animals have a

significantly reduced survival rate when exposed to *P. luminescens* as compared to control

animals (Fig 3C). We therefore conclude that CNC-2 plays a role in bacterial infection, in

addition to its known role against fungal pathogen [28].

Figure 3. DBL-1 regulates expression of AMPs *abf-2* and *cnc-2* through SMA-3

A. qRT-PCR results for *abf-2* and *cnc-2* expression in *dbl-1(wk70)* animals. *P. luminescens* used

as bacterial pathogen. B. qRT-PCR results for *abf-2* and *cnc-2* in *sma-3(wk30)* animals. *P.*

luminescens used as bacterial pathogen. C. Survival analysis of *cnc-2(ok3226)* animals on *P.*

luminescens pathogenic bacteria. N values: Control (91), *cnc-2* (74). Survival analysis was

repeated.

Data for qRT-PCR represents repeated analyses from two biological replicates. Statistical

analysis for qRT-PCR was done using One-way ANOVA with multiple comparison test. Error

bars represent standard deviation. Statistical analysis for survival was done using Log-rank

(Mantel-Cox) Test.

ns $P > 0.05$. * $P \le 0.05$. ** $P \le 0.01$. **** $P < 0.0001$.

DBL-1 pathway mediated regulation of AMP expression is cell non-autonomous

Bacterial infection such as that caused by *S. marcescens* or *P. luminescens* results in colonization and infection in the intestine of the worm [10, 29-32]. As our survival analyses demonstrated that intestine-specific *sma-3* expression was not sufficient to ameliorate the *sma-3* reduced survival phenotype, we next wanted to test whether DBL-1 mediated regulation of antimicrobial factors was cell autonomous or cell non-autonomous. We performed qRT-PCR analysis using our tissue-specific *sma-3* expressing strains to establish the expression patterns for *abf-2* after a 24-hour exposure to *P. luminescens* bacteria. Consistent with our survival analysis results, which indicated significance for pharyngeal *sma-3* expression in *C. elegans* survival, *sma-3* expression in the pharynx alone was sufficient to cause an induction in *abf-2* levels following *P. luminescens* exposure (Fig 4A). Interestingly, *sma-3* expressed in the epidermis also resulted in significant *abf-2* induction, although less significant than the increase seen in the pharynx expressing strain. We then tested the tissue specific *sma-3* expressing strains for expression patterns of *cnc-2* upon infection. We similarly saw a significant increase in *cnc-2* expression in either the epidermis or pharynx expressing strains when exposed to bacterial pathogen (Fig 4B). Together these results indicate that regulation of AMP expression by DBL-1 signaling is indeed cell non-autonomous.

Figure 4. SMA-3 mediated regulation of AMP expression is cell non-autonomous

A. qRT-PCR results for *abf-2* in tissue specific *sma-3* expressing animals. B. qRT-PCR results

for *cnc-2* in tissue specific *sma-3* expressing animals. *P. luminescens* used as bacterial pathogen.

Data for qRT-PCR represents repeated analyses from two biological replicates. Statistical

analysis for qRT-PCR was done using One-way ANOVA with multiple comparison test. Error

bars represent standard deviation. Relative Expression

ns $P > 0.05$. * $P ≤ 0.05$. ** $P ≤ 0.01$. *** $P < 0.001$.

Expression of *sma-3* in pharyngeal muscle or epidermis restores pumping rate in *sma-3* mutants

We next investigated whether *sma-3* expression in the pharynx resulted in any effect on the mechanical function of the pharynx. We assayed the pharyngeal pumping rate of animals. To match the conditions of our survival and gene expression studies, we transferred L4 animals from control plates to plates seeded with pathogenic *P. luminescens* bacteria and measured pumping rates in Day 1 adults after 24 hours of exposure to pathogen. We found that *sma-3* mutants had a reduction in pumping rate as compared to control animals on pathogen. The pumping rate of *sma-3* mutants was restored when *sma-3* was expressed solely in the pharynx (Fig 5A) or in the epidermis (Fig 5B), indicating both cell non-autonomous and potentially cell autonomous regulation of pumping rate by SMA-3.

Following these results, we decided to assess the survival patterns and pumping rates of two pharyngeal defect mutants to assess whether functionality of the pharynx contributes to survivability on pathogenic bacteria. Here we use *eat-2* and *phm-2,* both involved in pharynx grinder function, for these experiments. Mutation of *eat-2* results in reduced grinder function and a reduced rate of pharyngeal pumping [33-36]. Mutating *phm-2* similarly results in reduced grinder function [33, 37-38]. We found that when we exposed either strain to *P. luminescens* bacteria for 24 hours there was a significant decrease in pharyngeal pumping rate (Fig 6A), consistent with our expectations. When we exposed worms to *P. luminescens* bacteria from the L4 stage, we found that the *phm-2* mutants have a significantly reduced rate of survival compared to control animals. We saw a similar survival phenotype for *eat-2* mutants when exposed to *P. luminescens* as well (Fig 6B). Other studies have demonstrated that both mutations

253   result in sensitivity to bacterial infection, due to live bacteria reaching the intestine for

254   colonization [37-41]. Our results taken with these prior studies suggest that while disrupting

255   pharyngeal function contributes to susceptibility to infection, the effects we observe in our

256   pharyngeal and epidermal *sma-3* rescued animals are unlikely to be due solely to a cell-

257   autonomous effect on grinder function.

Figure 5. Expression of *sma-3* in pharyngeal muscle or epidermis restores pumping rate

from reduction in *sma-3* mutants

A. Pumping rate in Pumps/20 Seconds of *sma-3(wk30)* mutants and pharyngeal *sma-3* rescue

animals on *P. luminescens* bacteria. B. Pumping rate in Pumps/20 Seconds of *sma-3(wk30)*

mutants and epidermal *sma-3* rescue animals on *P. luminescens* bacteria.

10 worms per strain were counted and experiments were repeated on two separate biological

replicates. Statistical analysis done using One way ANOVA with multiple comparison test.

ns $P > 0.05$. *** $P \leq 0.001$. **** $P \leq 0.0001$

Figure 6. Mutants with pharyngeal defect have reduced survival and reduced pumping rate

on *P. luminescens* bacteria

A. Pharyngeal pumping rate in Pumps/20 Seconds for pharyngeal defect mutants *eat-2(ad1116)*

and *phm-2(ad597)* on *P. luminescens* bacteria. 10 worms per strain were counted and

experiments were repeated on two separate biological replicates. Statistical analysis done using

One way ANOVA with multiple comparison test. B. Survival analysis of *eat-2(ad1116)* and

phm-2(ad597) on *P. luminescens* bacteria. N values: Control (96), *eat-2* (58), Control (81),

phm-2 (99). Statistical analysis was done using Log-rank (Mantel-Cox) Test.

**** $P \leq 0.0001$

Discussion

The BMP-like ligand DBL-1 has previously been shown to play a significant role in the *C. elegans* immune response to bacterial pathogen [10]. Additionally, it has been demonstrated that the DBL-1 pathway effector, R-Smad SMA-3 is involved in a non-canonical signaling response to fungal infection, in which it acts without SMA-2 or the Co-Smad SMA-4 [28]. We were able to use survival analysis to determine where DBL-1 pathway effector SMA-3 plays the largest role in the *C. elegans* immune response by observing survival phenotypes in tissue-specific *sma-3* expressing animals. This question was of interest to us as it has already been shown that *sma-3* expressed solely in the epidermis is sufficient to restore normal body size and lipid stores in adult animals [17 and 26]. In this study we see that *sma-3* expression in the intestinal site of infection does not have any measurable effect on survival against bacterial pathogen. However, expression of *sma-3*, and therefore pathway activity, in the pharynx and to a lesser degree the epidermis, does restore survival towards that of control animals. Improved survival of pharynx specific and epidermis specific expressing strains as compared to *sma-3* mutants suggests the cell non-autonomous activity of the DBL-1 pathway on immunity.

We also looked to determine whether the effect of DBL-1 signaling on the immune response to bacterial pathogen was related to regulation of immune response related genes. The antibacterial factor related peptide ABF-2 has previously been shown to play a role in the response to bacterial pathogen [27]. Here we demonstrate through qRT-PCR analysis that DBL-1 signaling regulates the expression of *abf-2*. We also show that this effect is regulated through R-Smad SMA-3. We find that expressing *sma-3* in either the epidermis or pharynx results in a significant increase in *abf-2* and *cnc-2* expression upon exposure to bacterial pathogen. This supports the

idea that DBL-1 signaling to regulate AMP expression is cell non-autonomous. We note that

expression levels in some of the tissue-specific *sma-3* transgenic strains are higher than those

seen in N2 control animals. This is possibly caused by the multi copy number of *sma-3* present in

these transgenic animals. As CNC-2 has previously been correlated with fungal infection of the

epidermis and not with intestinal bacterial infection [28], we also looked to see how *cnc-2*

mutants survived on bacterial pathogen. We found that *cnc-2* mutant animals had a significantly

reduced survival pattern compared to control animals. This is consistent with our findings that

DBL-1 signaling regulates *cnc-2* expression levels in response to bacterial pathogen exposure as

part of the *C. elegans* immune response to bacterial infection.

We were also interested in determining if DBL-1 signaling activity in the pharynx influenced the

physical function of the *C. elegans* pharynx. We used pharyngeal pumping rate as a method for

measuring the effect of *sma-3* expression on pharyngeal mechanical action. We found that when

worms are grown on pathogenic bacteria, animals that are mutant for *sma-3* have a reduced

pharyngeal pumping rate as compared to control animals and that this can be rescued with

expression of *sma-3* in either pharyngeal or epidermal tissue.

Materials and Methods

Nematode Strains and Growth Conditions

C. elegans were maintained on *E. coli* (DA837) at 20°C on EZ worm plates containing

streptomycin. The N2 strain is used as a control unless otherwise specified. In tissue specific

sma-3 experiments, the strains used are: CS152 *qcIs6[sma-3::GFP + rol-6(su1006)]*, CS628

sma-3(wk30); qcIs55[vha-7p::GFP::sma-3 + rol-6(su1006)] ; CS640 *sma-3(wk30);*

qcIs53[myo-2p::GFP::sma-3 + rol-6(su1006)] ; CS619 *sma-3(wk30);*

qcIs59[vha-6p::GFP::sma-3 + rol-6(su1006)]; CS635 *sma-3(wk30); qcIs62[rol-6(su1006)].*

Remaining strains used in this study are*: sma-3(wk30), cnc-2(ok3226), sma-6(wk7),*

sma-2(e502), dbl-1(wk70), phm-2(ad597), eat-2(ad1116).

Bacteria

Control bacteria used in all experiments is *E. coli* strain DA837, cultured at 37°C. *S. marcescens*

strain Db11 cultured at 37°C and *P. luminescens* cultured at 30°C were used for pathogenic

bacteria exposure. *Serratia* (Db11) is seeded on EZ worm plates containing streptomycin and

grown at 37°C. *P. luminescens* is seeded on EZ worm plates with no antibiotic and grown at

30°C.

Survival Analysis

Survival plates were prepared at least one day prior to plating worms. Each plate was seeded

with 500 µl of pathogenic bacteria. FuDR (5-Fluoro-2'-deoxyuridine) was used to prevent

reproduction, at a concentration of 50 µM per plate. This treatment was done to reduce the

338   incidence of matricide by internal hatching of embryos during survival analysis. All survival

339   experiments were carried out at 20°C. All survival analyses were repeated. All graphs made

340   using GraphPad Prism software and statistical analysis performed using Logrank/Mantel Cox

341   test.

qRT-PCR Analysis

Worms were synchronized using overnight egg lay followed by four-hour synchronization.

When animals reached L4 stage, they were washed and moved to experimental plates for 24-

hour exposure to *P. luminescens*. Animals were collected and washed after 24-hour exposure

and RNA was extracted using previously published protocol [42] followed by Qiagen RNeasy

miniprep kit (Cat. No. 74104). cDNA was generated using Invitrogen SuperScript IV VILO

Master Mix (Cat. No.11756050). qRT-PCR analysis was done using Applied Biosystems *Power*

SYBR Green PCR Master Mix (Cat. No. 4367659). Delta delta Ct analysis was done using

Applied Biosystems and StepOne software. All qRT-PCR analysis was repeated on separate

biological replicates. All graphs were made using GraphPad Prism software and statistical

analysis was performed using One-way ANOVA with Multiple Comparison Test.

Pumping Rate Experiments

356   Pumping rate experiments were done with animals exposed at L4 stage for 24 hours on *P.*

357   *luminescens* bacteria. Pharyngeal pumps counted over 20 second timespan. All pumping

358   experiments were repeated. Graph made using GraphPad Prism software and statistical analysis

359   performed using One-way ANOVA with Multiple Comparisons Test.

ACKNOWLEDGEMENTS

This work was funded by R15GM112147 to CSD (National Institutes of Health/NIGMS) and by a PSC-CUNY grant to CSD. Some strains were provided by the Caenorhabditis Genetics Center, which is supported by the National Institute of Health - Office of Research Infrastructure Programs (P40 OD010440). The *cnc-2(ok3226)* strain was generated by the *C. elegans* Gene Knockout Project at the Oklahoma Medical Research Foundation, part of the International *C. elegans* Gene Knockout Consortium. This work was carried out in partial fulfillment of the requirements for the Ph.D. degree from the Graduate Center of City University of New York (EJC).

References

1. Medzhitov R, Janeway CA. Innate immune recognition and control of adaptive immune responses. Semin Immunol. 1998 Oct;10(5):351–3.

2. Hoffmann JA, Kafatos FC, Janeway CA, Ezekowitz RA. Phylogenetic perspectives in innate immunity. Science. 1999 May 21;284(5418):1313–8.

3. Kim DH, Feinbaum R, Alloing G, Emerson FE, Garsin DA, Inoue H, et al. A conserved p38 MAP kinase pathway in Caenorhabditis elegans innate immunity. Science. 2002 Jul 26;297(5581):623–6.

4. Schulenburg H, Kurz CL, Ewbank JJ. Evolution of the innate immune system: the worm perspective. Immunol Rev. 2004 Apr;198:36–58.

5. Millet ACM, Ewbank JJ. Immunity in Caenorhabditis elegans. Curr Opin Immunol. 2004 Feb;16(1):4–9.

6. Irazoqui JE, Ng A, Xavier RJ, Ausubel FM. Role for beta-catenin and HOX transcription factors in Caenorhabditis elegans and mammalian host epithelial-pathogen interactions. Proc Natl Acad Sci U S A. 2008 Nov 11;105(45):17469–74.

7. Partridge FA, Gravato-Nobre MJ, Hodgkin J. Signal transduction pathways that function in both development and innate immunity. Developmental Dynamics. 2010;239(5):1330–6.

8. Pukkila-Worley R, Ausubel FM. Immune defense mechanisms in the Caenorhabditis elegans intestinal epithelium. Curr Opin Immunol. 2012 Feb;24(1):3–9.

9. Kim DH, Ewbank JJ. Signaling in the innate immune response [Internet]. WormBook: The Online Review of C. elegans Biology [Internet]. WormBook; 2018 [cited 2023 Jan 16]. Available from: https://www.ncbi.nlm.nih.gov/books/NBK19673/

10. Mallo GV, Kurz CL, Couillault C, Pujol N, Granjeaud S, Kohara Y, et al. Inducible
antibacterial defense system in C. elegans. Curr Biol. 2002 Jul 23;12(14):1209–14.

11. Massagué J. TGF-beta signal transduction. Annu Rev Biochem. 1998;67:753–91.

12. Chen W, ten Dijke P. Immunoregulation by members of the TGFβ superfamily. Nat Rev
Immunol. 2016 Nov 25;16(12): 723-40.

13. Morita K, Flemming AJ, Sugihara Y, Mochii M, Suzuki Y, Yoshida S, et al. A
Caenorhabditis elegans TGF-β, DBL-1, controls the expression of LON-1, a PR-related
protein, that regulates polyploidization and body length. EMBO J. 2002 Mar
1;21(5):1063–73.

14. Suzuki Y, Yandell MD, Roy PJ, Krishna S, Savage-Dunn C, Ross RM, et al. A BMP
homolog acts as a dose-dependent regulator of body size and male tail patterning in
Caenorhabditis elegans. Development. 1999 Jan;126(2):241–50.

15. Inoue T, Thomas JH. Targets of TGF-β Signaling in Caenorhabditis elegans Dauer
Formation. Developmental Biology. 2000 Jan 1;217(1):192–204.

16. Yoshida S, Morita K, Mocchi M, Ueno N. Hypodermal expression of Caenorhabditis
elegans TGF-beta type I receptor SMA-6 is essential for the growth and maintenance of
body length. Developmental Biology. 2001 Dec 1;240(1):32–45.

17. Wang J, Tokarz R, Savage-Dunn C. The expression of TGFbeta signal transducers in the
hypodermis regulates body size in C. elegans. Development. 2002 Nov;129(21):4989–98.

18. Savage-Dunn C, Tokarz R, Wang H, Cohen S, Giannikas C, Padgett RW. SMA-3 Smad
Has Specific and Critical Functions in DBL-1/SMA-6 TGFβ-Related Signaling.
Developmental Biology. 2000 Jul 1;223(1):70–6.

19. Savage C, Das P, Finelli AL, Townsend SR, Sun CY, Baird SE, et al. Caenorhabditis

elegans genes sma-2, sma-3, and sma-4 define a conserved family of transforming growth

factor beta pathway components. Proc Natl Acad Sci U S A. 1996 Jan 23;93(2):790–4.

20. Gumienny TL, Savage-Dunn C. TGF-β signaling in C. elegans [Internet]. WormBook:

The Online Review of C. elegans Biology [Internet]. WormBook; 2018 [cited 2023 Jan

16]. Available from: https://www.ncbi.nlm.nih.gov/books/NBK19692/

21. Heldin CH, Miyazono K, ten Dijke P. TGF-beta signalling from cell membrane to

nucleus through SMAD proteins. Nature. 1997 Dec 4;390(6659):465–71.

22. Padgett RW, Das P, Krishna S. TGF-β signaling, Smads, and tumor suppressors.

BioEssays. 1998;20(5):382–90.

23. Whitman M. Smads and early developmental signaling by the TGFbeta superfamily.

Genes Dev. 1998 Aug 15;12(16):2445–62.

24. Attisano L, Wrana JL. Signal transduction by members of the transforming growth

factor-beta superfamily. Cytokine Growth Factor Rev. 1996 Dec;7(4):327–39.

25. Derynck R, Zhang Y, Feng XH. Smads: transcriptional activators of TGF-beta responses.

Cell. 1998 Dec 11;95(6):737–40.

26. Clark JF, Meade M, Ranepura G, Hall DH, Savage-Dunn C. Caenorhabditis elegans

DBL-1/BMP Regulates Lipid Accumulation via Interaction with Insulin Signaling. G3

(Bethesda). 2018 Jan 4;8(1):343–51.

27. Kato Y, Aizawa T, Hoshino H, Kawano K, Nitta K, Zhang H. abf-1 and abf-2, ASABF-

type antimicrobial peptide genes in Caenorhabditis elegans. Biochem J. 2002 Jan

15;361(Pt 2):221–30.

28. Zugasti O, Ewbank JJ. Neuroimmune regulation of antimicrobial peptide expression by a

noncanonical TGF-beta signaling pathway in Caenorhabditis elegans epidermis. Nat

Immunol. 2009 Mar;10(3):249–56.

29. Kurz CL, Ewbank JJ. Caenorhabditis elegans: an emerging genetic model for the study of

innate immunity. Nat Rev Genet. 2003 May;4(5):380–90.

30. Sato K, Yoshiga T, Hasegawa K. Activated and inactivated immune responses in

Caenorhabditis elegans against Photorhabdus luminescens TT01. Springerplus. 2014 Jun

1;3:274.

31. Tan MW, Mahajan-Miklos S, Ausubel FM. Killing of Caenorhabditis elegans by

Pseudomonas aeruginosa used to model mammalian bacterial pathogenesis. Proceedings

of the National Academy of Sciences. 1999 Jan 19;96(2):715–20.

32. Garsin DA, Sifri CD, Mylonakis E, Qin X, Singh KV, Murray BE, et al. A simple model

host for identifying Gram-positive virulence factors. Proc Natl Acad Sci U S A. 2001 Sep

11;98(19):10892–7.

33. Avery L. The genetics of feeding in Caenorhabditis elegans. Genetics. 1993

Apr;133(4):897–917.

34. Lakowski B, Hekimi S. The genetics of caloric restriction in Caenorhabditis elegans.

Proceedings of the National Academy of Sciences. 1998 Oct 27;95(22):13091–6.

35. Raizen DM, Lee RY, Avery L. Interacting genes required for pharyngeal excitation by

motor neuron MC in Caenorhabditis elegans. Genetics. 1995 Dec;141(4):1365–82.

36. Avery L, You YJ. C. elegans feeding. WormBook. 2012 May 21;1–23.

37. Kumar S, Egan BM, Kocsisova Z, Schneider DL, Murphy JT, Diwan A, et al. Lifespan

Extension in C. elegans Caused by Bacterial Colonization of the Intestine and Subsequent

Activation of an Innate Immune Response. Dev Cell. 2019 Apr 8;49(1):100-117.e6.

38. Scharf A, Pohl F, Egan BM, Kocsisova Z, Kornfeld K. Reproductive Aging in

Caenorhabditis elegans: From Molecules to Ecology. Front Cell Dev Biol. 2021 Sep

16;9:718522.

39. Labrousse A, Chauvet S, Couillault C, Kurz CL, Ewbank JJ. Caenorhabditis elegans is a

model host for Salmonella typhimurium. Curr Biol. 2000 Nov 30;10(23):1543–5.

40. Portal-Celhay C, Bradley ER, Blaser MJ. Control of intestinal bacterial proliferation in

regulation of lifespan in Caenorhabditis elegans. BMC Microbiol. 2012 Mar 27;12:49.

41. Smith MP, Laws TR, Atkins TP, Oyston PCF, de Pomerai DI, Titball RW. A liquid-

based method for the assessment of bacterial pathogenicity using the nematode

Caenorhabditis elegans. FEMS Microbiol Lett. 2002 May 7;210(2):181–5.

42. Yin J, Madaan U, Park A, Aftab N, Savage-Dunn C. Multiple cis elements and GATA

factors regulate a cuticle collagen gene in Caenorhabditis elegans. Genesis. 2015;53(3–

4):278–84.

Supporting Information

S1Fig. qRT-PCR analysis panel of 12 immune response genes in *dbl-1(wk70)* mutants

qRT-PCR analysis of 12 immune response genes in control and *dbl-1(wk70)* mutant animals,

comparing 24-hour bacterial pathogen exposure to control conditions. *P. luminescens* used as

bacterial pathogen. Data for *abf-2* and *cnc-2* represents repeated analyses for two biological

replicates. Remaining genes were analyzed for one biological sample. Graph made using

GraphPad Prism. Statistical analysis for *abf-2* and *cnc-2* done using One-way ANOVA with

multiple comparison test. ns P > 0.05. * P ≤ 0.05. ** P ≤ 0.01.