Immunoproximity biotinylation reveals the axon initial segment proteome

Wei Zhang[1,2,3], Yu Fu[1], Luxin Peng[1], Yuki Ogawa[3], Xiaoyun Ding[3], Anne Rasband[3], Xinyue

Zhou[2], Maya Shelly[5], Matthew N. Rasband[3*], Peng Zou[1,2,4*]

1. College of Chemistry and Molecular Engineering, Synthetic and Functional

Biomolecules Center, Beijing National Laboratory for Molecular Sciences, PKU-

IDG/McGovern Institute for Brain Research, Key Laboratory of Bioorganic Chemistry

and Molecular Engineering of Ministry of Education, Peking University, Beijing, 100871,

China

2. Academy for Advanced Interdisciplinary Studies, PKU-Tsinghua Center for Life Science,

Peking University, Beijing 100871, China

3. Department of Neuroscience, Baylor College of Medicine, Houston, TX, USA

4. Chinese Institute for Brain Research (CIBR), Beijing 102206, China

5. Department of Neurobiology and Behavior, Stony Brook University, NY, USA

[*] Correspondence: zoupeng@pku.edu.cn (P.Z.); rasband@bcm.edu (M.N.R)

ABSTRACT

The axon initial segment (AIS) is a specialized neuronal compartment required for action

potential generation and neuronal polarity. However, understanding the mechanisms

regulating AIS structure and function has been hindered by an incomplete knowledge of

its molecular composition. Here, using immuno-proximity biotinylation we further define

the AIS proteome and its dynamic changes during neuronal maturation. Among the many

AIS proteins identified, we show that SCRIB is highly enriched in the AIS both *in vitro* and

in vivo, and exhibits a periodic architecture like the axonal spectrin-based cytoskeleton.

We found that ankyrinG interacts with and recruits SCRIB to the AIS. However, loss of

SCRIB has no effect on ankyrinG. This powerful and flexible approach further defines the

AIS proteome and provides a rich resource to elucidate the mechanisms regulating AIS

structure and function.

KEYWORDS

axon initial segment, proteomics, proximity labeling, SCRIB, ankyrinG, neuron

INTRODUCTION

The axon initial segment (AIS) is the 20~60 µm long proximal region of the axon

responsible for action potential generation and maintenance of neuronal polarity [1, 2].

Changes in the molecular composition of the AIS, its length, or position can alter neuronal

excitability [3, 4, 5]. Disruption of the AIS causes axons to acquire dendritic characteristics [6,

7]. Recent studies show that AIS disruption occurs in many neurological diseases including

autism, Alzheimer's disease, stroke, bipolar disorder and schizophrenia [8]. Rescuing AIS

integrity and function can ameliorate neurological symptoms in Alzheimer's disease and

Angelman Syndrome mouse models [9, 10]. However, our incomplete knowledge of AIS

components hinders our understanding of the structural and functional regulation of AIS

in health and disease.

Axon initial segments consist of a specialized extracellular matrix, clustered cell

adhesion molecules (CAMs) and voltage gated ion channels, and a unique cytoskeleton.

Among the previously reported AIS proteins, the scaffolding protein ankyrinG (AnkG) is

the master organizer for AIS assembly and maintenance [8]. AnkG links membrane proteins

to the actin cytoskeleton through a tetramer consisting of βIV and αII-spectrin, and to the

microtubule cytoskeleton through the end binding proteins EB1 and EB3 [11]. At the AIS,

microtubules form parallel fascicles thought to be organized by TRIM46 [12, 13]. As the site

for axonal action potential initiation, AIS also have voltage gated sodium (e.g. Nav1.2),

potassium (e.g. Kv7) and calcium channels (e.g. Cav2) that regulate spike generation,

pattern and shape [14]. NF186, a neuron-specific isoform of the CAM neurofascin (NFASC),

assembles and links extracellular matrix molecules (e.g. Brevican) to the AIS cytoskeleton

[15]. In addition, some synaptic proteins (e.g. Gephyrin), cisternal organelle molecules (e.g.

Synaptopodin) and a Kv1 channel complex can also be found at the AIS [16].

The combination of enzyme-mediated proximity-dependent biotinylation and mass

spectrometry-based quantitative proteomics has emerged as a powerful tool to elucidate

endogenous protein complexes in subcellular domains. Two main categories of proximity

labeling methods have been developed based on enzymes used for catalysis: biotin ligase-

based proximity labeling (e.g. BioID and TurboID) and peroxidase-based proximity

labeling (e.g. HRP and APEX2). Pioneering experiments using AIS-targeted BioID were

recently used to uncover a partial AIS proteome. However, the experiments were

restricted to cytoplasmic AIS proteins and the labeling radius of BioID is confined to ~10

nm [17].

Here, we report the development of **i**mmuno **p**roximity **l**abeling in fixed neurons,

together with a multiple ratiometric analysis strategy to define the **AIS** proteome (IPL-

AIS). This method targets endogenous baits without requiring genetic manipulation, thus

avoiding potential artifacts from over-expression or fusion proteins with altered

subcellular localization [18, 19, 20]. This approach also has the advantage of a larger labeling

radius by peroxidase-mediated protein biotinylation in permeabilized and fixed samples

[21], allowing us to deeply mine AIS structural components. We applied quantitative

ratiometric proteomics with multiple controls to identify AIS proteins [22, 23], allowing us to

compare the relative expression of proteins in the AIS against other neuronal

compartments (axon, dendrites or soma).

IPL-AIS profiling at days *in vitro* (DIV) 7, 14 and 21 revealed the dynamic changes in

AIS components during neuronal development. Subsequent validation confirmed the

identification of novel AIS enriched proteins, including the tumor-suppressor protein

scribble (SCRIB encoded by the gene *Scrib*). We found that SCRIB is highly enriched in the

AIS *in vitro* and *in vivo*. In addition, AnkG interacts with SCRIB and is required for SCRIB

localization to the AIS. Together, our experiments define the AIS proteome, which paves

the way to understand the molecular mechanisms regulating AIS structure and function.

RESULTS

Development of antibody targeted proximity labeling at the AIS

The CAM NF186 is highly and stably localized in the AIS [24, 25], therefore we used

NF186 for antibody targeted proximity labeling at the AIS. As illustrated in Fig. 1a, cultured

cortical neurons were fixed and labeled using antibodies to restrict horseradish peroxidase

(HRP) to the AIS. HRP-mediated proximity-dependent biotinylation is triggered with the

addition of biotin-phenol substrates and hydrogen peroxide. Biotinylation can be

evaluated with fluorescent streptavidin and immunoblot analysis. Following enrichment

via affinity purification, biotinylated proteins are digested with trypsin and identified via

quantitative LC-MS/MS analysis.

We optimized antibody dilutions, H_2O_2 concentration, reaction time and different

biotin-phenol (BP) and biotin- aniline (BA) probes in the proximity labeling system using

mature fixed cortical neurons (Supplementary Fig. 1). We found that with optimized

conditions we could achieve robust labeling and specificity after just one minute. Previous

studies showed the peroxidase substrate BP2 performs well for labeling of cytosolic

protein complexes [26], while the substrates BA1 and BA2 exhibit higher reactivity towards

nucleic acids (Supplementary Fig. 1e) [27]. Our analysis revealed biotin-phenol (BP) as the

most efficient and specific substrate for *in vitro* AIS proximity labeling. We employed these

optimized parameters and found strongly biotinylated proteins localized in the AIS (Fig.

1b). Through imaging, we also tested other AIS-directed antibodies for proximity labeling,

including antibodies against AnkG and TRIM46. In our hands, NFASC targeted proximity

labeling exhibited the best performance, with less background and highly specific

biotinylation of AIS.

To remove background signal arising from non-specific binding of antibodies,

experiments omitting the primary antibody (i.e. anti-NFASC) were used as the negative

control. To quantify the level of protein enrichment at the AIS, we used NeuN and MAP2

to define soma and somatodendritic domains, respectively. NeuN localizes in the nuclei

and cytoplasm of neurons [28, 29], while MAP2 is a microtubule-associated protein that is

widely distributed throughout the soma and dendrites [5, 30]. Fluorescence microscopy

showed highly colocalized biotinylation and immunofluorescence labeling for both NeuN

and MAP2 (Fig. 1b). Thus, we applied the same experimental workflow to anti-NeuN and

anti-MAP2 defined compartments and used these as references for ratiometric analysis

of the AIS proteome.

Biochemical characterization by streptavidin-HRP blot and silver staining showed

successful protein biotinylation from anti-NFASC, anti-NeuN, and anti-MAP2 proximity

labeling experiments (Fig. 1c). As expected, the negative control sample omitting primary

antibody yielded significantly less signal; any background signal presumably arose from

detection of endogenously biotinylated proteins. Notably, the biotinylation signal was

stronger in anti-NeuN and anti-MAP2 reference samples compared to the anti-NFASC

sample, which may reflect the higher abundance of and larger volume occupied by NeuN

and MAP2.

To quantitatively compare protein abundance between samples, we designed the

following sets of dimethyl labeling-based ratiometric MS proteomic experiments: 1) anti-

NFASC vs. no primary antibody; 2) anti-NFASC vs. anti-NeuN; and 3) anti-NFASC vs. anti-

MAP2. The first experiment served to remove background labeling arising from non-

specific staining of secondary antibody-HRP conjugates, endogenous biotinylated proteins,

and non-specific protein adsorption on streptavidin-coated beads. The next two

experiments allowed distinction of AIS-specific proteins from those broadly distributed

across the cell. For each set of experiments, three biological replicates were performed

on cultured cortical neurons at DIV14 (Fig. 1d).

A total of 568 proteins were identified after a stringent cut-off analysis (see Methods),

among which several known AIS proteins, including TRIM46, AnkG, and NFASC were highly

enriched in our dataset (Fig. 1e and Supplementary Table 1). On the other hand, we

noticed the low enrichment of EB1 and SEPTIN5, which are also found at the AIS, but not

restricted to the AIS [11, 17], which explains their lack of specific enrichment in our

experiments. Taken together, these results demonstrate that the AIS proteome can be

defined using our IPL-AIS methods.

Identification of AIS proteome at DIV14

Building upon the success of our pilot IPL-AIS experiments, we next sought to

improve the quality of our proteomic dataset through implementing the following

changes in the workflow (Fig. 2a, b): 1) to improve protein abundance quantitation across

multiple samples, we introduced Tandem Mass Tag (TMT) 10-plex isobaric tags to label

tryptic peptides [31], 2) to achieve higher coverage in labeled peptides, we reduced sample

complexity by fractionating peptides prior to loading onto the LC-MS/MS, and 3) to further

improve spatial specificity, we added anti-SMI312 targeted axon proximity labeling to the

reference samples (in addition to NeuN and MAP2) (Fig. 2a). Confocal fluorescence

imaging analysis confirmed axonal localization of anti-SMI312 targeted protein

biotinylation (Supplementary Fig. 2).

Streptavidin-HRP blot analysis confirmed successful enrichment of biotinylated

proteins across antibody targeted proximity labeling samples, whereas negative control

samples yielded negligible signal (Supplementary Fig. 3). A total of 2,755 proteins with at

least 2 unique peptides were detected in the 10-plex TMT MS experiment (Fig. 2c). The

MS intensity was highly reproducible between replicates (Fig. 2d). We then used the

following data analysis pipeline to refine our AIS proteome (Fig. 2c). We first identified

biotinylated proteins by comparing the MS intensities of anti-NFASC samples against

negative controls (-BP and -1Ab). By applying a cutoff ratio of 2, we obtained a list of 1403

biotinylated proteins. For each protein in this raw list, we calculated the ratios of its MS

intensities between anti-NFASC samples versus those in the reference samples (+NeuN,

+MAP2, and +SMI312). We then ranked proteins according to their averaged MS intensity

ratios (+NF/+NeuN, +NF/+MAP2, and +NF/+SMI312) (Fig. 2e) and calculated their

averaged rank scores (Fig. 2f). Notably, our results included many known AIS proteins (Figs

2e, f and 4a), including extracellular proteins (e.g. Brevican and Versican), CAMs (e.g.

NFASC and NRCAM), cytoplasmic proteins (e.g. NDEL1 and LIS1), and cytoskeletal or

cytoskeleton-associated proteins (e.g. βIV-spectrin, αII-spectrin and TRIM46). This broad

coverage of both cytoplasmic and cell surface proteins reflects the large labeling radius of

optimized IPL-AIS. Importantly, our dataset included many voltage-gated ion channels,

such as sodium channels (e.g. Nav1.2 and Navβ2), potassium channels (e.g. Kv7.2, Kv1.2,

Kv2.1 and Kvβ2), and calcium channels (e.g. Cav2.1 and Cav2.2), consistent with their roles

in regulating action potential initiation and shape [14].

AIS candidates (1403) were compared against each reference control and showed

their relative amounts in the AIS versus soma, somatodendritic, or axonal domains (Fig.

2e and Supplementary Table 2). AIS highly expressed proteins (NFASC, Nav1.2, AnkG,

TRIM46 and βIV-spectrin) were always ranked at the top, while more widely expressed AIS

proteins differentiated according to their relative expression level in subcellular domains,

such as the microtubule protein TUBA4A or microtubule related protein MAP6 (Fig. 2e) [17].

To better define AIS enriched proteins, we applied averaged rank scores to re-list the

known and putative AIS proteins (Fig. 2f and Supplementary Table 2). Consistently, highly

expressed, known AIS proteins were found among the top 5. More widely expressed

proteins, such as F-actin monooxygenase MICAL3 (ranked 371) and Rho GTPase activating

protein ARHGAP21 (ranked 455), were not as highly ranked [17].

To obtain a high-confidence AIS proteome list, we took the overlap of the top 200

proteins in each list of MS intensity ratios (+NF/+NeuN, +NF/+MAP2, and +NF/+SMI312).

The resulting high-confidence list contained 71 proteins (Fig. 2c, g and Supplementary

Table 2). Gene Ontology cellular compartment analysis revealed an over-representation

of AIS and node of Ranvier terms due to the similar compositions of these two

compartments (Fig. 2h). The remaining top 10 GOCC terms were consistent with AIS

characteristics and its endocytosis functions [16, 32].

Mapping the AIS proteome across neuronal maturation

The AIS is developmentally and physiologically dynamic [25, 33]. To determine dynamic developmental changes in the AIS proteome, we performed IPL-AIS at DIV7, DIV14, and DIV21. Two parallel 10-plex TMT experiments were performed and DIV14 samples were used as a reference to bridge DIV7 and DIV21 samples (Fig. 3a).

We first analyzed the DIV7 AIS proteome. Parallel analysis of 1532 common proteins revealed highly reproducible protein quantifications (R^2= 0.94) (Fig. 3b). After stringent filtering (Supplementary Fig. 4a), we obtained 1407 biotinylated proteins. Rank analysis revealed the specificity of our DIV7 AIS dataset with top ranked AIS proteins including AnkG, TRIM46, βIV-spectrin, NFASC and Nav1.2 (Fig. 3c). Other AIS proteins were more widely dispersed, possibly due to different expression levels and labeling efficiencies. Through integrating average rank scores of AIS versus soma or somatodendrites, we obtained a final ranked putative AIS proteome including AIS proteins in top positions (Fig. 3d and Supplementary Table 3). Using the same workflow (Supplementary Fig. 4b), we identified 1738 biotinylated proteins at DIV21. Parallel analysis of anti-NFASC replicates showed highly reproducible protein quantifications (R^2= 0.98) (Fig. 3e), and rank scores of AIS versus soma or somatodendritic proteins (Fig. 3f) resulted in AIS proteins being consistently at the top (Fig. 3g and Supplementary Table 3). Some AIS proteins, such as Synaptopodin (SYNPO) and PSD-93 (PSD93) were ranked lower (Fig. 3g) [34, 35].

Comparing biotinylated proteins between DIV7 and DIV14 in 10-plex TMT revealed 173 proteins only present at DIV7 and 152 proteins only present at DIV14, among which there were only two previously reported AIS proteins: Gephyrin and microtubule protein TUBA4A (Supplementary Fig. 4d). DIV14 samples in the two parallel 10-plex TMT experiments presented a very high reproducibility among biological replicates (Supplementary Fig. 4e) and enrichment of known AIS proteins (Supplementary Fig. 4f), allowing us to compare relative protein expression in three stages together. To identify the core set of putative AIS proteins during development, DIV7, 14, 21 and the prior DIV14

datasets (Fig. 2) were combined, which generated a total of 549 common proteins (Fig.

3h). Among these, 534 proteins showed 20% changes in any one pair comparison of the

three time points and were used for heatmap clustering. Six clusters were generated with

distinct expression patterns (Fig. 3i and Supplementary Table 3). 51.7% of proteins were

gradually upregulated and 4.1% of proteins were gradually downregulated along neuronal

development (Fig. 3j). The cluster with the highest expression level at DIV21 included 83.5%

of proteins, while at DIV7 the cluster with the highest expression level included only 15.3%

of proteins.

Consistently, the previously reported AIS proteins exhibited the highest expression

level at DIV21, including NFASC (*Nfasc*), AnkG (*Ank3*), Nav1.2 (*Scn2a*), and βIV-spectrin

(*Sptbn4*), while TRIM46 (*Trim46*) decreased along neuronal development (Fig. 3k).

Quantitative analysis revealed the fold changes and statistical significance between stages

(Supplementary Table 3). For example, microtubule-associated proteins MAP1A (*Map1a*),

MAP6 (*Map6*), sodium/potassium-transporting ATPase subunit alpha-1 (*Atp1a1*) and

casein kinase II subunit alpha (*Csnk2a1*) exhibited a more than two-fold significant change

from DIV7 to 14 (Fig. 3l). From DIV14 to 21, more microtubule or microtubule-associated

proteins like TUBB5 (*Tubb5*), TUBB3 (*Tubb3*) and EB3 (*Mapre3*) significantly increased,

while calcium/calmodulin-dependent protein kinase KCC2D (*Camk2d*) and KCC2A

(*Camk2a*) also had more than two-fold changes (Fig. 3m). Comparison between DIV7 and

21 revealed more components with a two-fold significant increase, including the AIS

proteins NFASC (*Nfasc*), Nav1.2 (*Scn2a*) and GABA(A) receptor subunit gamma-2 (*Gabrg2*),

while TRIM46 (*Trim46*) showed a two-fold significant decrease (Fig. 3n), further

demonstrating the dynamic nature of AIS composition along neuronal development.

SCRIB is a bona fide AIS enriched protein

To identify novel AIS proteins, we focused on genes highly ranked in our DIV14

dataset but not previously reported at the AIS (Fig. 4a). Three of the top ranked hits *Wdr7*,

Scrib and *Wdr47* were selected for further analysis and to assess their subcellular

localization in neurons. WD repeat-containing protein 7 (WDR7) mediates V-ATPase

dependent vesicle acidification in kidney cell lines and neuroendocrine cells [36, 37]. In

neurons WDR7 may regulate synaptic vesicle acidification [38]. WDR47 is required for

neuronal polarization and axonal and dendritic development [39, 40]. The scaffold protein

SCRIB has been reported as an important regulator for apical dendrite development,

spine morphology, and synapse plasticity [41, 42, 43].

We first leveraged CRISPR/Cas9-based homology independent genome editing to

integrate spaghetti monster fluorescent protein with V5 tags (smFP-V5) into *Wdr7*, *Scrib*,

and *Wdr47* to create C-terminal fusion protein [44]. Cultured neurons were infected at DIV0

by AAV delivery and observed at DIV14. We found SCRIB was highly and specifically

enriched at the AIS, while WDR47 and WDR7 were present at the AIS, but not specifically

enriched there (Fig. 4b). Quantification of the smFP-V5 signal in the AIS reveals SCRIB is

17 ± 2.55 times higher in the AIS than in proximal dendrites, while WDR47 and WDR7 are

2.21 ± 0.42 and 1.12 ± 0.12 fold higher in the AIS than proximal dendrites, respectively

(Fig. 4c).

To further investigate AIS SCRIB, we treated neurons after endogenous tagging

(smFP-V5) of SCRIB using 0.5% TritonX-100 to reveal the detergent-resistant pool of SCRIB

(Fig. 5a). The detergent insoluble SCRIB was retained and highly enriched at the AIS where

it colocalized with the AIS cytoskeletal protein βIV-spectrin; these observations are

consistent with previous studies showing that AIS proteins are resistant to detergent

extraction [45, 46]. Some AIS proteins, including voltage gated sodium channels, βIV-spectrin,

NFASC and AnkG have a periodic spacing in the AIS with of approximately 180~190 nm [47,

48]. To test whether SCRIB also exhibits this pattern, we used stimulated emission

depletion (STED) nanoscopy to image smFP-V5-tagged SCRIB. We found that AIS SCRIB

has a periodicity of 189.4 ± 6.6 nm in DIV16 neurons (Fig. 5b). Together, these results

suggest a strong association of SCRIB with previously described components of the AIS

periodic cytoskeleton [47].

To determine whether SCRIB localizes to the AIS *in vivo*, we performed

intraventricular injection in Cas9 transgenic mice using AAV to express gRNA to target

endogenous mouse *Scrib* and introduce smFP-V5 into the c-terminus of SCRIB. We

confirmed the gRNA targeting mouse *Scrib* efficiently resulted in detection of AIS SCRIB in

cultured mouse hippocampal neurons (Supplementary Fig. 5a). Three weeks after

injection of AAV into P0 Cas9 mouse pups, we found strong V5 immunolabeling of AIS

that colocalized with βIV-spectrin (Fig. 5c). Thus, SCRIB is present at the AIS both *in vitro*

and *in vivo*.

To further confirm SCRIB localization at the AIS, we used commercial antibodies for

SCRIB labeling. We used 2 different antibodies targeting different antigens of human

SCRIB (amino acids 1100-1400 and 1568-1630; see methods). Consistent with our *Scrib*

knock-in results, antibody labeling also showed strong and specific SCRIB enrichment at

the AIS with and without detergent extraction (Fig. 6a and Supplementary Fig. 5b). We

validated the specificity of the immunostaining using CRISPR/Cas9 mediated *Scrib* knock-

out. We infected cultured neurons using AAV to express an HA-tag and three gRNA

targeting *Scrib,* or AAV expressing HA and template; transduction was performed at DIV0

and neurons were fixed 14 days later. SCRIB was still enriched in the AIS after transduction

with AAV expressing the template, but AIS labeling was lost in neurons transduced with

the *Scrib* gRNAs (Fig. 6b). We generated two different AAVs with 3 gRNA each targeting

Scrib. Quantification showed a ~75% and 82% reduction in AIS SCRIB positive neurons

after transduction with these AAVs (SCRIB positive AIS: 88.9% ± 1.5% for control, 13.7% ±

6.8% for *Scrib* gRNA1, and 6.7% ± 2.7% for *Scrib* gRNA2; Mean ± SEM, one-way ANOVA)

(Fig. 6c). In addition, loss of SCRIB did not affect AnkG clustering. We found AIS AnkG was

comparable in the presence or absence of SCRIB (Figs 6b, d). Finally, using the validated

antibodies, we found SCRIB is enriched in the AIS of neurons in cerebral cortex (Fig. 6e).

These results show that SCRIB is an AIS protein.

AnkG recruits SCRIB to the AIS

To determine when SCRIB becomes clustered at the AIS, we analyzed its expression

and distribution along neuronal development. At DIV3 we found AnkG was highly

enriched in the proximal axon, while SCRIB was not (Fig. 7a). By DIV7 we found clustered

and enriched SCRIB that colocalized with AnkG at the AIS (Fig. 7b). Quantification of SCRIB

and AnkG enrichment throughout development showed that AnkG was enriched in the

proximal axon in 52.8% ± 10.3% neurons at DIV3 and in more than 90% neurons after

DIV7 (Fig. 7c). In contrast, AIS SCRIB was present in only 15.1% ± 5.8% of neurons at DIV3

and increased to 67.8% ± 7.4%, 75.8% ± 4.4%, and 93.5% ± 2.0% at DIV7, 14, and 21,

respectively (Fig. 7c). These results suggest that SCRIB clustering at the AIS follows AnkG

clustering.

Previous studies show that AnkG is the earliest AIS protein and is required for AIS

formation and maintenance [8, 16]. To determine if AnkG also contributes to AIS clustering

of SCRIB, we used CRISPR/Cas9 mediated knockout of AnkG. Neurons were infected with

AAV expressing template or *Ank3* triple gRNA at DIV0 and then fixed at DIV14. In control

conditions SCRIB colocalized with AnkG. However, AIS SCRIB was not detected in AnkG

deficient neurons (Fig. 7d). We found a ~68% decrease of AIS AnkG positive neurons in

Ank3 gRNA conditions (97.5% ± 1.4% for control; 29.4% ± 5.0% for *Ank3* gRNA conditions;

Mean ± SEM, p= 0.0002, unpaired t test) (Fig. 7e). Similarly, neurons with AIS SCRIB also

showed a ~63% decrease after loss of AnkG (94.5% ± 1.7% for control; 31.7% ± 5.8% for

Ank3 gRNA conditions; Mean ± SEM, p= 0.0005, unpaired t test) (Fig. 7f). These

observations suggest that AnkG is necessary for SCRIB clustering at the AIS.

To further define the relationship between AnkG and SCRIB, we performed co-

immunoprecipitation (IP) experiments using HEK293T cells transfected with flag-tagged

SCRIB (Scrib-Flag) and GFP-tagged AnkG (AnkG270-EGFP) or pEGFP-N1. Using anti-Flag

antibodies, we successfully co-immunoprecipitated AnkG270-EGFP (Fig. 7g). Similarly,

Scrib-Flag co-immunoprecipitated with AnkG-EGFP when the latter was

immunoprecipitated using anti-GFP antibodies (Fig. 7h). Together, these results suggest

that SCRIB interacts directly with AnkG.

To further define how SCRIB interacts with AnkG, we used various *Scrib* truncated

constructs omitting N-terminal 16 leucine-rich repeats (ΔLRR), 4 PDZ domains (ΔPDZ), or

spacer regions between the last LRR and the first PDZ (ΔIMR) (Fig. 7i); the SCRIB was also

fused to dTomato. AnkG270-EGFP and truncated *Scrib* were co-transfected in HEK293T

cells and AnkG270-EGFP was immunoprecipitated. We found that SCRIB lacking its IMR

amino acids failed to interact with AnkG (Fig. 7j). In addition, compared with dTomato-

Scrib-ΔPDZ, dTomato-Scrib-ΔLRR had lower binding to AnkG (Fig. 7j), suggesting that the

N-terminus of SCRIB is critical for AnkG binding.

DISCUSSION

The AIS plays key roles in regulating action potential initiation and maintenance of

neuronal polarity. Functional and structural changes to the AIS in normal and pathological

conditions are often associated with changes in its molecular composition. Here, we used

IPL-AIS to identify the AIS proteome. Our experiments identified nearly all previously

reported AIS components including extracellular matrix proteins, membrane proteins, and

cytoskeleton-associated proteins. Notably, most voltage-gated ion channels and their

accessory subunits were also identified, including Nav1.2, Navβ2, Kv2.1, Kv7.2, Kv1.2, Kvβ2,

and Cav2.1 and Cav2.2 (Fig. 2f). The results reported here complement and extend a

previous proximity labeling study using BioID [17] especially as it relates to cell surface

proteins, since the prior study was biased towards cytoplasmic proteins.

Although we used stringent thresholding, there are likely non-AIS proteins in our

dataset owing to non-specific absorption of antibodies, signal amplification after

biotinylation, and the large labeling radius of the horse-radish peroxidase. Nevertheless,

our ratiometric analysis against multiple controls allowed for the strong enrichment of AIS

proteins including AnkG, βIV-spectrin, NFASC, Nav1.2 and TRIM46 among others (Figs 2

and 3). The dispersed distribution of some other known AIS proteins might be attributed

to their wide distribution in neurons, varied protein abundance at the AIS, or differences

in labeling efficiency. In addition, our control experiments may also be useful in future

studies to examine the relative expression of proteins in different neuronal compartments

(AIS, axon, dendrites and soma).

The AIS composition may change during neuronal development and AIS maturation.

For example, recent studies revealed that NuMA1 and the P2Y1 purinergic receptor

participate only in AIS assembly [25, 49]. We applied TMT-based quantitative mass

spectrometry to identify developmental changes in the AIS proteome, revealing a gradual

increase of most AIS proteins, including the known AIS proteins βIV-spectrin, AnkG, NFASC

and voltage-gated ion channels; these observations are consistent with previous analyses

of AIS development using immunostaining methods [33]. While as many as 83.5% of proteins

identified exhibited the highest level of enrichment at DIV21, we observed some proteins

that deviated from this general trend. For example, enrichment of the AIS protein TRIM46

reached its maximum at DIV7. This observation is in accordance with the proposed role of

TRIM46 on establishing early neuronal polarity and axon specification [13].

As an example of the utility of our approach to identify new AIS proteins, we

examined more closely the top three candidates that were not previously reported on the

AIS. Among them, we found SCRIB (ranked 8) is highly and specifically enriched at the AIS,

although all three showed some degree of enrichment. We further confirmed SCRIB's AIS

localization by 1) smFP-V5 tagging of endogenous SCRIB using CRISPR, 2) immunostaining,

3) knock-out, 4) detergent resistance, 5) periodicity like that of other AIS proteins, and 6)

binding to and dependence on AnkG. In the future we will perform similar analyses on

other candidates that are highly enriched in the AIS proteome reported here.

SCRIB was previously reported to regulate neuronal migration, apical dendrite

development, axonal connectivity and synaptic plasticity[41, 43, 50]. *SCRIB* deletion in humans

causes severe dysmorphic features [51], and other pathogenic variants cause neural tube

defects and craniorachischisis [52, 53]. Similarly, SCRIB mutants lacking the last two PDZ and

the C-terminal domains die before or at birth with severe brain malformation[41].

Heterozygous or conditional knock-out mice also show psychomotor deficits and autism-

like behaviors [41, 50]. However, no report has described SCRIB at the AIS. We found that

SCRIB's AIS localization is AnkG-dependent. Although loss of SCRIB did not affect AnkG,

altered AnkG expression is associated with epilepsy, and psychiatric disorders, including

bipolar spectrum disorder and schizophrenia [8]. Since AnkG maintains SCRIB at the AIS, we

speculate that loss of SCRIB function might be a core molecular pathology of AnkG-related

neurodevelopmental disorders. Exploring SCRIB's function at the AIS both physiologically

and pathologically will be of considerable interest in future studies.

In conclusion, the results reported here illustrate the power of IPL-AIS to define

molecular complexes in distinct subcellular domains. Furthermore, the results reported

here are a rich resource to identify other AIS-enriched proteins that may contribute to AIS

structure and function.

METHODS

Animals

P0 Sprague-Dawley rat pups were purchased from Peking University Health Science Center. E18 pregnant Sprague-Dawley rats were purchased from Charles River Laboratories. Rat pups and embryos were used for the neuron culture. P7 wild-type C57BL/6 mice were purchased from Beijing Vital River Laboratory Animal Technology Co., Ltd. Transgenic Cas9 mice (JAX stock #027650) were obtained from The Jackson Laboratory. P0 Cas9 pups were used for neuron culture and intraventricular injection to tag endogenous *Scrib in vitro* and *in vivo*. All experimental procedures were performed in accordance with the guidelines of the Institutional Animal Care and Use Committees of Peking University and Baylor College of Medicine (IACUC #AN-4634).

Probe synthesis

Synthesis of compound **1**

To a solution of D-biotin (1.76 g, 7.2 mmol) and EDCI (1.54 g, 8.0 mmol) in 50 ml DMF,

N-hydroxysuccinimide (0.92 g, 8.0 mmol) was added at room temperature (RT). The

reaction mixture was stirred at 50°C overnight. DMF was removed under vacuum to a

residue of 5 ml. Add 100 ml cold ethanol into the residue during which time a white

precipitate formed. Precipitate was filtered and washed with 15 ml ethanol twice, then

dried in vacuum to afford compound **1**. The yield was 73%.

^{1}H-NMR (400 MHz, d_6-DMSO): 6.43 (1H, s), 6.37 (1H, s), 4.30 (1H, m), 4.15 (1H, m),

3.11 (2H, m), 2.84 (1H, dd), 2.81 (4H, s), 2.67 (2H, t), 2.60 (1H, d), 1.75-1.30 (6H, m).

Synthesis of compound **2**

Trimethylamine (1.60 g, 16 mmol) was added to a solution of mono-Boc protected

ethylene diamine (1.66 g, 10 mmol) and compound **1** (1.76 g, 5.2 mmol) in 50 ml DMF.

The reaction mixture was stirred at RT overnight and concentrated by rotary evaporation.

The residue was dissolved in a solution of DCM and ^{i}PrOH (v/v = 4:1) and washed with 20

ml 1 M HCl followed by 20 ml water three times. The organic layers were combined and

evaporated, and the residue was purified by column chromatography, affording

compound **2**. The yield was 72%.

^{1}H-NMR (400 MHz, d_6-DMSO): 7.79 (1H, t), 6.79 (1H, t), 6.43 (1H, s), 6.36 (1H, s), 4.30

(1H, m), 4.13 (1H, m), 3.05 (3H, m), 2.96 (2H, m), 2.81 (1H, dd), 2.59 (1H, d), 2.04 (2H, t),

1.68-1.41 (4H, m), 1.37 (9H, s), 1.34- 1.19 (2H, m).

Synthesis of **BP** and **BA1**

To a solution of compound **1** (0.50 g, 1 e.q.) and corresponding primary amine (1.1

e.q.) in 50 ml DMF, trimethylamine (3 e.q.) was added to the mixture and stirred overnight

at RT. The reaction solvent was evaporated and the residue was purified by C18 reverse

phase column (Waters XBridge Prep C18 5 μm OBD 19x150 mm) on semi-preparative

UPLC (Waters 2998 Photodiode Array Detector and 2545 Binary Gradient Module) with a

gradient of 3% to 60% methanol in water over 25 min. The overall yields were 60-70%.

^{1}H-NMR for **BP** (400 MHz, d_6-DMSO): 9.13 (1H, s), 7.79 (1H, t), 6.97 (2H, d), 6.66 (2H,

d), 6.42 (1H, s), 6.35 (1H, s), 4.31 (1H, m), 4.12 (1H, m), 3.18 (2H, dd), 3.08 (1H, m), 2.83

(1H, dd), 2.57 (3H, m), 2.03 (2H, t), 1.65-1.39 (4H, m), 1.35-1.19 (2H, m).

^{1}H-NMR for **BA1** (400 MHz, d_6-DMSO):7.78 (1H, t), 6.84 (2H, d), 6.49 (2H, d),6.43 (1H,

s), 6.36 (1H, s), 4.85 (2H, s), 4.31 (1H, m), 4.12 (1H, m), 3.16 (2H, dd), 3.09 (2H, dd),2.84

(1H, dd), 2.56 (3H, m), 2.03 (2H, t), 1.38-1.66 (4H, m), 1.28 (2H, m).

Synthesis of **BP2** and **BP3**

To a solution of D-biotin (1.00 g, 4.0 mmol), EDCI (0.85 g, 4.5 mmol) in 30 ml DMF,

corresponding phenol or aniline (0.56 g, 4.5 mmol) was added and the reaction mixture

was stirred at RT overnight. The reaction mixture was evaporated, and the residue was

purified by column chromatography, affording compound **L1**.

To the solution of **L1** in 20 ml DCM, BBr$_3$ (1.0 ml, 10 mmol) was added to the mixture

dropwise at 0°C. After being stirred at 0°C for 4 h, the mixture was heated to RT and

continued to react for 12 h. Then, 50 ml water was added to the mixture, and the residue

was washed by 10 ml cold water for three times, and purified by semi-preparative column

to get the purified product **BP2** or **BP3**. The whole yield was 25%.

^{1}H-NMR for **BP2** (400 MHz, d_6-DMSO): 9.60 (s, 1H), 9.13 (s, 1H), 7.35 (d, 2H), 6.67 (d,

2H), 6.45 (s, 1H), 6.37 (s, 1H), 4.31 (m, 1H), 4.14 (m, 1H), 3.14 (m, 1H), 2.84 (dd, 1H), 2.59

(d, 1H), 2.25 (t, 2H), 1.69-1.46 (m, 4H), 1.42-1.27 (m, 2H).

^{1}H-NMR for **BP3** (400 MHz, d_6-DMSO): 9.44 (s, 1H), 6.90 (m, 2H), 6.76-6.74 (m, 2H),

6.47 (s, 1H), 6.38 (s, 1H), 4.31 (m, 1H), 4.15 (m, 1H), 3.13 (m, 1H), 2.83 (dd, 1H), 2.58 (d,

1H), 2.53 (m, 2H), 1.71-1.47 (m, 4H), 1.45-1.34 (m, 2H).

Synthesis of **BP4**

Compound **2** (0.45 g, 1.2 mmol) was dissolved in 15 ml TFA/DCM (v/v= 1:1) and

stirred at RT for 2 h. The solvent was removed under vacuum and the residue was

dissolved in 20 ml DMF. To this solution was added 4-Hydroxybenzoic acid (0.15 g, 1.2

mmol), EDCI (0.45 g, 2.4 mmol), HOBt (0.35 g, 2.4 mmol), and TEA (0.36 g, 3.5 mmol). The

reaction mixture was stirred at RT overnight. Then, the reaction mixture was evaporated,

and the residue was purified by C18 reverse phase column (Waters XBridge Prep C18 5

μm OBD 19x150 mm) on semi-preparative UPLC (Waters 2998 Photodiode Array Detector

and 2545 Binary Gradient Module) with a gradient of 3% to 80% methanol in water over

30 min. The overall yields were 50%.

^{1}H-NMR (400 MHz, d_6-DMSO): 8.21 (t, 1H), 7.90 (t, 1H), 7.69 (d, 2H), 6.85 (d, 2H),

6.39 (d, 2H), 4.29 (dd, 1H), 4.14 (m, 1H), 3.03 (ddd, 1H), 2.80 (dd, 1H), 2.57 (d, 1H), 2.06

(t, 2H), 1.67-1.19 (m, 6H).

Primary neuron culture

Neurons were prepared from cortices or hippocampi of embryos and P0 pups.

Tissues were dissected and incubated in a digestion solution (0.25% trypsin and 0.4 mg/ml

DNase in Ca^{2+}/Mg^{2+} free HBSS) at 37°C for 15 min. Followed by the dissociation, neurons

were plated onto 100 μg/ml poly-D-lysine (Sigma) coated 10-cm dishes at a density of

100,000 cells/cm^2 or on coverslips coated by 20 μg/ml poly-D-lysine and 10 μg/ml laminin

mouse protein (Gibco) at a density of 40,000 cells/cm^2 in plating medium (high glucose

DMEM medium containing L-glutamine and 10% fetal bovine serum). After 3 h, the

medium was changed into neuronal culture medium (Neurobasal, B27 supplement,

Glutamax-I and penicillin-streptomycin) to maintain neuronal long-term growth. 2 μM

cytosine β-D-arabinofuranoside hydrochloride (AraC) was added after 6 days to slow

down glia cells proliferation. One third of culture medium was replaced with fresh

neuronal culture medium every seven days.

Optimization of anti-NFASC antibody directed AIS proximity labeling

Neurons were fixed with 4% formaldehyde for 15 min at RT. The free aldehyde group

was quenched by 0.25 M Glycine for 10 min. After washing four times with PBS,

endogenous peroxidase of cells was deactivated in 1.5% H_2O_2 as long as 1.5 h. Washed

samples were blocked by 5% fetal bovine serum dissolved in PBS containing 0.1% Tween-

20 for 1 h at RT. Then neurons were stained with 1, 2 or 5 μg/ml mouse anti-Pan-

Neurofascin (anti-NFASC, NeuroMab, Cat# 75-172, stock 1 mg/ml) dissolved in antibody

dilution solution (1% FBS with 0.1% Tween-20 in PBS) for 1 h at RT. Followed by washes

in washing solutions (0.1% Tween-20 in PBS), samples were labelled with 0.1, 0.2, 0.5 or

1 μg/ml HRP conjugated anti-mouse IgG (Cell signaling technology, Cat# 7076S, stock 1

mg/ml) for 1 h. Samples were washed for another four times with washing solution and

incubated with 6 different probes (see Supplementary Fig. 1e) individually at a

concentration of 500 μM for 10 min. The proximity labeling was triggered with the

addition of H_2O_2 (1, 100, 500 or 1000 μM). After 1, 5 or 10-min reaction, the proximity

labeling was stopped by adding quenching solutions (50 mM sodium ascorbate and 5 mM

Trolox dissolved in PBS). Endogenous bait protein NFASC and biotinylated proteins were

detected by immunocytochemistry to compare proximity labeling specificity and

efficiency through microscopy.

Immunoproximity labeling

The above optimized parameters were applied to anti-NFASC, anti-NeuN, anti-MAP2

and anti-SMI312 directed proximity labeling to capture AIS, soma, somatodendrites and

axon components. The blocked samples were stained with primary antibodies dissolved

in antibody dilution solution (1% FBS with 0.1%Tween-20 in PBS) for 1 h at RT. After

washing, samples were labelled with HRP conjugated second antibodies for 1 h and then

incubated with 500 μM biotin-phenol (BP) for 10 min. The proximity labeling was

triggered with the addition of 100 μM H_2O_2. After 1 min reaction, free radicals were

quickly quenched twice by quenching solutions (50 mM sodium ascorbate and 5 mM

Trolox dissolved in PBS). Samples were finally washed with PBS.

In the case of anti-AnkG and anti-TRIM46 targeted proximity labeling, parameters

including antibody dilutions and H_2O_2 concentration were also individually tested.

Antibodies used for proximity labeling were listed as follows. Primary antibodies: mouse

anti-Pan-Neurofascin (anti-NFASC, NeuroMab, Cat# 75-172), mouse anti-AnkG

(NeuroMab, Cat# 75-146), mouse anti-NeuN (Abcam, Cat# ab104224), mouse anti-MAP2

(Sigma-Aldrich, Cat# M1406), mouse anti-Neurofilament (anti-SMI312, Biolegend, Cat#

837904), chicken anti-AnkG (Synaptic Systems, Cat# 386006) and chicken anti-TRIM46

(Synaptic Systems, Cat# 377006). HRP conjugated second antibodies: anti-mouse IgG (Cell

signaling technology, Cat# 7076S) or anti-chicken IgY (Invitrogen, Cat# A16054; Abcam,

Cat# ab6877).

536   Protein lysis of cortical neurons after proximity labeling

Dimethyl experiments include three pairs: anti-NFASC vs. no primary antibody; anti-

NFASC vs. anti-NeuN; and anti-NFASC vs. anti-MAP2. Three times independent replicates

were performed in each pair. The 10-plex TMT DIV14 MS experiment include: three

biological replicates for anti-NFASC targeted proximity labeling; no primary antibody or

BP probe negative controls; reference samples of anti-NeuN, anti-MAP2 (duplicates), and

anti-SMI312 (duplicates). Due to differences in the bait abundance, references samples

of anti-NeuN, anti-MAP2 and anti-SMI312 yielded higher amounts of labeled proteins

than the anti-NFASC sample. To balance peptide loading to LC-MS/MS analysis, we used

only a fraction of the reference samples for biotinylated proteins enrichment. Anti-NF186,

no primary antibody and no substrate BP shared the same protein quantity for

enrichment. 1/2 protein quantity was used for anti-NeuN enrichment and 1/4 amount

was used for anti-MAP2 and anti-SMI312 enrichment respectively.

In AIS developmental TMT experiments, two parallel 10-plex TMT experiments were

performed and DIV14 samples were used as a reference for DIV7 and DIV21 samples.

Proximity labelled samples at DIV14 were equally divided and used for TMT1 and TMT2

pipeline. Five conditions were designed for each stage: anti-NFASC targeted AIS proximity

labeling (duplicates); no primary antibody (-1Ab); anti-NeuN targeted soma proximity

labeling and anti-MAP2 targeted somatodendrites proximity labeling. An amount of 1.5

mg proteins in samples of anti-NFASC and no primary antibody (DIV7, 14 and 21) were

used for enrichment, while 1/2 and 1/4 protein quantity were used for anti-NeuN and

anti-MAP2 enrichment individually.

Proximity labeling samples were lysed with lysis solution (50 mM Tris-HCl pH 7.6, 150

mM NaCl, 1% SDS, 0.5% sodium deoxycholate, 1% Triton X-100 and protease inhibitors

cocktail) on ice for 10 min. 500 µl lysis solution was added for each 10-cm dish. In the

condition of anti-NFASC, no primary antibody and no substrate BP, three 10-cm dishes in

one replicate were needed for protein lysis. While for anti-NeuN, anti-MAP2 and anti-

SMI312 conditions, one or two 10-cm dishes were needed. Protein lysis were de-

crosslinked at 99ºC for 1 h. The lysed solution was sonicated on ice and then centrifuged

at 12,000 rpm for 10 min at 4ºC to collect supernatant. Proteins were precipitated in cold

methanol at -80ºC.

Enrichment of biotinylated proteins and on-bead digestion

Precipitated proteins were washed twice using cold methanol. Purified proteins

were dissolved completely in 0.5% SDS (w/v). Protein concentrations were measured

using a BCA Protein Assay Kit to adjust protein quantity for enrichment. 1% of pre-

enrichment proteins were kept to test biotinylated signal via Western blotting or silver

staining to assess proximity labeling efficiency. The remaining proteins were incubated

with 200 µl streptavidin beads for 3 h with gentle rotation at RT. 5% of post-enrichment

proteins were kept to test the enrichment efficiency via Western blotting or silver staining.

Then the protein-beads mixture was sequentially washed twice by 2% SDS (w/v), 8 M urea

and 2 M sodium chloride. The mixture was reduced by 10 mM dithiothreitol and then with

20 mM iodoacetamide for alkylation. After washing four times with 100 mM TEAB,

proteins on beads were digested with the addition of 2 μg trypsin (Promega, Cat# V511A)

for 18 h at 37°C.

Immunoblotting and silver staining

Protein samples were heated at 99°C for 10 min with the addition of reducing

Laemmli SDS loading buffer. In the case of post-enrichment biotinylated protein samples,

2 mM biotin was added to elute biotinylated proteins. In the case of co-

immunoprecipitation (IP) experiments, AnkG270-EGFP or pEGFPN1 were co-transfected

with Scrib-Flag or its truncation variants in HEK293T cells and lysed after 48 h. The Scrib-

Flag plasmid was generated by amplifying human *Scrib* coding sequence from MSCV Puro

SCRIB WT (Addgene, Cat# 88886) and fused with 3xFlag tag at the C-terminal in pcDNA3

backbone. AnkG270-GFP was a kind gift from Dr. Vann Bennett (Duke University). The *Scrib*

truncation mutants dTomato-Scrib-ΔLRR, dTomato-Scrib-ΔIMR, and dTomato-Scrib-ΔPDZ

were kind gifts from Maya Shelly (Stony Brook University).

Input and immunoprecipitated samples were dissolved in Laemmli buffer (62.5 mM

Tris-HCl (pH 6.8), 2% SDS, 10% Glycerol, 2% 2-Mercaptoethanol, and 0.005% Bromophenol

blue) for immunoblotting. Proteins were loaded for electrophoresis in 5% SDS-PAGE

stacking gel. 10% or 6% SDS-PAGE separating gels were used for biotinylated samples and

co-IP samples respectively. The blots were blocked with 5% BSA at RT for 1 h and then

sequentially labeled using primary antibodies and HRP conjugated secondary antibodies.

Antibodies used were: rabbit anti-GFP (Invitrogen, Cat# A11122), chicken anti-GFP (Aves

Labs, Cat# GFP-1020), mouse anti-Flag (MBL, Cat# M185-3L), chicken anti-mCherry (Aves

Labs, Cat# MCHERRY-0020), rabbit anti-Flag conjugated with HRP (Cell Signaling, Cat#

86861), streptavidin-HRP (Invitrogen, Cat# 21124), goat anti-mouse IgG (H+L) conjugated

with HRP (Jackson, Cat# 115-035-146), goat anti-rabbit IgG (H+L) conjugated with HRP

(Jackson, Cat# 111-035-003), and goat anti-chicken IgY conjugated with HRP (Aves, Cat#

H-1004). The blots were developed by Clarity western ECL substrate (Bio-Rad, Cat#

1705060) and imaged in ChemiDoc MP Imaging System (Bio-Rad). In addition, the SDS-

PAGE gels could be directly used for silver staining using the Fast Silver Stain Kit (Beyotime,

Cat# P0017S).

Dimethyl labeling

Enriched proteins were digested with trypsin and the resulting peptides were treated

with isotope-encoded formaldehyde (heavy D^{13}CDO for anti-NFASC samples and light

HCHO for negative control/reference samples) and NaBH$_3$CN for their $-$NH$_2$ groups

methylation, resulting in a mass shift from 34.0631 Da to 28.0313 Da. For pairs of anti-

NFASC and no primary antibody, peptides were combined directly and desalted by C18

tips (ThermoFisher, Cat#87784). For pairs of anti-NFASC/ anti-NeuN or anti-NFASC/ anti-

MAP2, peptides were desalted first separately and then equal amounts of peptides were

combined. Peptides concentration was measured by Quantitative Colorimetric Peptide

Assay Kit (ThermoFisher, Cat# 23275). Combined peptides were dried in a vacuum

concentrator and ready for liquid chromatography-tandem mass spectrometry (LC-

MS/MS).

10-plex TMT labeling and peptides fractionation

Digested peptides were desalted by C18 tips (ThermoFisher, Cat#87784). In the case

of 14 DIV independent TMT experiments, desalted total peptides of anti-NF186, no

primary antibody and no substrate BP were directly used for TMT labeling. Taking anti-

NF186 peptides amount as reference, the same quantity of peptides from anti-NeuN,

anti-MAP2 and anti-SMI312 conditions were prepared in advance. Peptide concentration

was measured by Quantitative Colorimetric Peptide Assay Kit (ThermoFisher, Cat# 23275).

In the case of AIS developmental TMT experiments, sample amount was controlled when

performing enrichment. Therefore, these peptides were directly applied to TMT labeling.

TMT labeling was performed as instructed by TMT 10-plex Mass Tag Labeling Kits

and Reagents (ThermoFisher, Cat# 90110). Briefly, TMT reagents were maintained at RT

and reconstituted with 41 µl anhydrous acetonitrile for 0.8 mg vial of each tag. Peptides

were reconstituted in 10 µl 100 mM TEAB and dissolved completely. Add 12.5 µl TMT

solution into dissolved peptides and label for 2 h in the dark. The labeling in each

condition was described in Figs 2b and 3a. The reaction was quenched by adding 8 µl 5%

hydroxylamine and which was incubated for 15 min. Peptides from ten conditions were

mixed and dried in a vacuum concentrator. Mixed peptides were re-dissolved in 300 µl

0.1% trifluoroacetic acid and fractionated according to the instructions of Pierce High pH

Reversed-Phase Peptide Fractionation Kit (ThermoFisher, Cat# 84868). Samples were

eluted by gradient acetonitrile of 10.0%, 12.5%, 15.0%, 17.5%, 20.0%, 22.5%, 25.0%, 50.0%

solution in 0.1% triethylamine and defined as fraction 1-8 respectively. Fractions were

combined as pairs of '1+5', '2+6', '3+7' and '4+8' and dried again in vacuum concentrators.

Peptides were ready for LC-MS/MS.

Liquid chromatography and mass spectrometry

Peptides were reconstituted in 0.1% formic acid and separated in a loading column

(100 µm × 2 cm) and a C18 separating capillary column (75 µm × 15 cm) packed in-house

with Luna 3 µm C18(2) bulk packing material (Phenomenex, USA). A 2-h liquid

chromatography (LC) was applied to peptides separation. The LC gradient was held at 2%

for the first 8 minutes of the analysis, followed by an increase from 2% to 10% B from 8

to 9 minutes, an increase from 10% to 44% B from 9 to 123 minutes, and an increase from

44% to 99% B from 123 to 128 minutes (A: 0.1% formic acid in water and B: 80%

acetonitrile with 0.1% formic acid).

Samples were analyzed by Orbitrap Fusion LUMOS Tribrid Mass Spectrometer. The

precursors were ionized using an EASY-Spray ionization source (Thermo Fisher Scientific).

Survey scans of peptide precursors were collected in the Orbitrap from 350-1600 Th with

an advance gain control (AGC) target of 400,000, a maximum injection time of 50 ms, RF

lens at 30%, and a resolution of 60,000 at 200 m/z. In the case of HCD scans, an automated

scan range determination was enabled. An isolation window of 1.6 Th was used to select

precursor ions with the quadrupole. Product ions were collected in the Orbitrap with the

first mass of 110 Th, an AGC target of 50,000, a maximum injection time of 30 ms, HCD

collision energy at 30%, and a resolution of 15,000. Monoisotopic precursor selection was

enabled for peptide isotopic distributions, precursors of $z = 2$-7 were selected for data-

dependent MS/MS scans for 3 s of cycle time, and dynamic exclusion was set to 15 s with

a ±10 ppm window set around the precursor mono-isotope.

Mass spectrometry data analysis

Raw data files were loaded in MaxQuant software (version 1.6.10.43) and searched

against Rattus norvegicus proteomes from uniprot database (Taxonomy, 10116,

downloaded on Nov 15th, 2020). Trypsin was selected as the digestion mode. In the case

of searching dimethyl proteomics, the composition of Hx(6)C(2)H(-2) was replaced by

Hx(4)Cx(2) in heavy labels of DimethLys6 and DimethNter6. Cysteine acetylation

(carbamidomethyl) was a fixed modification. Methionine oxidation and protein N-

terminal acetylation were variable modifications. Re-quantify button in Misc. module and

match between runs in Identification module were activated. In the case of 10-plex TMT

proteomics, reporter ion MS2 was selected and 10-plex TMT was activated. Correction

factors were added to the tags based on the reagent instructions (ThermoFisher, Cat#

90110, Lot: UL291038). In both cases, the false discovery rate was set to 0.01.

Proteome data analysis

Anti-NFASC targeted AIS proximity labeling was the positive group. Samples of no

primary antibody or no substrate BP were negative controls for cutoff analysis, while anti-

NeuN, anti-MAP2 or anti-SMI312 targeted soma, somatodendrites or axon proximity

labeling were reference controls for ranking AIS candidates. Top ranked proteins would

have more possibility to be expressed or enriched in the AIS.

In the case of dimethyl experiments, contaminants and proteins identified as

reversed hits were removed. Proteins containing at least 2 unique peptides and quantified

in at least two out of three independent replicates were retained for further analysis (anti-

NFASC vs. anti-NeuN: 965; anti-NFASC vs. anti-MAP2: 802). Additionally, an H/L ratio > 1.5

was set as the cutoff for each replicate in the experiment of anti-NFASC versus no primary

antibody, which resulted in the identification of 704 proteins. Proteins in three pairs were

intersected and a total of 568 proteins were obtained. We normalized the H/L ratio

against the medium in each column using the formula of 'log$_2$(H/L)-medium log$_2$(H/L)' in

+NFASC/+NeuN and +NFASC/+MAP2 replicates. Averaged normalized data was obtained

from three replicates and used for distribution analysis.

In the case of DIV14 TMT experiments, contaminants and proteins identified as

reversed hits were removed. Proteins with at least two unique peptides were kept. Ratios

were calculated by division using their MS intensity. Parallel analysis was analyzed using

Pearson correlation. The average ratio of replicates was used for the following analysis.

The cutoff ratio was set at 2 for +NFASC/-BP and +NFASC/-1Ab. A total of 1403 proteins

were retained for further analysis. Ratios of +NFASC/+NeuN, +NFASC/+MAP2 and

+NFASC/+SMI312 were normalized against the medium ratio in their corresponding

column. Proteins were ranked based on average normalized log$_2$(+NFASC/+References,

fold change) in a descending order. The final overall rank was based on the average rank

scores against soma, somatodendrites and axon. Proteins in the top 200 against each

reference were overlapped and a total of 71 proteins were obtained. Gene Oncology

analysis of these 71 proteins was performed PANTHER overrepresentation test. A total of

2755 neuronal proteins in this TMT were used as a reference. Among them, 2744 proteins

were valid data in Gene Oncology database. Top 10 cellular compartment terms were

presented.

In the case of AIS developmental TMT experiments, two parallel 10-plex TMT labeling

were employed to analyze AIS developing (DIV7), mature (DIV14) and completely mature

(DIV21) proteome changes. Contaminants and proteins identified as reversed hits were

removed. Proteins with at least two unique peptides were kept. Ratios were calculated

by division using their MS intensity. Parallel analysis was analyzed using Pearson

correlation. The average ratio of replicates was used for further analysis. Proteins were

filtered with the standard of 'average $\log_2$(+NFASC)-$\log_2$(-NFASC)>3.5' for DIV7 and 21,

while for DIV14 the standard was set as 'average $\log_2$(+NFASC)-$\log_2$(-NFASC)>1'. This

setting was based on the endogenous biotinylated proteins distribution. Proteins with

intensity in +NFASC condition, not in -NFASC condition were also retained. Ratios of

+NFASC/+NeuN and +NFASC/+MAP2 were normalized against the medium ratio in their

corresponding column in each time point. Proteins were ranked based on average

normalized $\log_2$(+NFASC/+References, fold change) in a descending order. The final

overall rank was based on the average rank scores against soma and somatodendrites.

For analysis of AIS protein dynamics along neuronal development, biotinylated

proteins at DIV7, 14 and 21 together with prior DIV14 biotinylated proteins were

overlapped and revealed 549 common proteins in three stages. These proteins in the

condition of anti-NFASC at DIV7, 14 and 21 were normalized based on endogenous

biotinylated protein PCCA [54]. Among 549 proteins, 534 proteins were presented more

than 20% changes of average MS intensity in at least one pair of time points. Heatmap

was performed using cutree script in R package of pheatmap. AIS molecules identified in

three stages were performed Z-scored normalized abundance analysis. Significant

analysis and fold changes were analyzed regarding these AIS proteins according to a guide

of statistics for proteomics data analysis [55].

Immunocytochemistry and immunohistochemistry

For immunocytochemistry, neurons were fixed in 4% formaldehyde for 15 min at RT

and washed three times in PBS. Samples were blocked by 5% fetal bovine serum dissolved

in PBS containing 0.1% Tween-20 for 1 h at RT. Then neurons were sequentially stained

with primary antibodies and fluorescent second antibodies for 1 h at RT. Nuclei were

stained with DAPI (Invitrogen, Cat# D1306). In the case of detergent extraction

experiments, neurons were treated with 0.5% Triton X-100 in PBS for 5 min at RT before

fixation.

Primary antibodies used for immunocytochemistry in this study were: mouse anti-

Pan-Neurofascin (NeuroMab, Cat# 75-172, 1:1000), mouse anti-AnkG (NeuroMab, Cat#

75-146, 1:1000), chicken anti-AnkG (Synaptic Systems, Cat# 386006, 1:500), guinea pig

anti-AnkG (Synaptic Systems, Cat# 386004, 1:1000), mouse anti-NeuN (Abcam, Cat#

ab104224, 1:1000), mouse anti-Neurofilament (Biolegend, Cat# 837904, 1:1000), mouse

anti-MAP2 (Sigma-Aldrich, Cat# M1406, 1:1000), chicken anti-MAP2 (Abcam, Cat# ab5392,

1:10000), chicken anti-MAP2 (EnCor Biotechnology, Cat# CPCA-MAP2, 1: 2000), chicken

anti-TRIM46 (Synaptic Systems, Cat# 377006, 1:500), rabbit anti-βIV-spectrin (laboratory

of Dr. Matthew Rasband in BCM, 1:1000), rabbit anti-SCRIB (Abclonal, Cat# A17450, 1:50),

rabbit anti-SCRIB (Invitrogen, Cat# PA5-28628, 1:100), mouse anti-V5 (Invitrogen, Cat#

R960CUS, 1:500) and rat anti-HA (Millipore Sigma, Cat# 11867423001, 1:1000).

Fluorescent second antibodies were: goat anti-mouse IgG2a, Alexa Fluor 488 (Invitrogen,

Cat# A21131); goat anti-mouse IgG (H+L), Alexa Flour 488 (Invitrogen, Cat# A11029), 568

(Invitrogen, Cat# A11031), and Plus 594 (Invitrogen, Cat# A32742); goat anti-rabbit IgG

(H+L), Alexa Flour 488 (Invitrogen, Cat# A11034), and 568 (Invitrogen, Cat# A11036); goat

anti-rat IgG (H+L), Alexa Flour 594 (Invitrogen, Cat# A11007); goat anti-chicken IgY (H+L),

Alexa Flour 488 (Invitrogen, Cat# A11039), 647 (Invitrogen, Cat# A21449), and AMCA

(Jackson, Cat# 103-155-155). In the case of proximity labeling samples, biotinylated

protein signal was detected by incubation with streptavidin conjugated with Alexa Fluor

568 (Invitrogen, Cat# S11226) or 647 (Invitrogen, Cat# S21374).

For immunohistochemistry, brain sections of P7 C57BL/6 mice or P23 Cas9 mice

tagged with smFP-V5 to *Scrib* were used for the staining. Cortical sections were blocked

for 2 h at RT in 10% fetal bovine serum dissolved in PBS containing 0.1% TritonX-100 and

then incubated with first antibodies at 4°C overnight. After washing, slices were incubated

with second antibodies and counterstained with DAPI. First antibodies used were: mouse

anti-AnkG (NeuroMab, Cat# 75-146, 1:500), rabbit anti-SCRIB (Abclonal, Cat# A17450,

1:50), mouse anti-V5 (Invitrogen, Cat# R960CUS, 1:500), and rabbit anti-βIV-spectrin

(laboratory of Dr. Matthew Rasband in BCM, 1:500). Second antibodies used were: goat

anti-mouse IgG (H+L), Alexa Flour 568 (Invitrogen, Cat# A11031), goat anti-rabbit IgG

(H+L), Alexa Flour 488 (Invitrogen, Cat# A11034), and 568 (Invitrogen, Cat# A11036), and

goat anti-mouse IgG2a, Alexa Fluor 488 (Invitrogen, Cat# A21131). Sections were

mounted to glass slides with Fluoromount-G (SouthernBiotech, Cat# 0100-01).

Images were acquired on an inverted fluorescence microscope (Nikon-TiE, Japan)

equipped with a 40x1.3 NA oil immersion objective lens, confocal laser lines (OBIS 405,

488, 532, 561 and 637 nm, Coherent, USA), spinning disk confocal unit (Yokogawa CSU-

X1, Japan) and scientific CMOS cameras (Hamamatsu ORCA-Flash 4.0 v2). The microscope,

camera and lasers were controlled with a customer-built software written in LabVIEW

(National Instruments). Sections were imaged in the focal position or in z-stacks with a

0.5 μm step size at a resolution of 1500x1000 pixels. Z-projection was obtained in Fiji-

ImageJ software. Figures were adjusted and prepared in Fiji-ImageJ and Adobe Photoshop

CC2018.

V5 tag knock-in and microscopy

V5 tag knock-in sgRNA and donor constructs were generated following previously

described strategies [44, 56]. The specific target sequences of guide RNA (gRNA) were

791 designed for knock-in: rat *Scrib* 5'-GGGAAGCACCTGGCCCTAGG-3'; rat *Wdr47* 5'-

792 TCTGGACTTACAGTGGCTAG-3'; rat *Wdr7* 5'-ACGGCAGTCAGACCATGAAG-3'; mouse *Scrib*

793 5'-GTCCCGGCCGCACAGACCGA-3'. Plasmids of gRNA or AAV-SpCas9 (Addgene# 60957, Dr.

794 Feng Zhang) together with pUCmini-iCAP-PHP.S (Addgene# 103002, Dr. Viviana Gradinaru)

795 and pHelper (Cat #240071, Agilent Technologies) were used for AAV production. Adeno-

796 associated virus (AAV) cell-lysates were produced using the AAVpro Purification Kit (All

797 Serotypes) (Takara) with slight modifications. Briefly, HEK293T cells were transfected with

798 AAV plasmid, helper plasmid, and serotype PHP.S plasmid and collected after three days.

799 The cells were lysed using the AAV Extraction Solution A plus and B, and this crude AAV

800 solution was used for neuronal *in vitro* transduction. An aliquot of 20 µl AAVs (10 µl of

801 Cas9 and 10 µl of knock-in sgRNA and donor) was added to a well of 12-well plates at DIV0.

802 The medium was replaced after two days of infection. Neurons were fixed at the indicated

803 time points for immunocytochemistry. Concentrated AAV solution of smFP-V5 knock-in

804 to mouse *Scrib* was used for AAV intraventricular injection on P0 Cas9 mice pups. 50~100

805 µl AAV was concentrated from 2x15-cm dishes and 2 µl was used for each hemisphere.

806  Imaging was performed using a Zeiss AxioImager Z2 fitted with apotome for

807 structured illumination. Images were captured using the Zeiss ZEN software. Super-

808 resolution images were performed by Stimulated Emission Depletion (STED) microscope

809 using a STEDyCON (Abberior) system fitted to a Nikon Eclipse Ti2 microscope.   The

810 periodicity of SCRIB was measured in Fiji-ImageJ after generating a line scan across the

811 AIS.   The distance between peak fluorescence intensities (maximum to maximum) was

812 then measured.

814 **Genes knock-out and imaging by microscopy**

815  Triple gRNAs for rat *Scrib* and *Ank3* were designed: *Scrib* gRNA1 (5'-

816 GCTATTGAACTTGCGGAAGT-3', 5'-CCAGGCATACCAGCCGCCGC-3', and 5'-

817 CTGTGAGGATCAGTTCCGAG-3'); *Scrib* gRNA2 (5'-AGCCACAGCTCCCGTAGGTT -3', 5'-

GGCTAACTTCATGCAACTGG-3', and 5'-TTATGCTCTCAGGTATCTCG-3'); *Ank3* gRNA (5'-

CTGCTCGAGAACGACACGAA -3', 5'-CGCTCGGTTTAACAGCAACG -3', and 5'-

CTTCACGCCGCTGTATATGG-3'). The plasmid backbone consists of a cassette of synapsin1

promoted tandem HA tag as indicator for transfection. Crude AAV virus were generated

by co-transfection of AAV plasmid, helper plasmid, and serotype PHP.S plasmid in

HEK293T cells. An aliquot of 20 µl AAVs (10 µl of Cas9 and 10 µl of knock-out sgRNAs) was

added to a well of 12-well plates at DIV0. Neurons were fixed at DIV14 and stained for

imaging using a Zeiss structured illumination microscope with apotome.

Image analysis:

Cortical neurons of DIV10-14 were processed for optimizing AIS proximity labeling

parameters. AIS was defined by endogenous NF186. Biotinylated proteins signal in the AIS

and soma was quantified. Images were taken using the same parameter settings in all the

conditions of every experiment. Correct mean intensity (CMF) was obtained using mean

intensity in target regions extracted by the background of the same image. Labeling

efficiency was evaluated by the AIS CMF. Labeling specificity was evaluated by the ratio

of AIS CMF/ Soma CMF.

smFP-V5 tag knock-in to SCRIB, WDR47 and WDR7 samples at DIV14 were used for

polarity analysis. Neurons were infected at DIV0. The AIS was defined by βIV-spectrin

labeling. V5 signal was measured in the AIS and proximal dendrites (similar width). A 10-

pixel line was drawn along the AIS, or the proximal dendrites to measure the intensity

using Fiji-ImageJ software (NIH). Background was extracted and the polarity index was

evaluated by the formula AIS/Dendrite ratio= AIS CMF/ Dendrite CMF. At least three

independent experiments were performed.

DIV14 neurons were used for the integrated AnkG intensity measurement at the AIS.

Images were acquired in the focal position for quantification. A 10-pixel line was drawn

along the AIS using Fiji-ImageJ software. Integrated AnkG intensity was obtained and

normalized against control. Three times independent experiments were performed.

Statistical analysis

All statistical analyses were performed in GraphPad Prism 8. The statistical details

including the number of experiments, number of cells and statistical tests can be found

in figure legends. Statistical analysis was performed by two tailed t-test for two group

comparisons and by one-way ANOVA for multiple group comparisons. Graphs are

presented as the mean ± SEM. Differences were considered significant when p-values

were less than 0.05.

DATA AVAILABILITY STATEMENT

Raw data of proteome files have been deposited in the ProteomeXchange database under

the accession code PXD045921.

859
860

REFERENCES

1. Eichel K, Shen K. The function of the axon initial segment in neuronal polarity. *Developmental biology* **489**, 47-54 (2022).

2. Kole MH, Stuart GJ. Signal processing in the axon initial segment. *Neuron* **73**, 235-247 (2012).

3. Grubb MS, Burrone J. Activity-dependent relocation of the axon initial segment fine-tunes neuronal excitability. *Nature* **465**, 1070-1074 (2010).

4. Kaphzan H, Buffington SA, Jung JI, Rasband MN, Klann E. Alterations in intrinsic membrane properties and the axon initial segment in a mouse model of Angelman syndrome. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **31**, 17637-17648 (2011).

5. Zhang W*, et al.* Formin Activity and mDia1 Contribute to Maintain Axon Initial Segment Composition and Structure. *Molecular neurobiology* **58**, 6153-6169 (2021).

6. Hedstrom KL, Ogawa Y, Rasband MN. AnkyrinG is required for maintenance of the axon initial segment and neuronal polarity. *The Journal of cell biology* **183**, 635-640 (2008).

7. Sobotzik JM*, et al.* AnkyrinG is required to maintain axo-dendritic polarity in vivo. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 17564-17569 (2009).

8. Huang CY, Rasband MN. Axon initial segments: structure, function, and disease. *Annals of the New York Academy of Sciences* **1420**, 46-61 (2018).

891   9.   Kaphzan H*, et al.* Genetic reduction of the α1 subunit of Na/K-ATPase corrects
892   multiple hippocampal phenotypes in Angelman syndrome. *Cell reports* **4**, 405-412
893   (2013).

895   10.   Sun X*, et al.* Selective filtering defect at the axon initial segment in Alzheimer's
896   disease mouse models. *Proceedings of the National Academy of Sciences of the*
897   *United States of America* **111**, 14271-14276 (2014).

899   11.   Leterrier C*, et al.* End-binding proteins EB3 and EB1 link microtubules to ankyrin G
900   in the axon initial segment. *Proceedings of the National Academy of Sciences of*
901   *the United States of America* **108**, 8826-8831 (2011).

903   12.   Harterink M*, et al.* TRIM46 Organizes Microtubule Fasciculation in the Axon Initial
904   Segment. *The Journal of neuroscience : the official journal of the Society for*
905   *Neuroscience* **39**, 4864-4873 (2019).

907   13.   van Beuningen SFB*, et al.* TRIM46 Controls Neuronal Polarity and Axon
908   Specification by Driving the Formation of Parallel Microtubule Arrays. *Neuron* **88**,
909   1208-1226 (2015).

911   14.   Bender KJ, Trussell LO. The physiology of the axon initial segment. *Annual review*
912   *of neuroscience* **35**, 249-265 (2012).

914   15.   Hedstrom KL*, et al.* Neurofascin assembles a specialized extracellular matrix at the
915   axon initial segment. *The Journal of cell biology* **178**, 875-886 (2007).

917   16.   Leterrier C. The Axon Initial Segment, 50Years Later: A Nexus for Neuronal
918   Organization and Function. *Current topics in membranes* **77**, 185-233 (2016).

920   17.   Hamdan H*, et al.* Mapping axon initial segment structure and function by
921   multiplexed proximity biotinylation. *Nature communications* **11**, 100 (2020).

18. Bar DZ, Atkatsh K, Tavarez U, Erdos MR, Gruenbaum Y, Collins FS. Biotinylation by antibody recognition-a method for proximity labeling. *Nature methods* **15**, 127-133 (2018).

19. Dopie J, Sweredoski MJ, Moradian A, Belmont AS. Tyramide signal amplification mass spectrometry (TSA-MS) ratio identifies nuclear speckle proteins. *The Journal of cell biology* **219**, (2020).

20. Li X*, et al.* Defining Proximity Proteome of Histone Modifications by Antibody-mediated Protein A-APEX2 Labeling. *Genomics, proteomics & bioinformatics* **20**, 87-100 (2022).

21. Chen Y*, et al.* Mapping 3D genome organization relative to nuclear compartments using TSA-Seq as a cytological ruler. *The Journal of cell biology* **217**, 4025-4048 (2018).

22. Hung V*, et al.* Proteomic mapping of the human mitochondrial intermembrane space in live cells via ratiometric APEX tagging. *Molecular cell* **55**, 332-341 (2014).

23. Loh KH*, et al.* Proteomic Analysis of Unbounded Cellular Compartments: Synaptic Clefts. *Cell* **166**, 1295-1307.e1221 (2016).

24. Fréal A*, et al.* Feedback-Driven Assembly of the Axon Initial Segment. *Neuron* **104**, 305-321.e308 (2019).

25. Torii T*, et al.* NuMA1 promotes axon initial segment assembly through inhibition of endocytosis. *The Journal of cell biology* **219**, (2020).

26. Ke M*, et al.* Spatiotemporal profiling of cytosolic signaling complexes in living cells by selective proximity proteomics. *Nature communications* **12**, 71 (2021).

954   27.   Zhou Y*, et al.* Expanding APEX2 Substrates for Proximity-Dependent Labeling of
955   Nucleic Acids and Proteins in Living Cells. *Angewandte Chemie (International ed in*
956   *English)* **58**, 11763-11767 (2019).

958   28.   Lind D, Franken S, Kappler J, Jankowski J, Schilling K. Characterization of the
959   neuronal marker NeuN as a multiply phosphorylated antigen with discrete
960   subcellular localization. *Journal of neuroscience research* **79**, 295-302 (2005).

962   29.   Wolf HK*, et al.* NeuN: a useful neuronal marker for diagnostic histopathology. *The*
963   *journal of histochemistry and cytochemistry : official journal of the Histochemistry*
964   *Society* **44**, 1167-1171 (1996).

966   30.   Dehmelt L, Halpain S. The MAP2/Tau family of microtubule-associated proteins.
967   *Genome biology* **6**, 204 (2005).

969   31.   Thompson A*, et al.* Tandem mass tags: a novel quantification strategy for
970   comparative analysis of complex protein mixtures by MS/MS. *Analytical chemistry*
971   **75**, 1895-1904 (2003).

973   32.   Eichel K*, et al.* Endocytosis in the axon initial segment maintains neuronal polarity.
974   *Nature* **609**, 128-135 (2022).

976   33.   Jones SL, Korobova F, Svitkina T. Axon initial segment cytoskeleton comprises a
977   multiprotein submembranous coat containing sparse actin filaments. *The Journal*
978   *of cell biology* **205**, 67-81 (2014).

980   34.   Bas Orth C, Schultz C, Müller CM, Frotscher M, Deller T. Loss of the cisternal
981   organelle in the axon initial segment of cortical neurons in synaptopodin-deficient
982   mice. *The Journal of comparative neurology* **504**, 441-449 (2007).

984   35.   Ogawa Y, Horresh I, Trimmer JS, Bredt DS, Peles E, Rasband MN. Postsynaptic
985   density-93 clusters Kv1 channels at axon initial segments independently of Caspr2.

The Journal of neuroscience : the official journal of the Society for Neuroscience **28**,
5731-5739 (2008).

36.   Merkulova M, Păunescu TG, Azroyan A, Marshansky V, Breton S, Brown D.
Mapping the H(+) (V)-ATPase interactome: identification of proteins involved in
trafficking, folding, assembly and phosphorylation. *Scientific reports* **5**, 14827
(2015).

37.   Crummy E, Mani M, Thellman JC, Martin TFJ. The priming factor CAPS1 regulates
dense-core vesicle acidification by interacting with rabconnectin3β/WDR7 in
neuroendocrine cells. *The Journal of biological chemistry* **294**, 9402-9415 (2019).

38.   Gowrisankaran S, Milosevic I. Regulation of synaptic vesicle acidification at the
neuronal synapse. *IUBMB life* **72**, 568-576 (2020).

1000

1001   39.   Chen Y*, et al.* Wdr47 Controls Neuronal Polarization through the Camsap Family
1002   Microtubule Minus-End-Binding Proteins. *Cell reports* **31**, 107526 (2020).

1003

1004   40.   Buijs RR*, et al.* WDR47 protects neuronal microtubule minus ends from katanin-
1005   mediated severing. *Cell reports* **36**, 109371 (2021).

1006

1007   41.   Moreau MM*, et al.* The planar polarity protein Scribble1 is essential for neuronal
1008   plasticity and brain function. *The Journal of neuroscience : the official journal of*
1009   *the Society for Neuroscience* **30**, 9738-9752 (2010).

1010

1011   42.   Roche JP, Packard MC, Moeckel-Cole S, Budnik V. Regulation of synaptic plasticity
1012   and synaptic vesicle dynamics by the PDZ protein Scribble. *The Journal of*
1013   *neuroscience : the official journal of the Society for Neuroscience* **22**, 6471-6479
1014   (2002).

1015

1016   43.   Szczurkowska J*, et al.* A Localized Scaffold for cGMP Increase Is Required for Apical
1017   Dendrite Development. *Cell reports* **31**, 107519 (2020).

1018

44. Gao Y*, et al.* Plug-and-Play Protein Modification Using Homology-Independent Universal Genome Engineering. *Neuron* **103**, 583-597.e588 (2019).

45. Garrido JJ*, et al.* A targeting motif involved in sodium channel clustering at the axonal initial segment. *Science (New York, NY)* **300**, 2091-2094 (2003).

46. Winckler B, Forscher P, Mellman I. A diffusion barrier maintains distribution of membrane proteins in polarized neurons. *Nature* **397**, 698-701 (1999).

47. Leterrier C, Potier J, Caillol G, Debarnot C, Rueda Boroni F, Dargent B. Nanoscale Architecture of the Axon Initial Segment Reveals an Organized and Robust Scaffold. *Cell reports* **13**, 2781-2793 (2015).

48. Xu K, Zhong G, Zhuang X. Actin, spectrin, and associated proteins form a periodic cytoskeletal structure in axons. *Science (New York, NY)* **339**, 452-456 (2013).

49. Zhang W, Bonadiman A, Ciorraga M, Benitez MJ, Garrido JJ. P2Y1 Purinergic Receptor Modulate Axon Initial Segment Initial Development. *Frontiers in cellular neuroscience* **13**, 152 (2019).

50. Ezan J*, et al.* Early loss of Scribble affects cortical development, interhemispheric connectivity and psychomotor activity. *Scientific reports* **11**, 9106 (2021).

51. Wells C*, et al.* First fetal case of the 8q24.3 contiguous genes syndrome. *American journal of medical genetics Part A* **170a**, 239-242 (2016).

52. Lei Y*, et al.* Mutations in planar cell polarity gene SCRIB are associated with spina bifida. *PloS one* **8**, e69262 (2013).

53. Robinson A*, et al.* Mutations in the planar cell polarity genes CELSR1 and SCRIB are associated with the severe neural tube defect craniorachischisis. *Human mutation* **33**, 440-447 (2012).

1051

1052   54.   Frankenfield AM, Fernandopulle MS, Hasan S, Ward ME, Hao L. Development and
1053   Comparative Evaluation of Endolysosomal Proximity Labeling-Based Proteomic
1054   Methods in Human iPSC-Derived Neurons. *Analytical chemistry* **92**, 15437-15444
1055   (2020).

1056

1057   55.   Aguilan JT, Kulej K, Sidoli S. Guide for protein fold change and p-value calculation
1058   for non-experts in proteomics. *Molecular omics* **16**, 573-582 (2020).

1059

1060   56.   Ogawa Y, Rasband MN. Endogenously expressed Ranbp2 is not at the axon initial
1061   segment. *Journal of cell science* **134**,   (2021).

1062

1063

ACKNOWLEDGMENTS

We thank Y. Li, G. Wang, F. Yuan and Y. Chen for helpful discussions. This work was supported by the Ministry of Science and Technology (2018YFA0507600, 2017YFA0503600), the National Natural Science Foundation of China (32088101, 21727806), and a grant from the National Institutes of Neurological Disorders and Stroke (NS122073 to M.N.R.). P.Z. is sponsored by Bayer Investigator Award. W.Z. was supported in part by the Postdoctoral Fellowship of Peking-Tsinghua Center for Life Sciences. We thank Ms. W. Zhou from the Analytical Instrumentation Center in Peking University for assistance with MS sample identification.

AUTHOR CONTRIBUTIONS

W.Z., P.Z. and M.N.R. conceived the project. W.Z., P.Z. and M.N.R. designed experiments. W.Z., Y.F., L.P., Y.O., X.D., A.R., X.Z., and M.S. performed experiments. W.Z. and P.Z. analyzed the data and wrote the paper with input from all authors. W.Z., P.Z. and M.N.R. revised the paper.

COMPETING INTERESTS

The authors declare no competing interests.

Wei et al.

FIGURES

1083

Figure 1. Immunoproximity labeling of AIS

a, Experimental scheme of IPL-AIS method. HRP is directed to the axon initial segment in

fixed cortical neurons through the specific binding of anti-NFASC antibodies (1Ab) and HRP

conjugated secondary antibodies (2Ab). HRP-mediated proximity biotinylation is

mediated by the addition of biotin-phenol substrates and hydrogen peroxide. Biotinylated

proteins are enriched via affinity purification, digested with trypsin, and identified via LC-

MS/MS analysis.

b, Biotinylated proteins colocalize with endogenous protein signals in the axon initial

segment, soma and somatodendrites. The proximity labeling was directed by anti-NFASC,

anti-NeuN and anti-MAP2 antibodies individually in DIV14 cortical neurons with one

minute of reaction time. Scale bar= 50 µm.

c, Streptavidin blots and silver staining of DIV12 cortical neuron lysates (pre-enrichment)

or streptavidin bead eluates (post-enrichment) in conditions of anti-NFASC, no primary

antibody (-1Ab), anti-NeuN and anti-MAP2 targeted proximity labeling. Samples of anti-

NFASC, -1Ab, anti-NeuN and anti-MAP2 were loaded as the ratio of 4:4:2:1.

d, Venn diagrams showing 568 overlapping biotinylated proteins from +NFASC/+NeuN,

+NFASC/+MAP2 and filtered +/-NFASC datasheet. Three independent dimethyl proteome

replicates were performed at DIV14 for each condition.

e, Scatterplot showing the enrichment of anti-NFASC captured biotinylated proteins over

soma (x-axis) and somatodendrites (y-axis) intensity in the reference of average

normalized $\log_2$(H/L ratios). Red dots indicate reported AIS components and gray dots are

proteins having no AIS information.

Figure 2. The axon initial segment proteome at DIV14.

a, Schematic of neurons biotinylated at the axon initial segment by targeting NFASC (red), soma by targeting NeuN (green), somatodendrites by targeting MAP2 (orange), and axon by targeting phosphorylated neurofilaments using SMI312 antibody (blue).

b, Design of 10-plex TMT for AIS quantitative proteomic experiments. Samples were

collected from at least three independent primary neuron cultures for each condition.

TMT tags were labelled on the dishes.

c, Workflow for AIS proteome analysis.

d, Correlation analysis of biological replicates showing high reproducibility. Pearson

correlation was applied to parallel analysis of 2377, 2691 and 2498 common elements in

anti-NFASC, anti-MAP2 and anti-SMI312 replicates, respectively.

e, Ranking of 1403 biotinylated proteins based on their averaged MS intensity ratios. Red

dots indicate previously reported AIS components and gray dots are proteins having no

AIS information.

f, Overall rank plot integrating three reference proteomes (e) listing potential AIS enriched

proteins in the top positions. Proteins are sorted by average rank scores from

+NFASC/+NeuN, +NFASC/+MAP2 and +NFASC/+SMI312 experiments. Red dots indicate

reported AIS components and gray dots are proteins having no AIS information. Some AIS

proteins are labelled.

g, Venn diagrams showing 71 high-confidence AIS proteins from the top 200 candidates

from +NFASC/+NeuN, +NFASC/+MAP2 and +NFASC/+SMI312 proteomes.

h, Top 10 cellular compartment terms. 71 high-confidence AIS candidates were applied to

Gene Ontology cellular compartment analysis. A total of 2744 valid proteins in this TMT

were used as a reference (2755 proteins identified and 2744 valid data in Gene Ontology).

Wei et al.

Figure 3. The axon initial segment proteome across development.

a, Experimental design to evaluate the AIS proteome during development using two parallel 10-plex TMT: DIV7 for early AIS, DIV14 for mature AIS and DIV21 for mature AIS.

b, Pearson correlation showing high reproducibility in anti-NFASC targeted AIS proximity

labeling for DIV7 cortical neurons (1532 common elements from 2 replicates).

c, Ranking DIV7 AIS biotinylated 1407 proteins according to their averaged MS intensity

ratios (+NFASC/+NeuN and +NFASC/+MAP2).

d, DIV7 overall rank plot listing potential AIS proteins. Proteins are sorted by the average

rank scores from +NFASC/+NeuN and +NFASC/+MAP2 experiments from (c). Some known

AIS proteins are labelled in the graph. Red dots indicate reported AIS components and

gray dots are proteins having no AIS information.

e, Pearson correlation showing high reproducibility in anti-NFASC targeted AIS proximity

labeling for DIV21 cortical neurons (1859 common elements from 2 replicates).

f, Ranking DIV21 AIS biotinylated 1738 proteins according to their averaged MS intensity

ratios (+NFASC/+NeuN and +NFASC/+MAP2).

g, DIV21 overall rank plot listing potential AIS proteins. Proteins are sorted by the average

rank scores from +NFASC/+NeuN and +NFASC/+MAP2 experiments from (f). Some known

AIS proteins are labelled in the graph. Red dots indicate reported AIS components and

gray dots are proteins having no AIS information.

h, Venn diagrams showing 549 common AIS candidates at DIV7, 14 and 21. The prior AIS

proteome at DIV14 was also included in this analysis.

i, Six clusters with distinct protein expression profiles were generated by pheatmap

analysis in R package along neuronal development.

j, The percentage of protein numbers in each cluster of the 534 analyzed proteins.

k, AIS protein expression patterns along neuronal development. Proteins are represented

by their gene names.

l-n, Volcano plots showing fold changes in AIS proteins between DIV7 and 14 (l), DIV 14

and 21 (m) or DIV7 and 21 (n). Red and blue points represent AIS proteins with significant

up-regulation and down-regulation respectively. Proteins are represented by their gene

names. Horizontal dashed lines indicate p= 0.05. Vertical dashed lines indicate the cutoff

of $\log_2$(fold change) = ±1.

Figure 4. Screening AIS candidates by tagging endogenous genes.

a, Top 50 ranked AIS candidates at DIV14. The fold change of proteins intensity in the AIS compared with soma, somatodendrites and the axon. Previously reported AIS proteins are shown in red, and the rest are shown in black. Proteins are represented by their gene names.

b, Representative images of smFP-V5 tag knock-in neurons for SCRIB, WDR47, and WDR7. Neurons were infected with two AAVs, one is for Cas9 expression and another for gRNA and donor expression for homology-independent knock-in at DIV0. Samples were fixed at DIV14 and labelled with βIV-spectrin (green, AIS) and V5 tag (red, targeted proteins). Scale bar= 20 μm.

c, Quantification of V5 tag mean intensity in the AIS versus in proximal dendrites from knock-in samples. Four independent experiments were performed for WDR7 (n= 23 neurons) and SCRIB (n= 15 neurons). Three independent experiments were performed for

WDR47 (n= 33 neurons). Data are shown as mean ± SEM.

Figure 5. Validation of SCRIB enrichment at the AIS *in vitro* and *in vivo* by knock-in.

a, Representative images of smFP-V5 tag knock-in to endogenous SCRIB. Live neurons were treated with or without 0.5% TritonX-100. Neurons were infected at DIV0. After 14 days, samples were immunolabeled for the V5 tag (red, endogenous SCRIB), βIV-spectrin (green, AIS), and MAP2 (blue, somatodendrites). Scale bar= 20 μm.

b, A representative STED image of smFP-V5 tagged SCRIB. Knock-in neurons were fixed at DIV16, and stained with V5 tag antibody. Regions indicated by red lines were used to generate the intensity profiles, and the average periodicities were calculated. Column is mean ± SEM. Scale bar= 1 μm.

c, Representative image of smFP-V5 tagged SCRIB in the cortex. P0 Cas9 pups were intraventricularly injected with mouse specific *Scrib* gRNA and donor AAV and sacrificed at postnatal day 23 (P23). Samples were stained for the V5 tag (green, endogenous SCRIB) and βIV-spectrin (red, AIS). Scale bar = 20 μm.

1190

1191

Figure 6. Validation of SCRIB enrichment at the AIS *in vitro* and *in vivo* by immunostaining.

a, Representative images of DIV14 neurons treated with or without 0.5% TritonX-100 extraction prior to fixation. Fixed hippocampal neurons were stained with SCRIB (red), AnkG (green, AIS), and MAP2 (blue, somatodendrites). Scale bar = 50 µm.

b, Representative images of SCRIB knock-out using the CRISPR-Cas9 system. Hippocampal neurons were infected at DIV0 with AAV to express *Scrib* triple gRNA or template plasmids;

both also harbor an HA expression cassette. Neurons were fixed at DIV14 and stained for

SCRIB (magenta), AnkG (green), HA (red), and MAP2 (blue). Arrows indicate the AIS. Scale

bar = 20 µm.

c, Quantification of the percentage of AIS SCRIB positive neurons. Seven independent

experiments were performed using two different *Scrib* triple gRNA to disrupt *Scrib*

expression. 30~50 transfected neurons in each condition for each experiment were

analyzed. Data are mean ± SEM. One-way ANOVA, p<0.0001.

d, Comparison of integrated AnkG intensity in the AIS 14 days after transduction with AAV

to disrupt *Scrib* expression. Three independent experiments were performed. Data are

mean ± SEM. Unpaired t test followed by Welch's correction; ns, not significant.

e, Coronal cortical sections from P7 mice were stained for SCRIB (green), AnkG (red) and

nuclei (DAPI, blue). Scale bar= 50 µm.

1211

1212

Figure 7. AnkG is required for SCRIB enrichment at the AIS.

a-b, Representative images of DIV3 (a) and DIV7 (b) hippocampal neurons stained for

SCRIB (green), AnkG (red), and MAP2 (blue). Arrows indicate the proximal axon (a) and

AIS (b). Scale bar= 20 μm.

c, Quantification of the percentage of neurons with AIS SCRIB and AnkG at DIV3, 7, 14 and

21. Four independent experiments were performed at each time point. Data are mean ±

SEM.

d, Representative images of AnkG knock-out using the CRISPR-Cas9 system. DIV0

hippocampal neurons were transduced with *Ank3* triple gRNA or template plasmids; the

plasmids also harbor an HA expression cassette. Neurons were fixed at DIV14 and stained

for SCRIB (magenta), AnkG (green), HA (red), and MAP2 (blue). Scale bar= 20 μm.

e-f, Quantification of the percentage of neurons with AIS AnkG (e) and SCRIB (f) after loss

transduction with AAV to disrupt AnkG expression. Three independent experiments were

performed with the number of neurons analyzed indicated on the figure. Data are mean

± SEM. Unpaired t test, p= 0.0002 (e) and p= 0.0005 (f).

g-h, Co-immunoprecipitation of Scrib-Flag with AnkG270-EGFP. Scrib-Flag and AnkG270-

EGFP or pEGFPN1 were co-transfected in HEK293T cells and immunoprecipitated by Flag

antibody (g) or GFP antibody (h). IP, immunoprecipitation. IB, immunoblotting. Rb-GFP,

rabbit anti-GFP. Rb-Flag, rabbit anti-Flag. Ms-Flag, mouse anti-Flag. Ck-GFP, chicken anti-

GFP.

i, Illustration of the full length and SCRIB truncation constructs used.

j, Co-immunoprecipitation of truncated SCRIB with AnkG270. IP, immunoprecipitation. IB,

immunoblotting. Ck-GFP, chicken anti-GFP. Ck-mCherry, chicken anti-mCherry. Rb-GFP,

rabbit anti-GFP.

1237

Supplementary Figure 1. Optimization of anti-NFASC directed proximity labeling.

a-f, Anti-NFASC antibody directed proximity labeling parameters were tested in DIV10-14

fixed cortical neurons. Tested parameters are as follows: first antibody (1Ab, a), HRP

conjugated second antibody (2Ab, b) and H_2O_2 concentration (c), reaction duration (d) and

reaction substrates (f). The stock concentration of 1Ab and 2Ab is 1 mg/ml. (e) Chemical

probes tested in this study. The labeling efficiency was evaluated by the mean intensity of

biotinylated proteins signal in the AIS. The labeling specificity was evaluated by the ratio

of biotinylated protein signal in the AIS against that in the soma. Each point represents

one neuron. Graphs are presented as mean ± SEM, one-way ANOVA, p-values are

indicated on the figure and ns, not significant.

1248

1249

1250 **Supplementary Figure 2. SMI312 directed proximity labeling.** Fixed DIV14 cortical

1251 neurons were processed for proximity labeling with or without SMI312 antibody.

1252 Biotinylated proteins are in red, endogenous proteins labelled by SMI312 is in green. Scale

1253 bar = 50 μm.

1254

Supplementary Figure 3. Proximity labeling and enrichment in DIV14 TMT experiments.
Streptavidin blots of DIV14 cortical neuron lysates (pre-enrichment) or streptavidin bead eluate (post-enrichment) from samples of anti-NFASC, NeuN, MAP2, SMI312 and no primary antibody (-1Ab) or substrate BP (-BP). Samples of anti-NFASC, -1Ab, -BP, anti-NeuN, anti-MAP2 and anti-SMI312 were loaded as the ratio of 3: 3: 3: 1: 1: 1, respectively.

Supplementary Figure 4. The AIS developmental proteome.

a-c, Workflow for DIV7 (a), 21 (b) and 14 (c) biotinylated protein analysis in anti-NFASC samples.

d, Venn diagram of anti-NFASC targeted biotinylated proteins between DIV7 and 14 in 10-

plex TMT1.

e-f, Analysis of DIV14 anti-NFASC targeted biotinylated proteins from two parallel TMT

experiments. (e) Correlation analysis of biological replicates showing high reproducibility

in anti-NFASC directed AIS proximity labeling from two 10-plex TMT experiments. (f)

Scatterplot showing the enrichment of anti-NFASC targeted biotinylated proteins against

soma (x-axis) and somatodendrites (y-axis). Red dots indicate reported AIS components

and gray dots are proteins having no AIS information.

1271

Supplementary Figure 5. SCRIB localizes at the AIS.

a, Representative image of smFP-V5-tagged SCRIB in Cas9 transgenic mouse neurons. Cortical neurons were infected at DIV0 with gRNA and donor AAV and fixed at DIV10. Samples were stained for the V5 tag (red, endogenous SCRIB), βIV-spectrin (green, AIS), and MAP2 (blue, somatodendrites). Scale bar = 20 μm.

b, Representative images of DIV14 rat neurons stained with the Invitrogen SCRIB antibody targeting amino acids 1568-1630. Hippocampal neurons were treated with or without 0.5% TritonX-100 before fixation and stained for SCRIB (green), AnkG (red, AIS), and MAP2 (blue, somatodendrites). Scale bar = 20 μm.