Myosin VIII and XI isoforms interact with *Agrobacterium* VirE2 protein and

help direct transport from the plasma membrane to the perinuclear region

during plant transformation

Nana Liu, Lan-Ying Lee, Fang-Yu Hsu, Yanjun Yu, Praveen Rao, and Stanton B. Gelvin[*]

Department of Biological Sciences, Purdue University, West Lafayette, IN 47907

*Corresponding author: Stanton B. Gelvin (gelvin@purdue.edu)

Short title: Myosins mediate plant transformation

NL, L-YL, and SBG designed the experiments; NL, L-YL, F-YH, YY, and PR carried out the

experiments; NL and SBG wrote the manuscript with the help of L-YL; SBG envisaged the

project and obtained funding support.

The author responsible for distribution of materials integral to the findings presented in this

article in accordance with the policy described in the Instructions for Authors

(https://academic.oup.com/plcell/pages/General-Instructions) is: Stanton B. Gelvin

(gelvin@purdue.edu)

Authors e-mail addresses:

NL: liu2333@purdue.edu

L-YL: lee34@purdue.edu

F-YH: hsuf@purdue.edu; g9680517@oz.nthu.edu.tw

YY: yyj74@163.com

PR: Praveen_Rao@URMC.Rochester.edu

SBG: gelvin@purdue.edu

ABSTRACT

Virulent *Agrobacterium* strains transfer single-strand T-DNA (T-strands) and virulence effector proteins into plant cells. VirE2, one of these virulence effectors, enters the plant cell and is thought to bind T-strands, protecting them from nuclease degradation and helping guide them to the nucleus. How VirE2 is trafficked inside the plant cell is not fully understood. Using bimolecular fluorescence complementation, *in vitro* pull-down, yeast two-hybrid, and *in vivo* co-immunoprecipitation assays, we found that VirE2 binds directly to the cargo binding domains of several myosin VIII family members, and to myosin XI-K. We observed reduced susceptibility of several *Arabidopsis* actin mutants and a myosin *VIII-1/2/a/b* quadruple mutant to *Agrobacterium*-mediated transformation. Expression of cargo binding domains of myosin VIII-1, VIII-2, VIII-A, or VIII-B in transgenic plants inhibits *Arabidopsis* root transformation. However, none of the myosin VIII proteins contribute to the intracellular trafficking of VirE2. Expression of myosin *VIII-2*, *-A*, *-B*, but not *VIII-1*, cDNAs in the myosin *VIII-1/2/a/b* mutant partially restored transformation. Furthermore, functional fluorescently-tagged VirE2, synthesized in plant cells, relocalized from the cellular periphery into the cytoplasm after delivery of T-strands from *Agrobacterium*. Surprisingly, mutation of myosin *XI-k* and expression of the myosin XI-K cargo binding domain had no effect on transformation, although it blocked VirE2 movement along actin filaments. We hypothesize that myosin VIII proteins facilitate VirE2 tethering to the plasma membrane and are required for efficient localization of VirE2 to membrane sites from which they bind incoming T-strands. Myosin XI-K is important for VirE2 movement through the cytoplasm towards the nucleus.

INTRODUCTION

The soil-borne phytopathogen *Agrobacterium tumefaciens* is the causative agent of crown gall disease in plants (Gelvin, 2000; Tzfira et al., 2004). The ability of *Agrobacterium* to genetically transform plant cells has made *Agrobacterium*-mediated transformation the most widely used platform for generating transgenic plants (Gelvin, 2017). During transformation, T-DNA, a defined segment of the resident tumor inducing (Ti) plasmid, is transferred to the plant cell along with several virulence (Vir) effector proteins (Gelvin, 2003; McCullen and Binns, 2006; Lacroix and Citovsky, 2013). Virulence effector proteins VirD5, VirE2, VirE3, and VirF, along with VirD2, which covalently links to T-strands (single-strand T-DNA molecules), are transferred from *Agrobacterium* to the plant through a type IV secretion system (T4SS) (Cascales and Christie, 2004; Christie et al., 2005).

VirE2, the most abundant of the virulence effector proteins (Engstrom et al., 1987), is important for efficient plant transformation as *virE2* mutant *Agrobacterium* strains are highly attenuated in virulence (Stachel and Nester, 1986; Rossi et al., 1996). *In vitro*, VirE2 binds to single-strand (ss)DNA in a sequence-independent manner and has been proposed to interact with single-strand T-strands, linked to VirD2, in the plant cytoplasm, forming a "T-complex" and protecting T-strands from nuclease degradation (Citovsky et al., 1989; Sen et al., 1989; Howard and Citovsky, 1990; Yusibov et al., 1994; Rossi et al., 1996; Gelvin, 1998). VirE2 does not interact with T-strands in *Agrobacterium* (Cascales and Christie, 2004) and its interaction with T-strands to form a T-complex *in planta* has only been inferred. In addition to its *in vitro* DNA binding activity, VirE2 can form membrane channels which permit the transport of ssDNA through artificial membranes (Dumas et al., 2001; Duckely et al., 2005). Recent research showed that VirE2 is anchored on the host plasma membrane through interaction with VirE3 at *Agrobacterium*-host contact sites. Such membrane associations may facilitate the proposed interactions between VirE2 and T-strands as the T-strands enter the plant cell (Li et al., 2018, 2020).

Efficient plant transformation by a *virE2* mutant *Agrobacterium* strain (the "T-DNA donor") can be restored by co-inoculation with a bacterial strain lacking T-DNA but expressing VirE2 (the "VirE2 donor"; Otten et al., 1984; Simone et al., 2001), and a *virE2* mutant *Agrobacterium* strain can be complemented to full virulence by expression of a *VirE2* transgene *in planta* (Citovsky et al., 1992; Lapham et al., 2021), indicating that VirE2 is functional in plants during

transformation. However, the subcellular localization of VirE2 remains controversial. Early

reports demonstrated nuclear localization of VirE2 (Citovsky et al., 1992), but other studies from

several laboratories indicated that VirE2 localizes in the cytoplasm (Grange et al., 2008; Gelvin,

2010; Roushan et al., 2018). A recent report claimed that a small amount of VirE2 can enter the

nucleus, but only in the presence of VirD2 and T-strands (Li et al., 2020). How VirE2, and

potentially associated T-strands, are trafficked from their entry point at the cell periphery to the

nucleus remains unknown.

Myosins, motor proteins that convert chemical energy into directed movement of cargo

proteins along actin filaments, are key players in trafficking proteins and organelles in plants

(Vale, 2003; Richards and Cavalier-Smith, 2005). Two classes of myosins, myosin VIII and XI,

are encoded by plant genomes. *Arabidopsis thaliana* encodes 13 myosin XI group members and

four myosin VIII group members (Bezanilla et al., 2003; Foth et al., 2006). Generally, myosin

proteins contain a highly conserved N-terminal motor domain for actin binding, a neck domain

with a number of IQ repeats for light-chain binding, a coiled-coil domain that is responsible for

myosin protein dimerization, and a specific C-terminal tail domain that binds cargo (Trybus,

2008). Class XI myosins, known as organelle transporters, colocalize with organelles and are

involved in the rapid trafficking of Golgi stacks, streaming of endoplasmic reticulum (ER), and

cellular remodeling (Lee and Liu, 2004; Reisen and Hanson, 2007; Avisar et al., 2008b; Sparkes

et al., 2009; Ueda et al., 2010). In addition, myosins XI-K and XI-2 play major and overlapping

roles in root hair development (Peremyslov et al., 2008). In contrast to class XI myosins,

considerably less is known about the intracellular functions of class VIII myosins. Class VIII

myosins colocalize with plasmodesmata, the endoplasmic reticulum, and the plasma membrane

(Golomb et al., 2008) and may function in intercellular protein and RNA delivery to the

plasmodesmata, in endocytosis, and in anchoring actin filaments at plasmodesmata sites

(Reichelt et al., 1999; Wojtaszek et al., 2005; Avisar et al., 2008a; Golomb et al., 2008;

Sattarzadeh et al., 2008). Moreover, myosin VIII is important for the cell-to-cell movement of

virus proteins, and is required for viral movement protein targeting to and virus trafficking

through plasmodesmata (Amari et al., 2014).

The actin cytoskeleton functions to organize the endomembrane system and trafficking

patterns within the cell (Smith and Oppenheimer, 2005; Hussey et al., 2006). Cargo-binding and

intracellular transport by myosin is proposed to exert force on the actin filaments (Staiger et al.,

2009). Yang et al. (2017) reported that trafficking of VirE2 inside plant cells is powered by

myosin XI-K via the endoplasmic reticulum and F-actin filaments. However, our knowledge of

the roles of plant myosins in specific steps of VirE2 trafficking and transformation remains

sparse. In the present study, we investigated the roles of myosin VIII family members and

myosin XI-K in *Agrobacterium*-mediated transformation and in VirE2 intracellular movement.

Using a dominant-negative approach by overexpressing individual myosin cargo-binding

domains, and by overexpressing various individual full-length myosin cDNAs, we show that

Arabidopsis myosins VIII-2, VIII-A, and VIII-B, but not XI-K, are important for transformation.

These three myosin VIII isoforms interact with VirE2 and retain it at the periphery of root cells.

When T-DNA enters the plant cell, a portion of peripherally localized VirE2 relocalizes into the

cytosol, especially the perinuclear region. However, class VIII myosins do not affect the

movement of VirE2 along actin filaments within the cytoplasm. In contrast, myosin XI-K

supports the intracellular movement of VirE2. We propose that myosins VIII-2, VIII-A, and

VIII-B tether VirE2 to the plasma membrane. Entry of VirD2/T-strands into the plant cell

releases VirE2 from the membrane, following which VirE2 translocates to the perinuclear region

along actin filaments, powered by myosin XI-K.

RESULTS

The actin cytoskeleton is important for *Agrobacterium*-mediated transformation

The actin cytoskeleton is conserved in diverse eukaryotic organisms and is important for a

variety of plant developmental and other processes, including cell growth, division, and

expansion; organelle motility; organization of the cellular interior; endomembrane trafficking;

and host defense responses (Smith and Oppenheimer, 2005; Embley and Martin, 2006; Hussey et

al., 2006; Szymanski and Cosgrove, 2009; Peremyslov et al., 2010). To test whether the actin

cytoskeleton plays a role in *Agrobacterium*-mediated transformation, we conducted transient and

stable *Arabidopsis* root transformation assays using wild-type plants and plants individually

mutant in two root-expressed actin genes, *actin2* (*act2*) and *actin7* (*act7*). These actin mutant

plants showed substantial reductions (three- to five-fold) in transformation. Introduction of a

wild-type *Act7* cDNA into the *act7-4* mutant restored both transient and stable transformation

(Figure 1A). These results show that root-expressed actin genes are important for transformation

of *Arabidopsis* roots. We also conducted root transformation assays using a mutant of the

pollen-expressed *actin12* gene (*act12*) and the *botero* (*bot1*) mutant that shows disorganization

of the root cortical microtubule network (Bichet et al., 2001). These mutants showed wild-type

levels of transformation (Figure 1A), indicating that neither a pollen-expressed actin gene nor the

correct organization of root microtubules is essential for *Agrobacterium*-mediated root

transformation.

Yang et al. (2017) showed that the *Agrobacterium* virulence effector protein VirE2 uses

myosin XI-K to traffic through the plant cytoplasm. Given the importance of VirE2 for

transformation, we individually tested *Arabidopsis* mutants in each of the 13 myosin XI and four

myosin VIII gene family members for their susceptibility to transient and stable *Agrobacterium*-

mediated transformation. Figures 1B and C, and Supplemental Figure S1 show that, apart from

the *myosin XI-h* mutant, mutation of no other single myosin gene affected transformation. In

particular, mutation of the *myosin XI-k* gene did not decrease transformation. Because of

potential functional redundancy among the plant myosin VIII and XI family members

(Prokhnevsky et al., 2008), we also tested several higher order myosin VIII and myosin XI

mutants. Higher order *myosin XI-k* mutants (*XI-k/i*, *XI-k/1/2*, and *XI-k/1/2/i*) showed decreased

transformation susceptibility (Figures 1B and 1C). Similarly, the higher order myosin VIII

mutants *1/a*, *2/b*, *1/2/a*, and *1/2/a/b* showed reduced transformation susceptibility compared to

wild-type Col-0 plants (Figures 1B and 1C).

We compared root and above-ground plant growth of each of the higher order myosin XI and

myosin VIII mutants. Similar to what was previously observed (Ojangu et al., 2007; Peremyslov

et al., 2008), the myosin *XI-k/1/2/i* quadruple mutant displayed a strong growth and

developmental phenotype compared with Col-0 plants: the main roots had fewer lateral branches

and shorter root hairs, and the plants were dwarf. However, the myosin *VIII 1/2/a/b* quadruple

mutant appeared similar to Col-0 plants in growth, root development, and the ability to form calli

from cut root segments (Supplemental Figures S2A and S2B). Roots and the crown of the plant

are the natural target tissues for *Agrobacterium*-mediated transformation. Because we were

conducting quantitative root transformation assays (Gelvin, 2006), we did not continue to

investigate transformation susceptibility of the abnormal roots of the myosin *XI-k/1/2/i* quadruple

mutant. We therefore turned our attention to investigating the importance of the various myosin

VIII isoforms for transformation, and to the myosin XI-K protein.

Expression of full- length myosin *VIII-2*, *VIII-A*, and *VIII-B* cDNAs, but not a myosin *VIII-1* cDNA, increases susceptibility of the myosin *VIII-1/2/a/b* quadruple mutant to *Agrobacterium*-mediated transformation

To determine the importance of each *myosin VIII* gene for *Agrobacterium*-mediated root transformation, we individually expressed cDNAs for each *myosin VIII* gene in the *myosin VIII-1/2/a/b* quadruple mutant. We placed each *myosin VIII* cDNA under the control of a strong Cauliflower Mosaic Virus (CaMV) double 35S promoter (hereafter reported in Supplemental Table 1 as the *Agrobacterium* strains used to make the transgenic lines; At2361-At2164; At2360 was used as an empty vector control) or a β-estradiol-inducible promoter (Zuo et al., 2000; At2389-2392) and generated multiple transgenic lines. Because of the importance of myosins in trafficking molecules and organelles through the plant cytoplasm (Avisar et al., 2008b), we hereafter limited our transformation assays to transient transformation which measures T-DNA and virulence effector protein movement through the cytoplasm, entry into the nucleus, and expression of non-integrated transgenes. Thus, these assays predominantly measure non-integrated transgenes.

Transient root transformation assays of five independent transgenic lines of each *myosin VIII* cDNA indicated that individual expression of the *myosin VIII-2*, *VIII-A*, and *VIII-B* cDNAs increased the transformation susceptibility of the *myosin VIII-1/2/a/b* quadruple mutant (Figure 2A and Supplemental Figures S3B-D), indicating a role for these myosins in *Agrobacterium*-mediated transformation. However, expression of the *myosin VIII-1* cDNA in the quadruple mutant background did not increase transformation susceptibility (Figure 2A and Supplemental Figure S3A). This latter result suggests either that myosin VIII-1 does not contribute to transformation or that expression of myosin VIII-1 is not sufficient to increase transformation in the absence of the other three myosin VIII isoforms. To distinguish between these two possibilities, we individually overexpressed each *myosin VIII* cDNA in wild-type Col-0 plants. Interestingly, overexpression of the *myosin VIII-1* cDNA, but not the *myosin VIII-2*, *VIII-A*, or *VIII-B* cDNAs, increased susceptibility of Col-0 plants to *Agrobacterium*-mediated transformation (Figure 1E and Supplemental Figures S3A-D). These results indicate that myosin VIII-1 can contribute to transformation, but that expression of myosin VIII-1 is not sufficient to do so in the absence of the other myosin VIII isoforms. Transgenic plants harboring inducible

myosin VIII cDNAs yielded results similar to that of plants constitutively expressing these

cDNAs (Supplemental Figures S4A-D).

Finally, we generated transgenic lines that expressed two *myosin VIII* cDNAs (*VIII-A* and

VIII-B [At2375], or *VIII-2* and *VIII-B* [At2376]) in both the *myosin VIII-1/2/a/b* quadruple

mutant (Figure 2C) and in wild-type Col-0 plants (Figure 2D). Overexpression of these two sets

of myosin cDNAs did not increase transformation of otherwise wild-type plants. However,

expression of these pairs of myosin cDNAs in the *myosin VIII-1/2/a//b* quadruple mutant

enhanced transformation to a greater extent than did expression of these cDNAs individually

(Figures 2C and Supplemental Figures S5A-D). These results indicate that *myosin VIII-A*, *VIII-B*,

and *VIII-2* genes contribute synergistically to transformation. These results further suggest that

the level of expression of the *myosin VIII-2*, *VIII-A*, and *VIII-B* genes in wild-type roots is

sufficient for maximal transformation, and that increasing expression of these genes cannot

further increase transformation.

Inducible expression of *myosin VIII*, but not *myosin XI-K*, cargo binding domain transgenes

affects *Agrobacterium*-mediated transient root transformation

Yang et al. (2017) demonstrated that expression of a myosin XI-K cargo binding (globular tail)

domain inhibited VirE2 movement in tobacco leaf cells. To determine whether specific myosin

VIII members are important for *Agrobacterium*-mediated transformation using a similar

dominant-negative approach, we generated transgenic *Arabidopsis* lines individually expressing

cDNAs encoding each of the four myosin VIII cargo binding domains (CBD; At2315-2318), or

the myosin XI-K CBD (At2319), under the control of a β-estradiol inducible promoter. Because

these CBD constructs lack the myosin motor domain, they are not able to transport their cargoes

along actin filaments. We tested the susceptibility of 3-4 independent transgenic lines for each

CBD construct to *Agrobacterium*-mediated transient transformation, either after 24 hr transgene

induction or without induction. Induction of each of the four *myosin VIII* CBD transgenes

decreased transformation two- to three-fold, whereas induction of the *myosin XI-K* CBD

transgene had no significant effect on transformation (Figure 3 and Supplemental Figures S6A-

E). These data further suggest that the four myosin VIII proteins contribute to *Agrobacterium*-

mediated transformation, whereas myosin XI-K is not required for transformation. However,

below we show that myosin XI-K is important for VirE2 movement through the cytoplasm.

241   Myosin VIII proteins are required for *Arabidopsis* roots expressing VirE2 to complement

242   the virulence of a *virE2* mutant *Agrobacterium* strain, and to mitigate VirE2 aggregation

Agrobacterium virE2 mutants are severely debilitated in virulence (Rossi et al., 1996). However,

expression of VirE2 or VirE2-Venus transgenes in plants restores virulence, indicating that the

major role of VirE2 in transformation occurs in the plant (Citovsky et al., 1992; Bhattacharjee et

al., 2008; Lapham et al., 2021). To determine the importance of myosin VIII proteins for such

"*in planta*" complementation, we generated numerous *Arabidopsis* transgenic lines expressing a

VirE2-Venus transgene, under the control of a β-estradiol-inducible promoter, in both wild-type

and in *myosin VIII* quadruple mutant plants. Twenty four hr after incubation of roots of these

lines in medium with or without inducer, we cut the roots into small segments and infected them

with *A. tumefaciens* At1879 containing the T-DNA binary vector pBISN2 with a *gusA*-intron

gene in the T-DNA (Narasimhulu et al., 1996). Six days later we assessed transient

transformation efficiency by conducting β-glucuronidase (GUS) activity assays using the

chromogenic substrate 5-bromo-4-chloro-3-indolyl-β-D-glucuronide (X-gluc). As previously

reported (Lapham et al., 2021), induction of the *VirE2-Venus* transgene in wild-type Col-0 plants

complemented the loss of VirE2 function when a *virE2 Agrobacterium* mutant was used.

However, induction of the *VirE2-Venus* transgene in *myosin VIII* quadruple mutant plants could

not compensate the loss of VirE2 function when a *virE2 Agrobacterium* mutant was used

(Figures 4A and 4B). Thus, *Myosin VIII* expression is required for *in planta* complementation of

an *Agrobacterium virE2* mutant.

VirE2 usually localizes in the plant cytoplasm, although in the presence of the virulence

effector protein VirD2 and T-strands a small amount may enter the nucleus (Gelvin, 2010; Li et

al., 2020). To determine whether myosin VIII affects the subcellular distribution of VirE2, we

used confocal microscopy to visualize β-estradiol induced VirE2-Venus in both the Col-0 and

myosin VIII quadruple mutant backgrounds containing an inducible *VirE2-Venus* transgene.

VirE2 is known to aggregate both *in vitro* and in plants (Bhattacharjee et al., 2008; Dym et al.,

2008). Under our induction conditions VirE2-Venus showed a pattern of aggregation in the

myosin VIII quadruple mutant distinct from that seen in the Col-0 background. In roots of the

quadruple mutant, VirE2-Venus aggregated to a greater extent than it did in wild-type roots

(Figure 4C). The sizes of VirE2 aggregates from two random areas of each of six images were

measured and the average sizes were calculated. The average VirE2 aggregate size in root cells

of the *myosin VIII* quadruple mutant was 6.2 ± 0.6 μm^2, compared to an average aggregate size of

2.0 ± 0.3 μm^2 in wild-type roots (Figure 4D). Furthermore, the total cellular area occupied by

these VirE2 aggregates was greater in root cells of the *myosin VIII* mutant ($17.0\pm1.1\%$) than in

similar wild-type cells ($4.7\pm0.9\%$) (Figure 4D). To determine whether this extensive VirE2

aggregation in myosin *VIII* mutant cells resulted from higher levels of VirE2-Venus expression

in roots, we extracted total proteins from β-estradiol-induced roots of these plants and

determined the relative amounts of VirE2-Venus expressed in roots using anti-GFP antibodies.

As a loading control for the protein blots, we used an antibody targeting the house-keeping

protein phosphoenolpyruvate carboxylase (PEPC). Strikingly, myosin *VIII* quadruple mutant

roots express, on average, less VirE2-Venus than do wild-type roots (Figure 4E). Therefore, the

more extensive aggregation of VirE2-Venus in *myosin VIII* mutant roots is unique and cannot be

explained by greater accumulation of this protein.

To determine which myosin protein influences VirE2-Venus aggregation, we generated

numerous transgenic *Arabidopsis* lines (using *Agrobacterium* strains At2315-At2319), in the

myosin VIII quadruple mutant background, expressing both VirE2-Venus and individual

mCherry-myosin VIII CBD cDNAs under the control of a β-estradiol-inducible promoter.

Following induction, we measured the size of the VirE2-Venus aggregates in root cells and the

percentage of root cell volume taken up by these aggregates. Figures 5A and 5B show that,

compared to expression of VirE2-Venus in the absence of expression of a myosin CBD

transgene, expression of myosin VIII-A, VIII-B, and VIII-2 CBDs reduced both the size of

VirE2-Venus aggregates (from 7.5 ± 1.9 μm^2 down to 2.4 ± 0.3 μm^2, 3.7 ± 0.5 μm^2, and 3.1 ± 0.3

μm^2, respectively) and the percentage of the cell volume (from $17.7\pm2.0\%$ down to $3.6\pm0.6\%$,

$6.3\pm0.4\%$, and $6.1\pm0.5\%$, respectively) occupied by these aggregates, due to their dominant

negative effects. However, the CBD of myosin VIII-1 did not substantially reduce aggregate

size or the percentage of the cell volume occupied by the aggregates (Figure 5A): the average

aggregates size was 6.7 ± 0.7 μm^2, and the percentage of the cell volume was $11.8\pm1.1\%$. These

results suggest that myosin VIII-A, VIII-B, and VIII-2 CBDs interact with VirE2 in plant cells

and that the myosin VIII-1 CBD may fundamentally differ from the CBDs of the other three

myosin VIII members.

VirE2 interacts directly with some, but not all, myosin VIII cargo binding domains and

with the CBD of myosin XI-K

Because myosin VIII proteins are required for VirE2 to function in virulence, and because

myosin VIII CBD expression affects the degree to which VirE2 aggregates *in planta*, we

investigated whether myosin protein cargo binding domains interact with VirE2. We first tested

the direct interaction of VirE2 with each of the myosin VIII and the myosin XI-K protein CBDs

using an *in vitro* pull-down assay. We individually incubated recombinant VirE2-Venus with

recombinant myc-tagged myosin CBDs. As a control, we substituted recombinant Venus protein

for VirE2-Venus. As further controls, we incubated VirE2-Venus with myc-tagged VIP1, a

protein known to interact with VirE2 (Tzfira et al., 2001; Djamei et al., 2007), or myc-tagged

Lamin C, a protein known not to interact with VirE2 (Ueki and Citovsky, 2005). We captured

VirE2-Venus, or Venus, and any interacting proteins on anti-GFP antibody beads, subjected the

bound proteins to SDS gel electrophoresis, then conducted Western blot analysis using anti-myc

antibodies. Figures 6A-F show that the CBDs of myosins VIII-A, VIII-B, VIII-2, and XI-K

directly interacted with VirE2-Venus, but not with Venus. However, we were not able to detect

interaction of VirE2-Venus with the CBD of myosin VIII-1. As expected, VirE2-Venus

interacted with myc-tagged VIP1 but not with myc-tagged Lamin C (Figure 6). These data were

confirmed using a yeast two-hybrid system. VirE2 interacted in yeast with the CBDs of myosins

VIII-A, VIII-B, VIII-2, and XI-K, as well as with VIP1. However, VirE2 did not interact in

yeast with the CBD of myosin VIII-1 or with the negative control Lamin C (Figures 7A and 7B).

To assess the interactions of VirE2 with myosin VIII or XI-K CBD proteins *in vivo*, we

conducted co-immunoprecipitation experiments using transgenic *Arabidopsis* roots expressing

VirE2-Venus and, individually, each of the myc-tagged myosin VIII or the myosin XI-K CBDs.

Using anti-GFP antibody beads to capture VirE2-Venus and associated proteins followed by

immunoblotting using anti-myc tag antibody to detect CBDs, we observed robust co-

immunoprecipitation between VirE2-Venus and the myosin VIII-2, VIII-A, VIII-B, and XI-K

CBDs (~4% of the input myc-myosin VIII-2, VIII-A, VIII-B, and XI-K CBDs were co-

immunoprecipitated with anti-GFP antibody). However, we also observed some co-

immunoprecipitation between VirE2 and the myc-tagged myosin VIII-1 CBD (~1.6-2% of the

input myosin VIII-1 CBD; Figures 8A and 8B). As expected, VirE2-Venus interacted with VIP1

(Figure 8A), and the myosin VIII-2 CBD did not interact with the control Venus protein (Figure

8B). The *in vivo* co-immunoprecipitation of VirE2-Venus with the myosin VIII-2, VIII-A, VIII-

B, and XI-K CBDs recapitulated the *in vitro* interaction and yeast two-hybrid results. However,

in vivo co-immunoprecipitation of the myosin VIII-1 CBD with VirE2-Venus, in the absence of

interaction *in vitro* and in yeast, suggests that this *in vivo* interaction may be indirect and

mediated by other factors within *Arabidopsis* roots.

VirE2 and Myosin VIII cargo binding domains co-localize at the cell periphery

The vast majority of VirE2 protein localizes within the cellular cytoplasm, and in wild-type

Arabidopsis roots is often found in the peripheral regions of the cell (Shi et al., 2014). VirE2 has

been proposed to form membrane channels through which VirD2/T-strands translocate from

Agrobacterium into the plant cell (Dumas et al., 2001; Duckely et al., 2005). Myosin VIII

proteins also localize to the plasma membrane (Golomb et al., 2008). To determine whether

VirE2 and myosin cargo binding domains co-localize at the cell periphery, we generated

transgenic *Arabidopsis* plants expressing VirE2-Venus and various mCherry-tagged myosin

CBD cDNAs under the control of a β-estradiol inducible promoter (using *Agrobacterium* strains

At2315-At2319). After β-estradiol induction, we imaged roots using confocal microscopy

(Figures 9A and 9B). The CBDs of myosins VIII-A and VIII-2 localized almost exclusively to

the cellular periphery, whereas the CBD of myosin VIII-B localized both to the cellular

periphery and to the nucleus (Figure 9A). The myosin VIII-1 and XI-K CBDs localized not only

at the cellular periphery but also throughout the cytoplasm (Figure 9A).

When co-expressed in the Col-0 background, VirE2 and the myosin VIII-2, VIII-A, VIII-B,

and XI-K CBDs partially co-localized at the cell periphery (Pearson's correlation coefficient =

0.56±0.04, 0.68±0.05, 0.63±0.04, and 0.69±0.03, respectively; overlap coefficient = 0.65±0.03,

0.72±0.06, 0.70±0.03, and 0.77±0.04, respectively; Figure 9B). However, co-localization of

VirE2 with the myosin VIII-1 CBD at the cell periphery was less extensive (Pearson's

correlation coefficient = 0.44±0.03; overlap coefficient = 0.48±0.01; Figure 9B). Staining

VirE2-Venus expressing *Arabidopsis* roots with the plasma membrane dye FM4-64 revealed that

VirE2 localized near to but not in the plasma membrane (Pearson's correlation coefficient =

0.42±0.07; overlap coefficient = 0.45±0.03; Supplemental Figures S7A and B). However, in the

myosin VIII quadruple mutant background, VirE2 localized less extensively at the cellular

periphery and more within the general cytoplasm (Supplemental Figures S8A and B). Co-

expression of mCherry-myosin VIII-1, VIII-A, or VIII-2 CBDs with the plasma membrane

marker PIP2A-Venus (using *Agrobacterium* strains At2370, At2371, and At2374, respectively)

revealed that these myosin VIII CBDs co-localize with PIP2A (Supplemental Figures S9A-C),

similar to what has previously been reported (Golomb et al., 2008). Taken together, these results

suggest that the myosin VIII-2, VIII-A, and VIII-B CBDs localize to the plasma membrane

where they may temporarily tether VirE2.

VirE2 and several myosin VIII cargo binding domains interact at the cell periphery

In order to test whether VirE2 may be a cargo of myosin VIII proteins, we conducted

bimolecular fluorescence complementation (BiFC) assays to visualize *in planta* protein-protein

interactions. We generated transgenic *Arabidopsis* plants expressing VirE2-cYFP and nVenus-

tagged myosin CBD cDNAs under the control of a β-estradiol inducible promoter (using

Agrobacterium strains At2320-At2324). The plants also constitutively expressed mCherry-actin

binding domain 2 (ABD2) as an actin marker. After inducing the plants with β-estradiol, we

detected YFP fluorescence mainly at the cellular periphery of root cells when VirE2-cYFP was

co-expressed with the nVenus-tagged CBDs of myosins VIII-A, VIII-B, VIII-2, or myosin XI-K,

but no fluorescence with the myosin VIII-1 CBD (Figures 10A-F), indicating a lack of direct

interaction. VirE2-cYFP additionally interacted with the CBD of myosin XI-K in the cytoplasm.

VirE2-cYFP did not interact with the negative control protein Lamin C. These results again

suggest that the CBDs of myosin VIII-2, VIII-A, VIII-B, and XI-K may temporarily tether VirE2

to the plasma membrane, but that only myosin XI-K may transport VirE2 along actin filaments

through the cytoplasm.

Some VirE2 molecules re-localize to the perinuclear region during infection by a virulent

***Agrobacterium* strain**

Some VirE2-Venus fluorescence, when constitutively expressed in *Arabidopsis* roots, localizes at the cellular periphery, especially at the cell poles (Shi et al., 2014). Li et al. (2020) showed that during agroinfiltraion, a small amount of VirE2 can enter the nucleus of tobacco leaf cells, and this entry requires the presence of VirD2 and T-strands. To determine whether transgenically-expressed VirE2 also relocalizes to the nucleus in the presence of VirD2 and T-strands, we incubated *Arabidopsis* roots containing an inducible *VirE2-Venus* transgene with β-estradiol. After 24 hr, we infected root segments with a *virE2*-deletion *Agrobacterium* strain either lacking T-DNA (At2403; no binary vector) or containing T-DNA on a binary vector (At2404). Twenty-four hours after inoculation, we observed VirE2-Venus subcellular localization by confocal microscopy. Root segments infected by *A. tumefaciens* At2403 did not show altered VirE2-Venus subcellular localization: fluorescence was distributed throughout the cytoplasm, including at the cellular periphery (Supplemental Figure S10A). However, infection by *A. tumefaciens* At2404 resulted in a partial redistribution of VirE2-Venus within the cytoplasm: fluorescence increased in the perinuclear area (Supplemental Figure S10B). Quantitative analysis of fluorescence intensity showed that perinuclear VirE2 increased ~5-fold in the presence of T-DNA (Figure 11A). However, we were unable to detect VirE2-Venus within the nucleus. These data suggest that the transfer of T-DNA to plant cells results in a partial redistribution of VirE2 to the area surrounding the nucleus.

T-strands delivered by *Agrobacterium* can relocalize VirE2 from the cellular periphery into the cytosol

Because much VirE2 produced *in planta* is already in the general cell cytosol, the data presented in Figure 11A and the movies in Supplemental Figure S11 were not able to show redistribution of VirE2-Venus specifically from the cell periphery into the perinuclear area. We therefore conducted the VirE2-Venus re-localization experiments, following *Agrobacterium* infection, on root segments overexpressing the various myosin VIII or myosin XI-K CBDs. We reasoned that when most VirE2-Venus is tethered to the plasma membrane by myosin VIII-A, VIII-B, VIII-2,

and (to a lesser extent) XI-K, as seen in Figure 9A, we may be able to detect re-distribution to

VirE2-Venus more easily when T-strands are introduced into the cells by *Agrobacterium.*

We generated transgenic plants expressing VirE2-Venus and individual myc-tagged myosin

CBD cDNAs (using *Agrobacterium* strains At2423-At2427), both under the control of a β-

estradiol-inducible promoter. After inducing the transgenes for 24 hr, we infected root segments

with *virE2* mutant *Agrobacterium* strains lacking or containing a T-DNA binary vector. The

bacteria (At2403 and At2404, respectively) additionally contained a plasmid expressing mCherry

to mark the location of the bacteria during infection. We observed fluorescence by confocal

microscopy eight hours after infection and calculated the percentage of fluorescence shift

between the cell periphery and the cytosol. When At2404, capable of transferring T-DNA, was

used, some of the VirE2-Venus fluorescence signal re-localized from the cell periphery into the

cytoplasm when plants expressed the myosin VIII-A, VIII-B, or VIII-2 CBDs (Figures 11C,

11D, and 11E, respectively, and movies in Supplemental Figures S13-15). No such

redistribution of the VirE2-Venus signal was observed when we infected roots with a strain,

At2403, lacking T-DNA. The distribution of VirE2-Venus signal was not altered in roots

infected with *A. tumefaciens* At2404 and expressing the CBDs of myosins VIII-1 or XI-K

(Figures 11 B and F; movies in Supplemental Figures S12 and S16). These results further

suggest that myosins VIII-A, VIII-B, and VIII-2 may temporarily tether VirE2 at the cell

periphery. However, upon transfer of T-DNA into the cell, VirE2 is released into the cellular

interior.

Following T-DNA transfer, VirE2 moves from the cellular periphery to the interior along

actin filaments

To address whether VirE2-Venus moves along actin filaments, we used *Arabidopsis* transgenic

lines expressing inducible VirE2-Venus and myc-tagged myosin VIII CBD cDNAs, and

expressing mCherry-ABD2 to mark actin filaments (using *Agrobacterium* strains At2365-

At2369). Time-lapse imaging of control plants, which expressed VirE2-Venus and mCherry-

ABD2 but without any myosin CBD, indicated that some VirE2-Venus moves along actin

filaments at a velocity of 0.353±0.06 µm/sec (Figure 12). In the presence of either one of the

four myosin VIII CBDs, the velocity of VirE2-Venus along actin filaments in leaf or root cells

remained unchanged (Table 1). Additionally, the velocity of VirE2-Venus movement along

actin filaments was not altered in the *myosin VIII-1/2/a/b* quadruple mutant (Figure 12, Table 1).

However, when the myosin XI-K CBD was expressed in the plant, no VirE2 movement was

detected in either leaf or root cells. These results indicate that only myosin XI-K, but not myosin

VIII proteins, are important for VirE2 trafficking through leaf and root cells.

To test if the blockage of VirE2-Venus movement along actin filaments by the myosin XI-K

CBD could be reversed by the presence of VirD2/T-strands, we infected roots with a *virE2*

mutant *Agrobacterium* strain harboring a T-DNA with a mCherry-intron-NLS expression

cassette (At2405). Time-lapse confocal microscopy still indicated no movement of VirE2-Venus

in the mCherry expressing cells. (Table 1 and Supplemental Figure S17).

Taken together, our results indicate that myosin XI-K but not myosin VIII proteins are

important for VirE2 intracellular trafficking. We suggest that myosin VIII proteins help tether

VirE2 to the plasma membrane/peripheral regions of the cell. Upon introduction of VirD2/T-

strands, VirE2 is released and is trafficked through the cell by myosin XI-K.

DISCUSSION

Importance of VirE2 in *Agrobacterium*-mediated plant transformation

Agrobacterium tumefaciens can transfer virulence (Vir) proteins and T-DNA into plant, fungal, and yeast cells (Chilton et al., 1977; Bundock et al., 1995; Piers et al., 1996; de Groot et al., 1998). During the transfer process, VirE2, the most abundant virulence protein (Engstrom et al., 1987), and four other Vir proteins exit the bacterium via a VirB/VirD4 type IV secretion system (T4SS) and facilitate transformation (Schrammeijer et al., 2003; Cascales and Christie, 2004; Vergunst et al., 2005). VirE2 is a single-strand (ss) DNA binding protein that binds ssDNA *in vitro* (Zupan et al., 1996). A popular theory proposes that VirE2 binds T-strands that enter the plant cell following infection by *Agrobacterium*, and that the resulting hypothetical T-complexes, composed of VirD2 covalently linked to T-strands and coated by VirE2 molecules, are the form of T-DNA that traffics within the cell (Li et al., 2020). However, T-complexes have not yet been identified in *Agrobacterium*-infected plants.

virE2 mutant *Agrobacterium* strains are highly attenuated in virulence (Stachel and Nester, 1986). Fewer T-strands can be isolated from plant cells infected by an *Agrobacterium virE2* mutant than from cells infected by a wild-type *Agrobacterium* strain (Yusibov et al., 1994), and the few transformants produced by a *virE2* mutant *Agrobacterium* strain generally harbor large T-DNA deletions, suggesting a protective function of VirE2 on T-strands (Rossi et al., 1996). However, a *virE2* mutant *Agrobacterium* strain can efficiently transform plants expressing VirE2 (*in planta* complementation), indicating both the importance of VirE2 in transformation and that the function of VirE2 in transformation occurs within the plant (Citovsky et al., 1992; Lapham et al., 2021). The efficient rescue of an *Agrobacterium virE2* mutant by *in planta* complementation indicates that VirE2 trafficking within a plant cell can be studied using plant lines that produce VirE2, rather than studying VirE2 trafficking after delivery from *Agrobacterium*. However, our *in planta* system is not appropriate for investigating initial VirE2 delivery into the plant.

As well as its proposed protective function, VirE2 may facilitate T-DNA entry into the plant by forming single-strand DNA-specific pores in the plant plasma membrane, as it can in artificial membranes (Dumas et al., 2001). Following transfer to the plant, VirE2 initially accumulates on the cytoplasmic side of host plasma membranes at points where bacteria contact the plant cell (Li and Pan, 2017). Accumulation of VirE2 on the membrane is facilitated by its interaction with

the *Agrobacterium* effector protein VirE3, which also accumulates in the host plasma membrane

at contact sites with the bacteria (Lacroix et al., 2005; Li et al., 2018; Li et al., 2020). Interaction

of VirE2 with importin alpha may help guide VirE2 towards the nucleus (Bhattacharjee et al.,

2008), although this interaction is weak (Chang et al., 2014). Therefore, VirE2 may be involved

in the trafficking of the T-complex inside the plant cell. However, our knowledge of the precise

trafficking process of VirE2 inside the host cells, and the proteins that coordinate this trafficking

to facilitate transformation, remains sparse.

Importance of the actin cytoskeleton for transformation and intracellular VirE2 movement

The actin cytoskeleton is essential for numerous biological processes in plants, including cell

morphogenesis, tip growth, organelle movement, vesicle trafficking, and maintenance of cell

polarity (Hussey et al., 2006; Staiger et al., 2009; Duan and Tominaga, 2018). Actin genes are

categorized according to their expression patterns. In *Arabidopsis*, the vegetative actins (*ACT2*,

ACT7, and *ACT8*) are mainly expressed in roots (Meagher et al., 1999), whereas *ACT4* and

ACT12 are primarily expressed during pollen development (Huang et al., 1996). We conducted

transient and stable root transformation assays of *act2* and *act7* mutants (Figure 1A). These

assays indicated that mutation of these genes greatly inhibited transformation. Restoration of

transformation-susceptibility by introducing an *ACT7* cDNA into *act7* mutant plants confirmed

that the transformation deficiency of the *act7* mutant was caused by mutation of this gene.

Mutation of the pollen-expressed *ACT12* gene had no effect on root transformation, as expected.

In addition, we measured the velocity of VirE2-Venus movement along actin filaments, which

were marked with mCherry-ABD2, in *Arabidopsis* leaf cells. VirE2 aggregates moved with an

average velocity of 0.353 ± 0.06 µm/sec in leaf cells and 0.312 ± 0.07 µm/sec in root cells (Table

1), similar to the velocity (0.434 µm/sec) recorded by Yang et al. (2017) within tobacco leaf cells.

Salman et al. (2005) showed that VirE2 binds to microtubules *in vitro*, and Sakalis et al.

(2014) showed colocalization of YFP-VirE2 with fluorescently tagged microtubules in yeast and

plant cells. However, they did not assay transformation in the presence of microtubule

disrupting agents. We observed no difference in transformation between wild-type and *bot-1*

mutant *Arabidopsis* roots, which have a disorganized cortical microtubule network, indicating

that correctly organized microtubules are not required for efficient transformation of plants.

Importance of *Arabidopsis* myosin isoforms for transformation

The movement of molecules, vesicles, and organelles along actin filaments relies on the highly

conserved myosin motors (Avisar et al., 2008b; Madison and Nebenfuhr, 2013; Ueda et al., 2015;

Citovsky and Liu, 2017). Myosins have actin-binding, ATPase, and cargo binding domains, and

these activities contribute to a wide range of physiological processes, including root hair growth,

organelle motility, tip growth of pollen tubes, and gamete nuclear migration (Prokhnevsky et al.,

2008; Peremyslov et al., 2010; Madison et al., 2015; Geeves, 2016). Plants contain two classes

of myosins, the plant-specific myosin VIII class comprising four members (myosins VIII-1, 2, A,

and B) and the myosin XI class comprising 13 members (myosins XI-A through K, and myosins

XI-1 and XI-2). Most of the myosin XI proteins are localized in the cytosol and are responsible

for intracellular movement of their cargos, including vesicles and organelles (Peremyslov et al.,

2008; Peremyslov et al., 2010). Myosin XI-K in particular is important for vesicular trafficking

(Yang et al., 2014). Myosin VIII proteins localize predominantly to the cellular periphery,

including the plasma membrane, plasmodesmata, and cortical microtubules (Golomb et al., 2008;

Haraguchi et al., 2014; Bar-Sinai et al., 2022) although we have found that myosin VIII-B also

localizes to the nucleus (Figure 9A). Myosin VIII proteins may be important for intercellular

movement of viral proteins and for helping guide plant cell division (Avisar et al., 2008a; Amari

et al., 2014; Wu and Bezanilla, 2014).

Upon entry into a plant cell, VirE2 may interact with VirE3 and associate with the plasma

membrane where myosin VIII proteins localize, then move towards the nucleus after associating

with T-DNA (Li et al., 2018; Tu et al., 2018). We therefore tested the importance of each

myosin XI and myosin VIII protein for transformation by conducting transient and stable root

transformation assays on individual myosin gene mutants (Figures 1B and 1C). Interestingly,

only disruption of the myosin *XI-h* gene substantially (greater than two-fold) inhibited

transformation (Supplemental Figure S1A). Myosin XI-H is moderately strongly expressed in

all root tissues and is phylogenetically most closely related to myosins XI-B and XI-G, that are

not well-expressed in roots, and to myosin XI-2 which does show root expression (Peremyslov et

al., 2011). No specific cellular function had previously been assigned to myosin XI-H (reviewed

by Duan and Tominaga, 2018). We were somewhat surprised not to see an effect of a myosin *xi-*

k mutation on transformation because Yang et al. (2017) showed that RNAi inhibition of *N.*

benthamiana myosin XI-K decreased *Agrobacterium*-mediated transformation by ~50%.

However, *Arabidopsis* may encode more myosin XI isoforms than does *N. benthamiana* (Avisar

et al., 2008b), permitting functional redundancy of the *Arabidopsis* myosin XI isoforms to

conceal a role for *Arabidopsis* myosin XI-K in transformation.

Because the vast majority of single myosin XI or myosin VIII mutants did not affect

transformation, we tested the effect of higher order myosin mutants on this process. Roots from

plants mutant in two, three, or four myosin XI genes showed increasing deficiency in

transformation susceptibility (Figures 1B and 1C). As previously noted, these higher order

myosin XI mutants showed increasing root developmental defects as more myosin XI genes

were mutated (Supplemental Figure S1B; Ojangu et al., 2007; Peremyslov et al., 2008). Because

our *Arabidopsis* transformation assays were conducted on roots, the natural target for

Agrobacterium-mediated transformation, we were concerned that these root developmental

problems may complicate the interpretation of our transformation data. We therefore turned our

attention to higher order *Arabidopsis* myosin VIII mutants whose roots appeared normal

(Supplemental Figure S1B). These higher order myosin VIII mutants also showed decreased

transformation susceptibility (Figures 1B and 1C).

To test the importance of each myosin *VIII* gene in transformation, we individually expressed

cDNAs for each of the myosin VIII proteins in the myosin *VIII-1/2/a/b* quadruple mutant. Three

of these genes, encoding myosins VIII-2, A, and B, could partially restore transformation

susceptibility to the quadruple mutant (Figure 2A), indicating that they play an important role in

transformation. Simultaneous expression of two of these cDNAs in the quadruple mutant further

increased transformation (Figure 2C), indicating that these myosin VIII proteins are not

completely redundant with regard to transformation. However, expression of the cDNA

encoding myosin VIII-1 did not affect transformation (Figure 2A), suggesting a fundamental

difference among these class VIII myosins for their roles in *Agrobacterium*-mediated

transformation. As shown in Figures 6, 7, 8, 10 and discussed below, myosins VIII-2, VIII-A,

and VIII-B can directly interact with *Agrobacterium* VirE2, an important effector protein for

transformation, whereas myosin VIII-1 does not directly interact with VirE2. However, it may

indirectly interact with VirE2 within a protein complex. Ectopically expressed myosin VIII-A,

VIII-B, and VIII-2 cargo binding domains (CBD) localize strongly to the peripheral regions of

root cells (myosin VIII-B also localizes in the nucleus), whereas myosin VIII-1 additionally

localizes substantially in the cytosol (Figure 9A). Furthermore, expression of the CBDs of

myosins VIII-2 and VIII-A decreased the size of VirE2 aggregates that form in the myosin *VIII-*

1/2/a/b quadruple mutant, whereas expression of the myosin VIII-1 CBD did not. These

observations indicate a fundamental difference among these myosin VIII isoforms in their

interaction with VirE2, their subcellular localization, and their effects on transformation.

Interestingly, overexpression of a myosin *VIII-1* cDNA in wild-type *Arabidopsis* increased

transformation, indicating both that myosin VIII-1 does play some role in transformation, and

that this protein is limiting in wild-type roots (Figure 2B). Overexpression of myosin *VIII-A*,

VIII-B, or *VIII-2* cDNAs in wild-type roots, either individually or in combination, did not alter

transformation susceptibility, indicating that these proteins are naturally found in amounts

maximal for transformation.

Some myosin VIII isoforms may tether VirE2 to the plasma membrane until T-strands

enter the plant cell

Several studies have indicated that myosin VIII proteins, or their globular tail domains, localize

at the plasma membrane, especially at plasmodesmata (Radford and White, 1998; Reichelt et al.,

1999; Avisar et al., 2008a; Golomb et al., 2008; Amari et al., 2014). Our results showed that

VirE2-Venus substantially co-localizes with each of the four transgenically expressed myosin

VIII protein CBDs at the plasma membrane of root cells, although co-localization with myosin

VIII-1 is less extensive (Figure 9A). When roots of these transgenic plants were incubated with

Agrobacterium lacking T-DNA, the association of VirE2 with the peripheral regions of the cell

remained largely unchanged. However, incubation of these roots with *Agrobacterium* containing

T-DNA relocalized much of the VirE2-Venus signal to the cytosol. In addition, incubation of

roots expressing VirE2-Venus (but not any myosin VIII CBD) with *Agrobacterium* containing T-

DNA relocalized a portion of the VirE2-Venus signal to the perinuclear area, but not into the

nucleus. These results are consistent with previous observations that VirE2 localizes mainly

within the cytosol and, especially, perinuclear regions upon infection of cells with an

Agrobacterium strain that can transfer VirD2/T-strands to plant cells, and that only a very small

quantity may enter the nucleus (Li et al., 2020).

To examine how various myosin isoforms may faciltate VirE2 movement from the plasma

membrane region to the cytosol, we measured the velocity of VirE2-Venus movement along

mCherry-ABD2-labeled actin filaments with and without expression of various myosin VIII or

myosin XI-K CBDs. For these experiments, it was assumed that the CBDs could act as domiant

negative molecules to bind VirE2, thereby inhibiting VirE2 movement along these actin

filaments. In both the absence and presence of myosin VIII CBD expression, VirE2 trafficked

along *Arabidopsis* actin filaments with a velocity of 0.31-0.40 µm/sec, a velocity similar (~0.502

µm/sec) to that calculated by Yang et al. (2017) in tobacco. These data indicate that myosin VIII

proteins are not important for intracellular VirE2 trafficking. However, co-expression of the

myosin XI-K CBD with VirE2-Venus completely blocked VirE2 movement along actin

filaments, suggesting that this myosin isoform is at least partially responsible for intracellular

VirE2 trafficking, as suggested by Yang et al. (2017).

A model for the role of myosin proteins in VirE2 plasma membrane tethering and

movement through plant cells

Research from several laboratories, and this study, has informed our revised model for how

Agrobacterium virulence effector proteins, such as VirE2, and T-DNA enter a plant cell, form

complexes, and traffic T-DNA to the nucleus (Figure 13). VirE2 has the ability to generate pores

in artificial lipid membranes through which single-strand DNA molecules, such as T-strands,

may penetrate (Dumas et al., 2001; Duckely et al., 2005; Grange et al., 2008). VirE2 delivered

from *Agrobacterium* localizes to the cytoplasmic side of the plant plasma membrane, held there

by interaction with another virulence effector protein, VirE3 (Li et al., 2018, 2020). Our data

similarly show that plant-expressed VirE2-Venus, which can effect transformation by a *virE2*

mutant *Agrobacterium* cell, also partially localizes to the cytoplasmic side of the plasma

membrane. We propose that VirE2 is tethered to the plasma membrane by various myosin VIII

isoforms. Upon entry of VirD2/T-strands into the plant, VirE2 is released from the plasma

membrane and associates with T-strands. This hypothetical T-complex is trafficked along an

actin/endoplasmic reticulum network (Yang et al., 2017) by myosin XI-K, or a functionally

similar myosin XI protein, towards the nucleus. Consistent with this mechanism of trafficking,

we have previously shown that a Rab GTPase, which modulates vesicle trafficking in

biosynthetic and endocytic pathways, is important for transformation (Hwang and Gelvin, 2004),

and that a VAP33-like SNARE protein, that localizes to ER-plasma membrane contact sites

(Wang et al., 2014; Wang et al., 2015), interacts with VirE2 (Lee et al., 2012).

We are aware that one aspect of our analysis does not easily fit this proposed model:

Expression of the myosin XI-K CBD completely inhibited detectable VirE2 movement through

the cytosol but had no effect on *Agrobacterium*-mediated transformation. Inhibition of VirE2

movement by the myosin XI-K CBD occurred regardless of whether *Agrobacterium* was present

to transfer T-DNA into the cells. We note that the requirement for VirE2 in *Agrobacterium*-

mediated transformation is not absolute: 1. *Agrobacterium*-mediated yeast transformation does

not require VirE2, although mutation of *virE2* results in a ~10-fold decrease in yeast

transformation (Bundock et al., 1995), and 2. Some *A. rhizogenes* strains do not encode VirE2,

although they are highly virulent. These strains encode two proteins, GALLS-FL and GALLS-

CT, that can functionally complement a *virE2*-mutant *A. tumefaciens* strain, although the amino

acid sequence and protein domains of VirE2 and GALLS proteins are highly dissimilar (Hodges

et al., 2004, 2006; Ream, 2009). With regard to the results of experiments conducted in this

study, we suggest either that a small, undetectable population of VirE2 molecules associates with

incoming T-strands and traffics towards the nucleus to effect transformation, or (a somewhat

heretical model) that VirE2 does not interact with T-strands in plant cells and that tracking VirE2

movement through the cell does not correlate with intracellular T-DNA movement. Future

experiments will attempt to solve this conundrum.

MATERIALS AND METHODS

Plant and bacterial materials and growth conditions. Seeds of wild-type (ecotype Col-0), mutant, and transgenic *Arabidopsis thaliana* were surface-sterilized with 50% bleach and 0.01% SDS, washed five times with sterile water, then placed on Gamborg's B5 medium (Gibco) with the appropriate antibiotics (when used, hygromycin 10 µg/mL). Seedlings were grown in 16 hr light/8 hr dark conditions at 25°C. *E. coli* strains were grown at 37°C in LB medium (per liter: Tryptone 10 g, NaCl 10 g, Yeast extract 5 g). *A. tumefaciens* strains were grown at 28°C in YEP medium (per liter: Bacto peptone 10 g, Yeast extract 10 g, NaCl 5 g). When used, antibiotic concentrations were: For *E. coli*, ampicillin, 100 µg/ml; kanamycin, 50 µg/ml; spectinomycin, 100 µg/ml; for *A. tumefaciens*, rifampicin, 10 µg/ml; spectinomycin, 100 µg/ml. All bacterial strains and plasmid constructs used in this study are listed in Supplemental Tables S1 and S2.

Construction of plasmids and binary vectors and generation of transgenic plants. A description of all plasmids used in this study, including details of their construction, is given in Supplemental Table S1. Primers used for these constructions are given in Supplemental Table S3. Final *E. coli* and *A. tumefaciens* strains are given in Supplemental Table S2. Recombinant constructions were transformed into *E. coli* TOP10 or DH10B before characterization and transfer to the appropriate *Agrobacterium* strains by electroporation. For protein expression, plasmids were transferred to *E. coli* BL21DE3. T-DNA binary vectors in *Agrobacterium* were used to transform plants of the Col-0 or the *myosin VIII-1/2/a/b* quadruple mutant background utilizing a floral dip transformation protocol (Clough and Bent, 1998).

***Agrobacterium*-mediated transient and stable transformation assays.** Overnight *Agrobacterium* cultures were diluted 1:10 into 20 ml of YEP medium containing the appropriate antibiotics and grown at 28°C until they reached an $A_{600} = 0.83$ (~10^9 cells/ml). The cells were collected by centrifugation, washed with sterile 0.9% NaCl, and serially diluted in 0.9% NaCl for transient and stable root transformation assays. For transient transformation assays, infection of *Arabidopsis* root segments was performed with *Agrobacterium* strains containing the T-DNA binary vector pBISN1 (Narasimhulu et al., 1996) as previously described (Tenea et al., 2009). Transformation efficiency was reported as the percentage of root segments stained blue with 5-

bromo-4-chloro-3-indolyl-β-D-glucuronic acid (X-gluc) after two days co-cultivation with

bacteria on MS medium plus an additional four days on CIM (Callus-Inducing Medium)

containing 100 µg/ml timentin. For stable transformation, after two days of cocultivation, root

segments were separated onto CIM containing 100 µg/ml timentin and the appropriate selection

agent. The plates were incubated at 23°C for four weeks, after which they were scored for callus

development on the ends of the roots. Stable transformation efficiency was calculated as the

percentage of total segments showing calli. Transient and stable transformation assays were

conducted in triplicate, with >100 root segments per replicate.

Bimolecular fluorescence complementation (BiFC). A VirE2 coding sequence was fused with

cEYFP (C-terminal 64 amino acid residues of an *EYFP* gene) and under the control of a β-

estradiol-inducer promoter in the vector pSAT1 to make pSAT1-Pi-VirE2-cEYFP. cDNAs

encoding myosin VIII CBDs were amplified using the appropriate primers (Supplemental Table

S3), fused with nVenus (N-terminal 174 amino acids of a *Venus* gene) and cloned, under the

control of a β-estradiol-inducer promoter in pSAT6, to make pSAT6-Pi-nVenus-myosin VIII

CBD. The Pi-VirE2-cEYFP expression cassette and the Pi-nVenus-myosin VIII CBD

expression cassettes were digested with *Asc*I and PI-*Psp*I, respectively, and cloned into the same

sites in the binary vector pE4437. pE4437 also contains a *hptII* plant selection marker, an

expression cassette for a nuclear marker (P_{35S}-Cerulean-NLS), and an actin cytoskeleton marker

(P_{35S}-mCherry-actin binding domain2, ABD2). These final constructs were introduced into *A.*

tumefaciens GV3101::pMP90 (Koncz and Schell, 1986) by electroporation to make *A.*

tumefaciens At2320-At2324, respectively, and used to generate transgenic Col-0 *Arabidopsis*

plants for *in vivo* BiFC assays.

Yeast two hybrid assay. An octopine-type VirE2 gene from pTiA6 was cloned into the bait

vector pGBKT7, and the myosin VIII and myosin XI CBD cDNAs were individually cloned into

the prey vector pGADT7 (Clontech). As negative controls, empty bait, empty prey, and Lamin

C prey plasmids were used (Krendel et al., 2007). A cDNA encoding the VirE2-interacting

protein 1 (VIP1) was used as a positive control (Tzfira et al., 2001). Yeast transformations were

performed using a lithium acetate method (Sato et al., 1994). Cells were plated onto synthetic

dropout (SD)-agar plates lacking tryptophan and leucine. Three days after transformation, yeast

cells from the resulting colonies were plated onto SD medium lacking tryptophan, leucine, and

histidine (-Trp-Leu-His) supplemented with 3-amino-1,2,4-triazole (3AT) to test for protein

interactions.

***In vitro* protein pull-down assay.** The plasmid pET28a (Novagen, San Diego, CA) was used to

generate recombinant proteins fused in-frame with a Venus tag or a myc tag. A full-length

cDNA encoding VirE2 fused with Venus was cloned into pET28a with *Nco*I+*Not*I. cDNAs

encoding the CBDs (amino acids 953-1,166 for myosin VIII-1; amino acids 1,003-1,220 for

myosin VIII-2; amino acids 948-1,153 for myosin VIII-A; amino acids 954-1,136 for myosin

VIII-B; amino acids 808-1,531 for myosin XI-K) of myosin VIII and XI-K were amplified and

cloned into pET28a with *Nco*I+*Bam*HI (myosin VIII CBDs) or *Nco*I+*Not*I (myosin XI-K CBD)

to generate myc fusions. The full-length cDNA of VIP1 was cloned into pET28a with

*Nco*I+*Bam*HI to use as a myc-tagged positive control, and for negative controls Lamin C fused

with a myc tag, or Venus only, were cloned into pET28a with *Sma*I+*Bam*HI. The constructs

were transformed into *E. coli* BL21 (DE3). Bacteria were cultured at 16^{o}C for 18 hours in the

presence of 0.5 mM isopropylthio-β-galactoside (IPTG). The cells were centrifuged and the

pellet was sonicated in cell lysis buffers (20 mM HEPES-KOH pH 7.2, 50 mM potassium acetate,

50 mM KCl, 1 mM EDTA, 1 mM EGTA, 1 mM DTT, 5% glycerol, 0.5% Triton X-100, 1 mM

phenylmethylsulfonylfluoride [PMSF]). VirE2-Venus and Venus cell lysates were incubated

with GFP-Trap magnetic agarose beads (Chromotek, USA) at 4°C for 2 hr with gentle shaking.

The beads were washed three times with cell lysate buffer (0.2% Triton X-100) and then

incubated with bacterial lysates from *E. coli* cells expressing myc-tagged VIP1, myc-tagged

Lamin C, or myc-tagged myosin VIII/XI-K CBDs proteins that were prepared as described

above. The binding reactions were incubated at 4°C for 1 hr with gentle shaking. The beads

were washed three times with cell lysate buffer (0.2% Triton X-100) followed by boiling in

sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) buffer for 10 min.

Protein samples were loaded onto 10% SDS polyacrylamide gels, subjected to electrophoresis

(60 Volts 1.5 hrs for the stacking gel and 100 Volts 4.5 hrs for the separating gel), then either

stained with Coomassie blue or subjected to western blot analysis using a 1:1000 dilution of

mouse anti-GFP or mouse anti-Myc antibodies (Cell Signaling Technology, USA). After

addition of a 1:5000 dilution of horseradish peroxidase-conjugated horse anti-mouse secondary

antibodies (Catalog #7076; Cell Signaling Technology, USA), proteins were detected using an

ECL kit (LI-COR Biosciences, Lincoln, NE, USA).

***In vivo* co-immunoprecipitation (co-IP) assay.** Transgenic *Arabidopsis* plants (generated using

Agrobacterium strains At2423-At2427) harboring an inducible expression cassette of either

VirE2-Venus, or Venus, and of each myc-tagged myosin VIII or XI-K CBDs were used for *in*

vivo protein-protein interaction assays. Transgenic lines expressing, upon induction, myc-tagged

CBDs and Venus were used as negative controls (At2428). Four-week old plants were treated

with 5 µM β-estradiol for 24 hr, and roots were harvested and ground in liquid nitrogen using a

mortar and pestle. A total of 0.25 g of the ground tissue was transferred into a 2 ml tube and 500

µl of CoIP buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 0.1% Triton X-100, 0.2% Nonidet

P-40, 0.6 mM PMSF, and 20 µM MG132 in a Roche protease inhibitor cocktail) was added. The

mixture was mixed thoroughly by repeated cycles of 30 sec vortexing followed by chilling on ice.

The samples were centrifuged at 4°C, 13000g for 5 min. A total of 20 µl of anti-GFP beads was

pre-washed with wash buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 0.6 mM PMSF, and 20

µM MG132 with Roche protease inhibitor cocktail) three times, the beads were added to the

extracted supernatant solution, and the reactions were incubated at 4°C for 1 hr with gentle

shaking. The beads were washed five times with wash buffer, proteins were eluted by incubation

with 4x LDS sample loading buffer (Invitrogen, CA, USA) at 95℃ for 3 min. The proteins were

fractionated by SDS-PAGE and analyzed using western blots, with a 1:1000 dilution of anti-GFP

and anti-Myc antibodies, respectively. The proteins were detected using an ECL kit (LI-COR

Biosciencces, Lincoln, NE, USA).

Localization of VirE2-Venus after infection of *Arabidopsis* roots with *Agrobacterium* strains

harboring or lacking T-DNA. Transgenic *A. thaliana* lines expressing a *VirE2-Venus* gene

under the control of a β-estradiol inducible promoter were treated with 5 µM β-estradiol for 24 hr.

Roots were infected with a *virE2* mutant strain *A. tumefaciens* strain (*A. tumefaciens* EHA105

containing a non-polar deletion of *virE2*; At1872) containing the plasmid pBBR1-

MCS2::mCherry to track red fluorescent bacteria (At2405). The *Agrobacterium* cells either

contained or lacked the T-DNA binary vector pE4672 (*A. tumefaciens* At2404 and At2403,

respectively). Overnight bacterial cultures were transferred to fresh AB-glucose medium and

grown until the cell density reached an A_{600}=0.83. The bacteria were centrifuged and

resuspended in induction medium (1x AB salts, 2% glucose, 30 mM MES, 2 mM Na$_2$HPO4, 2

mM NaH$_2$PO4) containing 100 µM acetosyringone and cultured for 16-20 hr. Six hr after

cocultivation with bacteria at 10^8 cfu/ml, root segments were observed by confocal microscopy

(Zesis 880 Upright Confocal microscope with a Plan-Apo 20×/0.8 objective). Images from 100

cells of each infection were analyzed using Image J.

Confocal microscopy imaging. For VirE2-Venus trafficking, transgenic plantlets harboring an

inducible *Venus-VirE2* transgene (generated using *A. tumefaciens* At2365-At2369) were placed

on B5 agar plates containing 100 µg/ml timentin and 20 µg/ml hygromycin and grown vertically

for one week. The plants were treated with 10 µM β-estradiol for 9 hr, washed with sterile water,

and time lapse images of the roots were taken using a Zeiss 880 confocal microscope equipped

with a 20 X water objective. For VirE2 relocalization assays, similarly grown plants were

treated with 5 µM β-estradiol for 24 hr, washed with sterile water, and infected with 10^8 cfu/ml

of a *virE2* mutant *Agrobacterium* strain containing or lacking a T-DNA binary vector. Infections

were conducted for 8 hr as described above, and images were taken using a Zeiss 880 confocal

microscope equipped with a Plan-Apo 20×/0.8 objective.

SUPPLEMENTAL DATA

Supplemental Figure S1. Most *myosin VIII* and *myosin XI* single mutants are not deficient in

stable transformation.

Supplemental Figure S2. Growth and morphology of wild-type and higher order *myosin VIII*

and *XI* mutant plants.

Supplemental Figure S3. Effect on transformation of expressing full-length myosin cDNAs in

wild-type and *myosin VIII-1/2/a/b* mutant plants.

Supplemental Figure S4. Effect on transformation of expressing full-length inducible myosin

cDNAs in wild-type and *myosin VIII-1/2/a/b* mutant plants.

Supplemental Figure S5. Effect on transformation of expressing pairs of full-length myosin

cDNAs in wild-type and *myosin VIII-1/2/a/b* mutant plants.

Supplemental Figure S6. Inducible expression of myosin VIII, but not myosin XI-K, CBDs

inhibits transformation.

Supplemental Figure S7. Co-localization of VirE2-Venus and FM4-64.

Supplemental Figure S8. Subcellular localization of VirE2-Venus in transgenic plants.

Supplemental Figure S9. Myosin VIII CBDs colocalize with the plasma membrane marker

PIP2A.

Supplemental Figure S10. VirE2 re-localizes to the perinuclear area after infection by an

Agrobacterium strain that can transfer T-DNA.

Supplemental Figure S11. Infection of root cells with an *Agrobacterium* strain capable of

transferring T-DNA re-localizes VirE2-Venus from the cellular periphery into the cytoplasm.

Supplemental Figure S12. Infection of root cells with an *Agrobacterium* strain capable of

transferring T-DNA does not re-localizes VirE2-Venus from the cellular periphery into the

cytoplasm when the myosin VIII-1 CBD is expressed.

Supplemental Figure S13. Infection of root cells with an *Agrobacterium* strain capable of

transferring T-DNA re-localizes VirE2-Venus from the cellular periphery into the cytoplasm.

Supplemental Figure S14. Infection of root cells with an *Agrobacterium* strain capable of

transferring T-DNA re-localizes VirE2-Venus from the cellular periphery into the cytoplasm.

Supplemental Figure S15. Infection of root cells with an *Agrobacterium* strain capable of

transferring T-DNA re-localizes VirE2-Venus from the cellular periphery into the cytoplasm.

Supplemental Figure S16. Expression of the myosin XI-K CBD does not affect VirE2-Venus

localization upon infection by *Agrobacterium.*

Supplemental Figure S17. Expression of the myosin XI-K CBD inhibits movement of VirE2-

Venus.

ACKNOWLEDGEMENTS

The authors thank Dr. Valerian V. Dolja, Oregon State University, for providing the *Arabidopsis*

mutant lines. We thank the Purdue University Imaging Facility for use of their confocal

microscope. We also thank Dr. Kiran Mysore for critical reading of the manuscript.

FUNDING

This project was supported by grants from the US National Science Foundation: IOS 1725122 and IOS-2006668. DNA sequence data were obtained with partial support from a P30 grant (CA023168) to the Purdue University Cancer Center.

REFERENCES

Amari K, Di Donato M, Dolja VV, Heinlein M. (2014) Myosins VIII and XI play distinct roles
in reproduction and transport of tobacco mosaic virus. PLoS Pathog 10: e1004448

Avisar D, Prokhnevsky AI, Dolja VV. (2008a) Class VIII myosins are required for
plasmodesmatal localization of a closterovirus Hsp70 homolog. J Virol 82: 2836-2843

Avisar D, Prokhnevsky AI, Makarova KS, Koonin EV, Dolja VV. (2008b) Myosin XI-K is
required for rapid trafficking of Golgi stacks, peroxisomes, and mitochondria in leaf cells
of *Nicotiana benthamiana*. Plant Physiol 146: 1098-1108

Bar-Sinai S, Belausov E, Dwivedi V, Sadot,E. (2022) Collisions of cortical microtubules with
membrane associated myosin VIII tail. Cells 11:145

Bezanilla M, Horton AC, Sevener HC, Quatrano RS. (2003) Phylogenetic analysis of new plant
myosin sequences. J Mol Evol 57: 229-239

Bhattacharjee S, Lee L-Y, Oltmanns H, Cao H, Veena, Cuperus J, Gelvin SB. (2008) IMPa-4, an
Arabidopsis importin alpha isoform, is preferentially involved in *Agrobacterium*-
mediated plant transformation. Plant Cell 20: 2661-2680

Bichet A, Desnos T, Turner S, Grandjean O, Hofte H. (2001) BOTERO1 is required for normal
orientation of cortical microtubules and anisotropic cell expansion in *Arabidopsis*. Plant J
25: 137-148

Bundock P, den Dulk-Ras A, Beijersbergen A, Hooykaas PJ. (1995) Trans-kingdom T-DNA
transfer from *Agrobacterium tumefaciens* to *Saccharomyces cerevisiae*. EMBO J 14:
3206-3214

Cascales E, Christie,PJ. (2004) Definition of a bacterial type IV secretion pathway for a DNA
substrate. Science 304: 1170-1173

Chang CW, Williams SJ, Counago RM, Kobe B. (2014) Structural basis of interaction of
bipartite nuclear localization signal from *Agrobacterium* VirD2 with rice importin-alpha.
Mol Plant 7: 1061-1064

Chilton M-D, Drummond MH, Merio DJ, Sciaky D, Montoya AL, Gordon MP, Nester EW.
(1977) Stable incorporation of plasmid DNA into higher plant cells: The molecular basis
of crown gall tumorigenesis. Cell 11: 263-271

Christie PJ, Atmakuri K, Krishnamoorthy V, Jakubowski S, Cascales E. (2005) Biogenesis,

architecture, and function of bacterial type IV secretion systems. Annu Rev Microbiol 59:

451-485

Citovsky V, Liu B. (2017) Myosin-driven transport network in plants is functionally robust and

distinctive. Proc Natl Acad Sci USA 114: 1756-1758

Citovsky V, Wong ML, Zambryski P. (1989) Cooperative interaction of *Agrobacterium* VirE2

protein with single-stranded DNA: Implications for the T-DNA transfer process. Proc

Natl Acad Sci USA 86: 1193-1197

Citovsky V, Zupan J, Warnick D, Zambryski P. (1992) Nuclear localization of *Agrobacterium*

VirE2 protein in plant cells. Science 256: 1802-1805

Clough SJ, Bent AF. (1998) Floral dip: A simplified method for *Agrobacterium*-mediated

transformation of *Arabidopsis thaliana*. Plant J 16: 735-743

de Groot MJ, Bundock P, Hooykaas PJ, Beijersbergen AG. (1998) *Agrobacterium tumefaciens*-

mediated transformation of filamentous fungi. Nat Biotechnol 16: 839-842

Djamei A, Pitzschke A, Nakagami H, Rajh I, Hirt H. (2007) Trojan horse strategy in

Agrobacterium transformation: abusing MAPK defense signaling. Science 318: 453-456

Duan Z, Tominaga M. (2018) Actin-myosin XI: An intracellular control network in plants.

Biochem Biophys Res Commun 506: 403-408

Duckely M, Oomen C, Axthelm F, Van Gelder P, Waksman G, Engel A. (2005) The

VirE1VirE2 complex of *Agrobacterium tumefaciens* interacts with single-stranded DNA

and forms channels. Mol Microbiol 58: 1130-1142

Dumas F, Duckely M, Pelczar P, Van Gelder P, Hohn B. (2001) An *Agrobacterium* VirE2

channel for transferred-DNA transport into plant cells. Proc Natl Acad Sci USA 98: 485-

490

Dym O, Albeck S, Unger T, Jacobovitch J, Branzburg A, Michael Y, Frenkiel-Krispin D, Wolf

S., Elbaum M. (2008) Crystal structure of the *Agrobacterium* virulence complex VirE1-

VirE2 reveals a flexible protein that can accommodate different partners. Proc Natl Acad

Sci USA 105: 11170-11175

Embley TM, Martin W. (2006) Eukaryotic evolution, changes and challenges. Nature 440: 623-

630

Engstrom P, Zambryski P, Van Montagu M, Stachel S. (1987) Characterization of
*Agrobacterium tumefacien*s virulence proteins induced by the plant factor acetosyringone.
J Mol Biol 197: 635-645
Foth BJ, Goedecke MC, Soldati D. (2006) New insights into myosin evolution and classification.
Proc Natl Acad Sci USA 103: 3681-3686
Geeves MA. (2016) Review: The ATPase mechanism of myosin and actomyosin. Biopolymers
105: 483-491
Gelvin SB. (1998) *Agrobacterium* VirE2 proteins can form a complex with T strands in the plant
cytoplasm. J Bacteriol 180: 4300-4302
Gelvin SB. (2000) *Agrobacterium* and plant genes involved in T-DNA transfer and integration.
Annu Rev Plant Physiol Plant Mol Biol 51: 223-256
Gelvin SB. (2003) *Agrobacterium* and plant transformation: The biology behind the "gene-
jockeying" tool. Microbiol Mol Biol Rev 67: 16-37
Gelvin, SB. (2006) *Agrobacterium* transformation of *Arabidopsis thaliana* roots: A quantitative
assay. Methods Mol Biol 343: 105-113
Gelvin SB. (2010) Finding a way to the nucleus. Curr Opin Microbiol 13: 53-58
Gelvin SB. (2017) Integration of *Agrobacterium* T-DNA into the plant genome. Annual review
of genetics 51: 22
Golomb L, Abu-Abied M, Belausov E, Sadot E. (2008) Different subcellular localizations and
functions of *Arabidopsis* myosin VIII. BMC Plant Biol 8: 3
Grange W, Duckely M, Husale S, Jacob S, Engel A, Hegner M. (2008) VirE2: A unique ssDNA-
compacting molecular machine. PLoS Biol 6: e44
Haraguchi T, Tominaga M, Matsumoto R, Sato K, Nakano A, Yamamoto K, Ito K. (2014)
Molecular characterization and subcellular localization of *Arabidopsis* class VIII myosin,
ATM1. J Biol Chem 289: 12343-12355
Hodges L, Cuperus J, Ream W. (2004) *Agrobacterium rhizogenes* GALLS protein
substitutes for *Agobacterium tumefaciens* single-stranded DNA-binding protein VirE2. J
Bacteriol 186: 3065-3077
Hodges LD, Vergunst AC, Neal-McKinney J, den Dulk-Ras A, Moyer DM,

Hooykaas PJJ, Ream W. (2006) *Agrobacterium rhizogenes* GALLS protein contains

domains for ATP binding, nuclear localization, and type IV secretion. J Bacteriol 188:

8222-8230

Howard E, Citovsky V. (1990) The emerging structure of the *Agrobacterium* T-DNA transfer

complex. Bioessays 12: 103-108

Huang S, An YQ, McDowell JM, McKinney EC, Meagher RB. (1996) The *Arabidopsis thaliana*

ACT4/ACT12 actin gene subclass is strongly expressed throughout pollen development.

Plant J 10: 189-202

Hussey,PJ, Ketelaar T, Deeks MJ. (2006) Control of the actin cytoskeleton in plant cell growth.

Annu Rev Plant Biol 57: 109-125

Hwang H-H, Gelvin SB. (2004) Plant proteins that interact with VirB2, the *Agrobacterium*

tumefaciens pilin protein, mediate plant transformation. Plant Cell 16: 3148-3167

Koncz C, Schell J. (1986) The promoter of TL-DNA gene 5 controls the tissue-specific

expression of chimaeric genes carried by a novel type of *Agrobacterium* binary vector.

Mol Gen Genet 204: 383-396

Krendel M, Osterweil EK, Mooseker MS. (2007) Myosin 1E interacts with synaptojanin-1 and

dynamin and is involved in endocytosis. FEBS Lett 581: 644-650

Lacroix B, Citovsky V. (2013) The roles of bacterial and host plant factors in *Agrobacterium*-

mediated genetic transformation. Int J Dev Biol 57: 467-481

Lacroix B, Vaidya M, Tzfira T, Citovsky V. (2005) The VirE3 protein of *Agrobacterium* mimics

a host cell function required for plant genetic transformation. EMBO J 24: 428-437

Lapham RA, Lee L-Y, Xhako E, Gomez EG, Nivya VM., and Gelvin, S.B. (2021).

Agrobacterium VirE2 protein modulates plant gene expression and mediates

transformation from its location outside the nucleus. Front Plant Sci 12, 684192

Lee L-Y, Wu F-H, Hsu C-T, She, S-C, Yeh,HY, Liao D-C, Fang M-J, Liu N-T, Yen Y-C,

Dokládal L, Sýkorová E, Gelvin SB, Lin C-S (2012) Screening a cDNA library for

protein-protein interactions directly *in planta*. Plant Cell. 24: 1746-1759

Lee YR, Liu B. (2004) Cytoskeletal motors in *Arabidopsis*. Sixty-one kinesins and seventeen

myosins. Plant Physiol 136: 3877-3883

Li X, Pan SQ. (2017) *Agrobacterium* delivers VirE2 protein into host cells via clathrin-mediated

endocytosis. Sci Adv 3: e1601528

Li X, Tu H, Pan SQ. (2018) *Agrobacterium* delivers anchorage protein VirE3 for companion

VirE2 to aggregate at host entry sites for T-DNA protection. Cell Rep 25: 302-311 e306

Li X, Yang Q, Peng L, Tu H, Lee L-Y, Gelvin SB, Pan SQ. (2020) *Agrobacterium*-delivered

VirE2 interacts with host nucleoporin CG1 to facilitate the nuclear import of VirE2-

coated T complex. Proc Natl Acad Sci USA 117: 26389-26397

Li X, Zhu T, Tu H, Pan SQ. (2020) *Agrobacterium* VirE3 uses its two tandem

domains at the C-terminus to retain its companion VirE2 on the cytoplasmic side of the

host plasma membrane. Front Plant Sci 11:464

Madison SL, Nebenfuhr A. (2013) Understanding myosin functions in plants: Are we there yet?

Curr Opin Plant Biol 16: 710-717

Madison SL, Buchanan ML, Glass JD, McClain TF, Park E, Nebenfuhr A. (2015) Class XI

myosins move specific organelles in pollen tubes and are required for normal fertility and

pollen tube growth in *Arabidopsis*. Plant Physiol 169: 1946-1960

McCullen CA, Binns AN. (2006) *Agrobacterium tumefaciens* and plant cell interactions and

activities required for interkingdom macromolecular transfer. Annu Rev Cell Dev Biol 22:

101-127

Meagher RB, McKinney EC, Kandasamy MK. (1999) Isovariant dynamics expand and buffer

the responses of complex systems: The diverse plant actin gene family. Plant Cell 11:

995-1006

Narasimhulu SB, Deng XB, Sarria R, Gelvin SB. (1996) Early transcription of *Agrobacterium* T-

DNA genes in tobacco and maize. Plant Cell 8: 873-886

Ojangu EL, Jarve K, Paves H, Truve E. (2007) *Arabidopsis thaliana* myosin XIK is involved in

root hair as well as trichome morphogenesis on stems and leaves. Protoplasma 230: 193-

202

Otten L, DeGreve H, Leemans J, Hain R, Hooykaas P, Schell J. (1984) Restoration of

virulence of *vir* region mutants of *Agrobacterium tumefaciens* strain B6S3 by coinfection

with normal and mutant *Agrobacterium* strains. Mol Gen Genet 195: 159-163

Peremyslov VV, Prokhnevsky AI, Dolja VV. (2010) Class XI myosins are required for

development, cell expansion, and F-Actin organization in *Arabidopsis*. Plant Cell 22:

1883-1897

Peremyslov VV, Prokhnevsky AI, Avisar D, Dolja VV. (2008) Two class XI myosins function in

organelle trafficking and root hair development in *Arabidopsis*. Plant Physiol 146: 1109-

1116

Peremyslov,VV, Mockler TC, Filichkin S., Fox SE, Jaiswal P, Makarova K., Koonin .V,

Dolja,VV. (2011) Expression, splicing, and evolution of the myosin gene family in plants.

Plant Physiol 155: 1191-1204

Piers KL, Heath JD, Liang X, Stephens KM, Nester EW. (1996) *Agrobacterium tumefaciens*-

mediated transformation of yeast. Proc Natl Acad Sci USA 93: 1613-1618

Prokhnevsky AI, Peremyslov,VV, Dolja VV. (2008) Overlapping functions of the four class XI

myosins in *Arabidopsis* growth, root hair elongation, and organelle motility. Proc Natl

Acad Sci USA 105: 19744-19749

Radford JE, White RG. (1998) Localization of a myosin-like protein to plasmodesmata. Plant J

14: 743-750

Ream W. (2009) *Agrobacterium tumefaciens* and *A. rhizogenes* use different proteins to transport

bacterial DNA into the plant cell nucleus. Micro Biotechnol 2: 416-42.

Reichelt S, Knight AE, Hodge TP, Baluska F, Samaj J, Volkmann D, Kendrick-Jones J. (1999)

Characterization of the unconventional myosin VIII in plant cells and its localization at

the post-cytokinetic cell wall. Plant J 19: 555-567

Reisen D, Hanson MR. (2007) Association of six YFP-myosin XI-tail fusions with mobile plant

cell organelles. BMC Plant Biol 7: 6

Richards TA, Cavalier-Smith T. (2005) Myosin domain evolution and the primary divergence of

eukaryotes. Nature 436: 1113-1118

Rossi L, Hohn B, Tinland B. (1996) Integration of complete transferred DNA units is dependent

on the activity of virulence E2 protein of *Agrobacterium tumefaciens*. Proc Natl Acad Sci

USA 93: 126-130

Roushan MR, deZeeuw MAM, Hooykaas PJJ, van Heusden GPH. (2018) Application

of phiLOV2.1 as a fluorescent marker for visualization of *Agrobacterium* effector protein

translocation. Plant J 96: 685-699

Sakalis,PA, van Heusden GP, Hooykaas PJ. (2014) Visualization of VirE2 protein translocation

by the *Agrobacterium* type IV secretion system into host cells. Microbiologyopen 3: 104-

117

Salman H, Abu-Arish A, Oliel S, Loyter A, Klafter J, Granek R, Elbaum M. (2005) Nuclear

localization signal peptides induce molecular delivery along microtubules. Biophys J 89:

2134-2145

Sato T, Hanada M, Bodrug S, Irie, S, Iwama N, Boise LH, Thompson CB, Golemis E, Fong L,

Wang HG. (1994) Interactions among members of the Bcl-2 protein family analyzed with

a yeast two-hybrid system. Proc Natl Acad Sci U S A 91: 9238-9242

Sattarzadeh A, Franzen R, Schmelzer E. (2008) The *Arabidopsis* class VIII myosin ATM2 is

involved in endocytosis. Cell Motil Cytoskeleton 65: 457-468

Schrammeijer B, den Dulk-Ras A, Vergunst AC, Jurado Jacome E, Hooykaas PJ. (2003)

Analysis of Vir protein translocation from *Agrobacterium tumefaciens* using

Saccharomyces cerevisiae as a model: Evidence for transport of a novel effector protein

VirE3. Nucleic Acids Res 31: 860-868

Sen P, Pazour GJ, Anderson D, Das A. (1989) Cooperative binding of *Agrobacterium*

tumefaciens VirE2 protein to single-stranded DNA. J Bacteriol 171: 2573-2580

Shi Y, Lee L-Y, Gelvin SB. (2014) Is VIP1 important for *Agrobacterium*-mediated

transformation? Plant J 79: 848-860

Simone M, McCullen CA, Stahl LE, Binns AN. (2001) The carboxy-terminus of VirE2 from

Agrobacterium tumefaciens is required for its transport to host cells by the virB-encoded

type IV transport system. Mol Microbiol 41: 1283-1293

Smith LG, Oppenheimer .G. (2005) Spatial control of cell expansion by the plant cytoskeleton.

Annu Rev Cell Dev Biol 21: 271-295

Sparkes I, Runions J, Hawes C, Griffing L. (2009) Movement and remodeling of the

endoplasmic reticulum in nondividing cells of tobacco leaves. Plant Cell 21: 3937-3949

Stachel SE, Nester EW. (1986) The genetic and transcriptional organization of the vir region of

the A6 Ti plasmid of *Agrobacterium tumefaciens*. EMBO J 5: 1445-1454

Staiger CJ, Sheahan MB, Khurana P, Wang X, McCurdy DW, Blanchoin L. (2009) Actin

filament dynamics are dominated by rapid growth and severing activity in the

Arabidopsis cortical array. J Cell Biol 184: 269-280

Szymanski DB, Cosgrove DJ. (2009) Dynamic coordination of cytoskeletal and cell wall systems

during plant cell morphogenesis. Curr Biol 19: R800-811

Tenea GN, Spantzel J, Lee L-Y, Zhu Y, Lin K, Johnson SJ, Gelvin SB. (2009) Over-expression

of several *Arabidopsis* histone genes increases *Agrobacterium*-mediated transformation

and transgene expression in plant cells. Plant Cell 21: 3350–3367

Trybus KM. (2008) Myosin V from head to tail. Cell Mol Life Sci 65: 1378-1389

Tu H, Li X, Yang Q, Peng L, Pan SQ. (2018) Real-time trafficking of *Agrobacterium* virulence

protein VirE2 inside host cells. Curr Top Microbiol Immunol 418: 261-286

Tzfira T, Vaidya M, Citovsky V. (2001) VIP1, an *Arabidopsis* protein that interacts with

Agrobacterium VirE2, is involved in VirE2 nuclear import and *Agrobacterium* infectivity.

EMBO J 20: 3596-3607

Tzfira T, Li J, Lacroix B, Citovsky V. (2004) *Agrobacterium* T-DNA integration: Molecules and

models. Trends Genet 20: 375-383

Ueda H, Tamura K, Hara-Nishimura I. (2015) Functions of plant-specific myosin XI: from

intracellular motility to plant postures. Curr Opin Plant Biol 28: 30-38

Ueda H, Yokota E, Kutsuna N, Shimada T, Tamura K, Shimmen T, Hasezawa S, Dolja VV.,

Hara-Nishimura I. (2010) Myosin-dependent endoplasmic reticulum motility and F-actin

organization in plant cells. Proc Natl Acad Sci USA 107: 6894-6899

Ueki S, Citovsky V. (2005) Identification of an interactor of cadmium ion-induced glycine-rich

protein involved in regulation of callose levels in plant vasculature. Proc Natl Acad Sci

USA 102: 12089-12094

Vale RD. (2003) The molecular motor toolbox for intracellular transport. Cell 112: 467-480

Vergunst AC, van Lier MC, den Dulk-Ras A, Stuve TA, Ouwehand A, Hooykaas PJ. (2005)

Positive charge is an important feature of the C-terminal transport signal of the VirB/D4-

translocated proteins of *Agrobacterium*. Proc Natl Acad Sci USA 102: 832-837

Wang H, Han S, Siao W, Song C, Xiang Y, Wu X, Cheng P, Li H, Jasik J, Micieta K, Turna J,

Voigt B, Baluska F, Liu J, Wang Y, Zhao H. (2015) *Arabidopsis* synaptotagmin 2

participates in pollen germination and tube growth and is delivered to plasma membrane

via conventional secretion. Mol Plant 8: 1737-1750

Wang P, Hawkins TJ, Richardson C, Cummins I, Deeks MJ, Sparkes I, Hawes C, Hussey PJ.

(2014) The plant cytoskeleton, NET3C, and VAP27 mediate the link between the plasma

membrane and endoplasmic reticulum. Curr Biol 24: 1397-1405

Wojtaszek P, Anielska-Mazur A, Gabrys H, Baluska F, Volkmann D. (2005) Recruitment of
myosin VIII towards plastid surfaces is root-cap specific and provides the evidence for
actomyosin involvement in root osmosensing. Funct Plant Biol 32: 721-736

Wu Z, Bezanilla M. (2014) Myosin VIII associates with microtubule ends and together with
actin plays a role in guiding plant cell division. Elife 3:e03498

Yang L, Qin L, Liu G, Peremyslov VV, Dolja VV, Wei Y. (2014) Myosins XI modulate host
cellular responses and penetration resistance to fungal pathogens. Proc Natl Acad Sci
USA 111: 13996-14001

Yang Q, Li X, Tu H, Pan SQ. (2017) *Agrobacterium*-delivered virulence protein VirE2 is
trafficked inside host cells via a myosin XI-K-powered ER/actin network. Proc Natl Acad
Sci USA 114: 2982-2987

Yusibov VM, Steck TR, Gupta V, Gelvin, SB. (1994) Association of single-stranded transferred
DNA from *Agrobacterium tumefaciens* with tobacco cells. Proc Natl Acad Sci USA 91:
2994-2998

Zuo J, Niu Q-W, Chua N-H. (2000) An estrogen receptor-based transactivator XVE mediates
highy inducible gene expression in transgenic plants. Plant J 24: 265-273

Zupan JR, Citovsky V, Zambryski P. (1996) *Agrobacterium* VirE2 protein mediates nuclear
uptake of single-stranded DNA in plant cells. Proc Natl Acad Sci USA 93: 2392-2397

1107

1108

FIGURE LEGENDS

Figure 1. Some *Arabidopsis* actin and myosin genes are important for *Agrobacterium*-mediated transformation. (A) Root segments from *Arabidopsis* wild-type and *act2-1*, *act7-1*, *act7-4*, *act12*, *act7*-4 comp (complemented), or *bot1* mutants were infected with *A. tumefaciens* A208 (tumorigenic strain for stable transformation; left panel) or *A. tumefaciens* At849 (T-DNA contains a plant-active *gusA*-intron gene for transient transformation; right panel). After two days, the root segments were moved to medium (MS lacking phytohormones for stable transformation, CIM for transient transformation) containing timentin to kill the bacteria. For stable transformation, the presence of tumors was scored one month after infection. For transient transformation, root segments were stained with X-gluc after an additional four days, and the percentage of roots showing GUS activity was calculated. *act7-4* comp, *act7-4* mutant complemented with an *ACT7* cDNA. **(B and C)** Root segments from *Arabidopsis* wild-type and single or higher order *myosin VIII* or *XI* mutants were infected with *A. tumefaciens* At849 transiently or *A. tumefaciens* A208 stably. A total of 10-15 plants and >100 segments per plant were tested for stable transformation. Values given are means ± SE. Asterisks indicate significant differences compared to wild-type plants. [*t-test*, *$P < 0.05$; **$P < 0.01$].

Figure 2. **All Myosin VIII family members can contribute to *Agrobacterium*-mediated transformation.** Root segments from transgenic *Arabidopsis* individually expressing either an empty vector or *myosin VIII* cDNAs in *myosin VIII-1/2/a/b* mutant plants **(A)**, or in wild-type plants **(B)**, were infected with *A. tumefaciens* At849. Transgenic *Arabidopsis* expressing cDNAs encoding *myosins VIII-2, VIII-A, VIII-B, VIII-2+VIII-B* or *VIII-A+VIII-B* in the *myosin VIII-1/2/a/b* mutant background **(C)** or in wild-type Col0 plants **(D)** were similarly infected with *A. tumefaciens* At849. Transient transformation efficiencies are indicated as the percentage of roots showing GUS activity six days after infection. A total of 10-15 plants and >100 segments per plant, from five independent transgenic lines, were tested for transient transformation. Values given are means ± SE Asterisks indicate significant differences compared to control plants that contain an empty vector (**A** and **B**), the *myosin VIII-1/2/a/b* quadruple mutant (**C**), or wild-type Col-0 plants (**D**). [*t-test*, *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$; ****$P < 0.0001$].

Figure 3. **Inducible expression of myosin VIII, but not the myosin XI-K, cargo binding domains (CBD) inhibits transient transformation.** Root segments from transgenic *Arabidopsis* expressing the cargo binding domain (CBD) of the indicated myosin protein, under the control of a β-estradiol inducible promoter, were treated with β-estradiol or control solutions for 24 hr before infection with *A. tumefaciens* At849. The percentage of roots showing GUS activity six days after infection indicates the transient transformation efficiency. A total of 10-15 plants and >100 segments per plant, from five independent transgenic lines, were tested for transient transformation. Values given are means ± SE. Asterisks indicate significant differences compared to uninduced plants. [*t-test*, **$P < 0.01$; ***$P < 0.001$].

Figure 4. **Expression of VirE2-Venus in the *myosin VIII-1/2/a/b* mutant cannot complement a *virE2* mutant *Agrobacterium* strain for transient transformation.** **(A)** *Agrobacterium*-mediated transient transformation assays were conducted on roots of four independent transgenic lines of wild-type (Lines 2, 4, 5 and 8) or **(B)** the *myosin VIII-1/2/a/b* mutant (Lines 2, 3, 12, and 13). Root segments were inoculated with 10^6, 10^7, or 10^8 cfu/ml of the *virE2* mutant strain *A. tumefaciens* At1879 containing the T-DNA binary vector pBISN2. Plants were treated with β-estradiol (gray bars) or control solution (black bars) for VirE2-Venus expression 24 hr before infection. The percentage of roots showing GUS activity was calculated as in Figure 1. A total of 10-15 plants from each line and >100 segments per plant were tested for transient transformation. Values given are means ± SE. Asterisks indicate significant differences compared to uninduced plants. [*t-test*, *$P < 0.05$; **$P < 0.01$]. **(C)** Confocal images showing aggregation of VirE2-Venus proteins in transgenic roots of either wild-type (top panel) or *myosin VIII 1/2/a/b* mutant backgrounds (bottom panel). **(D)** Quantitative analysis of the size of VirE2-Venus aggregates and the percentage of the cellular area occupied by the aggregates. Image J was used for analysis. The average VirE2-Venus aggregate size was 2.0±0.3 μm^2 in wild-type roots, and 6.2±0.6 μm^2 in *myosin VIII-1/2/a/b* quadruple mutant roots. **(E)** Western blot detection of VirE2-Venus proteins expressed in transgenic plants of either the Col-0 or the *myosin VIII-1/2/a/b* quadruple myosin mutant background. Mouse anti-GFP antibody was used to detect VirE2-Venus protein expressed after induction of individual transgenic lines. The

house-keeping protein phosphoenolpyruvate carboxylase (PEPC) was detected using a rabbit

anti-PEPC antibody and served as an internal control.

Figure 5. **Analysis of VirE2-Venus aggregates in roots of transgenic *Arabidopsis* expressing**

the CBDs of various Myosin VIII proteins. Roots were treated with β-estradiol for eight hr.

Random areas of confocal images from roots expressing **(A)** Myosin VIII-1 or VIII-2 CBDs, or

(B) Myosin VIII-A or VIII-B CBDs were analyzed using Image J. Both the sizes and the

percentage of the cellular area occupied by the VirE2-Venus aggregates are shown.

Figure 6. **VirE2-Venus interacts with some myosin cargo binding domains (CBDs) *in vitro*.**

VirE2-Venus or Venus proteins were incubated with the indicated myc-tagged myosin CBD,

VIP1, or Lamin C. Following binding to anti-GFP beads, proteins were eluted and subjected to

Western blot analysis using anti-myc antibodies. Coomassie stained gels are shown in panels **A**,

C, and **E**, and immunoblots in panels **B**, **D**, and **F**. VirE2-Venus interacts with the CBDs of

myosins VIII-2, VIII-A, VIII-B, and XI-K, and with VIP1 (panels **B**, **D**, and **F**), but not with the

CBD of myosin VIII-1 or with Lamin C (panels **D** and **F**). Venus, as a negative control, does not

interact with any myc-tagged protein.

Figure 7. **The cargo binding domains (CBD) of some myosins interact with VirE2 in yeast.**

Yeast two-hybrid analysis was conducted with VirE2 as the bait and individual myosin CBDs as

the prey. Following co-transformation of the bait and prey plasmids, single colonies were picked

and serial dilutions of yeast were grown on **(A)** SD-Trp-Leu (-TL) medium, or **(B)** SD-Trp-Leu-

His medium supplemented with 3 mM 3-amino-1,2,4-triazole (-TLH+3AT). Colony growth on

SD-TLH+3AT indicates interaction between VirE2 and the myosin CBD. VIP1 was used as a

positive control for interaction with VirE2. Lamin C and the empty vector were used as negative

controls. SD, synthetic dropout medium.

Figure 8. **Myosin cargo binding domains can form complexes with VirE2 when expressed**

in *Arabidopsis*. (A) Protein extracts from roots of transgenic *Arabidopsis* plants expressing

VirE2-Venus, or Venus, and the indicated myc-tagged myosin CBD transgenes were incubated

with anti-GFP beads, the bound proteins eluted, and subjected to Western blot analysis using

anti-myc antibodies. Myc-tagged VIP1 was used as a positive control. **(B)** Root extracts

expressing VirE2-Venus or Venus and myc-tagged myosin VIII-2 CBD were incubated with

anti-GFP beads, the bound proteins eluted, and subjected to Western blot analysis using anti-myc

antibodies.

Figure 9. **Co-localization of VirE2-Venus and mCherry-myosin CBDs in *Arabidopsis* roots.**

(A) Roots of transgenic *Arabidopsis* plants containing VirE2-Venus and mCherry-myosin CBD

genes, each under the control of a β-estradiol-inducible promoter, were treated with β-estradiol

for 8 hr and examined by confocal microscopy. Bars indicate 50 μm. **(B)** Pearson's correlation

and overlap coefficient of Venus and mCherry fluorescence. A total of 10-15 transgenic lines

and >50 cells per construct were used for quantitative analysis using Image J. Values >0.5

indicate co-localization of the two proteins for both methods of analysis. The red line indicates a

co-localization ratio of 0.5.

Figure 10. **Bimolecular fluorescence complementation (BiFC) of VirE2 with various**

myosin CBDs in *Arabidopsis* roots. Transgenic *Arabidopsis* plants containing VirE2-cEYFP

and the indicated myosin CBDs-nVenus, each under the control of a β-estradiol-inducible

promoter, were treated with β-estradiol for 48 hr and the fluorescence signal detected by

confocal microscopy. VirE2 interacts with the CBDs of **(A),** myosin VIII-2; **(B),** myosin VIII-A;

and **(D),** myosin VIII-B at the plasma membrane. VirE2 interacts with the myosin XI-K CBD at

the plasma membrane and within the cytoplasm **(E)**. VirE2 does not interact with the myosin

VIII-1 CBD **(C)** or Lamin C **(F)**. Yellow fluorescence indicates interaction of VirE2-Venus with

the myosin CBD. Actin filaments are labeled by mCherry-ABD2. The figure shows merged

images of the Venus and mCherry signals. Bars indicate 50 μm.

1219

Figure 11. **Incubation of *Arabidopsis* roots with *Agrobacterium* that can transfer T-DNA**

results in increased VirE2-Venus in the perinuclear region. (A) Transgenic *Arabidopsis*

plants containing an inducible *VirE2-Venus* transgene were treated with 5 μm β-estradiol for 24

hr, followed by inoculation with 10^8 cfu/ml of the *virE2* mutant *A. tumefaciens* At1872 without a

binary vector (At2403, left panel) or with T-DNA on a binary vector (At2404, right panel). The

percentage of total VirE2-Venus fluorescence in the perinuclear area was calculated. Images of

10-15 transgenic lines and >50 cells were analyzed. The data indicate means ± SE. Asterisks

1227 indicate significant differences between roots inoculated with the two *Agrobacterium* strains.

1228 **(B-F)** the relative distribution of VirE2-Venus at the cell poles or in the cytoplasm following

1229 infection with *A. tumefaciens* At2403 or At2404 of roots co-expressing CBDs of myosin VIII-1.

1230 myosin VIII-2, myosin VIII-A, myosin VIII-B, or myosin XI-K (B, C, D, E, F, respectively).

1231 Images of 10-15 transgenic lines and >50 cells were analyzed. The data indicate means ± SE.

1232 Asterisks indicate significant differences between roots inoculated with the two *Agrobacterium*

1233 strains. [*t-test*, *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$].

1234

1235 **Figure 12**. **Loss of function of myosin XI-K, but not myosin VIII, disrupts movement of**

1236 **VirE2-Venus along actin filaments.** Transgenic plants expressing mCherry-ABD2

1237 constitutively and VirE2-Venus inducibly in Col-0, *myosin VIII-1/2/a/b*, and myosin XI-K CBD

1238 plants were used in this experiment. Roots of these plants were treated with 10 μm β-estradiol

1239 for 8 hr. Time lapse images were taken by confocal microscopy. Four images of each field are

1240 shown at the indicated times (0, 60, 90, and 120 seconds). Blue arrows indicate the initial

1241 position of a particular VirE2-Venus aggregate; white arrows indicate the position of the same

1242 aggregate at various times thereafter. Bars indicate 5 μm.

1243

1244 **Figure 13**. **Model of myosin VIII and myosin XI-K involvement in VirE2 trafficking to the**

1245 **perinuclear region. (A)** *Agrobacterium* virulence effector proteins, including VirE2 and VirE3,

1246 are secreted into host cells through a Type IV secretion system. VirE2 and VirE3 interact at the

1247 plant plasma membrane. Myosin VIII proteins localize at the plasma membrane and help tether

1248 VirE2 to the plasma membrane when no T-DNA is present. **(B)** When T-strands (as a VirD2/T-

1249 strand complex) enter the host cell, VirE2 molecules are released and some may bind T-DNA,

1250 following which VirE2 interacts with myosin XI-K and moves along actin filaments to the

1251 perinuclear region. Not shown are the *Agrobacterium* virulence effector proteins VirD5 and

1252 VirF which also are transferred to the plant.

1253
1254

1255 **Table 1.** Velocity of VirE2-Venus movement in leaf and root cells of various *Arabidopsis*
1256 transgenic lines

Transgene(s) in plant/ *Agrobacterium* infection[a]	*Arabidopsis* background	Velocity in leaves (μm/sec)	Velocity in roots (μm/sec)
VirE2-Venus	Col-0	0.353± 0.06	0.312± 0.07
VirE2-Venus	myosin *VIII-1/2/a/b*	0.378± 0.02	0.318± 0.02
VirE2-Venus + Myosin VIII-1 CBD	Col-0	0.407 ± 0.06	0.341 ± 0.04
VirE2-Venus + Myosin VIII-2 CBD	Col-0	0.398 ± 0.07	0.321 ± 0.02
VirE2-Venus + Myosin VIII-A CBD	Col-0	0.387 ± 0.07	0.304 ± 0.01
VirE2-Venus + Myosin VIII-B CBD	Col-0	0.401 ± 0.05	0.328 ± 0.04
VirE2-Venus + Myosin XI-K CBD	Col-0	Not detectable	Not detectable
VirE2-Venus + Myosin XI-K CBD + *Agrobacterium* (- T-DNA)	Col-0	Not detectable	Not detectable
VirE2-Venus + Myosin XI-K CBD + *Agrobacterium* (+ T-DNA)	Col-0	Not detectable	Not detectable

1257 [a]All transgenes are under the control of a β-estradiol-inducible promoter; CBD, cargo binding
1258 domain. >50 cells of 10 independent transgenic lines were examined for each genotype of
1259 transgenic plant.
1260

Figure 1. Some *Arabidopsis* actin and myosin genes are important for *Agrobacterium*-mediated root transformation.
(A) Root segments from *Arabidopsis* wild-type and *act2-1*, *act7-1*, act7-4, *act12*, *act7*-4 comp, or *bot1* mutants were infected with *A. tumefaciens* A208 (tumorigenic strain for stable transformation; left panel) or *A. tumefaciens* At849 (T-DNA contains a plant-active *gusA*-intron gene for transient transformation; right panel). After two days, the root segments were moved to medium (MS lacking phytohormones for stable transformation, CIM for transient transformation) containing timentin to kill the bacteria. For stable transformation, the presence of tumors was scored one month after infection. For transient transformation, root segments were stained with X-gluc after an additional four days, and the percentage of roots showing GUS activity was calculated. *act7-4* comp, *act7-4* mutant complemented with an *ACT7* cDNA. **(B and C)** Root segments from *Arabidopsis* wild-type and single or higher order *myosin VIII* or *XI* mutants were infected with *A. tumefaciens* At849 transiently or *A. tumefaciens* A208 stably. A total of 10-15 plants and >100 segments per plant were tested for stable transformation. Values given are means ± SE. Asterisks indicate significant differences compared to wild-type plants. [*t-test*, *$P < 0.05$; **$P < 0.01$].

Figure 2. **All Myosin VIII family members can contribute to *Agrobacterium*-mediated transformation.** Root segments from transgenic *Arabidopsis* individually expressing either an empty vector or *myosin VIII* cDNAs in *myosin VIII-1/2/a/b* mutant plants **(A)**, or in wild-type plants **(B)**, were infected with *A. tumefaciens* At849. Transgenic *Arabidopsis* expressing cDNAs encoding *myosins VIII-2, VIII-A, VIII-B, VIII-2+VIII-B* or *VIII-A+VIII-B* in the *myosin VIII-1/2/a/b* mutant background **(C)** or in wild-type Col-0 plants **(D)** were similarly infected with *A. tumefaciens* At849. Transient transformation efficiencies are indicated as the percentage of roots showing GUS activity six days after infection. A total of 10-15 plants and >100 segments per plant, from five independent transgenic lines, were tested for transient transformation. Values given are means ± SE. Asterisks indicate significant differences compared to control plants that contain an empty vector **(A** and **B)**, the *myosin VIII-1/2/a/b* quadruple mutant **(C)**, or wild-type Col-0 plants **(D)**. [*t-test*, *$P<0.05$; **$P<0.01$; ***$P<0.001$; ****$P<0.0001$].

Figure 3. **Inducible expression of myosin VIII, but not the myosin XI-K, cargo binding domains (CBD) inhibits transient transformation.** Root segments from transgenic *Arabidopsis* expressing the cargo binding domain (CBD) of the indicated myosin protein, under the control of a β-estradiol inducible promoter, were treated with β-estradiol or control solutions for 24 hr before infection with *A. tumefaciens* At849. The percentage of roots showing GUS activity six days after infection indicates the transient transformation efficiency. A total of 10-15 plants and >100 segments per plant, from five independent transgenic lines, were tested for transient transformation. Values given are means ± SE. Asterisks indicate significant differences compared to uninduced plants. [*t-test*, **$P < 0.01$; ***$P < 0.001$].

- **Figure 4**. **Expression of VirE2-Venus in the *myosin VIII-1/2/a/b* mutant cannot complement a *virE2* mutant *Agrobacterium* strain for transient transformation.**   **(A)** *Agrobacterium*-mediated transient transformation assays were conducted on roots of four independent transgenic lines of wild-type (Lines 2, 4, 5 and 8) or **(B)** the *myosin VIII-1/2/a/b* mutant (Lines 2, 3, 12, and 13). Root segments were inoculated with 10^6, 10^7, or 10^8 cfu/ml of the *virE2* mutant strain *A. tumefaciens* At1879 containing the T-DNA binary vector pBISN2. Plants were treated with β-estradiol (gray bars) or control solution (black bars) for VirE2-Venus expression 24 hr before infection. The percentage of roots showing GUS activity was calculated as in Figure 1. A total of 10-15 plants from each line and >100 segments per plant were tested for transient transformation.   Values given are means ± SE.   Asterisks indicate significant differences compared to uninduced plants. [*t-test*, $*P < 0.05$; $**P < 0.01$]. **(C)** Confocal images showing aggregation of VirE2-Venus proteins in transgenic roots of either wild-type (top panel) or *myosin VIII 1/2/a/b* mutant backgrounds (bottom panel). **(D)** Quantitative analysis of the size of VirE2-Venus aggregates and the percentage of the cellular area occupied by the aggregates.   Image J was used for analysis. The average VirE2-Venus aggregate size was 2.0±0.3 μm^2 in wild-type roots, and 6.2±0.6 μm^2 in *myosin VIII-1/2/a/b* quadruple mutant roots. **(E)** Western blot detection of VirE2-Venus proteins expressed in transgenic plants of either the Col-0 or the *myosin VIII-1/2/a/b* quadruple myosin mutant background.   Mouse anti-GFP antibody was used to detect VirE2-Venus protein expressed after induction of individual transgenic lines.   The house-keeping protein phosphoenolpyruvate carboxylase (PEPC) was detected using a rabbit anti-PEPC antibody and served as an internal control.

Figure 5. **Analysis of VirE2-Venus aggregates in roots of transgenic *Arabidopsis* expressing the CBDs of various Myosin VIII proteins.** Roots were treated with β-estradiol for eight hr. Random areas of confocal images from roots expressing **(A)** Myosin VIII-1 or VIII-2 CBDs, or **(B)** Myosin VIII-A or VIII-B CBDs were analyzed using Image J. Both the sizes and the percentage of the cellular area occupied by the VirE2-Venus aggregates are shown.

Figure 6. **VirE2-Venus interacts with some myosin cargo binding domains (CBDs)** *in vitro*. VirE2-Venus or Venus proteins were incubated with the indicated myc-tagged myosin CBD, VIP1, or Lamin C. Following binding to anti-GFP beads, proteins were eluted and subjected to Western blot analysis using anti-myc antibodies. Coomassie stained gels are shown in panels **A**, **C**, and **E**, and immunoblots in panels **B**, **D**, and **F**. VirE2-Venus interacts with the CBDs of myosins VIII-2, VIII-A, VIII-B, and XI-K, and with VIP1 (panels **B**, **D**, and **F**), but not with the CBD of myosin VIII-1 or with Lamin C (panels **D** and **F**). Venus, as a negative control, does not interact with any myc-tagged protein.

Figure 7. The cargo binding domains (CBD) of some myosins interact with VirE2 in yeast. Yeast two-hybrid analysis was conducted with VirE2 as the bait and individual myosin CBDs as the prey. Following co-transformation of the bait and prey plasmids, single colonies were picked and serial dilutions of yeast were grown on **(A)** SD-Trp-Leu (-TL) medium, or **(B)** SD-Trp-Leu-His medium supplemented with 3 mM 3-amino-1,2,4-triazole (-TLH+3AT). Colony growth on SD-TLH+3AT indicates interaction between VirE2 and the myosin CBD. VIP1 was used as a positive control for interaction with VirE2. Lamin C and the empty vector were used as negative controls. SD, synthetic dropout medium.

Figure 8. **Myosin cargo binding domains can form complexes with VirE2 when expressed in *Arabidopsis*.** **(A)** Protein extracts from roots of transgenic *Arabidopsis* plants expressing VirE2-Venus, or Venus, and the indicated myc-tagged myosin CBD transgenes were incubated with anti-GFP beads, the bound proteins eluted, and subjected to Western blot analysis using anti-myc antibodies. Myc-tagged VIP1 was used as a positive control. **(B)** Root extracts expressing VirE2-Venus or Venus and myc-tagged myosin VIII-2 CBD were incubated with anti-GFP beads, the bound proteins eluted, and subjected to Western blot analysis using anti-myc antibodies.

Figure 9. **Co-localization of VirE2-Venus and mCherry-myosin CBDs in *Arabidopsis* roots.** **(A)** Roots of transgenic *Arabidopsis* plants containing VirE2-Venus and mCherry-myosin CBD genes, each under the control of a β-estradiol-inducible promoter, were treated with β-estradiol for 8 hr and examined by confocal microscopy. Bars indicate 50 μm. **(B)** Pearson's correlation and overlap coefficient of Venus and mCherry fluorescence. A total of 10-15 transgenic lines and >50 cells per construct were used for quantitative analysis using Image J. Values >0.5 indicate co-localization of the two proteins for both methods of analysis. The red line indicates a co-localization ratio of 0.5.

Figure 10. **Bimolecular fluorescence complementation (BiFC) of VirE2 with various myosin CBDs in** *Arabidopsis* **roots.** Transgenic *Arabidopsis* plants containing VirE2-cEYFP and the indicated myosin CBDs-nVenus, each under the control of a β-estradiol-inducible promoter, were treated with β-estradiol for 48 hr and the fluorescence signal detected by confocal microscopy. VirE2 interacts with the CBDs of **(A),** myosin VIII-2; **(B),** myosin VIII-A; and **(D),** myosin VIII-B at the plasma membrane. VirE2 interacts with the myosin XI-K CBD at the plasma membrane and within the cytoplasm **(E)**. VirE2 does not interact with the myosin VIII-1 CBD **(C)** or Lamin C **(F)**. Yellow fluorescence indicates interaction of VirE2-Venus with the myosin CBD. Actin filaments are labeled by mCherry-ABD2. The figure shows merged images of the Venus and mCherry signals. Bars indicate 50 μm.

Figure 11. **Incubation of *Arabidopsis* roots with *Agrobacterium* that can transfer T-DNA results in increased VirE2-Venus in the perinuclear region.** **(A)** Transgenic *Arabidopsis* plants containing an inducible *VirE2-Venus* transgene were treated with 5 μm β-estradiol for 24 hr, followed by inoculation with 10^8 cfu/ml of the *virE2* mutant *A. tumefaciens* At1872 without a binary vector (At2403, left panel) or with T-DNA on a binary vector (At2404, right panel). The percentage of total VirE2-Venus fluorescence in the perinuclear area was calculated. Images of 10-15 transgenic lines and >50 cells were analyzed. The data indicate means ± SE. Asterisks indicate significant differences between roots inoculated with the two *Agrobacterium* strains. **(B-F)** the relative distribution of VirE2-Venus at the cell poles or in the cytoplasm following infection with *A. tumefaciens* At2403 or At2404 of roots co-expressing CBDs of myosin VIII-1. myosin VIII-2, myosin VIII-A, myosin VIII-B, or myosin XI-K (B, C, D, E, F, respectively). Images of 10-15 transgenic lines and >50 cells were analyzed. The data indicate means ± SE. Asterisks indicate significant differences between roots inoculated with the two *Agrobacterium* strains. [*t-test*, *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$].

Figure 12. **Loss of function of myosin XI-K, but not myosin VIII, disrupts movement of VirE2-Venus along actin filaments.** Transgenic plants expressing mCherry-ABD2 constitutively and VirE2-Venus inducibly in Col-0, *myosin VIII-1/2/a/b*, and myosin XI-K CBD plants were used in this experiment. Roots of these plants were treated with 10 μM β-estradiol for 8 hr. Time lapse images were taken by confocal microscopy. Four images of each field are shown at the indicated times (0, 60, 90, and 120 seconds). Blue arrows indicate the initial position of a particular VirE2-Venus aggregate; white arrows indicate the position of the same aggregate at various times thereafter. Bars indicate 5 μm.

Figure 13. **Model of myosin VIII and myosin XI-K involvement in VirE2 trafficking to the perinuclear region.** (A) *Agrobacterium* virulence effector proteins, including VirE2 and VirE3, are secreted into host cells through a Type IV secretion system. VirE2 and VirE3 interact at the plant plasma membrane. Myosin VIII proteins localize at the plasma membrane and help tether VirE2 to the plasma membrane when no T-DNA is present. (B) When T-strands (as a VirD2/T-strand complex) enter the host cell, VirE2 molecules are released and some may bind T-DNA, following which VirE2 interacts with myosin XI-K and moves along actin filaments to the perinuclear region. Not shown are the *Agrobacterium* virulence effector proteins VirD5 and VirF which also are transferred to the plant.

Supplemental Table 1. Plasmids used in this study.

Plasmid	Important features/proteins	Plasmid description	Antibiotic resistance	Reference
pE886	Cloning vector	pBluescript KS+	Amp	Kozaki et al., 2004
pE3086	EYFP	pSAT6-P$_{35S}$-cEYFP-N1	Amp	Citovsky et al., 2006
pE3129	PIP2A	pSAT6-PIP2A-mRFP	Amp	Gelvin lab strain
pE3183	myc	pSAT6-P$_{35S}$-myc-MCS	Amp	Gelvin lab strain
pE3230	nVenus	pSAT6-P$_{35S}$-nVenus-C	Amp	Lee and Gelvin 2014
pE3275	mCherry	pSAT6-P$_{35S}$-mCherry-C1-B	Amp	Lee and Gelvin 2014
pE3341	mCherry	pBBR1-MCS2::mCherry	Kan	From Clay Fuqua
pE3447	Lamin C	pBSKS-Lamin C	Amp	Gelvin lab strain
pE3519	Binary vector	pPZP-*bar*-RCS2	Spec	Gelvin lab strain
pE3522	Yeast bait vector	pGBKT7	Kan	Lu et al., 2010
pE3542	Venus	pSAT1-Venus-C	Amp	Lee and Gelvin 2014
pE3745	Yeast prey vector	pGADT7	Amp	Chien et al., 1991
pE3802	For N-terminal nVenus tagging of myosin XI-K CBD	pSAT6-P$_{35S}$-nVenus-Myosin XI-K CBD	Amp	Gelvin lab strain

pE3982	VirE2-cEYFP	pSAT1-P35S-VirE2-cEYFP	Amp	Gelvin lab strain
pE4014	VirE2-cCFP	*Nco*I-*Bam*HI fragment from pSAT4-VirE2-cCFP cloned into the same sites in pGBKT7	Kan	Gelvin lab strain
pE4224	β-estradiol-inducible promoter	pSAT1-β-estradiol-inducible promoter	Amp	Gelvin lab strain
pE4282	VirE2-Venus	pSAT1-Pi-VirE2-Venus	Amp	Gelvin lab strain
pE4292	VirE2-Venus, mCherry-ABD2	pPZP-Pi-VirE2-Venus-P_{nos}-mCherry-ABD2	Spec	Gelvin lab strain
pE4317	Binary vector; VirE2-Venus, hptII	pPZP-Pi-VirE2-Venus-*hptII*	Spec	Gelvin lab strain
pE4381	mCherry-intron-NLS	pPZP-P35S-mCherry-intron-NLS	Spec	Gelvin lab strain
pE4437	Binary vector; Cerulean-NLS, mCherry-ABD2	pPZP -P_{nos}-Cerulean-NLS-P35S-mCherry-ABD2	Spec	Gelvin lab strain
pE4438	VirE2-Venus, Cerulean-NLS	pPZP-Pi-VirE2-Venus-P_{nos}-Cerulean-NLS	Spec	Gelvin lab strain
pE4515	pSAT1-P35S	pSAT1-MCS	Kan	Gelvin lab strain
pE4672	Binary vector; *hptII*	pPZP-P_{nos}-*hptII*	Spec	Lee and Gelvin, 2014

pE4712	Myosin VIII-1 CBD	PCR fragment of Myosin VIII-1 CBD with *Kpn*I/*Bam*HI sites, cloned into the *Eco*RV site of pBluescript KS+ to make pBS-Myosin VIII-1 CBD	Amp	This study
pE4713	Myosin VIII-2 CBD	PCR fragment of Myosin VIII-2 CBD with *Kpn*I/*Bam*HI sites, cloned into the *Eco*RV site of pBluescript KS+ to make pBS-Myosin VIII-2 CBD	Amp	This study
pE4714	Myosin VIII-A CBD	PCR fragment of Myosin VIII-A CBD with *Kpn*I/*Bam*HI sites, cloned into the *Eco*RV site of pBluescript KS+ to make pBS-Myosin VIII-A CBD	Amp	This study
pE4715	Myosin VIII-B CBD	PCR fragment of Myosin VIII-B CBD with *Kpn*I/*Bam*HI sites, cloned into the *Eco*RV site of pBluescript KS+ to make pBS-Myosin VIII-B CBD	Amp	This study
pE4716	β-estradiol-inducible promoter	PCR fragment of inducible promoter from pE4224 with *Age*I/*Eco*RV sites, cloned into the *Eco*RV site of pBluescript KS+ to make pBS-Pi promoter	Amp	This study
pE4720	pSAT6-Pi-mCherry-C	*Age*I+*Eco*RV digested pE4716, cloned into *Age*I+*Eco*RV digested pE3275, to make pSAT6-Pi-mCherry	Amp	This study
pE4721	mCherry-Myosin VIII-1 CBD	*Kpn*I+*Bam*HI digested pE4712 cloned into *Kpn*I+*Bam*HI digested pE4720 to make pSAT6-Pi-mCherry-Myosin VIII-1 CBD	Amp	This study
pE4722	mCherry-Myosin VIII-2 CBD	*Kpn*I+*Bam*HI digested pE4713 cloned into *Kpn*I+*Bam*HI digested pE4720 to make pSAT6-Pi-mCherry-Myosin VIII-2 CBD	Amp	This study
pE4723	mCherry-Myosin VIII-A CBD	*Kpn*I+*Bam*HI digested pE4714 cloned into *Kpn*I+*Bam*HI digested pE4720 to make pSAT6-Pi-mCherry-Myosin VIII-A CBD	Amp	This study
pE4724	mCherry-Myosin VIII-B CBD	*Kpn*I+*Bam*HI digested pE4715 cloned into *Kpn*I+*Bam*HI digested pE4720 to make pSAT6-Pi-mCherry-Myosin VIII-B CBD	Amp	This study

pE4725	mCherry-Myosin XI-K CBD	*Kpn*I+*Pst*I (Blunt) digested pE3802 cloned into *Kpn*I+*Bam*HI (Blunt) digested pE4720 to make pSAT6-Pi-mCherry-Myosin XI-K CBD	Amp	This study
pE4726	mCherry-Myosin VIII-1 CBD, VirE2-Venus, Cerulean-NLS	PI-*Psp*I digested pE4721 cloned into PI-*Psp*I digested pE4438 to make pPZP-Pi-VirE2-Venus-P$_{nos}$-Cerulean-NLS-Pi-mCherry-Myosin VIII-1 CBD	Spec	This study
pE4727	mCherry-Myosin VIII-2 CBD, VirE2-Venus, Cerulean-NLS	PI-*Psp*I digested pE4722 cloned into PI-*Psp*I digested pE4438 to make pPZP-Pi-VirE2-Venus-P$_{nos}$-Cerulean-NLS-Pi-mCherry-Myosin VIII-2 CBD	Spec	This study
pE4728	mCherry-Myosin VIII-A CBD, VirE2-Venus, Cerulean-NLS	PI-*Psp*I digested pE4723 cloned into PI-*Psp*I digested pE4438 to make pPZP-Pi-VirE2-Venus-P$_{nos}$-Cerulean-NLS-Pi-mCherry-Myosin VIII-A CBD	Spec	This study
pE4729	mCherry-Myosin VIII-B CBD, VirE2-Venus, Cerulean-NLS	PI-*Psp*I digested pE4724 cloned into PI-*Psp*I digested pE4438 to make pPZP-Pi-VirE2-Venus-P$_{nos}$-Cerulean-NLS-Pi-mCherry-Myosin VIII-B CBD	Spec	This study
pE4730	mCherry-Myosin XI-K CBD, VirE2-Venus, Cerulean-NLS	PI-*Psp*I digested pE4725 cloned into PI-*Psp*I digested pE4438 to make pPZP-Pi-VirE2-Venus-P$_{nos}$-Cerulean-NLS-Pi-mCherry-Myosin XI-K CBD	Spec	This study
pE4731	pSAT6-Pi-nVenus-C	*Age*I+*Eco*RV digested pE4716 cloned into *Age*I+*Eco*RV digested pE3230 to make pSAT6-Pi-nVenus	Amp	This study
pE4732	VirE2-cEYFP	*Age*I+*Swa*I digested pE4731 cloned into *Age*I+*Swa*I digested pE3982 to make pSAT1-Pi-VirE2-cEYFP	Amp	This study
pE4733	nVenus-Myosin VIII-1 CBD	*Kpn*I+*Bam*HI digested pE4712 cloned into *Kpn*I+*Bam*HI digested pE4731 to make pSAT6-Pi-nVenus-Myosin VIII-1 CBD	Amp	This study
pE4734	nVenus-Myosin VIII-2 CBD	*Kpn*I+*Bam*HI digested pE4713 cloned into *Kpn*I+*Bam*HI digested pE4731 to make pSAT6-Pi-nVenus-Myosin VIII-2 CBD	Amp	This study
pE4735	nVenus-Myosin VIII-A CBD	*Kpn*I+*Bam*HI digested pE4714 cloned into *Kpn*I+*Bam*HI digested pE4731 to make pSAT6-Pi-nVenus-Myosin VIII-A CBD	Amp	This study
pE4736	nVenus-Myosin VIII-B CBD	*Kpn*I+*Bam*HI digested pE4715 cloned into *Kpn*I+*Bam*HI digested pE4731 to make pSAT6-Pi-nVenus-Myosin VIII-B CBD	Amp	This study

pE4737	nVenus-Myosin XI-K CBD	*Kpn*I+*Not*I digested pE4725 cloned into *Kpn*I+*Not*I digested pE4731 to make pSAT6-Pi-nVenus-Myosin XI-K CBD	Amp	This study
pE4738	VirE2-cYFP, Cerulean-NLS, mCherry-ABD2	*Asc*I digested pE4732 cloned into *Asc*I digested pE4437, to make pPZP-Pi-VirE2-cYFP-P_{nos}-Cerulean-NLS-P_{35S}-mCherry-ABD2	Spec	This study
pE4739	pSAT1-Pi-mCherry	*Xho*I (Blunt)/*Not*I digested pE4424 cloned into *Nco*I (Blunt)/*Not*I digested pE3275 to make pSAT1-inducible promoter-mCherry	Amp	This study
pE4755	nVenus-Myosin VIII-A CBD, VirE2-cYFP, Cerulean-NLS, mCherry-ABD2	PI-*Psp*I digested pE4735 cloned into PI-*Psp*I digested pE4738 to make pPZP-Pi-VirE2-cYFP-P_{nos}-Cerulean-NLS-P_{35S}-mCherry-ABD2-Pi-nVenus-myosin VIII-A CBD	Spec	This study
pE4756	nVenus-Myosin VIII-B CBD, VirE2-cYFP, Cerulean-NLS, mCherry-ABD2	PI-*Psp*I digested pE4736 cloned into PI-*Psp*I digested pE4738 to make pPZP-Pi-VirE2-cYFP-P_{nos}-Cerulean-NLS-P_{35S}-mCherry-ABD2-Pi-nVenus-myosin VIII-B CBD	Spec	This study
pE4757	nVenus-Myosin XI-K CBD, VirE2-cYFP, Cerulean-NLS, mCherry-ABD2	PI-*Psp*I digested pE4737 cloned into PI-*Psp*I digested pE4738 to make pPZP-Pi-VirE2-cYFP-P_{nos}-Cerulean-NLS-P_{35S}-mCherry-ABD2-Pi-nVenus-myosin XI-K CBD	Spec	This study
pE4758	nVenus-Myosin VIII-1 CBD, VirE2-cYFP, Cerulean-NLS, mCherry-ABD2	PI-*Psp*I digested pE4733 cloned into PI-*Psp*I digested pE4738 to make pPZP-Pi-VirE2-cYFP-P_{nos}-Cerulean-NLS-P_{35S}-mCherry-ABD2-Pi-nVenus-myosin VIII-1 CBD	Spec	This study
pE4759	nVenus-Myosin VIII-2 CBD, VirE2-cYFP, Cerulean-NLS, mCherryABD2	PI-*Psp*I digested pE4734 cloned into PI-*Psp*I digested pE4738 to make pPZP-Pi-VirE2-cYFP-P_{nos}-Cerulean-NLS-P_{35S}-mCherry-ABD2-Pi-nVenus-myosin VIII-2 CBD	Spec	This study
pE4774	Myc	PCR fragment of myc tag (pE3183 as template) with *Nco*I and *Not*I (35S terminator), cloned into *Eco*RV site of pBluesescript KS+ (pE886) to make pBS-Myc	Amp	This study
pE4775	Myc	*Nco*I+*Not*I digested pBS-Myc cloned into *Nco*I+*Not*I digested pE4720, to make Pi-pSAT6-myc	Amp	This study

pE4776	Myc-Myosin VIII-1 CBD	*Kpn*I+*Bam*HI digested pE4712 cloned into *Kpn*I+*Bam*HI digested pE4775 to make pSAT6-Pi-myc-Myosin VIII-1 CBD	Amp	This study
pE4777	Myc-Myosin VIII-2 CBD	*Kpn*I+*Bam*HI digested pE4713 cloned into *Kpn*I+*Bam*HI digested pE4775 to make pSAT6-Pi-myc-Myosin VIII-2 CBD	Amp	This study
pE4778	Myc-Myosin VIII-A CBD	*Kpn*I+*Bam*HI digested pE4714 cloned into *Kpn*I+*Bam*HI digested pE4775 to make pSAT6-Pi-myc-Myosin VIII-A CBD	Amp	This study
pE4779	Myc-Myosin VIII-B CBD	*Kpn*I+*Bam*HI digested pE4715 cloned into *Kpn*I+*Bam*HI digested pE4775 to make pSAT6-Pi-myc-Myosin VIII-B CBD	Amp	This study
pE4780	Myc-Myosin XI-K CBD	*Kpn*I+*Not*I digested pE4725 cloned into *Kpn*I+*Not*I digested pE4775 to make pSAT6-Pi-myc-Myosin XI-K CBD	Amp	This study
pE4781	mCherry-Myosin VIII-1 CBD, Cerulean-NLS	*Asc*I digested pE4726 self-ligated to make pPZP-Pnos-Cerulean-NLS-Pi-mCherry-Myosin VIII-1 CBD	Spec	This study
pE4782	mCherry-Myosin VIII-2 CBD, Cerulean-NLS	*Asc*I digested pE4727 self-ligated to make pPZP-Pnos-Cerulean-NLS-Pi-mCherry-Myosin VIII-2 CBD	Spec	This study
pE4783	mCherry-Myosin VIII-A CBD, Cerulean-NLS	*Asc*I digested pE4728 self-ligated to make pPZP-Pnos-Cerulean-NLS-Pi-mCherry-Myosin VIII-A CBD	Spec	This study
pE4784	mCherry-Myosin VIII-B CBD, Cerulean-NLS	*Asc*I digested pE472 self-ligated to make pPZP-Pnos-Cerulean-NLS-Pi-mCherry-Myosin VIII-B CBD	Spec	This study
pE4785	mCherry-Myosin XI-K CBD, Cerulean-NLS	*Asc*I digested pE4730 self-ligated to make pPZP-Pnos-Cerulean-NLS-Pi-mCherry-Myosin XI-K CBD	Spec	This study
pE4786	Myc-Myosin VIII-1 CBD, VirE2-Venus	PI-*Psp*I digested pE4776 cloned into PI-*Psp*I digested pE4317 to make pPZP-Pi-VirE2-Venus-Pi-myc-myosin VIII-1 CBD	Spec	This study
pE4787	Myc-Myosin VIII-2 CBD, VirE2-Venus	PI-*Psp*I digested pE4777 cloned into PI-*Psp*I digested pE4317 to make pPZP-Pi-VirE2-Venus-Pi-myc-myosin VIII-2 CBD	Spec	This study

pE4788	Myc-Myosin VIII-A CBD, VirE2-Venus	PI-*Psp*I digested pE4778 cloned into PI-*Psp*I digested pE4317 to make pPZP-Pi-VirE2-Venus-Pi-myc-myosin VIII-A CBD	Spec	This study
pE4789	Myc-Myosin VIII-B CBD, VirE2-Venus	PI-*Psp*I digested pE4779 cloned into PI-*Psp*I digested pE4317 to make pPZP-Pi-VirE2-Venus-Pi-myc-myosin VIII-B CBD	Spec	This study
pE4790	Myc-Myosin XI-K CBD, VirE2-Venus	PI-*Psp*I digested pE4780 cloned into PI-*Psp*I digested pE4317 to make pPZP-Pi-VirE2-Venus-Pi-myc-myosin XI-K CBD	Spec	This study
pE4791	New inducible promoter	*Age*I+*Swa*I digested pE4754 cloned into *Age*I+*Swa*I digested pE4731 to make pSAT6-Pi(New)-nVenus	Amp	This study
pE4792	nVenus-Lamin C	*Sma*I+*Bam*HI digested pE3447 cloned into *Sma*I+*Bam*HI digested pE4791 to make pSAT6-Pi(new)-nVenus-Lamin C	Amp	This study
pE4814	Venus	*Nco*I+*Not*I digested pE3542 cloned into *Nco*I+*Not*I digested pET28a, to make pET28a-Venus	Kan	This study
pE4815	VirE2-Venus	*Nco*I+*Not*I digested pE4282 cloned into *Nco*I/+*Not*I digested pET28a, to make pET28a-VirE2-Venus	Kan	This study
pE4816	Myc-Myosin VIII-1 CBD	*Nco*I+*Bam*HI digested pE4776 cloned into *Nco*I+*Bam*HI digested pET28a to make pET28a-myc-myosin VIII-1 CBD	Kan	This study
pE4817	Myc-Myosin VIII-2 CBD	*Nco*I+*Bam*HI digested pE4777 cloned into *Nco*I+*Bam*HI digested pET28a to make pET28a-myc-myosin VIII-2 CBD	Kan	This study
pE4818	Myc-Myosin VIII-A CBD	*Nco*I+*Bam*HI digested pE4778 cloned into *Nco*I+*Bam*HI digested pET28a to make pET28a-myc-myosin VIII-A CBD	Kan	This study
pE4819	Myc-Myosin VIII-B CBD	*Nco*I+*Bam*HI digested pE4779 cloned into *Nco*I+*Bam*HI digested pET28a to make pET28a-myc-myosin VIII-B CBD	Kan	This study
pE4820	Myc-Myosin XI-K CBD	*Nco*I+*Not*I digested pE4780 cloned into *Nco*I+*Not*I digested pET28a to make pET28a-myc-myosin XI-K CBD	Kan	This study
pE4821	Myc-VIP1	*Xho*I+*Bam*HI digested pE3395 cloned into *Xho*I+*Bam*HI digested pE4775 to make Pi-pSAT6-myc-VIP1	Amp	This study

pE4822	Myc-VIP1	*Nco*I+*Bam*HI digested pE4821 cloned into *Nco*I+*Bam*HI digested pET28a to make pET28a-myc-VIP1	Kan	This study
pE4831	Gal4BD-Myosin VIII-1 CBD	*Nco*I+*Bam*HI digested pE4776 cloned into *Nco*I+*Bam*HI digested pE3745 to make pGADT7-myosin VIII-1 CBD	Amp	This study
pE4832	Gal4BD-Myosin VIII-2 CBD	*Nco*I+*Bam*HI digested pE4777 cloned into *Nco*I+*Bam*HI digested pE3745 to make pGADT7-myosin VIII-2 CBD	Amp	This study
pE4833	Gal4BD-Myosin VIII-A CBD	*Nco*I+*Bam*HI digested pE4778 cloned into *Nco*I+*Bam*HI digested pE3745 to make pGADT7-myosin VIII-A CBD	Amp	This study
pE4834	Gal4BD-Myosin VIII-B CBD	*Nco*I+*Bam*HI digested pE4779 cloned into *Nco*I+*Bam*HI digested pE3745 to make pGADT7-myosin VIII-B CBD	Amp	This study
pE4835	Gal4BD-Myosin XI-K CBD	*Pst*I-Blunt+*Bam*HI digested pE3082 cloned into *Eco*R53KI+*Bam*HI digested pE3745 to make pGADT7-myosin XI-K CBD	Amp	This study
pE4836	Gal4BD-VIP1	*Nco*I+*Bam*HI digested pE4821 cloned into *Nco*I+*Bam*HI digested pE3745 to make pGADT7-VIP1	Amp	This study
pE4837	Myc-Lamin C	*Sma*I+*Bam*HI digested pE4822 cloned into *Sma*I+*Bam*HI digested pE4792 to make pET28a-myc-Lamin C	Kan	This study
pE4843	Myosin VIII-1	PCR fragment of Myosin VIII-1 full length with *Kpn*I/*Bam*HI sites, cloned into *Eco*RV site of pBlusescript KS+ to make pBS-Myosin VIII-1 full length	Amp	This study
pE4844	Myosin VIII-2	PCR fragment of Myosin VIII-2 full length with *Kpn*I/*Sma*I sites, cloned into *Eco*RV site of pBlusescript KS+ to make pBS-Myosin VIII-2 full length	Amp	This study
pE4845	Myosin VIII-A	PCR fragment of Myosin VIII-A full length with *Kpn*I/*Bam*HI sites, cloned into *Eco*RV site of pBlusescript KS+ to make pBS-Myosin VIII-A full length	Amp	This study

pE4846	Myosin VIII-B	PCR fragment of Myosin VIII-B full length with *Kpn*I/*Sma*I sites, cloned into *Eco*RV site of pBluescript KS+ to make pBS-Myosin VIII-B full length	Amp	This study
pE4847	Myosin VIII-1	*Kpn*I+*Bam*HI digested pE4843 cloned into *Kpn*I+*Bam*HI digested pE3086 to make pSAT6-myosin VIII-1 full length	Amp	This study
pE4848	Myosin VIII-B	*Kpn*I+*Sma*I digested pE4846 cloned into *Kpn*I+*Sma*I digested pE3086 to make pSAT6-myosin VIII-B full length	Amp	This study
pE4849	Myc-Myosin VIII-1 CBD, Venus	*Asc*I digested pE3542 cloned into *Asc*I digested pE4786 to make pPZP-35S-Venus-Pi-myc-myosin VIII-1 CBD	Spec	This study
pE4850	Myc-Myosin VIII-2 CBD, Venus	*Asc*I digested pE3542 cloned into *Asc*I digested pE4787 to make pPZP-35S-Venus-Pi-myc-myosin VIII-2 CBD	Spec	This study
pE4851	Myc-Myosin VIII-A CBD, Venus	*Asc*I digested pE3542 cloned into *Asc*I digested pE4788 to make pPZP-35S-Venus-Pi-myc-myosin VIII-A CBD	Spec	This study
pE4852	Myc-Myosin VIII-2 CBD, VirE2-Venus, mCherry-ABD2	PI-*Psp*I digested pE4777 cloned into PI-*Psp*I digested pE4292 to make pPZP-Pi-VirE2-Venus-P_{nos}-mCherry-ABD2-Pi-myc-myosin VIII-2 CBD	Spec	This study
pE4853	Myc-Myosin VIII-A CBD, VirE2-Venus, mCherry-ABD2	PI-*Psp*I digested pE4778 cloned into PI-*Psp*I digested pE4292 to make pPZP-Pi-VirE2-Venus-P_{nos}-mCherry-ABD2-Pi-myc-myosin VIII-A CBD	Spec	This study
pE4854	Myosin VIII-1, hptII	PI-*Psp*I digested pE4847 cloned into PI-*Psp*I digested pE4672 to make pPZP-hpt-XVE-P35S-myosin VIII-1 full length	Spec	This study
pE4855	Myc-Myosin VIII-1 CBD, VirE2-Venus, mCherry-ABD2	PI-*Psp*I digested pE4776 cloned into PI-*Psp*I digested pE4292 to make pPZP-Pi-VirE2-Venus-P_{nos}-mCherry-ABD2-Pi-myc-myosin VIII-1 CBD	Spec	This study
pE4856	Myc-Myosin VIII-B CBD, VirE2-Venus, mCherry-ABD2	PI-*Psp*I digested pE4779 cloned into PI-*Psp*I digested pE4292 to make pPZP-Pi-VirE2-Venus-P_{nos}-mCherry-ABD2-Pi-myc-myosin VIII-B CBD	Spec	This study

pE4857	Myosin VIII-B, hptII	PI-*Psp*I digested pE4848 cloned into PI-*Psp*I digested pE4672 to make pPZP-*hptII*-XVE-P35S-myosin VIII-B full length	Spec	This study
pE4858	Myosin VIII-A	*Kpn*I+*Bam*HI digested pE4845 cloned into *Kpn*I+*Bam*HI digested pE4515 to make pSAT1-Myosin VIII-A full length	Kan	This study
pE4859	PIP2A-Venus	*Xho*I+*Bam*HI digested pE3129 (PIP2A) cloned into *Xho*I+*Bam*HI digested pE3758 to make pSAT1-PIP2A-Venus	Amp	This study
pE4874	Myosin VIII-A, *hptII*	*Asc*I digested pE4858 cloned into *Asc*I digested pE4672 to make pPZP-XVE-P35S-myosin VIII-A full length	Spec	This study
pE4875	Myosin VIII-2	*Kpn*I+*Sma*I digested pE4844 cloned into *Kpn*I+*Sma*I digested pE4515 to make pSAT1-Myosin VIII-2 full length	Kan	This study
pE4876	Myosin VIII-2, *hptII*	*Asc*I digested pE4875 cloned into *Asc*I digested pE4672 to make pPZP-XVE-P35S-myosin VIII-2 full length	Spec	This study
pE4877	mCherry-Myosin VIII-1 CBD, Cerulean, PIP2A-Venus	*Asc*I digested pE4859 cloned into *Asc*I digested pE4781 to make pPZP-P$_{nos}$-Cerulean-P35S-PIP2A-Venus-Pi-mCherry-myosin VIII-1 CBD	Spec	This study
pE4878	mCherry-Myosin VIII-A CBD, Cerulean, PIP2A-Venus	*Asc*I digested pE4859 cloned into *Asc*I digested pE4783 to make pPZP-P$_{nos}$-Cerulean-P35S-PIP2A-Venus-Pi-mCherry-myosin VIII-A CBD	Spec	This study
pE4879	mCherry-Myosin VIII-B CBD, Cerulean, PIP2A-Venus	*Asc*I digested pE4859 cloned into *Asc*I digested pE4784 to make pPZP-P$_{nos}$-Cerulean-P35S-PIP2A-Venus-Pi-mCherry-myosin VIII-B CBD	Spec	This study
pE4880	mCherry-Myosin XI-K CBD, Cerulean, PIP2A-Venus	*Asc*I digested pE4859 cloned into *Asc*I digested pE4785 to make pPZP-P$_{nos}$-Cerulean-P35S-PIP2A-Venus-Pi-mCherry-myosin XI-K CBD	Spec	This study
pE4889	Myc-Myosin XI-K CBD, VirE2-Venus, mCherry-ABD2	PI-*Psp*I digested pE4780 cloned into PI-*Psp*I digested pE4292 to make pPZP-Pi-VirE2-Venus-P$_{nos}$-mCherry-ABD2-Pi-myc-myosin XI-K CBD	Spec	This study

pE4890	mCherry-Myosin VIII-2 CBD, Cerulean, PIP2A-Venus	*Asc*I digested pE4859 cloned into *Asc*I digested pE4782 to make pPZP-P$_{nos}$-Cerulean-P$_{35S}$-PIP2A-Venus-Pi-mCherry-myosin VIII-2 CBD	Spec	This study
pE4891	Myosin VIII-A, Myosin VIII-B, hptII	*Asc*I digested pE4858 cloned into *Asc*I digested pE4857 to make pPZP-hpt-XVE-P$_{35S}$-myosin VIII-B full length-myosin VIII-A full length	Spec	This study
pE4892	Myosin VIII-2, Myosin VIII-B, hptII	*Asc*I digested pE4875 cloned into *Asc*I digested pE4857 to make pPZP-hpt-XVE-P$_{35S}$-myosin VIII-B full length-myosin VIII-2 full length	Spec	This study
pE4908	Myosin VIII-B	*Age*I+*Swa*I digested pE4848 cloned into *Age*I+*Swa*I digested pE4754 to make pSAT6-Pi-myosin VIII-B FL	Amp	This study
pE4909	Myosin VIII-1	*Swa*I+*Bam*HI digested pE4847 cloned into *Swa*I+*Bam*HI digested pE4754 to make pSAT1-Pi-myosin VIII-1 FL	Amp	This study
pE4910	Myosin VIII-2	*Kpn*I+*Sma*I digested pE4875 cloned into *Kpn*I+*Sma*I digested pE4908 to make pSAT6-Pi-Myosin VIII-2 full length	Amp	This study
pE4911	Myosin VIII-A	*Kpn*I+*Bam*HI digested pE4858 cloned into *Kpn*I+*Bam*HI digested pE4909 to make pSAT1-Pi-Myosin VIII-A full length	Amp	This study
pE4912	Myosin VIII-1, *hptII*	*Asc*I digested pE4909 cloned into *Asc*I digested pE4672 to make pPZP-*hptII*-XVE-Pi-myosin VIII-1 full length	Spec	This study
pE4913	Myosin VIII-A, *hptII*	*Asc*I digested pE4911 cloned into *Asc*I digested pE4672 to make pPZP-*hptII*-XVE-Pi-myosin VIII-A full length	Spec	This study
pE4914	Myosin VIII-B, *hptII*	PI-*Psp*I digested pE4908 cloned into PI-*Psp*I digested pE4672 to make pPZP-*hptII*-XVE-Pi-myosin VIII-B full length	Spec	This study
pE4915	Myosin VIII-2, *hptII*	PI-*Psp*I digested pE4910 cloned into PI-*Psp*I digested pE4672 to make pPZP-*hptII*-XVE-Pi-myosin VIII-2 full length	Spec	This study

Amp, ampicillin; Gent, gentamicin; Kan, kanamycin; Rif, rifampicin; Spec, spectinomycin; hpt/hptII, hyromycin phosphotransferase II

P$_{35S}$, CaMV 35S promoter; P$_{nos}$, nopaline synthase promoter; Pi, inducible promoter
CBD, cargo binding domain

Supplemental Table 2. Bacterial strains used in this study.
Agrobacterium strains

Stock number	Bacterial host	Plasmid harbored	Important features	Antibiotic resistance	Reference
At1872	EHA105ΔVirE2	-----	Non-polar *virE2* deletion of pTiEHA105	Rif	Gelvin lab strain
At1879	At1872	pBISN2	pBISN2 *gusA-intron* gene	Kan, Rif, Spec	Gelvin lab strain
At2315	GV3101	pE4726	mCherry-Myosin VIII-1 CBD VirE2-Venus Cerulean-NLS	Rif, Spec	This study
At2316	GV3101	pE4727	mCherry-Myosin VIII-2 CBD VirE2-Venus Cerulean-NLS	Rif, Spec	This study
At2317	GV3101	pE4728	mCherry-Myosin VIII-A CBD VirE2-Venus Cerulean-NLS	Rif, Spec	This study
At2318	GV3101	pE4729	mCherry-Myosin VIII-B CBD VirE2-Venus Cerulean-NLS	Rif, Spec	This study
At2319	GV3101	pE4730	mCherry-Myosin XI-K CBD VirE2-Venus Cerulean-NLS	Rif, Spec	This study
At2320	GV3101	pE4758	nVenus-Myosin VIII-1 CBD VirE2-cYFP Cerulean-NLS mCherry-ABD2	Rif, Spec	This study
At2321	GV3101	pE4759	nVenus-Myosin VIII-2 CBD VirE2-cYFP Cerulean-NLS mCherryABD2	Rif, Spec	This study

At2322	GV3101	pE4755	nVenus-Myosin VIII-A CBD VirE2-cYFP Cerulean-NLS mCherry-ABD2	Rif, Spec	This study
At2323	GV3101	pE4756	nVenus-Myosin VIII-B CBD VirE2-cYFP Cerulean-NLS mCherry-ABD2	Rif, Spec	This study
At2324	GV3101	pE4757	nVenus-Myosin XI-K CBD VirE2-cYFP Cerulean-NLS mCherry-ABD2	Rif, Spec	This study
At2360	GV3101	pE4672	Binary vector; *hptII*	Rif, Spec	This study
At2361	GV3101	pE4854	Myosin VIII-1 *hptII*	Rif, Spec	This study
At2362	GV3101	pE4857	Myosin VIII-B *hptII*	Rif, Spec	This study
At2363	GV3101	pE4874	Myosin VIII-A *hptII*	Rif, Spec	This study
At2364	GV3101	pE4876	Myosin VIII-2 *hptII*	Rif, Spec	This study
At2365	GV3101	pE4852	myc-Myosin VIII-2 CBD VirE2-Venus mCherry-ABD2	Rif, Spec	This study
At2366	GV3101	pE4853	myc-Myosin VIII-A CBD VirE2-Venus mCherry-ABD2	Rif, Spec	This study
At2367	GV3101	pE4855	myc-Myosin VIII-1 CBD VirE2-Venus mCherry-ABD2	Rif, Spec	This study
At2368	GV3101	pE4856	myc-Myosin VIII-B CBD VirE2-Venus mCherry-ABD2	Rif, Spec	This study
At2369	GV3101	pE4889	myc-Myosin XI-K CBD VirE2-Venus mCherry-ABD2	Rif, Spec	This study
At2370	GV3101	pE4877	mCherry-Myosin VIII-1 CBD Cerulean PIP2A-Venus	Rif, Spec	This study

At2371	GV3101	pE4878	mCherry-Myosin VIII-A CBD Cerulean PIP2A-Venus	Rif, Spec	This study
At2372	GV3101	pE4879	mCherry-Myosin VIII-B CBD Cerulean PIP2A-Venus	Rif, Spec	This study
At2373	GV3101	pE4880	mCherry-Myosin XI-K CBD Cerulean PIP2A-Venus	Rif, Spec	This study
At2374	GV3101	pE4890	mCherry-Myosin VIII-2 CBD Cerulean PIP2A-Venus	Rif, Spec	This study
At2375	GV3101	pE4891	Myosin VIII-A Myosin VIII-B *hptII*	Rif, Spec	This study
At2376	GV3101	pE4892	Myosin VIII-2 Myosin VIII-B *hptII*	Rif, Spec	This study
At2389	GV3101	pE4912	Myosin VIII-1 *hptII*	Rif, Spec	This study
At2390	GV3101	pE4915	Myosin VIII-2 *hptII*	Rif, Spec	This study
At2391	GV3101	pE4913	Myosin VIII-A *hptII*	Rif, Spec	This study
At2392	GV3101	pE4914	Myosin VIII-B *hptII*	Rif, Spec	This study
At2403	At1872	pE3341	pBBR1-MCS2::mCherry	Kan	This study
At2404	At2403	pE4672	pPZP-RCS-P_{nos}-*hptII*	Kan, Spec	This study
At2405	At1872	pE4381	pPZP-P_{35S}-mCherry-intron-NLS	Spec	This study

At2423	GV3101	pE4786	Myc-Myosin VIII-1 CBD VirE2-Venus	Rif, Spec	This study
At2424	GV3101	pE4787	Myc-Myosin VIII-2 CBD VirE2-Venus	Rif, Spec	This study
At2425	GV3101	pE4788	Myc-Myosin VIII-A CBD VirE2-Venus	Rif, Spec	This study
At2426	GV3101	pE4789	Myc-Myosin VIII-B CBD VirE2-Venus	Rif, Spec	This study
At2427	GV3101	pE4790	Myc-Myosin XI-K CBD VirE2-Venus	Rif, Spec	This study
At2428	GV3101	pE4850	Myc-Myosin VIII-2 CBD Venus	Rif, Spec	This study

Kan, kanamycin; Rif, rifampicin; Spec, spectinomycin
CBD, cargo binding domain; FL, full-length

E. coli strains

Stock number	Host	Description of strain	Antibiotic resistance	Reference
E4823	BL21(DE3)	pE4814 in BL21DE3	Kan	This study
E4824	BL21(DE3)	pE4815 in BL21DE3	Kan	This study
E4825	BL21(DE3)	pE4816 in BL21DE3	Kan	This study
E4826	BL21(DE3)	pE4817 in BL21DE3	Kan	This study
E4827	BL21(DE3)	pE4818 in BL21DE3	Kan	This study
E4828	BL21(DE3)	pE4819 in BL21DE3	Kan	This study
E4829	BL21(DE3)	pE4820 in BL21DE3	Kan	This study
E4830	BL21(DE3)	pE4822 in BL21DE3	Kan	This study
Top10	N/A	F⁻ *mcr*A Δ(*mrr-hsd*RMS-*mcr*BC) φ80 *lac*ZΔM15Δ *lac*X74 *rec*A1 *ara*Δ139Δ(*ara-leu*)7697 *gal*U *gal*K *rps*L (StrR) *end*A1 *nup*G	Str	Invitrogen
BL21(DE3)	N/A	*E. coli* B F⁻ *dcm ompT hsdS* (*rB⁻mB⁻*) *gal* λ(DE3)	None	Studier and Moffatt, 1986

Kan, kanamycin; Str, streptomycin; N/A, not applicable

Supplemental Table 3. Primers used in this study.

Purpose	Forward primer	Reverse primer
myosin cDNA amplification		
VIII-1 CBD	5'-CCGGGGTACCAAAGCATCA GTACTTTCCGAGCT-3'	5'-CCGCGGATCCTCAATACC TGGTGCTATTTCTCC-3'
VIII-2 CBD	5'-CCGGGGTACCTCTATGT CTGATCTCCAGAAACG-3'	5'-CCGCGGATCCCTAGCC TCTTTTTCCCCACCA-3'
VIII-A CBD	5'-CCGGGGTACCTATCTTT CCGATCTTCAGCGT-3'	5'-CCGCGGATCCTTAATACC TAGTACTCCTCAACCTC-3'
VIII-B CBD	5'-CCGGGGTACCGTTCTTG CTGATCTCCAGAGCC-3'	5'-CCGCGGATCCTCAATAAC TTTTCTTGCACCACCAA-3'
XI-K CBD	5'-CCGGGGTACCGCAAGG AAAGCTATTGAAGAAGC-3'	5'-CCGCGGATCCTTACGAT GTACTGCCTTCTTTAC-3'
VIII-1 FL	5'-AATTGGTACCATGTCTC AGAAGGTTACTCCATT-3'	5'-TTAAGGATCCTCAATAC CTGGTGCTATTTCTCC-3'
VIII-2 FL	5'-AATTGGTACCATGATGT TATCGGCATCGCCGAAC-3'	5'-TTAACCCGGGCTAGCCT CTTTTTCCCCACCA-3'
VIII-A FL	5'-AATTGGTACCATGGCAC ACAAGGTTAAGGCATC-3'	5'-TTAAGGATCCTTAATAC CTAGTACTCCTCAACCTC-3'
VIII-B FL	5'-AATTGGTACCATGATGA AAAGTTCAGTGAAGG-3'	5'-TTAACCCGGGTCAATAA CTTTTCTTGCACCACC-3'

Purpose	Forward primer	Reverse primer
Plant genotyping		
Myosin VIII-1	5'-TTCGTGTGAACGTTGA TTCTG-3'	5'-TCCAGCTTGAATAGAT GACGG-3'
Myosin VIII-2	5'-CTGAGTTCAGGAGTGT TTCCG-3'	5'-TGTACCTCCGAACTGGA CAAG-3'
Myosin VIII-A	5'-AGGTTGTACAACACTG CTGGC-3'	5'-ACGAGAAATGGTCTTGT GCTG-3'
Myosin VIII-B	5'-AGTAAAGCAGGGCCA GTTTTG-3'	5'-CTACACTTTGCTTCAGC AGGG-3'
Myosin XI-1	5'-TCAAAACGTTGAACTA ACCGG-3'	5'- TTGTTTGGACGGGTATC TCAG-3'
Myosin XI-2	5'-GTGCTCCCGAAGATCC TATT-3'	5'-CAGTGCAACCACATGAA GATG-3'
Myosin XI-B	5'-CAGAAGCAACCCACTT CAGTC-3'	5'-GCTACCGAAGGAAGGAC TGTC-3'
Myosin XI-C	5'-TTATGATCACACCCGA GGAAG-3'	5'-AAAGACCCACAACAAAG GGAC-3'
Myosin XI-H	5'-AAATGCTTGAGCTGGA ACATG -3'	5'-TGCCCTGATTTTTCATT TGTC-3'
Myosin XI-J	5'-CTCACCTTGCAAAGTG GAGTC-3'	5'-CTCAACTTGCAATAAGG CCTG-3'
Myosin XI-E	5'-TTGGGATGACCCAACT TGTAC-3'	5'-GCCTGGAATACACTGAA GCTG-3'
Myosin XI-I	5'-TTCTGCAATTTCAATT CAGGC-3'	5'-CAGCAGACTTCTCCTTC ATGG-3'
Myosin XI-K	5'-GGGTAGCAAGATACTC CTCGG-3'	5'-GCAAGAGCAACTCAATT CTGG-3'
Salk T-DNA left border	5'-TGGTTCACGTAGTGGG CCATCG-3'	

CBD, cargo binding domain; FL, full-length

REFERENCES

Chien CT, Bartel PL, Sternglanz R, Fields F. (1991) The two-hybrid system: a method to identify and clone genes for proteins that interact with a protein of interest. Proc Natl Acad Sci USA **88:** 9578-9582

Citovsky V, Lee L-Y, Vyas S, Glick E, Chen M-H, Vainstein A, Gafni Y, Gelvin SB, Tzfira T. (2006) Subcellular localization of interacting proteins by bimolecular fluorescence complementation *in planta*. J Mol Biol **362**: 1120-1131

Koncz C, Schell J. (1986) The promoter of TL-DNA gene 5 controls the tissue-specific expression of chimaeric genes carried by a novel type of *Agrobacterium* binary vector. Mol Gen Genet **204**: 383-396

Kozaki A, Hake S, Colasanti J. (2004) The maize ID1 flowering time regulator is a zinc finger protein with novel DNA binding properties. Nucl Acids Res **32**: 1710-1720

Lee L-Y, Gelvin SB. (2014) Bimolecular fluorescence complementation for imaging protein-protein interactions in plant hosts of microbial pathogens. In Methods in Molecular Biology: Host-Bacteria Interactions. (A.C. Vergunst and D. O'Callaghan, eds.). Springer, Humana Press. Vol. 1197. pp. 185-208

Lu Q, Tang X, Tian G, Wang F, Liu K, Nguyen V, Kohalmi SE, Keller WA, Tsang EW, Harada JJ, Rothstein SJ, Cui Y. (2010) *Arabidopsis* homolog of the yeast TREX-2 mRNA export complex: Components and anchoring nucleoporin. Plant J **61:** 259-70

Studier FW, Moffatt BA. (1986) Use of bacteriophage T7 RNA polymerase to direct selective high-level expression of cloned genes. J Mol Biol **189**: 113-130

Supplemental Figure S1. Most *myosin VIII* and *myosin XI* single mutants are not deficient in stable transformation. Root segments from the indicated *Arabidopsis myosin VIII* or *myosin XI* single mutants, as well as control wild-type plants, were infected with the tumorigenic strain *A. tumefaciens* A208 at the concentration of 10^6 cfu/ml and the tumors were scored after 30 days of infection. A total of 10-15 plants and >100 segments per plant were tested for stable transformation. Values given are means ± SE. No significant differences were observed compared to wild-type plants.

Supplemental Figure S2. Growth and morphology of wild-type and higher order *myosin VIII* and *XI* mutant plants. (A) Roots; **(B)** Above-ground tissues.

Supplemental Figure S3. Effect on transformation of expressing full-length myosin cDNAs in wild-type and *myosin VIII-1/2/a/b* mutant plants. Root segments of five transgenic plants of each independent line, expressing the indicated *myosin VIII* cDNAs in the indicated genetic backgrounds, were infected with *A. tumefaciens* At849 containing the binary vector pBISN1 with a *gusA*-intron gene in the T-DNA region, at 10^7 cfu/ml. Six days after infection the root segments were stained with X-gluc and the percentage of roots showing GUS activity was calculated. (**A**) *myosin VIII-1* cDNA; (**B**) *myosin VIII-2* cDNA; (**C**) *myosin VIII-A* cDNA; (**D**) *myosin VIII-B* cDNA. Each bar indicates the mean of five plants and >100 root segments/plant ± SE for each transgenic line. Asterisks indicate significant differences of the indicated *myosin VIII* cDNA transgenic plant compared with plants harboring an empty vector (EV). L, Independent transgenic line.
[*t-test*, *$P < 0.05$; **$P < 0.01$].

Supplemental Figure S4. Effect on transformation of expressing full-length inducible myosin cDNAs in wild-type and *myosin VIII-1/2/a/b* mutant plants. Root segments of five transgenic plants of each independent line, expressing the indicated inducible *myosin VIII* cDNAs in the indicated genetic backgrounds, treated with β-estradiol or control solution for 24 hr, then infected with *A. tumefaciens* At849 containing the binary vector pBISN1 with a *gusA*-intron gene in the T-DNA region, at 10^7 cfu/ml. Six days after infection the root segments were stained with X-gluc and the percentage of roots showing GUS activity was calculated. (**A**) *myosin VIII-1* cDNA; (**B**) *myosin VIII-2* cDNA; (**C**) *myosin VIII-A* cDNA; (**D**) *myosin VIII-B* cDNA. Each bar indicates the mean of five plants and >100 root segments/plant ± SE for each transgenic line. Asterisks indicate significant differences of the indicated *myosin VIII* cDNA transgenic plant compared with plants treated with control solution. [*t-test*, *$P < 0.05$; **$P < 0.01$].

Supplemental Figure S5. Effect on transformation of expressing pairs of full-length myosin cDNAs in wild-type and *myosin VIII-1/2/a/b* mutant plants. Root segments of five transgenic plants of each independent line, expressing the indicated single and pairs of *myosin VIII* cDNAs in the indicated genetic backgrounds, were infected with *A. tumefaciens* At849 as mentioned above. The percentage of roots showing GUS activity was calculated. **(A)** *myosin VIII-2*, *myosin VIII-B*, and *myosin VIII-2+VIII-B* cDNAs in the *myosin VIII-1/2/a/b* mutant; **(B)** *myosin VIII-2*, *myosin VIII-B*, and *myosin VIII-2+VIII-B* cDNAs in Col-0; **(C)** *myosin VIII-A*, *myosin VIII-B*, and *myosin VIII-A+VIII-B* cDNAs in the *myosin VIII-1/2/a/b* mutant; **(D)** *myosin VIII-A*, *myosin VIII-B*, and *myosin VIII-A+VIII-B* cDNAs in Col-0. Each bar indicates the mean of five plants and >100 root segments/plant ± SE for each transgenic line. Except where otherwise indicated, asterisks indicate significant differences of the indicated *myosin VIII* cDNA transgenic plant compared with plants harboring an empty vector.
[*t-test*, *$P < 0.05$; **$P < 0.01$].

Supplemental Figure S6. Inducible expression of myosin VIII, but not myosin XI-K, CBDs inhibits transformation. Root segments of transgenic plants of each independent line, expressing the indicated inducible *myosin VIII* or *myosin XI-K* CBD cDNA, were treated with β-estradiol or control solution for 24 hr, then infected with *A. tumefaciens* At849 containing the binary vector pBISN1 with a *gusA*-intron gene in the T-DNA region, at the indicated 10^6, 10^7 and 10^8 cfu/ml. Six days after infection the root segments were stained with X-gluc and the percentage of roots showing GUS activity was calculated. **(A)** myosin VIII-1 CBD; **(B)** myosin VIII-2 CBD; **(C)** myosin VIII-A CBD; **(D)** myosin VIII-B CBD; **(E)** myosin XI-K CBD. Each bar indicates the mean of five plants and >100 root segments/plant ± SE for each transgenic line. Asterisks indicate significant differences of the indicated *myosin VIII* cDNA transgenic plant compared with plants treated with control solution. [*t-test*, *$P < 0.05$; **$P < 0.01$].

A

B

Supplemental Figure S7. Lack of co-localization of VirE2-Venus and FM4-64. **(A)** Inducible VirE2-Venus transgenic plants were treated with β-estradiol for 8 hr, then the root cells were stained with FM4-64 for 5 minutes before imaging. The Z-stack image was taken using a Zesis LSM 880 Upright Confocal Plan-Apo microscope with a 20×/0.8 objective. **(B)** Pearson's and overlap correlations of Venus and red FM4-64 fluorescence. A total of 10-15 transgenic lines and >50 cells were used for quantitative analysis using Image J. Values <0.5 indicate lack of co-localization of the two proteins.

Supplemental Figure S8. Subcellular localization of VirE2-Venus in transgenic plants. (A) Inducible VirE2-Venus expressed in wild-type plants. **(B)** Inducible VirE2-Venus expressed in *myosin VIII-1/2/a/b* quadruple mutant plants. Roots from three independent transgenic plants were treated with β-estradiol for 8 hr, and the images were taken using a Zesis LSM 880 Upright Confocal Plan-Apo microscope with a 20×/0.8 objective.

| PIP2A-Venus | mCherry-myosin VIII CBD | Merge |

A — Myosin VIII-1 CBD

B — Myosin VIII-2 CBD

C — Myosin VIII-A CBD

Supplemental Figure S9. Myosin VIII CBDs colocalize with the plasma membrane marker PIP2A. (A) Transgenic *Arabidopsis* plant roots that inducibly express an mCherry-myosin VIII-1 CBD and constitutively express PIP2A-Venus. **(B)** Transgenic *Arabidopsis* plants roots that inducibly express an mCherry-myosin VIII-2 CBD and constitutively express PIP2A-Venus. **(C)** Transgenic *Arabidopsis* plants roots that inducibly express an mCherry-myosin VIII-A CBD and constitutively express PIP2A-Venus. A total of 10 independent transgenic lines were treated with β-estradiol for 8 hr, and representative images were taken using a Zesis LSM 880 Upright Confocal Plan-Apo microscope with a 20×/0.8 objective.

Supplemental Figure S10. VirE2 re-localizes to the perinuclear area after infection by an *Agrobacterium* strain that can transfer T-DNA. VirE2-Venus transgenic plants were treated with 5 µM β-estradiol for 24 hr. Root segments were inoculated with 10^8 cfu/ml of the *virE2* mutant strain *A. tumefaciens* At1872 **(A)** lacking or **(B)** containing a T-DNA region. Images of the elongation zone of the root were taken 8 hr after infection using a Zesis LSM 880 Upright Confocal microscope with a Plan-Apo 20×/0.8 objective. Statistical analysis of VirE2 localization is shown in Figure 11A. Red arrows indicate several examples of perinuclear VirE2-Venus. Bars indicate 50 µm.

Supplemental Figure S11. Infection of root cells with an *Agrobacterium* strain capable of transferring T-DNA re-localizes VirE2-Venus from the cellular periphery into the cytoplasm. Transgenic plants inducibly expressing VirE2-Venus were treated with 5 μM β-estradiol for 24 hr, then inoculated with 10^8 cfu/ml of the *virE2* mutant strain *A. tumefaciens* At1872 without (left panel) or with (right panel) T-DNA. The *Agrobacterium* cells also expressed an mCherry protein. Z-stack images of cells in the root elongation zone were taken after 8 hr of infection using a Zesis LSM 880 Upright Confocal Plan-Apo microscope using a 20×/0.8 objective.

Supplemental Figure S12. Infection of root cells with an *Agrobacterium* strain capable of transferring T-DNA does not re-localize VirE2-Venus from the cellular periphery into the cytoplasm when the myosin VIII-1 CBD is expressed. Transgenic plants inducibly expressing the myosin VIII-1 CBD and VirE2-Venus were treated with 5 μM β-estradiol for 24 hr, then inoculated with 10^8 cfu/ml of the *virE2* mutant strain *A. tumefaciens* At1872 without (left panel) or with (right panel) T-DNA. Z-stack images of cells in the root elongation zone were taken after 8 hr of infection using a Zesis LSM 880 Upright Confocal Plan-Apo microscope using a 20×/0.8 objective.

Supplemental Figure S13. Infection of root cells expressing the myosin VIII-2 CBD with an *Agrobacterium* strain capable of transferring T-DNA relocalizes VirE2-Venus from the cellular periphery into the cytoplasm. Transgenic plants inducibly expressing the myosin VIII-2 CBD and VirE2-Venus were treated with 5 μM β-estradiol for 24 hr, then inoculated with 10^8 cfu/ml of the *virE2* mutant strain *A. tumefaciens* At1872 without (left panel) or with (right panel) T-DNA. Z-stack images of cells in the root elongation zone were taken after 8 hr of infection using a Zesis LSM 880 Upright Confocal Plan-Apo microscope using a 20×/0.8 objective.

Supplemental Figure S14. Infection of root cells expressing the myosin VIII-A CBD with an *Agrobacterium* strain capable of transferring T-DNA relocalizes VirE2-Venus from the cellular periphery into the cytoplasm. Transgenic plants inducibly expressing the myosin VIII-A CBD and VirE2-Venus were treated with 5 μM β-estradiol for 24 hr, then inoculated with 10^8 cfu/ml of the *virE2* mutant strain *A. tumefaciens* At1872 without (left panel) or with (right panel) T-DNA. Z-stack images of cells in the root elongation zone were taken after 8 hr of infection using a Zesis LSM 880 Upright Confocal Plan-Apo microscope using a 20×/0.8 objective.

Supplemental Figure S15. Infection of root cells expressing the myosin VIII-B CBD with an *Agrobacterium* strain capable of transferring T-DNA relocalizes VirE2-Venus from the cellular periphery into the cytoplasm. Transgenic plants inducibly expressing the myosin VIII-B CBD and VirE2-Venus were treated with 5 μM β-estradiol for 24 hr, then inoculated with 10^8 cfu/ml of the *virE2* mutant strain *A. tumefaciens* At1872 without (left panel) or with (right panel) T-DNA. Z-stack images of cells in the root elongation zone were taken after 8 hr of infection using a Zesis LSM 880 Upright Confocal Plan-Apo microscope using a 20×/0.8 objective.

Supplemental Figure S16. Expression of the myosin XI-K CBD does not affect VirE2-Venus localization upon infection by *Agrobacterium*. Transgenic plants inducibly expressing the myosin XI-K CBD and VirE2-Venus were treated with 5 µM β-estradiol for 24 hr, then inoculated with 10^8 cfu/ml of the *virE2* mutant strain *A. tumefaciens* At1872 without (left panel) or with (right panel) T-DNA. Z-stack images of cells in the root elongation zone were taken after 8 hr of infection using a Zesis LSM 880 Upright Confocal Plan-Apo microscope using a 20×/0.8 objective.

Supplemental Figure 17. Expression of the myosin XI-K CBD inhibits movement of VirE2-Venus. Transgenic plants inducibly expressing Myc-myosin XI-K CBD and VirE2-Venus were treated with 5 μM β-estradiol for 24 hr. Root segments were inoculated with 10^8 cfu/ml of the *virE2* mutant strain *A. tumefaciens* At2405 harboring a T-DNA which constitutively expressed mCherry-NLS (to mark nucleus of transformed cells) (right panel) or without T-DNA (left panel). Time lapse images were taken after 24 hr of infection using a Zesis LSM 880 Upright Confocal Plan-Apo microscope with a 20×/0.8 objective. White arrows indicate nuclei, yellow arrow indicates a non-transformed cell. Bars indicate 50 μm.

Parsed Citations

Amari K, Di Donato M, Dolja VV, Heinlein M. (2014) Myosins VIII and XI play distinct roles in reproduction and transport of tobacco mosaic virus. PLoS Pathog 10: e1004448
  Google Scholar: Author Only Title Only Author and Title

Avisar D, Prokhnevsky AI, Dolja VV. (2008a) Class VIII myosins are required for plasmodesmatal localization of a closterovirus Hsp70 homolog. J Virol 82: 2836-2843
  Google Scholar: Author Only Title Only Author and Title

Avisar D, Prokhnevsky AI, Makarova KS, Koonin EV, Dolja VV. (2008b) Myosin XI-K is required for rapid trafficking of Golgi stacks, peroxisomes, and mitochondria in leaf cells of Nicotiana benthamiana. Plant Physiol 146: 1098-1108
  Google Scholar: Author Only Title Only Author and Title

Bar-Sinai S, Belausov E, Dwivedi V, Sadot,E. (2022) Collisions of cortical microtubules with membrane associated myosin VIII tail. Cells 11:145
  Google Scholar: Author Only Title Only Author and Title

Bezanilla M, Horton AC, Sevener HC, Quatrano RS. (2003) Phylogenetic analysis of new plant myosin sequences. J Mol Evol 57: 229-239
  Google Scholar: Author Only Title Only Author and Title

Bhattacharjee S, Lee L-Y, Oltmanns H, Cao H, Veena, Cuperus J, Gelvin SB. (2008) IMPa-4, an Arabidopsis importin alpha isoform, is preferentially involved in Agrobacterium-mediated plant transformation. Plant Cell 20: 2661-2680
  Google Scholar: Author Only Title Only Author and Title

Bichet A, Desnos T, Turner S, Grandjean O, Hofte H. (2001) BOTERO1 is required for normal orientation of cortical microtubules and anisotropic cell expansion in Arabidopsis. Plant J 25: 137-148
  Google Scholar: Author Only Title Only Author and Title

Bundock P, den Dulk-Ras A, Beijersbergen A, Hooykaas PJ. (1995) Trans-kingdom T-DNA transfer from Agrobacterium tumefaciens to Saccharomyces cerevisiae. EMBO J 14: 3206-3214
  Google Scholar: Author Only Title Only Author and Title

Cascales E, Christie,PJ. (2004) Definition of a bacterial type IV secretion pathway for a DNA substrate. Science 304: 1170-1173
  Google Scholar: Author Only Title Only Author and Title

Chang CW, Williams SJ, Counago RM, Kobe B. (2014) Structural basis of interaction of bipartite nuclear localization signal from Agrobacterium VirD2 with rice importin-alpha. Mol Plant 7: 1061-1064
  Google Scholar: Author Only Title Only Author and Title

Chilton M-D, Drummond MH, Merio DJ, Sciaky D, Montoya AL, Gordon MP, Nester EW. (1977) Stable incorporation of plasmid DNA into higher plant cells: The molecular basis of crown gall tumorigenesis. Cell 11: 263-271
  Google Scholar: Author Only Title Only Author and Title

Christie PJ, Atmakuri K, Krishnamoorthy V, Jakubowski S, Cascales E. (2005) Biogenesis, architecture, and function of bacterial type IV secretion systems. Annu Rev Microbiol 59: 451-485
  Google Scholar: Author Only Title Only Author and Title

Citovsky V, Liu B. (2017) Myosin-driven transport network in plants is functionally robust and distinctive. Proc Natl Acad Sci USA 114: 1756-1758
  Google Scholar: Author Only Title Only Author and Title

Citovsky V, Wong ML, Zambryski P. (1989) Cooperative interaction of Agrobacterium VirE2 protein with single-stranded DNA: Implications for the T-DNA transfer process. Proc Natl Acad Sci USA 86: 1193-1197
  Google Scholar: Author Only Title Only Author and Title

Citovsky V, Zupan J, Warnick D, Zambryski P. (1992) Nuclear localization of Agrobacterium VirE2 protein in plant cells. Science 256: 1802-1805
  Google Scholar: Author Only Title Only Author and Title

Clough SJ, Bent AF. (1998) Floral dip: A simplified method for Agrobacterium-mediated

transformation of Arabidopsis thaliana. Plant J 16: 735-743

de Groot MJ, Bundock P, Hooykaas PJ, Beijersbergen AG. (1998) Agrobacterium tumefaciens-mediated transformation of filamentous fungi. Nat Biotechnol 16: 839-842
  Google Scholar: Author Only Title Only Author and Title

Djamei A, Pitzschke A, Nakagami H, Rajh I, Hirt H. (2007) Trojan horse strategy in Agrobacterium transformation: abusing MAPK

defense signaling. Science 318: 453-456
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Duan Z, Tominaga M. (2018) Actin-myosin XI: An intracellular control network in plants. Biochem Biophys Res Commun 506: 403-408
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Duckely M, Oomen C, Axthelm F, Van Gelder P, Waksman G, Engel A. (2005) The VirE1VirE2 complex of Agrobacterium tumefaciens interacts with single-stranded DNA and forms channels. Mol Microbiol 58: 1130-1142
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Dumas F, Duckely M, Pelczar P, Van Gelder P, Hohn B. (2001) An Agrobacterium VirE2 channel for transferred-DNA transport into plant cells. Proc Natl Acad Sci USA 98: 485-490
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Dym O, Albeck S, Unger T, Jacobovitch J, Branzburg A, Michael Y, Frenkiel-Krispin D, Wolf S., Elbaum M. (2008) Crystal structure of the Agrobacterium virulence complex VirE1-VirE2 reveals a flexible protein that can accommodate different partners. Proc Natl Acad Sci USA 105: 11170-11175
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Embley TM, Martin W. (2006) Eukaryotic evolution, changes and challenges. Nature 440: 623-630
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Engstrom P, Zambryski P, Van Montagu M, Stachel S. (1987) Characterization of Agrobacterium tumefaciens virulence proteins induced by the plant factor acetosyringone. J Mol Biol 197: 635-645
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Foth BJ, Goedecke MC, Soldati D. (2006) New insights into myosin evolution and classification. Proc Natl Acad Sci USA 103: 3681-3686
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Geeves MA. (2016) Review: The ATPase mechanism of myosin and actomyosin. Biopolymers 105: 483-491
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Gelvin SB. (1998) Agrobacterium VirE2 proteins can form a complex with T strands in the plant cytoplasm. J Bacteriol 180: 4300-4302
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Gelvin SB. (2000) Agrobacterium and plant genes involved in T-DNA transfer and integration. Annu Rev Plant Physiol Plant Mol Biol 51: 223-256
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Gelvin SB. (2003) Agrobacterium and plant transformation: The biology behind the "gene-jockeying" tool. Microbiol Mol Biol Rev 67: 16-37
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Gelvin, SB. (2006) Agrobacterium transformation of Arabidopsis thaliana roots: A quantitative assay. Methods Mol Biol 343: 105-113
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Gelvin SB. (2010) Finding a way to the nucleus. Curr Opin Microbiol 13: 53-58
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Gelvin SB. (2017) Integration of Agrobacterium T-DNA into the plant genome. Annual review of genetics 51: 22
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Golomb L, Abu-Abied M, Belausov E, Sadot E. (2008) Different subcellular localizations and functions of Arabidopsis myosin VIII. BMC Plant Biol 8: 3
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Grange W, Duckely M, Husale S, Jacob S, Engel A, Hegner M. (2008) VirE2: A unique ssDNA-compacting molecular machine. PLoS Biol 6: e44
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Haraguchi T, Tominaga M, Matsumoto R, Sato K, Nakano A, Yamamoto K, Ito K. (2014) Molecular characterization and subcellular localization of Arabidopsis class VIII myosin, ATM1. J Biol Chem 289: 12343-12355
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Hodges L, Cuperus J, Ream W. (2004) Agrobacterium rhizogenes GALLS protein

substitutes for Agobacterium tumefaciens single-stranded DNA-binding protein VirE2. J Bacteriol 186: 3065-3077

Hodges LD, Vergunst AC, Neal-McKinney J, den Dulk-Ras A, Moyer DM,

Hooykaas PJJ, Ream W. (2006) Agrobacterium rhizogenes GALLS protein contains domains for ATP binding, nuclear localization, and type IV secretion. J Bacteriol 188: 8222-8230
  Google Scholar: Author Only Title Only Author and Title

Howard E, Citovsky V. (1990) The emerging structure of the Agrobacterium T-DNA transfer complex. Bioessays 12: 103-108
  Google Scholar: Author Only Title Only Author and Title

Huang S, An YQ, McDowell JM, McKinney EC, Meagher RB. (1996) The Arabidopsis thaliana ACT4/ACT12 actin gene subclass is strongly expressed throughout pollen development. Plant J 10: 189-202
  Google Scholar: Author Only Title Only Author and Title

Hussey,PJ, Ketelaar T, Deeks MJ. (2006) Control of the actin cytoskeleton in plant cell growth. Annu Rev Plant Biol 57: 109-125
  Google Scholar: Author Only Title Only Author and Title

Hwang H-H, Gelvin SB. (2004) Plant proteins that interact with VirB2, the Agrobacterium tumefaciens pilin protein, mediate plant transformation. Plant Cell 16: 3148-3167
  Google Scholar: Author Only Title Only Author and Title

Koncz C, Schell J. (1986) The promoter of TL-DNA gene 5 controls the tissue-specific

expression of chimaeric genes carried by a novel type of Agrobacterium binary vector. Mol Gen Genet 204: 383-396

Krendel M, Osterweil EK, Mooseker MS. (2007) Myosin 1E interacts with synaptojanin-1 and

dynamin and is involved in endocytosis. FEBS Lett 581: 644-650

Lacroix B, Citovsky V. (2013) The roles of bacterial and host plant factors in Agrobacterium-mediated genetic transformation. Int J Dev Biol 57: 467-481
  Google Scholar: Author Only Title Only Author and Title

Lacroix B, Vaidya M, Tzfira T, Citovsky V. (2005) The VirE3 protein of Agrobacterium mimics a host cell function required for plant genetic transformation. EMBO J 24: 428-437
  Google Scholar: Author Only Title Only Author and Title

Lapham RA, Lee L-Y, Xhako E, Gomez EG, Nivya VM., and Gelvin, S.B. (2021). Agrobacterium VirE2 protein modulates plant gene expression and mediates transformation from its location outside the nucleus. Front Plant Sci 12, 684192
  Google Scholar: Author Only Title Only Author and Title

Lee L-Y, Wu F-H, Hsu C-T, She, S-C, Yeh,HY, Liao D-C, Fang M-J, Liu N-T, Yen Y-C, Dokládal L, Sýkorová E, Gelvin SB, Lin C-S (2012) Screening a cDNA library for protein-protein interactions directly in planta. Plant Cell. 24: 1746-1759
  Google Scholar: Author Only Title Only Author and Title

Lee YR, Liu B. (2004) Cytoskeletal motors in Arabidopsis. Sixty-one kinesins and seventeen myosins. Plant Physiol 136: 3877-3883
  Google Scholar: Author Only Title Only Author and Title

Li X, Pan SQ. (2017) Agrobacterium delivers VirE2 protein into host cells via clathrin-mediated endocytosis. Sci Adv 3: e1601528
  Google Scholar: Author Only Title Only Author and Title

Li X, Tu H, Pan SQ. (2018) Agrobacterium delivers anchorage protein VirE3 for companion VirE2 to aggregate at host entry sites for T-DNA protection. Cell Rep 25: 302-311 e306
  Google Scholar: Author Only Title Only Author and Title

Li X, Yang Q, Peng L, Tu H, Lee L-Y, Gelvin SB, Pan SQ. (2020) Agrobacterium-delivered VirE2 interacts with host nucleoporin CG1 to facilitate the nuclear import of VirE2-coated T complex. Proc Natl Acad Sci USA 117: 26389-26397
  Google Scholar: Author Only Title Only Author and Title

Li X, Zhu T, Tu H, Pan SQ. (2020) Agrobacterium VirE3 uses its two tandem

domains at the C-terminus to retain its companion VirE2 on the cytoplasmic side of the host plasma membrane. Front Plant Sci 11:464

Madison SL, Nebenfuhr A. (2013) Understanding myosin functions in plants: Are we there yet? Curr Opin Plant Biol 16: 710-717
  Google Scholar: Author Only Title Only Author and Title

Madison SL, Buchanan ML, Glass JD, McClain TF, Park E, Nebenfuhr A. (2015) Class XI myosins move specific organelles in pollen tubes and are required for normal fertility and pollen tube growth in Arabidopsis. Plant Physiol 169: 1946-1960
  Google Scholar: Author Only Title Only Author and Title

McCullen CA, Binns AN. (2006) Agrobacterium tumefaciens and plant cell interactions and activities required for interkingdom

macromolecular transfer. Annu Rev Cell Dev Biol 22: 101-127
Google Scholar: Author Only Title Only Author and Title

Meagher RB, McKinney EC, Kandasamy MK. (1999) Isovariant dynamics expand and buffer the responses of complex systems: The diverse plant actin gene family. Plant Cell 11: 995-1006
Google Scholar: Author Only Title Only Author and Title

Narasimhulu SB, Deng XB, Sarria R, Gelvin SB. (1996) Early transcription of Agrobacterium T-DNA genes in tobacco and maize. Plant Cell 8: 873-886
Google Scholar: Author Only Title Only Author and Title

Ojangu EL, Jarve K, Paves H, Truve E. (2007) Arabidopsis thaliana myosin XIK is involved in root hair as well as trichome morphogenesis on stems and leaves. Protoplasma 230: 193-202
Google Scholar: Author Only Title Only Author and Title

Otten L, DeGreve H, Leemans J, Hain R, Hooykaas P, Schell J. (1984) Restoration of

virulence of vir region mutants of Agrobacterium tumefaciens strain B6S3 by coinfection with normal and mutant Agrobacterium strains. Mol Gen Genet 195: 159-163

Peremyslov VV, Prokhnevsky AI, Dolja VV. (2010) Class XI myosins are required for development, cell expansion, and F-Actin organization in Arabidopsis. Plant Cell 22: 1883-1897
Google Scholar: Author Only Title Only Author and Title

Peremyslov VV, Prokhnevsky AI, Avisar D, Dolja VV. (2008) Two class XI myosins function in organelle trafficking and root hair development in Arabidopsis. Plant Physiol 146: 1109-1116
Google Scholar: Author Only Title Only Author and Title

Peremyslov,VV, Mockler TC, Filichkin S., Fox SE, Jaiswal P, Makarova K., Koonin .V, Dolja,VV. (2011) Expression, splicing, and evolution of the myosin gene family in plants. Plant Physiol 155: 1191-1204
Google Scholar: Author Only Title Only Author and Title

Piers KL, Heath JD, Liang X, Stephens KM, Nester EW. (1996) Agrobacterium tumefaciens-mediated transformation of yeast. Proc Natl Acad Sci USA 93: 1613-1618
Google Scholar: Author Only Title Only Author and Title

Prokhnevsky AI, Peremyslov,VV, Dolja VV. (2008) Overlapping functions of the four class XI myosins in Arabidopsis growth, root hair elongation, and organelle motility. Proc Natl Acad Sci USA 105: 19744-19749
Google Scholar: Author Only Title Only Author and Title

Radford JE, White RG. (1998) Localization of a myosin-like protein to plasmodesmata. Plant J 14: 743-750
Google Scholar: Author Only Title Only Author and Title

Ream W. (2009) Agrobacterium tumefaciens and A. rhizogenes use different proteins to transport bacterial DNA into the plant cell nucleus. Micro Biotechnol 2: 416-42.
Google Scholar: Author Only Title Only Author and Title

Reichelt S, Knight AE, Hodge TP, Baluska F, Samaj J, Volkmann D, Kendrick-Jones J. (1999) Characterization of the unconventional myosin VIII in plant cells and its localization at the post-cytokinetic cell wall. Plant J 19: 555-567
Google Scholar: Author Only Title Only Author and Title

Reisen D, Hanson MR. (2007) Association of six YFP-myosin XI-tail fusions with mobile plant cell organelles. BMC Plant Biol 7: 6
Google Scholar: Author Only Title Only Author and Title

Richards TA, Cavalier-Smith T. (2005) Myosin domain evolution and the primary divergence of eukaryotes. Nature 436: 1113-1118
Google Scholar: Author Only Title Only Author and Title

Rossi L, Hohn B, Tinland B. (1996) Integration of complete transferred DNA units is dependent on the activity of virulence E2 protein of Agrobacterium tumefaciens. Proc Natl Acad Sci USA 93: 126-130
Google Scholar: Author Only Title Only Author and Title

Roushan MR, deZeeuw MAM, Hooykaas PJJ, van Heusden GPH. (2018) Application

of phiLOV2.1 as a fluorescent marker for visualization of Agrobacterium effector protein translocation. Plant J 96: 685-699

Sakalis,PA, van Heusden GP, Hooykaas PJ. (2014) Visualization of VirE2 protein translocation by the Agrobacterium type IV secretion system into host cells. Microbiologyopen 3: 104-117
Google Scholar: Author Only Title Only Author and Title

Salman H, Abu-Arish A, Oliel S, Loyter A, Klafter J, Granek R, Elbaum M. (2005) Nuclear localization signal peptides induce molecular delivery along microtubules. Biophys J 89: 2134-2145

Google Scholar: Author Only Title Only Author and Title

Sato T, Hanada M, Bodrug S, Irie, S, Iwama N, Boise LH, Thompson CB, Golemis E, Fong L, Wang HG. (1994) Interactions among members of the Bcl-2 protein family analyzed with a yeast two-hybrid system. Proc Natl Acad Sci U S A 91: 9238-9242
  Google Scholar: Author Only Title Only Author and Title

Sattarzadeh A, Franzen R, Schmelzer E. (2008) The Arabidopsis class VIII myosin ATM2 is involved in endocytosis. Cell Motil Cytoskeleton 65: 457-468
  Google Scholar: Author Only Title Only Author and Title

Schrammeijer B, den Dulk-Ras A, Vergunst AC, Jurado Jacome E, Hooykaas PJ. (2003) Analysis of Vir protein translocation from Agrobacterium tumefaciens using Saccharomyces cerevisiae as a model: Evidence for transport of a novel effector protein VirE3. Nucleic Acids Res 31: 860-868
  Google Scholar: Author Only Title Only Author and Title

Sen P, Pazour GJ, Anderson D, Das A. (1989) Cooperative binding of Agrobacterium tumefaciens VirE2 protein to single-stranded DNA. J Bacteriol 171: 2573-2580
  Google Scholar: Author Only Title Only Author and Title

Shi Y, Lee L-Y, Gelvin SB. (2014) Is VIP1 important for Agrobacterium-mediated transformation? Plant J 79: 848-860
  Google Scholar: Author Only Title Only Author and Title

Simone M, McCullen CA, Stahl LE, Binns AN. (2001) The carboxy-terminus of VirE2 from Agrobacterium tumefaciens is required for its transport to host cells by the virB-encoded type IV transport system. Mol Microbiol 41: 1283-1293
  Google Scholar: Author Only Title Only Author and Title

Smith LG, Oppenheimer .G. (2005) Spatial control of cell expansion by the plant cytoskeleton. Annu Rev Cell Dev Biol 21: 271-295
  Google Scholar: Author Only Title Only Author and Title

Sparkes I, Runions J, Hawes C, Griffing L. (2009) Movement and remodeling of the endoplasmic reticulum in nondividing cells of tobacco leaves. Plant Cell 21: 3937-3949
  Google Scholar: Author Only Title Only Author and Title

Stachel SE, Nester EW. (1986) The genetic and transcriptional organization of the vir region of the A6 Ti plasmid of Agrobacterium tumefaciens. EMBO J 5: 1445-1454
  Google Scholar: Author Only Title Only Author and Title

Staiger CJ, Sheahan MB, Khurana P, Wang X, McCurdy DW, Blanchoin L. (2009) Actin filament dynamics are dominated by rapid growth and severing activity in the Arabidopsis cortical array. J Cell Biol 184: 269-280
  Google Scholar: Author Only Title Only Author and Title

Szymanski DB, Cosgrove DJ. (2009) Dynamic coordination of cytoskeletal and cell wall systems during plant cell morphogenesis. Curr Biol 19: R800-811
  Google Scholar: Author Only Title Only Author and Title

Tenea GN, Spantzel J, Lee L-Y, Zhu Y, Lin K, Johnson SJ, Gelvin SB. (2009) Over-expression of several Arabidopsis histone genes increases Agrobacterium-mediated transformation and transgene expression in plant cells. Plant Cell 21: 3350–3367
  Google Scholar: Author Only Title Only Author and Title

Trybus KM. (2008) Myosin V from head to tail. Cell Mol Life Sci 65: 1378-1389
  Google Scholar: Author Only Title Only Author and Title

Tu H, Li X, Yang Q, Peng L, Pan SQ. (2018) Real-time trafficking of Agrobacterium virulence protein VirE2 inside host cells. Curr Top Microbiol Immunol 418: 261-286
  Google Scholar: Author Only Title Only Author and Title

Tzfira T, Vaidya M, Citovsky V. (2001) VIP1, an Arabidopsis protein that interacts with Agrobacterium VirE2, is involved in VirE2 nuclear import and Agrobacterium infectivity. EMBO J 20: 3596-3607
  Google Scholar: Author Only Title Only Author and Title

Tzfira T, Li J, Lacroix B, Citovsky V. (2004) Agrobacterium T-DNA integration: Molecules and models. Trends Genet 20: 375-383
  Google Scholar: Author Only Title Only Author and Title

Ueda H, Tamura K, Hara-Nishimura I. (2015) Functions of plant-specific myosin XI: from intracellular motility to plant postures. Curr Opin Plant Biol 28: 30-38
  Google Scholar: Author Only Title Only Author and Title

Ueda H, Yokota E, Kutsuna N, Shimada T, Tamura K, Shimmen T, Hasezawa S, Dolja VV., Hara-Nishimura I. (2010) Myosin-dependent endoplasmic reticulum motility and F-actin organization in plant cells. Proc Natl Acad Sci USA 107: 6894-6899
  Google Scholar: Author Only Title Only Author and Title

Ueki S, Citovsky V. (2005) Identification of an interactor of cadmium ion-induced glycine-rich protein involved in regulation of callose levels in plant vasculature. Proc Natl Acad Sci USA 102: 12089-12094

Google Scholar: Author Only Title Only Author and Title

Vale RD. (2003) The molecular motor toolbox for intracellular transport. Cell 112: 467-480

Google Scholar: Author Only Title Only Author and Title

Vergunst AC, van Lier MC, den Dulk-Ras A, Stuve TA, Ouwehand A, Hooykaas PJ. (2005) Positive charge is an important feature of the C-terminal transport signal of the VirB/D4-translocated proteins of Agrobacterium. Proc Natl Acad Sci USA 102: 832-837

Google Scholar: Author Only Title Only Author and Title

Wang H, Han S, Siao W, Song C, Xiang Y, Wu X, Cheng P, Li H, Jasik J, Micieta K, Turna J, Voigt B, Baluska F, Liu J, Wang Y, Zhao H. (2015) Arabidopsis synaptotagmin 2 participates in pollen germination and tube growth and is delivered to plasma membrane via conventional secretion. Mol Plant 8: 1737-1750

Google Scholar: Author Only Title Only Author and Title

Wang P, Hawkins TJ, Richardson C, Cummins I, Deeks MJ, Sparkes I, Hawes C, Hussey PJ. (2014) The plant cytoskeleton, NET3C, and VAP27 mediate the link between the plasma membrane and endoplasmic reticulum. Curr Biol 24: 1397-1405

Google Scholar: Author Only Title Only Author and Title

Wojtaszek P, Anielska-Mazur A, Gabrys H, Baluska F, Volkmann D. (2005) Recruitment of myosin VIII towards plastid surfaces is root-cap specific and provides the evidence for actomyosin involvement in root osmosensing. Funct Plant Biol 32: 721-736

Google Scholar: Author Only Title Only Author and Title

Wu Z, Bezanilla M. (2014) Myosin VIII associates with microtubule ends and together with actin plays a role in guiding plant cell division. Elife 3:e03498

Google Scholar: Author Only Title Only Author and Title

Yang L, Qin L, Liu G, Peremyslov VV, Dolja VV, Wei Y. (2014) Myosins XI modulate host cellular responses and penetration resistance to fungal pathogens. Proc Natl Acad Sci USA 111: 13996-14001

Google Scholar: Author Only Title Only Author and Title

Yang Q, Li X, Tu H, Pan SQ. (2017) Agrobacterium-delivered virulence protein VirE2 is trafficked inside host cells via a myosin XI-K-powered ER/actin network. Proc Natl Acad Sci USA 114: 2982-2987

Google Scholar: Author Only Title Only Author and Title

Yusibov VM, Steck TR, Gupta V, Gelvin, SB. (1994) Association of single-stranded transferred DNA from Agrobacterium tumefaciens with tobacco cells. Proc Natl Acad Sci USA 91: 2994-2998

Google Scholar: Author Only Title Only Author and Title

Zuo J, Niu Q-W, Chua N-H. (2000) An estrogen receptor-based transactivator XVE mediates

highy inducible gene expression in transgenic plants. Plant J 24: 265-273

Zupan JR, Citovsky V, Zambryski P. (1996) Agrobacterium VirE2 protein mediates nuclear uptake of single-stranded DNA in plant cells. Proc Natl Acad Sci USA 93: 2392-2397

Google Scholar: Author Only Title Only Author and Title