DNAJB8 oligomerization is mediated by an aromatic-rich motif that is dispensable for substrate activity

Bryan D. Ryder[1,*], David R. Boyer[2], Elizaveta Ustyantseva[3], Ayde Mendoza-Oliva[1], Mikołaj I. Kuska[1], Paweł M. Wydorski[1], Michael Sawaya[2], Marc I. Diamond[1], David S. Eisenberg[2], Harm H. Kampinga[3], Lukasz A. Joachimiak[1,4,#]

[1]Center for Alzheimer's and Neurodegenerative Diseases, Peter O'Donnell Jr. Brain Institute, University of Texas Southwestern Medical Center, Dallas, Texas 75390
[2]Molecular Biology Institute, University of California, Los Angeles; Howard Hughes Medical Institute, Los Angeles, California 90095
[3]Department of Biomedical Sciences of Cells & Systems, University Medical Center Groningen, University of Groningen, Groningen, 9713 AV, The Netherlands
[4]Department of Biochemistry, University of Texas Southwestern Medical Center, Dallas, Texas 75390
[*]Current address: Institute for Neurodegenerative Diseases, University of California San Francisco, San Francisco, California 94158

[#]Corresponding Author: Lukasz.Joachimiak@utsouthwestern.edu

SUMMARY

J-domain protein (JDP) molecular chaperones have emerged as central players that maintain a healthy proteome. The diverse members of the JDP family function as monomers/dimers and a small subset assemble into micron-sized oligomers. The oligomeric JDP members have eluded structural characterization due to their low-complexity, intrinsically disordered middle domains. This in turn, obscures the biological significance of these larger oligomers in protein folding processes. Here, we identified a short, aromatic motif within DNAJB8, that drives self-assembly through π-π stacking and determined its X-ray structure. We show that mutations in the motif disrupt DNAJB8 oligomerization *in vitro* and in cells. DNAJB8 variants that are unable to assemble bind to misfolded tau seeds more specifically and retain capacity to reduce protein aggregation *in vitro* and in cells. We propose a new model for DNAJB8 function in which the sequences in the low-complexity domains play distinct roles in assembly and substrate activity.

INTRODUCTION

Cellular protein homeostasis is maintained by a complex network of molecular chaperones that regulate protein folding, refolding[1-3] and degradation[4-5]. Disruption of this network increases the accumulation risk of proteins with folds that are prone to form proteotoxic β-sheet-rich amyloids, which are the hallmarks of many neurodegenerative diseases[6-8]. For these amyloidogenic proteins, short sequence motifs contribute to the formation of aggregates[9-10]. The challenge for the cell is to disseminate protein conformations with these exposed motifs from non-pathogenic conformations that bury or sequester these motifs. Despite the critical role that chaperones play in preventing the formation of misfolded proteins, there are several lingering questions as to how they can discriminate between normal vs. disease conformations of substrates.

One of the primary chaperones involved in the cell proteostasis network is the highly conserved protein chaperone, Hsp70, but one of the major weaknesses of this chaperone is its broad recognition of substrates[2, 4, 11]. This lack of specificity makes it a weak target in aggregation prevention. Given that there is a vast repertoire of possible unique sequence motifs that are exposed in amyloidogenic substrates, Hsp70 requires additional co-chaperones to facilitate specific recognition of sequences and substrates[2-3]. This additional specificity is imparted by a family of J-domain proteins (JDPs) that classically coordinate substrate activities with Hsp70s[1-2, 12-14]. In the human proteome, JPDs are encoded by 47 known members and possess a diverse range of specialized domains for substrate recognition[1-2, 15-18]. Members of the JDP family bind to Hsp70 through a highly conserved J-domain (JD), which in turn triggers ATP hydrolysis within the nucleotide binding domain of Hsp70 and the closure of the substrate binding domain lid onto the protein substrate[1-2, 19]. The substrate is then refolded by Hsp70 and either restored to its native state or passed along for further processing by other chaperones such as Hsp90[1, 20-21]. The initial substrate recognition step by JDPs is crucial in the protein refolding network given the wide breadth of unique proteins in the cell. It is important that JDPs not only recognize unique protein sequences, but also identify irregular conformations that render proteins dysfunctional or potentially toxic. This is especially true for amyloidogenic proteins, where specific JDPs are able to recognize multiple distinct conformations or oligomeric states of a single protein substrate.

The microtubule-associated protein tau (herein; tau), which forms amyloids in a myriad of neurodegenerative diseases[3, 22-25], is an example of one of these substrates. Tau is associated with several neurodegenerative diseases that are categorized as tauopathies[23, 26]. One of the most striking features of each tauopathy is that while all tauopathies are defined by the presence of tau aggregates in the brain, each tauopathy arises from several different tau polymorphs. Recently, cryo-EM has been used to solve the structures of several different tau fibril cores that are defined by arrangements of the tau repeat domains (RD)[27-28]. As revealed by these structures, the variations in how the RDs are arranged lead to different sequence motifs becoming exposed or buried depending on the specific conformation. This means that a single protein recognition mechanism of tau is insufficient to recognize all possible polymorphs across different tauopathies. The wide diversity of the JDP family compensates for this complication, and provides motif-specific diversity in targeting aggregation-prone tau[29-30].

There are a number of JDPs that have been shown to recognize different conformations of tau monomers or small oligomers (DNAJA1, DNAJA2, DNAJC5)[29-33], as well as larger oligomers and fibrils (DNAJB1, DNAJB4)[29-30, 34]. Recent work from our lab has shown that tau can adopt alternative aggregation-competent conformations (i.e. seeds) of monomer[35-36] that are formed early in disease prior to the detection of oligomers and insoluble fibrils[37]. Derived from these studies, we showed that DNAJC7 preferentially interacts with natively folded conformations of tau over pathogenic mutants or monomeric seeds[38]. However, the high sequence variability and diversity across different JDPs[2] has made identifying JDPs that interact with different tau species difficult. Thus, it remains unknown how JDPs recognize the diversity of species adopted by a single protein.

A number of JDPs are poor candidates for use in classical structural biology methods because they often encode domains that are largely unstructured and dynamic[2]. Some JDPs assemble as dimers (DNAJA1/A2 and DNAJB1)[29] while others remain monomeric. For a few, the unstructured regions in some JDPs can lead to self-assembly, such as DNAJB6 and DNAJB8, which are known to form polydisperse oligomers[39-43]. DNAJB6 and DNAJB8 evolutionarily belong to a distinct subgroup of the

DNAJB1 subclass. Whilst encoding similar architectures that include an N-terminal JD and a flanking unstructured glycine/phenylalanine-rich (G/F) domain, the regions C-terminal of this G/F region are entirely different, with DNAJB6 and DNAJB8 containing serine-/threonine-rich (S/T) domains and a CTD that can interact with the JD, thereby regulating recruitment of Hsp70[39, 43]. Previous work has shown that in DNAJB8, phenylalanines in the G/F and S/T domains are required for its assembly into polydisperse oligomers[39, 42], which reach sizes larger than what is reported for DNAJB6 oligomers[40]. Additionally, we and others have shown that the folded JD and CTD can interact, thereby regulating recruitment of Hsp70[39, 43]. It remains unclear what role oligomerization of DNAJB6 and DNAJB8 plays in their activity, but both were initially discovered as potent inhibitors of amyloid protein assembly[40-41, 44-47], with DNAJB8 showing a slight advantage in cells[41].

Initial genetic studies showed that DNAJB6 is essential for prevention of poly-glutamine (poly-Q) aggregation of the Huntington protein (Htt)[42]. Later studies showed that these oligomeric JDPs are active across a wide range of substrates including α-synuclein[46], Aβ[47], TDP-43 and others [45,48]. Studies on DNAJB6 and DNAJB8 interactions with Htt uncovered that the S/T domain is essential for substrate activity and its deletion[42] or mutation[44] interferes with Htt substrate activity. Studies to determine mechanism indicated that DNAJB6 may function to block nucleation of Htt aggregation consistent with chaperone activity at substoichiometric ratios[44]. Ultimately, the detailed mechanism of DNAJB6 and DNAJB8 substrate activity and the role of oligomerization in this process remain poorly understood.

Here, we used multi-disciplinary methods to understand the minimal driving forces that govern DNAJB8 self-assembly and how DNAJB8 oligomerization may influence substrate activity. We identified a [147]AFSSFN[152] motif within the S/T-rich domain that drives assembly. Mutations of F151 to serine in both the peptide and FL DNAJB8 yields a loss of self-assembly. And yet, using a series of DNAJB8 mutants, we provide *in vitro* and biological evidence that self-assembly is not essential for the anti-amyloidogenesis functionality of DNAJB8. Our data provide mechanistic insight into the detailed mode of DNAJB8 assembly, allowing us to decouple JDP self-assembly and substrate activity.

RESULTS

Phenylalanine residues in the S/T-rich domain drive self-assembly

DNAJB8 contains four domains (Figure 1A): the N-terminal conserved JD (Figure 1A; red), the disordered G/F-rich (Figure 1A; blue) and S/T-rich (Figure 1A; cyan) domains, and a β-stranded auto-inhibitory CTD (Figure 1A; green). In a previous study, we mutated all 18 phenylalanines to serine in the DNAJB8 disordered G/F-rich and S/T-rich domains yielding a variant that is monomeric and unable to self-assemble[39]. We set out to understand how the different 18 phenylalanines in the G/F and S/T domains of DNAJB8 influence its self-assembly. We first considered the distribution and density of phenylalanines within the G/F and S/T domains. We find that the phenylalanines in the middle of the G/F-rich domain contains two dense clusters of phenylalanine residues, while the periphery of the G/F-rich and S/T-rich domains possess a less dense but more evenly distributed arrangement of phenylalanine residues (Figure 1A). Guided by this analysis, we divided the G/F and S/T domains into four segments (DNAJB8$_{S1}$: F87S, F93S, F100S, F103S, F104S; DNAJB8$_{S2}$: F110S, F112S, F114S, F119S, F134S; DNAJB8$_{S3}$: F138S, F141S, F144S, F148S, F151S; and DNAJB8$_{S4}$: F164S, F169S, F179S) to capture the phenylalanine clusters while maintaining similar numbers of phenylalanines in each segment (Figure 1A). We next tested the capacity of FL DNAJB8 S1-S4 mutants to self-assemble in cells and *in vitro* (Figure 1B).

To determine if the serine mutations are changing the size and assembly of recombinant WT and mutant DNAJB8 *in vitro*, we used dynamic light scattering (DLS) to measure changes in the distribution of hydrodynamic radii (R_h) over time. We observed that DNAJB8$_{WT}$ demonstrated evidence of assembly into larger oligomers (>100nm) over 30 hours, with an initial average R_h of 7.9 ± 0.2 nm, indicating the presence of dimers and small oligomers (Figure 1C; blue). This was comparable to the R_h distribution of DNAJB8$_{S1}$ 8.8 ± 0.1 nm and DNAJB8$_{S2}$ 7.3 ± 0.2 nm (Figure 1C; green, gold). However, the average particle size of DNAJB8$_{S3}$ and DNAJB8$_{S4}$ were smaller at 2.4 ± 0.1 nm and 4.5 ± 0.6 nm, respectively (Figure 1C; cyan, grey). These values are comparable to the size of monomeric DNAJB8 with all phenylalanines mutated to serine that we previously characterized[39]. Based on the initial sizes of each mutant, the

similarity of DNAJB8$_{S1}$ and DNAJB8$_{S2}$ to DNAJB8$_{WT}$ are due to their capacity to form dimers that lead to larger assemblies as we previously characterized[39]. By contrast, DNAJB8$_{S3}$ and DNAJB8$_{S4}$, remain small, consistent with sizes expected for a monomer[39]. These data show that the S/T-rich domain contains the specific phenylalanines that drive DNAJB8 oligomerization.

Recently, alternative classes of nonpolar motifs enriched in aromatic side chains have been implicated in protein self-assembly described as low complexity aromatic-rich kinked segments (LARKS)[9, 49]. These motifs stabilize β-sheet conformations that self-assemble through nonpolar side chain contacts and β-strand hydrogen bonds often detectible with Thioflavin T (ThT). Based on this information, we hypothesized that a short, LARKS-like motif could be driving DNAJB8 oligomerization based on the evenly spaced arrangement of Phe residues in the S/T-rich domain (Figure 1A). To consider this possibility, we used a ThT fluorescence aggregation assay to compare β-sheet-mediated assembly of our DNAJB8 mutants. DNAJB8$_{WT}$ and DNAJB8$_{S1}$ showed the highest ThT fluorescence signal at 181.3 ± 23.6 F.U. and 196.3 ± 15.2 F.U., respectively (Figure 1D; blue and green). DNAJB8$_{S2}$ and DNAJB8$_{S4}$ showed an approximate 2-fold decrease in fluorescence signal relative to WT with 104.0 ± 3.0 F.U. and 97.0 ± 9.9 F.U., respectively (Figure 1D; gold and grey). However, DNAJB8$_{S3}$ did not yield any fluorescent signal, consistent with the inability to form β-sheet-rich oligomers (Figure 1D; cyan). These results support that mutations in DNAJB8$_{S3}$ are indeed the most disruptive to DNAJB8 self-assembly, and further strengthen the hypothesis that a short sequence motif may drive the self-assembly of DNAJB8 via β-sheet structures.

While our DLS and ThT data have shown that DNAJB8 oligomerization can be disrupted by serine substitutions within the S/T-rich domain, the next question is whether or not this behavior translates into cells. One known marker of DNAJB8 assembly in cells is the presence of puncta[39, 43], which could be an indication of large oligomers or inclusions that interact with potential substrates. To test the effect DNAJB8 serine segment mutants on puncta formation in cells, we expressed DNAJB8$_{WT}$, DNAJB8$_{S1}$, DNAJB8$_{S2}$, DNAJB8$_{S3}$, and DNAJB8$_{S4}$ mutants as C-terminal fusions to

mRuby3 in HEK293T cells. Cells were transfected, incubated for 48 hours at 37°C and imaged on a high content microscope. The images were analyzed to quantify the frequency, size and intensity of DNAJB8 puncta based on mRuby3 fluorescence (see methods). Quantification of fluorescence intensity showed even levels of protein mRuby3-based expression (Figure S1A) with no effects on viability (Figure S1B). As previously observed[39], expression of DNAJB8$_{WT}$-mClover3 led to the formation of cytoplasmic and juxtanuclear puncta with an average diameter of 1.1 ± 0.3 µm with signal excluded from the nucleus (Figure 1E and Figure S1C). While a similar frequency of puncta was observed for cells expressing DNAJB8$_{S1}$ and DNAJB8$_{S2}$, there was a 5-fold and 3-fold decrease in the number of puncta in cells expressing DNAJB8$_{S3}$ and DNAJB8$_{S4}$, respectively (Figure 1F). Furthermore, while cells expressing DNAJB8 that encoded the S1, S2 and S4 mutations displayed the same cytoplasmic localization as DNAJB8$_{WT}$-mRuby3 (Figure S1E-G+I), we observed that DNAJB8$_{S3}$ was present in both the cytoplasm and the nucleus (Figure 1E and Figure S1H). Mutations of phenylalanine to serine in S3 segment reduce the number of DNAJB8 puncta and promote nuclear localization (Figure 1F). These data show that as observed *in vitro*, phenylalanine to serine substitutions within the S/T-rich domain are changing how DNAJB8 assembles in the cell.

A steric zipper motif drives local S/T-rich domain self-assembly

Based on the results of the cell and *in vitro* experiments, we identified the phenylalanine residues in DNAJB8 self-assembly are among 5 residues in segment 3: F138, F141, F144, F148, and F151. Guided by the presence of detectible ThT signal in oligomeric DNAJB8 (Figure 1F), we designed a ThT aggregation-based screen of peptides across the disordered G/F and S/T domains (Figure 2A; Supplementary Table 1). All peptides used in this experiment were non-overlapping 20 amino acids fragments and spanned from the beginning of the G/F domain to the end of the C-terminal domain. Each peptide was monitored for its capacity to self-assemble using a ThT fluorescence assay over 10 hours. Only one peptide, B8_5, within the screen showed evidence of self-assembly by ThT, which includes 3/5 phenylalanines found in segment 3 of DNAJB8 (Figure 2B).

Negative stain TEM confirmed the presence of fibril-like assemblies for B8_5 (Figure 2C). Interestingly, sequence-based prediction of amyloid motifs[10] in the B8_5 fragment did not predict amyloidogenic properties (Figure S2A) suggesting that this fragment may adopt assemblies distinct from previously described steric zippers. Nevertheless, our experimental results indicate that a minimal motif found in segment 3 may be responsible for ordered self-assembly of DNAJB8.

To gain structural insight into how DNAJB8 fragments may assemble, we generated a library of hexapeptides derived from the B8_5 fragment and screened them for the formation of crystals suitable for X-ray structure determination. Only one hexapeptide, [147]AFSSFN[152], yielded diffracting crystals. We determined a 0.75 Å resolution crystal structure of the [147]AFSSFN[152] assembly revealing a class 6 steric zipper motif[9] (Figure 2D, Figure S2B, Supplementary Table 2). The assembly is defined by two pairs of anti-parallel β-strands stabilized through hydrophobic interactions along opposite faces (face-to-back)[9]. The β-strand dimers show 2-fold symmetry within the assembly (Figure 2D and Figure S2C). The hydrophobic core of the assembly contains F151 on one dimer of β-strands surrounded on the edges by F148 from a second dimer. Within the core of the assembly, both F151 residues on a single dimer are oriented towards the center of the tetramer, but only one residue of F148 is oriented towards the center (Figure 2D).

To distinguish which phenylalanine position is the most important in stabilizing the dimer, we minimized the structures in Rosetta[50] and calculated the solvent-accessible surface area (SASA) of each residue. The analysis was run in the context of the peptide dimer that was derived from the crystal symmetry (Figure 2E). We calculated the SASA for every residue across each peptide unit (labelled as "chains"). We first ran each chain as independent monomers and found that both F148 and F151 are relatively exposed (Figure S2E). In the context of the dimer, F148 remains exposed while F151 becomes more buried (Figure 2E; Figure S2F). When considering the orientation of the aromatic rings, we find that while the F151 pair interacts with each other across 2 chains to stabilize the structure, only one residue of F148 is oriented towards the center of the dimer, while the other is exposed to the side (Figure S2D, left). This leads to a lower

SASA for F151, where both instances of F151 contribute to each other's burial within the minimal unit, while F148 does not (Figure 2E; Figure S2F). This implies that F151 burial is an important interaction to stabilize intermolecular contacts within the dimer.

To analyze the importance of F148 and F151 in the context of oligomeric assemblies, we used the dimer as a base unit and expanded the assemblies in a horizontal and vertical direction. Upon expansion of the assembly in a horizontal direction, we find that both F148 and F151 are buried, where F148-F148 interactions are buried in the horizontal plane (Figure 2F and Figure S2D, right). Meanwhile, F151 burial is independent of this horizontal axis and does not require additional dimer-dimer interactions to stabilize itself (Figure 2F; Figure S2G). In the vertical expansion of the dimer assembly, F148 shows higher SASA than the horizontal assembly due to a greater exposure of the side chains to the solvent (Figure 2G; Figure S2H). However, partial burial is observed from the chains above and below due to the hydrophobic F151 face (Figure 2G; Figure S2D, center). Analysis of the different interfaces formed in the crystal lattice reveals differential burial modes of F148 and F151. The horizontal assembly is more dependent on the F148-F148 interactions, while the vertical assembly is more dependent on the formation of the F151 hydrophobic face. Therefore, we can conclude that F151 is crucially important for the initial dimer formation and subsequent oligomeric assemblies, while two possible modes of oligomeric assemblies could exist that are dependent on the aromatic ring orientation of F148.

Burial of phenylalanine at position 151 dictates assembly of DNAJB8 fragments

Given the importance of phenylalanine burial in the different geometries of the steric zipper, we hypothesized that mutations to these positions would either weaken or disrupt the formation of this motif, leading to disruption of self-assembly. To test this, we introduced mutations to the B8_5 fragment to determine the role of aromatic side chains (F148S, F151S, F148S_F151S) and β-sheet conformation (S149P, S149P_S150P). As before, we used a ThT fluorescence aggregation assay to monitor self-assembly of mutant peptides compared to the WT B8_5 sequence (Figure 2H). The results showed that self-assembly of the B8_5 with the F148S mutation was reduced, and self-

assembly was abolished with the F151S and F148S_F151S mutations. Furthermore, all B8_5 peptides containing a single or double serine to proline mutation in B8_5 were not able to self-assemble suggesting that backbone contacts are important for oligomerization.

We next determined how phenylalanine to serine mutations may influence the total SASA of the dimer and the horizontal and vertical hexamers. The F148S, F151S, and F148S_F151S mutations were introduced into the ^{147}AFSSFN^{152} structure using Rosetta and minimized as previously described[50]. The ΔSASA values were calculated by comparing the SASA between WT SASA and each mutant across the dimer, horizontal and vertical hexamer assemblies (Figure S2E). Compared to the WT sequence, the F148S mutation led to a loss of SASA in both the dimer and horizontal assemblies, while F151S resulted in a gain of SASA (Figure S2F, salmon and brown). Given the disengagement of F148 in the structure, it is expected that a mutation to this position would lead to a greater loss of SASA due to the loss of the side chain. However, the mutation to F151 does not result in such a loss since both F151 residues are more buried in the structure due to their greater role in stabilizing the packing of the zipper. The double mutant, F148S_F151S showed additive effects from both mutations, which resulted in a net loss within the dimer, but no change within the horizontal assembly (Figure S2F, green). However, when considering the effect of the mutations on the SASA of the vertical assembly, we observed an overall gain of SASA from both single mutants and the double mutant (Figure S2F, right). These results are more consistent with the hypothesis that both F148 and F151 participate in SASA burial to a certain degree within the larger peptide assembly if the directionality is vertical. However, one explanation that is consistent with our data is that while F148 plays a role in stabilizing larger oligomers, F151 is critical for the initial peptide dimer formation that precedes the initiation of self-assembly.

We further tested the homologous peptide sequences within DNAJB6b to see if the S/T domain sequences spontaneously self-assemble similar to the B8_5 fragment in DNAJB8. No similar mode of assembly was observed for DNAJB6b S/T domain fragments or CTD sequences that were previously implicated in DNAJB6b assembly[51-52]

(Figure S2G). These data indicate that the observed steric zipper structure resulting from the [147]AFSSFN[152] motif is unique to DNAJB8 and not shared by the DNAJB6b homolog, with DNAJB6b self-assembly driven through other molecular forces. Overall, these data validate that the [147]AFSSFN[152] steric zipper motif controls B8_5 fragment assembly.

Phenylalanine at position 151 is the primary driver of FL DNAJB8 self-assembly

We next tested whether mutations designed to disrupt the steric zipper motif formed by [147]AFSSFN[152] also inhibited assembly of FL DNAJB8 in vitro and in cells. To test this, recombinant FL DNAJB8 constructs containing F148S, F151S, or F148S_F151S (herein DNAJB8$_{F148S}$, DNAJB8$_{F151S}$, DNAJB8$_{F148S\_F151S}$, respectively) were purified and their assembly tested in ThT fluorescence and DLS assays *in vitro* (Figure 3A). As observed in the peptide experiments, DNAJB8$_{F148S}$ yielded a similar fluorescence signal as DNAJB8$_{WT}$, but DNAJB8$_{F151S}$ and the double mutant DNAJB8$_{F148S\_F151S}$ showed a decrease in ThT signal with the double mutant approaching the same fluorescence amplitude as DNAJB8$_{S3}$ (Figure 3B). The capacity of single and double mutants to self-assemble was next confirmed using DLS. The initial average R_h of DNAJB8$_{F148S}$ was 13.9 ± 0.7 nm, which is comparable to WT (Figure 3C). Furthermore, evidence of DNAJB8 oligomers was observed after 30 hours in both DNAJB8$_{F148S}$ and DNAJB8$_{WT}$ (Figure 3C). However, DNAJB8$_{F151S}$ and DNAJB8$_{F148S\_F151S}$ showed an inability to self-assemble over time and remained monomeric at 2.5 ± 0.5 nm and 2.5 ± 0.05 nm respectively, comparable to DNAJB8$_{S3}$ and the previously reported mutant with all phenylalanines mutated[39]. These data suggest that the F151S mutation disrupts β-strand-dependent self-assembly *in vitro*, with the inclusion of F148S yielding an additive effect.

We next measured puncta formation of DNAJB8-mRuby3 encoding F148S, F151S, and F148S F151S mutations (herein DNAJB8$_{F148S}$-mRuby3, DNAJB8$_{F151S}$-mRuby3, and DNAJB8$_{F148S\_F151S}$-mRuby3, respectively) in HEK293T cells. After transfection, the cells were incubated for 48 hours at 37°C before image collection on a high content microscope and the images were analyzed as before. The average mRuby3

fluorescence intensity was similar across all the variants tested and expression of the proteins did not affect cellular viability (Figure S3A-B). While DNAJB8$_{F148S}$-mRuby3 remained primarily within the cytoplasm and similar to DNAJB8$_{WT}$-mRuby3, we observed that DNAJB8$_{F151S}$-mRuby3 and DNAJB8$_{F148S\_F151S}$-mRuby3 both showed a decrease in puncta and both mutants were now able to enter the nucleus, similar to DNAJB8$_{S3}$-mRuby3 (Figure 3D). While decreases in frequency of puncta were observed across all mutants, DNAJB8$_{F148S}$-mRuby3 showed the smallest decrease with 9.7 ± 1.8% of cells with puncta, while DNAJB8$_{F151S}$-mRuby3 and DNAJB8$_{F148S\_F151S}$-mRuby3 showed a decrease with 6.9 ± 2.4% and 3.8 ± 0.4% of cells with puncta, respectively (Figure 3E). Consistent with the peptide experiments, our results on FL JDP support the importance of F151 in DNAJB8 self-assembly, with evidence that additional phenylalanine to serine substitutions in tandem with F151 decrease the oligomeric propensity of DNAJB8, yielding predominantly monomers. The fact that this motif is found within the S/T-rich domain is further supported by previous data that indicated the importance of the S/T-rich domain in self-assembly[42, 44].

DNAJB8 preferentially binds a misfolded tau seed

The M$_s$ form of tau can self-assemble into amyloids and can induce aggregation of recombinant tau. Comparison of the of M$_s$ and normal inert tau (M$_i$) ensembles uncovered differences in local interactions but also folding of the N-terminal acidic tail[35-37]. We have also previously developed assays to measure chaperone activity against both M$_i$ and M$_s$ forms of tau to show differences in DNAJC7 binding and activity[38]. While oligomeric JDPs, such as DNAJB6 and DNAJB8, have not previously been reported to interact with tau, other class B JDPs are known to interact with tau fibrils[16, 29-30]. Here, we tested directly whether DNAJB8$_{WT}$ or DNAJB8$_{S3}$ can discriminate between M$_i$ and M$_s$ to influence tau aggregation.

One of the most challenging complications in measuring DNAJB8$_{WT}$ substrate binding affinity is the difficulty to parse out whether monomers or oligomers are binding to substrates. However, our DNAJB8$_{S3}$ mutant remains monomeric, which eliminates this problem. Using this advantage, we first leveraged our engineered monomeric

DNAJB8$_{S3}$ to measure binding to M$_i$ and M$_s$ using microscale thermophoresis (MST) (Figure 4A). Titration of unlabeled M$_i$ against Cy5 labeled DNAJB8$_{S3}$ yielded a K$_d$ > 400 µM indicating that the binding is very weak (Figure 4B, black and Figure S4A). Similarly, we also measured binding between M$_s$ and DNAJB8$_{S3}$ yielding a substantially tighter K$_d$ of 24 µM (Figure 4B, red and Figure S4B). This change in affinity may be caused by differences in M$_i$ and M$_s$ conformations as was previously shown[35-36, 53], but most importantly, DNAJB8 is showing preferential binding to the aggregation-prone conformation of tau.

Based on these data, we were interested in understanding whether the WT and mutant JDP may influence assembly of different forms of tau. We developed an assay to monitor crosslinking-based assembly of FITC-labeled M$_i$ (FITC-M$_i$) tau and quantify whether DNAJB8 can influence this process. By treating reactions with a crosslinker, depletion of monomer and formation of covalent adducts can be visualized on an SDS-PAGE gel. Based on prior crosslinking experiments on tau[36, 53] and DNAJB8[39], we optimized the reactions using 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methyl-morpholinium chloride (DMTMM) which traps contacts between primary amines and carboxylates. 1µM FITC-M$_i$, 20µM DNAJB8$_{WT}$ and FITC-M$_i$:DNAJB8$_{WT}$ were treated with crosslinker for 0, 7.5, and 15 minutes, quenched and visualized on an SDS-PAGE gel. DNAJB8$_{WT}$ shows a slow reduction of monomer as crosslinker incubation time increases, consistent with its capacity to self-assemble (Figure S4C-D). Meanwhile, M$_i$ in the presence of DMTMM shows evidence of rapid assembly, with monomers being depleted rapidly into larger oligomers (Figure S4C-D). Quantification of the FITC-M$_i$ band intensities reveals a near 10-fold depletion of tau (Figure 4C; black). By contrast, addition of DNAJB8$_{WT}$ to M$_i$ slows down the depletion of M$_i$ by 5-fold (Figure 4C and Figure S4C-D). We also performed a parallel experiment with FITC-M$_i$:DNAJB8$_{S3}$. Consistent with DNAJB8$_{S3}$ being monomeric, reactions containing only the mutant JDP, show little assembly (Figure S4E). Despite the inability to assemble, co-incubation of FITC-M$_i$ with DNAJB8$_{S3}$ yielded a 6-fold reduction of M$_i$ depletion (Figure 4D). The close similarities in tau monomer retention between DNAJB8$_{WT}$ and DNAJB8$_{S3}$ show that both JDPs are active in our M$_i$ tau assembly assay despite weak affinities.

To gain more resolution into the surfaces involved in the interactions between the JDP and tau we employed crosslinking mass spectrometry. While other structural approaches are not suitable for low affinity interactions, crosslinking mass spectrometry (XL-MS) can uncover binding surfaces for even weak interactions[39, 54]. In addition to probing interactions between DNAJB8$_{WT}$ or DNAJB8$_{S3}$ and M$_i$, we also probed interactions between the two chaperones and tau M$_s$. Triplicate samples were analyzed by mass spectrometry and processed using the Xquest pipeline[54-55]. In the DNAJB8$_{WT}$:M$_i$ and DNAJB8$_{S3}$:M$_i$ containing data, the JD and CTD interact with different tau domains indiscriminately, suggesting weak non-specific interactions (Figure 4E-F; Figure S4G-H). By contrast, in the DNAJB8$_{WT}$:M$_s$ and DNAJB8$_{S3}$:M$_s$ datasets the patterns of interaction are different with both chaperones preferentially crosslinking to the PRD domains (Figure 4G-H; Figure S4I-J). Calculating the frequency of the contacts as a function of the entire dataset shows that the percentage of contacts to the PRD in the M$_i$ experiments is 20% and increases to 60% in the M$_s$ datasets (Figure 4E-H). Our XL-MS data support that M$_s$ is folded differently than M$_i$ consistent with our prior work[53] and that this change in conformation leads to more specific contacts between the JDPs (both WT and S3) to M$_s$.

To test whether we can observe aggregation suppression of tau in the presence of DNAJB8$_{WT}$, we used a thioflavin T fluorescence-based aggregation assay where a recombinantly derived minimal construct of tau containing only the repeat domains (tauRD) is seeded using tau M$_s$. Our results show that at substoichiometric concentrations, tau M$_s$ induces rapid tauRD aggregation (Supplementary Figure 4k; orange, t$_{1/2max}$= 6.3 ± 0.2 hours). Addition of DNAJB8$_{WT}$ slows down M$_s$-seeded tauRD aggregation nearly 10-fold (Figure S4K; black, t$_{1/2max}$= 61.6 ± 2.9 hours). While these results show that the presence of DNAJB8$_{WT}$ does delay tau aggregation, it is difficult to comment on how DNAJB8:M$_s$ interactions specifically effect the nucleation stages of an aggregation experiment. Therefore, to test whether the chaperones can influence M$_s$ self-assembly directly, we designed a DLS-based assay using low concentrations of M$_s$ monomers (0.1µM) that are invisible to detect by scattering but once they form assemblies they scatter and can be detected directly. This experiment also removes M$_i$ from the system, and unlike a ThT assay, directly reports on the isolated M$_s$

conformation assembly at a concentration where ThT signal cannot be observed. Importantly, parallel reactions with 10μM WT and mutant JDPs remain flat in the assay allowing us to quantify their effect on M_s assembly by comparing the $t_{1/2max}$ of the reactions (Figure 4I). M_s alone assembled with a $t_{1/2max}$ of 24.7 ± 0.2 hours (Figure 4I, red), addition of DNAJB8$_{WT}$ delays the increase in M_s assembly with a $t_{1/2max}$ of 33.8 ± 0.3 hours (Figure 4I, blue). By contrast, reactions that included DNAJB8$_{S3}$ delayed the M_s assembly even further yielding a $t_{1/2max}$ of 54.3 ± 7.5 hours (Figure 4I, light blue). The addition of other mutants DNAJB8$_{S1}$, DNAJB8$_{S2}$, and DNAJB8$_{S4}$ delayed the increase of M_s assembly similarly to DNAJB8$_{WT}$ (Figure S4L). Since DNAJB8$_{S3}$ is a monomer at these concentrations, these results show that self-assembly is not a pre-requisite capacity for DNAJB8 to act as a holdase.

Engineered monomeric DNAJB8 mutants retain anti-aggregation substrate activity in cells

Our data show that DNAJB8 is able to influence aggregation of tau *in vitro* despite weak binding affinities. We next tested whether our engineered DNAJB8 mutants are active against substrate aggregation in cells using tau and Htt as model substrates. We first used a cellular model of tau aggregation that co-express P301S tauRD-mCerulean3 and P301S tauRD-mClover3 in HEK293T cells (Figure 5A). Transduction of tau aggregates into these cells induces aggregation of the intracellular tau which can be quantified using FRET$_{Clover/Cerulean}$[23, 56-57]. To test the effect of WT and mutant DNAJB8 on tau aggregation in this system, we co-transfected the tau biosensor cells with WT and mutant DNAJB8 C-terminally fused to mRuby3, followed by seeding the cells with pre-formed tau fibrils using lipofectamine (Figure 5A). We next interpreted changes in tau aggregation as measured by FRET$_{Clover/Cerulean}$ but also changes in DNAJB8 oligomerization state as described above (Figure 1D and 3E). Parallel experiments were setup for imaging analysis and flow cytometry. After 48 hours following tau fibril treatment, one plate of cells was imaged to analyze tau and DNAJB8 inclusions, while a second plate was analyzed by flow cytometry to quantify tau FRET$_{Clover/Cerulean}$ in each condition. Using DRAQ5 for nuclear staining, we were able to produce images showing

DRAQ5, tau-mClover3, and DNAJB8$_{WT}$-mRuby3/DNAJB8$_{S3}$-mRuby3 for seeded and unseeded conditions (Figure S5A). In unseeded cells, the tau expression is diffuse throughout the cells (Figure S5A). Treatment of cells with 20nM fibrils leads to the formation of mClover3 positive tau puncta (Figure S5A). Consistent with our earlier experiments in HEK293T cells expressing only the chaperone, cytoplasmic DNAJB8 puncta were observed in cells expressing DNAJB8$_{WT}$-mRuby3 (Figure S5A; yellow arrows), while DNAJB8$_{S3}$-mRuby3 remained diffuse in the tau biosensor cell lines. In cells that have both DNAJB8$_{WT}$-mRuby3 and tauRD-mClover3 puncta, the two inclusions do not colocalize (Figure S5A).

To evaluate the effect of DNAJB8 variants on tau seeding, we quantified the FRET$_{Clover/Cerulean}$ signal in mRuby3 positive cells (Figure S5B). Cell lines not treated with tau fibrils had no detectable FRET$_{Clover/Cerulean}$ (Figure 5B). For cells expressing mRuby3 alone, 16.5 ± 2.5% of the cells were FRET positive for tau aggregates, while cell lines expressing DNAJB6b$_{WT}$-mRuby3 or DNAJB8$_{WT}$-mRuby3 lead to a statistically significant 2-fold reduction in signal compared to mRuby3 alone (Figure 5B). We compared the %FRET$_{Clover/Cerulean}$ in the cell populations expressing DNAJB8 mutants and reported the results normalized to DNAJB8$_{WT}$ (Figure 5C). All seeded tau biosensor cell lines expressing DNAJB8 mutants yielded small increases in seeding compared to DNAJB8$_{WT}$ but were not statistically significant, which suggests these mutations are not important for DNAJB8 holdase activity on tau. Cell viability was comparable across all DNAJB8-expressing tau biosensor lines including controls (Figure S5C), thus confirming that the processes were not affected by toxicity in response to co-expression of tau and DNAJB8. Consistent with our *in vitro* results, the designed DNAJB8 mutants that are monomeric retain activity similar to DNAJB8$_{WT}$.

Using the mRuby3 tag on our DNAJB8 mutants, we performed an image-based analysis on each tau biosensor cell line to determine if the frequency of DNAJB8 puncta has changed in the presence of tau as described in Figure 1. First, we quantified mRuby3 signal across all lines transfected with a mRuby3-fusion construct to look at baseline expression. We observed similar variations in expression regardless of seeding, where relative to DNAJB8$_{WT}$, the only mutants that showed a significant increase in mRuby3

signal per population were DNAJB8$_{S2}$, DNAJB8$_{S3}$, DNAJB8$_{S4}$, DNAJB8$_{F148S}$, and DNAJB8$_{F148S\_F151S}$ (Figure S5D). Overall, the average size of puncta and signal intensity was consistent across all cell lines (Figure S5E-F). Control cell lines expressing mRuby3 alone under unseeded and seeded with 20nM tau fibrils conditions yielded low frequency of puncta (Figure 5D). Expression of DNAJB6b$_{WT}$-mRuby3 yielded 2-fold higher frequencies of puncta in unseeded and seeded conditions, and DNAJB8$_{WT}$-mRuby3 yielded 6-fold higher puncta frequency compared to the control (Figure 5D). Under similar conditions, DNAJB8 has a higher propensity to form large assemblies in cells compared to DNAJB6b. When we compared our DNAJB8 variants to the puncta frequency observed in DNAJB8$_{WT}$-mRuby3 expressing tau biosensor cells, we observed the same trend as our HEK293 cells where mutations reduced puncta frequency for all mutants, with DNAJB8$_{S3}$ reaching the same frequency as the mRuby3 control (Figure 5E). Our results suggest that the oligomeric assembly propensity of these two oligomeric JDPs is not related to their protein aggregation function, as supported by our data with mutant DNAJB8. Additionally, our data support that while oligomerization is not required for activity, JDP oligomerization may be a mechanism to promote persistence of activity on longer time scales which may reduce cellular toxicity.

Finally, we asked whether our results on tau aggregation suppression by DNAJB8 in cells would translate to the well-established Htt substrate[40-42, 44]. We tested DNAJB8$_{WT}$ and mutants on (non-seeded) HttQ119 amyloidisation in cells using a Filter trap assay (FTA). HEK293T cells were co-transfected with a 1:3 ratio of Htt Q119-GFP: DNAJB8-mRuby3 and the amount of Htt aggregates were quantified by FTA (Figure 5F). Expression of DNAJB8$_{WT}$ and mutants were confirmed to be similar by western blot (Figure S5G). Htt aggregate band intensity in the FTA was quantified and normalized to the mRuby3 negative control and reported as a percentage of aggregates of Q119 in this condition. As reported previously, DNAJB8$_{WT}$ is a potent suppressor of Q119 aggregation yielding no signal (Figure 5G and Figure S5H). Similarly, little signal was detected in cells expressing DNAJB8$_{F148S}$, DNAJB8$_{F151S}$, DNAJB8$_{F148S\_F151S}$, and DNAJB8$_{S1}$ (Figure 5G and Figure S5L). A minor loss in activity was found for DNAJB8$_{S2}$, DNAJB8$_{S4}$, and DNAJB8$_{S3}$ (Figure 5G and Figure S5H), suggesting that most activity is retained even though some DNAJB8 mutants are unable to self-

assemble. To ensure that the observed anti-aggregation activity is not influenced by the presence of endogenous DNAJB6b, we repeated the experiment using a HEK293T cell line in which DNAJB6b was knocked out[58]. Also here, all DNAJB8 variants were able to reduce polyQ aggregation with DNAJB8$_{S2}$, DNAJB8$_{S3}$, and DNAJB8$_{S4}$ relatively being slightly more active than in cells that did express endogenous DNAJB6b (Figure S5I-J).

The minor loss of activity of DNAJB8$_{S2}$, DNAJB8$_{S4}$ and DNAJB8$_{S3}$ is not the result of changes in the oligomeric assembly of DNAJB8 since the monomeric minimal mutants DNAJB8$_{F151}$ and DNAJB8$_{F148S\_F151S}$ retain equal chaperone activity relative to DNAJB8$_{WT}$. Therefore, this loss of activity in DNAJB8$_{S3}$ must be attributed to the loss of hydrophobic residues in the S/T-rich domain which seem more important for Htt activity compared to tau. Our combined data suggest that oligomeric assembly of DNAJB8 is not essential for substrate activity.

DISCUSSION

The capacity of DNAJB6 and DNAJB8 to oligomerize has been established *in vitro* and in cells[39,41-42,43]. However, their mechanism of assembly has been poorly understood and remained intertwined with substrate activity. Previous studies suggested that DNAJB6 assembly may be driven by sequences along the C-terminus of the protein, similar to dimerization observed in other JDPs[51-52]. This is not true for DNAJB8, where the CTD domain is a stable monomer that is unable to assemble[39]. Instead, DNAJB8 oligomerization was abolished by mutating aromatic residues in the disordered G/F-rich and S/T-rich domains[39]. By combining cellular and *in vitro* studies, we identified a small phenylalanine-rich region (i.e. S3) within the S/T-rich domain that is responsible for DNAJB8 self-assembly. We determined an X-ray structure of the minimal assembly motif and by using mutagenesis verified that this motif ([147]AFSSFN[152]) is important for assembly of FL DNAJB8 *in vitro* and in cells. F→S mutations to positions in this motif yields monomeric DNAJB8 mutants *in vitro* with a more exposed S/T domain that retains activity against substrate aggregation *in vitro* and in cells. Our data builds a new paradigm for oligomerization of JDPs and for the first time decouples oligomerization

from substrate activity underscoring that oligomerization is not a prerequisite for holdase activity. The S/T domain of oligomeric JDPs has been implicated in substrate binding activity, but as we discovered, other regions may play important specific substrate conformation and sequence dependent roles.

Molecular interactions that drive DNAJB8 assembly

An emerging concept in the field of biology, is the versatile way in which weak molecular nonpolar or charge complementary interactions can drive reversible (and irreversible) protein assembly to control spatial organization of compartments in cells. Proteins that encode low complexity domains, such as FUS, TDP-43 or HNRP1, use charge complementary or nonpolar residues to drive weak intermolecular interactions that promote phase separation, which result in dynamic and reversible assemblies[59-62]. In particular, aromatic-aromatic residue interactions are especially stabilizing through π-π interactions[62-63]. Intrinsically disordered proteins such as tau, amyloidβ, or α-synuclein have been shown to initiate self-assembly through the formation of steric zippers, which are mediated by a combination of side chain and backbone hydrogen bonds [9, 49]. In addition, the role of backbone interactions in phase separation is unclear, but recent studies have suggested that a combination of backbone and side chain interactions may be critical for phase separation of low-complexity sequences[64]. Our data support that DNAJB8 oligomerization is dependent on both the side chain interactions as well β-strand mediated backbone hydrogen bonds (Figure 6A). The combination of both of these intermolecular forces promotes the initial dimer assembly, followed by the hydrophobic interactions that yield larger oligomeric assemblies. We demonstrate that in both peptides and FL DNAJB8 disruptions to either the side chains (F→S) or the backbone (S→P) result in a loss of oligomeric assembly, indicating that a combination of hydrophobic "stickers" and backbone hydrogen bonds are equally important for DNAJB8 assembly (Figure 6B). While low-complexity sequence motifs in other proteins may not necessarily share the same amino acid identities as DNAJB8, our findings present the idea that similar combinations of forces may be required for controlling reversible large-scale oligomeric assembly of a chaperone in cells. It is perhaps

surprising that a chaperone would encode the capacity to adopt reversible assemblies, but one possible explanation is that the assemblies may be a storage conformation that persists in the cytosol, which can harbor monomers for future release to provide active chaperone conformations on demand.

Oligomeric DNAJB8 interactions with substrate are dynamic

Chaperones often bind to substrates with micromolar affinities, which by design promotes substrate release and refolding[1-2]. This is true for large molecular chaperones such as chaperonins, Hsp90, as well as many JDPs[13, 21, 65-66]. For globular substrates, chaperones interact with unfolded intermediates with weak affinities to prevent aggregation and, upon release, allow refolding into globular functional states. For intrinsically disordered proteins that do not adopt a defined globular fold, how do molecular chaperones regulate their folding? Our lab has uncovered that DNAJC7 binds tightly and specifically to the inert aggregation-resistant conformation of tau with nanomolar affinity[38]. How oligomeric JDPs interact with substrates has remained an open question. The S/T-rich domain in oligomeric JDPs has been implicated in interactions with HttEx1and amyloidβ[44, 47]. Studying how oligomeric JDPs binding to substrates has been hampered by measuring interactions between polydisperse species of chaperone with substrates. This is further confounded by the proposed role of JDPs to interact with misfolded conformations of substrates. In this study we have uncoupled oligomerization of DNAJB8 from substrate activity to create a monomeric form of DNAJB8 that retains activity (Figure 6C). Our methods to produce recombinant seeds of the tau protein[35-36] allowed us to probe whether DNAJB8 mutants have differential activity against normal aggregation-resistant tau compared to a pathogenic monomer seed. Measuring affinities between the monomeric chaperone and these two forms of tau yield dramatic differences in binding affinity consistent with the chaperone preferentially interacting with misfolded seeds[44]. Crosslinking of the mutant JDP to the two different conformations of tau reveals different patterns of interaction, and validates that the seed and normal tau differ in their shape and thus present different surfaces for JDP binding. Perhaps the largest difference is that the chaperone appears to bind more

favorably with the basic PRD domain of tau, which may be stabilized by cation-π interactions. This is consistent with our prior work that the N-terminus rearranges its interaction with the repeat domain in the seeding form compared to M_i[53]. Based on the presently described π-π interactions that drive DNAJB8 oligomerization, it could be possible that large, positively charged surfaces along the tau PRD could interact with phenylalanine rings in a similar manner to cation-π interactions, resulting in a weak-charge based interaction between DNAJB8 and tau. In turn, this minor interaction would promote the extension of the DNAJB8 monomer and disengage the JD from the CTD in preparation for Hsp70 binding.

Despite the low affinities the monomeric JDP retains substrate aggregation prevention activity in vitro and in cells. When considering our activity data with HttEx1 Q119 in cells, the activity differences observed with DNAJB8$_{S3}$ are likely due to the specific substitutions of five Phe positions as opposed to whether or not the protein is capable of self-assembly. This is consistent with previously reported data, where it has been proposed that the S/T-rich domain plays an important role in substrate recognition[42, 44]. This leads to new questions about why the presence of Phe residues in the S/T-rich domain is important for DNAJB8 recognition of HttEx1, but the absence of those same residues improves *in vitro* recognition of tau. One possible explanation may rely upon previous work suggesting that the tau PRD is a common recognition domain to JDPs[30]. This data for the first time uncovers differences in substrate activity and suggests that different surfaces on the JDP may play a role in substrate recognition. This data also raises another question, how does the chaperone influence aggregation of a substrate through low affinity interactions? Intrinsically disordered proteins sample different conformations and this is certainly true for the M_i conformation of tau. Misfolded seeds (i.e M_s) on the other hand adopts more defined alternate conformations. Based on previous experiments, we proposed that M_s preferentially exposes amyloid motifs[35, 53] and more recently we showed that the N-terminus of tau folds differently on the basic repeat domain[36]. The higher binding affinity between the JDP and M_s suggest that the flexibility of normal inert tau lends to low affinities while the more defined and ordered

M_s conformation leads to higher affinity possibly gained through PRD and amyloid motif interactions. Higher resolution structural experiments between engineered JDPs and substrates could uncover the details of these interactions.

Substrate selectivity by oligomeric JDPs

DNAJB6 is ubiquitously expressed, including in the human brain but its expression decreases with age[42, 44, 58]. DNAJB8 expression on the other hand is limited to the testes[67]. Thus, the link between these JDPs and neurodegenerative diseases is weak because the onset of symptoms manifests later in life. Mutations in DNAJB6 are linked to muscular dystrophy but the disease age onset varies[68]. Initial genetic screens discovered these JDPs are regulators of Htt aggregation[40-42, 44] and subsequent studies demonstrated their anti-aggregation activity against other IDPs[43, 45-47, 58]. Why are these JDPs not expressed later in life when neurodegenerative diseases typically develop? Perhaps they only function at early stages of aggregation or alternatively their expression later in life has deleterious effects on other processes. What is the role of these JDPs? New evidence suggests that both DNAJB6 and DNAJB8 interact with FG nucleoporins to help assemble nuclear pores[69-70]. FG Nups and IDPs belong to functionally different molecules and encode distinct sequence features. It is possible that these JDPs function on these two different classes of substrates using different folding mechanisms. Understanding in more detail how they function on substrates may allow us to engineer the different properties to increase their specificity, perhaps allowing them to be developed as therapeutics targeting specific proteins. Our study highlights that engineering DNAJB8 into a monomer allows it to maintain substrate activity but begins to change specificity between tau and Htt suggesting that distinct sequence features of the chaperone are used for different substrates. Future studies must be focused on understanding the role of this family of JDPs in maintaining cellular homeostasis, defining a clearer substrate interactome and understanding the mechanism of activity against substrates with different properties.

METHODS

Cell biological analysis of DNAJB8-mRuby3 lines

All plasmids for cell experiments were prepared by Twist Bioscience. The coding sequences for human DNAJB8$_{WT}$, and the mutants DNAJB8$_{S1}$, DNAJB8$_{S2}$, DNAJB8$_{S3}$, DNAJB8$_{S4}$, DNAJB8$_{F148S}$, DNAJB8$_{F151S}$, and DNAJB8$_{F148S\_F151S}$ were cloned into FM5 lentiviral expression plasmids, in which the UbC promoter was replaced with a CMV promoter, the linker sequence was replaced by "GSAGSAAGSGEF," and the YFP was replaced by mRuby3. The resulting genes produced a DNAJB8-mruby3 fusion protein along with all mutant variants. In parallel, we produced a construct that expresses the fluorescent protein (mRuby3) but lacks any DNAJB8 gene. All plasmids were separately co-transfected into HEK293T cells using lipofectamine 2000 (Invitrogen). 100,000 cells were plated per well in media (10% FBS, 1% Pen/Strep, 1% GlutaMax in Dulbecco's modified Eagle's medium) in a 24-well glass bottom plate (Cellvis, P24-1.5-N). After 48h, cells were stained with Hoescht33342 at a final concentration of 2 µg/mL in cell media for 30 min at 37 °C and 5% CO2. The plate was placed on an IN Cell 6000 Analyzer (GE Healthcare) with a heated stage and 50 fields of view were imaged under 4′,6-diamidino-2-phenylindole (DAPI) and TxRed channels at ×60 magnification (Nikon ×60/0.95, Plan Apo, Corr Collar 0.11–0.23, CFI/60 lambda). Images were exported as TIFF files for downstream analysis. DNAJB8$_{WT}$-mRuby3, all DNAJB8-mRuby3 mutants, mRuby3 alone, and HEK293T cells were plated and imaged in duplicates. Total cell counting, total mRuby3 expression, and total puncta counting was done using the CellProfiler v4.2.1 software.

Recombinant DNAJB8 expression and purification

DNAJB8$_{WT}$ and all DNAJB8 mutants were cloned into pET-29b(+) vectors and prepared by Twist Bioscience. DNAJB8-containing plasmids were transformed into *E.coli* BL21(DE3) competent cells, streaked, plated, and inoculated into 1L 2xLB , 0.05 mg/mL kanamycin and incubated at 37°C shaking at 220 rpm. Once OD$_{600}$=0.6-0.8 AU, expression was induced upon adding 1mL of 1M IPTG to the 1L culture, and incubation

continued for 4 hours at 37°C shaking at 220 rpm. Cells were harvested by spinning down the culture at 4,000xg for 20 min. The media was discarded and the cells were resuspended in 50mL lysis buffer (8M guanidinium HCl, 50mM HEPES, 20mM imidazole, 1mM DTT, pH 7.5). The cells were sonicated at 30% power, 5x pulse, for 10 min using an Omni Sonic Ruptor 4000 (Omni International). The lysed cells were pelleted at 10,000xg for 30 min and supernatant was retained. The lysate was mixed with 2mL HisPur™ Ni-NTA Resin (Thermo Scientific) for 1 hour before being loaded onto a gravity column. The column was washed with an additional 50mL of lysis buffer followed by 50mL of 50mM HEPES, 20mM imidazole, 1mM DTT, pH 7.5. The protein was eluted with 30mL elution buffer (50mM HEPES, 500mM imidazole, 1mM DTT, pH 7.5) and collected in 2mL fractions. High purity DNAJB8-containing fractions were pooled and loaded into 3.5 kDa Biotech CE Dialysis Tubing (Spectrum Labs) to be dialyzed overnight in 50mM ammonium formate at 4°C to minimize assembly formation while removing the imidazole. DNAJB8 was then lyophilized and stored at -80°C for future use.

Dynamic light scattering

All DNAJB8 samples were prepared at 1.2 mg/mL in 1× PBS, 1 mM DTT pH 7.4. All protein samples were filtered through a 0.22-µm cellulose acetate sterile filter and loaded in triplicate onto a 384-well clear flat-bottom plate. The plate was loaded into a Wyatt DynaPro Plate Reader III and set to run continuously at room temperature at a scanning rate of 1 scan/15 min, with 1 scan composed of ten acquisitions. The data were analyzed using the Wyatt Dynamics software version 7.8.2.18. Light-scattering results were filtered by the sum of squares (SOS) < 20 to eliminate statistical outlier acquisitions within each scan. R_h of observed particles for two time points (0 and 30 h) was reported as histograms as a function of mass% for DNAJB8$_{WT}$ and all DNAJB8 mutants. DNAJB8 samples were prepared at 10µM and tau M_s was prepared at 0.1µM for M_s assembly experiments. Data for M_s assembly experiments were plotted as average R_h of total particles over time. Data was not collected for some time points of the M_s experiment due to high SOS values and heterogeneity. It is expected during the

aggregation phase of M_s that assembling fibrils exist at varying sizes until the population reaches equilibrium.

Thioflavin T fluorescence aggregation assays

For all endpoint assays, DNAJB8$_{WT}$ and all DNAJB8 mutants were resuspended in 1xPBS, 1 mM DTT pH 7.4 to a final concentration of 50µM and incubated at room temperature for 30h. All peptides were analyzed at a concentration of 100µM. All samples were mixed with 25µM ThT and plated into a 384-well clear flat bottom plate in triplicate as technical replicates. ThT fluorescence was measured using a Tecan M1000 plate reader at 446 nm Ex (5 nm bandwidth), 482 nm Em (5 nm bandwidth). For peptide time course experiments, all peptides were suspended in a 1:1 mixture (v/v) of TFA (Pierce) and incubated at room temperature (RT) for 1h to monomerize all peptides. In a chemical fume hood, the peptide solution was dried under a stream of nitrogen gas, and then immediately placed under vacuum to remove any residual volatile solvents. The peptide residue was resuspended in 1xPBS, 1mM DTT and the pH restored to pH 7 using NaOH with a final concentration of 100µM. All peptides were mixed with 25µM ThT and plated into a 384-well clear flat bottom plate in triplicate as technical replicates. ThT fluorescence was measured using a Tecan M1000 plate reader at 446 nm Ex (5 nm bandwidth), 482 nm Em (5 nm bandwidth). For the tau aggregation experiment, DNAJB8 was resuspended in 1xPBS, 1 mM DTT pH 7.4 to a final concentration of 10µM, tau M_s was diluted in1xPBS, 1 mM DTT pH 7.4 to a final concentration of 0.05µM. tauRD was diluted in 1xPBS, 1 mM DTT pH 7.4 to a final concentration of 17µM followed by boiling at 100 °C for 5 min before being diluted further in 1xPBS, 1 mM DTT pH 7.4 to a final concentration of 5µM. 25µM ThT was mixed with all samples before the addition of M_s to all M_s-containing reactions. All reactions were plated into a 384-well clear flat bottom plate in triplicate as technical replicates. ThT fluorescence was measured using a Tecan M1000 plate reader at 446 nm Ex (5 nm bandwidth), 482 nm Em (5 nm bandwidth).

Transmission electron microscopy

Peptide samples were resuspended in 1xPBS, 1 mM DTT pH 7.4 to a final concentration of 100µM. 5 µL of sample was loaded onto a glow discharged Formvar-coated 300-mesh copper grid for 30 s and was blotted by filter paper followed by washing the grid with 5 µL of ddH2O. After another 30 seconds, 2% uranyl acetate was loaded on the grid and blotted again. The grid was dried for 1 min and loaded into a FEI Tecnai G2 Spirit Biotwin TEM. All images were captured using a Gatan 2K × 2K multiport readout post-column CCD at the UT Southwestern EM Core Facility.

AFSSFN crystallization and structure determination

AFSSFN synthetic peptide (Genscript) was dissolved to 30 mg/mL in water and crystallized by hanging drop at a 1:2 ratio with 20% PEG 8000, 100 mM HEPES pH 7.5. X-ray diffraction data was collected at APS Beamline 24-ID-C. Diffraction data were indexed and scaled using XDS and XSCALE[71]. The SHELX macromolecular structure determination suite was used for phasing the measured intensities[72]. Model-building and manual real-space refinement was performed in COOT[73]. Automated reciprocal-space and real-space refinement was performed using Refmac and Phenix[74-75]. A summary of data collection and refinement statistics is given in Supplementary Table 2.

Tau Expression and Purification and Production of Seeding Monomer.

Full-length recombinant tau was cloned into a pET28b vector as described previously[53]. This vector was transformed into BL21(DE3) competent cells, and cultured at 37°C shaking at 220 rpm until OD_{600}=1.4 AU. Expression was induced in each flask by the addition of 1mL 0.5M IPTG and continued incubation for 3.25 hours at 37°C shaking at 220 rpm. The cells were harvested by spinning down the cells at 5,000xg for 12 min and discarding the media. The cell pellets were combined and resuspended in 225mL lysis buffer (50mM Tris, 500mM NaCl, 20mM imidazole, 1mM βME, 1mM PMSF, pH 7.5) and sonicated at 30% power, 5x pulse, for 10 min on ice using an Omni Sonic Ruptor 4000 (Omni International). The lysate was spun down at 15,000xg for 20 min and the pellets were discarded. 6mL of HisPur™ Ni-NTA Resin (Thermo Scientific) was added to the

lysate and incubated on a orbital shaker for 1 hour at 4°C. The lysate was loaded onto a gravity column and washed with 40CV of lysis buffer. The sample was eluted with 10mL (3CV) of elution buffer (50mM Tris, 250mM NaCl, 300mM imidazole, 1mM βME, pH 7.5) into 1mL fractions. Tau-containing fractions were pooled and desalted into 5mL 50mM MES, 50mM NaCl, 1mM βME, pH 6.0 using a PD10 desalting column containing Sephadex 25 resin (Cytiva). The sample was filtered and injected onto a 5x5mL HiTrap SP-HP cation exchange column (Cytiva) over a NaCl gradient of 50-200mM and collecting 2mL fractions for 80mL of elution volume. Tau-containing fractions were pooled and concentrated down to 500μL using a 10kDa MW cutoff Amicon Ultra-15 Centrifugal Filter (EMD Millipore). The sample was filtered and loaded onto a 10/300 Superdex 200 column (Cytiva) running either on 10mM HEPES, 100mM NaCl, 1mM DTT, pH 7.4 (Tau M_i elutions) or 30mM MOPS, 50mM KCl, 5mM $MgCl_2$, 1mM DTT, pH 6.5 (Tau pre-M_s elutions).

Tau that was eluted into pre-M_s elution buffer was prepared in 2 tubes of 500μL at 8μM in pre-M_s elution buffer. One tube was injected onto a 10/300 Superdex 200 column (Cytiva) running 1xPBS, 1mM DTT, pH 7.4 to determine the elution profile for tau M_i. The other tube was mixed with heparin at a 1:10 molar ratio (tau:heparin) and incubated at room temperature for 30 minutes to generate tau M_s. The sample was then filtered and injected onto a 10/300 Superdex 200 column (Cytiva) running 1xPBS, 1mM DTT, pH 7.4, where fractions containing tau M_s were isolated from remaining tau M_i and oligomers. Samples were immediately flash frozen in liquid nitrogen and stored at -80°C for future experiments.

Microscale thermophoresis

All MST experiments were performed on a Nanotemper Monolith NT.115 in the Molecular Biophysics Core at UTSW and analyzed with a standard protocol[76]. All binding measurements were done as technical triplicates. 1μM tau M_i and M_s was labelled with 4mM Cy5-NHS by incubating in the dark for 30 minutes at room temperature. Excess dye was removed using a Microspin™ G-25 desalting column (Cytiva). Labelled tau was mixed with a 16-step titration of

DNAJB8$_{S3}$ between 0.006-100µM. Data were fit in PALMIST using a 1:1 binding model[76] and analyzed using GUSSI[77].

Crosslinking reagents

DMTMM (Sigma-Aldrich) is commercially available. For all cross-linking experiments, DMTMM (Sigma-Aldrich) was prepared at a 120 mg/mL concentration in 1× PBS pH 7 and stored at -80°C as a stock solution.

Cross-linking mass spectrometry

For all DNAJB8 experiments, DNAJB8$_{WT}$, DNAJB8$_{S1}$, DNAJB8$_{S2}$, DNAJB8$_{S3}$, and DNAJB8$_{S4}$ were all resuspended in 1xPBS, 1mM DTT pH 7.4 buffer to a final concentration of 25µM. For DNAJB8 with M$_s$ experiments, DNAJB8$_{WT}$ and DNAJB8$_{S3}$ were resuspended in 1xPBS, 1mM DTT pH 7.4 buffer to a final concentration of 10µM and M$_s$ was diluted in 1xPBS, 1mM DTT pH 7.4 buffer to a final concentration of 0.1µM. All experiments were performed in triplicate. All samples were incubated at 37 °C while shaking at 350 r.p.m. for 30 min. Final concentrations of 36mM DMTMM (Sigma-Aldrich) were added to the protein samples and incubated at 37 °C with shaking at 350 r.p.m. for 30 min. The reactions were quenched with 100mM ammonium bicarbonate and incubated at 37 °C for 30 min. Samples were lyophilized and resuspended in 8M urea. Samples were reduced with 2.5mM tris(2-carboxyethyl)phosphine (TCEP) incubated at 37 °C for 30 min, followed by alkylation with 5mM iodoacetimide for 30 min in the dark. Samples were diluted to 1M urea using a stock of 50mM ammonium bicarbonate and trypsin (Promega) was added at a 1:50 enzyme-tosubstrate ratio and incubated overnight at 37 °C while shaking at 600 r.p.m. Two percent (v/v) formic acid was added to acidify the samples following overnight digestion. All samples were run on reverse-phase Sep-Pak tC18 cartridges (Waters) eluted in 50% acetonitrile with 0.1% formic acid. Ten microliters of the purified peptide fractions were injected for liquid Chromatography with tandem mass spectrometry analysis on an Eksigent 1D-NanoLC-Ultra HPLC system coupled to a Thermo Orbitrap Fusion Tribrid System. The MS was

operated in data-dependent mode by selecting the five most abundant precursor ions (m/z 350–1600, charge state 3+ and above) from a preview scan and subjecting them to collision-induced dissociation (normalized collision energy = 35%, 30 ms activation). Fragment ions were detected at low resolution in the linear ion trap. Dynamic exclusion was enabled (repeat count 1, exclusion duration 30 s).

Analysis of MS results

All MS experiments were carried out on an Orbitrap Fusion Lumos Tribrid instrument available through the UTSW proteomics core facility. Each Thermo.raw file was converted to .mzXML format for analysis using an in-house installation of xQuest. Score thresholds were set through xProphet, which uses a target/decoy model. The search parameters were set as follows: maximum number of missed cleavages = 2, peptide length = 5–50 residues, fixed modifications carbamidomethyl-Cys (mass shift = 57.02146 Da), mass shift of cross-linker = −18.010595 Da, no monolink mass specified, MS1 tolerance = 15 p.p.m., and MS2 tolerance = 0.2 Da for common ions and 0.3 Da for cross-link ions; search in enumeration mode.

Cell biological analysis of tau biosensor lines expressing DNAJB8-mRuby3

Original FM5 plasmids used for generating the tauRD(P301S) cells were previously described[23, 56], and modified for these experiments as previously described[57]. HEK293T cells were co-transfected with lentivirus encoding tauRD(P301S)-mClover3/mCerulean3 and sorted as previously described[57] to generate stable cell lines. All DNAJB8-containing plasmids for cell experiments were prepared by Twist Bioscience and constructed as described (see Cell biological analysis of DNAJB8-mRuby3 lines). tauRD(P301S) biosensor cells were plated at 10,000 cells/well in media (10% FBS, 1% Pen/Strep, 1% GlutaMax in Dulbecco's modified Eagle's medium) in a 96-well glass bottom plate (Cellvis, P96-1.5-N). 75ng DNAJB8-mRuby3 plasmids were separately co-transfected into the cells using lipofectamine 2000 (Invitrogen), with each DNAJB8 construct plated in 6 replicate wells. At either 24 or 72 hours after DNAJB8-mRuby3

transfection, 20nM sonicated full-length tau fibrils were transfected into the cells using lipofectamine 2000 (Invitrogen). Tau fibrils were transfected into 3/6 wells per condition, resulting in triplicate wells for each DNAJB8 construct with and without tau seeding. After 24 hours, cells were stained with DRAQ5 at a final concentration of 5μM in cell media for 15 min at 37 °C and 5% CO2. The plate was loaded into an ImageXpress Confocal HT.ai (Molecular Dynamics) with a heated stage and 64 fields of view were taken using far red, FITC, CFP, and TxRed channels at 20x magnification (Nikon 20x/0.80, Plan Apo λD, WD 0.8, CS 0.17, MRD70270). Images were acquired in the MetaXpress software (Molecular Dynamics) and exported as TIFF files for downstream analysis. Total cell counting, total mClover3 expression, total mRuby3 expression, and total DNAJB8 puncta counting was done using the CellProfiler v4.2.1 software.

Flow cytometry analysis of tau biosensor lines

The LSRFortessa SORP (BD Biosciences) was used to perform FRET flow cytometry. We used a gating strategy to isolate the cell population that expressed the three fluorescent proteins: mCerulean, mClover and mRuby (Fig 5a). First, from the single cells we isolated the cells that were mClover and mCerulean positive, and then sub-gating for mRuby positive cells. The gating strategy for this selection is highlighted in Supplementary Fig 5c. Subsequently, within this population, FRET was calculated as previously described[56, 78]. FRET signal is defined as the percentage of FRET-positive cells in all analyses. For each experiment, 10,000 cells per replicate were analyzed and each condition was analyzed as biological triplicates. Data analysis was performed using FlowJo v10 software (Treestar).

Q119 cell culture and transient transfection

HEK293 were cultured at 37° C and 5% CO2 in DMEM (Gibco) supplemented with 10% FBS (Fetal bovine serum, Bodinco) and 100 U/ml penicillin/streptomycin (Invitrogen). For transient transfections, 5×10^5 cells were plated on 35 mm cell culture dishes (Nunc) one the day before. Cells were transfected with 2 ug of total plasmid DNA (0.2 μg

pQ119-GFP, 1.8 μg pDNAJB8 variant/empty vector) using polyethyleneimine (PEI) in 1:3 ratio.

Protein extraction for FTA and western blot

24 h after transfection, cells were washed twice with cold phosphate-buffer saline (PBS), scraped in 230 μl lysis buffer (50 mM TRIS-HCl pH 7.4, 100 mM NaCl, 1 mM $MgCl_2$, 0.5% SDS, EDTA free complete protease inhibitors cocktail (Roche), and 50 units/ml Denarase (c-LEcta)), and left on ice for 30 minutes with occasional vortexing. After incubation, SDS concentration was increased to 2%. Protein concentrations were measured with DC protein assay (Bio-Rad), equalized using dilution buffer (50 mM TRIS-HCl pH 7.4, 100 mM NaCl, 1 mM $MgCl_2$, 2% SDS, 50 mM DTT), boiled for 5 minutes, and stores at -20° C.

Filter trap assay

For filter trap assay (FTA), prepared samples were diluted 5-fold in dilution buffer. Then, both original (1x) and dilute (0.4x) samples were applied onto a 0.2 μm pore Cellulose acetate membrane, prewashed with FTA buffer (10 mM TRIS-HCl pH 8.0, 150 mM NaCl, 0.1% SDS). The membrane was washed under mild suction three times with FTA buffer, blocked in 10% non-fat milk, and blotted with anti-GFP/YFP antibody (mouse, Clontech) at 1:5000 dilution overnight at 4° C on a rocking platform. After, the membrane was incubated with an HPR-conjugated secondary antibody and visualized with enhanced chemiluminescence using ChemiDoc Imaging System (Bio-Rad). Signal intensities were measured in ImageLab Software, and the results of two dilutions (where possible) were averaged. Values were normalized to the control sample, analyzed with GraphPad Prism, and plotted in a graph. The resulting graph represents an average of three separate experiments.

Western Blot analysis

For WB, samples aliquots were mixed with 4x Sample buffer (50 mM Tris-HCl pH 6.7, 2% SDS, 10% glycerol, 12.5 mM EDTA, 0.02% Bromophenol blue). Samples were loaded on 10% SDS-PAGE gel and ran at 90 V. Proteins were transferred to nitrocellulose membrane (Schleicher and Schuell, PerkinElmer, Waltham, MA, USA), blocked in 10% non-fat milk, and blotted with primary anti-GAPDH antibody (Sigma) diluted 1:5000, or with anti-DNAJB8 antibody (monoclonal, mouse, clone #EMR-DNAJB8.214-8) overnight at 4° C on a rocking platform. The anti-DNAJB8 antibody was prepared as previously described[67, 79]. After, the membrane was incubated with an HPR-conjugated secondary antibody and visualized with enhanced chemiluminescence using ChemiDoc Imaging System (Bio-Rad).

ACKNOWLEDGEMENTS

L.A.J is supported by a Effie Marie Cain Scholarship in Medical Research and a grant from the NIH-NIA (R01AG065407). Mass spectrometry experiments were carried out at the UTSW Proteomics core. Transmission electron microscopy was performed at the Electron Microscopy Core Facility at UTSW, which is supported by the National Institutes of Health (NIH) (1S10OD021685-01A1 and 1S10OD020103-01). We would like to thank Dr. Andrew Lemoff from the UTSW Proteomics Core for his valuable insights on troubleshooting procedures in cross-linking mass spectrometry. We want to thank Vaibhav Bommareddy for preparing the plasmids of CMV FM5 tau P301S. We thank all members of the Joachimiak lab for discussions and input on the manuscript.

CONTRIBUTIONS

B.D.R. and L.A.J. initiated the project. B.D.R purified all proteins involved in the study. B.D.R. performed all peptide and protein assembly experiments. B.D.R performed all cell experiments and image analyses. B.D.R and A.O. performed the flow cytometry analysis. D.R.B. and M.S. determined the X-ray structure of the peptide. B.D.R and

M.I.K. performed the gel crosslinking experiments. P.M.W. performed the MST binding experiments. B.D.R performed the XL-MS analysis. E.U. performed polyQ cell experiments. M.I.D and H.H.K. provided guidance on cell experiments. Finally, B.D.R and L.A.J. conceived of and directed the research as well as wrote the manuscript. All authors contributed to the revisions of the manuscript.

DATA AVAILABILITY

The structural datasets for the AFSSFN peptide assembly generated during the current study are available in the Protein Data Bank repository (https://www.rcsb.org/) under PDB id: 8DTS. Analysis of WT and mutant DNAJB8-mRuby3 expression in HEK293T cells and HEK293T cells expressing tauRD are available in Source Data 1. Raw DLS data is available as Source Data 2. XL-MS data are available in Source Data 3. Raw data from fluorescence assays, SASA calculations, FTA, SDS-PAGE, and western blots are available in Source Data 4. All other data generated during the current study are available upon reasonable request.

DECLARATION OF INTERESTS

The authors declare that they have no competing interests.

REFERENCES

1.   Hartl, F. U., Molecular chaperones in cellular protein folding. *Nature* **1996,** *381* (6583), 571-9.
2.   Kampinga, H. H.; Craig, E. A., The HSP70 chaperone machinery: J proteins as drivers of functional specificity. *Nat Rev Mol Cell Biol* **2010,** *11* (8), 579-92.
3.   Ryder, B. D.; Wydorski, P. M.; Hou, Z.; Joachimiak, L. A., Chaperoning shape-shifting tau in disease. *Trends Biochem Sci* **2022,** *47* (4), 301-313.
4.   Fernandez-Fernandez, M. R.; Gragera, M.; Ochoa-Ibarrola, L.; Quintana-Gallardo, L.; Valpuesta, J. M., Hsp70 - a master regulator in protein degradation. *FEBS Lett* **2017,** *591* (17), 2648-2660.
5.   Quintana-Gallardo, L.; Martin-Benito, J.; Marcilla, M.; Espadas, G.; Sabido, E.; Valpuesta, J. M., The cochaperone CHIP marks Hsp70- and Hsp90-bound substrates for degradation through a very flexible mechanism. *Sci Rep* **2019,** *9* (1), 5102.

6.   Abrams, J.; Arhar, T.; Mok, S. A.; Taylor, I. R.; Kampmann, M.; Gestwicki, J. E., Functional genomics screen identifies proteostasis targets that modulate prion protein (PrP) stability. *Cell Stress Chaperones* **2021,** *26* (2), 443-452.

7.   Hohn, A.; Tramutola, A.; Cascella, R., Proteostasis Failure in Neurodegenerative Diseases: Focus on Oxidative Stress. *Oxid Med Cell Longev* **2020,** *2020*, 5497046.

8.   Morawe, T.; Hiebel, C.; Kern, A.; Behl, C., Protein homeostasis, aging and Alzheimer's disease. *Mol Neurobiol* **2012,** *46* (1), 41-54.

9.   Sawaya, M. R.; Sambashivan, S.; Nelson, R.; Ivanova, M. I.; Sievers, S. A.; Apostol, M. I.; Thompson, M. J.; Balbirnie, M.; Wiltzius, J. J.; McFarlane, H. T.; Madsen, A. O.; Riekel, C.; Eisenberg, D., Atomic structures of amyloid cross-beta spines reveal varied steric zippers. *Nature* **2007,** *447* (7143), 453-7.

10.   Goldschmidt, L.; Teng, P. K.; Riek, R.; Eisenberg, D., Identifying the amylome, proteins capable of forming amyloid-like fibrils. *Proc Natl Acad Sci U S A* **2010,** *107* (8), 3487-92.

11.   Rosenzweig, R.; Nillegoda, N. B.; Mayer, M. P.; Bukau, B., The Hsp70 chaperone network. *Nat Rev Mol Cell Biol* **2019,** *20* (11), 665-680.

12.   Glover, J. R.; Lindquist, S., Hsp104, Hsp70, and Hsp40: A Novel Chaperone System that Rescues Previously Aggregated Proteins. *Cell* **1998,** *94*, 73-82.

13.   Qiu, X. B.; Shao, Y. M.; Miao, S.; Wang, L., The diversity of the DnaJ/Hsp40 family, the crucial partners for Hsp70 chaperones. *Cell Mol Life Sci* **2006,** *63* (22), 2560-70.

14.   Nillegoda, N. B.; Kirstein, J.; Szlachcic, A.; Berynskyy, M.; Stank, A.; Stengel, F.; Arnsburg, K.; Gao, X.; Scior, A.; Aebersold, R.; Guilbride, D. L.; Wade, R. C.; Morimoto, R. I.; Mayer, M. P.; Bukau, B., Crucial HSP70 co-chaperone complex unlocks metazoan protein disaggregation. *Nature* **2015,** *524* (7564), 247-51.

15.   Cheetham, M. E.; Caplan, A. J., Structure, function and evolution of DnaJ: conservation and adaptation of chaperone function. *Cell Stress Chaperones* **1998,** *3* (1), 28-36.

16.   Faust, O.; Abayev-Avraham, M.; Wentink, A. S.; Maurer, M.; Nillegoda, N. B.; London, N.; Bukau, B.; Rosenzweig, R., HSP40 proteins use class-specific regulation to drive HSP70 functional diversity. *Nature* **2020,** *587* (7834), 489-494.

17.   Piette, B. L.; Alerasool, N.; Lin, Z. Y.; Lacoste, J.; Lam, M. H. Y.; Qian, W. W.; Tran, S.; Larsen, B.; Campos, E.; Peng, J.; Gingras, A. C.; Taipale, M., Comprehensive interactome profiling of the human Hsp70 network highlights functional differentiation of J domains. *Mol Cell* **2021,** *81* (12), 2549-2565 e8.

18.   Zhang, R.; Malinverni, D.; Cyr, D. M.; Rios, P. L.; Nillegoda, N. B., J-domain protein chaperone circuits in proteostasis and disease. *Trends Cell Biol* **2023,** *33* (1), 30-47.

19.   Tsai, J.; Douglas, M. G., A conserved HPD sequence of the J-domain is necessary for YDJ1 stimulation of Hsp70 ATPase activity at a site distinct from substrate binding. *J Biol Chem* **1996,** *271* (16), 9347-54.

20.   Lackie, R. E.; Maciejewski, A.; Ostapchenko, V. G.; Marques-Lopes, J.; Choy, W. Y.; Duennwald, M. L.; Prado, V. F.; Prado, M. A. M., The Hsp70/Hsp90 Chaperone Machinery in Neurodegenerative Diseases. *Front Neurosci* **2017,** *11*, 254.

21.   Schopf, F. H.; Biebl, M. M.; Buchner, J., The HSP90 chaperone machinery. *Nat Rev Mol Cell Biol* **2017,** *18* (6), 345-360.

22.   Clavaguera, F.; Bolmont, T.; Crowther, R. A.; Abramowski, D.; Frank, S.; Probst, A.; Fraser, G.; Stalder, A. K.; Beibel, M.; Staufenbiel, M.; Jucker, M.; Goedert, M.; Tolnay, M., Transmission and spreading of tauopathy in transgenic mouse brain. *Nat Cell Biol* **2009,** *11* (7), 909-13.

23.   Sanders, D. W.; Kaufman, S. K.; DeVos, S. L.; Sharma, A. M.; Mirbaha, H.; Li, A.; Barker, S. J.; Foley, A. C.; Thorpe, J. R.; Serpell, L. C.; Miller, T. M.; Grinberg, L. T.; Seeley, W. W.; Diamond, M. I., Distinct tau prion strains propagate in cells and mice and define different tauopathies. *Neuron* **2014,** *82* (6), 1271-88.

24.   Hong, M.; Zhukareva, V.; Vogelsberg-Ragaglia, V.; Wszolek, Z.; Reed, L.; Miller, B. I.; Geschwind, D. H.; Bird, T. D.; McKeel, D.; Goate, A.; Morris, J. C.; Wilhelmsen, K. C.; Schellenberg, G. D.; Trojanowski,

J. Q.; Lee, V. M.-Y., Mutation-Specific Functional Impairments in Distinct Tau Isoforms of Hereditary FTDP-17. *Science* **1998,** *282*, 1914–1917.

25. Goedert, M.; Eisenberg, D. S.; Crowther, R. A., Propagation of Tau Aggregates and Neurodegeneration. *Annu Rev Neurosci* **2017,** *40*, 189-210.

26. Kaufman, S. K.; Sanders, D. W.; Thomas, T. L.; Ruchinskas, A. J.; Vaquer-Alicea, J.; Sharma, A. M.; Miller, T. M.; Diamond, M. I., Tau Prion Strains Dictate Patterns of Cell Pathology, Progression Rate, and Regional Vulnerability In Vivo. *Neuron* **2016,** *92* (4), 796-812.

27. Scheres, S. H.; Zhang, W.; Falcon, B.; Goedert, M., Cryo-EM structures of tau filaments. *Curr Opin Struct Biol* **2020,** *64*, 17-25.

28. Fitzpatrick, A. W. P.; Falcon, B.; He, S.; Murzin, A. G.; Murshudov, G.; Garringer, H. J.; Crowther, R. A.; Ghetti, B.; Goedert, M.; Scheres, S. H. W., Cryo-EM structures of tau filaments from Alzheimer's disease. *Nature* **2017,** *547* (7662), 185-190.

29. Irwin, R.; Faust, O.; Petrovic, I.; Wolf, S. G.; Hofmann, H.; Rosenzweig, R., Hsp40s play complementary roles in the prevention of tau amyloid formation. *Elife* **2021**.

30. Mok, S. A.; Condello, C.; Freilich, R.; Gillies, A.; Arhar, T.; Oroz, J.; Kadavath, H.; Julien, O.; Assimon, V. A.; Rauch, J. N.; Dunyak, B. M.; Lee, J.; Tsai, F. T. F.; Wilson, M. R.; Zweckstetter, M.; Dickey, C. A.; Gestwicki, J. E., Mapping interactions with the chaperone network reveals factors that protect against tau aggregation. *Nat Struct Mol Biol* **2018,** *25* (5), 384-393.

31. Abisambra, J. F.; Jinwal, U. K.; Suntharalingam, A.; Arulselvam, K.; Brady, S.; Cockman, M.; Jin, Y.; Zhang, B.; Dickey, C. A., DnaJA1 antagonizes constitutive Hsp70-mediated stabilization of tau. *J Mol Biol* **2012,** *421* (4-5), 653-61.

32. Burré, J.; Sharma, M.; Tsetsenis, T.; Buchman, V.; Etherton, M.; Südhof, T. C., α-Synuclein Promotes SNARE-Complex Assembly in vivo and in vitro. *Science* **2010,** *329* (5999), 1663-1667.

33. Sharma, M.; Burre, J.; Bronk, P.; Zhang, Y.; Xu, W.; Sudhof, T. C., CSPalpha knockout causes neurodegeneration by impairing SNAP-25 function. *EMBO J* **2012,** *31* (4), 829-41.

34. Nachman, E.; Wentink, A. S.; Madiona, K.; Bousset, L.; Katsinelos, T.; Allinson, K.; Kampinga, H.; McEwan, W. A.; Jahn, T. R.; Melki, R.; Mogk, A.; Bukau, B.; Nussbaum-Krammer, C., Disassembly of Tau fibrils by the human Hsp70 disaggregation machinery generates small seeding-competent species. *J Biol Chem* **2020,** *295* (28), 9676-9690.

35. Mirbaha, H.; Chen, D.; Morazova, O. A.; Ruff, K. M.; Sharma, A. M.; Liu, X.; Goodarzi, M.; Pappu, R. V.; Colby, D. W.; Mirzaei, H.; Joachimiak, L. A.; Diamond, M. I., Inert and seed-competent tau monomers suggest structural origins of aggregation. *Elife* **2018,** *7*.

36. Hou, Z.; Chen, D.; Ryder, B. D.; Joachimiak, L. A., Biophysical properties of a tau seed. *Sci Rep* **2021,** *11* (1), 13602.

37. Mirbaha, H.; Chen, D.; Mullapudi, V.; Terpack, S. J.; White, C. L., 3rd; Joachimiak, L. A.; Diamond, M. I., Seed-competent tau monomer initiates pathology in a tauopathy mouse model. *J Biol Chem* **2022,** *298* (8), 102163.

38. Hou, Z.; Wydorski, P. M.; Perez, V. A.; Mendoza-Oliva, A.; Ryder, B. D.; Mirbaha, H.; Kashmer, O.; Joachimiak, L. A., DnaJC7 binds natively folded structural elements in tau to inhibit amyloid formation. *Nat. Comm.* **2021,** *12*.

39. Ryder, B. D.; Matlahov, I.; Bali, S.; Vaquer-Alicea, J.; van der Wel, P. C. A.; Joachimiak, L. A., Regulatory inter-domain interactions influence Hsp70 recruitment to the DnaJB8 chaperone. *Nat Commun* **2021,** *12* (1), 946.

40. Månsson, C.; Kakkar, V.; Monsellier, E.; Sourigues, Y.; Härmark, J.; Kampinga, H. H.; Melki, R.; Emanuelsson, C., DNAJB6 is a peptide-binding chaperone which can suppress amyloid fibrillation of polyglutamine peptides at substoichiometric molar ratios. *Cell Stress and Chaperones* **2014,** *19*, 227-239.

41. Gillis, J.; Schipper-Krom, S.; Juenemann, K.; Gruber, A.; Coolen, S.; van den Nieuwendijk, R.; van Veen, H.; Overkleeft, H.; Goedhart, J.; Kampinga, H. H.; Reits, E. A., The DNAJB6 and DNAJB8 protein

chaperones prevent intracellular aggregation of polyglutamine peptides. *J Biol Chem* **2013,** *288* (24), 17225-37.

42. Hageman, J.; Rujano, M. A.; van Waarde, M. A.; Kakkar, V.; Dirks, R. P.; Govorukhina, N.; Oosterveld-Hut, H. M.; Lubsen, N. H.; Kampinga, H. H., A DNAJB chaperone subfamily with HDAC-dependent activities suppresses toxic protein aggregation. *Mol Cell* **2010,** *37* (3), 355-69.

43. McMahon, S.; Bergink, S.; Kampinga, H. H.; Ecroyd, H., DNAJB chaperones suppress destabilised protein aggregation via a region distinct from that used to inhibit amyloidogenesis. *J Cell Sci* **2021,** *134* (7).

44. Kakkar, V.; Mansson, C.; de Mattos, E. P.; Bergink, S.; van der Zwaag, M.; van Waarde, M. A.; Kloosterhuis, N. J.; Melki, R.; van Cruchten, R. T.; Al-Karadaghi, S.; Arosio, P.; Dobson, C. M.; Knowles, T. P.; Bates, G. P.; van Deursen, J. M.; Linse, S.; van de Sluis, B.; Emanuelsson, C.; Kampinga, H. H., The S/T-Rich Motif in the DNAJB6 Chaperone Delays Polyglutamine Aggregation and the Onset of Disease in a Mouse Model. *Mol Cell* **2016**.

45. Aprile, F. A.; Kallstig, E.; Limorenko, G.; Vendruscolo, M.; Ron, D.; Hansen, C., The molecular chaperones DNAJB6 and Hsp70 cooperate to suppress alpha-synuclein aggregation. *Sci Rep* **2017,** *7* (1), 9039.

46. Arkan, S.; Ljungberg, M.; Kirik, D.; Hansen, C., DNAJB6 suppresses alpha-synuclein induced pathology in an animal model of Parkinson's disease. *Neurobiol Dis* **2021,** *158*, 105477.

47. Osterlund, N.; Lundqvist, M.; Ilag, L. L.; Graslund, A.; Emanuelsson, C., Amyloid-beta oligomers are captured by the DNAJB6 chaperone: Direct detection of interactions that can prevent primary nucleation. *J Biol Chem* **2020,** *295* (24), 8135-8144.

48. Udan-Johns, M.; Bengoechea, R.; Bell, S.; Shao, J.; Diamond, M. I.; True, H. L.; Weihl, C. C.; Baloh, R. H., Prion-like nuclear aggregation of TDP-43 during heat shock is regulated by HSP40/70 chaperones. *Hum Mol Genet* **2014,** *23* (1), 157-70.

49. Hughes, M. P.; Sawaya, M. R.; Boyer, D. R.; Goldschmidt, L.; Rodriguez, J. A.; Cascio, D.; Chong, L.; Gonen, T.; Eisenberg, D. S., Atomic structures of low-complexity protein segments reveal kinked b sheets that assemble networks. *Science* **2018,** *359*, 698-701.

50. Mullapudi, V.; Vaquer-Alicea, J.; Bommareddy, V.; Vega, A. R.; Ryder, B. D.; White, C. L.; Diamond, M. I.; Joachimiak, L. A., Network of hotspot interactions cluster tau amyloid folds. *Biorxiv* **2022**.

51. Karamanos, T. K.; Tugarinov, V.; Clore, G. M., Unraveling the structure and dynamics of the human DNAJB6b chaperone by NMR reveals insights into Hsp40-mediated proteostasis. *Proc Natl Acad Sci U S A* **2019,** *116* (43), 21529-21538.

52. Karamanos, T. K.; Tugarinov, V.; Clore, G. M., An S/T motif controls reversible oligomerization of the Hsp40 chaperone DNAJB6b through subtle reorganization of a beta sheet backbone. *Proc Natl Acad Sci U S A* **2020,** *117* (48), 30441-30450.

53. Chen, D.; Drombosky, K. W.; Hou, Z.; Sari, L.; Kashmer, O. M.; Ryder, B. D.; Perez, V. A.; Woodard, D. R.; Lin, M. M.; Diamond, M. I.; Joachimiak, L. A., Tau local structure shields an amyloid-forming motif and controls aggregation propensity. *Nat Commun* **2019,** *10* (1), 2493.

54. Leitner, A.; Faini, M.; Stengel, F.; Aebersold, R., Crosslinking and Mass Spectrometry: An Integrated Technology to Understand the Structure and Function of Molecular Machines. *Trends Biochem Sci* **2016,** *41* (1), 20-32.

55. Walzthoeni, T.; Joachimiak, L. A.; Rosenberger, G.; Rost, H. L.; Malmstrom, L.; Leitner, A.; Frydman, J.; Aebersold, R., xTract: software for characterizing conformational changes of protein complexes by quantitative cross-linking mass spectrometry. *Nat Methods* **2015,** *12* (12), 1185-90.

56. Holmes, B. B.; Furman, J. L.; Mahan, T. E.; Yamasaki, T. R.; Mirbaha, H.; Eades, W. C.; Belaygorod, L.; Cairns, N. J.; Holtzman, D. M.; Diamond, M. I., Proteopathic tau seeding predicts tauopathy in vivo. *Proc Natl Acad Sci U S A* **2014,** *111* (41), E4376-85.

57.   Hitt, B. D.; Vaquer-Alicea, J.; Manon, V. A.; Beaver, J. D.; Kashmer, O. M.; Garcia, J. N.; Diamond, M. I., Ultrasensitive tau biosensor cells detect no seeding in Alzheimer's disease CSF. *Acta Neuropathol Commun* **2021,** *9* (1), 99.

58.   Thiruvalluvan, A.; de Mattos, E. P.; Brunsting, J. F.; Bakels, R.; Surlidaki, D.; Barazzuol, L.; Conforti, P.; Fatima, A.; Koyuncu, S.; Cattaneo, E.; Vilchez, D.; Bergink, S.; Boddeke, E. H. W. G.; Copray, S.; Kampinga, H. H., DNAJB6, a Key Factor in Neuronal Sensitivity to Amyloidogenesis. *Mol Cell* **2020,** *78*, 346-358.

59.   Pak, C. W.; Kosno, M.; Holehouse, A. S.; Padrick, S. B.; Mittal, A.; Ali, R.; Yunus, A. A.; Liu, D. R.; Pappu, R. V.; Rosen, M. K., Sequence Determinants of Intracellular Phase Separation by Complex Coacervation of a Disordered Protein. *Mol Cell* **2016,** *63* (1), 72-85.

60.   Krainer, G.; Welsh, T. J.; Joseph, J. A.; Espinosa, J. R.; Wittmann, S.; de Csillery, E.; Sridhar, A.; Toprakcioglu, Z.; Gudiskyte, G.; Czekalska, M. A.; Arter, W. E.; Guillen-Boixet, J.; Franzmann, T. M.; Qamar, S.; George-Hyslop, P. S.; Hyman, A. A.; Collepardo-Guevara, R.; Alberti, S.; Knowles, T. P. J., Reentrant liquid condensate phase of proteins is stabilized by hydrophobic and non-ionic interactions. *Nat Commun* **2021,** *12* (1), 1085.

61.   Dignon, G. L.; Best, R. B.; Mittal, J., Biomolecular Phase Separation: From Molecular Driving Forces to Macroscopic Properties. *Annu Rev Phys Chem* **2020,** *71*, 53-75.

62.   Vernon, R. M.; Chong, P. A.; Tsang, B.; Kim, T. H.; Bah, A.; Farber, P.; Lin, H.; Forman-Kay, J. D., Pi-Pi contacts are an overlooked protein feature relevant to phase separation. *Elife* **2018,** *7*.

63.   McGaughey, G. B.; Gagne, M.; Rappe, A. K., pi-Stacking interactions. Alive and well in proteins. *J Biol Chem* **1998,** *273* (25), 15458-63.

64.   Zhou, X.; Sumrow, L.; Tashiro, K.; Sutherland, L.; Liu, D.; Qin, T.; Kato, M.; Liszczak, G.; McKnight, S. L., Mutations linked to neurological disease enhance self-association of low-complexity protein sequences. *Science* **2022,** *377* (6601).

65.   Zarouchlioti, C.; Parfitt, D. A.; Li, W.; Gittings, L. M.; Cheetham, M. E., DNAJ Proteins in neurodegeneration: essential and protective factors. *Phil. Trans. R. Soc. B* **2017,** *373*.

66.   Johnson, J. L., Evolution and function of diverse Hsp90 homologs and cochaperone proteins. *Biochim Biophys Acta* **2012,** *1823* (3), 607-13.

67.   Nishizawa, S.; Hirohashi, Y.; Torigoe, T.; Takahashi, A.; Tamura, Y.; Mori, T.; Kanaseki, T.; Kamiguchi, K.; Asanuma, H.; Morita, R.; Sokolovskaya, A.; Matsuzaki, J.; Yamada, R.; Fujii, R.; Kampinga, H. H.; Kondo, T.; Hasegawa, T.; Hara, I.; Sato, N., HSP DNAJB8 controls tumor-initiating ability in renal cancer stem-like cells. *Cancer Res* **2012,** *72* (11), 2844-54.

68.   Sarparanta, J.; Jonson, P. H.; Golzio, C.; Sandell, S.; Luque, H.; Screen, M.; McDonald, K.; Stajich, J. M.; Mahjneh, I.; Vihola, A.; Raheem, O.; Penttila, S.; Lehtinen, S.; Huovinen, S.; Palmio, J.; Tasca, G.; Ricci, E.; Hackman, P.; Hauser, M.; Katsanis, N.; Udd, B., Mutations affecting the cytoplasmic functions of the co-chaperone DNAJB6 cause limb-girdle muscular dystrophy. *Nat Genet* **2012,** *44* (4), 450-5, S1-2.

69.   Kuiper, E. F. E.; Gallardo, P.; Bergsma, T.; Mari, M.; Kolbe Musskopf, M.; Kuipers, J.; Giepmans, B. N. G.; Steen, A.; Kampinga, H. H.; Veenhoff, L. M.; Bergink, S., The chaperone DNAJB6 surveils FG-nucleoporins and is required for interphase nuclear pore complex biogenesis. *Nat Cell Biol* **2022,** *24* (11), 1584-1594.

70.   Prophet, S. M.; Rampello, A. J.; Niescier, R. F.; Gentile, J. E.; Mallik, S.; Koleske, A. J.; Schlieker, C., Atypical nuclear envelope condensates linked to neurological disorders reveal nucleoporin-directed chaperone activities. *Nat Cell Biol* **2022,** *24* (11), 1630-1641.

71.   Kabsch, W., Xds. *Acta Crystallogr D Biol Crystallogr* **2010,** *66* (Pt 2), 125-32.

72.   Sheldrick, G. M., A short history of SHELX. *Acta Crystallogr A* **2008,** *64* (Pt 1), 112-22.

73.   McCoy, A. J.; Grosse-Kunstleve, R. W.; Adams, P. D.; Winn, M. D.; Storoni, L. C.; Read, R. J., Phaser crystallographic software. *J Appl Crystallogr* **2007,** *40* (Pt 4), 658-674.

74.   Murshudov, G.; Vagin, A. A.; Dodson, E. J., Refinement of macromolecular structures by the maximum-likelihood method *Acta Crystallogr D Biol Crystallogr.* **1997,** *53*, 240-255.

75.   Afonine, P. V.; Grosse-Kunstleve, R. W.; Echols, N.; Headd, J. J.; Moriarty, N. W.; Mustyakimov, M.; Terwilliger, T. C.; Urzhumtsev, A.; Zwart, P. H.; Adams, P. D., Towards automated crystallographic structure refinement with phenix.refine. *Acta Crystallogr D Biol Crystallogr* **2012,** *68* (Pt 4), 352-67.

76.   Scheuermann, T. H.; Padrick, S. B.; Gardner, K. H.; Brautigam, C. A., On the acquisition and analysis of microscale thermophoresis data. *Anal Biochem* **2016,** *496*, 79-93.

77.   Brautigam, C. A., Calculations and Publication-Quality Illustrations for Analytical Ultracentrifugation Data. *Methods Enzymol* **2015,** *562*, 109-33.

78.   Furman, J. L.; Holmes, B. B.; Diamond, M. I., Sensitive Detection of Proteopathic Seeding Activity with FRET Flow Cytometry. *J Vis Exp* **2015,** (106), e53205.

79.   Inoda, S.; Hirohashi, Y.; Torigoe, T.; Nakatsugawa, M.; Kiriyama, K.; Nakazawa, E.; Harada, K.; Takasu, H.; Tamura, Y.; Kamiguchi, K.; Asanuma, H.; Tsuruma, T.; Terui, T.; Ishitani, K.; Ohmura, T.; Wang, Q.; Greene, M. I.; Hasegawa, T.; Hirata, K.; Sato, N., Cep55/c10orf3, a Tumor Antigen Derived From a Centrosome Residing Protein in Breast Carcinoma. *J. Immunother.* **2009,** *32* (5), 474-485.

Figures and Legends

Figure 1. Phenylalanine residues in the S/T-rich domain drive self-assembly. (A) Domain map of DNAJB8 is colored according to domain annotation: JD (red), G/F-rich (blue), S/T-rich (cyan), and CTD (green). A histogram below the G/F-rich and S/T-rich domains shows the distribution of phenylalanine amino acids. Each position is given an arbitrary value to reflect local phenylalanine densities. Based on phenylalanie distributions and density characterizations, we arbitrarily divided the disordered domains into 4 segments (shown as S1, S2, S3, S4). We designed Phe→Ser substitution mutants (termed DNAJB8$_{S1}$, DNAJB8$_{S2}$, DNAJB8$_{S3}$, and DNAJB8$_{S4}$) for each segment with domain maps indicated which Phe residues (black bars) were mutated to Ser (red bars). **(B)** Domain maps for DNAJB8 used in cell experiments (DNAJB8-mRuby3) and in vitro experiments (DNAJB8). mRuby3 is colored pale red. **(C)** DLS histogram reporting R_h distributions of DNAJB8$_{WT}$ (blue), DNAJB8$_{S1}$ (green), DNAJB8$_{S2}$ (gold), DNAJB8$_{S3}$ (cyan), and DNAJB8$_{S4}$ (grey) at time points t=0hrs (solid lines) and t=30hrs (dashed lines). While mutations to all mutants resulted in slower assembly, DNAJB8$_{S1}$ and DNAJB8$_{S2}$ showed size distributions consistent with DNAJB8$_{WT}$, while DNAJB8$_{S3}$ and DNAJB8$_{S4}$ were more compact, consistent with stable monomers with buried hydrophobic residues. Shown values were calculated with SOS < 20. **(D)** Bar plot showing Thioflavin T endpoint fluorescence after 30h incubation at room temperature. Average fluorescence signal across triplicate technical replicates was reported for DNAJB8$_{WT}$ (blue), DNAJB8$_{S1}$ (green), DNAJB8$_{S2}$ (gold), DNAJB8$_{S3}$ (cyan), and DNAJB8$_{S4}$ (grey). DNAJB8$_{S1}$ fluorescence signal was comparable to self-assembly prone DNAJB8$_{WT}$, while DNAJB8$_{S2}$ and DNAJB8$_{S4}$ showed a ~2-fold reduction in fluorescence signal. DNAJB8$_{S3}$ was the only protein to remain ThT negative. One-way ANOVA was used to report significance where F= 101.2 and P-value<0.0001. For multiple comparisons to DNAJB8$_{WT}$: *** $p<0.001$, and **** $p<0.0001$. **(E)** Representative images from cells transfected with mRuby3 (top right), DNAJB8 (bottom left), and DNAJB8$_{S3}$ (bottom right). DAPI is colored blue and mRuby3 signal is shown in red. DNAJB8 is expressed in the cytoplasm and forms cytoplasmic and juxtanuclear puncta. DNAJB8$_{S3}$ shows few instances of puncta, but is able to be expressed in both the cytoplasm and nucleus. **(f)** Histogram of puncta frequency across cell lines expressing mRuby3 (red), DNAJB8$_{WT}$-mRuby3 (blue), DNAJB8$_{S1}$-mRuby3 (green), DNAJB8$_{S2}$-

mRuby3 (gold), DNAJB8$_{S3}$-mRuby3 (cyan), and DNAJB8$_{S4}$-mRuby3 (grey). Phenylalanine to serine mutations within the S/T-rich domain in DNAJB8$_{S3}$ and DNAJB8$_{S4}$ resulted in a decrease of puncta. All values were calculated from CellProfiler v.4.2.1. One-way ANOVA was used to report significance where F=21.94 and P-value=0.0009. For multiple comparisons to DNAJB8$_{WT}$, ** p<0.01.

Figure 2. A steric zipper motif drives local S/T-rich domain self-assembly. (**A**) Domain map of DNAJB8 using the same color scheme as in Figure 1.1a. All peptides used in the peptide library are shown underneath their respective locations within the DNAJB8 sequence. (**B**) Thioflavin T time course experiment and TEM image of B8_5 fragment (blue). Out of all peptides used in the screen, only the B8_5 fragment showed

evidence of assembly through increasing ThT fluorescence. Likewise, the B8_5 fragment was the only peptide to show visual evidence of large assemblies. Sale bar: 200nm. (**C**) TEM image of oligomeric B8_5 peptide assemblies after 10 hrs of incubation at room temperature. Scale bar: 200 nm. (**D**) 0.75Å microED structure of the ^{147}AFSSFN^{152} class 6 steric zipper motif identified within the B8_5 peptide. Four peptides of ^{147}AFSSFN^{152} are shown within the base subunit of the assembly (left). The structure is stabilized through hydrophobic packing of F148 and F151 (center), and anti-parallel β-stranded backbones (right). Crystallography statistics shown in Supplementary Table 2. (**E**) SASA analysis of F148 and F151 in a peptide dimer based on the ^{147}AFSSFN^{152} structure in panel d. A graphical representation shows the peptide arrangement used in the simulation. Higher SASA values correspond to greater solvent exposure (blue), while lower values correspond to less solvent exposure (red). (**F**) SASA analysis of F148 and F151 in a horizontally aligned hexameric assembly based on the ^{147}AFSSFN^{152} structure in panel d. A graphical representation shows the peptide arrangement used in the simulation. Higher SASA values correspond to greater solvent exposure (blue), while lower values correspond to less solvent exposure (red). (**G**) SASA analysis of F148 and F151 in a vertically aligned hexameric assembly based on the ^{147}AFSSFN^{152} structure in panel d. A graphical representation shows the peptide arrangement used in the simulation. Higher SASA values correspond to greater solvent exposure (blue), while lower values correspond to less solvent exposure (red). (**H**) Thioflavin T time course of B8_5 WT (blue) derived peptides where mutations are introduced to disrupt assembly based on Phe-Phe interactions: F148S (salmon), F151S (gold), F148S F151S (teal), or the β-strand backbone: S149P (orange), S149P S150P (black).

Figure 3. Phenylalanine at position 151 is the primary driver of FL DNAJB8 self-assembly. (**A**) Domain maps of DNAJB8WT, DNAJB8F148S, DNAJB8F151S, and DNAJB8F148S_F151S using the same color scheme as in Figure 1a. Red hashes are shown to indicate the relative positions of the mutations. (**B**) Bar plot showing Thioflavin T endpoint fluorescence after 30 h incubation at room temperature. Average fluorescence signal across triplicate technical replicates was reported for DNAJB8WT (blue), DNAJB8S3 (cyan), DNAJB8F148S (lime), DNAJB8F151S (purple) and DNAJB8F148S_F151S (teal). Data for DNAJB8WT and DNAJB8S3 (gray box) was previously shown in Figure 1D. While F148S in the context of FL DNAJB8 did not show any loss of

self-assembly by ThT fluorescence compared to DNAJB8$_{WT}$, both DNAJB8$_{F148S}$ and DNAJB8$_{F148S\_F151S}$ showed considerably less ThT fluorescence, akin to decreased β-stranded self-assembly. Statistics calculated by one-way ANOVA. ** $p<0.01$, **** $p<0.0001$. (**C**) DLS histogram of R_h distributions across DNAJB8$_{WT}$ (blue), DNAJB8$_{S3}$ (cyan), DNAJB8$_{F148S}$ (lime), DNAJB8$_{F151S}$ (purple) and DNAJB8$_{F148S\_F151S}$ (teal) at two time points: 0 h (solid lines) and 30 h (dashed lines). DNAJB8$_{WT}$ and DNAJB8$_{S3}$ are the same data as shown in Figure 1C (grey box). The size distribution of F148S matched the size of DNAJB8$_{WT}$ at t=0 h, with DNAJB8$_{F148S}$ showing evidence of self-assembly after t=30h. DNAJB8$_{F151S}$ and DNAJB8$_{F148S\_F151S}$ both remained monomeric throughout the experiment with average R_h values indicating sufficient burial of hydrophobic surfaces. (**D**) Representative images of cell lines expressing DNAJB8$_{F148S}$-mRuby3 (top right), DNAJB8$_{F151S}$-mRuby3 (bottom left), and DNAJB8$_{F148S\_F151S}$-mRuby3 (bottom right). DAPI is shown in blue and mRuby3 signal from fusion proteins is shown in red. The single F148S mutation resulted in a cell population with less observed puncta than DNAJB8$_{WT}$, with expression primarily in the cytoplasm and trace expression within the nucleus. However, the DNAJB8$_{F151S}$ and DNAJB8$_{F148S\_F151S}$ mutants showed few instances of puncta and were expressed in both the nucleus and cytoplasm, sharing the same phenotype as DNAJB8 S3. (**E**) Histogram of puncta frequency across cell lines expressing mRuby3 (red), DNAJB8$_{WT}$-mRuby3 (blue), DNAJB8$_{S3}$-mRuby3 (cyan), DNAJB8$_{F148S}$-mRuby3 (lime), DNAJB8$_{F151S}$-mRuby3 (purple) and DNAJB8$_{F148S\_F151S}$-mRuby3 (teal). Data for mRuby3, DNAJB8 $_{WT}$-mRuby3, and DNAJB8$_{S3}$-mRuby3 (gray box) were previously shown in Figure 1F. Single and double phenylalanine to serine mutations at positions F148 and F151 decreased the abundance of puncta within each cell line, with the double mutant F148S/F151S reaching comparable levels to DNAJB8$_{S3}$. This shows a relative decrease in self-assembly, with complete loss observed in DNAJB8$_{F151S}$ and DNAJB8$_{F148S\_F151S}$. All values were calculated from CellProfiler v.4.2.1. One-way ANOVA was used to report significance where F=24.7 and P-value=0.0006. ** $p<0.01$, *** $p<0.001$.

Figure 4. DNAJB8 preferentially binds a misfolded tau seed. (**A**) Domain maps of DNAJB8_WT/DNAJB8_S3, and full-length tau M_i/M_s used in *in vitro* activity experiments. Domain color scheme for DNAJB8 is the same as shown in Fig 1a. Tau domains are shown from N to C termini as N-terminal domains (N1 and N2; orange), proline rich domain (PRD; cyan), and all repeat domains (R1; red, R2; green, R3; blue, R4; purple, R'; dark grey). (**B**) MST binding curves for S3 B8 binding to M_i tau (black, circles) or M_s tau (red, squares). Fn values calculated from thermophoretic time traces, as a function of titrated protein. Baseline for unbound thermophoresis has been subtracted for the clarity of representation. Error bars represent the range of three replicates. K_d difference between both tau species is ~17-fold. (**C**) Relative coomassie signal intensity of the tau

monomer band on a SDS-PAGE gel where DMTMM was added and quenched immediately (0 min), after 7.5 min, or after 15 min and either in samples of 1μM tau M_i (black) or 1μM tau M_i with 20μM DNAJB8$_{WT}$ (blue). (**D**) Relative coomassie signal intensity of the tau monomer band on a SDS-PAGE gel where DMTMM was added and quenched immediately (0 min), after 7.5 min, or after 15 min and either in samples of 1μM tau M_i (black) or 1μM tau M_i with 20μM DNAJB8$_{S3}$ (cyan). (**E**) Contribution of different tau domains in the DNAJB8$_{WT}$:tau M_i XL-MS low-resolution (id score >10) results. Out of a total 173 pairs, the majority of cross-links are evenly distributed across the N-terminal domain, PRD, R1, and C-terminal domains. This is indicative of a mix of different interactions across multiple states of tau M_i. (**F**) Contribution of different tau domains in the DNAJB8$_{S3}$:tau M_i XL-MS low-resolution (id score >10) results. Out of a total 180 pairs, the majority of cross-links are evenly distributed across the N-terminal domain, PRD, R1, and C-terminal domains. These results closely mirror the tau domain contributions when crosslinked to DNAJB8$_{WT}$, showing a similar mode of interaction. (**G**) Contribution of different tau domains in the DNAJB8$_{WT}$:tau M_s XL-MS low-resolution (id score >10) results. Out of a total of 37 crosslinks, a majority of 62.12% localize to the PRD, with contributions from all other domains being minimal. The decrease in unique contacts combined with the increased frequency around a specific domain indicates that the DNAJB8$_{WT}$:tau M_s interaction is more defined than the DNAJB8$_{WT}$: tau M_i interaction. (**H**) Contribution of different tau domains in the DNAJB8$_{S3}$:tau M_s XL-MS low-resolution (id score >10) results. Out of a total of 37 crosslinks, a majority of 59.7% localize to the PRD, with contributions from all other domains being minimal. (**I**) Dynamic light scattering time course experiment reporting average R_h as time points for DNAJB8$_{WT}$ (brown), tau M_s (red), DNAJB8 $_{WT}$ and tau M_s (blue), and DNAJB8$_{S3}$ and tau M_s (cyan). After 20 hrs, tau M_s is able to rapidly aggregate in vitro at the specified concentration of 0.1μM. Meanwhile, at 10μM concentrations, DNAJB8$_{WT}$ is unable to self-assemble within the time course of the experiment. Each DNAJB8 mutant was mixed with tau M_s at a 100:1 ratio. At these conditions, DNAJB8$_{WT}$ is able to slow the rate of tau aggregation by ~20 hrs. However, the presence of DNAJB8$_{S3}$ with tau M_s delays the onset of tau M_s assembly by ~30 hrs.

Figure 5. Engineered monomeric DNAJB8 mutants retain anti-aggregation substrate activity in cells.

(**A**) Schematic of experiments involving stable HEK293T P301S tau biosensor lines. The tau biosensor cells are a stable line endogenously co-expressing P301S tauRD-mClover3 and P301S tauRD-mCerulean3. Upon transient transfection with sonicated tau fibrils, aggregation can be quantified through $FRET_{Clover/Cerulean}$. Before adding sonicated fibrils, DNAJB8-mRuby3 constructs are first transfected into the stable cell lines for expression 24 or 72 hours before seeding with fibrils. Changes to tau aggregation are then quantified by $FRET_{Clover/Cerulean}$. (**B**) Tau biosensor lines where transfected with mRuby3 (red), DNAJB6b$_{WT}$-mRuby3 (pink), and DNAJB8$_{WT}$-mRuby3 (blue) and expressed for 24 hours before either treatment with lipofectamine (solid bars) or seeded with 20nM sonicated tau fibrils (dashed lines). These lines were prepared as biological triplicate populations. Aggregation was quantified as a percentage of the cell line with positive $FRET_{Clover/Cerulean}$ signal. Relative to the mRuby3 control, both cell lines expressing DNAJB6b$_{WT}$ or DNAJB8$_{WT}$ showed a 2-fold decrease in $FRET_{Clover/Cerulean}$ positivity within the cell populations. (**C**) Tau biosensor lines where transfected with DNAJB8$_{WT}$-mRuby3 (blue), DNAJB8$_{S1}$-mRuby3 (green), DNAJB8$_{S2}$-mRuby3 (yellow), DNAJB8$_{S3}$-mRuby3 (cyan), DNAJB8$_{S4}$-mRuby3 (grey), DNAJB8$_{F148S}$-mRuby3 (lime), DNAJB8$_{F151S}$-mRuby3 (purple), and DNAJB8$_{F148S\_F151S}$-mRuby3 (teal) and expressed for 24 hours before either treatment with lipofectamine (solid bars) or seeded with 20nM sonicated tau fibrils (dashed lines). % $FRET_{Clover/Cerulean}$ was normalized to DNAJB8$_{WT}$ as shown in panel 5b to directly compare DNAJB8 mutants to DNAJB8$_{WT}$ activity (dashed line). Two-way ANOVA was used to calculate statistics across unseeded and 20nM tau fibrils lines, where no significant changes could be reported across the different mutants for either condition. (**D**) Percentage of cells containing mRuby3 puncta across tau biosensor cells transfected with mRuby3 (red), DNAJB6b$_{WT}$-mRuby3 (pink), and DNAJB8$_{WT}$-mRuby3 (blue) and expressed for 24 hours before either treatment with lipofectamine (solid bars) or seeded with 20nM sonicated tau fibrils (dashed lines). A 5-fold increase of puncta is observed for DNAJB8$_{WT}$-mRuby3 compared to the mRuby3 control under both unseeded and seeded conditions, indicating that this effect is driven by DNAJB8$_{WT}$. Statistics were calculated using two-way ANOVA, where * $p < 0.1$, **

p<0.01, **** p<0.0001. Values taken from image analysis with CellProfiler v.4.2.1. (**E**) Normalized number of puncta within tau biosensor cell populations transfected with DNAJB8$_{WT}$-mRuby3 (blue), DNAJB8$_{S1}$-mRuby3 (green), DNAJB8$_{S2}$-mRuby3 (yellow), DNAJB8$_{S3}$-mRuby3 (cyan), DNAJB8$_{S4}$-mRuby3 (grey), DNAJB8$_{F148S}$-mRuby3 (lime), DNAJB8$_{F151S}$-mRuby3 (purple), and DNAJB8$_{F148S\_F151S}$-mRuby3 (teal) and expressed for 24 hours before either treatment with lipofectamine (solid bars) or seeded with 20nM sonicated tau fibrils (dashed lines). The data were normalized to DNAJB8$_{WT}$-mRuby3 as shown in panel 5d to directly compare each mutant DNAJB8 construct to DNAJB8$_{WT}$-mRuby3 (dashed line). Similar to HEK293T cell experiments in Figure 1d and Figure 3e, monomeric DNAJB8 mutants DNAJB8$_{S3}$, DNAJB8$_{F151S}$, and DNAJB8$_{F148S\_F151S}$ show a significant decrease in puncta frequency that is dependent on the DNAJB8 mutations, but no changes observed due to the effect of tau aggregation. Statistics were calculated using two-way ANOVA, where * p<0.1, ** p<0.01, *** p<0.001, **** p<0.0001. Values taken from image analysis with CellProfiler v.4.2.1. (**F**) Schematic of experiments involving HEK293T cells with tetracycline-inducible HttEx1 Q119-GFP expression. DNAJB8-mRuby3 constructs were transfected into cells before inducing HttEx1 Q119-GFP expression. The cells were lysed and HttEx1 Q119-GFP aggregates were quantified via a filter trap assay. (**G**) FTA results showing HttEx1 Q119 aggregation for cells expressing mRuby3 (red), DNAJB8$_{WT}$-mRuby3 (blue), DNAJB8$_{S1}$-mRuby3 (green), DNAJB8$_{S2}$-mRuby3 (yellow), DNAJB8$_{S3}$-mRuby3 (cyan), DNAJB8$_{S4}$-mRuby3 (grey), DNAJB8$_{F148S}$-mRuby3 (lime), DNAJB8$_{F151S}$-mRuby3 (purple), and DNAJB8$_{F148S\_F151S}$-mRuby3 (teal). All values were normalized to the mRuby3 negative control. A small loss of DNAJB8 activity was reported for DNAJB8$_{S2}$-mRuby3, DNAJB8$_{S3}$-mRuby3, and DNAJB8$_{S4}$-mRuby3 expressing cells. Statistics were calculated using Welch's T test for each pair relative to DNAJB8$_{WT}$, where * p<0.1.

Figure 6. DNAJB8 oligomerization is not a prerequisite for activity.

(**A**) Model of DNAJB8 oligomeric assembly. The [147]AFSSFN[152] motif within the S/T-rich domain forms the core of the DNAJB8 oligomer with β-stranded dimers and Phe-Phe interactions stabilizing the hydrophobic core of the oligomer. (**B**) Model showing how Phe→Ser mutations in the S/T-rich domain result in a mutant of DNAJB8 that is incapable of self-assembling. This results in DNAJB8 remaining a stable monomer in solution in the absence of Phe-Phe interactions. (**C**) Model summarizing that DNAJB8 chaperone activity is independent on whether or not DNAJB8 exists as a monomer or oligomer. DNAJB8 mutants where oligomerization was inhibited showed equal activity

to DNAJB8$_{WT}$ and other mutants that were still capable of forming oligomers in cells. Therefore, the role of DNAJB8 oligomeric assemblies in cells is unrelated to its activity as a JDP chaperone.