Transcriptional adaptation of drug-tolerant *Mycobacterium tuberculosis* in mice

Elizabeth A Wynn,[1,2,3]* Christian Dide-Agossou,[1,3,4]* Matthew Reichlen,[3,5] Karen Rossmassler,[1,3,4] Reem Al Mubarak,[1,3,4] Justin J Reid,[1,3,4] Samuel T Tabor,[1,3,4] Sarah E M Born,[3,5] Monica R Ransom,[6] Rebecca M Davidson,[7] Kendra N Walton,[7] Jeanne B Benoit,[7] Amanda Hoppers,[7] Allison A Bauman,[8] Lisa M Massoudi,[8] Gregory Dolganov,[9] Payam Nahid,[3,10,11] Martin I Voskuil,[3,5] Gregory T Robertson,[3,8] Camille M Moore,[2,3,7]† Nicholas D Walter[1,3,4]†‡

* These authors contributed equally to this work (EAW, CD-A)
† These authors contributed equally to this work (NDW, CMM)
‡ Corresponding authors (NDW)

1. Rocky Mountain Regional VA Medical Center, Aurora, CO, USA
2. Department of Biostatistics and Informatics, University of Colorado, Anschutz Medical Campus, Aurora, CO, USA
3. Consortium for Applied Microbial Metrics, Aurora, CO, USA
4. Division of Pulmonary Sciences and Critical Care Medicine, University of Colorado Anschutz Medical Campus, Aurora, CO, USA
5. Department of Immunology and Microbiology, University of Colorado Anschutz Medical Campus, Aurora, CO, USA
6. Division of Hematology, Department of Medicine, University of Colorado School of Medicine, Aurora, CO, USA
7. Center for Genes, Environment and Health, National Jewish Health, Denver, CO, USA
8. Mycobacteria Research Laboratories, Department of Microbiology, Immunology, and Pathology, Colorado State University, Fort Collins, CO, USA
9. Division of Infectious Diseases and Geographic Medicine, Stanford University, Palo Alto, CA, USA
10. Division of Pulmonary and Critical Care Medicine, University of California San Francisco, CA, USA
11. UCSF Center for Tuberculosis, University of California, San Francisco, CA, USA

ABSTRACT

Transcriptome evaluation of *Mycobacterium tuberculosis* in the lungs of laboratory animals during long-term treatment has been limited by extremely low abundance of bacterial mRNA relative to eukaryotic RNA. Here we report a targeted amplification RNA sequencing method called SEARCH-TB. After confirming that SEARCH-TB recapitulates conventional RNA-seq *in vitro*, we applied SEARCH-TB to *Mycobacterium tuberculosis*-infected BALB/c mice treated for up to 28 days with the global standard isoniazid, rifampin, pyrazinamide, and ethambutol regimen. We compared results in mice with 8-day exposure to the same regimen *in vitro*. After treatment of mice for 28 days, SEARCH-TB suggested broad suppression of genes associated with bacterial growth, transcription, translation, synthesis of rRNA proteins and immunogenic secretory peptides. Adaptation of drug-stressed *Mycobacterium tuberculosis* appeared to include a metabolic transition from ATP-maximizing respiration towards lower-efficiency pathways, modification and recycling of cell wall components, large-scale regulatory reprogramming, and reconfiguration of efflux pumps expression. Despite markedly different expression at pre-treatment baseline, murine and *in vitro* samples had broadly similar transcriptional change during treatment. The differences observed likely indicate the importance of immunity and pharmacokinetics in the mouse. By elucidating the long-term effect of tuberculosis treatment on bacterial cellular processes *in vivo*, SEARCH-TB represents a highly granular pharmacodynamic monitoring tool with potential to enhance evaluation of new regimens and thereby accelerate progress towards a new generation of more effective tuberculosis treatment.

INTRODUCTION

Tuberculosis (TB) is an ongoing public health crisis, killing approximately 1.2 million people each year.[1] A key challenge for global TB control is the length of treatment required to cure TB, 4-6 months for infection with drug-sensitive genotypes. The need for prolonged treatment is thought to be due to a transition of the *M. tuberculosis* (*Mtb*) population to harder-to-kill phenotypes as therapy progresses.[2] At the start of treatment, the *Mtb* population is highly susceptible to killing. When mice or humans initiate the global standard isoniazid, rifampin, pyrazinamide, ethambutol (HRZE) regimen, ~99% of the culturable bacterial population is eliminated during the initial days to weeks of the bactericidal phase. Thereafter, the rate of killing slows considerably[3–7] as drug-tolerant phenotypes come to dominate the residual *Mtb* population. Drug-tolerant phenotypes have been defined as *Mtb* with decreased susceptibility to killing despite an absence of drug resistance conferring mutations.[2,8]

Eliminating drug-tolerant *Mtb* phenotypes is considered crucial to shortening the duration of TB treatment.[2] The central focus of contemporary drug discovery and regimen development is drug-tolerant *Mtb* phenotypes that cause relapse and dictate the need for prolonged treatment.[2,9–11] Unfortunately, the physiologic state of the *Mtb* subpopulation that withstands drug treatment *in vivo* remains obscure, limiting our ability to rationally design regimens that target specific cellular processes.

Transcriptional profiling has been used extensively as a readout of how drugs affect bacterial cellular processes *in vitro*.[12–18] By contrast, we are not aware of previous genome-wide transcriptional profiling of *Mtb* during prolonged treatment of laboratory animals. *In vivo* analyses are important because the pathogen adapts its cellular processes to the physicochemical

conditions and immune responses it encounters in the host, resulting in a phenotype and

transcriptome distinct from that measured *in vitro*.[19–27] Since phenotypic characteristics of the

Mtb population strongly influence drug sensitivity,[28] understanding the effect of drugs on the

bacterial populations that cause disease *in vivo* would enable us to better "know the enemy,"

thereby informing drug and regimen development.

The central impediment to transcriptional profiling *in vivo* is that *Mtb* mRNA is generally

present at extremely low abundance relative to host RNA even in a treatment-naïve animal.

When the pathogen burden decreases due to treatment, quantification of *Mtb* mRNA becomes

progressively more difficult. To elucidate *Mtb* signal from the overwhelming eukaryotic

background, diverse methods have been used to first enrich and then quantify *Mtb* mRNA (**Fig.**

1). Enrichment methods have included various combinations of differential cell lysis to separate

Mtb and eukaryotic material during sample preparation, reverse transcription with pathogen-

specific primers, depletion of host and/or bacterial rRNA, various hybridization capture methods,

and selective PCR amplification of target *Mtb* transcripts. Enrichment is followed by

quantification via microarray,[19,20,27,29] multiplex PCR,[7,21,25,30–33] or RNA-seq.[22,26,34–37]

Unfortunately, even the most sophisticated and contemporary platforms have limited sensitivity

that has precluded transcriptional evaluation of the effect of long-term treatment with potent TB

regimens. In this work, we demonstrate SEquencing after Amplicon enRiCHment for TB

(SEARCH-TB), a targeted, highly sensitive, pathogen-specific sequencing method for enriching

and quantifying the *Mtb* transcriptome in tissues of infected animals. Using a novel combination

of enrichment (differential cell lysis during sample preparation + targeted amplification) and

quantification (RNA-seq), SEARCH-TB achieves sensitivity that will enable routine use of

transcriptional profiling for drug evaluation in animal models, expanding understanding of drug

effect and accelerating new regimen development.

Here, we used SEARCH-TB to evaluate gene expression of the *Mtb* population that

persists in the lungs of BALB/c mice treated with HRZE for up to four weeks. We further used

SEARCH-TB to compare the effect of HRZE *in vivo* and *in vitro*. SEARCH-TB has the potential

to serve as a highly granular readout of drug effect by providing a molecular readout of cellular

adaptations of *in vivo* drug-tolerant *Mtb*.

METHODS

Design of the SEARCH-TB assay

The SEARCH-TB assay uses an AmpliSeq for Illumina custom pool designed to amplify

coding sequence (CDS) of *Mtb* complex (MTBC) organisms. To assure amplification across

diverse lineages, the design used eight MTBC reference genomes (**Table S1**). Annotations were

prepared for *Mtb* H37Rv (NC_000962) and *Mtb* Erdman (AP012340.1) as described in the

Supplemental Information. To avoid off-target amplification (*i.e.,* of non-MTBC organisms),

primers were cross-referenced during design with 12 "exclusion" genomes that included

phylogenetically diverse bacteria as well as human and mouse (**Table S2**).

Evaluation of amplification bias

We tested for amplification bias (*i.e.,* differences in amplification efficiency between

primer pairs targeting different *Mtb* sequences) using replicate human lung RNA samples spiked

with *Mtb* genomic DNA (gDNA) (Supplemental Information). Since all targeted sequences are

present as single copies in gDNA, an entirely unbiased assay would hypothetically result in the

same copy number for all targets, indicating that all primer pairs amplified with identical

efficiency. We defined amplification bias as deviation from this ideal by comparing the observed

expression for a gene to the expected expression assuming no amplification bias (Supplemental

Information).

Evaluation of repeatability of amplification

To evaluate repeatability, we spiked 1pg *Mtb* RNA into 1ng human lung RNA

(Supplemental Information). We compared counts per million (CPM) values for each gene

between technical replicates. We also evaluated repeatability over time by prepping and

sequencing 19 of the *in vitro* and murine samples described below two times with up to a 4-

month intervening interval. We quantified batch effect by calculating the difference in

expression (normalized with DESeq2's variance stabilizing transformation[38]) of each gene

between replicate pairs, then averaging across replicates. We compared the magnitude of the

batch effect with the observed treatment effect.

Concordance of SEARCH-TB with conventional RNA-seq

We evaluated whether SEARCH-TB identified the same transcriptional changes as a

conventional RNA-seq method without *Mtb* targeted amplification (Illumina TrueSeq) after 24-

hour *in vitro* isoniazid (INH) exposure (Supplemental Information). We first evaluated RNA

from control (N=4) and INH-treated samples (N=4) via conventional RNA-seq to serve as a

reference standard. We then spiked the same RNA from control and INH-treated *Mtb* into human

lung RNA at a ratio of 1:1,000 and sequenced via SEARCH-TB. After calculating differential

expression between control and INH-treated samples separately for conventional RNA-seq and

SEARCH-TB using edgeR,[39] we compared the significant genes and fold-changes identified by

the two platforms.

In vitro experiments

Mtb strains H37Rv and Erdman were cultured *in vitro* using Middlebrook 7H9 broth

(Difco) supplemented with 0.085 g/l NaCl, 0.2% glucose, 0.2% glycerol, 0.5% BSA, and 0.05%

Tween-80. All culturing was performed at 36.5°C and 5.0% CO_2. Single use frozen *Mtb*

aliquots were revived in 7H9 and grown to mid-log phase then cultures were diluted to

OD_{600}=0.05, dispensed in 5.0 ml aliquots into sterile glass tubes (20 by 125 mm) containing

sterile stir bars (12 by 4.5 mm), and outgrown for 18 h under rapid agitation (~200 rpm stirring

speed using a rotary magnetic tumble stirrer) prior to the initiation of drug exposure. RNA was

collected from *Mtb* H37Rv exposed to INH or *Mtb* Erdman exposed to HRZE *in vitro*

(Supplemental Information).

Murine drug experiments

All animal procedures were conducted according to relevant national and international

guidelines and approved by the Colorado State University Animal Care and Use Committee as

described in the Supplemental Information. Briefly, female BALB/c mice, 6 to 8 weeks old, were

aerosol infected (Glas-Col) with *Mtb* Erdman strain resulting in the deposition of 4.55±0.03

(SEM) log10 CFU in lungs one day following aerosol. After 11 days, five mice were euthanized

to serve as the pre-treatment control group. Groups of five mice each were treated with HRZE at

standard doses five days a week for 14 or 28 days before euthanasia. Lungs were aseptically

dissected, and flash frozen in liquid nitrogen before processing.

RNA extraction, sequencing, and data preparation

RNA extraction, library preparation, sequencing and data preparation are detailed in the

Supplemental Information.

Statistical analysis of murine and *in vitro* experiments

Murine and *in vitro* sequence data were analyzed together using edgeR[39,40] to identify the

effect of HRZE treatment in and compare gene expression between murine and *in vitro*

experiments. We fit negative binomial generalized linear models to each gene and included terms

for murine and *in vitro* time points (control, day 14 and day 28 in mice; control, day 4 and day 8

in vitro). Likelihood ratio tests were performed to compare expression between murine time

points, between *in vitro* time points, and between murine and *in vitro* experiments before and at

the end of treatment. Genes with Benjamini-Hochberg adjusted *P*-value[41] less than 0.05 were

considered significant. For murine experiments, we used hierarchical clustering to identify

groups of genes with similar changes in gene expression over the course of treatment as follows.

First, for genes that were differentially expressed between at least two time points, we calculated

the expected expression at each time point using the edgeR models. Then, the expected

expression values were hierarchically clustered based on Euclidian distance using Ward's

method[42] to find clusters of genes with similar patterns of expression over time.

Principal component analysis (PCA) was performed using data for the 500 most variable

genes after normalization with DESeq2's variance stabilizing transformation.[38]

Using hypergeometric tests in the hypeR R package,[43] we performed functional

enrichment for each pairwise combination of murine and *in vitro* time points to evaluate whether

differentially expressed genes were overrepresented in gene categories established by Cole[44] or

curated from the literature (**Table S3**). Enrichment analysis was run twice for each pairwise

combination, first using significantly upregulated genes and then using significantly

downregulated genes. Gene categories with <8 genes were excluded. Gene categories with

Benjamini-Hochberg adjusted *P*-values[41] less than 0.05 were considered significant. All analysis

used R (v4.1.1).[45]

Online analysis tool

Differential expression, functional enrichment, and visualizations can be evaluated

interactively using an Online Analysis Tool [https://microbialmetrics.org/analysis-tools/] created

using the R package Shiny.[46]

RESULTS

Validation of SEARCH-TB assay

Results of SEARCH-TB design

Primers were designed to amplify 3,733 (92.6%) of 4,031 CDS in the *Mtb* H37Rv genome.

Primers for 95 genes failed to amplify in initial testing and were excluded from analysis.

Additionally, we excluded 70 genes that were not present in the *Mtb* Erdman strain used in

murine and *in vitro* experiments. The final panel analyzed targeted 3,568 *Mtb* genes (**Fig. S2**).

Evaluation of amplification bias

Using gDNA (a matrix in which each target sequence should be present in equal abundance), we

quantified deviation from the ideal value that would be expected if all primers had identical

amplification efficiency (**Fig. 2a**). Of all primer pairs, 78% were within one $\log_2$ fold change of

the ideal value.

Repeatability of amplification

SEARCH-TB results were highly repeatable among spike-in technical replicates (**Fig. 2b**). The

batch effect between replicates prepared and sequenced over time was minimal relative to the

treatment effect (**Fig. S3**).

Concordance of SEARCH-TB with conventional RNA-seq

RNA from *in vitro* exposure showed a similar direction and scale of differential expression when

sequenced via SEARCH-TB and conventional RNA-seq (**Fig. 2c-d**). The gene sets identified as

differentially expressed by the two platforms strongly overlapped (**Fig. 2e**). The fold-changes

quantified by the two platforms showed strong agreement (R^2=0.82) (**Fig. 2f**), indicating that

while SEARCH-TB is uniquely capable of profiling in drug-treated animals, both platforms

provide the same biological information in *in vitro* RNA.

Transcriptional differences of between *Mtb* in murine and *in vitro* controls

In the absence of HRZE, a PCA plot showed the transcriptional profiles of the pre-

treatment murine control and the *in vitro* early log-phase growth control were distinct (**Fig. 3a**)

with 2,444 (68%) differentially expressed genes (**Fig. 3b**). Compared with the *in vitro* control,

untreated mice appeared to harbor a less active *Mtb* phenotype with lower expression of genes

associated with rRNA protein synthesis (adj-*P*=1.7x10^{-18}), aerobic metabolism (adj-*P*=2.1x10^{-5}),

fatty acid synthesis (adj-P=0.002) and other growth-associated processes (Online Analysis Tool).

Conversely, *Mtb* in untreated mice had higher expression of processes previously associated with

adaptation to the host[47], including genes in the DosR regulon that responds to hypoxia, nitric

oxide, and carbon monoxide[48,49] (adj-P=1.4x10^{-12}), the KstR1 and KstR2 regulons involved in

cholesterol catabolism[50] (adj-P=7.2x10^{-5} and 3.9x10^{-4}, respectively), mycobactin genes that

respond to iron scarcity[51] (adj-P=6.4x10^{-4}) and other processes (Online Analysis Tool).

SEARCH-TB recapitulated specific previously described adaptations for persistence *in vivo,*

including higher expression in untreated mice compared to the *in vitro* control of *icl1* (adj-

P=3.8x10^{-13}), the first gene of the glyoxylate bypass used during catabolism of fatty acids[52], and

increased expression of *tgs1* (adj-P=8.1x10^{-35}) involved in triacylglycerol synthesis under

environmental stress[53].

Efficacy of drug treatment

In mice, treatment with HRZE reduced the CFU burden by 99.8% by day 28 (7.63 to 4.82

log$_{10}$ CFU). A newer marker, the RS ratio,[10] declined more rapidly than CFU, indicating that

interruption of rRNA synthesis occurs more rapidly than killing (**Fig. 3c**).

Global effect of HRZE on the *Mtb* transcriptome

In mice, treatment with HRZE transformed the *Mtb* transcriptome, significantly altering

expression of 2,049 (57%) and 2,329 (65%) genes at day 14 and day 28 relative to pre-treatment

control on day 0, respectively (**Fig. 3d-e**). A smaller set of genes (159 (4%)) was differentially

expressed at day 28 relative to day 14. Hierarchical clustering identified clusters of genes with

similar expression changes over time (**Fig. 3f**). The greatest transcriptional change occurred by

day 14 (**Fig. 3g**, Clusters 2 and 4). A smaller subset of genes changed more gradually (**Fig. 3g**,

Clusters 1 and 3). *In vitro* HRZE exposure also had a large impact on the *Mtb* transcriptome,

altering expression of 2,748 (77%) and 2,800 (78%) of genes at day 4 and day 8 relative to

control, respectively.

There were broad similarities as well as important differences between the effects of

HRZE in mice and *in vitro*. Considering the effect of HRZE on gene expression at the latest

treatment time points (8 days *in vitro* and 28 days in mice), differentially expressed genes were

largely concordant between the mouse and *in vitro* samples (**Fig 3h**). However, eight days of

HRZE *in vitro* induced overall larger fold-changes than 28 days of HRZE in mice (**Fig. 3i**). A

notable subset of genes changed in discordant directions (colored gold in **Fig. 3i**), indicating

differences between the effect of HRZE in mice and *in vitro* that are further described below.

Effect of HRZE on bacterial cellular processes

Since our primary focus is *Mtb* phenotypes that survive prolonged drug exposure, the

remainder of this manuscript describes only expression changes at the latest time point (28 days

in mice or 8 days *in vitro*) relative to control, unless otherwise noted. Differences between other

time points can be evaluated via the Online Analysis Tool.

Genes significantly impacted by 28-day HRZE treatment in mice were enriched for 36 of

the 124 functional gene sets (**Table S3**) evaluated in enrichment analysis. Figure 4a provides a

global portrait of change in cellular processes.

Growth and synthesis of macromolecules

A dominant effect of HRZE was decreased expression of categories associated with

growth and synthesis of macromolecules. Genes coding for ribosomal proteins were profoundly

suppressed after treatment in mice (adj-P=2.9x10^{-15}) and *in vitro* (adj-P=1.0x10^{-14}) relative to

their respective controls (**Fig. 4b**), consistent with decreased ribosome synthesis, a process

fundamentally coupled with bacterial replication.[54,55] An exception to this pattern were the four

alternative C-ribosomal protein paralogs lacking the zinc-binding CXXC motif that are a

component of ribosomal remodeling. As shown in red in Figure 4b, genes for the alternative

ribosomal proteins (*rpmB1, rpsR2, rpsN2, rpmG1*) had either sustained or significantly increased

expression relative to control, consistent with ribosomal remodeling by a slowly replicating *Mtb*

population.

239   Reduced protein synthesis was additionally suggested by decreased expression of the

protein translation and modification category that includes genes responsible for translational

initiation, promotion of tRNA binding, elongation, termination, and protein folding (adj-P=0.006

in mice, adj-P=0.007 *in vitro*). Concordant with downregulation of translation, transcription

appeared suppressed, as evidenced by significant decrease in genes coding for gyrase A and B

subunits (*gyrA, gyrB*) in mice and *in vitro* (least significant adj-P=1.8x10^{-5}).

Cell-wall synthesis, remodeling, and recycling

Mycolic acids. Slowing of the initial step of mycolic acid synthesis was suggested by decreased

expression of Rv2524c (*fas),* the gene coding for fatty acid synthetase I (FAS-I), in mice (adj-

P=1.7x10^{-9}) and *in vitro* (adj-P=3.2x10^{-9}) and by decreased expression of the *fas* transcriptional

regulator Rv3208[56] in mice (adj-P=1.9x10^{-4}) and *in vitro* (adj-P=1.1x10^{-11}). Genes coding for the

second step of elongation of acyl-coenzyme A to long chain fatty acids by fatty acid synthetase II

(FAS-II) were also significantly decreased in both mice (adj-P=0.011) and *in vitro* (adj-

P=0.027) (**Fig. 4c**). Finally, in *Mtb* treated with HRZE in mice and *in vitro,* there was decreased

expression of the set of genes associated with elongation, desaturation, modification, and

transport of the mature mycolic acids to the cell wall.

Phthiocerol dimycocerosates (PDIM). Genes coding for PDIM biosynthesis, the outer surface

glycolipids that are important for intracellular survival and virulence, were significantly

downregulated in mice (adj-*P*=9.9×10^{-5}) and *in vitro* (adj-*P*=0.017).

Peptidoglycan. In contrast to the downregulation observed in mycolic acid and PDIM genes, key

genes involved in peptidoglycan synthesis, modification, and recycling had increased expression.

Specifically, all five peptidoglycan synthesis genes assayed in the division cell wall operon (*ftsQ,*

murC, ftsW, murD, murR, murE) had significantly increased expression at 14 days of HRZE and

three remained significantly upregulated at day 28 after HRZE treatment in mice (Online

Analysis Tool). Active recycling was also suggested by significantly increased expression with

HRZE treatment in mice and *in vitro* of genes coding for the UspABC amino-sugar importer and

the DppABCD dipeptide importer that transport peptidoglycan breakdown products (**Fig. 4d**).[57]

Trehalose. The TreY/Z genes that synthesize free trehalose from glycogen were significantly

upregulated at all post-treatment time points in mice and *in vitro.* The other two trehalose

synthesis pathways (OtsA/B and GlgE/TreS) were not differentially expressed with HRZE

treatment in mice or *in vitro.* Remodeling of the trehalose component of the cell envelope was

suggested by significantly increased expression of Rv3451 (*cut3*) under the test conditions in

mice (adj-*P*=0.002) and *in vitro* (adj-*P*=4.2×10^{-4}), which codes for a stress-responsive trehalose

dimycolate hydrolase.[58] Additionally, both mice and *in vitro* samples had significantly increased

expression of four of the five genes for the LpqY/SugABC importer that is specific for the

transport of trehalose.[59]

Metabolic readjustment

Electron transport and aerobic respiration. The eight genes that collectively code for ATP synthetase were strongly suppressed in mice (adj-*P*=7.6x10^{-4}) and *in vitro* (adj-*P*=.003) (**Fig. 4e**). Oxidative phosphorylation appeared to transition from the primary cytochrome *bcc/aa3* supercomplex (downregulated) to the alternative less efficient cytochrome *bd* oxidase (upregulated) that has been implicated in persistence under environmental and drug stress[60] (**Fig 4f**). Genes of the TCA cycle were downregulated with HRZE treatment in mice (adj-*P*=0.001) and *in vitro* (adj-*P*=0.010). Genes coding for NADH dehydrogenase types I and II and succinate dehydrogenase types I and II were also downregulated (Online Analysis Tool). By contrast, three of the four fumarate reductase genes were significantly induced by HRZE in mice, and all were induced *in vitro*.

Central carbon metabolism. Slowing of the TCA cycle was not accompanied with increased expression of glyoxylate bypass genes, an alternative pathway previously implicated in drug tolerance.[52] Instead, the gene for isocitrate lyase (*icl1*), the first step of the glyoxylate bypass, was among the most profoundly suppressed by HRZE both in mice (adj-*P*=1.9x10^{-37}) and *in vitro* (adj-*P*=2.0x10^{-14}) (**Fig. 4g**). The gene coding for the alternative isocitrate lyase (*aceAa*) was also significantly downregulated during HRZE treatment in mice (adj-*P*=7.3.x10^{-13}) and *in vitro* (adj-*P*=3.x10^{-15}). Genes associated with carbon storage as triacylglycerol had discordant regulation in mice and *in vitro*. Specifically, *tgs1,* a gene in the DosR regulon which codes for triacylglycerol synthase, decreased 4.4-fold (adj-P=7.0x10^{-13}) in mice but increased 10.9-fold (adj-P=3.2x10^{-16}) *in vitro.*

Cholesterol degradation

Although more highly expressed in mice than *in vitro* prior to drug exposure, genes

related to cholesterol breakdown were suppressed by HRZE in mice and increased by HRZE *in*

vitro (**Fig 5a,** significance of individual genes in Online Analysis Tool).

Glycerophosphodiesters. The four genes coding for the UgpABCE transporter responsible for

import of glycerophosphodiesters that are a major source of carbon and phosphate[61] were

strongly upregulated at both treatment time points in mice and *in vitro* (Online Analysis Tool).

Lipolytic enzymes

The complex repertoire of lipolytic enzymes that are implicated in *Mtb* dormancy and

virulence includes Lipase family genes predicted based on sequences to have lipase/esterase

activity. In mice, 11 of the lipase genes included in the SEARCH-TB assay had significantly

increased expression, including *lipY* that codes for triacylglycerol lipase LipY. There was

particularly strong induction of *lipY in vitro* (15.5-fold increase, $P=1.2 \times 10^{-35}$). Additionally,

three of the four phospholipase C genes (*plcA, plcB, plcC*) were significantly upregulated in

mice.

Canonical stress responses

Although the 14 heat shock proteins (HSPs) that act as chaperones in protein folding are

often described as a stress response, expression of HSP genes largely decreased following HRZE

exposure. A notable discordance between mice and *in vitro* results is the hypoxia-responsive

hspX (*arc*) (noted in red in **Fig. 5b**) that had the greatest negative fold-change of all genes

evaluated in mice (48.8-fold decrease, adj-$P=4.8 \times 10^{-57}$) yet was significantly upregulated *in vitro*

(4.6-fold increase, adj-$P=2.0 \times 10^{-8}$). Concordant with the *hspX* result, we found that HRZE

significantly decreased expression of the DosR regulon in mice (adj-P=0.032) but induced the

DosR regulon *in vitro* (adj-P=2.2×10^{-7}) (**Fig. 5c**). Genes of the stringent response, which is

typically induced by nutrient deprivation and other environmental stresses, were significantly

downregulated with HRZE treatment in mice (adj-P=2.9×10^{-15}) and *in vitro* (adj-P=1.6×10^{-14})

(Online Analysis Tool). Similarly, *relA,* the gene coding for the stringent response regulator was

significantly decreased in mice (adj-P=2.0×10^{-6}) and *in vitro* (adj-P=7.0×10^{-6}).

Secretion of peptides and immunogenic proteins

Genes coding for the Antigen 85 complex, the major secreted protein that is essential for

intracellular survival within macrophages[62], were profoundly downregulated under the test

conditions in mice (smallest fold-decrease=6.5, least-significant adj-P=3.1×10^{-24}) and *in vitro*

(smallest fold-decrease=14.2, least-significant adj-P=2.6×10^{-31}) (**Fig. 5d**). Expression of the

ESX-1 locus was downregulated (adj-P=9.4×10^{-5} in mice, adj-P=1.3×10^{-5} *in vitro*) with

particularly strong suppression of *esxA* and *esxB,* genes coding for highly-immunogenic early

secretory antigenic 6 kDa (ESAT-6) and culture filtrate protein 10 (CFP-10), in mice (smallest

fold-decrease=10.0, least-significant adj-P=4.7×10^{-24}) and *in vitro* (smallest fold-decrease=55.7,

least-significant adj-P=2.4×10^{-35}) (**Fig. 5e**). Expression of all 9 genes assayed in the ESX-3 locus

was significantly downregulated with particularly strong suppression of immunogenic *esxG* and

esxH in mice (smallest fold-decrease=18.5, least-significant adj-P=1.1×10^{-34}) and *in vitro*

(smallest fold-decrease=44.3, least-significant adj-P=4.0×10^{-32}). In contrast, six of the seven

genes in the ESX-4 locus that a recent review[63] described as having "wholly unknown" function

had significantly increased expression. Interestingly, the Sec-independent Tat system that exports

pre-folded proteins had significantly increased expression of genes coding for the TatBC

complex (*tatB, tatC*) but significantly decreased expression of the gene for the PMF-dependent

TatA pore protein (*tatA*).

Efflux pumps

Most of the diverse set of transporters that includes putative drug efflux pumps had

altered expression after HRZE exposure, but the direction of their regulation varied (**Table S4**).

As examples, Rv2686c-2688c that is associated with fluoroquinolone tolerance[64] was

upregulated in mice and *in vitro* (**Fig. 5f**) but DrrABC that is associated with daunorubicin

tolerance and appears in clinical drug-resistant strains[64] was downregulated (**Fig. 5g**).

Transcriptional and post-transcriptional regulation

SEARCH-TB indicated large-scale regulatory reprogramming. For example, most of the

188 transcription factors assayed in mice changed after 28 days of HRZE treatment, with 66

significantly increased and 47 significantly decreased. The regulatory perturbation was even

more pronounced *in vitro* with 87 significantly increased and 55 significantly decreased

transcription factors after 8 days of HRZE treatment. Of the 12 sigma factor genes included in

SEARCH-TB, *sigF, sigI* and *sigM* were significantly upregulated following HRZE treatment in

both mice and *in vitro. sigA, sigB, sigD* and *sigK* were significantly downregulated in both mice

and *in vitro.* Expression of *sigC* and *sigL* was significant in both mice and *in vitro* but in

discordant directions, suggesting differing regulatory responses *in vivo* and *in vitro.*

HRZE appeared to activate the post-transcriptional toxin-antitoxin system that modulates

the concentration of existing transcripts. Toxin genes had significantly increased expression in

mice (adj-*P*=0.011) and *in vitro* (adj-*P*=0.027). The counter-regulatory antitoxins that restrict

toxin activity were not categorically altered in mice but were significantly suppressed *in vitro*

(adj-*P*=0.037).

Drug targets

Because processes that are upregulated following prolonged drug exposure may represent

survival mechanisms that could be targeted to eradicate persisting *Mtb,* we evaluated genes that

code for drug targets. Evaluation of targets of 31 existing drugs or investigational compounds

(36 genes) (**Table S5**) were predominantly suppressed in mice (19 down-regulated and 3

upregulated) and *in vitro* (26 down-regulated and 2 upregulated). Notable exceptions to

downregulation of drug targets were increased expression of the genes for Mur ligases B and C

that initiate peptidoglycan synthesis in mice[65] and increased expression of *rfe,* the gene for

phosphoglycosyltransferases WecA that initiates arabinogalactan synthesis in mice and *in vitro.*[66]

Transcriptional differences between *Mtb* in mice and *in vitro* at final time points

After 8 and 28 days of HRZE treatment, the *in vitro* and murine *Mtb* transcriptomes were

more similar than they were prior to HRZE exposure (**Fig 3a**). Nonetheless, 1,014 genes (28%)

remained differentially expressed between the final murine and *in vitro* time points.

DISCUSSION

This work established the SEARCH-TB platform, a method for elucidating the *Mtb*

transcriptome during drug exposure *in vivo*. After 28 days of treatment with the global standard

4-drug combination (HRZE) and 99.8% reduction in the burden of culturable *Mtb* in mouse

lungs, SEARCH-TB indicated broad suppression of cellular activity including slowing of

metabolism, synthesis of macromolecules and secretion of immune-modulating peptides.

SEARCH-TB also suggested bacterial adaptation to drug stress, including a shift in electron

transport to the alternative less efficient cytochrome *bd* oxidase, ribosomal remodeling, cell wall

remodeling and recycling, and reprogramming of regulatory and efflux pump activity. The

effects of HRZE in mice and *in vitro* had both broad similarities and notable differences that

likely reflect effects of immunity and environment. As the first platform capable of nearly

genome-wide quantification of extremely low abundance *Mtb* transcripts in mice, SEARCH-TB

should enable highly granular evaluation of the effect of drugs and regimens *in vivo*.

Unsurprisingly, at baseline, prior to the start of HRZE treatment, SEARCH-TB indicated

that *Mtb* cellular processes differed substantially between mice and early log phase growth,

reflecting bacterial adaptation to immunity and the lung environment. These differences between

untreated mouse and *in vitro* control are consistent with a recent review[47] of the treatment-naïve

in vivo Mtb transcriptome, both in terms of broad cellular processes (downregulation of genes

associated with transcription, translation and metabolism *in vivo*) and specific adaptations (*e.g.,*

increased expression of genes of the DosR regulon, glyoxylate bypass genes). Importantly, the

transcriptome of the untreated mouse was the starting point for our current analysis of drug

effect. Combination antimicrobial treatment is a categorically different type of stress than

environmental conditions such as pH, hypoxia, and nutrient starvation. Correspondingly, as

highlighted in the discussion below, SEARCH-TB showed that drug exposure elicited

transcriptional adaptations that often diverge from well-established transcriptional responses to

environmental stress.

SEARCH-TB results indicated that a major effect of HRZE treatment in mice is

suppression of *Mtb* growth and metabolism. Consistent with previous *in vitro* analyses of drug

effects, transcription results suggested decreased synthesis of all major macromolecules,

including rRNA, protein, lipids, and cell wall constituents. An additional manifestation of

decreased bacterial activity was decreased protein and peptide secretion, including the ESX1

Type VII secretion system that exports the highly immunogenic ESAT-6 and CFP-10 proteins.

This suggests that drug stress might alter the pathogen's capacity to modulate host immunity.

Despite the broad downregulation of activity, SEARCH-TB provided evidence that drug-

stressed *Mtb* is not inert or incapacitated. Increased expression of genes associated with

peptidoglycan and trehalose synthesis and recycling suggested active cell wall modification.

SEARCH-TB indicated broad transcriptional and post-transcriptional regulatory reprogramming

and reconfiguration of efflux pump expression. HRZE appears to have induced reconfiguration

of metabolism and energy generation away from high respiratory activity that maximizes ATP

generation, and towards reduced respiratory efficiency and oxidative phosphorylation activity.

A striking finding was that HRZE led to significantly decreased expression of the DosR

regulon in mice. DosR is the *Mtb* response to impaired aerobic respiration induced by hypoxia,

nitric oxide, or carbon monoxide exposure.[67] DosR was termed the "dormancy survival regulon"

because impaired respiration leads to growth arrest. In mice, HRZE appeared to arrest growth yet

expression of the DosR regulon was suppressed. We hypothesize that once 99.8% of culturable

Mtb have been eliminated and secretion of immunogenic peptides, including Ag85, ESAT-6 and

CFP-10, is diminished in the residual population, host immunity may be modulated. A less-

intense inflammatory response would be predicted to decrease increase localized oxygen

availability and decrease nitric oxide production, thereby diminishing induction of DosR-

inducing conditions that restrict aerobic respiration. The hypothesis that DosR may be an indirect

readout of immune response is consistent with our previous observation that the DosR regulation

has significantly lower expression in TB patients with AIDS than in immunocompetent

patients.[33] Also consistent with this hypothesis was our finding that *in vitro*, in the absence of

immunity, HRZE significantly increased rather than decreased expression of DosR genes.

Also notable was expression of the gene for isocitrate lyase, the first step of the

glyoxylate bypass that is essential to establishing infection in animals.[68–70] Consistent with these

prior results, we found that *icl1* was highly expressed in untreated mice relative to log phase

growth. *icl1* was also shown to have increased expression following sublethal exposure to

rifampin, INH or streptomycin *in vitro* and was identified as a mediator of drug tolerance.[52] By

contrast, we found that treatment with lethal doses of HRZE strongly suppressed, rather than

induced, expression of both *icl1* and *aceAa* which codes for an alternative isocitrate lyase. This

highlights, first, that adaptations associated with prolonged tolerance to HRZE differ from

adaptations to host environments that enable persistent infection in the absence of drug therapy

and, second, that the response to exposure to a combination regimen *in vivo* may differ from

single drug exposure *in vivo*.

Drugs have historically targeted transcription, translation, and other growth-associated

cellular processes[71] that SEARCH-TB shows are downregulated after treatment with HRZE.

While transcriptional downregulation of drug targets does not necessarily mean a drug will be

ineffective, genes such as the Mur ligases that have increased expression after 1-month of HRZE

treatment *in vitro* are noteworthy as they might indicate adaptations that enable *Mtb* to withstand

drug exposure. Similarly, genes coding for the alternative bd oxidase that is a proposed drug

target[72] were upregulated at a time when nearly all other metabolism-associated genes had

suppressed expression.

A novel comparison in this report is the effect of HRZE in the mouse versus *in vitro.*

Although the physiologic state of *Mtb* appeared quite different in mice and *in vitro* at pre-

treatment baseline, HRZE induced broadly similar changes *in vivo* and *in vitro*. The magnitude

of fold-change was greater *in vitro,* potentially indicating a lower effective drug exposure in the

mouse. Nonetheless, we also observed discordant transcriptional changes such as in expression

of the DosR regulon, indicating that drug effect *in vitro* is not a simple surrogate for effect in the

mouse*.* Furthermore, at the latest treatment endpoints, a large number of genes remained

differentially expressed between *in vitro* and the mouse, likely indicating the importance of

immunity and pharmacokinetics in the mouse.

As a highly granular readout of the effect of TB treatment on bacterial cellular processes,

SEARCH-TB may improve precision of pharmacodynamic evaluation, providing greater

information than the existing standard method of assessing drug effectiveness *in vivo* (*i.e.,*

culture-based enumeration of bacterial burden[73]). We have previously shown that CFU burden

does not capture the entirety of complex drug effects *in vivo*. For example, regimens that have

identical effects on CFU in mice can have different long-term relapse outcomes.[10,74] In previous

studies, we demonstrated that quantification of rRNA synthesis via the RS ratio showed proof of

concept that molecular measures of bacterial cellular processes *in vivo* can distinguish regimens

that are indistinguishable based on CFU.[10,74] SEARCH-TB advances molecular characterization

of drug effects *in vivo* to a higher level of granularity. Others have demonstrated the power of

Mtb transcriptional readouts of drug effect to predict drug interactions *in vitro*.[75] We believe that

SEARCH-TB has potential to enable a new era of pharmacodynamic evaluation in which drug

interactions and regimens are assessed based on molecular effects on cellular process *in vivo.*

This report has several limitations. First, between-target variation in the amplification

efficiency of SEARCH-TB primers likely affects the rank-order of gene counts, meaning that,

for individual genes, a higher absolute count does not necessarily indicate greater expression.

However, because amplification is highly repeatable, the modest amplification bias identified

does not affect estimation of differential expression between groups which is the primary

purpose of SEARCH-TB. Indeed, we showed that SEARCH-TB provides the same biological

interpretation as conventional RNA-seq. Second, specific and efficient primers could not be

designed for ~12% of *Mtb* transcripts. Third, transcriptional profiling is inherently unable to

resolve the enduring question of whether the drug-tolerant subpopulation that persists late into

treatment results from selection (*i.e.,* elimination of easily-killed *Mtb*) or physiological

transformation of *Mtb* that were present at baseline. Finally, this initial demonstration of

SEARCH-TB evaluated a single regimen (HRZE) in a single murine model. Important next steps

include evaluation of individual drugs and diverse regimens in additional animal models.

SEARCH-TB enabled what to our knowledge is the first evaluation of the effect of

prolonged drug treatment on *Mtb* transcription in animal models, revealing adaptations distinct

from those observed under environmental stress. The *Mtb* subpopulation that survived one

month of HRZE treatment appeared substantially less active than prior to treatment but was not

inert with transcriptional changes suggesting adaptation for survival. SEARCH-TB should

enable a new era of *in vivo* molecular pharmacodynamics with potential to accelerate

identification of new highly potent regimens.

Data availability. All raw sequencing data have been deposited in the Sequence Read Archive

(SRA) under BioProject accession PRJNA939248. Individual samples have BioSample

accession numbers SAMN33461189 through SAMN33461251.

Funding. GR acknowledges funding from the Bill and Melinda Gates Foundation (INV-

009105). NDW and RD acknowledge funding from the Bill and Melinda Gates Foundation

(OPP1170003). NDW and PN acknowledge funding from the US National Institutes of Health

(1R01AI127300-01A1). NDW acknowledges funding from Veterans Affairs (1I01BX004527-

01A1).

Acknowledgements. We are grateful to the Illumina Design Team that developed the custom

primer pool used in SEARCH-TB.

REFERENCES

1.   World Health Organization. *Global tuberculosis report 2020*. (2020).

2.   Connolly, L. E., Edelstein, P. H. & Ramakrishnan, L. Why is long-term therapy required

to cure tuberculosis? *PLoS Med.* **4**, e120 (2007).

3.   Mitchison, D. A. Basic mechanisms of chemotherapy. *Chest* **76**, 771–780 (1979).

4.   Jindani, A., Aber, V. R., Edwards, E. A. & Mitchison, D. A. The early bactericidal activity

of drugs in patients with pulmonary tuberculosis. *Am. Rev. Respir. Dis.* **121**, 939–949

(1980).

5.   Jindani, A., Doré, C. J. & Mitchison, D. A. Bactericidal and sterilizing activities of

antituberculosis drugs during the first 14 days. *Am. J. Respir. Crit. Care Med.* **167**, 1348–

1354 (2003).

6.   Davies, G. R., Brindle, R., Khoo, S. H. & Aarons, L. J. Use of nonlinear mixed-effects

analysis for improved precision of early pharmacodynamic measures in tuberculosis

treatment. *Antimicrob. Agents Chemother.* **50**, 3154–3156 (2006).

7.   Walter, N. D. *et al.* Transcriptional adaptation of drug-tolerant *Mycobacterium*

tuberculosis during treatment of human tuberculosis. *J. Infect. Dis.* **212**, 990–8 (2015).

8.   Balaban, N. Q., Gerdes, K., Lewis, K. & McKinney, J. D. A problem of persistence: Still

more questions than answers? *Nature Reviews Microbiology* **11**, 587–591 (2013).

9.   Mitchison, D. A. Role of individual drugs in the chemotherapy of tuberculosis. *Int. J.*

Tuberc. Lung Dis. **4**, 796–806 (2000).

10.   Walter, N. D. *et al.* *Mycobacterium tuberculosis* precursor rRNA as a measure of

treatment-shortening activity of drugs and regimens. *Nat. Commun.* **12**, 1–11 (2021).

11. Davies, G. R. Early clinical development of anti-tuberculosis drugs: Science, statistics and

sterilizing activity. *Tuberculosis* **90**, 171–176 (2010).

12. De Knegt, G. J. *et al.* Rifampicin-induced transcriptome response in rifampicin-resistant

Mycobacterium tuberculosis. *Tuberculosis* **93**, 96–101 (2013).

13. Waddell, S. J. *et al.* The use of microarray analysis to determine the gene expression

profiles of *Mycobacterium tuberculosis* in response to anti-bacterial compounds.

Tuberculosis **84**, 263–274 (2004).

14. Boshoff, H. I. M. *et al.* The transcriptional responses of *Mycobacterium tuberculosis* to

inhibitors of metabolism: Novel insights into drug mechanisms of action. *J. Biol. Chem.*

279, 40174–40184 (2004).

15. Wilson, M. *et al.* Exploring drug-induced alterations in gene expression in *Mycobacterium*

tuberculosis by microarray hybridization. *Proc. Natl. Acad. Sci. U. S. A.* **96**, 12833–12838

(1999).

16. Deb, C. *et al.* A novel in vitro multiple-stress dormancy model for *Mycobacterium*

tuberculosis generates a lipid-loaded, drug-tolerant, dormant pathogen. *PLoS One* **4**,

e6077 (2009).

17. Betts, J. C., Lukey, P. T., Robb, L. C., McAdam, R. A. & Duncan, K. Evaluation of a

nutrient starvation model of *Mycobacterium tuberculosis* persistence by gene and protein

expression profiling. *Mol. Microbiol.* **43**, 717–731 (2002).

18. Keren, I., Minami, S., Rubin, E. & Lewis, K. Characterization and transcriptome analysis

of *Mycobacterium tuberculosis* persisters. *MBio* **2**, (2011).

19. Honeyborne, I. *et al.* Profiling persistent tubercule bacilli from patient sputa during

therapy predicts early drug efficacy. *BMC Med.* **14**, 1–13 (2016).

20.   Sharma, S. *et al.* Transcriptome analysis of mycobacteria in sputum samples of pulmonary

tuberculosis patients. *PLoS One* **12**, e0173508 (2017).

21.   Gautam, U. S., Mehra, S. & Kaushal, D. In-vivo gene signatures of Mycobacterium

tuberculosis in C3HeB/FeJ mice. *PLoS One* **10**, e0135208 (2015).

22.   Lai, R. P. J. *et al.* Transcriptomic characterization of tuberculous sputum reveals a host

Warburg effect and microbial cholesterol catabolism. *MBio* **12**, (2021).

23.   Garton, N. J. *et al.* Cytological and transcript analyses reveal fat and lazy persister-like

bacilli in tuberculous sputum. *PLoS Med.* **5**, 0634–0645 (2008).

24.   Talaat, A. M., Lyons, R., Howard, S. T. & Johnston, S. A. The temporal expression profile

of *Mycobacterium tuberculosis* infection in mice. *Proc. Natl. Acad. Sci. U. S. A.* **101**,

4602–4607 (2004).

25.   Rockwood, N., Lai, R. P. J., Seldon, R., Young, D. B. & Wilkinson, R. J. Variation in pre-

therapy levels of selected *Mycobacterium tuberculosis* transcripts in sputum and their

relationship with 2-month culture conversion. *Wellcome Open Res.* **4**, 106 (2019).

26.   Pisu, D., Huang, L., Grenier, J. K. & Russell, D. G. Dual RNA-seq of *Mtb*-infected

macrophages in vivo reveals ontologically distinct host-pathogen interactions. *Cell Rep.*

30, 335-350.e4 (2020).

27.   Hudock, T. A. *et al.* Hypoxia sensing and persistence genes are expressed during the

intragranulomatous survival of *Mycobacterium tuberculosis*. *Am. J. Respir. Cell Mol. Biol.*

56, 637–647 (2017).

28.   Karakousis, P. C., Williams, E. P. & Bishai, W. R. Altered expression of isoniazid-

regulated genes in drug-treated dormant *Mycobacterium tuberculosis*. *J. Antimicrob.*

Chemother. **61**, 323–331 (2008).

29.   Bukka, A., Price, C. T. D., Kernodle, D. S. & Graham, J. E. *Mycobacterium tuberculosis*

RNA expression patterns in sputum bacteria indicate secreted Esx factors contributing to

growth are highly expressed in active disease. *Front. Microbiol.* **2**, (2012).

30.   Gautam, U. S. *et al.* DosS is required for the complete virulence of *Mycobacterium*

tuberculosis in mice with classical granulomatous lesions. *Am. J. Respir. Cell Mol. Biol.*

52, 708–716 (2015).

31.   Coppola, M. *et al.* New genome-wide algorithm identifies novel in-vivo expressed

Mycobacterium tuberculosis antigens inducing human t-cell responses with classical and

unconventional cytokine profiles. *Sci. Rep.* **6**, (2016).

32.   Garcia, B. J. *et al.* Sputum is a surrogate for bronchoalveolar lavage for monitoring

Mycobacterium tuberculosis transcriptional profiles in TB patients. *Tuberculosis* **100**, 89–

94 (2016).

33.   Walter, N. D. *et al.* Adaptation of *Mycobacterium tuberculosis* to impaired host immunity

in HIV-infected patients. *J. Infect. Dis.* **214**, 1205–1211 (2016).

34.   Skvortsov, T. A., Ignatov, D. V., Majorov, K. B., Apt, A. S. & Azhikina, T. L.

Mycobacterium tuberculosis transcriptome profiling in mice with genetically different

susceptibility to tuberculosis. *Acta Naturae* **5**, 62–69 (2013).

35.   Shaikh, A. *et al.* Early phase of effective treatment induces distinct transcriptional changes

in *Mycobacterium tuberculosis* expelled by pulmonary tuberculosis patients. *Sci. Rep.* **11**,

1–13 (2021).

36.   Cornejo-Granados, F. *et al.* Targeted RNA-seq reveals the *M. tuberculosis transcriptome*

from an in vivo infection model. *Biology (Basel).* **10**, 848 (2021).

37.   Betin, V. *et al.* Hybridization-based capture of pathogen mRNA enables paired host-

pathogen transcriptional analysis. *Sci. Rep.* **9**, (2019).

38.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

39.   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: A Bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–

140 (2010).

40.   McCarthy, D. J., Chen, Y. & Smyth, G. K. Differential expression analysis of multifactor

RNA-Seq experiments with respect to biological variation. *Nucleic Acids Res.* **40**, 4288–

4297 (2012).

41.   Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: A practical and

powerful approach to multiple testing. *J. R. Stat. Soc. Ser. B* **57**, 289–300 (1995).

42.   Murtagh, F. & Legendre, P. Ward's hierarchical agglomerative clustering method: Which

algorithms implement Ward's criterion? *J. Classif.* **31**, 274–295 (2014).

43.   Federico, A. & Monti, S. HypeR: An R package for geneset enrichment workflows.

Bioinformatics **36**, 1307–1308 (2020).

44.   Cole, S. T. *et al.* Erratum: Deciphering the biology of Mycobacterium tuberculosis from

the complete genome sequence. *Nature* **396**, 190 (1998).

45.   R Core Team. R: A language and environment for statistical computing. (2021).

46.   Chang, W. *et al.* shiny: Web application framework for R. (2022).

47.   Kundu, M. & Basu, J. Applications of transcriptomics and proteomics for understanding

dormancy and resuscitation in *Mycobacterium tuberculosis*. *Frontiers in Microbiology* **12**,

642487 (2021).

48.   Voskuil, M. I. *et al.* Inhibition of respiration by nitric oxide induces a *Mycobacterium*

tuberculosis dormancy program. *J. Exp. Med.* **198**, 705–713 (2003).

49.   Voskuil, M. I., Bartek, I. L., Visconti, K. & Schoolnik, G. K. The response of

Mycobacterium tuberculosis to reactive oxygen and nitrogen species. *Front. Microbiol.* **2**,

105 (2011).

50.   Kendall, S. L. *et al.* Cholesterol utilization in mycobacteria is controlled by two TetR-type

transcriptional regulators: *kstR* and *kstR2*. *Microbiology* **156**, 1362–1371 (2010).

51.   Reddy, P. V. *et al.* Disruption of mycobactin biosynthesis leads to attenuation of

Mycobacterium tuberculosis for growth and virulence. *J. Infect. Dis.* **208**, 1255–1265

(2013).

52.   Nandakumar, M., Nathan, C. & Rhee, K. Y. Isocitrate lyase mediates broad antibiotic

tolerance in *Mycobacterium tuberculosis*. *Nat. Commun.* **5**, (2014).

53.   Sirakova, T. D. *et al.* Identification of a diacylglycerol acyltransferase gene involved in

accumulation of triacylglycerol in *Mycobacterium tuberculosis* under stress. *Microbiology*

152, 2717–2725 (2006).

54.   Gourse, R. L., Gaal, T., Bartlett, M. S., Appleman, J. A. & Ross, W. rRNA transcription

and growth rate-dependent regulation of ribosome synthesis in *Escherichia coli*. *Annual*

Review of Microbiology **50**, 645–677 (1996).

55.   Maitra, A. & Dill, K. A. Bacterial growth laws reflect the evolutionary importance of

energy efficiency. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 406–411 (2015).

56.   Mondino, S., Gago, G. & Gramajo, H. Transcriptional regulation of fatty acid biosynthesis

in mycobacteria. *Mol. Microbiol.* **89**, 372–387 (2013).

57.   Fullam, E., Prokes, I., Fütterer, K. & Besra, G. S. Structural and functional analysis of the

solute-binding protein UspC from *Mycobacterium tuberculosis* that is specific for amino

sugars. *Open Biol.* **6**, (2016).

58.   Yang, Y. *et al.* A hydrolase of trehalose dimycolate induces nutrient influx and stress

sensitivity to balance intracellular growth of *Mycobacterium tuberculosis. Cell Host*

Microbe **15**, 153–163 (2014).

59.   Kalscheuer, R., Weinrick, B., Veeraraghavan, U., Besra, G. S. & Jacobs, W. R. Trehalose-

recycling ABC transporter LpqY-SugA-SugB-SugC is essential for virulence of

Mycobacterium tuberculosis. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 21761–21766 (2010).

60.   Mascolo, L. & Bald, D. Cytochrome bd in *Mycobacterium tuberculosis*: A respiratory

chain protein involved in the defense against antibacterials. *Progress in Biophysics and*

Molecular Biology **152**, 55–63 (2020).

61.   Fenn, J. S. *et al.* Structural basis of glycerophosphodiester recognition by the

Mycobacterium tuberculosis substrate-binding protein ugpb. *ACS Chem. Biol.* **14**, 1879–

1887 (2019).

62.   Karbalaei Zadeh Babaki, M., Soleimanpour, S. & Rezaee, S. A. Antigen 85 complex as a

powerful Mycobacterium tuberculosis immunogene: Biology, immune-pathogenicity,

applications in diagnosis, and vaccine design. *Microbial Pathogenesis* **112**, 20–29 (2017).

63.   Gröschel, M. I., Sayes, F., Simeone, R., Majlessi, L. & Brosch, R. ESX secretion systems:

Mycobacterial evolution to counter host immunity. *Nature Reviews Microbiology* **14**, 677–691 (2016).

64. Remm, S., Earp, J. C., Dick, T., Dartois, V. & Seeger, M. A. Critical discussion on drug efflux in *Mycobacterium tuberculosis*. *FEMS Microbiology Reviews* **46**, 1–15 (2022).

65. Shinde, Y., Ahmad, I., Surana, S. & Patel, H. The Mur enzymes chink in the armour of *Mycobacterium tuberculosis* cell wall. *European Journal of Medicinal Chemistry* **222**, 113568 (2021).

66. Huszár, S. *et al.* N-acetylglucosamine-1-phosphate transferase, WecA, as a validated drug target in *Mycobacterium tuberculosis*. *Antimicrob. Agents Chemother.* **61**, (2017).

67. Voskuil, M. I. *Mycobacterium tuberculosis* gene expression during environmental conditions associated with latency. *Tuberculosis* **84**, (2004).

68. Gould, T. A., Van De Langemheen, H., Muñoz-Elías, E. J., McKinney, J. D. & Sacchettini, J. C. Dual role of isocitrate lyase 1 in the glyoxylate and methylcitrate cycles in *Mycobacterium tuberculosis*. *Mol. Microbiol.* **61**, 940–947 (2006).

69. McKinney, J. D. *et al.* Persistence of *Mycobacterium tuberculosis* in macrophages and mice requires the glyoxylate shunt enzyme isocitrate lyase. *Nature* **406**, 735–738 (2000).

70. Muñoz-Elías, E. J. & McKinney, J. D. Mycobacterium tuberculosis isocitrate lyases 1 and 2 are jointly required for in vivo growth and virulence. *Nat. Med.* **11**, 638–644 (2005).

71. Shetye, G. S., Franzblau, S. G. & Cho, S. New tuberculosis drug targets, their inhibitors, and potential therapeutic impact. *Translational Research* **220**, 68–97 (2020).

72. Lee, B. S., Sviriaeva, E. & Pethe, K. Targeting the cytochrome oxidases for drug development in mycobacteria. *Progress in Biophysics and Molecular Biology* **152**, 45–54

(2020).

73. Gumbo, T., Lenaerts, A. J., Hanna, D., Romero, K. & Nuermberger, E. Nonclinical

models for antituberculosis drug development: A landscape analysis. *J. Infect. Dis.* **211**,

S83–S95 (2015).

74. Dide-Agossou, C. *et al.* Combination of *Mycobacterium tuberculosis* RS ratio and CFU

improves the ability of murine efficacy experiments to distinguish between drug

treatments. *Antimicrob. Agents Chemother.* **66**, (2022).

75. Ma, S. *et al.* Transcriptomic signatures predict regulators of drug synergy and clinical

regimen efficacy against tuberculosis. *MBio* **10**, (2019).

76. Maitra, A. *et al.* Cell wall peptidoglycan in *Mycobacterium tuberculosis*: An Achilles'

heel for the TB-causing pathogen. *FEMS Microbiology Reviews* **43**, 548–575 (2019).

77. Pawełczyk, J. *et al.* Cholesterol-dependent transcriptome remodeling reveals new insight

into the contribution of cholesterol to *Mycobacterium tuberculosis* pathogenesis. *Sci. Rep.*

11, 1–16 (2021).

FIGURES

Fig. 1. **Visual summary of methods previously used to quantify the *Mtb* transcriptome *in vivo*.** Each horizontal arrow represents a distinct combination of enrichment and quantification. Varying enrichment methods are represented via the symbols shown in the key. SEARCH-TB is a unique combination of enrichment (eukaryotic cell lysis + targeted amplification) followed by quantification via RNA-seq that has enabled transcriptome evaluation in mice treated for weeks with a potent combination regimen. Image created with Biorender.com.

Fig. 2. Evaluation of SEARCH-TB platform. a. Distribution of log₂ fold differences of normalized gDNA SEARCH-TB expression data relative to the value expected if there were no amplification bias. Zero (dashed vertical line) represents no amplification bias. **b.** Evaluation of repeatability of SEARCH-TB showing normalized expression data (log counts per million) for two technical replicates in which *Mtb* RNA was spiked into human lung RNA. **c-d.** Volcano plot showing log₂ fold-changes and -log₁₀ *p*-values induced by *in vitro* INH exposure as quantified by RNA-seq (**c**) and SEARCH-TB (**d**). Genes significantly down- and upregulated with INH exposure relative to control (adj. p-value< 0.05) are shown in blue and red. **e.** Comparison of differential expression between INH treated samples and control samples from RNA-seq or SEARCH-TB data. Purple shading indicates genes with concordant fold-change direction and significance between RNA-seq and SEARCH-TB. green shading indicates genes that were significant in RNA-seq or SEARCH-TB results, but not both. Gold shading indicates genes that were significant for both RNA-seq and SEARCH-TB, but in opposite directions. Gray shading indicates genes that were not significantly differentially expressed in either RNA-seq or SEARCH-TB. **f.** Comparison of INH vs. control fold-changes from RNA-seq data vs. SEARCH-TB data. Purple, green, gold and gray colors have the same meaning as in **e**.

Fig. 3. Overview of transcriptional response to HRZE in mice and *in vitro* experiments. a. Principal components plot including all mouse (triangles) and *in vitro* (squares) samples. Time points are shown by color. **b.** Volcano plot summarizing the differential expression between *Mtb* in mice and *in vitro* prior to treatment. Genes significantly down- (blue) or upregulated (red) in mice relative to *in vitro* (adj-P-value< 0.05) are shown. **c.** *Mtb* CFU and RS ratio values for control mouse samples and 14- or 28- days after HRZE treatment initiation. **d-e.** Volcano plots summarizing the differential expression between *Mtb* in 14-day HRZE treated and control mouse samples (**d**) and 28-day HRZE treated and control mouse samples (**e**). **f.** Estimated gene expression over time in mice. Genes that were significantly differentially expressed between at least two treatment time points are shown (N=2,429). Values are row-scaled, with red and blue indicating higher and lower expression, respectively. Hierarchical clustering of genes identified four broad patterns. **g.** Average $\log_2$ fold change for each of the four clusters relative to control. Values above and below zero represent up- and downregulation relative to control, respectively. **h.** Comparison of differential expression between mouse (day 28) or *in vitro* (day 8) relative to respective controls. Purple shading indicates genes with concordant fold-change direction and significance between mouse and *in vitro* experiments. Green shading indicates genes that were significant for either mouse or *in vitro* experiments but not both. Gold shading indicates genes that were significant for both mouse and *in vitro* experiments but in opposite directions. Gray shading indicates the genes that were not differentially expressed with HRZE treatment either in mouse or *in vitro* experiments. **i.** Comparison of fold-changes between mouse (day 28) or *in vitro* (day 8) relative to respective controls. Purple, green, gold, and gray colors have the same meaning as in **h**.

Fig. 4 Summary of gene set enrichment and transcriptional changes in biological processes. a. Gene categories significantly enriched for genes differentially expressed between day 28 and control murine samples. The percentage of genes in each category significantly up- (red) or down- (blue) regulated for each comparison is illustrated. Asterisks indicate statistical significance (adj-*P*<0.05). **b.** Fold-change of ribosomal protein genes in mice at day 28 (left) and *in vitro* (right) at day 8, relative to control. Red bars indicate the four alternative C-ribosomal protein paralogs. **c, e-g** Fold-change values in mice at days 14 and 28 (left) and *in vitro* (right) at days 4 and 8, relative to control for FAS-II (**c**), ATP synthetase (**e**), cytochrome bcc/aa3 supercomplex and the bd oxidase (**f**), and glyoxylate bypass (**g**) gene sets. **d.** Graphical representation of changes in the peptidoglycan synthesis, modification, and recycling pathways in mice at day 28 relative to control. Log$_2$ fold-change values for genes in the process are indicated by the color of each box and the bolded outlines of boxes represent genes that are significantly differentially expressed. Figure adapted from Maitra et al., 2019.[76] Image created with Biorender.com.

Fig. 5 Summary of transcriptional changes in biological processes. a. Log$_2$ fold-change values for genes in the cholesterol degradation pathway for mice at day 28 and *in vitro* at day 8, relative to control. Deeper red values represent higher upregulation of genes after drug treatment relative to control, while deeper blue values represent higher downregulation. Figure adapted from Pawełczyk et al., 2021[77]. **b-g.** Fold-change in mice (left) and *in vitro* (right), relative to control for heat shock proteins (red bar represents *hspX*) (**b**), dosR (**c**), antigen 85 (**d**), ESX-1 (red bar represents *esxA*, blue bar represents *esxB*) (**e**), the Rv2686c-2688c efflux pump (**f**), and the DrrABC efflux pump (**g**) gene sets.