Comparative transcriptomics reveals divergence in pathogen response gene families amongst twenty forest tree species

[1*]Mengmeng Lu, [2]Min Cao, [2,3]Jie Yang, [1]Nathan G. Swenson

[1] Department of Biological Sciences, University of Notre Dame, 100 Galvin Life Sciences, Notre Dame, IN 46556, USA

[2] CAS Key Laboratory of Tropical Forest Ecology, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Mengla, Yunnan 666303, China.

[3] Center of Plant Ecology, Core Botanical Gardens, Chinese Academy of Sciences, Mengla, Yunnan 666303, China.

[*]Corresponding author: Department of Biological Sciences, University of Notre Dame, 100 Galvin Life Sciences, Notre Dame, IN 46556, USA. Email: mlu@nd.edu

21   **Abstract**
22
Forest trees provide critical ecosystem services for humanity that are under threat due to ongoing
global change. Measuring and characterizing genetic diversity is key to understanding adaptive
potential and developing strategies to mitigate negative consequences arising from climate
change. In the area of forest genetic diversity, genetic divergence caused by large-scale changes
at the chromosomal level has been largely understudied. In this study, we used the RNA-seq data
of twenty co-occurring forest trees species from genera including *Acer*, *Alnus*, *Amelanchier*,
Betula, *Cornus*, *Corylus*, *Dirca*, *Fraxinus*, *Ostrya*, *Populus*, *Prunus*, *Quercus*, *Ribes*, *Tilia*, and
Ulmus sampled from Upper Peninsula of Michigan. These data were used to infer the origin and
maintenance of gene family variation, species divergence time, as well as gene family expansion
and contraction. We identified a signal of common whole genome duplication events shared by
core eudicots. We also found rapid evolution, namely fast expansion or fast contraction of gene
families, in plant-pathogen interaction genes amongst the studied diploid species. Finally, the
results lay the foundation for further research on the genetic diversity and adaptive capacity of
forest trees, which will inform forest management and conservation policies.
37
38   **Keywords:** angiosperms,gene duplication,gene family expansion and contraction, Ks plot,
39   Ka/Ks, plant-pathogen interaction

Introduction

Forest trees occupy nearly a third of the Earth's land surface and provide key ecological benefits and services such as water, air purification and carbon sequestration (FAO 2020; Gamfeldt et al. 2013; Miura et al. 2015). They are critical components of terrestrial biodiversity in terms of their own diversity, but also with respect to the habitats and renewable materials such as wood, cellulose, and lignin (Cortés et al. 2020). It is estimated that there are ~73,000 tree species globally with most of being rare and regionally endemic, implying vulnerability of global tree species to anthropogenic and climate changes (Cazzolla Gatti et al. 2022). Indeed, climate change has already impacted forests and their species composition via range migration, wildfires, droughts, pest, and pathogens (Allen et al. 2010; Hammond et al. 2022; Kirilenko and Sedjo 2007; Rizvi et al. 2015). The genetic composition and diversity of the tree species in forests is foundational not only to our understanding of the abiotic and biotic factors governing the distribution and dynamics of species populations, but also to how they will respond to current and future environmental change. Thus, measuring and characterizing intra- and inter-specific genetic diversity is the key for understanding the current structure and dynamics of forests and their adaptive potential and for developing strategies to mitigate the damages brought along with environmental change.

Gene duplication and gene loss are major sources of genetic diversity. Gene duplication, generated through either whole genome duplication (WGD) or small-scale duplication (SSD), is the primary source of new genetic material among flowering plants (Cui et al. 2006; Panchy et al. 2016). WGD contributes to the majority of duplicate genes (Amborella Genome Project 2013; Soltis and Soltis 2016). The ancient WGDs can lead to competitive advantage, increased species diversity, and evolutionary innovations (Van de Peer et al. 2009). Most flowering species have a polyploid history either through autopolyploidy or allopolyploidy. Specifically, flowering species are either currently polyploid, or have a polyploid ancestor and have reverted to being diploid (Blanc and Wolfe 2004). Earlier studies placed at least three WGD events during the radiation of angiosperms, the oldest WGD occurred before the monocot-eudicot divergence, a second WGD occurred specifically to eudicots, a more recent WGD, gamma (γ) triplication event, occurred before the separation of rosids and asterids (Jiao et al. 2012; Tang et al. 2008). Alternatively, pervasive SSD, including tandem duplication (Zhang 2003), transposon-mediated duplication (Jiang et al. 2004), segmental duplication (Bailey et al. 2002), and retroduplication (Brosius 1991), also generates a great number of duplicate genes. While the majority of duplicate genes end up being silenced and eliminated (Lynch and Conery 2000), a few preserved duplicate genes may be retained by selection due to neo-functionalization, relative dosage, absolute dosage or sub- functionalization (Conant et al. 2014), and play important role in generating genetic diversity and building up reproductive isolation barriers. Recent studies explained retention of duplicate genes as some paralogous genes depend on or compensate for their cognate copy (Diss et al. 2017), while some paralogous gene pairs are stuck in a functional and structural entanglement status over evolutionary time scale (Kuzmin et al. 2020).

Compared to gene duplication, the mechanism that underlie gene loss are not yet well understood. It is commonly assumed that gene loss is realized by deletion or pseudogenization, as a result of neutral or adaptive processes (Demuth and Hahn 2009). Most nonsense mutations are subject to drift to fixation. Some deleterious mutations could be accumulated and eventually become

86 silenced (Johri et al. 2022). Gene loss might be selected to retain dosage balance, as evidenced in
87 a set of single-copy genes with essential housekeeping functions, which are typically restored to
88 become singletons after WGD or SSD (De Smet et al. 2013).
89
Differential duplication and loss of genes among evolutionary lineages results in gene family
expansion and contraction (Demuth and Hahn 2009). Gene families comprise a set of similar
nucleotide or amino acid sequences. They arise as a result of duplication events and have similar
cellular and biochemical functions. Gene family expansions and contractions related to metabolic,
regulatory and signalling gene networks contribute to the adaptations of plants to various abiotic
habitats and biotic interactions (One Thousand Plant Transcriptomes Initiative 2019). They are
sources of plant functional evolution and innovation (Force et al. 1999; Ohno 1970), exhibiting a
clear functional basis, which are influenced by the mechanisms of duplication (Hanada et al.
2008; Maere et al. 2005). Comparisons of gene family content across species, therefore, may
provide insight into the ecological interactions and evolutionary pressures that have shaped
adaptation and diversity. This is particularly true when we conduct comparative transcriptomic
analyses of species that currently co-occur and that, potentially have co-occurred for long periods
of time (Swenson and Jones 2017).
103
Genetic investigations of forest trees have, generally, focused on nucleotide substitutions in
specific genes (Isabel et al. 2020; Neale and Kremer 2011), while comparative analyses of
genetic divergence caused by large-scale changes at the chromosomal levels have been largely
understudied. However, the magnitude, rate, and distribution of changes in gene family size are
key drivers of the functional differentiation and adaptation of species (Demuth and Hahn 2009;
Lynch and Conery 2003). Large segmental duplication and changes in copy number are known
to be associated with disease response and environmental adaptation (Bailey et al. 2002; Reid et
al. 2016). Therefore, gene family size change is as much important as nucleotide substitution to
understanding genetic divergence and evolutionary forces responsible for these changes.
113
Such knowledge on genetic divergence and associated evolutionary forces in forest trees can
help to promote breeding healthy trees (i.e. disease resistant trees) in a changing climate (Allen
et al. 2010; Dodds and Rathjen 2010). The increasing occurrences and severity of insect and
disease outbreaks have threatened forest trees (Hurley et al. 2016; Linnakoski and Forbes 2019),
but we lack a thorough understanding to the disease related genes in forest trees. Trouern-Trend
et al. (2020) revealed contraction in disease associated gene families in *Juglans* species. Lu et al.
(2021) reported differential gene expression patterns between tolerant and susceptible lodgepole
pine trees to a fungal pathogen. It would be interesting to explore the potential link between
duplication events and disease response in expressed genes of forest trees, as the duplicate genes
can quickly lead to changes in gene expression with the acquisition of increased resistance to
different pathogens (Picart-Picolo et al. 2020).
125
In this study, we used the RNA-seq data for twenty co-occurring forest trees species to infer the
origin and maintenance of gene family variation as well as species divergence time, gene
expansion and contraction. We sought to address the following questions: What is the magnitude
of gene duplication? What factors impact duplicate gene evolution? Are the gene gain and loss
process random or controlled by natural selection? The answers to these questions are the key to

understanding the basis of adaptive divergence in forest tree species and will help to predict
adaptive capacity of these species in a changing climate.
133
134
Materials and methods
Sample collection
Leaves of each of twenty forest species were sampled in August, 2021, at the University of Notre
Dame Environmental Research Center (UNDERC) near Land O' Lakes, Wisconsin, U.S.A. (Fig.
1). The UNDERC site is located at latitude: 46.23391, longitude: -89.537254, and encompasses
approximately ~8,000 acres of mixed deciduous forest on both sides of the state line between
Wisconsin and Michigan's upper peninsula. The twenty species used in this study are dominate
components of the woody vegetation at the site and are: *Acer rubrum* (red maple), *Acer*
saccharum (sugar maple), *Acer spicatum* (mountain maple), *Alnus incana* (Gary alder),
Amelanchier laevis (smooth serviceberry), *Betula alleghaniensis* (yellow birch), *Betula*
papyrifera (paper birch), *Cornus alternifolia* (pagoda dogwood), *Corylus cornuta* (beaked
hazelnut), *Dirca palustris* (Eastern leatherwood), *Fraxinus nigra* (black ash), *Ostrya virginiana*
(American hophornbeam), *Populus grandidentata* (bigtooth aspen), *Populus tremuloides*
(quaking aspen), *Prunus serotina* (black cherry), *Prunus virginiana* (chokecherry), *Quercus*
rubra (northern red oak), *Ribes cynosbati* (prickly gooseberry), *Tilia americana* (American
basswood), *Ulmus americana* (American elm). They are all angiosperms (flowering plants) and
eudicots, belonging to the clades of asterids, saxifragales, and rosids. These species were chosen
as they are the most dominant at the study site and in large national forests in region (i.e. Ottawa,
Nicolet, and Hiawatha National Forests). Specifically, these species compose >95% of the tree
individuals and above-ground biomass in these forests (Swenson et al. 2017). We focus on the
most dominant species in order to provide insights into the ecological and evolutionary processes
driving transcriptomic variation within and across populations and communities (Swenson and
Jones 2017). Upon collection, the samples were flash frozen in liquid nitrogen, then transported
and stored in an ultracold freezer at the field station.
159
RNA-sequencing, de novo transcriptome assembly, and annotation
The samples were sent to Novogene corporation inc. (Chula Vista, CA, USA) for mRNA
extraction, library preparation, mRNA sequencing, and quality control. The samples were
sequenced on an Illumina NovaSeq 6000 platform to produce 150 bp paired-end reads. Raw
reads filtering was performed by Novogene to get the clean reads including removing reads
containing adapters, removing reads containing N >10%, and removing reads containing low
quality base (Phred quality score ≤5) which is over 50% of the total base. An average of 6 Gb
paired-end clean reads were acquired for each sample and deposited in NCBI Sequence Read
Archive (SRA accession number: PRJNA896247). *De novo* transcriptome assembly was
performed for each sample using the software TransLiG v1.3 with default parameters (Liu et al.
2019). The clean reads were then mapped to the assembled transcriptome using the software
RSEM v1.2.25 with default parameters (Li and Dewey 2011). The longest transcript (isoform)
with a transcript per million (TPM) value ≥1.5 for each gene was retained. Redundancy was
removed from the retained transcripts using the software CD-HIT v4.8.1 with parameters of -c
0.95 -n 8 -T 32 -M 1540 (Fu et al. 2012). To remove potential contamination, the retained
transcripts were aligned against the non-redundant (nr) protein database (downloaded on January
22, 2022) using the blastx function implemented by the software DIAMOND v2.0.13.151

(Buchfink et al. 2021) with parameter of --outfmt 102 to assign the best known taxon ID to each
query transcript. The file that was used to convert accession numbers to taxonomy was
downloaded from https://ftp.ncbi.nlm.nih.gov/pub/taxonomy/accession2taxid/, and the taxon ID
file was downloaded from https://ftp.ncbi.nlm.nih.gov/pub/taxonomy/new_taxdump/. The query
transcripts that could be assigned to green plant (Viridiplantae) taxon IDs were retained. The
filtered transcripts were aligned to the nr database that only includes green plant subjects using
the blastx function implemented by the software DIAMOND v2.0.13.151 with default
parameters. The aligned transcripts were combined with those transcripts assigned with green
plant taxon IDs to form the final transcriptomes. The completeness of the final transcriptomes
was examined using the 1,614 orthologues in the BUSCO (Benchmarking Universal Single Copy
Orthologs) v5.2.2 set of embryophyta_odb10 (Manni et al. 2021).
188
The transcriptomes were annotated with Gene Ontology (GO) and Kyoto Encyclopedia of Genes
and Genomes (KEGG) terms. The transcripts were aligned to the nr database that only includes
green plant subjects using the blastx function implemented by the software DIAMOND
v2.0.13.151 with parameters of -taxonlist 33090 --very-sensitive -f5. The .xml outputs were used
as input for mapping and annotation using the software Blast2GO (Götz et al. 2008). The
transcripts were translated to protein sequences using the software TransDecoder v5.5.0
(https://github.com/TransDecoder/TransDecoder/wiki) with default parameters to get the longest
open reading frame (ORF) per transcript, and then were annotated by the software InterProScan
v5.54-87.0 (Jones et al. 2014) with the parameters of -dra -f XML -goterms -pa. The final GO
terms were acquired by merging Blast2GO and InterProScan annotation outputs using the
software Blast2GO. To get the KEGG terms, the longest ORF per transcript were also annotated
using EggNOG-mapper v2.1.7 (Cantalapiedra et al. 2021) with parameters of -m diamond -
sensmode more-sensitive -tax_scope 33090 -tax_scope_mode 33090. The assembled
transcriptome and annotation were used in the downstream analyses as shown in the flowchart
(Supplementary Fig. S1).
204
Ks analyses
The level of synonymous substitutions (Ks) between two homologous sequences increases
approximately linearly with time. Thus pairs of paralogous genes within a species can be sorted
to estimate the relative ages of gene duplication (Blanc and Wolfe 2004). To identify paralogous
gene pairs within a species, an all-versus-all blastp was performed within each species. The
peptide sequences (the longest ORF per transcript) of each species were used as input to run all-
versus-all blastp using the software DIAMOND v2.0.13.151 with parameters of -p 16 --max-
target-seqs 20 --evalue 1e-10 --outfmt 6 qseqid sseqid pident nident evalue. The blastp output
was filtered by removing self blastp results, keeping outputs with pident (percent of identical
matches) ≥20 and nident (number of identical matches) ≥50, and removing sequences with more
than 10 hits. The filtered paralogous gene pairs were aligned using the software MAFFT v7.490
(Katoh et al. 2002) with default parameters. The alignments were used as input for estimating the
number of synonymous substitutions per synonymous site (Ks) and the number of non-
synonymous substitutions per nonsynonymous site (Ka) using the program yn00 in the software
PAML v4.9j (Yang 2007) with parameters of icode = 0 weighting = 0 commonf3×4 = 0. The Ks
of each paralogous gene pairs were acquired using the method by Yang and Nielsen (2000).
221

Ks can also be calculated using the orthologous gene pairs between species to estimate speciation
events (Blanc and Wolfe 2004). To identify orthologous gene pairs between species, reciprocal
best hit blastp were run between the compared pairs. The peptide sequences (the longest ORF per
transcript) of the paired species were used as input to run blastp using the software DIAMOND
v2.0.13.151 with parameters of -p 16 --max-target-seqs 20 --evalue 1e-10 --outfmt 6 qseqid
sseqid pident nident evalue. The best hits between forward and reverse blastp for each compared
sequence pairs were retained, which were then used for sequence alignment and estimating Ka
and Ks as abovementioned.
230
The Ks distribution is proportional to the time lapse. Thus, the Ks distribution within species can
be used to infer the timing of WGD, and the Ks distribution between two different species can
infer time since speciation (Blanc and Wolfe 2004). To plot the Ks distribution, the mixture
model was fit and plot for the Ks values ≥ 0.01 and ≤ 2 using the R package mixR (Yu 2022).
The best model from a candidate of mixture models was selected using the "select" function
based on the information criterion BIC.
237
Estimating duplication events and divergence time
To infer duplication events, the species tree and gene trees were first inferred using STAG and
STRIDE algorithms (Emms and Kelly 2017; 2018) implemented by the program OrthoFinder
v2.5.4 (Emms and Kelly 2019). The inferred species tree was similar to that expected given
current knowledge of the Angiosperm phylogeny with the exception of the placement of *Populus*
tremuloides and *P. grandidentata*. Thus, for downstream analysis and discussion, we removed
these two species for the species tree inference. The inferred species tree and gene trees were
used to estimate duplication events and divergence time. The gene duplication events on the
species tree node were reported if, at least 50% of the descendant species have retained both
copies of the duplicate gene. The divergence time was estimated using the software r8s v1.81
(Sanderson 2003) with calibration points of 115 million years (MYA) between *Fraxinus nigra*
and *Corylus cornuta*, which were obtained in November, 2022 from http://www.timetree.org/
(Kumar et al. 2017).
251
Gene families and their evolutionary rate
To infer gene families, the peptide sequences (the longest ORF per transcript) of the twenty
studied transcriptomes were used to construct orthogroups (gene families) using the program
OrthoFinder v2.5.4 with parameters of -S diamond -t 16 -a 16. Genes assigned to each gene
family were counted for each species. The evolutionary rate of protein coding genes was
represented by the ratio of nonsynonymous substitutions per nonsynonymous site to synonymous
substitutions per synonymous site (Ka/Ks) with *Dillenia indica* (elephant apple) being the
outgroup and the twenty studied species being the ingroup. *D. indica* is a eudicot species,
belonging to the order Dilleniales and the RNA-seq data for this species were downloaded from
https://trace.ncbi.nlm.nih.gov/Traces/index.html?view=run_browser&acc=ERR2040190&disp
lay=data-access (One Thousand Plant Transcriptomes Initiative 2019). The raw sequences were
filtered using the program fastp with default parameters (Chen et al. 2018). The transcriptome
assembly and protein sequence prediction were performed as abovementioned, yielding 22,753
transcripts and 19,345 protein sequences. Then the reciprocal best hit blastp was performed
between *D. indica* and each of the twenty studied species to identify the orthologous pairs. The
best hits between forward and reverse blastp for each compared sequence pairs were retained,

which were then used for sequence alignment and estimating Ka/Ks as abovementioned. The
mean Ka/Ks across genes for each gene family per species was used to represent the
evolutionary rate for each gene family. Gene families with a mean Ka/Ks >10 were filtered out.
The graphs were plotted using the R package ggplot2 and ggpubr (Kassambara 2023; Wickham
2016). The size and evolutionary rate of gene families were correlated using scatter plot. All
statistical analyses were performed in R (R Core Team 2021). Gene families with a size ≥5 and
Ka/Ks >1 were highlighted for further analysis. The enriched GO and KEGG terms were
identified in these highlighted gene families. The enriched GO terms of each category (biological
process, molecular function, cellular component) were separately detected using the "weight"
algorithm with the Fisher ratio test implemented by the R package "topGO" (Alexa and
Rahnenfuhrer 2023). Enriched KEGG terms were detected using the hypergeometric method
implemented by the R package "plyr" (Wickham 2011). Heatmaps were plot using the R
package pheatmap (Kolde 2019).
281
Estimating gene family expansion and contraction
To further study gene family changes, species tree and gene family counts were predicted and
used to estimate gene family expansion and contraction. All-versus-all blastp was performed to
find the most similar sequences for each sequence in the studied species using the program
DIAMOND v2.0.13.151. The program MCL (Enright et al. 2002) was used to find clusters of
similar sequences (gene families). The species tree was estimated using the program FastTree
(Price et al. 2009). The programs DIAMOND, MCL, and FastTree were implemented by the
program OrthoFinder v2.5.4, default parameters were used. Then, the species tree was converted
to a format for estimating ultrametric tree using the program CAFE5 (Mendes et al. 2021) with
the command python prep_r8s.py -s 60164 -p 'Fraxinus_nigra,Corylus_cornuta' -c 115. "60164"
was the number of sites in the alignment. "-c 115" represents a median of 115 MYA divergence
time, it is the calibration points between *Fraxinus nigra* and *Corylus cornuta*. Then the
ultrametric tree was estimated using the program r8s v1.81 with default parameters. The
ultrametric tree and gene family counts were used as input for the program CAFE5. Since
CAFE5 was not recommended to analyze recent polyploid species, only diploid species were
used to run CAFE5, including *Cornus alternifolia*, *Fraxinus nigra*, *Dirca palustris*, *Tilia*
americana, *Acer saccharum*, *Acer spicatum*, *Amelanchier laevis*, *Quercus rubra*, *Betula*
papyrifera, *Alnus incana*, *Corylus cornuta*, *Ostrya virginiana*, and *Ribes cynosbati*. Since
assembly errors might cause the observed number of gene copies in gene families to deviate
from the true ones and overestimation of the birth-death parameter λ (number of gene gains and
losses per gene per million years), an error model was estimated first with the ultrametric tree
and gene family counts using CAFE5 with the parameters of "-c 16 -p -e". This error model
along with the ultrametric tree and gene family counts were used to estimate the birth-death
parameter λ, which represents the probability that any gene will be gained or lost. Multiple λ
mode was used with three different rates of gene family evolution assigned as below:
307
308 ((Cornus_alternifolia:3,Fraxinus_nigra:3):3,(((((Dirca_palustris:2,Tilia_americana:2):2,(Acer_sac
309 charum:2,Acer_spicatum:2):2):2,(((((Betula_papyrifera:2,Alnus_incana:2):2,(Corylus_cornuta:2,
310 Ostrya_virginiana:2):2):2,Quercus_rubra:2):2,Amelanchier_laevis:2):2):2,Ribes_cynosbati:1):1);
311
The numbers 1, 2, 3 represent three different gene family evolutionary rates that were shared by
different clades. This tree file was used as input along with the ultrametric tree, gene family

counts, and error model using CAFE5 with the parameters of -c 16 -p -k 3. Gene families or
parent nodes with a significantly greater rate of evolution (family-wide p-value <0.01), in other
words, rapidly evolving gene families, were counted for each species and each parent node.
These gene families significantly violated a random gene birth and death model. Thus, the null
hypothesis of neutral evolution was rejected and these families could be recognized as fast
evolving gene families (Hahn et al. 2005). Enriched GO and KEGG terms were detected in the
fast expanding and fast contracting gene families, respectively.
321
322
323 **Results**
324
Ks analyses
We acquired an average of 41 million raw reads for each sample. The total number of assembled
transcripts ranged between 19,459 and 31,804 (Supplementary Table S1). We retrieved 76 ~ 88%
of the BUSCOs in the assembled transcriptomes (Supplementary Fig. S2). Most species have an
initial Ks peak around 0.02~0.04, except *Fraxinus nigra* and *Tilia americana*, which have the
initial peak around 0.1 and 0.09, respectively (Fig. 2). Such initial peaks represent most recently
duplicate genes. Another discernable separate peak shared by the studied species is around 0.5 ~
0.7. Such shared peaks by all species may correspond to large-scale duplication events, perhaps
the common WGDs across core eudicots. Some species have other discernable peaks, like
Populus grandidentata and *Populus tremuloides*, which have discernable peaks around 0.18.
Such unique peaks may represent ancient duplication events specific to certain species and
different evolutionary patterns across species. The two *Prunus* species have similar Ks shapes, so
do the two *Populus* species, suggestive of common WGDs before the two species diverged from
the same ancestors. By examining the orthologous Ks of the two *Prunus* and of the two *Populus*
species, respectively, we found a high peak close to zero, which may represent the duplication
events after the WGDs and divergence (Supplementary Fig. S3).
341
Duplication events and divergence time
Using a species tree with estimated branch lengths, we inferred divergence times and duplication
events (with at least 50% support rate, Fig. 3). Most duplication events (n=1843) occurred at the
divergence of rosids and asterids, which may correspond to common WGDs across eudicots
species. The divergence time of the two asterids species *Cornus alternifolia* and *Fraxinus nigra*
from a common ancestor was about 104 MYA (Fig. 3). The divergence time of the studied
malvids and fabids from a common ancestor with Saxifragales was about 100 MYA.
349
350 *Relationship between gene family size and evolutionary rate*
351 OrthoFinder assigned 393,796 genes (96.2% of total) to 21,900 orthogroups (gene families). The
352 mean and median gene family size is 17.3 and 12, respectively. We found that 3,060 families
353 present in all species, whereas 19 consisted entirely of single-copy genes.
354
After identifying reciprocal best blastp hits with the outgroup *D. indica*, we retained a total of
6903 ~ 7302 gene families for the studied species to calculate the relationship between gene
family size and evolutionary rate. Gene families with large sizes tended to have low Ka/Ks, yet
there are a few exceptions. We highlighted the gene families with a size ≥5 and Ka/Ks >1 in red
(Fig. 4a). We tested the enriched KEGG and GO terms in these highlighted gene families (Fig.

4b and Supplementary Fig. S4, see the whole list of GO enrichment in Supplementary Table S2).
The most highly shared enriched KEGG and GO terms are related to plant-pathogen interactions,
signaling, and metabolite biosynthesis.

Gene family expansion and contraction
We inferred the birth-death parameter λ (number of gene gains and losses per gene per million
years), which is specified in different clades, using the program CAFE5. $\lambda 1$ (saxifragales)
=0.0034, $\lambda 2$ (rosids) =0.0042, $\lambda 3$ (asterids) =0.0027, which means that the lineages leading to
rosids have the highest gene family turnover rates, followed by saxifragales, and then by asterids.
The percentage and numbers of gene family expansion and contraction are shown in Fig. 5a and
Supplementary Fig. S5, respectively. All of the studied species have more rapidly expanding
than rapidly contracting gene families (family-wide p-value <0.01, Fig. 5b), which may
contribute to their lineage-specific novelty.

We tested the enriched GO and KEGG terms in the fast evolving (expanding or contracting)
gene families. All of the studied species shared the enriched KEGG pathway term plant-pathogen
interaction (Fig. 6). This term was detected in the fast-expanding gene families of *Cornus*
alternifolia, *Tilia americana*, *Acer spicatum*, *Betula papyrifera*, *Quercus rubra*, *Amelanchier*
laevis, *Ribes cynosbati*; and this term was detected in the fast-contracting gene families of
Fraxinus nigra, *Dirca palustris*, *Alnus incana*, *Corylus cornuta*, *Ostrya virginiana*; also, this
term was detected in both of the fast expanding and fast contracting gene families of *Acer*
saccharum. The gene families assigned to the plant-pathogen interaction term encode proteins
(Supplementary Fig. S6) related to disease resistance protein, leucine-rich repeat receptor like
protein kinase and others. Some species shared the enriched KEGG pathway terms Tryptophan
metabolism and MAPK signaling pathway (Fig. 6).

Discussion
Gene duplication events are prevalent in all organisms and provide genetic variation for
evolutionary innovation and adaption to fluctuating conditions (Panchy et al. 2016; Van de Peer
et al. 2009; Zhang 2003). Research on the mechanisms of gene duplication as well as the origin,
maintenance, and functional roles of genetic diversity in forest tree species are few due to the
limited genome and transcriptome data resources. In the present study, we assembled twenty
transcriptomes from forest trees that co-occur. We identified the signal of common WGDs
shared by core eudicots, and the signal of recent duplication events specific to certain species.
We found gene families involved in plant-pathogen interaction, signaling, and metabolite
biosynthesis actively remain extensive variation. We also found fast expansion or fast
contraction in plant-pathogen interaction genes amongst the studied diploid species. In the
following sections, we will discuss these results in detail.

Tempo and mode of gene duplication
The Ks distribution plots confirmed the widespread paleopolyploidy in plants, that's said,
today's diploid species might be polyploid (paleopolyploid) at some points in evolutionary
history (Blanc and Wolfe 2004). The abundant duplication events provide resources for genetic
variation. It is worth noting that Ks distribution is not suitable to infer very ancient events due to
saturation with multiple substitutions (Blanc and Wolfe 2004), moreover, different peaks might

not be separable from each other, thus it is possible that we missed identifying some duplication
events in the present study. Collinearity- or synteny-based methods along with chromosome-
level reference genome assembly is an alternative method to identify ancient WGD, yet high
quality genome assembly is not easy to obtain (Zwaenepoel and Van de Peer 2019).
410
The predicted species tree (Fig. 3) generally agrees with current phylogenetic hypotheses for the
angiosperms with the exception of the two *Populus* species, which is probably caused by the
incomplete gene sets obtained from one leaf RNA-seq sample and/or interspecific gene flow.
The duplication events at the root (Fig. 3) of the tree is likely to be related to the gamma WGD
before the divergence of rosids and asterids. Such large-scale duplication may contribute to
genome structure variation, organismal complexity, and rapid speciation in angiosperm species.
WGD events followed by different evolutionary events might also occur in some clades. For
Acer, it is likely that *Acer spicatum* experienced a diploidization event to revert to diploid status,
while *Acer rubrum* maintained a polyploid status.
420
Divergence time mostly agrees with the previous reports, which stated rosids and asterids
diverged about 125 MYA, malvids and fabids diverged about 110 MYA (Zeng et al. 2017).
However, divergence estimate in the present study is rather rough. Further studies need to
consider using dense taxon sampling, diverse markers like nuclear internal transcribed spacer or
chloroplast DNA fragments, and molecular dating with fossil corrections methods to better
resolve the evolutionary relationships.
427
Sustained large size gene families
Though most duplicate genes were eliminated or pseudonized (Panchy et al. 2016), a small
portion of gene families can actively maintain duplicate genes as selection favors the fitness
related neo-functionalization. The main functions of these sustained large size gene families are
mainly involved in plant-pathogen interaction, signaling, and metabolite biosynthesis (Fig. 4b &
Supplementary Fig. S4). This is consistent with previous report that genes encoding functions
like signal transduction and stress response tend to have paralogs (Panchy et al. 2016), and genes
related to the function like transcription regulation and signal transduction had higher possibility
to survive in multiple rounds of genome duplication (Seoighe and Gehring 2004). Plants
constantly combat with pathogens and pests. The major class of plant disease resistance (R)
genes contain a nucleotide-binding site (NBS) and C-terminal leucin-rich repeats (LRRs).
Tandem and segmental gene duplication explain much of the diversification among R genes
(Andersen et al. 2020; Leister 2004; Meyers et al. 2003), possibly because they have experienced
frequent deletion and duplication throughout evolution, and thus are highly variable (Gao et al.
2018; Hanada et al. 2008). Plant secondary metabolites, such as monoterpenoid, flavonoid,
benzoxazinoid, play important roles in defense against predation and pathogens (Bergman et al.
2020; de Bruijn et al. 2018; Falcone Ferreyra et al. 2012). The regulation of plant secondary
metabolism requires the intimation of signal transduction network that leads to activation of
biosynthetic genes (Zhao et al. 2005).
447
Gene duplication and deletion give rise to gene family expansion and contraction. The high gene
gain rate likely reflects the occurrence of multiple WGDs and frequent SSDs (Carretero-Paulet et
al. 2015). While for the other species with contraction outnumbered expansion, the pattern is
mainly caused by gene loss and retention of specific gene families followed the common WGDs

preceded their common ancestor (Fig. 3). We found a high gene family turnover rate in rosids,
but we must caution that the unbalanced taxon sampling in our study species might cause this
high turnover value. Moreover, future studies should consider distinguishing the branches with
or without recent WGDs, as the branches with recent WGDs tend to have a high gene turnover
rate (Casola and Koralewski 2018).
457
Rapid evolution in plant-pathogen interaction genes
We found rapid evolution of genes involved in plant-pathogen interaction, tryptophan
metabolism and MAPK signaling pathway (Fig. 6) in the adaptive expansion and adaptive
contraction gene families across the studied species. The specific expanding or contracting gene
families involved in plant-pathogen interaction in each species differed from each other,
implying lineage-specificity in changed gene families (Supplementary Fig. S6). This result is
consistent with previous studies, which found that fast evolving gene families were mostly
groups with functions related with immune defense/stress response, metabolism, cell signaling,
chemoreception, and reproduction (Demuth and Hahn 2009). Additionally, a study on *Juglans*
(walnuts) found notable contractions in gene families that are involved in disease resistance and
other abiotic stress response (Trouern-Trend et al. 2020). Both metabolism and MAPK signaling
pathway are associated with plant defense. MAPK cascades sense pathogen/microbe-associated
molecular patters and pathogen effectors, and thus play pivotal roles in signaling plant defense
against pathogen attack (Meng and Zhang 2013). Plant secondary metabolites contribute to plant
immunity as antibiotics, or to control callose deposition and programmed cell death (Piasecka et
al. 2015). The biotic stress exerted by pathogen or pests is a major source of selection pressure
on the studied trees, which drive the rapid radiation of diverse and large plant immune receptor
gene repertoires. Expansion and/or contraction of cell-surface receptor genes correlate with the
expansion and/or contraction of intracellular receptor genes (Ngou et al. 2022).
477
Limitation of the current study
Our conclusions are drawn from samples collected from co-occurring species from a single
forest. The forest type sampled and the species are typical for the entire region including the
Ottawa, Nicolet, and Hiawatha National Forests in northern Wisconsin and the Upper Peninsula
of Michigan, USA. As individuals within species vary in their adaptive and plastic responses to
the environment, our results are, therefore, most applicable to the study region and not the entire
geographic range of the species. Furthermore, the assembled transcripts of each species in this
study are derived from leaf material and, therefore, cannot represent the whole gene set
expressed by the organism. Lastly, we also note that we filtered out genes that had low
expression levels to ensure the quality of the assembled transcriptome. Thus, our study does omit
some genes, and it is possible that we might end up with different conclusions if we had more
genes.
490
Conclusions
In summary, we found the common WGDs shared by all the studied species as well as signals of
recent duplication events specific to certain species, which are suggestive of different
evolutionary patterns. Gene families involved in plant-pathogen interaction, signaling, and
metabolite biosynthesis actively remain extensive variation and are fast expanding or contracting
amongst the studied species. This knowledge calls for an emphasis of gene family size variation,
i.e. copy number variation, when studying forest genetic diversity, which will inform forest

adaptive strategy. Future studies may focus on classification of duplications in forest tree species,
and investigate how the duplications give rise to species-specific innovation and adaption.
500
501

Data availability
The raw RNA-seq data were deposited in NCBI SRA (accession number: PRJNA896247;
http://www.ncbi.nlm.nih.gov/sra). The assembled transcriptomes and their annotation were
deposited in Dryad https://doi.org/10.5061/dryad.sj3tx968b. The scripts used for analyses were
deposited in https://github.com/Mengmeng-Lu/Gene-duplication-and-gene-expansion-analyses-
in-twenty-tree-species.
508

Acknowledgements
We thank the Center for Research Computing at University of Notre Dame for high performance
computing support. We specially acknowledge the assistance of Dr. Dodi Heryadi, Dr. Scott
Hampton, and Júlia Simon. We appreciate Dr. Sam Yeaman and Dr. James R. Whiting for
constructive suggestions. The research was performed at the University of Notre Dame
Environmental Research Center that has been generously supported by the Bernard J. Hank
Family Endowment. The research was permitted under UNDERC research permit
number: UNDERC-2021-2.
517
518

519 **Funding**
520 MM and NGS were supported by the US National Science Foundation Dimensions of
521 Biodiversity grant (DEB-2124466). JY and MC were supported by a NSFC China-US
522 Dimensions of Biodiversity Grant (DEB-32061123003).
523
524

525 **Conflicts of interest**
526 The authors declare no conflict of interest.
527
528
529

Literature cited
Alexa A, Rahnenfuhrer J. 2023. topGO: Enrichment analysis for gene ontology. R package
version 2.52.0.
Allen CD, Macalady AK, Chenchouni H, Bachelet D, McDowell N, Vennetier M, Kitzberger T,
Rigling A, Breshears DD, Hogg EHT, et al. 2010. A global overview of drought and heat-
induced tree mortality reveals emerging climate change risks for forests. Forest Ecol
Manag. 259(4):660-684. doi:10.1016/j.foreco.2009.09.001.
Amborella Genome Project. 2013. The Amborella genome and the evolution of flowering plants.
Science. 342(6165):1241089. doi:10.1126/science.1241089.
Andersen EJ, Nepal MP, Purintun JM, Nelson D, Mermigka G, Sarris PF. 2020. Wheat disease
resistance genes and their diversification through integrated domain fusions. Front
Genet. 11:898. doi:10.3389/fgene.2020.00898.

Bailey JA, Gu Z, Clark RA, Reinert K, Samonte RV, Schwartz S, Adams MD, Myers EW, Li PW,
Eichler EE. 2002. Recent segmental duplications in the human genome. Science.
297(5583):1003-1007. doi:10.1126/science.1072047.
Bergman ME, Chávez Á, Ferrer A, Phillips MA. 2020. Distinct metabolic pathways drive
monoterpenoid biosynthesis in a natural population of *Pelargonium graveolens*. J Exp
Bot. 71(1):258-271. doi:10.1093/jxb/erz397.
Blanc G, Wolfe KH. 2004. Widespread paleopolyploidy in model plant species inferred from age
distributions of duplicate genes[W]. Plant Cell. 16(7):1667-1678.
doi:10.1105/tpc.021345.
Brosius J. 1991. Retroposons—seeds of evolution. Science. 251(4995):753.
doi:10.1126/science.1990437.
Buchfink B, Reuter K, Drost H-G. 2021. Sensitive protein alignments at tree-of-life scale using
DIAMOND. Nat Methods. 18(4):366-368. doi:10.1038/s41592-021-01101-x.
Cantalapiedra CP, Hernández-Plaza A, Letunic I, Bork P, Huerta-Cepas J. 2021. eggNOG-mapper
v2: Functional annotation, orthology assignments, and domain prediction at the
metagenomic scale. Mol Biol Evol. 38(12):5825-5829. doi:10.1093/molbev/msab293.
Carretero-Paulet L, Librado P, Chang T-H, Ibarra-Laclette E, Herrera-Estrella L, Rozas J, Albert VA.
2015. High gene family turnover rates and gene space adaptation in the compact
genome of the Carnivorous plant *Utricularia gibba*. Mol Biol Evol. 32(5):1284-1295.
doi:10.1093/molbev/msv020.
Casola C, Koralewski TE. 2018. Pinaceae show elevated rates of gene turnover that are robust to
incomplete gene annotation. Plant J. doi:10.1111/tpj.13994.
Cazzolla Gatti R, Reich PB, Gamarra JGP, Crowther T, Hui C, Morera A, Bastin J-F, de-Miguel S,
Nabuurs G-J, Svenning J-C, et al. 2022. The number of tree species on Earth. Proc Natl
Acad Sci USA. 119(6):e2115329119. doi:10.1073/pnas.2115329119.
Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: An ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics. 34(17):i884-i890. doi:10.1093/bioinformatics/bty560.
Conant GC, Birchler JA, Pires JC. 2014. Dosage, duplication, and diploidization: clarifying the
interplay of multiple models for duplicate gene evolution over time. Curr Opin Plant Biol.
19:91-98. doi:10.1016/j.pbi.2014.05.008.
Cortés AJ, Restrepo-Montoya M, Bedoya-Canas LE. 2020. Modern strategies to assess and
breed forest tree adaptation to changing climate. Front Plant Sci. 11:583323.
doi:10.3389/fpls.2020.583323.
Cui L, Wall PK, Leebens-Mack JH, Lindsay BG, Soltis DE, Doyle JJ, Soltis PS, Carlson JE,
Arumuganathan K, Barakat A, et al. 2006. Widespread genome duplications throughout
the history of flowering plants. Genome Res. 16(6):738-749. doi:10.1101/gr.4825606.
de Bruijn WJC, Gruppen H, Vincken J-P. 2018. Structure and biosynthesis of benzoxazinoids:
Plant defence metabolites with potential as antimicrobial scaffolds. Phytochemistry.
155:233-243. doi:10.1016/j.phytochem.2018.07.005.
De Smet R, Adams KL, Vandepoele K, Van Montagu MCE, Maere S, Van de Peer Y. 2013.
Convergent gene loss following gene and genome duplications creates single-copy
families in flowering plants. Proc Natl Acad Sci USA. 110(8):2898-2903.
doi:10.1073/pnas.1300127110.

Demuth JP, Hahn MW. 2009. The life and death of gene families. Bioessays. 31(1):29-39.
doi:10.1002/bies.080085.
Diss G, Gagnon-Arsenault I, Dion-Coté A-M, Vignaud H, Ascencio DI, Berger CM, Landry CR. 2017.
Gene duplication can impart fragility, not robustness, in the yeast protein interaction
network. Science. 355(6325):630-634. doi:10.1126/science.aai7685.
Dodds PN, Rathjen JP. 2010. Plant immunity: towards an integrated view of plant–pathogen
interactions. Nat Rev Genet. 11(8):539-548. doi:10.1038/nrg2812.
Emms DM, Kelly S. 2017. STRIDE: Species tree root inference from gene duplication events. Mol
Biol Evol. 34(12):3267-3278. doi:10.1093/molbev/msx259.
Emms DM, Kelly S. 2018. STAG: Species tree inference from all genes. bioRxiv.
doi:10.1101/267914.
Emms DM, Kelly S. 2019. OrthoFinder: Phylogenetic orthology inference for comparative
genomics. Genome Biol. 20(1):238. doi:10.1186/s13059-019-1832-y.
Enright AJ, Van Dongen S, Ouzounis CA. 2002. An efficient algorithm for large-scale detection of
protein families. Nucleic Acids Res. 30(7):1575-1584. doi:10.1093/nar/30.7.1575.
Falcone Ferreyra ML, Rius SP, Casati P. 2012. Flavonoids: biosynthesis, biological functions, and
biotechnological applications. Front Plant Sci. 3:222. doi:10.3389/fpls.2012.00222.
FAO. 2020. Global Forest Resources Assessment 2020: Main report. Rome.
https://doi.org/10.4060/ca9825en.
Force A, Lynch M, Pickett FB, Amores A, Yan YL, Postlethwait J. 1999. Preservation of duplicate
genes by complementary, degenerative mutations. Genetics. 151(4):1531-1545.
doi:10.1093/genetics/151.4.1531.
Fu L, Niu B, Zhu Z, Wu S, Li W. 2012. CD-HIT: accelerated for clustering the next-generation
sequencing data. Bioinformatics. 28(23):3150-3152. doi:10.1093/bioinformatics/bts565.
Gamfeldt L, Snäll T, Bagchi R, Jonsson M, Gustafsson L, Kjellander P, Ruiz-Jaen MC, Fröberg M,
Stendahl J, Philipson CD, et al. 2013. Higher levels of multiple ecosystem services are
found in forests with more tree species. Nat Commun. 4(1):1340.
doi:10.1038/ncomms2328.
Gao Y, Wang W, Zhang T, Gong Z, Zhao H, Han G-Z. 2018. Out of water: The origin and early
diversification of plant R-genes. Plant Physiol. 177(1):82-89. doi:10.1104/pp.18.00185.
Götz S, García-Gómez JM, Terol J, Williams TD, Nagaraj SH, Nueda MJ, Robles M, Talón M,
Dopazo J, Conesa A. 2008. High-throughput functional annotation and data mining with
the Blast2GO suite. Nucleic Acids Res. 36(10):3420-3435. doi:10.1093/nar/gkn176.
Hahn MW, De Bie T, Stajich JE, Nguyen C, Cristianini N. 2005. Estimating the tempo and mode
of gene family evolution from comparative genomic data. Genome Res. 15(8):1153-1160.
doi:10.1101/gr.3567505.
Hammond WM, Williams AP, Abatzoglou JT, Adams HD, Klein T, López R, Sáenz-Romero C,
Hartmann H, Breshears DD, Allen CD. 2022. Global field observations of tree die-off
reveal hotter-drought fingerprint for Earth's forests. Nat Commun. 13(1):1761.
doi:10.1038/s41467-022-29289-2.
Hanada K, Zou C, Lehti-Shiu MD, Shinozaki K, Shiu SH. 2008. Importance of lineage-specific
expansion of plant tandem duplicates in the adaptive response to environmental stimuli.
Plant Physiol. 148(2):993-1003. doi:10.1104/pp.108.122457.

Hurley BP, Garnas J, Wingfield MJ, Branco M, Richardson DM, Slippers B. 2016. Increasing
629   numbers and intercontinental spread of invasive insects on eucalypts. Biol Invasions.
630   18(4):921-933. doi:10.1007/s10530-016-1081-x.
631   Isabel N, Holliday JA, Aitken SN. 2020. Forest genomics: Advancing climate adaptation, forest
632   health, productivity, and conservation. Evol Appl. 13(1):3-10. doi:10.1111/eva.12902.
633   Jiang N, Bao Z, Zhang X, Eddy SR, Wessler SR. 2004. Pack-MULE transposable elements mediate
634   gene evolution in plants. Nature. 431(7008):569-573. doi:10.1038/nature02953.
635   Jiao Y, Leebens-Mack J, Ayyampalayam S, Bowers JE, McKain MR, McNeal J, Rolf M, Ruzicka DR,
636   Wafula E, Wickett NJ, et al. 2012. A genome triplication associated with early
637   diversification of the core eudicots. Genome Biol. 13(1):R3. doi:10.1186/gb-2012-13-1-
638   r3.
639   Johri P, Gout J-F, Doak TG, Lynch M. 2022. A population-genetic lens into the process of gene
640   loss following whole-genome duplication. Mol Biol Evol. 39(6):msac118.
641   doi:10.1093/molbev/msac118.
642   Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A,
643   Nuka G, et al. 2014. InterProScan 5: Genome-scale protein function classification.
644   Bioinformatics. 30(9):1236-1240. doi:10.1093/bioinformatics/btu031.
645   Kassambara A. 2023. ggpubr: 'ggplot2' based publication ready plots. R package version 0.6.0.
646   https://rpkgs.datanovia.com/ggpubr/.
647   Katoh K, Misawa K, Kuma Ki, Miyata T. 2002. MAFFT: A novel method for rapid multiple
648   sequence alignment based on fast Fourier transform. Nucleic Acids Res. 30(14):3059-
649   3066. doi:10.1093/nar/gkf436.
650   Kirilenko AP, Sedjo RA. 2007. Climate change impacts on forestry. Proc Natl Acad Sci USA.
651   104(50):19697-19702. doi:10.1073/pnas.0701424104.
652   Kolde R. 2019. pheatmap: Pretty Heatmaps. R package version 1.0.12.
653   Kumar S, Stecher G, Suleski M, Hedges SB. 2017. TimeTree: A resource for timelines, timetrees,
654   and divergence times. Mol Biol Evol. 34(7):1812-1819. doi:10.1093/molbev/msx116.
655   Kuzmin E, VanderSluis B, Nguyen Ba AN, Wang W, Koch EN, Usaj M, Khmelinskii A, Usaj MM,
656   van Leeuwen J, Kraus O, et al. 2020. Exploring whole-genome duplicate gene retention
657   with complex genetic interaction analysis. Science. 368(6498):eaaz5667.
658   doi:10.1126/science.aaz5667.
659   Leister D. 2004. Tandem and segmental gene duplication and recombination in the evolution of
660   plant disease resistance gene. Trends Genet. 20(3):116-122.
661   doi:10.1016/j.tig.2004.01.007.
662   Li B, Dewey CN. 2011. RSEM: Accurate transcript quantification from RNA-Seq data with or
663   without a reference genome. BMC Bioinformatics. 12(1):323. doi:10.1186/1471-2105-
664   12-323.
665   Linnakoski R, Forbes KM. 2019. Pathogens—The hidden face of forest invasions by wood-boring
666   insect pests. Front Plant Sci. 10:90. doi:10.3389/fpls.2019.00090.
667   Liu J, Yu T, Mu Z, Li G. 2019. TransLiG: a de novo transcriptome assembler that uses line graph
668   iteration. Genome Biol. 20(1):81. doi:10.1186/s13059-019-1690-7.
669   Lu M, Feau N, Obreht Vidakovic D, Ukrainetz N, Wong B, Aitken SN, Hamelin RC, Yeaman S.
670   2021. Comparative gene expression analysis reveals mechanism of *Pinus contorta*

671   response to the fungal pathogen *Dothistroma septosporum*. Mol Plant Microbe Interact.
672   34(4):397-409. doi:10.1094/mpmi-10-20-0282-r.

673 Lynch M, Conery JS. 2000. The evolutionary fate and consequences of duplicate genes. Science.
674   290(5494):1151-1155. doi:10.1126/science.290.5494.1151.

675 Lynch M, Conery JS. 2003. The evolutionary demography of duplicate genes. J Struct Funct
676   Genomics. 3(1-4):35-44. doi:10.1023/A:1022696612931.

677 Maere S, De Bodt S, Raes J, Casneuf T, Van Montagu M, Kuiper M, Van de Peer Y. 2005.
678   Modeling gene and genome duplications in eukaryotes. Proc Natl Acad Sci USA.
679   102(15):5454-5459. doi:10.1073/pnas.0501102102.

680 Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO Update: novel and
681   streamlined workflows along with broader and deeper phylogenetic coverage for
682   scoring of eukaryotic, prokaryotic, and viral genomes. Mol Biol Evol. 38(10):4647-4654.
683   doi:10.1093/molbev/msab199.

684 Mendes FK, Vanderpool D, Fulton B, Hahn MW. 2021. CAFE 5 models variation in evolutionary
685   rates among gene families. Bioinformatics. 36(22-23):5516-5518.
686   doi:10.1093/bioinformatics/btaa1022.

687 Meng X, Zhang S. 2013. MAPK cascades in plant disease resistance signaling. Annu Rev
688   Phytopathol. 51:245-266. doi:10.1146/annurev-phyto-082712-102314.

689 Meyers BC, Kozik A, Griego A, Kuang H, Michelmore RW. 2003. Genome-wide analysis of NBS-
690   LRR-encoding genes in Arabidopsis. Plant Cell. 15(4):809-834. doi:10.1105/tpc.009308.

691 Miura S, Amacher M, Hofer T, San-Miguel-Ayanz J, Ernawati, Thackway R. 2015. Protective
692   functions and ecosystem services of global forests in the past quarter-century. Forest
693   Ecol Manag. 352(7):35-46. doi:10.1016/j.foreco.2015.03.039.

694 Neale DB, Kremer A. 2011. Forest tree genomics: growing resources and applications. Nat Rev
695   Genet. 12(2):111-122. doi:10.1038/nrg2931.

696 Ngou BPM, Heal R, Wyler M, Schmid MW, Jones JDG. 2022. Concerted expansion and
697   contraction of immune receptor gene repertoires in plant genomes. Nat Plants.
698   8(10):1146-1152. doi:10.1038/s41477-022-01260-5.

699 Ohno S. 1970. Evolution by gene duplication. New York: Springer.

700 One Thousand Plant Transcriptomes Initiative. 2019. One thousand plant transcriptomes and
701   the phylogenomics of green plants. Nature. 574(7780):679-685. doi:10.1038/s41586-
702   019-1693-2.

703 Panchy N, Lehti-Shiu M, Shiu SH. 2016. Evolution of gene duplication in plants. Plant Physiol.
704   171(4):2294-2316. doi:10.1104/pp.16.00523.

705 Piasecka A, Jedrzejczak-Rey N, Bednarek P. 2015. Secondary metabolites in plant innate
706   immunity: Conserved function of divergent chemicals. New Phytol. 206(3):948-964.
707   doi:10.1111/nph.13325.

708 Picart-Picolo A, Grob S, Picault N, Franek M, Llauro C, Halter T, Maier TR, Jobet E, Descombin J,
709   Zhang P, et al. 2020. Large tandem duplications affect gene expression, 3D organization,
710   and plant-pathogen response. Genome Res. 30(11):1583-1592.
711   doi:10.1101/gr.261586.120.

712 Price MN, Dehal PS, Arkin AP. 2009. FastTree: Computing large minimum evolution trees with
713   profiles instead of a distance matrix. Mol Biol Evol. 26(7):1641-1650.
714   doi:10.1093/molbev/msp077.

R Core Team. 2021. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. https://www.R-project.org/.
Reid NM, Proestou DA, Clark BW, Warren WC, Colbourne JK, Shaw JR, Karchner SI, Hahn ME,
Nacci D, Oleksiak MF, et al. 2016. The genomic landscape of rapid repeated evolutionary
adaptation to toxic pollution in wild fish. Science. 354(6317):1305-1308.
doi:10.1126/science.aah4993.
Rizvi AR, Baig S, Barrow E, Kumar C. 2015. Synergies between climate mitigation and adaptioon
in forest landscape restoration. Gland, Switzerland: IUCN.
Sanderson MJ. 2003. r8s: Inferring absolute rates of molecular evolution and divergence times
in the absence of a molecular clock. Bioinformatics. 19(2):301-302.
doi:10.1093/bioinformatics/19.2.301.
Seoighe C, Gehring C. 2004. Genome duplication led to highly selective expansion of the
Arabidopsis thaliana proteome. Trends Genet. 20(10):461-464.
doi:10.1016/j.tig.2004.07.008.
Soltis PS, Soltis DE. 2016. Ancient WGD events as drivers of key innovations in angiosperms.
Curr Opin Plant Biol. 30:159-165. doi:10.1016/j.pbi.2016.03.015.
Swenson NG, Iida Y, Howe R, Wolf A, Umaña MN, Petprakob K, Turner BL, Ma K. 2017. Tree co-
occurrence and transcriptomic response to drought. Nat Commun. 8(1):1996.
doi:10.1038/s41467-017-02034-w.
Swenson NG, Jones FA. 2017. Community transcriptomics, genomics and the problem of species
co-occurrence. J Ecol. 105(3):563-568. doi:10.1111/1365-2745.12771.
Tang H, Bowers JE, Wang X, Ming R, Alam M, Paterson AH. 2008. Synteny and collinearity in
plant genomes. Science. 320(5875):486-488. doi:10.1126/science.1153917.
Trouern-Trend AJ, Falk T, Zaman S, Caballero M, Neale DB, Langley CH, Dandekar AM, Stevens
KA, Wegrzyn JL. 2020. Comparative genomics of six Juglans species reveals disease-
associated gene family contractions. Plant J. 102(2):410-423. doi:10.1111/tpj.14630.
Van de Peer Y, Maere S, Meyer A. 2009. The evolutionary significance of ancient genome
duplications. Nat Rev Genet. 10(10):725-732. doi:10.1038/nrg2600.
Wickham H. 2011. The Split-Apply-Combine strategy for data analysis. J Stat Softw. 40(1):1-29.
doi:10.18637/jss.v040.i01.
Wickham H. 2016. ggplot2: Elegant graphics for data analysis. New York: Springer.
Yang Z. 2007. PAML 4: Phylogenetic analysis by maximum likelihood. Mol Biol Evol. 24(8):1586-
1591. doi:10.1093/molbev/msm088.
Yang Z, Nielsen R. 2000. Estimating synonymous and nonsynonymous substitution rates under
realistic evolutionary models. Mol Biol Evol. 17(1):32-43.
doi:10.1093/oxfordjournals.molbev.a026236.
Yu Y. 2022. mixR: An R package for finite mixture modeling for both raw and binned data.
Journal of Open Source Software. 7(69):4031. doi:10.21105/joss.04031.
Zeng L, Zhang N, Zhang Q, Endress PK, Huang J, Ma H. 2017. Resolution of deep eudicot
phylogeny and their temporal diversification using nuclear genes from transcriptomic
and genomic datasets. New Phytol. 214(3):1338-1354. doi:10.1111/nph.14503.
Zhang J. 2003. Evolution by gene duplication: an update. Trends Ecol Evol. 18(6):292-298.
doi:10.1016/S0169-5347(03)00033-8.

Zhao J, Davis LC, Verpoorte R. 2005. Elicitor signal transduction leading to production of plant
759   secondary metabolites. Biotechnol Adv. 23(4):283-333.
760   doi:10.1016/j.biotechadv.2005.01.003.
761   Zwaenepoel A, Van de Peer Y. 2019. Inference of ancient whole-genome duplications and the
762   evolution of gene duplication and loss rates. Mol Biol Evol. 36(7):1384-1404.
763   doi:10.1093/molbev/msz088.

764
765
766
767 **Figure legends**
768
769 **Fig. 1. The location (red dot) of the University of Notre Dame Environmental Research**
770 **Center (UNDERC) where all samples used in this study were collected.**
771
772 **Fig. 2. Distribution of Ks (the level of synonymous substitutions). A mixture model was fit**
773 **and plot for the Ks values ≥0.01 and ≤2 of each species using the R package mixR.**
774
775 **Fig. 3. Divergence time in million years and gene duplication events with 50% support rate.**
776 **Node numbers were followed by number of gene duplication events.**
777
778 **Fig. 4. Scatter plot between gene size and Ka/Ks per gene family (a) and KEGG pathway**
779 **terms enrichment (b) in gene families with a size ≥5 and Ka/Ks >1. The color bar in b**
780 **represents the adjusted p-values of KEGG enrichment test.**
781
782 **Fig. 5. The percentage of gene family expansion and contraction (a) and the number of fast**
783 **evolving gene families (b), with + representing fast expanding gene family and –**
784 **representing fast contracting gene family.**
785
786 **Fig. 6. Enriched KEGG pathway terms in the studied diploid species. Red represents fast**
787 **expanding gene family; blue represents fast contracting gene family; white represents both**
788 **fast expanding and contracting gene family. Dark grey represents the KEGG terms were**
789 **not enriched in fast expanding or contracting gene family.**
790
791