# Comparative transcriptomics reveals divergence in pathogen response gene families amongst twenty forest tree species

[1*]Mengmeng Lu, [2]Min Cao, [2,3]Jie Yang, [1]Nathan G. Swenson

[1] Department of Biological Sciences, University of Notre Dame, 100 Galvin Life Sciences, Notre Dame, IN 46556, USA

[2] CAS Key Laboratory of Tropical Forest Ecology, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Mengla, Yunnan 666303, China.

[3]Center of Plant Ecology, Core Botanical Gardens, Chinese Academy of Sciences, Mengla, Yunnan 666303, China.

[*]Corresponding author: Department of Biological Sciences, University of Notre Dame, 100 Galvin Life Sciences, Notre Dame, IN 46556, USA. Email: mlu@nd.edu

21 **Abstract**
22

Forest trees provide critical ecosystem services for humanity that are under threat due to ongoing
global change. Measuring and characterizing genetic diversity is key to understanding adaptive
potential and developing strategies to mitigate negative consequences arising from climate
change. In the area of forest genetic diversity, genetic divergence caused by large-scale changes
at the chromosomal level has been largely understudied. In this study, we used the RNA-seq data
of twenty co-occurring forest trees species from genera including *Acer*, *Alnus*, *Amelanchier*,
*Betula*, *Cornus*, *Corylus*, *Dirca*, *Fraxinus*, *Ostrya*, *Populus*, *Prunus*, *Quercus*, *Ribes*, *Tilia*, and
*Ulmus* sampled from Upper Peninsula of Michigan. These data were used to infer the origin and
maintenance of gene family variation, species divergence time, as well as gene family expansion
and contraction. We identified a signal of common whole genome duplication events shared by
core eudicots. We also found rapid evolution, namely fast expansion or fast contraction of gene
families, in plant-pathogen interaction genes amongst the studied diploid species. Finally, the
results lay the foundation for further research on the genetic diversity and adaptive capacity of
forest trees, which will inform forest management and conservation policies.
37

38 **Keywords:** angiosperms, gene duplication, gene family expansion and contraction, Ks plot,
39 Ka/Ks, plant-pathogen interaction

## Introduction

Forest trees occupy nearly a third of the Earth's land surface and provide key ecological benefits and services such as water, air purification and carbon sequestration (Gamfeldt et al., 2013; Miura et al., 2015; FAO, 2020). They are critical components of terrestrial biodiversity in terms of their own diversity, but also with respect to the habitats and renewable materials such as wood, cellulose, and lignin (Cortés et al., 2020). It is estimated that there are ~73,000 tree species globally with most of being rare and regionally endemic, implying vulnerability of global tree species to anthropogenic and climate changes (Cazzolla Gatti et al., 2022). Indeed, climate change has already impacted forests and their species composition via range migration, wildfires, droughts, pest, and pathogens (Kirilenko and Sedjo, 2007; Allen et al., 2010; Rizvi et al., 2015; Hammond et al., 2022). The genetic composition and diversity of the tree species in forests is foundational not only to our understanding of the abiotic and biotic factors governing the distribution and dynamics of species populations, but also to how they will respond to current and future environmental change. Thus, measuring and characterizing intra- and inter-specific genetic diversity is the key for understanding the current structure and dynamics of forests and their adaptive potential and for developing strategies to mitigate the damages brought along with environmental change.

Gene duplication and gene loss are major sources of genetic diversity. Gene duplication, generated through either whole genome duplication (WGD) or small-scale duplication (SSD), is the primary source of new genetic material among flowering plants (Cui et al., 2006; Panchy et al., 2016). WGD contributes to the majority of duplicate genes (AMBORELLA GENOME PROJECT et al., 2013; Soltis and Soltis, 2016). The ancient WGDs can lead to competitive advantage, increased species diversity, and evolutionary innovations (Van de Peer et al., 2009). Most flowering species have a polyploid history either through autopolyploidy or allopolyploidy. Specifically, flowering species are either currently polyploid, or have a polyploid ancestor and have reverted to being diploid (Blanc and Wolfe, 2004). Earlier studies placed at least three WGD events during the radiation of angiosperms, the oldest WGD occurred before the monocot-eudicot divergence, a second WGD occurred specifically to eudicots, a more recent WGD, gamma ($\gamma$) triplication event, occurred before the separation of rosids and asterids (Tang et al., 2008; Jiao et al., 2012). Alternatively, pervasive SSD, including tandem duplication (Zhang, 2003), transposon-mediated duplication (Jiang et al., 2004), segmental duplication (Bailey et al., 2002b), and retroduplication (Brosius, 1991), also generates a great number of duplicate genes. While the majority of duplicate genes end up being silenced and eliminated (Lynch and Conery, 2000), a few preserved duplicate genes may be retained by selection due to neo-functionalization, relative dosage, absolute dosage or sub- functionalization (Conant et al., 2014), and play important role in generating genetic diversity and building up reproductive isolation barriers. Recent studies explained retention of duplicate genes as some paralogous genes depend on or compensate for their cognate copy (Diss et al., 2017), while some paralogous gene pairs are stuck in a functional and structural entanglement status over evolutionary time scale (Kuzmin et al., 2020).

Compared to gene duplication, the mechanism that underlie gene loss are not yet well understood. It is commonly assumed that gene loss is realized by deletion or pseudogenization, as a result of neutral or adaptive processes (Demuth and Hahn, 2009). Most nonsense mutations

86  are subject to drift to fixation. Some deleterious mutations could be accumulated and eventually
87  become silenced (Johri et al., 2022). Gene loss might be selected to retain dosage balance, as
88  evidenced in a set of single-copy genes with essential housekeeping functions, which are
89  typically restored to become singletons after WGD or SSD (De Smet et al., 2013).
90
Differential duplication and loss of genes among evolutionary lineages results in gene family
expansion and contraction (Demuth and Hahn, 2009). Gene families comprise a set of similar
nucleotide or amino acid sequences. They arise as a result of duplication events and have similar
cellular and biochemical functions. Gene family expansions and contractions related to
metabolic, regulatory and signalling gene networks contribute to the adaptations of plants to
various abiotic habitats and biotic interactions (Leebens-Mack et al., 2019). They are sources of
plant functional evolution and innovation (Ohno, 1970; Force et al., 1999), exhibiting a clear
functional basis, which are influenced by the mechanisms of duplication (Maere et al., 2005;
Hanada et al., 2008). Comparisons of gene family content across species, therefore, may provide
insight into the ecological interactions and evolutionary pressures that have shaped adaptation
and diversity. This is particularly true when we conduct comparative transcriptomic analyses of
species that currently co-occur and that, potentially have co-occurred for long periods of time
(Swenson and Jones, 2017).
104
Genetic investigations of forest trees have, generally, focused on nucleotide substitutions in
specific genes (Neale and Kremer, 2011; Isabel et al., 2020), while comparative analyses of
genetic divergence caused by large-scale changes at the chromosomal levels have been largely
understudied. However, the magnitude, rate, and distribution of changes in gene family size are
key drivers of the functional differentiation and adaptation of species (Lynch and Conery, 2003;
Demuth and Hahn, 2009). Large segmental duplication and changes in copy number are known
to be associated with disease response and environmental adaptation (Bailey et al., 2002a; Reid
et al., 2016). Therefore, gene family size change is as much important as nucleotide substitution
to understanding genetic divergence and evolutionary forces responsible for these changes.
114
Such knowledge on genetic divergence and associated evolutionary forces in forest trees can
help to promote breeding healthy trees (i.e. disease resistant trees) in a changing climate (Allen
et al., 2010; Dodds and Rathjen, 2010). The increasing occurrences and severity of insect and
disease outbreaks have threatened forest trees (Hurley et al., 2016; Linnakoski and Forbes,
2019), but we lack a thorough understanding to the disease related genes in forest trees. Trouern-
Trend et al. (2020) revealed contraction in disease associated gene families in *Juglans* species.
Lu et al. (2021) reported differential gene expression patterns between tolerant and susceptible
lodgepole pine trees to a fungal pathogen. It would be interesting to explore the potential link
between duplication events and disease response in expressed genes of forest trees, as the
duplicate genes can quickly lead to changes in gene expression with the acquisition of increased
resistance to different pathogens (Picart-Picolo et al., 2020).
126
In this study, we used the RNA-seq data for twenty co-occurring forest trees species to infer the
origin and maintenance of gene family variation as well as species divergence time, gene
expansion and contraction. We sought to address the following questions: What is the magnitude
of gene duplication? What factors impact duplicate gene evolution? Are the gene gain and loss
process random or controlled by natural selection? The answers to these questions are the key to

understanding the basis of adaptive divergence in forest tree species and will help to predict
adaptive capacity of these species in a changing climate.
134
135
**Materials and methods**
*Sample collection*
Leaves of each of twenty forest species were sampled in August, 2021, at the University of Notre
Dame Environmental Research Center (UNDERC) near Land O' Lakes, Wisconsin, U.S.A. (Fig.
1). The UNDERC site is located at latitude: 46.23391, longitude: -89.537254, and encompasses
approximately ~8,000 acres of mixed deciduous forest on both sides of the state line between
Wisconsin and Michigan's upper peninsula. The twenty species used in this study are dominate
components of the woody vegetation at the site and are: *Acer rubrum* (red maple), *Acer*
*saccharum* (sugar maple), *Acer spicatum* (mountain maple), *Alnus incana* (Gary alder),
*Amelanchier laevis* (smooth serviceberry), *Betula alleghaniensis* (yellow birch), *Betula*
*papyrifera* (paper birch), *Cornus alternifolia* (pagoda dogwood), *Corylus cornuta* (beaked
hazel), *Dirca palustris* (Eastern leatherwood), *Fraxinus nigra* (black ash), *Ostrya virgniana*
(American hophornbeam), *Populus grandientata* (bigtooth aspen), *Populus tremuloides* (quaking
aspen), *Prunus serotina* (black cherry), *Prunus virginiana* (chokecherry), *Quercus rubra*
(Northern red oak), *Ribes cynosbati* (prickly gooseberry), *Tilia americana* (basswood), *Ulmus*
*americana* (American elm). They are all angiosperms (flowering plants) and eudicots, belonging
to the clades of asterids, saxifragles, and rosids.  These species were chosen as they are the most
dominant at the study site and in large national forests in region (i.e. Ottawa, Nicolet, and
Hiawatha National Forests). Specifically, these species compose >95% of the tree individuals
and above-ground biomass in these forests (Swenson et al., 2017). We focus on the most
dominant species in order to provide insights into the ecological and evolutionary processes
driving transcriptomic variation within and across populations and communities (Swenson and
Jones, 2017). Upon collection, the samples were flash frozen in liquid nitrogen, then transported
and stored in an ultracold freezer at the field station.
160
161

162
163  **Fig. 1. The location (red dot) of the University of Notre Dame Environmental Research**
164  **Center (UNDERC) where all samples used in this study were collected.**
165

166 *RNA-sequencing, de novo transcriptome assembly, and annotation*
167 The samples were sent to Novogene corporation inc. (Chula Vista, CA, USA) for mRNA
168 extraction, library preparation, mRNA sequencing, and quality control. The samples were
169 sequenced on an Illumina NovaSeq 6000 platform to produce 150bp paired-end reads. Raw reads
170 filtering was performed by Novogene to get the clean reads including removing reads containing
171 adapters, removing reads containing N >10%, and removing reads containing low quality (Phred
172 quality score ≤5) base which is over 50% of the total base. An average of 6 Gb paired-end clean
173 reads were acquired for each sample and deposited in NCBI Sequence Read Archive (SRA
174 accession number: PRJNA896247). *De novo* transcriptome assembly was performed for each
175 sample using the software TransLiG v1.3 with default parameters (Liu et al., 2019). The clean
176 reads were then mapped to the assembled transcriptome using the software RSEM v1.2.25 with
177 default parameters (Li and Dewey, 2011). The longest transcript (isoform) with a transcript per
178 million (TPM) value ≥1.5 for each gene was retained. Redundancy was removed from the
179 retained transcripts using the software CD-HIT v4.8.1 with parameters of -c 0.95 -n 8 -T 32 -M
180 1540 (Fu et al., 2012). To remove potential contamination, the retained transcripts were aligned
181 against the non-redundant (nr) protein database (downloaded on January 22, 2022) using the
182 blastx function implemented by the software DIAMOND v2.0.13.151 (Buchfink et al., 2021)
183 with parameter of --outfmt 102 to assign the best known taxon ID to each query transcript. The
184 file that was used to convert accession numbers to taxonomy was downloaded from
185 https://ftp.ncbi.nlm.nih.gov/pub/taxonomy/accession2taxid/, and the tax ID file was downloaded
186 from https://ftp.ncbi.nlm.nih.gov/pub/taxonomy/new_taxdump/. The query transcripts that could
187 be assigned to green plant (Viridiplantae) taxon IDs were retained. The filtered transcripts were
188 aligned to the nr database that only includes green plant subjects using the blastx function
189 implemented by the software DIAMOND v2.0.13.151 with default parameters. The aligned
190 transcripts were combined with those transcripts assigned with green plant taxon IDs to form the
191 final transcriptomes. The completeness of the final transcriptomes was examined using the 1,614
192 orthologues in the BUSCO (Benchmarking Universal Single Copy Orthologs) v5.2.2 set of
193 embryophyta_odb10 (Manni et al., 2021).
194
195 The transcriptomes were annotated with Gene Ontology (GO) and Kyoto Encyclopedia of Genes
196 and Genomes (KEGG) terms. The transcripts were aligned to the nr database that only includes
197 green plant subjects using the blastx function implemented by the software DIAMOND
198 v2.0.13.151 with parameters of -taxonlist 33090  --very-sensitive -f5. The .xml outputs were
199 used as input for mapping and annotation using the software Blast2GO (Götz et al., 2008). The
200 transcripts were translated to protein sequences using the software TransDecoder v5.5.0
201 (https://github.com/TransDecoder/TransDecoder/wiki) with default parameters to get the longest
202 ORF (open reading frame) per transcript, and then were annotated by the software InterProScan
203 v5.54-87.0 (Jones et al., 2014) with the parameters of -dra -f XML -goterms -pa. The final GO
204 terms were acquired by merging Blast2GO and InterProScan annotation outputs using the
205 software Blast2GO. To get the KEGG terms, the longest ORF per transcript were also annotated
206 using EggNOG-mapper v2.1.7 (Cantalapiedra et al., 2021) with parameters of -m diamond -
207 sensmode more-sensitive -tax_scope 33090 -tax_scope_mode 33090. The assembled
208 transcriptome and annotation were used in the downstream analyses as shown in the flowchart
209 (Supplementary Fig. S1).
210
211 *Ks analyses*

The level of synonymous substitutions (Ks) between two homologous sequences increases
approximately linearly with time. Thus pairs of paralogous genes within a species can be sorted
to estimate the relative ages of gene duplication (Blanc and Wolfe, 2004). To identify paralogous
gene pairs within a species, an all-versus-all blastp was performed within each species. The
peptide sequences (the longest ORF per transcript) of each species were used as input to run all-
versus-all blastp using the software DIAMOND v2.0.13.151 with parameters of -p 16 --max-
target-seqs 20 --evalue 1e-10 --outfmt 6 qseqid sseqid pident nident evalue. The blastp output
was filtered by removing self blastp results, keeping outputs with pident (percent of identical
matches) ≥20 and nident (number of identical matches) ≥50, and removing sequences with more
than 10 hits. The filtered paralogous gene pairs were aligned using the software MAFFT v7.490
(Katoh et al., 2002) with default parameters. The alignments were used as input for estimating
the number of synonymous substitutions per synonymous site (Ks) and the number of non-
synonymous substitutions per nonsynonymous site (Ka) using the program yn00 in the software
PAML v4.9j (Yang, 2007) with parameters of icode=0 weighting=0 commonf3×4=0. The Ks of
each paralogous gene pairs were acquired using the method by Yang and Nielsen (2000).
227
Ks can also be calculated using the orthologous gene pairs between species to estimate speciation
events (Blanc and Wolfe, 2004). To identify orthologous gene pairs between species, reciprocal
best hit blastp were run between the compared pairs. The peptide sequences (the longest ORF per
transcript) of the paired species were used as input to run blastp using the software DIAMOND
v2.0.13.151 with parameters of -p 16 --max-target-seqs 20 --evalue 1e-10 --outfmt 6 qseqid
sseqid pident nident evalue. The best hits between forward and reverse blastp for each compared
sequence pairs were retained, which were then used for sequence alignment and estimating Ka
and Ks as abovementioned.
236
The Ks distribution is proportional to the time lapse. Thus, the Ks distribution within species can
be used to infer the timing of WGD, and the Ks distribution between two different species can
infer time since speciation (Blanc and Wolfe, 2004). To plot the Ks distribution, the mixture
model was fit and plot for the Ks values ≥0.01 and ≤2 using the R package mixR (Yu, 2022).
The best model from a candidate of mixture models was selected using the "select" function
based on the information criterion BIC.
243
*Estimating duplication events and divergence time*
To infer duplication events, the species tree and gene trees were first inferred using STAG and
STRIDE algorithms implemented by the program OrthoFinder v2.5.4 (Emms and Kelly, 2017,
2018). The inferred species tree was similar to that expected given current knowledge of the
Angiosperm phylogeny with the exception of the placement of *Populus tremuloides* and *P.*
*grandidentata*. Thus, for downstream analysis and discussion, we removed these two species for
the species tree inference. The inferred species tree and gene trees were used to estimate
duplication events and divergence time. The gene duplication events on the species tree node
were reported if, at least 50% of the descendant species have retained both copies of the
duplicated gene. The divergence time was estimated using the software r8s v1.81 (Sanderson,
2003) with calibration points of 115 million years (MYA) between *Fraxinus nigra* and *Corylus*
*cornuta*, which were obtained from http://www.timetree.org/ (Kumar et al., 2017).
256
*Gene families and their evolutionary rate*

To infer gene families, the peptide sequences (the longest ORF per transcript) of the 20 studied
transcriptomes were used to construct orthogroups (gene families) using the program
OrthoFinder v2.5.4 with parameters of -S diamond  -t 16  -a 16. Genes assigned to each gene
family were counted for each species. The evolutionary rate of protein coding genes was
represented by the ratio of nonsynonymous substitutions per nonsynonymous site to synonymous
substitutions per synonymous site (Ka/Ks) with *Dillenia indica* (elephant apple) being the
outgroup and the twenty studied species being the ingroup. *D. indica* is an eudicot species,
belonging to the order Dilleniales and the RNA-seq data for this species were downloaded from
https://trace.ncbi.nlm.nih.gov/Traces/index.html?view=run_browser&acc=ERR2040190&disp
lay=data-access (Leebens-Mack et al., 2019). The raw sequences were filtered using the program
fastp with default parameters (Chen et al., 2018). The transcriptome assembly and protein
sequence prediction were performed as abovementioned, yielding 22,753 transcripts and 19,345
protein sequences. Then the reciprocal best hit blastp was performed between *D.indica* and each
of the twenty studied species to identify the orthologous pairs. The best hits between forward and
reverse blastp for each compared sequence pairs were retained, which were then used for
sequence alignment and estimating Ka/Ks as abovementioned. The mean Ka/Ks across genes for
each gene family per species was used to represent the evolutionary rate for each gene family.
Gene families with mean Ka/Ks >10 were filtered out. The graphs were plotted using the R
package ggplot2 and ggpubr (Wickham, 2016; Kassambara, 2020). The size and evolutionary
rate of gene families were correlated using scatter plot. All statistical analyses were performed in
R (R Core Team, 2021). Gene families with a size ≥5 and Ka/Ks >1 were highlighted for further
analysis. The enriched GO and KEGG terms were identified in these highlighted gene families.
The enriched GO terms of each category (biological process, molecular function, cellular
component) were separately detected using the "weight" algorithm with the Fisher ratio test
implemented by the R package "topGO" (Alexa and Rahnenfuhrer, 2018). Enriched KEGG
terms were detected using the hypergeometric method implemented by the R package "plyr"
(Wickham, 2011). Heatmaps were plot using the R package pheatmap (Kolde, 2019).
285
*Estimating gene family expansion and contraction*
To further study gene family changes, species tree and gene family counts were predicted and
used to estimate gene family expansion and contraction. All-versus-all blastp was performed to
find the most similar sequences for each sequence in the studied species using the program
DIAMOND (Buchfink et al., 2021). The program MCL (Enright et al., 2002) was used to find
clusters of similar sequences (gene families). The species tree was estimated using the program
FastTree (Price et al., 2009). The programs DIAMOND, MCL, and FastTree were implemented
by the program OrthoFinder v2.5.4, default parameters were used. Then, the species tree was
converted to a format for estimating ultrametric tree using the program CAFE5 (Mendes et al.,
2020) with the command python prep_r8s.py -s 60164 -p 'Fraxinus_nigra,Corylus_cornuta' -c
120. "60164" was the number of sites in the alignment. "-c 120" represents a median of 120
MYA divergence time, it is the calibration points between *Fraxinus nigra* and *Corylus cornuta*
checked from http://www.timetree.org/. Then the ultrametric tree was estimated using the
program r8s (Sanderson, 2003) with default parameters. The ultrametric tree and gene family
counts were used as input for the program CAFE5 (Mendes et al., 2020). Since CAFE5 was not
recommended to analyze recent polyploid species, only diploid study species were retained for
running CAFE5, including *Cornus alternifolia*, *Fraxinus nigra*, *Dirca palustris*, *Tilia americana*,
*Acer saccharum*, *Acer spicatum*, *Amelanchier laevis*, *Quercus rubra*, *Betula papyrifera*, *Alnus*

*incana*, *Corylus cornuta*, *Ostrya virginiana*, and *Ribes cynosbati*. Since assembly errors might
cause the observed number of gene copies in gene families to deviate from the true ones and
overestimation of the birth-death parameter $\lambda$ (number of gene gains and losses per gene per
million years), an error model was estimated first with the ultrametric tree and gene family
counts using CAFE5 with the parameters of "-c 16 -p -e". This error model along with the
ultrametric tree and gene family counts were used to estimate the birth-death parameter $\lambda$, which
represents the probability that any gene will be gained or lost. Multiple $\lambda$ mode was used with
three different rates of gene family evolution assigned as below:

((Cornus_alternifolia:3,Fraxinus_nigra:3):3,((((Dirca_palustris:2,Tilia_americana:2):2,(Acer_sac
charum:2,Acer_spicatum:2):2):2,((((Betula_papyrifera:2,Alnus_incana:2):2,(Corylus_cornuta:2,
Ostrya_virginiana:2):2):2,Quercus_rubra:2):2,Amelanchier_laevis:2):2):2,Ribes_cynosbati:1):1);

The numbers 1, 2, 3 represent three different gene family evolutionary rates that were shared by
different clades. This tree file was used as input along with the ultrametric tree, gene family
counts, and error model using CAFE5 with the parameters of -c 16 -p -k 3. Gene families or
parent nodes with a significantly greater rate of evolution (family-wide p-value <0.01), in other
words, rapidly evolving gene families, were counted for each species and each parent node.
These gene families significantly violated a random gene birth and death model. Thus, the null
hypothesis of neutral evolution was rejected and these families could be recognized as fast
evolving gene families (Hahn et al., 2005). Enriched GO and KEGG terms were detected in the
fast expanding and fast contracting gene families, respectively.

**Results**

*Ks analyses*

We acquired an average of 41 million raw reads for each sample. The total number of assembled
transcripts ranged between 19,459 and 31,804 (Supplementary Table S1). We retrieved 76 ~
88% of the BUSCOs in the assembled transcriptomes (Supplementary Fig. S2). Most species
have an initial Ks peak around 0.02~0.04, except *Fraxinus nigra* and *Tilia americana*, which
have the initial peak around 0.1 and 0.09, respectively (Fig. 2). Such initial peaks represent most
recently duplicate genes. Another discernable separate peak shared by the studied species is
around 0.5 ~ 0.7. Such shared peaks by all species may correspond to large-scale duplication
events, perhaps the common WGDs across core eudicots. Some species have other discernable
peaks, like *Populus grandidentata* and *Populus tremuloides*, which have discernable peaks
around 0.18. Such unique peaks may represent ancient duplication events specific to certain
species and different evolutionary patterns across species. The two *Prunus* species and the two
*Populus* species have similar Ks shapes, suggestive of common WGDs before the two species
diverged from the same ancestors. By examining the orthologous Ks between the two species,
we found a high peak close to zero, which may represent the duplication events after the WGDs
and divergence (Supplementary Fig. S3).

Substituions per synonymous site (Ks)

**Fig. 2. Distribution of Ks (the level of synonymous substitutions). A mixture model was fit and plot for the Ks values ≥0.01 and ≤2 of each species using the R package mixR.**

*Duplication events and divergence time*

Using a species tree with estimated branch lengths, we inferred divergence times and duplication events (with at least 50 % support rate, Fig. 3). Most duplication events (n=1843) occurred at the divergence of rosids and asterids, which may correspond to common WGD across eudicots species. The ancestor of *Populus tremuloides* and *Populus grandidentata* had 1,625 duplication events (Supplementary Fig. S4), which is consistent with the recent Ks peaks within these two species. The divergence time of the two asterids species *Cornus alternifolia* and *Fraxinus nigra* from a common ancestor was about 104 MYA (Fig. 3). The divergence time of the studied malvids and fabids from a common ancestor with Saxifragles was about 100 MYA.

**Fig. 3. Divergence time in million years and gene duplication events with 50 % support rate. Node numbers were followed by number of gene duplication events.**

*Relationship between gene family size and evolutionary rate*

OrthoFinder assigned 393,796 genes (96.2 % of total) to 21,900 orthogroups (gene families). The mean and median gene family size is 17.3 and 12, respectively. We found that 3,060 families present in all species, whereas 19 consisted entirely of single-copy genes.

After identifying reciprocal best blastp hits with the outgroup *D.indica*, we retained a total of 6903 ~ 7302 gene families for the studied species to calculate the relationship between gene family size and evolutionary rate. Gene families with large sizes tended to have low Ka/Ks, yet there are a few exceptions. We highlighted the gene families with a size ≥5 and Ka/Ks >1 in red (Fig. 4a). We tested the enriched KEGG and GO terms in these highlighted gene families (Fig. 4b and Supplementary Fig. S5, see the whole list of GO enrichment in Supplementary Table S2). The most highly shared enriched KEGG and GO terms are related to plant-pathogen interactions, signaling and secondary metabolite biosynthesis.

385
386 **Fig. 4. Scatter plot between gene size and Ka/Ks per gene family (a) and KEGG pathway**
387 **terms enrichment (b) in gene families with a size ≥5 and Ka/Ks >1. The color bar in b)**
388 **represents the adjusted p-values of KEGG enrichment test.**

389
390

391 *Gene family expansion and contraction*

392
393

We inferred the birth-death parameter $\lambda$ (number of gene gains and losses per gene per million
years), which is specified in different clades, using the program CAFE5. $\lambda 1$ (saxifragales)
=0.0034, $\lambda 2$ (rosids) =0.0042, $\lambda 3$ (asterids) =0.0027, which means that the lineages leading to
rosids have the highest gene family turnover rates, followed by saxifragales, and then by asterids.
The percentage and numbers of gene family expansion and contraction are shown in Fig. 5a and
Supplementary Fig. S6. All of the studied species have more rapidly expanding than rapidly
contracting gene families (family-wide p-value <0.01, Fig. 5b), which may contribute to their
lineage-specific novelty.
402

403
404 **Fig. 5. The percentage of gene family expansion and contraction (a) and the number of fast**
405 **evolving gene families (b), with + representing fast expanding gene family and –**
406 **representing fast contracting gene family.**

407
408
We tested the enriched GO and KEGG terms in the fast evolving (expanding or contracting)
gene families. All of the studied species shared the enriched KEGG pathway term plant-pathogen
interaction (Fig. 6). This term was detected in the fast-expanding gene families of *Cornus*
*alternifolia*, *Tilia americana*, *Acer spicatum*, *Betula papyrifera*, *Quercus rubra*, *Amelanchier*
*laevis*, *Ribes cynosbati*; and this term was detected in the fast-contracting gene families of
*Fraxinus nigra*, *Dirca palustris*, *Alnus incana*, *Corylus cornuta*, *Ostrya virginiana*; also, this
term was detected in both of the fast expanding and fast contracting gene families of *Acer*
*saccharum*. The gene families assigned to the plant-pathogen interaction term encode proteins
(Supplementary Fig. S7) related to disease resistance protein, leucine-rich repeat receptor like
protein kinase and others. Some species shared the enriched KEGG pathway terms Tryptophan
metabolism and MAPK signaling pathway (Fig. 6).

420
421

422
423
424  **Fig. 6. Enriched KEGG pathway terms in the studied diploid species. Red represents fast**
425  **expanding gene families; blue represents fast contracting gene families; white represents**
426  **both fast expanding and contracting gene families.**
427
428
**Discussion**
Gene duplication events are prevalent in all organisms and provide genetic variation for
evolutionary innovation and adaption to fluctuating conditions (Zhang, 2003; Van de Peer et al.,
2009; Panchy et al., 2016). Research on the mechanisms of gene duplication as well as the
origin, maintenance, and functional roles of genetic diversity in forest tree species are few due to
the limited genome and transcriptome data resources. In the present study, we assembled twenty
transcriptomes from forest trees that co-occur. We identified the signal of common WGDs
shared by core eudicots, and the signal of recent duplication events specific to certain species.
We found gene families involved in plant-pathogen interaction, signaling, and metabolite
biosynthesis actively remain extensive variation. We also found fast expansion or fast
contraction in plant-pathogen interaction genes amongst the studied diploid species. In the
following sections, we will discuss these results in detail.
441
*Tempo and mode of gene duplication*
The Ks distribution plots confirmed the widespread paleopolyploidy in plants, that's said,
today's diploid species might be polyploid (paleopolyploid) at some points in evolutionary
history (Blanc and Wolfe, 2004). The abundant duplication events provide resources for genetic
variation. It is worth noting that Ks distribution is not suitable to infer very ancient events due to
saturation with multiple substitutions (Blanc and Wolfe, 2004), moreover, different peaks might
not be separable from each other, thus it is possible that we missed identifying some duplication
events in the present study. Collinearity- or synteny-based methods along with chromosome-
level reference genome assembly is alternative method to identify ancient WGD, yet high quality
genome assembly is not easy to obtain (Zwaenepoel and Van de Peer, 2019).
452

The predicted species tree (Fig. 3) generally agrees with current phylogenetic hypotheses for the
angiosperms with the exception of the two *Populus* species, which is probably caused by the
incomplete gene sets obtained from one leaf RNA-seq sample and/or interspecific gene flow.
The duplication events at the root (Fig. 3) of the tree is likely to be related to the gamma WGD
before the divergence of rosids and asterids. Such large-scale duplication may contribute to
genome structure variation, organismal complexity, and rapid speciation in angiosperm species.
WGD events followed by different evolutionary events might also occur in the common
ancestors for *Acer*, *Prunus*, and *Populus* species. For *Acer*, it is likely that *Acer spicatum*
experienced a diploidization event to revert to diploid status, while *Acer rubrum* maintained a
polyploid status.
463
Divergence time mostly agrees with the previous reports, which stated rosids and asterids
diverged about 125 MYA, malvids and fabids diverged about 110 MYA (Zeng et al., 2017).
However, divergence estimate in the present study is rather rough. Further studies need to
consider using dense taxon sampling, diverse markers like nuclear internal transcribed spacer or
chloroplast DNA fragments, and molecular dating with fossil corrections methods to better
resolve the evolutionary relationships.
470
*Sustained large size gene families*
Though most duplicate genes were eliminated or pseudonized (Panchy et al., 2016), a small
portion of gene families can actively maintain duplicate genes as selection favors the fitness
related neofunctionalization. The main functions of these sustained large size gene families are
mainly involved in plant-pathogen interaction, signaling, and metabolite biosynthesis (Fig. 4b &
Supplementary Fig. S5). This is consistent with previous report that genes encoding functions
like signal transduction and stress response tend to have paralogs (Panchy et al., 2016), and
genes related to the function like transcription regulation and signal transduction had higher
possibility to survive in multiple rounds of genome duplication (Seoighe and Gehring, 2004).
Plants constantly combat with pathogens and pests. The major class of plant disease resistance
(R) genes contain a nucleotide-binding site (NBS) and C-terminal leucin-rich repeats (LRRs).
Tandem and segmental gene duplication explain much of the diversification among R genes
(Meyers et al., 2003; Leister, 2004; Andersen et al., 2020), possibly because they have
experienced frequent deletion and duplication throughout evolution, and thus are highly variable
(Hanada et al., 2008; Gao et al., 2018). Plant secondary metabolites, such as monoterpenoid,
flavonoid, benzoxazinoid, play important roles in defense against predation and pathogens
(Falcone Ferreyra et al., 2012; de Bruijn et al., 2018; Bergman et al., 2020). The regulation of
plant secondary metabolism requires the intimation of signal transduction network that leads to
activation of biosynthetic genes (Zhao et al., 2005).
490
Gene duplication and deletion give rise to gene family expansion and contraction. The high gene
gain rate likely reflects the occurrence of multiple WGDs and frequent SSDs (Carretero-Paulet et
al., 2015). While for the other species with contraction outnumbered expansion, the pattern is
mainly caused by gene loss and retention of specific gene families followed the common WGDs
preceded their common ancestor (Supplementary Fig. S4). We found a high gene family turnover
rate in rosids, but we must caution that the unbalanced taxon sampling in our study species might
cause this high turnover value. Moreover, future studies should consider distinguishing the

branches with or without recent WGDs, as the branches with recent WGDs tend to have a high
gene turnover rate (Casola and Koralewski, 2018).
500
*Rapid evolution in plant-pathogen interaction genes*
We found rapid evolution of genes involved in plant-pathogen interaction, tryptophan
metabolism and MAPK signaling pathway (Fig. 6) in the adaptive expansion and adaptive
contraction gene families across the studied species. The specific expanding or contracting gene
families involved in plant-pathogen interaction in each species differed from each other,
implying lineage-specificity in changed gene families (Supplementary Fig. S7). This result is
consistent with previous studies, which found that fast evolving gene families were mostly
groups with functions related with immune defense/stress response, metabolism, cell signaling,
chemoreception, and reproduction (Demuth and Hahn, 2009). Additionally, a study on *Juglans*
(walnuts) found notable contractions in gene families that are involved in disease resistance and
other abiotic stress response (Trouern-Trend et al., 2020). Both metabolism and MAPK signaling
pathway are associated with plant defense. MAPK cascades sense pathogen/microbe-associated
molecular patters and pathogen effectors, and thus play pivotal roles in signaling plant defense
against pathogen attack (Meng and Zhang, 2013). Plant secondary metabolites contribute to plant
immunity as antibiotics, or to control callose deposition and programmed cell death (Piasecka et
al., 2015). The biotic stress exerted by pathogen or pests is a major source of selection pressure
on the studied trees, which drive the rapid radiation of diverse and large plant immune receptor
gene repertoires. Expansion and/or contraction of cell-surface receptor genes correlate with the
expansion and/or contraction of intracellular receptor genes (Ngou et al., 2022).
520
**Conclusions**
In summary, we found the common WGDs shared by all the studied species as well as signals of
recent duplication events specific to certain species, which are suggestive of different
evolutionary patterns. Gene families involved in plant-pathogen interaction, signaling, and
metabolite biosynthesis actively remain extensive variation and are fast expanding or contracting
amongst the studied species. This knowledge calls for an emphasis of gene family size variation,
i.e. copy number variation, when studying forest genetic diversity, which will inform forest
adaptive strategy. Future studies may focus on classification of duplications in forest tree
species, and investigate how the duplications give rise to species-specific innovation and
adaption.
531
532
**Data availability**
The raw RNA-seq data were deposited in NCBI SRA (accession number: PRJNA896247;
http://www.ncbi.nlm.nih.gov/sra). The assembled transcriptomes and their annotation were
deposited in Dryad https://doi.org/10.5061/dryad.sj3tx968b. The scripts used for analyses were
deposited in https://github.com/Mengmeng-Lu/Gene-duplication-and-gene-expansion-analyses-
in-twenty-tree-species.
539

**Acknowledgements**
We thank the Center for Research Computing at University of Notre Dame for high
performance computing support. We specially acknowledge the assistance of Dr. Dodi
Heryadi, Dr. Scott Hampton, and Júlia Simon. The research was performed at the University of

Notre Dame Environmental Research Center that has been generously supported by the
Bernard J. Hank Family Endowment. The research was permitted under UNDERC research
permit number: UNDERC-2021-2.

547
548

549 **Funding**
550 MM and NGS were supported by the US National Science Foundation Dimensions of
551 Biodiversity grant (DEB-2124466). JY and MC were supported by a NSFC China-US
552 Dimensions of Biodiversity Grant (DEB-32061123003).

553
554

555 **Conflicts of interest**
556 The authors declare no conflict of interest.

557
558
559

**Literature cited**

Alexa, A., and Rahnenfuhrer, J. 2018. topGO: Enrichment Analysis for Gene Ontology. R package
version 2.34.0.
Allen, C.D., Macalady, A.K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M.,
Kitzberger, T., Rigling, A., Breshears, D.D., Hogg, E.H.T., Gonzalez, P., Fensham, R., Zhang,
Z., Castro, J., Demidova, N., Lim, J.-H., Allard, G., Running, S.W., Semerci, A., and Cobb,
N. 2010. A global overview of drought and heat-induced tree mortality reveals emerging
climate change risks for forests. For Ecol Manage 259:660-684.
AMBORELLA GENOME PROJECT, Albert, V.A., Barbazuk, W.B., dePamphilis, C.W., Der, J.P.,
Leebens-Mack, J., Ma, H., Palmer, J.D., Rounsley, S., Sankoff, D., Schuster, S.C., Soltis,
D.E., Soltis, P.S., Wessler, S.R., Wing, R.A., Albert, V.A., Ammiraju, J.S.S., Barbazuk, W.B.,
Chamala, S., Chanderbali, A.S., dePamphilis, C.W., Der, J.P., Determann, R., Leebens-
Mack, J., Ma, H., Ralph, P., Rounsley, S., Schuster, S.C., Soltis, D.E., Soltis, P.S., Talag, J.,
Tomsho, L., Walts, B., Wanke, S., Wing, R.A., Albert, V.A., Barbazuk, W.B., Chamala, S.,
Chanderbali, A.S., Chang, T.-H., Determann, R., Lan, T., Soltis, D.E., Soltis, P.S., Arikit, S.,
Axtell, M.J., Ayyampalayam, S., Barbazuk, W.B., Burnette, J.M., Chamala, S., De Paoli, E.,
dePamphilis, C.W., Der, J.P., Estill, J.C., Farrell, N.P., Harkess, A., Jiao, Y., Leebens-Mack,
J., Liu, K., Mei, W., Meyers, B.C., Shahid, S., Wafula, E., Walts, B., Wessler, S.R., Zhai, J.,
Zhang, X., Albert, V.A., Carretero-Paulet, L., dePamphilis, C.W., Der, J.P., Jiao, Y.,
Leebens-Mack, J., Lyons, E., Sankoff, D., Tang, H., Wafula, E., Zheng, C., Albert, V.A.,
Altman, N.S., Barbazuk, W.B., Carretero-Paulet, L., dePamphilis, C.W., Der, J.P., Estill,
J.C., Jiao, Y., Leebens-Mack, J., Liu, K., Mei, W., Wafula, E., Altman, N.S., Arikit, S., Axtell,
M.J., Chamala, S., Chanderbali, A.S., Chen, F., Chen, J.-Q., Chiang, V., De Paoli, E.,
dePamphilis, C.W., Der, J.P., Determann, R., Fogliani, B., Guo, C., Harholt, J., Harkess, A.,
Job, C., Job, D., Kim, S., Kong, H., Leebens-Mack, J., Li, G., Li, L., Liu, J., Ma, H., Meyers,
B.C., Park, J., Qi, X., Rajjou, L., Burtet-Sarramegna, V., Sederoff, R., Shahid, S., Soltis, D.E.,
Soltis, P.S., Sun, Y.-H., Ulvskov, P., Villegente, M., Xue, J.-Y., Yeh, T.-F., Yu, X., Zhai, J.,
Acosta, J.J., Albert, V.A., Barbazuk, W.B., Bruenn, R.A., Chamala, S., de Kochko, A.,

dePamphilis, C.W., Der, J.P., Herrera-Estrella, L.R., Ibarra-Laclette, E., Kirst, M., Leebens-
Mack, J., Pissis, S.P., Poncet, V., Schuster, S.C., Soltis, D.E., Soltis, P.S., and Tomsho, L.
2013. The Amborella Genome and the Evolution of Flowering Plants. Science
342:1241089.
Andersen, E.J., Nepal, M.P., Purintun, J.M., Nelson, D., Mermigka, G., and Sarris, P.F. 2020.
Wheat Disease Resistance Genes and Their Diversification Through Integrated Domain
Fusions. Front Genet 11:898.
Bailey, J.A., Gu, Z., Clark, R.A., Reinert, K., Samonte, R.V., Schwartz, S., Adams, M.D., Myers,
E.W., Li, P.W., and Eichler, E.E. 2002a. Recent segmental duplications in the human
genome. Science 297:1003-1007.
Bailey, J.A., Gu, Z., Clark, R.A., Reinert, K., Samonte, R.V., Schwartz, S., Adams, M.D., Myers,
E.W., Li, P.W., and Eichler, E.E. 2002b. Recent Segmental Duplications in the Human
Genome. Science 297:1003-1007.
Bergman, M.E., Chávez, Á., Ferrer, A., and Phillips, M.A. 2020. Distinct metabolic pathways drive
monoterpenoid biosynthesis in a natural population of Pelargonium graveolens. Journal
of Experimental Botany 71:258-271.
Blanc, G., and Wolfe, K.H. 2004. Widespread Paleopolyploidy in Model Plant Species Inferred
from Age Distributions of Duplicate Genes[W]. The Plant Cell 16:1667-1678.
Brosius, J. 1991. Retroposons—Seeds of Evolution. Science 251:753-753.
Buchfink, B., Reuter, K., and Drost, H.-G. 2021. Sensitive protein alignments at tree-of-life scale
using DIAMOND. Nature Methods 18:366-368.
Cantalapiedra, C.P., Hernández-Plaza, A., Letunic, I., Bork, P., and Huerta-Cepas, J. 2021.
eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain
Prediction at the Metagenomic Scale. Mol Biol Evol 38:5825-5829.
Carretero-Paulet, L., Librado, P., Chang, T.-H., Ibarra-Laclette, E., Herrera-Estrella, L., Rozas, J.,
and Albert, V.A. 2015. High Gene Family Turnover Rates and Gene Space Adaptation in
the Compact Genome of the Carnivorous Plant Utricularia gibba. Molecular Biology and
Evolution 32:1284-1295.
Casola, C., and Koralewski, T.E. 2018. Pinaceae show elevated rates of gene turnover that are
robust to incomplete gene annotation. Plant J.
Cazzolla Gatti, R., Reich, P.B., Gamarra, J.G.P., Crowther, T., Hui, C., Morera, A., Bastin, J.-F., de-
Miguel, S., Nabuurs, G.-J., Svenning, J.-C., Serra-Diaz, J.M., Merow, C., Enquist, B.,
Kamenetsky, M., Lee, J., Zhu, J., Fang, J., Jacobs, D.F., Pijanowski, B., Banerjee, A.,
Giaquinto, R.A., Alberti, G., Almeyda Zambrano, A.M., Alvarez-Davila, E., Araujo-
Murakami, A., Avitabile, V., Aymard, G.A., Balazy, R., Baraloto, C., Barroso, J.G., Bastian,
M.L., Birnbaum, P., Bitariho, R., Bogaert, J., Bongers, F., Bouriaud, O., Brancalion, P.H.S.,
Brearley, F.Q., Broadbent, E.N., Bussotti, F., Castro da Silva, W., César, R.G., Češljar, G.,
Chama Moscoso, V., Chen, H.Y.H., Cienciala, E., Clark, C.J., Coomes, D.A., Dayanandan,
S., Decuyper, M., Dee, L.E., Del Aguila Pasquel, J., Derroire, G., Djuikouo, M.N.K., Van Do,
T., Dolezal, J., Đorđević, I.Đ., Engel, J., Fayle, T.M., Feldpausch, T.R., Fridman, J.K., Harris,
D.J., Hemp, A., Hengeveld, G., Herault, B., Herold, M., Ibanez, T., Jagodzinski, A.M.,
Jaroszewicz, B., Jeffery, K.J., Johannsen, V.K., Jucker, T., Kangur, A., Karminov, V.N.,
Kartawinata, K., Kennard, D.K., Kepfer-Rojas, S., Keppel, G., Khan, M.L., Khare, P.K.,
Kileen, T.J., Kim, H.S., Korjus, H., Kumar, A., Kumar, A., Laarmann, D., Labrière, N., Lang,

M., Lewis, S.L., Lukina, N., Maitner, B.S., Malhi, Y., Marshall, A.R., Martynenko, O.V.,
Monteagudo Mendoza, A.L., Ontikov, P.V., Ortiz-Malavasi, E., Pallqui Camacho, N.C.,
Paquette, A., Park, M., Parthasarathy, N., Peri, P.L., Petronelli, P., Pfautsch, S., Phillips,
O.L., Picard, N., Piotto, D., Poorter, L., Poulsen, J.R., Pretzsch, H., Ramírez-Angulo, H.,
Restrepo Correa, Z., Rodeghiero, M., Rojas Gonzáles, R.D.P., Rolim, S.G., Rovero, F.,
Rutishauser, E., Saikia, P., Salas-Eljatib, C., Schepaschenko, D., Scherer-Lorenzen, M.,
Šebeň, V., Silveira, M., Slik, F., Sonké, B., Souza, A.F., Stereńczak, K.J., Svoboda, M.,
Taedoumg, H., Tchebakova, N., Terborgh, J., Tikhonova, E., Torres-Lezama, A., van der
Plas, F., Vásquez, R., Viana, H., Vibrans, A.C., Vilanova, E., Vos, V.A., Wang, H.-F.,
Westerlund, B., White, L.J.T., Wiser, S.K., Zawiła-Niedźwiecki, T., Zemagho, L., Zhu, Z.-X.,
Zo-Bi, I.C., and Liang, J. 2022. The number of tree species on Earth. Proceedings of the
National Academy of Sciences 119:e2115329119.
Chen, S., Zhou, Y., Chen, Y., and Gu, J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics 34:i884-i890.
Conant, G.C., Birchler, J.A., and Pires, J.C. 2014. Dosage, duplication, and diploidization:
clarifying the interplay of multiple models for duplicate gene evolution over time.
Current Opinion in Plant Biology 19:91-98.
Cortés, A.J., Restrepo-Montoya, M., and Bedoya-Canas, L.E. 2020. Modern Strategies to Assess
and Breed Forest Tree Adaptation to Changing Climate. Front Plant Sci 11:583323.
Cui, L., Wall, P.K., Leebens-Mack, J.H., Lindsay, B.G., Soltis, D.E., Doyle, J.J., Soltis, P.S., Carlson,
J.E., Arumuganathan, K., Barakat, A., Albert, V.A., Ma, H., and dePamphilis, C.W. 2006.
Widespread genome duplications throughout the history of flowering plants. Genome
Res 16:738-749.
de Bruijn, W.J.C., Gruppen, H., and Vincken, J.-P. 2018. Structure and biosynthesis of
benzoxazinoids: Plant defence metabolites with potential as antimicrobial scaffolds.
Phytochemistry 155:233-243.
De Smet, R., Adams, K.L., Vandepoele, K., Van Montagu, M.C.E., Maere, S., and Van de Peer, Y.
2013. Convergent gene loss following gene and genome duplications creates single-copy
families in flowering plants. Proceedings of the National Academy of Sciences 110:2898-
2903.
Demuth, J.P., and Hahn, M.W. 2009. The life and death of gene families. Bioessays 31:29-39.
Diss, G., Gagnon-Arsenault, I., Dion-Coté, A.-M., Vignaud, H., Ascencio, D.I., Berger, C.M., and
Landry, C.R. 2017. Gene duplication can impart fragility, not robustness, in the yeast
protein interaction network. Science 355:630-634.
Dodds, P.N., and Rathjen, J.P. 2010. Plant immunity: towards an integrated view of plant–
pathogen interactions. Nature Reviews Genetics 11:539-548.
Emms, D.M., and Kelly, S. 2017. STRIDE: Species Tree Root Inference from Gene Duplication
Events. Mol Biol Evol 34:3267-3278.
Emms, D.M., and Kelly, S. 2018. STAG: Species Tree Inference from All Genes (bioRxiv).
Enright, A.J., Van Dongen, S., and Ouzounis, C.A. 2002. An efficient algorithm for large-scale
detection of protein families. Nucleic Acids Research 30:1575-1584.
Falcone Ferreyra, M.L., Rius, S., and Casati, P. 2012. Flavonoids: biosynthesis, biological
functions, and biotechnological applications. Frontiers in Plant Science 3.
FAO. 2020. Global Forest Resources Assessment 2020: Main report. Rome.

Force, A., Lynch, M., Pickett, F.B., Amores, A., Yan, Y.L., and Postlethwait, J. 1999. Preservation
of duplicate genes by complementary, degenerate mutations. Genetics 151:1531-
1545.
Fu, L., Niu, B., Zhu, Z., Wu, S., and Li, W. 2012. CD-HIT: accelerated for clustering the next-
generation sequencing data. Bioinformatics 28:3150-3152.
Gamfeldt, L., Snäll, T., Bagchi, R., Jonsson, M., Gustafsson, L., Kjellander, P., Ruiz-Jaen, M.C.,
Fröberg, M., Stendahl, J., Philipson, C.D., Mikusiński, G., Andersson, E., Westerlund, B.,
Andrén, H., Moberg, F., Moen, J., and Bengtsson, J. 2013. Higher levels of multiple
ecosystem services are found in forests with more tree species. Nature Communications
4:1340.
Gao, Y., Wang, W., Zhang, T., Gong, Z., Zhao, H., and Han, G.-Z. 2018. Out of Water: The Origin
and Early Diversification of Plant R-Genes. Plant Physiology 177:82-89.
Götz, S., García-Gómez, J.M., Terol, J., Williams, T.D., Nagaraj, S.H., Nueda, M.J., Robles, M.,
Talón, M., Dopazo, J., and Conesa, A. 2008. High-throughput functional annotation and
data mining with the Blast2GO suite. Nucleic Acids Res. 36:3420-3435.
Hahn, M.W., De Bie, T., Stajich, J.E., Nguyen, C., and Cristianini, N. 2005. Estimating the tempo
and mode of gene family evolution from comparative genomic data. Genome Res
15:1153-1160.
Hammond, W.M., Williams, A.P., Abatzoglou, J.T., Adams, H.D., Klein, T., López, R., Sáenz-
Romero, C., Hartmann, H., Breshears, D.D., and Allen, C.D. 2022. Global field
observations of tree die-off reveal hotter-drought fingerprint for Earth's forests. Nature
Communications 13:1761.
Hanada, K., Zou, C., Lehti-Shiu, M.D., Shinozaki, K., and Shiu, S.H. 2008. Importance of lineage-
specific expansion of plant tandem duplicates in the adaptive response to
environmental stimuli. Plant Physiol 148:993-1003.
Hurley, B.P., Garnas, J., Wingfield, M.J., Branco, M., Richardson, D.M., and Slippers, B. 2016.
Increasing numbers and intercontinental spread of invasive insects on eucalypts.
Biological Invasions 18:921-933.
Isabel, N., Holliday, J.A., and Aitken, S.N. 2020. Forest genomics: Advancing climate adaptation,
forest health, productivity, and conservation. Evolutionary Applications 13:3-10.
Jiang, N., Bao, Z., Zhang, X., Eddy, S.R., and Wessler, S.R. 2004. Pack-MULE transposable
elements mediate gene evolution in plants. Nature 431:569-573.
Jiao, Y., Leebens-Mack, J., Ayyampalayam, S., Bowers, J.E., McKain, M.R., McNeal, J., Rolf, M.,
Ruzicka, D.R., Wafula, E., Wickett, N.J., Wu, X., Zhang, Y., Wang, J., Zhang, Y., Carpenter,
E.J., Deyholos, M.K., Kutchan, T.M., Chanderbali, A.S., Soltis, P.S., Stevenson, D.W.,
McCombie, R., Pires, J.C., Wong, G.K., Soltis, D.E., and Depamphilis, C.W. 2012. A
genome triplication associated with early diversification of the core eudicots. Genome
Biol 13:R3.
Johri, P., Gout, J.-F., Doak, T.G., and Lynch, M. 2022. A Population-Genetic Lens into the Process
of Gene Loss Following Whole-Genome Duplication. Molecular Biology and Evolution
39:msac118.
Jones, P., Binns, D., Chang, H.Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J.,
Mitchell, A., Nuka, G., Pesseat, S., Quinn, A.F., Sangrador-Vegas, A., Scheremetjew, M.,

Yong, S.Y., Lopez, R., and Hunter, S. 2014. InterProScan 5: Genome-scale protein
function classification. Bioinformatics 30:1236-1240.
Kassambara, A. 2020. ggpubr: 'ggplot2' Based Publication Ready Plots. https://CRAN.R-
project.org/package=ggpubr.
Katoh, K., Misawa, K., Kuma, K.i., and Miyata, T. 2002. MAFFT: a novel method for rapid
multiple sequence alignment based on fast Fourier transform. Nucleic Acids Research
30:3059-3066.
Kirilenko, A.P., and Sedjo, R.A. 2007. Climate change impacts on forestry. Proceedings of the
National Academy of Sciences 104:19697-19702.
Kolde, R. 2019. pheatmap: Pretty Heatmaps. R package version 1.0.12, https://CRAN.R-
project.org/package=pheatmap.
Kumar, S., Stecher, G., Suleski, M., and Hedges, S.B. 2017. TimeTree: A Resource for Timelines,
Timetrees, and Divergence Times. Molecular Biology and Evolution 34:1812-1819.
Kuzmin, E., VanderSluis, B., Nguyen Ba, A.N., Wang, W., Koch, E.N., Usaj, M., Khmelinskii, A.,
Usaj, M.M., van Leeuwen, J., Kraus, O., Tresenrider, A., Pryszlak, M., Hu, M.-C., Varriano,
B., Costanzo, M., Knop, M., Moses, A., Myers, C.L., Andrews, B.J., and Boone, C. 2020.
Exploring whole-genome duplicate gene retention with complex genetic interaction
analysis. Science 368:eaaz5667.
Leebens-Mack, J.H., Barker, M.S., Carpenter, E.J., Deyholos, M.K., Gitzendanner, M.A., Graham,
S.W., Grosse, I., Li, Z., Melkonian, M., Mirarab, S., Porsch, M., Quint, M., Rensing, S.A.,
Soltis, D.E., Soltis, P.S., Stevenson, D.W., Ullrich, K.K., Wickett, N.J., DeGironimo, L.,
Edger, P.P., Jordon-Thaden, I.E., Joya, S., Liu, T., Melkonian, B., Miles, N.W., Pokorny, L.,
Quigley, C., Thomas, P., Villarreal, J.C., Augustin, M.M., Barrett, M.D., Baucom, R.S.,
Beerling, D.J., Benstein, R.M., Biffin, E., Brockington, S.F., Burge, D.O., Burris, J.N., Burris,
K.P., Burtet-Sarramegna, V., Caicedo, A.L., Cannon, S.B., Çebi, Z., Chang, Y., Chater, C.,
Cheeseman, J.M., Chen, T., Clarke, N.D., Clayton, H., Covshoff, S., Crandall-Stotler, B.J.,
Cross, H., dePamphilis, C.W., Der, J.P., Determann, R., Dickson, R.C., Di Stilio, V.S., Ellis,
S., Fast, E., Feja, N., Field, K.J., Filatov, D.A., Finnegan, P.M., Floyd, S.K., Fogliani, B.,
García, N., Gâteblé, G., Godden, G.T., Goh, F., Greiner, S., Harkess, A., Heaney, J.M.,
Helliwell, K.E., Heyduk, K., Hibberd, J.M., Hodel, R.G.J., Hollingsworth, P.M., Johnson,
M.T.J., Jost, R., Joyce, B., Kapralov, M.V., Kazamia, E., Kellogg, E.A., Koch, M.A., Von
Konrat, M., Könyves, K., Kutchan, T.M., Lam, V., Larsson, A., Leitch, A.R., Lentz, R., Li, F.-
W., Lowe, A.J., Ludwig, M., Manos, P.S., Mavrodiev, E., McCormick, M.K., McKain, M.,
McLellan, T., McNeal, J.R., Miller, R.E., Nelson, M.N., Peng, Y., Ralph, P., Real, D., Riggins,
C.W., Ruhsam, M., Sage, R.F., Sakai, A.K., Scascitella, M., Schilling, E.E., Schlösser, E.-M.,
Sederoff, H., Servick, S., Sessa, E.B., Shaw, A.J., Shaw, S.W., Sigel, E.M., Skema, C., Smith,
A.G., Smithson, A., Stewart, C.N., Stinchcombe, J.R., Szövényi, P., Tate, J.A., Tiebel, H.,
Trapnell, D., Villegente, M., Wang, C.-N., Weller, S.G., Wenzel, M., Weststrand, S.,
Westwood, J.H., Whigham, D.F., Wu, S., Wulff, A.S., Yang, Y., Zhu, D., Zhuang, C., Zuidof,
J., Chase, M.W., Pires, J.C., Rothfels, C.J., Yu, J., Chen, C., Chen, L., Cheng, S., Li, J., Li, R.,
Li, X., Lu, H., Ou, Y., Sun, X., Tan, X., Tang, J., Tian, Z., Wang, F., Wang, J., Wei, X., Xu, X.,
Yan, Z., Yang, F., Zhong, X., Zhou, F., Zhu, Y., Zhang, Y., Ayyampalayam, S., Barkman, T.J.,
Nguyen, N.-p., Matasci, N., Nelson, D.R., Sayyari, E., Wafula, E.K., Walls, R.L., Warnow,
T., An, H., Arrigo, N., Baniaga, A.E., Galuska, S., Jorgensen, S.A., Kidder, T.I., Kong, H., Lu-

Irving, P., Marx, H.E., Qi, X., Reardon, C.R., Sutherland, B.L., Tiley, G.P., Welles, S.R., Yu,
R., Zhan, S., Gramzow, L., Theißen, G., Wong, G.K.-S., and One Thousand Plant
Transcriptomes, I. 2019. One thousand plant transcriptomes and the phylogenomics of
green plants. Nature 574:679-685.
Leister, D. 2004. Tandem and segmental gene duplication and recombination in the evolution of
plant disease resistance gene. Trends Genet 20:116-122.
Li, B., and Dewey, C.N. 2011. RSEM: accurate transcript quantification from RNA-Seq data with
or without a reference genome. BMC Bioinformatics 12:323.
Linnakoski, R., and Forbes, K.M. 2019. Pathogens—The Hidden Face of Forest Invasions by
Wood-Boring Insect Pests. Frontiers in Plant Science 10.
Liu, J., Yu, T., Mu, Z., and Li, G. 2019. TransLiG: a de novo transcriptome assembler that uses line
graph iteration. Genome Biology 20:81.
Lu, M., Feau, N., Vidakovic, D.O., Ukrainetz, N., Wong, B., Aitken, S.N., Hamelin, R.C., and
Yeaman, S. 2021. Comparative Gene Expression Analysis Reveals Mechanism of Pinus
contorta Response to the Fungal Pathogen Dothistroma septosporum. Mol Plant
Microbe Interact:Mpmi10200282r.
Lynch, M., and Conery, J.S. 2000. The evolutionary fate and consequences of duplicate genes.
Science 290:1151-1155.
Lynch, M., and Conery, J.S. 2003. The evolutionary demography of duplicate genes. Journal of
Structural and Functional Genomics 3:35-44.
Maere, S., De Bodt, S., Raes, J., Casneuf, T., Van Montagu, M., Kuiper, M., and Van de Peer, Y.
2005. Modeling gene and genome duplications in eukaryotes. Proceedings of the
National Academy of Sciences 102:5454-5459.
Manni, M., Berkeley, M.R., Seppey, M., Simão, F.A., and Zdobnov, E.M. 2021. BUSCO Update:
Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic
Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. Molecular Biology
and Evolution 38:4647-4654.
Mendes, F.K., Vanderpool, D., Fulton, B., and Hahn, M.W. 2020. CAFE 5 models variation in
evolutionary rates among gene families. Bioinformatics 36:5516-5518.
Meng, X., and Zhang, S. 2013. MAPK cascades in plant disease resistance signaling. Annu Rev
Phytopathol 51:245-266.
Meyers, B.C., Kozik, A., Griego, A., Kuang, H., and Michelmore, R.W. 2003. Genome-Wide
Analysis of NBS-LRR–Encoding Genes in Arabidopsis[W]. The Plant Cell 15:809-834.
Miura, S., Amacher, M., Hofer, T., San-Miguel-Ayanz, J., Ernawati, and Thackway, R. 2015.
Protective functions and ecosystem services of global forests in the past quarter-
century. Forest Ecology and Management 352:35-46.
Neale, D.B., and Kremer, A. 2011. Forest tree genomics: growing resources and applications.
Nat Rev Genet 12:111-122.
Ngou, B.P.M., Heal, R., Wyler, M., Schmid, M.W., and Jones, J.D.G. 2022. Concerted expansion
and contraction of immune receptor gene repertoires in plant genomes. Nature Plants
8:1146-1152.
Ohno, S. 1970. Evolution by gene duplication (Springer-Verlag).
Panchy, N., Lehti-Shiu, M., and Shiu, S.H. 2016. Evolution of Gene Duplication in Plants. Plant
Physiol 171:2294-2316.

Piasecka, A., Jedrzejczak-Rey, N., and Bednarek, P. 2015. Secondary metabolites in plant innate
immunity: conserved function of divergent chemicals. New Phytologist 206:948-964.
Picart-Picolo, A., Grob, S., Picault, N., Franek, M., Llauro, C., Halter, T., Maier, T.R., Jobet, E.,
Descombin, J., Zhang, P., Paramasivan, V., Baum, T.J., Navarro, L., Dvořáčková, M.,
Mirouze, M., and Pontvianne, F. 2020. Large tandem duplications affect gene
expression, 3D organization, and plant-pathogen response. Genome Res 30:1583-1592.
Price, M.N., Dehal, P.S., and Arkin, A.P. 2009. FastTree: computing large minimum evolution
trees with profiles instead of a distance matrix. Mol Biol Evol 26:1641-1650.
R Core Team. 2021. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. https://www.R-project.org/.
Reid, N.M., Proestou, D.A., Clark, B.W., Warren, W.C., Colbourne, J.K., Shaw, J.R., Karchner, S.I.,
Hahn, M.E., Nacci, D., Oleksiak, M.F., Crawford, D.L., and Whitehead, A. 2016. The
genomic landscape of rapid repeated evolutionary adaptation to toxic pollution in wild
fish. Science 354:1305-1308.
Rizvi, A.R., Baig, S., Barrow, E., and Kumar, C. 2015. Synergies between climate mitigation and
adaptioon in forest landscape restoration. Gland, Switzerland: IUCN.
Sanderson, M.J. 2003. r8s: inferring absolute rates of molecular evolution and divergence times
in the absence of a molecular clock. Bioinformatics 19:301-302.
Seoighe, C., and Gehring, C. 2004. Genome duplication led to highly selective expansion of the
<em>Arabidopsis thaliana</em> proteome. Trends in Genetics 20:461-464.
Soltis, P.S., and Soltis, D.E. 2016. Ancient WGD events as drivers of key innovations in
angiosperms. Curr Opin Plant Biol 30:159-165.
Swenson, N.G., and Jones, F.A. 2017. Community transcriptomics, genomics and the problem of
species co-occurrence. Journal of Ecology 105:563-568.
Swenson, N.G., Iida, Y., Howe, R., Wolf, A., Umaña, M.N., Petprakob, K., Turner, B.L., and Ma, K.
2017. Tree co-occurrence and transcriptomic response to drought. Nature
Communications 8:1996.
Tang, H., Bowers, J.E., Wang, X., Ming, R., Alam, M., and Paterson, A.H. 2008. Synteny and
Collinearity in Plant Genomes. Science 320:486-488.
Trouern-Trend, A.J., Falk, T., Zaman, S., Caballero, M., Neale, D.B., Langley, C.H., Dandekar,
A.M., Stevens, K.A., and Wegrzyn, J.L. 2020. Comparative genomics of six Juglans species
reveals disease-associated gene family contractions. The Plant Journal 102:410-423.
Van de Peer, Y., Maere, S., and Meyer, A. 2009. The evolutionary significance of ancient
genome duplications. Nature Reviews Genetics 10:725-732.
Wickham, H. 2011. The Split-Apply-Combine strategy for data analysis. J. Stat. Softw. 40:1-29.
Wickham, H. 2016. ggplot2: Elegant graphics for data analysis. Springer-Verlag New York, NY,
US.
Yang, Z. 2007. PAML 4: Phylogenetic analysis by maximum likelihood. Mol. Biol. Evol. 24:1586-
1591.
Yang, Z., and Nielsen, R. 2000. Estimating synonymous and nonsynonymous substitution rates
under realistic evolutionary models. Mol Biol Evol 17:32-43.
Yu, Y. 2022. mixR: An R package for finite mixture modeling for both raw and binned data.
Journal of Open Source Software 7:4031.

Zeng, L., Zhang, N., Zhang, Q., Endress, P.K., Huang, J., and Ma, H. 2017. Resolution of deep
851            eudicot phylogeny and their temporal diversification using nuclear genes from
852            transcriptomic and genomic datasets. New Phytol 214:1338-1354.
853    Zhang, J. 2003. Evolution by gene duplication: an update. Trends in Ecology & Evolution 18:292-
854            298.
855    Zhao, J., Davis, L.C., and Verpoorte, R. 2005. Elicitor signal transduction leading to production of
856            plant secondary metabolites. Biotechnol Adv 23:283-333.
857    Zwaenepoel, A., and Van de Peer, Y. 2019. Inference of Ancient Whole-Genome Duplications
858            and the Evolution of Gene Duplication and Loss Rates. Molecular Biology and Evolution
859            36:1384-1404.
860