Comparative transcriptomics reveals divergence in pathogen response gene families amongst twenty forest tree species

[1*]Mengmeng Lu, [2]Min Cao, [2,3]Jie Yang, [1]Nathan G. Swenson

[1] Department of Biological Sciences, University of Notre Dame, 100 Galvin Life Sciences, Notre

Dame, IN, USA 46556

[2] CAS Key Laboratory of Tropical Forest Ecology, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Mengla, 666303, Yunnan, China.

[3] Center of Plant Ecology, Core Botanical Gardens, Chinese Academy of Sciences, Mengla, 666303, Yunnan, China.

[*]Correspondance: Email: mlu@nd.edu

Funding: This research was supported by the National Science Foundation (NSF DEB-2124466) and NSFC China-US Dimensions of Biodiversity Grant (DEB-32061123003).

Abstract

Forest trees provide critical ecosystem services for humanity that are under threat due to ongoing global change. Measuring and characterizing genetic diversity is key to understanding adaptive potential and developing strategies to mitigate negative consequences arising from climate change. In the area of forest genetic diversity, genetic divergence caused by large-scale changes at the chromosomal level have been largely understudied. In this study, we used the RNA-seq data of twenty co-occurring forest trees species from genera including *Acer*, *Alnus*, *Amelanchier*, *Betula*, *Cornus*, *Corylus*, *Dirca*, *Fraxinus*, *Ostrya*, *Populus*, *Prunus*, *Quercus*, *Ribes*, *Tilia*, and *Ulmus* sampled from Upper Peninsula of Michigan. These data were used to infer the origin and maintenance of gene family variation, species divergence time, as well as gene family expansion and contraction. We identified a signal of common whole genome duplication events shared by core eudicots, and a signal of recent duplication events specific to particular species. We also found rapid evolution, namely fast expansion or fast contraction of gene families, in plant-pathogen interaction genes amongst the diploid species studied. Finally, the results lay the foundation for further research on the genetic diversity and adaptive capacity of forest trees, which will inform forest management and conservation policies.

44 **Introduction**
45
Forest trees occupy nearly a third of the Earth's land surface and provide key ecological benefits
and services such as water, air purification and carbon sequestration (Gamfeldt et al., 2013;
Miura et al., 2015; FAO, 2020). They are critical components of terrestrial biodiversity in terms
of their own diversity, but also with respect to the habitats and renewable materials such as
wood, cellulose, and lignin they supply other organisms (Cortés et al., 2020). It is estimated that
there are ~73,000 tree species globally with most of being rare and regionally endemic, implying
vulnerability of global tree species to anthropogenic and climate changes (Cazzolla Gatti et al.,
2022). Indeed, climate change has already impacted forests and their species composition via
range migration, wildfires, droughts, pest, and pathogens (Kirilenko and Sedjo, 2007; Allen et
al., 2010; Rizvi et al., 2015; Hammond et al., 2022). The genetic composition and diversity of
the tree species in forests is foundational to our understanding of the abiotic and biotic factors
governing the distribution and dynamics of species populations, but also how they will respond
to current and future environmental change. Thus, measuring and characterizing intra- and inter-
specific genetic diversity is the key for understanding the current structure and dynamics of
forests and their adaptive potential and for developing strategies to mitigate the damages brought
along with environmental change.
62
Gene duplication and gene loss are major sources of genetic diversity. Gene duplication,
generated through either whole genome duplication (WGD) or small-scale duplication (SSD), is
the primary source of new genetic material among flowering plants (Cui et al., 2006; Panchy et
al., 2016). WGD contributes to the majority of duplicate genes (AMBORELLA GENOME
PROJECT et al., 2013; Soltis and Soltis, 2016), where ancient WGDs can lead to competitive
advantage, increased species diversity, and evolutionary innovations (Van de Peer et al., 2009).
Most flowering species have a polyploid history either through autopolyploidy or allopolyploidy.
Specifically, flowering species are either currently polyploid, or have a polyploid ancestor and
have reverted to being diploid (Blanc and Wolfe, 2004). Alternatively, pervasive SSD, including
tandem duplication (Zhang, 2003), transposon-mediated duplication (Jiang et al., 2004),
segmental duplication (Bailey et al., 2002b), and retroduplication (Brosius, 1991), also generates
a great number of duplicate genes. While the majority of duplicate genes end up being silenced
and eliminated (Lynch and Conery, 2000), a few preserved duplicate genes may be retained by
selection due to neo-functionalization, relative dosage, absolute dosage or sub- functionalization
(Conant et al., 2014), and play important role in generating genetic diversity and building up
reproductive isolation barriers.
79
Compared to gene duplication, the mechanism that underlie gene loss are not yet well
understood. It is commonly assumed that gene loss is realized by deletion or pseudogenization,
as a result of neutral or adaptive processes (Demuth and Hahn, 2009). Most nonsense mutations
are subject to drift to fixation. Some deleterious mutations could be accumulated and eventually
become silenced (Johri et al., 2022). Gene loss might be selected to retain dosage balance, as
evidenced in a set of single-copy genes with essential housekeeping functions, which are
typically restored to become singletons after WGD or SSD (De Smet et al., 2013).
87
Differential duplication and loss of genes among evolutionary lineages results in gene family
expansion and contraction (Demuth and Hahn, 2009). Gene families comprise a set of similar

nucleotide or amino acid sequences. They arise as a result of duplication events and have similar
cellular and biochemical functions. Gene family expansions and contractions related to
metabolic, regulatory and signalling gene networks contribute to the adaptations of plants to
various abiotic habitats and biotic interactions (Leebens-Mack et al., 2019). They are sources of
plant functional evolution and innovation (Ohno, 1970; Force et al., 1999), exhibiting a clear
functional basis, which are influenced by the mechanisms of duplication (Maere et al., 2005;
Hanada et al., 2008). Comparisons of gene family content across species, therefore, may provide
insight into the ecological interactions and evolutionary pressures that have shaped adaptation
and diversity. This is particularly true when we conduct comparative transcriptomic analyses of
species that currently co-occur and that, potentially have co-occurred for long periods of time
(Swenson and Jones, 2017).

Genetic investigations of forest trees have, generally, focused on nucleotide substitutions in
specific genes (Neale and Kremer, 2011; Isabel et al., 2020), while comparative analyses of
genetic divergence caused by large-scale changes at the chromosomal levels have been largely
understudied. However, the magnitude, rate, and distribution of changes in gene family size are
key drivers of the functional differentiation and adaptation of species (Lynch and Conery, 2003;
Demuth and Hahn, 2009). Large segmental duplication and changes in copy number are known
to be associated with disease response and environmental adaptation (Bailey et al., 2002a; Reid
et al., 2016). Therefore, gene family size change is as much important as nucleotide substitution
to understanding genetic divergence and evolutionary forces responsible for these changes.

Such knowledge on genetic divergence and associated evolutionary forces in forest trees can
help to promote breeding healthy trees (i.e. disease resistant trees) in a changing climate (Allen
et al., 2010; Dodds and Rathjen, 2010). The increasing occurrences and severity of insect and
disease outbreaks have threatened forest trees (Hurley et al., 2016; Linnakoski and Forbes,
2019), but we lack a thorough understanding to the disease related genes in forest trees. Trouern-
Trend et al. (2020) revealed contraction in disease associated gene families in *Juglans* species.
Lu et al. (2021) reported differential gene expression patterns between tolerant and susceptible
lodgepole pine trees to a fungal pathogen. It would be interesting to explore the potential link
between duplication events and disease response in expressed genes of forest trees, as the
duplicate genes can quickly lead to changes in gene expression with the acquisition of increased
resistance to different pathogens (Picart-Picolo et al., 2020).

In this study, we used the RNA-seq data for twenty co-occurring forest trees species to infer the
origin and maintenance of gene family variation as well as species divergence time, gene
expansion and contraction. We sought to address the following questions: What is the magnitude
of gene duplication? What factors impact duplicate gene evolution? Are the gene gain and loss
process random or controlled by natural selection? The answers to these questions are the key to
understanding the basis of adaptive divergence in forest tree species and will help to predict
adaptive capacity of these species in a changing climate.

Material and methods
Samples, sequencing data, de novo assembly, and annotation

Leaves of each of twenty forest species were sampled in August, 2021 at the University of Notre
Dame Environmental Research Center (UNDERC) near Land O' Lakes, Wisconsin, U.S.A.
(Figure 1). The UNDERC site is located at latitude: 46.23391, longitude: -89.537254, and
encompasses approximately ~8,000 acres of mixed deciduous forest on both sides of the state
line between Wisconsin and Michigan's upper peninsula. The twenty species used in this study
are dominate components of the woody vegetation at the site and are: *Acer rubrum* (red maple),
Acer saccharum (sugar maple), *Acer spicatum* (mountain maple), *Alnus incana* (Gary alder),
Amelanchier laevis (smooth serviceberry), *Betula alleghaniensis* (yellow birch), *Betula*
papyrifera (paper birch), *Cornus alternifolia* (pagoda dogwood), *Corylus cornuta* (beaked
hazel), *Dirca palustris* (Eastern leatherwood), *Fraxinus nigra* (black ash), *Ostrya virgniana*
(American hophornbeam), *Populus grandientata* (bigtooth aspen), *Populus tremuloides* (quaking
aspen), *Prunus serotina* (black cherry), *Prunus virginiana* (chokecherry), *Quercus rubra*
(Northern red oak), *Ribes cynosbati* (prickly gooseberry), *Tilia americana* (basswood), *Ulmus*
americana (American elm). They are all angiosperms (flowering plants) and eudicots, belong to
the clades of asterids, saxifragles, and rosids. While these species all co-occur at the UNDERC
site, their geographic ranges largely overlap on the continental scale and the species have largely
co-occurred in North America since at least the Pleistocene. Upon collection, the samples were
flash frozen in liquid nitrogen, then transported and stored in an ultracold freezer at the field
station. The samples were sent to Novogene corporation inc. (Chula Vista, CA, USA) for mRNA
extraction, library preparation, mRNA sequencing, and quality control. The samples were
sequenced on an Illumina NovoSeq 6000 platform to produce 150bp paired-end reads.
157

158
159 **Figure 1. The location (red dot) of the University of Notre Dame Environmental Research**
160 **Center (UNDERC) where all samples used in this study were collected.**
161
Raw reads filtering was performed by Novogene to get the clean reads including removing reads
containing adapters, removing reads containing N >10%, and removing reads containing low
quality base which is over 50% of the total base. An average of 6Gb paired-end clean reads were
acquired for each sample and deposited in NCBI Sequence Read Archive (SRA accession
number: PRJNA896247). *De novo* transcriptome assembly was performed for each sample using
the software TransLiG v1.3 with default parameters (Liu et al., 2019). The clean reads were then
mapped to the assembled transcriptome using the software RSEM v1.2.25 with default
parameters (Li and Dewey, 2011). The longest transcript (isoform) with a transcript per million

(TPM) value ≥ 1.5 for each gene was retained. Redundancy was removed from the retained
transcripts using the software CD-HIT v4.8.1 with parameters of -c 0.95 -n 8 -T 32 -M 1540 (Fu
et al., 2012). To remove potential contamination, the retained transcripts were aligned against the
non-redundant (nr) protein database (downloaded on January 22, 2022) using the blastx function
implemented by the software DIAMOND v2.0.13.151 (Buchfink et al., 2021) with parameter of -
-outfmt 102 to assign the best known taxon ID to each query transcript. The file that was used to
convert accession numbers to taxonomy was downloaded from
https://ftp.ncbi.nlm.nih.gov/pub/taxonomy/accession2taxid/, and the tax ID file was downloaded
from https://ftp.ncbi.nlm.nih.gov/pub/taxonomy/new_taxdump/. The query transcripts that could
be assigned to green plant (Viridiplantae) taxon IDs were retained. The filtered transcripts were
aligned to the nr database that only includes green plant subjects using the blastx function
implemented by the software DIAMOND v2.0.13.151 with default parameters. The aligned
transcripts were also retained and then were combined with the previously retained transcripts to
form the final transcriptomes. The completeness of the final transcriptomes was examined using
the 1,614 orthologues in the BUSCO (Benchmarking Universal Single Copy Orthologs) v5.2.2
set of embryophyta_odb10 (Manni et al., 2021).
186
The transcriptomes were annotated with Gene Ontology (GO) and Kyoto Encyclopedia of Genes
and Genomes (KEGG) terms. The transcripts were aligned to the nr database that only includes
green plant subjects using the blastx function implemented by the software DIAMOND
v2.0.13.151 with parameters of -taxonlist 33090 --very-sensitive -f5. The .xml outputs were
used as input for mapping and annotation using the software Blast2GO (Götz et al., 2008). The
transcripts were translated to protein sequences using the software TransDecoder v5.5.0
(https://github.com/TransDecoder/TransDecoder/wiki) with default parameters to get the longest
ORF (open reading frame) per transcript, and then were annotated by the software InterProScan
v5.54-87.0 (Jones et al., 2014) with the parameters of -dra -f XML -goterms -pa. The final GO
terms were acquired by merging Blast2GO and InterProScan annotation outputs using the
software Blast2GO. To get the KEGG terms, the longest ORF per transcript were also annotated
using EggNOG-mapper v2.1.7 (Cantalapiedra et al., 2021) with parameters of -m diamond -
sensmode more-sensitive -tax_scope 33090 -tax_scope_mode 33090.
200
Ks analyses
The level of synonymous substitutions (Ks) between two homologous sequences increases
approximately linearly with time. Thus pairs of paralogous genes within a species can be sorted
to estimate the relative ages of gene duplication (Blanc and Wolfe, 2004). To identify paralogous
gene pairs within a species, an all-versus-all blastp was performed within each species. The
peptide sequences (the longest ORF per transcript) of each species were used as input to run all-
versus-all blastp using the software DIAMOND v2.0.13.151 with parameters of -p 16 --max-
target-seqs 20 --evalue 1e-10 --outfmt 6 qseqid sseqid pident nident evalue. The blastp output
was filtered by removing self blastp results, keeping outputs with pident (percent of identical
matches) ≥20 and nident (number of identical matches) ≥50, and removing sequences with more
than 10 hits. The filtered paralogous gene pairs were aligned using the software MAFFT v7.490
(Katoh et al., 2002) with default parameters. The alignments were used as input for estimating
the number of synonymous substitutions per synonymous site (Ks) and the number of non-
synonymous substitutions per nonsynonymous site (Ka) using the program yn00 in the software

PAML v4.9j (Yang, 2007) with parameters of icode=0 weighting=0 commonf3×4=0. The Ks of
each paralogous gene pairs were acquired using the method by Yang and Nielsen (2000).
217
Ks can also be calculated using the orthologous gene pairs between species to estimate speciation
events (Blanc and Wolfe, 2004). To identify orthologous gene pairs between species, reciprocal
best hit blastp were run between the compared pairs. The peptide sequences (the longest ORF per
transcript) of the paired species were used as input to run blastp using the software DIAMOND
v2.0.13.151 with parameters of -p 16 --max-target-seqs 20 --evalue 1e-10 --outfmt 6 qseqid
sseqid pident nident evalue. The best hits between forward and reverse blastp for each compared
sequence pairs were retained, which were then used for sequence alignment and estimating Ka
and Ks as abovementioned.
226
The Ks distribution is proportional to the time lapse. Thus, the Ks distribution within species can
be used to infer the timing of WGD, and the Ks distribution between two different species can
infer time since speciation (Blanc and Wolfe, 2004). To plot the Ks distribution, the mixture
model was fit and plot for the Ks values ≥0.01 and ≤2 using the R package mixR (Yu, 2022). In
case there are discernably separate peaks (multiple components), bootstrap likelihood ratio test
(LRT) was used to select the models.
233
Estimating duplication events and divergence time
To infer duplication events, the species tree and gene trees were first inferred using STAG and
STRIDE algorithms implemented by the program OrthoFinder v2.5.4 (Emms and Kelly, 2017,
2018). The inferred species tree was similar to that expected given current knowledge of the
Angiosperm phylogeny with the exception of the placement of *Populus tremuloides* and *P.*
grandidentata. Thus, for downstream analysis and discussion, we removed these two species for
the species tree inference. The inferred species tree and gene trees were used to estimate
duplication events and divergence time. The gene duplication events on the species tree node
were reported if, at least 50% of the descendant species have retained both copies of the
duplicated gene. The divergence time was estimated using the software r8s v1.81 (Sanderson,
2003) with calibration points of 115 million years (MYA) between *Fraxinus nigra* and *Corylus*
cornuta, which were obtained from http://www.timetree.org/ (Kumar et al., 2017).
246
Gene family and their evolutionary rate
To infer gene families, the peptide sequences (the longest ORF per transcript) of the 20 study
transcriptomes were used to construct orthogroups (gene families) using the program
OrthoFinder v2.5.4 with parameters of -S diamond -t 16 -a 16. Genes assigned to each gene
family were counted for each species. The evolutionary rate of protein coding genes was
represented by the ratio of nonsynonymous substitutions per nonsynonymous site to synonymous
substitutions per synonymous site (Ka/Ks) with *Dillenia indica* (elephant apple) being the
outgroup and the 20 study species being the ingroup. *D. indica* is an eudicot species, belonging
to the order Dilleniales and the RNA-Seq data for this species were downloaded from
https://trace.ncbi.nlm.nih.gov/Traces/index.html?view=run_browser&acc=ERR2040190&disp
lay=data-access (Leebens-Mack et al., 2019). The raw sequences were filtered using the program
fastp with default parameters (Chen et al., 2018). The transcriptome assembly and protein
sequence prediction were performed as abovementioned, yielding 22,753 transcripts and 19,345
protein sequences. Then the reciprocal best hit blastp was performed between *D.indica* and each

of the twenty study species to identify the orthologous pairs. The best hits between forward and
reverse blastp for each compared sequence pairs were retained, which were then used for
sequence alignment and estimating Ka/Ks as abovementioned. The mean Ka/Ks across genes for
each gene family per species was used to represent the evolutionary rate for each gene family.
Gene families with mean Ka/Ks >10 were filtered out. The graphs were plotted using the R
package ggplot2 and ggpubr (Wickham, 2016; Kassambara, 2020). We studied the relationship
between gene family size and evolutionary rate, which is represented by mean Ka/Ks per gene
family, using scatter plot. All statistical analyses were performed in R (R Core Team, 2021).
Gene families with a family size ≥5, and Ka/Ks ≥1.5 were highlighted for further analysis. The
enriched GO and KEGG terms were identified in these highlighted gene families. The enriched
GO terms of each category (biological process, molecular function, cellular component) were
separately detected using the "weight" algorithm with the Fisher ratio test implemented by the R
package "topGO" (Alexa and Rahnenfuhrer, 2018). Enriched KEGG terms were detected using
the hypergeometric method implemented by the R package "plyr" (Wickham, 2011). Heatmaps
were plot using the R package pheatmap (Kolde, 2019).
276
Estimating gene family expansion and contraction
To further study gene family changes, species tree and gene family counts were predicted and
used to estimate gene family expansion and contraction. All-versus-all blastp was performed to
find the most similar sequences for each sequence in the study species using the program
DIAMOND (Buchfink et al., 2021). The program MCL (Enright et al., 2002) was used to find
clusters of similar sequences (gene families). The species tree was estimated using the program
FastTree (Price et al., 2009). The programs DIAMOND, MCL, and FastTree were implemented
by the program OrthoFinder v2.5.4, default parameters were used. Then, the species tree was
converted to a format for estimating ultrametric tree using the program CAFE5 (Mendes et al.,
2020) with the command python prep_r8s.py -s 60164 -p 'Fraxinus_nigra,Corylus_cornuta' -c
120. "60164" was the number of sites in the alignment. "-c 120" represents a median of 120
MYA divergence time, it is the calibration points between *Fraxinus nigra* and *Corylus cornuta*
checked from http://www.timetree.org/. Then the ultrametric tree was estimated using the
program r8s (Sanderson, 2003) with default parameters. The ultrametric tree and gene family
counts were used as input for the program CAFE5 (Mendes et al., 2020). Since CAFE5 was not
recommended to analyze recent polyploid species, only diploid study species were retained for
running CAFE5, including *Cornus alternifolia, Fraxinus nigra, Dirca palustris, Tilia americana,*
Acer saccharum, Acer spicatum, Amelanchier laevis, Quercus rubra, Betula papyrifera, Alnus
incana, Corylus cornuta, Ostrya virginiana, and *Ribes cynosbati.* Since assembly errors might
cause the observed number of gene copies in gene families to deviate from the true ones and
overestimation of the birth-death parameter λ (number of gene gains and losses per gene per
million years), an error model was estimated first with the ultrametric tree and gene family
counts using CAFE5 with the parameters of "-c 16 -p -e". This error model along with the
ultrametric tree and gene family counts were used to estimate the birth-death parameter λ, which
represents the probability that any gene will be gained or lost. Multiple λ mode was used with
three different rates of gene family evolution assigned as below:
303
304 ((Cornus_alternifolia:3,Fraxinus_nigra:3):3,((((Dirca_palustris:2,Tilia_americana:2):2,(Acer_sac
305 charum:2,Acer_spicatum:2):2):2,((((Betula_papyrifera:2,Alnus_incana:2):2,(Corylus_cornuta:2,
306 Ostrya_virginiana:2):2):2,Quercus_rubra:2):2,Amelanchier_laevis:2):2):2,Ribes_cynosbati:1):1);

307
The numbers 1, 2, 3 represent three different gene family evolution rates that were shared by
different clades. This tree file was used as input along with the ultrametric tree, gene family
counts, and error model using CAFE5 with the parameters of -c 16 -p -k 3. Gene families or
parent nodes with a significantly greater rate of evolution (family-wide p-value <0.01), in other
words, rapidly evolving gene families, were counted for each species and each parent node
(Figure 5b). These gene families significantly violated a random gene birth and death model.
Thus, the null hypothesis of neutral evolution was rejected and these families could be
recognized as fast evolving gene families (Hahn et al., 2005). Enriched GO and KEGG terms
were detected in the fast expanding and fast contracting gene families, respectively.
317
318
319   **Results**
320   We acquired an average of 41 million raw reads for each sample. The total number of assembled
321   transcripts ranged between 19,459 and 31,804 (Table S1). We retrieved 76 ~ 88 % of the
322   BUSCOs in the assembled transcriptomes (Figure S1).
323
We found a large Ks peak around 0.71~0.75 shared by all the study species (Figure S2), and
discernable separate peaks on the left of the major peak can be observed from Ks plots of
Amelanchier laevis, *Dirca palustris*, *Fraxinus nigra*, *Populus grandidentata*, *Prunus serotina*,
Populus tremuloides, *Prunus virginiana*, and *Tilia americana* (Figure 2). Bootstrap LRT
indicates that considering these small peaks as a separate component could fit the mixture model
better. The shared large peaks by all species may correspond to large-scale duplication events,
perhaps the common WGDs across core eudicots, and the small peaks may represent recent
duplication events specific to certain species. The two *Prunus* species have the small peaks with
similar shapes at a similar position (Ks=0.10 or 0.11) and so do the two *Populus* species
(Ks=0.12). This pattern may represent a common WGD before the two species diverged from the
same ancestors. By examining the orthologous Ks between the two species, we found a high
peak closer to zero, which may represent the duplication events after the WGDs and divergence.
336

Figure 2. The Ks distributions for the studied species with discernable separate peaks on the left of the major peak.

Using a species tree with estimated branch lengths, we inferred divergence times and duplication events (with at least 50 % support rate, Figure 3). Most duplication events (n=1843) occurred at the divergence of rosids and asterids, which may correspond to common WGD across eudicots species. The ancestor of *Populus tremuloides* and *Populus grandidentata* had 1,625 duplication events (Figure S3), which is consistent with the recent Ks peaks within these two species. The divergence time of the two asterids species *Cornus alternifolia* and *Fraxinus nigra* from a common ancestor was about 104 MYA (Figure 3). The divergence time of the studied malvids and fabids from a common ancestor with Saxifragles was about 100 MYA.

Figure 3. Divergence time in million years and gene duplication events with 50 % support
rate. Node numbers were followed by number of gene duplication events.

OrthoFinder assigned 393,796 genes (96.2 % of total) to 21,900 orthogroups (gene families). The
mean and median gene family size is 17.3 and 12, respectively. We found that 3,060 families
present in all species, whereas 19 consisted entirely of single copy-genes. Most gene families
have a size <5, only 1.8~5.9 % of the total gene families across all species have a size ≥5
(Figure 4a). Gene families with large sizes tended to have low Ka/Ks, yet there are a few
exceptions. We highlighted the gene families with a size ≥5 and Ka/Ks ≥1.5 in red (Figure 4a).
We tested the KEGG and GO terms that are enriched in these highlighted gene families (Figures
4b &4c & S4, see the whole list of GO enrichment in Table S2). The most highly shared
enriched KEGG and GO terms are related to plant-pathogen interactions and secondary
metabolite biosynthesis. Across all species, percentage of gene families with a size ≥5 and
Ka/Ks ≥1.5 is 4.4 % among plant-pathogen interaction gene families, and is 0.7 % among all
gene families.

Figure 4. Scatter plot between gene size and Ka/Ks per gene family (a), KEGG pathway (b) and GO terms enrichment (c) in gene families with a size ≥5 and Ka/Ks ≥1.5.

We inferred the birth-death parameter, λ (number of gene gains and losses per gene per million years), which is specified in different clades, using the program CAFE. $\lambda1$ (saxifragales) =0.0034, $\lambda2$ (rosids) =0.0042, $\lambda3$ (asterids) =0.0027, which means that the lineage leading to rosids have the highest gene family turnover rates, followed by saxifragales, and then by asterids. The four species with recent duplication events (Figure 2) were *Fraxinus nigra*, *Dirca palustris*, *Tilia americana*, and *Amelanchier laevis*. Specifically, these species have more expanding than contracting gene families (Figure 5a, S5). All of the study species have more rapidly expanding than rapidly contracting gene families (family-wide p-value <0.01, Figure 5b), which may contribute to their lineage-specific novelty.

We tested the enriched GO and KEGG terms in the fast evolving (expanding or contracting) gene families. All of the study species shared the enriched KEGG pathway term plant-pathogen interaction (Figure 6). This term was detected in the fast-expanding gene families of *Cornus alternifolia*, *Tilia americana*, *Acer spicatum*, *Betula papyrifera*, *Quercus rubra*, *Amelanchier laevis*, *Ribes cynosbati*; and this term was detected in the fast-contracting gene families of *Fraxinus nigra*, *Dirca palustris*, *Alnus incana*, *Corylus cornuta*, *Ostrya virginiana*; also, this term was detected in both of the fast expanding and fast contracting gene families of *Acer saccharum*. The gene families assigned to the plant-pathogen interaction term encode proteins (Figure S6) related to disease resistance protein, leucine-rich repeat receptor like protein kinase and others. Some species shared the enriched KEGG pathway term Tryptophan metabolism and MAPK signaling pathway (Figure 6).

Figure 5. The percentage of gene family expansion and contraction (a) and the number of fast evolving gene families (b), with + representing fast expanding gene family and – representing fast contracting gene family.

Figure 6. Enriched KEGG pathway terms in the study diploid species. Red represents fast expanding gene families; blue represents fast contracting gene families; white represents both fast expanding and contracting gene families.

Discussion

Gene duplication events are prevalent in all organisms and provide genetic variation for evolutionary innovation and adaption to fluctuating conditions (Zhang, 2003; Van de Peer et al., 2009; Panchy et al., 2016). Research on the mechanisms of gene duplication as well as the

origin, maintenance, and functional roles of genetic diversity in forest tree species are few due to
the limited genome and transcriptome data resources. In the present study, we assembled twenty
transcriptomes from forest trees that co-occur. We identified the signal of common WGDs
shared by core eudicots, and the signal of recent duplication events specific to certain species.
We found gene families involved in plant-pathogen interaction and metabolite biosynthesis
actively remain extensive variation, and fast expansion or fast contraction in plant-pathogen
interaction genes amongst the study diploid species. In the following sections, we will discuss
these results in detail.
420
Tempo and mode of gene duplication
The Ks distribution plots confirmed the widespread paleopolyploidy in plants, that's said,
today's diploid species might be polyploid (paleopolyploid) at some points in evolutionary
history (Blanc and Wolfe, 2004). All the study species shared a duplication event as seen by the
large Ks peaks around 0.71~0.75 (Figure S2), which probably correspond to the common WGDs
across core eudicots. Besides the large peak, some species had a discernable recent duplication
event as seen by the small peaks on the left of the major peak (Figure 2). These abundant
duplication events provide resources for genetic variation, although most duplicate genes were
lost by the process of deletion of the entire duplicate sequence and/or pseudogenization through
loss-of-function mutations, the retained duplicate genes lead to lineage-specific innovation
(Panchy et al., 2016). Earlier studies placed at least three WGD events in angiosperm radiation,
the oldest WGD occurred before the monocot-eudicot divergence, a second WGD occurred
specifically to eudicots, a more recent WGD, gamma (γ) triplication event, occurred before the
separation of rosids and asterids (Tang et al., 2008; Jiao et al., 2012). Besides, the lineage-
specific WGDs and numerous SSDs continuously lead to new and complex gene contents. It is
worth noting that Ks distribution is not suitable to infer very ancient events due to saturation with
multiple substitutions (Blanc and Wolfe, 2004), moreover, different peaks might not be separable
from each other, thus it is possible that we missed identifying some duplication events in the
present study. Collinearity- or synteny-based methods along with chromosome-level reference
genome assembly is alternative method to identify ancient WGD, yet high quality genome
assembly is not easy to obtain (Zwaenepoel and Van de Peer, 2019).
442
The predicted species tree (Figure 3) generally agrees with current phylogenetic hypotheses for
the angiosperms with the exception of the two *Populus* species, which is probably caused by the
incomplete gene sets obtained from one leaf RNA-seq sample and/or interspecific gene flow. We
identified the most (1,843) duplication events at the root (Figure 3) of the tree, which is likely to
be related to the gamma WGD before the divergence of rosids and asterids. Such large-scale
duplication may contribute to genome structure variation, organismal complexity, and rapid
speciation in angiosperm species. The parental node for *Acer*, *Prunus*, *Populus* species (Figure 3
& S3) also have a large number of duplication events. The most-likely scenario is WGD event
occurred in the common ancestor, followed by different evolutionary events. For *Acer*, *Acer*
spicatum experienced a diploidization event to revert to diploid status, while *Acer rubrum*
maintained a polyploid status. Similarly, the recent Ks peaks on the similar location in two
Prunus species and two *Populus* species are in line with their large number of duplication events
in parental nodes.
456

Divergence time mostly agrees with the previous reports, which stated rosids and asterids
diverged about 125 MYA, malvids and fabids diverged about 110 MYA (Zeng et al., 2017).
However, divergence estimate in the present study is rather rough. Further studies need to
consider using dense taxon sampling, diverse markers like nuclear internal transcribed spacer
(ITS) or chloroplast DNA fragments, and molecular dating with fossil corrections methods to
better resolve the evolutionary relationships.
463
Sustained large size gene families
A small portion of gene families can actively maintain duplicate genes. Gene family evolution is
dynamic process resulting from gene or genome duplication events where most of these
duplicate genes are eventually eliminated or pseudonized (Panchy et al., 2016). The extreme
example is diploidization of paleopolyploidy species, which revert to one copy of genes after
WGDs. Nonetheless, some duplicate genes can be retained, perhaps due to the gene balance and
dosage effect (Birchler and Veitia, 2012). Alternatively, selection favors the neofunctionalization
in the duplicate genes, as these duplicate genes help organisms adapt to variable environments
(Cortesi et al., 2015). Recent studies explained retention of duplicate genes as some paralogous
genes depend on or compensate for their cognate copy (Diss et al., 2017), while some paralogous
gene pairs are stuck in a functional and structural entanglement status over evolutionary time
scale (Kuzmin et al., 2020).
476
What are the main functions of these sustained large size gene families? They are mainly
involved in plant-pathogen interaction and metabolite biosynthesis (Figure 4b & 4c). This is
consistent with previous report that genes encoding functions like signal transduction and stress
response tend to have paralogs (Panchy et al., 2016), and genes related to the function like
transcription regulation and signal transduction had higher possibility to survive in multiple
rounds of genome duplication (Seoighe and Gehring, 2004). Plants constantly combat with
pathogens and pests. The major class of plant disease resistance (R) genes contain a nucleotide-
binding site (NBS) and C-terminal leucin-rich repeats (LRRs). Tandem and segmental gene
duplication explain much of the diversification among R genes (Meyers et al., 2003; Leister,
2004; Andersen et al., 2020), possibly because they have experienced frequent deletion and
duplication throughout evolution, and thus are highly variable (Hanada et al., 2008; Gao et al.,
2018). Plant secondary metabolites, such as monoterpenoid, flavonoid, benzoxazinoid, play
important roles in defense against predation and pathogens (Falcone Ferreyra et al., 2012; de
Bruijn et al., 2018; Bergman et al., 2020). The regulation of plant secondary metabolism requires
the intimation of signal transduction network that leads to activation of biosynthetic genes (Zhao
et al., 2005).
493
Gene duplication and deletion give rise to gene family expansion and contraction. The greater
number in gene family expansions than contractions (Figure 5a) in *Fraxinus nigra*, *Dirca*
palustris, *Tilia americana*, and *Amelanchier laevis* is associated with their recent duplication
events as shown in the Ks distribution (Figure 2). A previous study showed that estimates of
birth and death rate was consistent with the estimates based on number of recent gene duplicates
(Demuth and Hahn, 2009), thus it is reasonable to see the expansion numbers parallel the Ks
distributions. The high gene gain rate likely reflects the occurrence of multiple WGDs and
frequent SSDs (Carretero-Paulet et al., 2015). While for the other species with contraction
outnumbered expansion, the pattern is mainly caused by gene loss and retention of specific gene

families followed the common WGDs preceded their common ancestor (Figure S2). We found a
high gene family turnover rate in rosids, but we must caution that the unbalanced taxon sampling
in our study species might cause this high turnover value. Moreover, future studies should
consider distinguishing the branches with or without recent WGDs, as the branches with recent
WGDs tend to have a high gene turnover rate (Casola and Koralewski, 2018).
508
Rapid evolution in plant-pathogen interaction genes
We found rapid evolution of genes involved in plant-pathogen interaction in the adaptive
expansion and adaptive contraction gene families across the study species (Figure 6). The
specific expanding or contracting gene families involved in plant-pathogen interaction in each
species differed from each other, implying lineage-specificity in changed gene families (Figure
S6). Other rapidly evolving gene families are related with functions such as tryptophan
metabolism and MAPK signaling pathway (Figure 6). This result is consistent with previous
studies, which found that fast evolving gene families were mostly groups with functions related
with immune defense/stress response, metabolism, cell signaling, chemoreception, and
reproduction (Demuth and Hahn, 2009). Additionally, a study on *Juglans* (walnuts) found
notable contractions in gene families that are involved in disease resistance and other abiotic
stress response (Trouern-Trend et al., 2020). Both metabolism and MAPK signaling pathway are
associated with plant defense. MAPK cascades sense pathogen/microbe-associated molecular
patters and pathogen effectors, and thus play pivotal roles in signaling plant defense against
pathogen attack (Meng and Zhang, 2013). Plant secondary metabolites contribute to plant
immunity as antibiotics, or to control callose deposition and programmed cell death (Piasecka et
al., 2015). The biotic stress exerted by pathogen or pests is a major source of selection pressure
on the study trees, which drive the rapid radiation of diverse and large plant immune receptor
gene repertoires. Expansion and/or contraction of cell-surface receptor genes correlate with the
expansion and/or contraction of intracellular receptor genes (Ngou et al., 2022).
529
Conclusions
In summary, we found the common WGDs shared by all the study species with some species
having recent duplication events, which contribute to the gene family expansions. Gene families
involved in plant-pathogen interaction and metabolite biosynthesis actively remain extensive
variation and are fast expanding or contracting amongst the study species. This knowledge calls
for an emphasis of gene family size variation, i.e. copy number variation, when studying forest
genetic diversity, which will inform forest adaptive strategy. Future studies may focus on
classification of duplications in forest tree species, and investigate how the duplications give rise
to species-specific innovation and adaption.
539
540
541
542 **Data availability**
543 The raw RNA-seq data were deposited in NCBI SRA (accession number: PRJNA896247;
544 http://www.ncbi.nlm.nih.gov/sra). The assembled transcriptomes and their annotation were
545 deposited in Dryad. The scripts used for analyses were deposited in
546 https://github.com/Mengmeng-Lu.
547
548

References

Alexa, A., and Rahnenfuhrer, J. 2018. topGO: Enrichment Analysis for Gene Ontology. R
package version 2.34.0.
Allen, C.D., Macalady, A.K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M.,
Kitzberger, T., Rigling, A., Breshears, D.D., Hogg, E.H.T., Gonzalez, P., Fensham, R.,
Zhang, Z., Castro, J., Demidova, N., Lim, J.-H., Allard, G., Running, S.W., Semerci, A.,
and Cobb, N. 2010. A global overview of drought and heat-induced tree mortality reveals
emerging climate change risks for forests. For Ecol Manage 259:660-684.
AMBORELLA GENOME PROJECT, Albert, V.A., Barbazuk, W.B., dePamphilis, C.W., Der,
J.P., Leebens-Mack, J., Ma, H., Palmer, J.D., Rounsley, S., Sankoff, D., Schuster, S.C.,
Soltis, D.E., Soltis, P.S., Wessler, S.R., Wing, R.A., Albert, V.A., Ammiraju, J.S.S.,
Barbazuk, W.B., Chamala, S., Chanderbali, A.S., dePamphilis, C.W., Der, J.P.,
Determann, R., Leebens-Mack, J., Ma, H., Ralph, P., Rounsley, S., Schuster, S.C., Soltis,
D.E., Soltis, P.S., Talag, J., Tomsho, L., Walts, B., Wanke, S., Wing, R.A., Albert, V.A.,
Barbazuk, W.B., Chamala, S., Chanderbali, A.S., Chang, T.-H., Determann, R., Lan, T.,
Soltis, D.E., Soltis, P.S., Arikit, S., Axtell, M.J., Ayyampalayam, S., Barbazuk, W.B.,
Burnette, J.M., Chamala, S., De Paoli, E., dePamphilis, C.W., Der, J.P., Estill, J.C.,
Farrell, N.P., Harkess, A., Jiao, Y., Leebens-Mack, J., Liu, K., Mei, W., Meyers, B.C.,
Shahid, S., Wafula, E., Walts, B., Wessler, S.R., Zhai, J., Zhang, X., Albert, V.A.,
Carretero-Paulet, L., dePamphilis, C.W., Der, J.P., Jiao, Y., Leebens-Mack, J., Lyons, E.,
Sankoff, D., Tang, H., Wafula, E., Zheng, C., Albert, V.A., Altman, N.S., Barbazuk,
W.B., Carretero-Paulet, L., dePamphilis, C.W., Der, J.P., Estill, J.C., Jiao, Y., Leebens-
Mack, J., Liu, K., Mei, W., Wafula, E., Altman, N.S., Arikit, S., Axtell, M.J., Chamala,
S., Chanderbali, A.S., Chen, F., Chen, J.-Q., Chiang, V., De Paoli, E., dePamphilis, C.W.,
Der, J.P., Determann, R., Fogliani, B., Guo, C., Harholt, J., Harkess, A., Job, C., Job, D.,
Kim, S., Kong, H., Leebens-Mack, J., Li, G., Li, L., Liu, J., Ma, H., Meyers, B.C., Park,
J., Qi, X., Rajjou, L., Burtet-Sarramegna, V., Sederoff, R., Shahid, S., Soltis, D.E., Soltis,
P.S., Sun, Y.-H., Ulvskov, P., Villegente, M., Xue, J.-Y., Yeh, T.-F., Yu, X., Zhai, J.,
Acosta, J.J., Albert, V.A., Barbazuk, W.B., Bruenn, R.A., Chamala, S., de Kochko, A.,
dePamphilis, C.W., Der, J.P., Herrera-Estrella, L.R., Ibarra-Laclette, E., Kirst, M.,
Leebens-Mack, J., Pissis, S.P., Poncet, V., Schuster, S.C., Soltis, D.E., Soltis, P.S., and
Tomsho, L. 2013. The Amborella Genome and the Evolution of Flowering Plants.
Science 342:1241089.
Andersen, E.J., Nepal, M.P., Purintun, J.M., Nelson, D., Mermigka, G., and Sarris, P.F. 2020.
Wheat Disease Resistance Genes and Their Diversification Through Integrated Domain
Fusions. Front Genet 11:898.
Bailey, J.A., Gu, Z., Clark, R.A., Reinert, K., Samonte, R.V., Schwartz, S., Adams, M.D., Myers,
E.W., Li, P.W., and Eichler, E.E. 2002a. Recent segmental duplications in the human
genome. Science 297:1003-1007.
Bailey, J.A., Gu, Z., Clark, R.A., Reinert, K., Samonte, R.V., Schwartz, S., Adams, M.D., Myers,
E.W., Li, P.W., and Eichler, E.E. 2002b. Recent Segmental Duplications in the Human
Genome. Science 297:1003-1007.
Bergman, M.E., Chávez, Á., Ferrer, A., and Phillips, M.A. 2020. Distinct metabolic pathways
drive monoterpenoid biosynthesis in a natural population of Pelargonium graveolens.
Journal of Experimental Botany 71:258-271.

Birchler, J.A., and Veitia, R.A. 2012. Gene balance hypothesis: Connecting issues of dosage
sensitivity across biological disciplines. Proceedings of the National Academy of
Sciences 109:14746-14753.

Blanc, G., and Wolfe, K.H. 2004. Widespread Paleopolyploidy in Model Plant Species Inferred
from Age Distributions of Duplicate Genes[W]. The Plant Cell 16:1667-1678.

Brosius, J. 1991. Retroposons—Seeds of Evolution. Science 251:753-753.

Buchfink, B., Reuter, K., and Drost, H.-G. 2021. Sensitive protein alignments at tree-of-life scale
using DIAMOND. Nature Methods 18:366-368.

Cantalapiedra, C.P., Hernández-Plaza, A., Letunic, I., Bork, P., and Huerta-Cepas, J. 2021.
eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain
Prediction at the Metagenomic Scale. Mol Biol Evol 38:5825-5829.

Carretero-Paulet, L., Librado, P., Chang, T.-H., Ibarra-Laclette, E., Herrera-Estrella, L., Rozas,
J., and Albert, V.A. 2015. High Gene Family Turnover Rates and Gene Space Adaptation
in the Compact Genome of the Carnivorous Plant Utricularia gibba. Molecular Biology
and Evolution 32:1284-1295.

Casola, C., and Koralewski, T.E. 2018. Pinaceae show elevated rates of gene turnover that are
robust to incomplete gene annotation. Plant J.

Cazzolla Gatti, R., Reich, P.B., Gamarra, J.G.P., Crowther, T., Hui, C., Morera, A., Bastin, J.-F.,
de-Miguel, S., Nabuurs, G.-J., Svenning, J.-C., Serra-Diaz, J.M., Merow, C., Enquist, B.,
Kamenetsky, M., Lee, J., Zhu, J., Fang, J., Jacobs, D.F., Pijanowski, B., Banerjee, A.,
Giaquinto, R.A., Alberti, G., Almeyda Zambrano, A.M., Alvarez-Davila, E., Araujo-
Murakami, A., Avitabile, V., Aymard, G.A., Balazy, R., Baraloto, C., Barroso, J.G.,
Bastian, M.L., Birnbaum, P., Bitariho, R., Bogaert, J., Bongers, F., Bouriaud, O.,
Brancalion, P.H.S., Brearley, F.Q., Broadbent, E.N., Bussotti, F., Castro da Silva, W.,
César, R.G., Češljar, G., Chama Moscoso, V., Chen, H.Y.H., Cienciala, E., Clark, C.J.,
Coomes, D.A., Dayanandan, S., Decuyper, M., Dee, L.E., Del Aguila Pasquel, J.,
Derroire, G., Djuikouo, M.N.K., Van Do, T., Dolezal, J., Đorđević, I.Đ., Engel, J., Fayle,
T.M., Feldpausch, T.R., Fridman, J.K., Harris, D.J., Hemp, A., Hengeveld, G., Herault,
B., Herold, M., Ibanez, T., Jagodzinski, A.M., Jaroszewicz, B., Jeffery, K.J., Johannsen,
V.K., Jucker, T., Kangur, A., Karminov, V.N., Kartawinata, K., Kennard, D.K., Kepfer-
Rojas, S., Keppel, G., Khan, M.L., Khare, P.K., Kileen, T.J., Kim, H.S., Korjus, H.,
Kumar, A., Kumar, A., Laarmann, D., Labrière, N., Lang, M., Lewis, S.L., Lukina, N.,
Maitner, B.S., Malhi, Y., Marshall, A.R., Martynenko, O.V., Monteagudo Mendoza,
A.L., Ontikov, P.V., Ortiz-Malavasi, E., Pallqui Camacho, N.C., Paquette, A., Park, M.,
Parthasarathy, N., Peri, P.L., Petronelli, P., Pfautsch, S., Phillips, O.L., Picard, N., Piotto,
D., Poorter, L., Poulsen, J.R., Pretzsch, H., Ramírez-Angulo, H., Restrepo Correa, Z.,
Rodeghiero, M., Rojas Gonzáles, R.D.P., Rolim, S.G., Rovero, F., Rutishauser, E.,
Saikia, P., Salas-Eljatib, C., Schepaschenko, D., Scherer-Lorenzen, M., Šebeň, V.,
Silveira, M., Slik, F., Sonké, B., Souza, A.F., Stereńczak, K.J., Svoboda, M., Taedoumg,
H., Tchebakova, N., Terborgh, J., Tikhonova, E., Torres-Lezama, A., van der Plas, F.,
Vásquez, R., Viana, H., Vibrans, A.C., Vilanova, E., Vos, V.A., Wang, H.-F.,
Westerlund, B., White, L.J.T., Wiser, S.K., Zawiła-Niedźwiecki, T., Zemagho, L., Zhu,
Z.-X., Zo-Bi, I.C., and Liang, J. 2022. The number of tree species on Earth. Proceedings
of the National Academy of Sciences 119:e2115329119.

Chen, S., Zhou, Y., Chen, Y., and Gu, J. 2018. fastp: an ultra-fast all-in-one FASTQ
preprocessor. Bioinformatics 34:i884-i890.

Conant, G.C., Birchler, J.A., and Pires, J.C. 2014. Dosage, duplication, and diploidization:
clarifying the interplay of multiple models for duplicate gene evolution over time.
Current Opinion in Plant Biology 19:91-98.
Cortés, A.J., Restrepo-Montoya, M., and Bedoya-Canas, L.E. 2020. Modern Strategies to Assess
and Breed Forest Tree Adaptation to Changing Climate. Front Plant Sci 11:583323.
Cortesi, F., Musilová, Z., Stieb, S.M., Hart, N.S., Siebeck, U.E., Malmstrøm, M., Tørresen, O.K.,
Jentoft, S., Cheney, K.L., Marshall, N.J., Carleton, K.L., and Salzburger, W. 2015.
Ancestral duplications and highly dynamic opsin gene evolution in percomorph fishes.
Proceedings of the National Academy of Sciences 112:1493-1498.
Cui, L., Wall, P.K., Leebens-Mack, J.H., Lindsay, B.G., Soltis, D.E., Doyle, J.J., Soltis, P.S.,
Carlson, J.E., Arumuganathan, K., Barakat, A., Albert, V.A., Ma, H., and dePamphilis,
C.W. 2006. Widespread genome duplications throughout the history of flowering plants.
Genome Res 16:738-749.
de Bruijn, W.J.C., Gruppen, H., and Vincken, J.-P. 2018. Structure and biosynthesis of
benzoxazinoids: Plant defence metabolites with potential as antimicrobial scaffolds.
Phytochemistry 155:233-243.
De Smet, R., Adams, K.L., Vandepoele, K., Van Montagu, M.C.E., Maere, S., and Van de Peer,
Y. 2013. Convergent gene loss following gene and genome duplications creates single-
copy families in flowering plants. Proceedings of the National Academy of Sciences
110:2898-2903.
Demuth, J.P., and Hahn, M.W. 2009. The life and death of gene families. Bioessays 31:29-39.
Diss, G., Gagnon-Arsenault, I., Dion-Coté, A.-M., Vignaud, H., Ascencio, D.I., Berger, C.M.,
and Landry, C.R. 2017. Gene duplication can impart fragility, not robustness, in the yeast
protein interaction network. Science 355:630-634.
Dodds, P.N., and Rathjen, J.P. 2010. Plant immunity: towards an integrated view of plant–
pathogen interactions. Nature Reviews Genetics 11:539-548.
Emms, D.M., and Kelly, S. 2017. STRIDE: Species Tree Root Inference from Gene Duplication
Events. Mol Biol Evol 34:3267-3278.
Emms, D.M., and Kelly, S. 2018. STAG: Species Tree Inference from All Genes (bioRxiv).
Enright, A.J., Van Dongen, S., and Ouzounis, C.A. 2002. An efficient algorithm for large-scale
detection of protein families. Nucleic Acids Research 30:1575-1584.
Falcone Ferreyra, M.L., Rius, S., and Casati, P. 2012. Flavonoids: biosynthesis, biological
functions, and biotechnological applications. Frontiers in Plant Science 3.
FAO. 2020. Global Forest Resources Assessment 2020: Main report. Rome.
Force, A., Lynch, M., Pickett, F.B., Amores, A., Yan, Y.L., and Postlethwait, J. 1999.
Preservation of duplicate genes by complementary, degenerative mutations. Genetics
151:1531-1545.
Fu, L., Niu, B., Zhu, Z., Wu, S., and Li, W. 2012. CD-HIT: accelerated for clustering the next-
generation sequencing data. Bioinformatics 28:3150-3152.
Gamfeldt, L., Snäll, T., Bagchi, R., Jonsson, M., Gustafsson, L., Kjellander, P., Ruiz-Jaen, M.C.,
Fröberg, M., Stendahl, J., Philipson, C.D., Mikusiński, G., Andersson, E., Westerlund, B.,
Andrén, H., Moberg, F., Moen, J., and Bengtsson, J. 2013. Higher levels of multiple
ecosystem services are found in forests with more tree species. Nature Communications
4:1340.
Gao, Y., Wang, W., Zhang, T., Gong, Z., Zhao, H., and Han, G.-Z. 2018. Out of Water: The
Origin and Early Diversification of Plant R-Genes. Plant Physiology 177:82-89.

Götz, S., García-Gómez, J.M., Terol, J., Williams, T.D., Nagaraj, S.H., Nueda, M.J., Robles, M.,
Talón, M., Dopazo, J., and Conesa, A. 2008. High-throughput functional annotation and
data mining with the Blast2GO suite. Nucleic Acids Res. 36:3420-3435.
Hahn, M.W., De Bie, T., Stajich, J.E., Nguyen, C., and Cristianini, N. 2005. Estimating the
tempo and mode of gene family evolution from comparative genomic data. Genome Res
15:1153-1160.
Hammond, W.M., Williams, A.P., Abatzoglou, J.T., Adams, H.D., Klein, T., López, R., Sáenz-
Romero, C., Hartmann, H., Breshears, D.D., and Allen, C.D. 2022. Global field
observations of tree die-off reveal hotter-drought fingerprint for Earth's forests. Nature
Communications 13:1761.
Hanada, K., Zou, C., Lehti-Shiu, M.D., Shinozaki, K., and Shiu, S.H. 2008. Importance of
lineage-specific expansion of plant tandem duplicates in the adaptive response to
environmental stimuli. Plant Physiol 148:993-1003.
Hurley, B.P., Garnas, J., Wingfield, M.J., Branco, M., Richardson, D.M., and Slippers, B. 2016.
Increasing numbers and intercontinental spread of invasive insects on eucalypts.
Biological Invasions 18:921-933.
Isabel, N., Holliday, J.A., and Aitken, S.N. 2020. Forest genomics: Advancing climate
adaptation, forest health, productivity, and conservation. Evolutionary Applications 13:3-
10.
Jiang, N., Bao, Z., Zhang, X., Eddy, S.R., and Wessler, S.R. 2004. Pack-MULE transposable
elements mediate gene evolution in plants. Nature 431:569-573.
Jiao, Y., Leebens-Mack, J., Ayyampalayam, S., Bowers, J.E., McKain, M.R., McNeal, J., Rolf,
M., Ruzicka, D.R., Wafula, E., Wickett, N.J., Wu, X., Zhang, Y., Wang, J., Zhang, Y.,
Carpenter, E.J., Deyholos, M.K., Kutchan, T.M., Chanderbali, A.S., Soltis, P.S.,
Stevenson, D.W., McCombie, R., Pires, J.C., Wong, G.K., Soltis, D.E., and Depamphilis,
C.W. 2012. A genome triplication associated with early diversification of the core
eudicots. Genome Biol 13:R3.
Johri, P., Gout, J.-F., Doak, T.G., and Lynch, M. 2022. A Population-Genetic Lens into the
Process of Gene Loss Following Whole-Genome Duplication. Molecular Biology and
Evolution 39:msac118.
Jones, P., Binns, D., Chang, H.Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J.,
Mitchell, A., Nuka, G., Pesseat, S., Quinn, A.F., Sangrador-Vegas, A., Scheremetjew, M.,
Yong, S.Y., Lopez, R., and Hunter, S. 2014. InterProScan 5: Genome-scale protein
function classification. Bioinformatics 30:1236-1240.
Kassambara, A. 2020. ggpubr: 'ggplot2' Based Publication Ready Plots. https://CRAN.R-
project.org/package=ggpubr.
Katoh, K., Misawa, K., Kuma, K.i., and Miyata, T. 2002. MAFFT: a novel method for rapid
multiple sequence alignment based on fast Fourier transform. Nucleic Acids Research
30:3059-3066.
Kirilenko, A.P., and Sedjo, R.A. 2007. Climate change impacts on forestry. Proceedings of the
National Academy of Sciences 104:19697-19702.
Kolde, R. 2019. pheatmap: Pretty Heatmaps. R package version 1.0.12, https://CRAN.R-
project.org/package=pheatmap.
Kumar, S., Stecher, G., Suleski, M., and Hedges, S.B. 2017. TimeTree: A Resource for
Timelines, Timetrees, and Divergence Times. Molecular Biology and Evolution 34:1812-
1819.

Kuzmin, E., VanderSluis, B., Nguyen Ba, A.N., Wang, W., Koch, E.N., Usaj, M., Khmelinskii,
A., Usaj, M.M., van Leeuwen, J., Kraus, O., Tresenrider, A., Pryszlak, M., Hu, M.-C.,
Varriano, B., Costanzo, M., Knop, M., Moses, A., Myers, C.L., Andrews, B.J., and
Boone, C. 2020. Exploring whole-genome duplicate gene retention with complex genetic
interaction analysis. Science 368:eaaz5667.
Leebens-Mack, J.H., Barker, M.S., Carpenter, E.J., Deyholos, M.K., Gitzendanner, M.A.,
Graham, S.W., Grosse, I., Li, Z., Melkonian, M., Mirarab, S., Porsch, M., Quint, M.,
Rensing, S.A., Soltis, D.E., Soltis, P.S., Stevenson, D.W., Ullrich, K.K., Wickett, N.J.,
DeGironimo, L., Edger, P.P., Jordon-Thaden, I.E., Joya, S., Liu, T., Melkonian, B.,
Miles, N.W., Pokorny, L., Quigley, C., Thomas, P., Villarreal, J.C., Augustin, M.M.,
Barrett, M.D., Baucom, R.S., Beerling, D.J., Benstein, R.M., Biffin, E., Brockington,
S.F., Burge, D.O., Burris, J.N., Burris, K.P., Burtet-Sarramegna, V., Caicedo, A.L.,
Cannon, S.B., Çebi, Z., Chang, Y., Chater, C., Cheeseman, J.M., Chen, T., Clarke, N.D.,
Clayton, H., Covshoff, S., Crandall-Stotler, B.J., Cross, H., dePamphilis, C.W., Der, J.P.,
Determann, R., Dickson, R.C., Di Stilio, V.S., Ellis, S., Fast, E., Feja, N., Field, K.J.,
Filatov, D.A., Finnegan, P.M., Floyd, S.K., Fogliani, B., García, N., Gâteblé, G., Godden,
G.T., Goh, F., Greiner, S., Harkess, A., Heaney, J.M., Helliwell, K.E., Heyduk, K.,
Hibberd, J.M., Hodel, R.G.J., Hollingsworth, P.M., Johnson, M.T.J., Jost, R., Joyce, B.,
Kapralov, M.V., Kazamia, E., Kellogg, E.A., Koch, M.A., Von Konrat, M., Könyves, K.,
Kutchan, T.M., Lam, V., Larsson, A., Leitch, A.R., Lentz, R., Li, F.-W., Lowe, A.J.,
Ludwig, M., Manos, P.S., Mavrodiev, E., McCormick, M.K., McKain, M., McLellan, T.,
McNeal, J.R., Miller, R.E., Nelson, M.N., Peng, Y., Ralph, P., Real, D., Riggins, C.W.,
Ruhsam, M., Sage, R.F., Sakai, A.K., Scascitella, M., Schilling, E.E., Schlösser, E.-M.,
Sederoff, H., Servick, S., Sessa, E.B., Shaw, A.J., Shaw, S.W., Sigel, E.M., Skema, C.,
Smith, A.G., Smithson, A., Stewart, C.N., Stinchcombe, J.R., Szövényi, P., Tate, J.A.,
Tiebel, H., Trapnell, D., Villegente, M., Wang, C.-N., Weller, S.G., Wenzel, M.,
Weststrand, S., Westwood, J.H., Whigham, D.F., Wu, S., Wulff, A.S., Yang, Y., Zhu, D.,
Zhuang, C., Zuidof, J., Chase, M.W., Pires, J.C., Rothfels, C.J., Yu, J., Chen, C., Chen,
L., Cheng, S., Li, J., Li, R., Li, X., Lu, H., Ou, Y., Sun, X., Tan, X., Tang, J., Tian, Z.,
Wang, F., Wang, J., Wei, X., Xu, X., Yan, Z., Yang, F., Zhong, X., Zhou, F., Zhu, Y.,
Zhang, Y., Ayyampalayam, S., Barkman, T.J., Nguyen, N.-p., Matasci, N., Nelson, D.R.,
Sayyari, E., Wafula, E.K., Walls, R.L., Warnow, T., An, H., Arrigo, N., Baniaga, A.E.,
Galuska, S., Jorgensen, S.A., Kidder, T.I., Kong, H., Lu-Irving, P., Marx, H.E., Qi, X.,
Reardon, C.R., Sutherland, B.L., Tiley, G.P., Welles, S.R., Yu, R., Zhan, S., Gramzow,
L., Theißen, G., Wong, G.K.-S., and One Thousand Plant Transcriptomes, I. 2019. One
thousand plant transcriptomes and the phylogenomics of green plants. Nature 574:679-
685.
Leister, D. 2004. Tandem and segmental gene duplication and recombination in the evolution of
plant disease resistance gene. Trends Genet 20:116-122.
Li, B., and Dewey, C.N. 2011. RSEM: accurate transcript quantification from RNA-Seq data
with or without a reference genome. BMC Bioinformatics 12:323.
Linnakoski, R., and Forbes, K.M. 2019. Pathogens—The Hidden Face of Forest Invasions by
Wood-Boring Insect Pests. Frontiers in Plant Science 10.
Liu, J., Yu, T., Mu, Z., and Li, G. 2019. TransLiG: a de novo transcriptome assembler that uses
line graph iteration. Genome Biology 20:81.

Lu, M., Feau, N., Vidakovic, D.O., Ukrainetz, N., Wong, B., Aitken, S.N., Hamelin, R.C., and
778   Yeaman, S. 2021. Comparative Gene Expression Analysis Reveals Mechanism of Pinus
779   contorta Response to the Fungal Pathogen Dothistroma septosporum. Mol Plant Microbe
780   Interact:Mpmi10200282r.
781   Lynch, M., and Conery, J.S. 2000. The evolutionary fate and consequences of duplicate genes.
782   Science 290:1151-1155.
783   Lynch, M., and Conery, J.S. 2003. The evolutionary demography of duplicate genes. Journal of
784   Structural and Functional Genomics 3:35-44.
785   Maere, S., De Bodt, S., Raes, J., Casneuf, T., Van Montagu, M., Kuiper, M., and Van de Peer, Y.
786   2005. Modeling gene and genome duplications in eukaryotes. Proceedings of the
787   National Academy of Sciences 102:5454-5459.
788   Manni, M., Berkeley, M.R., Seppey, M., Simão, F.A., and Zdobnov, E.M. 2021. BUSCO
789   Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic
790   Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. Molecular Biology
791   and Evolution 38:4647-4654.
792   Mendes, F.K., Vanderpool, D., Fulton, B., and Hahn, M.W. 2020. CAFE 5 models variation in
793   evolutionary rates among gene families. Bioinformatics 36:5516-5518.
794   Meng, X., and Zhang, S. 2013. MAPK cascades in plant disease resistance signaling. Annu Rev
795   Phytopathol 51:245-266.
796   Meyers, B.C., Kozik, A., Griego, A., Kuang, H., and Michelmore, R.W. 2003. Genome-Wide
797   Analysis of NBS-LRR–Encoding Genes in Arabidopsis[W]. The Plant Cell 15:809-834.
798   Miura, S., Amacher, M., Hofer, T., San-Miguel-Ayanz, J., Ernawati, and Thackway, R. 2015.
799   Protective functions and ecosystem services of global forests in the past quarter-century.
800   Forest Ecology and Management 352:35-46.
801   Neale, D.B., and Kremer, A. 2011. Forest tree genomics: growing resources and applications.
802   Nat Rev Genet 12:111-122.
803   Ngou, B.P.M., Heal, R., Wyler, M., Schmid, M.W., and Jones, J.D.G. 2022. Concerted
804   expansion and contraction of immune receptor gene repertoires in plant genomes. Nature
805   Plants 8:1146-1152.
806   Ohno, S. 1970. Evolution by gene duplication (Springer-Verlag).
807   Panchy, N., Lehti-Shiu, M., and Shiu, S.H. 2016. Evolution of Gene Duplication in Plants. Plant
808   Physiol 171:2294-2316.
809   Piasecka, A., Jedrzejczak-Rey, N., and Bednarek, P. 2015. Secondary metabolites in plant innate
810   immunity: conserved function of divergent chemicals. New Phytologist 206:948-964.
811   Picart-Picolo, A., Grob, S., Picault, N., Franek, M., Llauro, C., Halter, T., Maier, T.R., Jobet, E.,
812   Descombin, J., Zhang, P., Paramasivan, V., Baum, T.J., Navarro, L., Dvořáčková, M.,
813   Mirouze, M., and Pontvianne, F. 2020. Large tandem duplications affect gene expression,
814   3D organization, and plant-pathogen response. Genome Res 30:1583-1592.
815   Price, M.N., Dehal, P.S., and Arkin, A.P. 2009. FastTree: computing large minimum evolution
816   trees with profiles instead of a distance matrix. Mol Biol Evol 26:1641-1650.
817   Prunier, J., Caron, S., Lamothe, M., Blais, S., Bousquet, J., Isabel, N., and MacKay, J. 2017.
818   Gene copy number variations in adaptive evolution: The genomic distribution of gene
819   copy number variations revealed by genetic mapping and their adaptive role in an
820   undomesticated species, white spruce (*Picea glauca*). Mol Ecol 26:5989-6001.

821   Puttick, M.N., Clark, J., and Donoghue, P.C.J. 2015. Size is not everything: rates of genome size
822   evolution, not C-value, correlate with speciation in angiosperms. Proceedings of the
823   Royal Society B: Biological Sciences 282:20152289.
824   R Core Team. 2021. R: A language and environment for statistical computing. R Foundation for
825   Statistical Computing, Vienna, Austria. https://www.R-project.org/.
826   Reid, N.M., Proestou, D.A., Clark, B.W., Warren, W.C., Colbourne, J.K., Shaw, J.R., Karchner,
827   S.I., Hahn, M.E., Nacci, D., Oleksiak, M.F., Crawford, D.L., and Whitehead, A. 2016.
828   The genomic landscape of rapid repeated evolutionary adaptation to toxic pollution in
829   wild fish. Science 354:1305-1308.
830   Rizvi, A.R., Baig, S., Barrow, E., and Kumar, C. 2015. Synergies between climate mitigation
831   and adaptioon in forest landscape restoration. Gland, Switzerland: IUCN.
832   Rodgers-Melnick, E., Mane, S.P., Dharmawardhana, P., Slavov, G.T., Crasta, O.R., Strauss,
833   S.H., Brunner, A.M., and Difazio, S.P. 2012. Contrasting patterns of evolution following
834   whole genome versus tandem duplication events in Populus. Genome Res 22:95-105.
835   Sanderson, M.J. 2003. r8s: inferring absolute rates of molecular evolution and divergence times
836   in the absence of a molecular clock. Bioinformatics 19:301-302.
837   Seoighe, C., and Gehring, C. 2004. Genome duplication led to highly selective expansion of the
838   Arabidopsis thaliana proteome. Trends in Genetics 20:461-464.
839   Soltis, P.S., and Soltis, D.E. 2016. Ancient WGD events as drivers of key innovations in
840   angiosperms. Curr Opin Plant Biol 30:159-165.
841   Swenson, N.G., and Jones, F.A. 2017. Community transcriptomics, genomics and the problem of
842   species co-occurrence. Journal of Ecology 105:563-568.
843   Tang, H., Bowers, J.E., Wang, X., Ming, R., Alam, M., and Paterson, A.H. 2008. Synteny and
844   Collinearity in Plant Genomes. Science 320:486-488.
845   Trouern-Trend, A.J., Falk, T., Zaman, S., Caballero, M., Neale, D.B., Langley, C.H., Dandekar,
846   A.M., Stevens, K.A., and Wegrzyn, J.L. 2020. Comparative genomics of six Juglans
847   species reveals disease-associated gene family contractions. The Plant Journal 102:410-
848   423.
849   Van de Peer, Y., Maere, S., and Meyer, A. 2009. The evolutionary significance of ancient
850   genome duplications. Nature Reviews Genetics 10:725-732.
851   Wickham, H. 2011. The Split-Apply-Combine strategy for data analysis. J. Stat. Softw. 40:1-29.
852   Wickham, H. 2016. ggplot2: Elegant graphics for data analysis. Springer-Verlag New York, NY,
853   US.
854   Yang, Z. 2007. PAML 4: Phylogenetic analysis by maximum likelihood. Mol. Biol. Evol.
855   24:1586-1591.
856   Yang, Z., and Nielsen, R. 2000. Estimating synonymous and nonsynonymous substitution rates
857   under realistic evolutionary models. Mol Biol Evol 17:32-43.
858   Yu, Y. 2022. mixR: An R package for finite mixture modeling for both raw and binned data.
859   Journal of Open Source Software 7:4031.
860   Zeng, L., Zhang, N., Zhang, Q., Endress, P.K., Huang, J., and Ma, H. 2017. Resolution of deep
861   eudicot phylogeny and their temporal diversification using nuclear genes from
862   transcriptomic and genomic datasets. New Phytol 214:1338-1354.
863   Zhang, J. 2003. Evolution by gene duplication: an update. Trends in Ecology & Evolution
864   18:292-298.
865   Zhao, J., Davis, L.C., and Verpoorte, R. 2005. Elicitor signal transduction leading to production
866   of plant secondary metabolites. Biotechnol Adv 23:283-333.

867   Zwaenepoel, A., and Van de Peer, Y. 2019. Inference of Ancient Whole-Genome Duplications
868   and the Evolution of Gene Duplication and Loss Rates. Molecular Biology and Evolution
869   36:1384-1404.
870