# Sex chromosome – autosome fusions in *Heliconius* butterflies

Nicol Rueda - M[1], Chris D. Jiggins[2], Carolina Pardo-Diaz[1], W. Owen McMillan[3],

Shane McCarthy[4], Krzysztof M. Kozak[3,7], Carlos F. Arias[3,8], Jonathan Ready[5,6],

Richard Durbin[2,4], Gabriela Montejo-Kovacevich[2], Joana I. Meier[2,4*], Camilo

Salazar[1*✉]

* These authors contributed equally to this work.

[1]Biology Program, Faculty of Natural Sciences, Universidad del Rosario, Bogotá, Colombia. 111221

[2]Department of Zoology, University of Cambridge, Cambridge, CB2 3EJ, United Kingdom

[3]Smithsonian Tropical Research Institute, Panama

[4]Tree of Life Programme, Welcome Sanger Institute, United Kingdom

[5]Centre for Advanced Studies of Biodiversity - CEABIO, Guamá Science Park, Lot 11, Avenida Perimetral da Ciência km 01 - Guamá, Belém - PA, 66075-750, Brazil

[6]Institute for Biological Sciences, Federal University of Pará - UFPA, R. Augusto Corrêa, 1 - Guamá, Belém - PA, 66075-110, Brazil

[7]Museum of Vertebrate Zoology, UC Berkeley, USA

[8]Data Science Lab, Office of the Chief Information Officer, Smithsonian Institution, Washington, DC, United States

✉Correspondence: Camilo Salazar (camilo.salazar@urosario.edu.co)

## Abstract

Sex chromosomes are evolutionarily labile in many animals, and fusion with an autosome is one of the mechanisms by which they can evolve. Sex chromosome-autosome (SA) fusions can reduce sexual conflict and promote adaptation and reproductive isolation among species. Recently, advances in genomics have fuelled the discovery of SA fusions in multiple vertebrates and in some invertebrates such as Lepidoptera. Here, we discovered multiple SA fusions in the *sara/sapho* clade of the classical adaptive radiation of *Heliconius* butterflies. While the vast majority of *Heliconius* species have 21 chromosomes, species of this particularly rapidly diversifying clade have up to 60 chromosomes. We found a sex chromosome fusion with chromosome 4 at the base of the *sapho* clade and two additional sex chromosome fusions, each shared by two species. These sequential fusions between autosomes and sex chromosomes make the *Heliconius sapho* clade an ideal system to study the role of neo-sex chromosomes in adaptive radiations and the degeneration of sex chromosomes over time. Our study adds to a growing number of examples in butterflies that, in the future, may help to unravel the importance of such rearrangements in the evolution of Lepidoptera and animals in general.

## Keywords

Sex chromosome-autosome fusions, *Heliconius*, sex chromosome evolution, adaptation.

## Introduction

**Introduction**

Sex chromosome-autosome (SA) fusions contribute to the turnover of sex-

determination loci or the evolution of neo-sex chromosomes (Ross et al. 2009;

Yoshido et al. 2011; Šíchová et al. 2016), but it remains unclear what promotes

them. Sexually antagonistic selection, the sheltering of deleterious mutations, direct

selection, genetic drift or meiotic drive have all been suggested as possible drivers

of SA fusions (Kitano et al. 2009; Pennell et al. 2015; Sigeman et al. 2019; Jay et

al. 2022). Sexually antagonistic selection is thought to favour the fusion of sex

chromosomes with autosomes harbouring genes under sexually antagonistic

selection (Charlesworth and Charlesworth 1980). There is limited evidence for this

theory e.g. in sticklebacks (Ross et al. 2009) and warblers (Pala et al. 2012). An

alternative hypothesis is deleterious mutation sheltering, where SA fusions are

favoured because they prevent the expression of recessive deleterious alleles in

the heterogametic sex (Jay et al. 2022). SA fusions can also become fixed due to

meiotic drive, such as female meiotic drive elements on W/X-A fusions that

preferentially end up in the egg instead of the polar bodies (Pardo-Manuel de

Villena and Sapienza 2001).

As with other chromosomal rearrangements, SA fusions can suppress

recombination and potentially strengthen reproductive isolation if they bring

together barrier loci into regions with reduced recombination (Guerrero and

Kirkpatrick 2014; Lucek 2018; Hora et al. 2019; de Vos et al. 2020). For instance,

in the Japanese threespine stickleback *Gasterosteus aculeatus,* an SA fusion

resulted in a neo-X chromosome that linked loci underlying behavioural isolation

traits and hybrid sterility (Kitano et al. 2009). SA fusions can also facilitate

adaptation, such as the SA fusion in *Cydia pomonella* (Tortricidae), which linked

two insecticide-resistance genes and genes involved in detoxifying plant

metabolites (Nguyen et al. 2013).

Genomic studies have revealed that SA fusions have occurred multiple times

across vertebrates (Schmid et al. 2003; Matsumoto and Kitano 2016; Sigeman et al.

2019), and in some invertebrates such as mygalomorph spiders (Král et al. 2013),

*Drosophila* (Flores et al. 2008) and true bugs of the genus *Dysdercus* (Bressa et al.

2009). In Lepidoptera, examples of SA fusions include *Danaus* (Smith et al. 2016;

Mongue et al. 2017) and *Leptidea* butterflies (Šíchová et al. 2015; Šíchová et al.

2016), among others (Yoshido et al. 2005; Nguyen et al. 2013; Picq et al. 2018;

Carabajal Paladino et al. 2019). Compared to taxa with centromeres, the holocentric

chromosomes of Lepidoptera may facilitate fusions, as they are less likely to cause

segregation problems during cytokinesis and thus reduce hybrid fitness (Lucek et al.

2022). Nonetheless, butterflies and moths have remarkably constrained

chromosome evolution (White 1977), with most species having a haploid

chromosome number ranging between 28 and 32 except for a few groups that have

experienced extensive fission and fusion events (Hill et al. 2019). One of such cases

are the *Heliconius* butterflies, which mostly have 21 chromosomes due to ancestral

fusions (Davey et al. 2016). Only a few species in the genus differ from this ancestral

chromosome number, especially species in the *sara/sapho* clade, with some having

up to 60 chromosomes (Brown 1981). This clade comprises 12 species (Jiggins and

Lamas 2016), and differs from other *Heliconius* by showing an inability to synthesise

cyanogens, leading to reliance on toxins sequestered from larval host plants (Beltrán

et al. 2007; Engler-Chaouat and Gilbert 2007). A subclade of seven species within

the *sara/sapho* clade shows particularly high diversification rates (Kozak et al., 2015)

and high number of chromosomes (Brown 1981).

We generated whole genome resequencing data for 114 individuals from

these seven species in the *sara/sapho* clade to: (i) investigate genome-wide

phylogenetic relationships, (ii) determine the degree of genomic differentiation

between species and subspecies, and (iii) explore the impact of chromosomal

rearrangements . Despite a high number of chromosomes in this clade, suggesting

many chromosomal fissions, we find at least three SA fusions across five species,

making them a prime study system for the evolution of neo-sex chromosomes.

**Materials and Methods**

**Genome assembly of *Heliconius sara***

We used a female of *H. sara* (SAMEA5394385; Table S1) to generate an

assembled genome for this species. We assembled the genome by combining

PacBio, HiC and 10X data, all generated by the Tree of Life Programme and the

Wellcome Sanger Institute (https://www.sanger.ac.uk/programme/tree-of-life/). The

libraries were produced following the manufacturer's guidelines. We sequenced

four SMRT (single-molecule real-time) cells with the PacBio Sequel II system.

Dovetail Hi-C data was sequenced on a HiSeq X Ten platform. A 10X Genomics

Chromium linked-read sequencing library was sequenced with 150-bp paired-end

reads on four lanes on an Illumina HiSeq X Ten platform.

An Initial contig assembly was generated from the PacBio CLR data using wtdbg2

v2.2 (Ruan and Li 2020) . The PacBio data was then used to polish the contigs

using Arrow (https://github.com/PacificBiosciences/GenomicConsensus). The 10X

Chromium linked-reads were then mapped to the assembly using Longranger v2.2.

(10X Genomics), variants called using freebayes v1.1.0-3-g961e5f3 (Garrison and

Marth 2012) . Next, the assembly was polished using BCFtools consensus v1.9

(Danecek et al. 2021) by applying homozygous non-reference calls as edits. The

10X linked-reads were then used to scaffold contigs using Scaff10X v2.3

(https://github.com/wtsi-hpag/Scaff10X). A round of manual curation was

performed on these polished scaffolds using gEVAL (Chow et al. 2016) . Lastly,

Dovetail Genomics Hi-C data was used to scaffold the assembly further using

SALSA v2.2 (Ghurye et al. 2019), followed by another round of manual curation

with gEVAL (Chow et al. 2016). The chromosome-scale scaffolds were named by

synteny to the *Heliconius melpomene melpomene* assembly Hmel2.5 in LepBase.

We assessed the genome contiguity with gnx-tools (https://github.com/mh11/gnx-

tools/blob/master/README) and genome completeness with BUSCO v5.1.2

(Manni et al. 2021) using the Lepidoptera gene set.

**Sample collection for genome resequencing**

We collected 114 individuals of *Heliconius* from 7 species and 18 subspecies in the

*sara/sapho* clade across their distribution range: 48 *H. sara*, 2 *H. leucadia*, 21 *H.*

*antiochus*, 13 *H. sapho*, 3 *H. hewitsoni*, 17 *H. eleuchia* and 10 *H. congener* (Table

S1). The body of each was preserved in NaCl-saturated DMSO solution and stored

at -80°C; wings were kept for phenotype reference.

**Whole-genome resequencing and genotype calling**

Genomic DNA was extracted from thoracic tissue using a DNeasy Blood and

Tissue Kit (Qiagen). Library preparation and whole-genome Illumina resequencing

(PE reads) was carried out on Illumina's HiSeq X system by Novogene (Beijing,

China), with 30X coverage per individual. We also downloaded two samples of *H.*

*charithonia* (SRR4032025 – SRR4032026) from SRA

([https://www.ncbi.nlm.nih.gov/sra](https://www.ncbi.nlm.nih.gov/sra)) to include them as outgroups in phylogenetic

analyses. The *H. sara* genome (HelSar1) was used as a reference to map the

reads of each individual using BWA mem v0.7.12 (Li 2013) with default

parameters. We then used samtools v1.12 to sort and index the alignment files (Li

et al. 2009). PCR-duplicate reads were identified and removed using Picard tools

v2.9.2 (Broad 2019), and variant calling was conducted with Haplotype Caller

(GATK, v3.7.0) in BP-resolution mode (DePristo et al. 2011). Then, samples were

jointly genotyped using GATK's GenotypeGVCFs (DePristo et al. 2011). We used

vcftools v0.1.14 (Danecek et al. 2011) and the final VCF to calculate: (i) mean

depth per individual and site, (ii) quality per site, (iii) the proportion of missing data

per individual and (iv) the proportion of missing data per site, and (v) percentage

mapping per individual. Based on these results, we kept sites with quality value (--

minQ) ≥ 30 and less than 5% missing data. We also excluded sites with a depth

below 5 and mean depth per individual less than 1.5 times the mean to exclude

paralogous regions (Meier et al. 2018). For this, we used the custom script

*removeTooHighDepthSites.sh* from Joana Meier's Github repository

([https://github.com/joanam](https://github.com/joanam)). We removed sites with excess heterozygosity using

the vcftools option --*hardy* and a p-value cut-off of <1e-5. Finally, we checked our

samples for contamination or other issues generating erroneous heterozygotes

using the python script *checkHetsIndvVCF.sh* by David Marques, and modified by

Jessica Rick (https://github.com/jessicarick/lates-popgen/blob/master/scripts/).

**Phylogenetic analysis**

We generated a whole-genome ML tree using a vcf containing all sites as input in

RAXML v8.2.9 (Stamatakis et al. 2008), with the GTRGAMMA model and 100

bootstrap replicates. We applied the same procedure to obtain ML trees for each

chromosome. In all cases, *H. charitonia* was set as outgroup.

**Population structure and shared ancestry**

We performed a principal component analysis (PCA) to study the genetic structure

of populations. We filtered out monomorphic or multiallelic sites, and sites with

minor allele frequency (MAF) smaller than 0.1 with vcftools (Danecek et al. 2011).

To reduce the linkage disequilibrium effect, we used the python script *ldPruning.sh*

(https://github.com/joanam/scripts/blob/master/ldPruning.sh), which removes sites

with $r^2$>0.2 in windows of 50 Kbp sliding by 10 Kbp. We conducted the PCA using

Plink v2.0 with default parameters (Purcell and Chang 2007; Chang et al. 2015).

We used a vcf with 3,685,916 high-quality SNPs sites.

We analysed patterns of population structure using the R package *lostruct*

(Li and Ralph 2019) on chromosomes revealing sex-clustering phylogenies. We

performed this analysis between pairs of species (*H. eleuchia - H. congener* and *H.*

*sapho - H. hewitsoni*), and between subspecies of *H. antiochus*. We first divided

the chromosomes into non-overlapping windows of 100 SNPs and then performed

a PCA for each window to identify regions that might indicate variation in the

chromosome structure. We next used a two-dimension space multidimensional

scaling (MDS) analysis to illustrate how dissimilar the patterns of relatedness are

between each pair of windows. We then manually selected sets of windows to

calculate PCAs and test which regions do not group samples by species. Finally,

we used R (R Core Team 2013) to visualise the PCA results and plot the first two

MDS coordinates against the midpoint of each window to visualise variation across

the genome.

The genetic ancestry of each sample was estimated using ADMIXTURE v.

1.3.0 (Alexander et al. 2009). Input data were prepared using the same procedure

for the genome wide PCA analysis. We used K = 1-6 clusters and the cross-

validation method to estimate the best K value. All other parameters were set to

default.

## Patterns of genetic differentiation

We calculated $F_{ST}$ and $D_{XY}$ by pairs of sister species and by subspecies along

chromosomes in non-overlapping 50 Kbp windows. Because *H. antiochus* did not

have a sister species, we calculated these statistics between Andean and

Amazonian subspecies. Windows that contained less than 2,500 high-quality

genotyped variable sites were rejected. We used a dataset including SNPs and

monomorphic sites and the *popgenWindows.py* script from Simon Martin's General

Genomics GitHub repository

(https://github.com/simonhmartin/genomics_general/blob/master/popgenWindows.

py).

**Patterns of heterozygosity and mean depth by chromosome**

To study chromosomes with $F_{ST}$ and $D_{XY}$ patterns different from the genome

average, we used the options --het and --depth of vcftools v. 0.1.14 (Danecek et al.

2011), and calculated heterozygosity and mean depth per chromosome for each

individual of each species. We also calculated these statistics in 50 Kbp non-

overlapping sliding windows along the 'outlier' chromosomes identified. On these,

we calculated π specifying each individual as its own population, so π became a

measure of heterozygosity. This was done with the Python script

*popgenWindows.py* from the Simon Martin's General Genomics GitHub repository

(https://github.com/simonhmartin/genomics_general/blob/master/popgenWindows.

py). We then averaged these values across all individuals of the same sex and

species. For depth, we first generated a file containing the mean depth per site

averaged across all individuals of the same sex and species using the --site-mean-

depth option of vcftools v. 0.1.14 (Danecek et al. 2011). We then used the R

package windowscanr v. 0.1 from Hugo Tavares' GitHub repository

(https://github.com/tavareshugo/WindowScanR) to calculate the mean of the mean

depth per species, per sex, and per window. The few individuals of *H. hewitsoni*

and *H. leucadia* were not included in the sliding windows analysis. Statistical tests

were applied to assess significant differences in heterozygosity and mean depth

between sexes and between chromosomes. As the data were not normally

distributed, we performed a Wilcoxon signed rank test to compare sexes. To

assess differences between chromosomes, we applied a Kruskal-Wallis test and a

*post hoc* test (pairwise Wilcoxon test for Kruskal-Wallis).

## Results

### Genome assembly

The final assembled genome was 349 Mb (with 46,300 bp of Ns) in 399 scaffolds,

with scaffold size increasing from 2,007 bp to 22,4 Mb, and with a N50 of 17,8 Mb

which is better than the currently best *Heliconius* reference genomes (*H.*

*melpomene*: 14.3 Mb, *H. erato demophoon*: 10.7 Mb). Hi-C assigned 36 scaffolds

to 20 chromosomes and the Z chromosome, whilst the remaining 363 scaffolds

were unassigned. The BUSCO completeness is 98.4%, of which 98.2% are single-

copy BUSCOs.

### Whole-genome resequencing dataset

A total of 114 individuals were successfully sequenced and 112 were included in

downstream analyses. The average mapping percentage to the *H. sara* genome

was 95.56% (range: 77.38% - 99.14%; Table S1). Two individuals with a

particularly high proportion of missing data (19.5% and 39.5% respectively) and

low mean coverage (7.4X and 9.1X, respectively) were excluded from further

analyses (Table S1, Figure S1). In contrast, *H. sara* and *H. leucadia* had ≤10%

missing data per individual, much less than any other species (Figure S1) and

likely explained by close similarity to the reference genome. The mean coverage of

all other samples ranged between 10X and 27X (Table S1, Figure S1).

**Phylogenetic relationships**

The ML phylogenetic tree obtained with 183,282,470 sites separated individuals

into groups consistent with both PCA and ADMIXTURE analyses, and revealed

two main clades: (i) the *sara* clade and (ii) the *sapho* clade. The *sara* clade is

composed of two species, namely *H. sara* and *H. leucadia*, where *H. sara* is

subdivided into an Andean subgroup (*H. s. magdalena, H. s. sprucei,* and *H. s.*

*elektra*) and an Amazonian subgroup (*H. s. sara*). The *sapho* clade is split into two

well-resolved lineages (*H. antiochus* and a clade composed of two monophyletic

groups: *H. eleuchia/H. congener* and *H. sapho/H. hewitsoni*). In this case, *H.*

*antiochus* appeared as a monophyletic group split into an Andean clade (*H. a.*

*aranea* and *H. a. araneides*), and an Amazonian clade (*H. a. antiochus* and *H. a.*

*salvini*; Figure 1). *H. antiochus* nested into the *sapho* clade whereas *H. hewitsoni*

was found as sister to *H. sapho* thus resolving the previously undetermined

position of these species (Kozak et al. 2015).

We found strong phylogenetic incongruence across chromosomes. For

instance, the whole-genome topology was only recovered in eight chromosomes

(Chr1, Chr2, Chr10, Chr11, Chr13, Chr15, Chr18, and Chr19; Figure 2a and S2-

S22), while nine chromosomes showed *H. congener* varying its position in the

phylogeny and appearing as sister either to *H. e. eleuchia* or to a clade composed

of *H. e. eleusinus + H. e. primularis* (Chr3, Chr4, Chr6, Chr8, Chr9, Chr12, Chr17,

Chr20, and Chr21; Figure 2a and S2-S22). Similarly, the position of *H. hewitsoni* in

relation to *H. sapho* varied in eight chromosomes (Chr5, Chr6, Chr7, Chr14, Chr16,

Chr17, Chr20, Chr21; Figure 2a and S2-S22).

**Figure 1. Phylogeny and distribution of the sara/sapho clade.** Genome-wide ML phylogeny, highlighting

the position of the Sara and the Sapho clade, with bootstrap support values on the branches. Orange triangles

highlight clades where SA fusions were detected, and the SA fusion is specified. The distribution of all 17

subspecies in the clade from (Rueda-M et al., 2021) is shown, where dots represent sampling locations for the

present study. Each of the 17 subspecies is represented by a unique colour, that is the same in the phylogeny

and the distribution.

Interestingly, we found sex-specific clustering on Chr4, Chr9 and Chr14

(Figure 2b). In Chr4, all species in the *sapho* clade showed females and males

clustering separately, while males of *H. congener* and *H. eleuchia* formed a clade

apart from their conspecific females (Figure 2b). This sex clustering pattern was

also observed on Chr9 for *H. sapho* and *H. hewitsoni* (Figure 2b). In addition,

males and females of *H. eleuchia* and *H. congener* also formed a clade in Chr14

(Figure 2b)**.**

**Population structure and shared ancestry**

The PCA grouped samples into three main groups: (i) *H. sara and leucadia*

(hereafter *sara* clade), (ii) *H. antiochus*, and (iii) *H. eleuchia, H. congener, H. sapho*

and *H. hewitsoni* (hereafter *sapho* clade). The first two principal components

explained 60% (PC1) and 10% (PC2) of the total variance. PC1 separated the *sara*

clade and *H. charithonia* (outgroup) from *H. antiochus* and the *sapho* clade,

whereas PC2 separated *H. antiochus* from the *sapho* clade. *H. sapho* was closer

to *H. hewitsoni,* whereas *H. eleuchia* was closer to *H. congener* (Figure S23).

Results from ADMIXTURE were largely consistent with the PCA recovering *H.*

*antiochus*, *H. eleuchia, H. sapho* and *H. congener* as independent groups. We also

identified Andean subspecies of *H. sara* (*H. s. magdalena, H. s. sprucei,* and *H. s.*

*elektra*) and Amazonian subspecies of *H. sara* (*H. s. sara*) as separate groups. *H.*

*hewitsoni* and *H. leucadia* could not be correctly assigned as a separate group due

to the low number of samples. ADMIXTURE analysis did not reveal any evidence

of recent shared ancestry among species (Figure S24).

**Figure 2. ML phylogenies inferred genome-wide and per chromosome.** (a) Topologies found across the genome. Purple: genome-wide topology. Green: *H. congener* within *H. eleuchia*. Orange: *H. hewitsoni* as sister to *H. congener + H. eleuchia*. Brown: *H. hewitsoni* as sister to *H. congener + H. eleuchia + H. sapho*. Chromosomes are shown in the bottom, with coloured triangles indicating the topology revealed by each of them. (b) Topologies showing sex-specific grouping within some species, which is indicative of SA fusions in Chr4, Chr9 and Chr14. In these species, females are coloured in red and males in blue.

The local PCA confirmed the sex specific patterns on Chr4, Chr 9 and Chr14 (Figure 3 and Figure S25-S29). The MDS analysis summarising the PCA results of each 100 SNP window on Chr4 in *H. eleuchia*, *H. congener*, *H. sapho* and *H. hewitsoni* revealed two clusters, one of windows grouping the individuals by species, and the second grouping the individuals by sex (Figure 3a-3b and 3d-3e). In *H. antiochus*, all windows in Chr4 cluster individuals by sex (Figure S27). Chr14 behaves like Chr4 but only in *H. eleuchia* and *H. congener,* where sex-grouping is observed across the entire chromosome except for the ends, where grouping is by species (Figure S28). Finally, Chr9 shows a pattern where PC1 of each window groups *H. sapho* and *H. hewitsoni* individuals by species and PC2 groups them by

sex (Figure S29). Strikingly, in all species, sex-grouping windows are scattered along

Chr4, Chr9 and Chr14 (Figure 3c, 3f and S28c).

**Figure 3. Local PCA in 100 SNP windows on Chr4 reveals grouping by sex across the entire**

**chromosome. (a)** two-dimensional MDS plot of the distribution of 100 SNP windows along Chr4 for the *H.*

*congener/H. eleuchia* clade, **(b)** PCA plots for the black and grey windows in (a). **(c)** Actual distribution for

the black and grey windows in (a) along Chr4. **(d)** two-dimensional MDS plot of the distribution of 100 SNP

windows along Chr4 for the *H. sapho/H. hewitsoni* clade. **(e)** PCA plots for the black and grey windows in

(d). **(f)** Actual distribution for the black and grey windows in (d) along Chr4. Each point in (a) and (d)

represents one window. Black: windows that cluster by sex in the local PCA. Grey: windows that cluster by

species in the local PCA. Each individual in (b) and (e) is represented by a point with a species-specific

symbol and colour according to its sex.

## Patterns of genetic differentiation

Genomic differentiation ($F_{ST}$) was strongest between *H. sapho* vs. *H. hewitsoni*

followed by *H. congener* vs. *H. eleuchia,* and Andean vs. Amazonian *H. antiochus*

(average $F_{ST}$ = 0.33, 0.26, and 0.07 respectively; Figure 4). *H. sara* vs. *H. leucadia*

were the least differentiated pair (average $F_{ST}$ = 0.05, Figure 4). In contrast,

absolute divergence ($D_{XY}$) was highest in *H. sara* vs. *H. leucadia* while all other

species pairs exhibited a similar degree of divergence (Figure 4). We also

observed elevated $F_{ST}$ values on the Z chromosome compared to autosomes in all

but one comparison (*H. eleuchia* vs. *H. congener*; Figure 4). Interestingly, Chr4

shows lower $F_{ST}$ values and increased $D_{XY}$ values (in all comparisons but *H. sara*

vs. *H. leucadia*). Only in *H. sapho* vs. *H. hewitsoni* Chr9 showed lower $F_{ST}$ but did

not show higher $D_{XY}$ (Figure 4). Chr14 behaves like Chr9 but only in *H. eleuchia*

and *H. congener* (Figure 4). The same pattern was true when we compared $F_{ST}$

and $D_{XY}$ between subspecies (Fig. S30 – S33). Interestingly, $F_{ST}$ values peaked at

the end of Chr14 when comparing between subspecies of *H. congener*, in the

same region that groups individuals by species and not sex in the local PCA

(Figure S30).

**Figure 4. Genome-wide divergence ($F_{ST}$) between pairs of species in the *sara/sapho* clade.** Each dot represents a chromosome, and chromosomes with evidence of SA fusions are colour coded. Analyses for pairs of subspecies are shown in Figure S30 and S33.

## Sex-specific differences in heterozygosity and coverage on multiple chromosome

The grouping by sex we observed in the local PCA and the trees exclusively of Chr4, Chr9 and Chr14 suggest possible fusions between these autosomes and the Z or W chromosome. Because females of Lepidoptera lack crossing over, and their meiosis is achiasmatic, they do not recombine (Suomalainen et al. 1973). This means that, if the Z chromosome is involved in the fusion (Figure 5a), only females would have unfused autosomes that would tend to accumulate mutations and/or structural variants leading to divergence from its (now fused) homologous. Alternatively, if the W chromosome is involved in the fusion (Figure 5b), the SA fusion would be restricted to females and the fused SA chromosome would tend to accumulate mutations and/or structural variants leading to divergence from its unfused homologous. Both scenarios would lead to high heterozygosity and low depth in females (Figure 5). Consistently, Chr4, Chr9 and Chr14 showed striking sex-specific differences in heterozygosity in multiple species (Figure 6). The strongest difference in heterozygosity was observed on Chr4, where females of *H. eleuchia, H. congener, H. sapho, H. hewitsoni and H. antiochus* showed much higher heterozygosity than males (Figure 6). In *H. eleuchia* and *H. congener*, females also showed very high heterozygosity on Chr14, and in *H. hewitsoni* and *H. sapho,* on Chr9 (Figure 6). Differences between males and females were significant on these 3 chromosomes for all species of the sapho clade (Wilcoxon test, p <0.01, Figures S34-38), except

376     for *H. congener* and *H. hewitsoni* where differences could not be tested due to low

377     sample size. Heterozygosity of Chr4 for females was also significantly higher than

378     for the other chromosomes in *H. eleuchia*, *H. sapho*, *H. antiochus* (Wilcoxon test, *p*

379     <0.01, Figure S36a). The same was true for Chr14 in *H. eleuchia* and Chr9 in *H.*

380     *sapho* (Wilcoxon test, *p* <0.01, Figure S36a).

381             Females of *H. eleuchia*, *H. sapho*, *H. hewitsoni* and *H. antiochus* have

reduced mean depth on chromosome 4, whereas the mean depth on that

chromosome in males is normal (Figure 6). These differences were significant in *H.*

*eleuchia*, *H. sapho, H. antiochus* (Wilcoxon test, p ≤ 0.01; Figure S34). This pattern

was not true for Chr14 in *H. eleuchia* and *H. congener,* nor Chr9 for *H. sapho* and

*H. hewitsoni* (Figure 6 and S35). The mean depth of Chr4 was also lower than that

of all other chromosomes in females of *H. eleuchia*, *H. sapho* and *H. antiochus*

(Wilcoxon test, p ≤ 0.01; Figure S36a). However, this was not true for Chr14 and

Chr9 (Figure S36a).

**Figure 5. Scenarios of SA fusions involving either the W or Z chromosome.** Scenario of **(a)** ZA fusion and

**(b)** WA fusion for Chr4, Chr9 and Chr14 differentiating the expected pattern by sex. Yellow: Z chromosome.

The sliding window analyses on Chr4, Chr9 and Chr14 revealed that the

excess heterozygosity in females is present in most windows along the entire

chromosomes (i.e., it is not concentrated in a single continuous region; Wilcoxon

test, $p$ <0.01, Figure 7 and S37-S39). Also, mean depth values were significantly

lower for females than males in most windows on Chr4 (*H. eleuchia, H. congener,*

*H. sapho* and *H. antiochus*), Chr9 (*H. sapho*) and Chr14 (*H. eleuchia* and *H.*

*congener*) (Wilcoxon test, $p$ <0.01, Figure 7 and S37-S39). Interestingly, one

female of *H. congener* has a region of eight windows on Chr4 with very high

sequencing depth (Figure 7). There are also few windows on Chr4, Chr9 and

Chr14 where females exhibited both lower heterozygosity and mean depth than

males (Figure 7). *H. sara* and *H. leucadia* were the only species in the clade that

did not show sex-specific patterns in heterozygosity and mean depth in Chr4, Chr9

and Chr14 (Figure 6 and 7). Interestingly, the heterozygosity in females of *H.*

*congener* dropped at the end of Chr14 to values similar to those of the males

(Figure S39).

**Figure 6. Patterns of heterozygosity and depth across the genome.** Standardised proportion of heterozygosity sites and standardised mean depth per chromosome in each species (**p<0.01). Each line corresponds to one individual.

**Figure 7. Patterns of heterozygosity and depth across chromosome 4**. Proportion of heterozygosity sites and mean depth in 50Kb sliding windows in each species. Males are shown in blue and females in red, and their *n* is shown in the top right corner of each panel. Shadowed regions are those where females show very low sequencing depth and no heterozygous sites, suggesting that the female-limited haplotype diverged enough to become unmappable to the *H. sara* reference genome.

**Discussion**

We find evidence for the fusion of three autosomes to sex chromosomes in the evolution of the *sara/sapho* clade. The patterns of (i) sex-specific clustering in the phylogenies and local PCA, and (ii) high female heterozygosity seen on certain chromosomes suggest fusions of a sex chromosome with these autosomes. As females are the heterogametic sex in butterflies and show no recombination, SA fusions should generate female-specific haplotypes that do not recombine with the unfused chromosomes. Chr4 shows sex-specific clustering and high female heterozygosity throughout the chromosome in all five species of the *sapho* group, indicating that either the Z or W fused with Chr4 in the ancestor of the *sapho* group. In line with an old fusion and increasing degeneration, females show a very high proportion of heterozygous sites in most parts of the chromosome, whereas in other parts of this chromosome, females show very low sequencing depth and no heterozygous sites, indicating that the female-limited haplotype diverged enough to become unmappable to the *H. sara* reference genome. Similar, but less strong patterns of female-biased heterozygosity and sex-specific clustering in phylogenies are found on Chr9 in *H. sapho* and *H. hewitsoni* and on Chr14 in *H. eleuchia* and *H. congener.* In line with younger fusions with only weak degeneration, the sequencing depth of females matches that of the males on those chromosomes, indicating that the female-specific haplotypes still map well to the *H. sara* genome. In summary, we find three independent SA fusions in the *sapho* clade, each shared by two to five species (Figure 1). It is unknown if the autosomes are fused with the Z or W chromosome, as both types of SA fusions would lead to female-limited

haplotypes that do not recombine anymore (Chr4/9/14 in the ZA fusion and Chr W

+ Chr.4/9/14 in the WA fusion; Figure 5). Long-read sequencing data will be

required to distinguish between these possibilities. No sex chromosome fusions

have been reported previously for any *Heliconius* butterflies.

The finding of multiple sex chromosome fusions in the *sapho* group is

particularly striking, since this group is known for its high number of chromosomes

compared to all other *Heliconius* species. While most *Heliconius* species have 21

chromosomes, *H. eleuchia eleusinus* have 56-57 chromosomes, and *H. eleuchia*

and *H. congener* have 37 and 33 chromosomes, respectively (Brown et al. 1992).

This clearly suggests many chromosomal fission events, but surprisingly, our

results show that SA fusions also occurred in this group. In all species, the sex-

linked chromosomes show even patterns of increased heterozygosity and

clustering by sex across, except for the end of Chr14 in *H. congener*. This indicates

that the many chromosome fission events in the *sapho* group appear to have had

little impact on the chromosomes that were fused with the sex chromosomes.

While we do not have evidence for an adaptive role of the SA fusions in the

*sapho* group, the fact that SA fusions occurred three times and remained fixed in

multiple species, suggests that they are at least not deleterious, or that any

deleterious effect is masked. In the latter scenario, low recombination around the

SA fusions would force their sex-specific transmission and result in a permanent

heterozygosity that protects against the expression of deleterious recessive

mutations load and favours the accumulation of adaptive mutations (Jay et al.

2022). Alternatively, selection for suppressed recombination as a result of SA

fusions could lead to sex-determining genes being tightly linked to sexually

antagonistic loci on Chr4/9/14, thus coupling sex-specific beneficial alleles (Ross et

al. 2009; Pala et al. 2012; Sigeman et al. 2019). Also, the SA fusions may have

contributed to the particularly high diversification rate in this clade if they linked

together barrier loci in regions with reduced recombination (Guerrero and

Kirkpatrick 2014).

This is the first genomic study focused on the s*ara/sapho* clade. The

inclusion of multiple species and subspecies of this clade from a broad geographic

range allowed us to redefine some of the relations previously reported (Kozak et al.

2015), and to identify the effect of geography in shaping diversity. The phylogenetic

position we found for *H. antiochus* and *H. hewitsoni* contrasts with previous

amplicon based phylogenies (Beltrán et al. 2007; Kozak et al. 2015) but agrees

with a recent whole genome phylogeny based on *de novo* genome assemblies

(Cicconardi et al. 2022), suggesting that the phylogenetic relations we describe for

these two species are the most plausible. We also identified *cis* and *trans*-Andean

lineages for *H. sara* and *H. antiochus*, as well as *H. congener* and *H. eleuchia*

structured by the Andes. In addition, *H. sara* was the only species in which we

identified Andean and Amazonian lineages. The driving forces behind the SA

fusions identified here, as well as their role (if any) in the speciation or adaptation

of this clade, remain unknown. Our study adds to a growing number of examples in

butterflies that, in the future, may help to unravel the role of these rearrangements

in the evolution of Lepidoptera and other organisms.

**Data and Resource Availability**

The sequence data generated in this study have been deposited in the European

Nucleotide Archive database under accession code PRJEB60296 or elsewhere on

ENA if obtained from previous studies, as specified in Supplementary Data Table

S1 for each individual. Reference genomes for *H. sara* can be found in

https://tolqc.cog.sanger.ac.uk/durbin/jiggins/Heliconius_sara/. All records

associated to the individuals used for this study are available in the *Heliconius*

Earthcape database (https://heliconius.ecdb.io/106).

**Acknowledgments**

We also thank "Autoridad Nacional de Licencias Ambientales—ANLA" in Colombia

for granting Universidad del Rosario the collecting permit 530 and the Instituto

Chico Mendes de Conservação da Biodiversidade in Brazil for granting SISBIO

collection licence 59194-1 under which we performed our collecting activities. The

HPC Service of Universidad del Rosario (CALDAS) for computing time. This work

was funded by Universidad del Rosario (BigGrant IV-FGD005, Fondos

Concursables IV-FPD004) and NRM was funded by the Colombian Ministry of

Science and Technology (MinCiencias – doctoral scholarship 727).

**Author contributions**

NR-M, JM and CS conceived the study. NR-M, CS, CP-D, CJ, OMC, KK, CFA, and

JR collected the specimens. SMC, RD, CJ assembled the *H. sara* genome, GM

mapped the individuals to the *H. sara* genome reference. NR-M, CS and JM

analyzed the data. NR-M, CS, JM, and CP-D drafted the manuscript. All authors

read and approved the final version.

**References**

Alexander DH, Novembre J, Lange K. 2009. Fast model-based estimation of
ancestry in unrelated individuals. *Genome Res.* 19:1655–1664.

Beltrán M, Jiggins CD, Brower AVZ, Bermingham E, Mallet J. 2007. Do pollen
feeding, pupal-mating and larval gregariousness have a single origin in
Heliconius butterflies? Inferences from multilocus DNA sequence data. *Biol.*
*J. Linn. Soc.* 92:221–239.

Bressa MJ, Papeschi AG, Vítková M, Kubícková S, Fuková I, Pigozzi MI, Marec F.
2009. Sex chromosome evolution in cotton stainers of the genus Dysdercus
(Heteroptera: Pyrrhocoridae). *Cytogenet. Genome Res.* 125:292–305.

Broad I. 2019. Picard Toolkit. Available from: https://github.com/broadinstitute/picard

Brown KS. 1981. The biology of Heliconius and related genera. *Annu. Rev. Entomol.*
26:27–56.

Brown KS, Emmel TC, Eliazar PJ, Suomalainen E. 1992. Evolutionary patterns in
chromosome numbers in neotropical Lepidoptera. I. Chromosomes of the
Heliconiini (family Nymphalidae: subfamily Nymphalinae). *Hereditas*
117:109–125.

Carabajal Paladino LZ, Provazníková I, Berger M, Bass C, Aratchige NS, López SN,
Marec F, Nguyen P. 2019. Sex Chromosome Turnover in Moths of the
Diverse Superfamily Gelechioidea.Barluenga M, editor. *Genome Biol. Evol.*
11:1307–1319.

Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. 2015. Second-
generation PLINK: rising to the challenge of larger and richer datasets.
*GigaScience* 4:s13742-015-0047–0048.

Charlesworth D, Charlesworth B. 1980. Sex differences in fitness and selection for
centric fusions between sex-chromosomes and autosomes. *Genet. Res.*
35:205–214.

Chow W, Brugger K, Caccamo M, Sealy I, Torrance J, Howe K. 2016. gEVAL — a
web-based browser for evaluating genome assemblies. *Bioinformatics*
32:2508–2510.

Cicconardi F, Milanetti E, Castro ÉCP de, Mazo-Vargas A, Belleghem SMV, Ruggieri
AA, Rastas P, Hanly J, Evans E, Jiggins CD, et al. 2022. Evolutionary
dynamics of genome size and content during the adaptive radiation of
Heliconiini    butterflies.    :2022.08.12.503723.    Available    from:
https://www.biorxiv.org/content/10.1101/2022.08.12.503723v1

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker
RE, Lunter G, Marth GT, Sherry ST, et al. 2011. The variant call format and
VCFtools. *Bioinformatics* 27:2156–2158.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A,
Keane T, McCarthy SA, Davies RM, et al. 2021. Twelve years of SAMtools
and BCFtools. *GigaScience* 10:giab008.

Davey JW, Chouteau M, Barker SL, Maroja L, Baxter SW, Simpson F, Merrill RM,
Joron M, Mallet J, Dasmahapatra KK, et al. 2016. Major Improvements to the
Heliconius melpomene Genome Assembly Used to Confirm 10 Chromosome
Fusion    Events    in    6 Million    Years    of    Butterfly    Evolution.    *G3*
*GenesGenomesGenetics* 6:695–708.

DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, Philippakis AA,
del Angel G, Rivas MA, Hanna M, et al. 2011. A framework for variation
discovery and genotyping using next-generation DNA sequencing data. *Nat.*
*Genet.* 43:491–498.

Engler-Chaouat HS, Gilbert LE. 2007. De novo synthesis vs. sequestration:
negatively correlated metabolic traits and the evolution of host plant
specialization in cyanogenic butterflies. *J. Chem. Ecol.* 33:25–42.

Flores SV, Evans AL, McAllister BF. 2008. Independent Origins of New Sex-Linked
Chromosomes in the melanica and robusta Species Groups of Drosophila.
*BMC Evol. Biol.* 8:33.

Garrison E, Marth G. 2012. Haplotype-based variant detection from short-read
sequencing. Available from: http://arxiv.org/abs/1207.3907

Ghurye J, Rhie A, Walenz BP, Schmitt A, Selvaraj S, Pop M, Phillippy AM, Koren S.
2019. Integrating Hi-C links with assembly graphs for chromosome-scale
assembly. *PLOS Comput. Biol.* 15:e1007273.

Guerrero RF, Kirkpatrick M. 2014. Local adaptation and the evolution of
chromosome fusions. *Evolution* 68:2747–2756.

Hill J, Rastas P, Hornett E, Neethiraj R, Clark N, Morehouse N, La M, Celorio-
Mancera M de la P, Cols J, Dircksen H, et al. 2019. Unprecedented

reorganization of holocentric chromosomes provides insights into the enigma
of lepidopteran chromosome evolution. *Sci. Adv.* 5:eaau3648.

Hora KH, Marec F, Roessingh P, Menken SBJ. 2019. Limited intrinsic postzygotic
reproductive isolation despite chromosomal rearrangements between closely
related sympatric species of small ermine moths (Lepidoptera:
Yponomeutidae). *Biol. J. Linn. Soc.* 128:44–58.

Jay P, Tezenas E, Véber A, Giraud T. 2022. Sheltering of deleterious mutations
explains the stepwise extension of recombination suppression on sex
chromosomes and other supergenes. *PLOS Biol.* 20:e3001698.

Jiggins CD, Lamas G. 2016. The Ecology and Evolution of Heliconius Butterflies.
Oxford University Press Available from:
https://oxford.universitypressscholarship.com/view/10.1093/acprof:oso/9780
199566570.001.0001/acprof-9780199566570

Kitano J, Ross JA, Mori S, Kume M, Jones FC, Chan YF, Absher DM, Grimwood J,
Schmutz J, Myers RM, et al. 2009. A role for a neo-sex chromosome in
stickleback speciation. *Nature* 461:1079–1083.

Kozak KM, Wahlberg N, Neild AFE, Dasmahapatra KK, Mallet J, Jiggins CD. 2015.
Multilocus Species Trees Show the Recent Adaptive Radiation of the Mimetic
Heliconius Butterflies. *Syst. Biol.* 64:505–524.

Král J, Kořínková T, Krkavcová L, Musilová J, Forman M, Herrera IMÁ, Haddad CR,
Vítková M, Henriques S, Vargas JGP, et al. 2013. Evolution of karyotype, sex
chromosomes, and meiosis in mygalomorph spiders (Araneae:
Mygalomorphae). *Biol. J. Linn. Soc.* 109:377–408.

Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with
BWA-MEM. Available from: http://arxiv.org/abs/1303.3997

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,
Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The
Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25:2078–
2079.

Li H, Ralph P. 2019. Local PCA Shows How the Effect of Population Structure Differs
Along the Genome. *Genetics* 211:289–304.

Lucek K. 2018. Evolutionary Mechanisms of Varying Chromosome Numbers in the
Radiation of Erebia Butterflies. *Genes* 9:166.

Lucek K, Augustijnen H, Escudero M. 2022. A holocentric twist to chromosomal
speciation? *Trends Ecol. Evol.*:S0169534722000854.

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO Update:
Novel and Streamlined Workflows along with Broader and Deeper
Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral
Genomes. *Mol. Biol. Evol.* 38:4647–4654.

Matsumoto T, Kitano J. 2016. The intricate relationship between sexually
antagonistic selection and the evolution of sex chromosome fusions. *J. Theor.*
*Biol.* 404:97–108.

Meier JI, Marques DA, Wagner CE, Excoffier L, Seehausen O. 2018. Genomics of
Parallel Ecological Speciation in Lake Victoria Cichlids. *Mol. Biol. Evol.*
35:1489–1506.

Mongue AJ, Nguyen P, Voleníková A, Walters JR. 2017. Neo-sex Chromosomes in
the Monarch Butterfly, *Danaus plexippus*. *G3 GenesGenomesGenetics*
7:3281–3294.

Nguyen P, Sýkorová M, Šíchová J, Kůta V, Dalíková M, Čapková Frydrychová R,
Neven LG, Sahara K, Marec F. 2013. Neo-sex chromosomes and adaptive
potential in tortricid pests. *Proc. Natl. Acad. Sci. U. S. A.* 110:6931–6936.

Pala I, Naurin S, Stervander M, Hasselquist D, Bensch S, Hansson B. 2012.
Evidence of a neo-sex chromosome in birds. *Heredity* 108:264–272.

Pardo-Manuel de Villena F, Sapienza C. 2001. Female meiosis drives karyotypic
evolution in mammals. *Genetics* 159:1179–1189.

Pennell MW, Kirkpatrick M, Otto SP, Vamosi JC, Peichel CL, Valenzuela N, Kitano
J. 2015. Y Fuse? Sex Chromosome Fusions in Fishes and Reptiles. *PLOS*
*Genet.* 11:e1005237.

Picq S, Lumley L, Šíchová J, Laroche J, Pouliot E, Brunet BMT, Levesque RC,
Sperling FAH, Marec F, Cusson M. 2018. Insights into the Structure of the
Spruce Budworm ( *Choristoneura fumiferana* ) Genome, as Revealed by
Molecular Cytogenetic Analyses and a High-Density Linkage Map. *G3*
*GenesGenomesGenetics* 8:2539–2549.

Purcell S, Chang CC. 2007. PLINK: Whole genome data analysis toolset. Available
from: www.cog-genomics.org/plink/2.0/

R Core Team. 2013. R: A language and environment for statistical computing. R
Foundation for Statistical Computing. Available from: http://www.R-
project.org/

Ross JA, Urton JR, Boland J, Shapiro MD, Peichel CL. 2009. Turnover of Sex
Chromosomes in the Stickleback Fishes (Gasterosteidae). *PLOS Genet.*
5:e1000391.

Ruan J, Li H. 2020. Fast and accurate long-read assembly with wtdbg2. *Nat.*
*Methods* 17:155–158.

Schmid M, Feichtinger W, Steinlein C, Visbal García R, Fernández Badillo A. 2003.
Chromosome banding in Amphibia. XXVIII. Homomorphic XY sex
chromosomes and a derived Y-autosome translocation in Eleutherodactylus
riveroi (Anura, Leptodactylidae). *Cytogenet. Genome Res.* 101:62–73.

Šíchová J, Ohno M, Dincă V, Watanabe M, Sahara K, Marec F. 2016. Fissions,
fusions, and translocations shaped the karyotype and multiple sex
chromosome constitution of the northeast-Asian wood white butterfly,
*Leptidea amurensis. Biol. J. Linn. Soc.* 118:457–471.

Šíchová J, Voleníková A, Dincă V, Nguyen P, Vila R, Sahara K, Marec F. 2015.
Dynamic karyotype evolution and unique sex determination systems in
Leptidea wood white butterflies. *BMC Evol. Biol.* 15:89.

Sigeman H, Ponnikas S, Chauhan P, Dierickx E, Brooke M de L, Hansson B. 2019.
Repeated sex chromosome evolution in vertebrates supported by expanded
avian sex chromosomes. *Proc. R. Soc. B Biol. Sci.* 286:20192051.

Smith DAS, Gordon IJ, Traut W, Herren J, Collins S, Martins DJ, Saitoti K, Ireri P,
ffrench-Constant R. 2016. A neo-W chromosome in a tropical butterfly links
colour pattern, male-killing, and speciation. *Proc. R. Soc. B Biol. Sci.*
283:20160821.

Stamatakis A, Hoover P, Rougemont J. 2008. A Rapid Bootstrap Algorithm for the
RAxML Web Servers.Renner S, editor. *Syst. Biol.* 57:758–771.

Suomalainen E, Cook LM, Turner JRG. 1973. Achiasmatic oogenesis in the
Heliconiine butterflies. *Hereditas* 74:302–304.

de Vos JM, Augustijnen H, Bätscher L, Lucek K. 2020. Speciation through
chromosomal fusion and fission in Lepidoptera. *Phylosophical Trans. B*:11.

White MJD. 1977. Animal Cytology and Evolution. 3rd ed. Available from:
https://www.cambridge.org/co/academic/subjects/life-sciences/cell-biology-
and-developmental-biology/animal-cytology-and-evolution-3rd-edition,
https://www.cambridge.org/co/academic/subjects/life-sciences/cell-biology-
and-developmental-biology

Yoshido A, Marec F, Sahara K. 2005. Resolution of sex chromosome constitution by
683            genomic in situ hybridization and fluorescence in situ hybridization with
684            (TTAGG) n telomeric probe in some species of Lepidoptera. *Chromosoma*
685            114:193–202.

686    Yoshido A, Sahara K, Marec F, Matsuda Y. 2011. Step-by-step evolution of neo-sex
687            chromosomes in geographical populations of wild silkmoths, Samia cynthia
688            ssp. *Heredity* 106:614–624.

689