Three sequential sex chromosome – autosome fusions in *Heliconius* butterflies

Nicol Rueda - M[1], Chris D. Jiggins[2], Carolina Pardo-Diaz[1], Gabriela Montejo-Kovacevich[2], W. Owen McMillan[3], Shane McCarthy[4], Jonathan Ready[5,6], Krzysztof M. Kozak[3,7], Carlos F. Arias[3,8], Richard Durbin[2,4], Joana I. Meier[2,4¶], Camilo Salazar[1*¶]

¶ These authors contributed equally to this work.

[1]Biology Program, Faculty of Natural Sciences, Universidad del Rosario, Bogotá, Colombia. 111221

[2]Department of Zoology, University of Cambridge, Cambridge, CB2 3EJ, United Kingdom

[3]Smithsonian Tropical Research Institute, Panama

[4]Tree of Life Programme, Wellcome Sanger Institute, United Kingdom

[5]Centre for Advanced Studies of Biodiversity - CEABIO, Guamá Science Park, Lot 11, Avenida Perimetral da Ciência km 01 - Guamá, Belém - PA, 66075-750, Brazil

[6]Institute for Biological Sciences, Federal University of Pará - UFPA, R. Augusto Corrêa, 1 - Guamá, Belém - PA, 66075-110, Brazil

[7]Museum of Vertebrate Zoology, UC Berkeley, USA

[8]Data Science Lab, Office of the Chief Information Officer, Smithsonian Institution, Washington, DC, United States

*camilo.salazar@urosario.edu.co (CS)

Abstract

Sex chromosomes are evolutionarily labile in many animals, and fusion with an autosome is one of the mechanisms by which they can evolve. Sex chromosome-autosome (SA) fusions can reduce sexual conflict and promote adaptation and reproductive isolation among species. Recently, advances in genomics have fuelled the discovery of SA fusions in multiple vertebrates and in some invertebrates such as Lepidoptera. Here, we discovered multiple SA fusions in the *sara/sapho* clade of the classical adaptive radiation of *Heliconius* butterflies. While the vast majority of *Heliconius* species have 21 chromosomes, species of the particularly rapidly diversifying *sapho* clade have up to 60 chromosomes. We found a sex chromosome fusion with chromosome 4 at the base of the *sapho* clade and two additional sex chromosome fusions, each shared by two species. These sequential fusions between autosomes and sex chromosomes make the *Heliconius sapho* clade an ideal system to study the role of neo-sex chromosomes in adaptive radiations and the degeneration of sex chromosomes over time. Our study adds to a small but growing number of examples in butterflies with sex chromosome fusions that will help to unravel the importance of such rearrangements in the evolution of Lepidoptera and eukaryotes in general.

Author Summary

Chromosome number and structure are fundamental characteristics of a species. However, chromosomal rearrangements can occur spontaneously within a species, which can have significant consequences for recombination, segregation, speciation, and adaptation. Despite the importance of chromosomal rearrangements in evolution, we still have limited knowledge of the drivers and consequences of chromosomal fusions and fissions in natural populations, especially in adaptive radiations such as *Heliconius* butterflies. For the first time, we document chromosomal fusions between sex chromosomes and autosomes in *Heliconius*, particularly in species from the *sara/sapho* clade. Our research provides evidence for sex-autosome fusions involving autosomes 4, 9, and 14. All of these fusions seem to be associated with speciation events in this clade, with the sex-autosome 4 fusion being the oldest one. Although we do not yet understand the role or evolutionary consequences of these fusions, our study shows that chromosomal structure can evolve rapidly within a clade and generate chromosomal diversity. Overall, our findings contribute to a better understanding of chromosome evolution in highly diverse species.

Introduction

Sex chromosome-autosome (SA) fusions contribute to the turnover of sex-determination loci or the evolution of neo-sex chromosomes (1–3), but it remains unclear what promotes them. Sexually antagonistic selection, the sheltering of deleterious mutations, direct selection, genetic drift or meiotic drive have all been suggested as possible drivers of SA fusions (4–7). Sexually antagonistic selection is thought to favour the fusion of sex chromosomes with autosomes harbouring genes under sexually antagonistic selection (8). There is limited evidence for this theory e.g. in sticklebacks (1) and warblers (9). An alternative hypothesis is deleterious mutation sheltering, where SA fusions are favoured because they prevent the expression of recessive deleterious alleles in the heterogametic sex (4). SA fusions can also become fixed due to meiotic drive (or holocentric drive in holocentric organisms (10)), such as female meiotic drive elements on W/X-A fusions that preferentially end up in the egg instead of the polar bodies (11).

As with other chromosomal rearrangements, SA fusions can reduce recombination and potentially strengthen reproductive isolation if they bring together barrier loci into regions with reduced recombination (12,13). For instance, in the Japanese threespine stickleback *Gasterosteus aculeatus,* an SA fusion resulted in a neo-X chromosome that linked loci underlying behavioural isolation traits and hybrid sterility (5). SA fusions may also facilitate adaptation, such as the SA fusion in *Cydia pomonella* (Tortricidae), which apparently linked two insecticide-resistance genes and genes involved in detoxifying plant metabolites (14).

Genomic studies have revealed that SA fusions have occurred multiple times

across vertebrates (7,15,16), and in some invertebrates such as mygalomorph spiders

(17), *Drosophila* (18) and true bugs of the genus *Dysdercus* (19). In Lepidoptera,

examples of SA fusions include *Danaus* (20,21) and *Leptidea* butterflies (3,22), among

others (14,23–25). Compared to taxa with a single centromere per chromosome, the

holocentric chromosomes of Lepidoptera may facilitate fusions, as they are less likely to

cause segregation problems during cytokinesis and thus reduce hybrid fitness (26).

Nonetheless, butterflies and moths have remarkably constrained chromosome evolution

(27), with most species having a ZW or Z0 sex determination system and a haploid

chromosome number ranging between 28 and 32 except for a few groups that have

experienced extensive fission and fusion events (28). One of such cases are the

Heliconius butterflies, which mostly have 21 chromosomes due to ancestral fusions (29).

Only a few species in the genus differ from this ancestral chromosome number, especially

species in the *sara/sapho* clade, with some having up to 60 chromosomes (30). This clade

comprises 12 species (31), and differs from other *Heliconius* by showing an inability to

synthesise cyanogens, leading to reliance on toxins sequestered from larval host plants

(32,33). A subclade of seven species within the *sara/sapho* clade shows particularly high

diversification rates (34) and high number of chromosomes (30).

We generated whole genome resequencing data for 114 individuals from these

seven species in the *sara/sapho* clade to: (i) investigate genome-wide phylogenetic

relationships, (ii) determine the degree of genomic differentiation between species and

subspecies, and (iii) explore the impact of chromosomal rearrangements. Despite a high

number of chromosomes in this clade, suggesting many chromosomal fissions, we find

at least three SA fusions across five species, making them a prime study system for the

evolution of neo-sex chromosomes.

Results

Genome assembly

The final assembled genome was 349 Mb (with 46,300 bp of Ns) in 399 scaffolds, with

scaffold size increasing from 2,007 bp to 22.4 Mb, and with a N50 of 17.8 Mb which is

better than the currently best *Heliconius* reference genomes (*H. melpomene*: 14.3 Mb,

H. erato demophoon: 10.7 Mb). Hi-C assigned 36 scaffolds to 20 chromosomes and the

Z chromosome, whilst the remaining 363 scaffolds were unassigned. The BUSCO

completeness is 98.4%, of which 98.2% are single-copy BUSCOs.

Whole-genome resequencing dataset

A total of 114 individuals were successfully sequenced and 112 were included in

downstream analyses. The average mapping percentage to the *H. sara* genome was

95.56% (range: 77.38% - 99.14%; S1 Table). Two individuals with a particularly high

proportion of missing data (19.5% and 39.5% respectively) and low mean coverage

(7.4X and 9.1X, respectively) were excluded from further analyses (S1 Table and S1

Fig). In contrast, *H. sara* and *H. leucadia* had ≤10% missing data per individual, much

less than any other species (S1 Fig) and likely explained by close similarity to the

reference genome. The mean coverage of all other samples ranged between 10X and

27X (S1 Table and S1 Fig).

Phylogenetic relationships

The ML phylogenetic tree obtained with 183,282,470 sites separated individuals into

groups consistent with both PCA and ADMIXTURE analyses, and revealed two main

clades: (i) the *sara* clade and (ii) the *sapho* clade (Fig 1). The *sara* clade is composed of

two species, namely *H. sara* and *H. leucadia*, where *H. sara* is subdivided into an

Andean subgroup (*H. s. magdalena, H. s. sprucei,* and *H. s. elektra*) and an Amazonian

subgroup (*H. s. sara*). The *sapho* clade is split into two well-resolved lineages (*H.*

antiochus and a clade composed of two monophyletic groups: *H. eleuchia/H. congener*

and *H. sapho/H. hewitsoni*). In this case, *H. antiochus* appeared as a monophyletic

group split into an Andean clade (*H. a. aranea* and *H. a. araneides*), and an Amazonian

clade (*H. a. antiochus* and *H. a. salvini*; Fig 1). *H. antiochus* nested into the *sapho* clade

whereas *H. hewitsoni* was found as sister to *H. sapho* thus resolving the previously

undetermined position of these species **(34)**.

We found strong phylogenetic incongruence across chromosomes. For instance,

the whole-genome topology was only recovered in eight chromosomes (Chr1, Chr2,

Chr10, Chr11, Chr13, Chr15, Chr18, and Chr19; Fig 2A and S2-S22), while nine

chromosomes showed *H. congener* varying its position in the phylogeny and appearing

as sister either to *H. e. eleuchia* or to a clade composed of *H. e. eleusinus + H. e.*

primularis (Chr3, Chr4, Chr6, Chr8, Chr9, Chr12, Chr17, Chr20, and Chr21; Fig 2A and

S2-S22). Similarly, the position of *H. hewitsoni* in relation to *H. sapho* varied in eight

chromosomes (Chr5, Chr6, Chr7, Chr14, Chr16, Chr17, Chr20, Chr21; Fig 2A and S2-

S22).

Fig 1. Phylogeny and distribution of the *sara/sapho* clade. Genome-wide ML phylogeny, highlighting the position
of the *Sara* and the *Sapho* clade, with bootstrap support values on the branches. The distribution of all 17 subspecies
in the clade from (35) is shown, where dots represent sampling locations for the present study. Each of the 17
subspecies is represented by a unique colour, that is the same in the phylogeny and the distribution. The chromosome
number of each species is shown in gray rectangles. These values were taken from (36).

Interestingly, we found sex-specific clustering on Chr4, Chr9 and Chr14 (Fig 2B).

In Chr4, all species in the *sapho* clade showed females and males clustering separately,

while males of *H. congener* and *H. eleuchia* formed a clade apart from their conspecific

females (Fig 2B). In the case of *H. antiochus*, *H. sapho*, *H. hewitsoni*, females do not

group by sex among themselves because, like *H. congener* and *H. eleuchia*, they are

heterozygous for sex-specific haplotypes. However, in these species, the non-female

specific copy of Chr4 is highly divergent between them. This sex clustering pattern was

also observed on Chr9 for *H. sapho* and *H. hewitsoni* (Fig 2B). In addition, males and

females of *H. eleuchia* and *H. congener* also formed a clade in Chr14 (Fig 2B)**.**

Population structure and shared ancestry

The PCA grouped samples into three main groups: (i) *H. sara and leucadia* (hereafter

sara clade), (ii) *H. antiochus*, and (iii) *H. eleuchia, H. congener, H. sapho* and *H.*

hewitsoni (hereafter *sapho* clade). The first two principal components explained 60%

(PC1) and 10% (PC2) of the total variance. PC1 separated the *sara* clade and *H.*

charithonia (outgroup) from *H. antiochus* and the *sapho* clade, whereas PC2 separated

H. antiochus from the *sapho* clade. *H. sapho* was closer to *H. hewitsoni,* whereas *H.*

eleuchia was closer to *H. congener* (S23 Fig)*.* Results from ADMIXTURE were largely

consistent with the PCA recovering *H. antiochus*, *H. eleuchia, H. sapho* and *H. congener*

as independent groups. We also identified the Andean subspecies of *H. sara* (*H. s.*

magdalena, H. s. sprucei, and *H. s. elektra*) and the Amazonian subspecies of *H. sara*

(*H. s. sara*) as separate groups. *H. hewitsoni* and *H. leucadia* could not be correctly

assigned as a separate group due to the low number of samples. ADMIXTURE analysis

did not reveal any evidence of recent introgression among species (S24 Fig).

Fig 2. ML phylogenies inferred genome-wide and per chromosome. A) Topologies found across the genome.

Purple: genome-wide topology. Green: *H. congener* within *H. eleuchia*. Orange: *H. hewitsoni* as sister to *H.*

congener + H. eleuchia. Brown: *H. hewitsoni* as sister to *H. congener + H. eleuchia + H. sapho*. Chromosomes are

shown in the bottom, with coloured triangles indicating the topology revealed by each of them. B) Topologies

showing sex-specific grouping within some species, which is indicative of SA fusions in Chr4, Chr9 and Chr14. In

these species, females are coloured in red and males in blue. These trees are not based on phased haplotypes, so

heterozygotes are encoded as ambiguous codes.

The local PCA confirmed the sex specific patterns on Chr4, Chr 9 and Chr14 (Fig

3 and S25-S29 Fig). The MDS analysis summarising the PCA results of each 100 SNP

window on Chr4 in *H. eleuchia*, *H. congener*, *H. sapho* and *H. hewitsoni* revealed two

clusters, one of windows grouping the individuals by species, and the second grouping

the individuals by sex (Fig 3A-3B and 3D-3E). In *H. antiochus*, all windows in Chr4 cluster

individuals by sex (S27 Fig). Chr14 behaves like Chr4 but only in *H. eleuchia* and *H.*

congener, where sex-grouping is observed across the entire chromosome except for the

ends, where grouping is by species (S28 Fig). Finally, Chr9 shows a pattern where PC1

of each window groups *H. sapho* and *H. hewitsoni* individuals by species and PC2 groups

them by sex (S29 Fig). Strikingly, in all species, sex-grouping windows are scattered along

Chr4, Chr9 and Chr14 (Fig 3C, 3F and S28C).

Fig 3. Local PCA in 100 SNP windows on Chr4 reveals grouping by sex across the entire chromosome. A) two-

dimensional MDS plot of the distribution of 100 SNP windows along Chr4 for the *H. congener/H. eleuchia* clade, **B)**

PCA plots for the black and grey windows in A). **C)** Actual distribution for the black and grey windows in A) along

Chr4. **D)** two-dimensional MDS plot of the distribution of 100 SNP windows along Chr4 for the *H. sapho/H.*

hewitsoni clade. **E)** PCA plots for the black and grey windows in D). **F)** Actual distribution for the black and grey

windows in D) along Chr4. Each point in A) and D) represents one window. Black: windows that cluster by sex in the local PCA. Grey: windows that cluster by species in the local PCA. Each individual in B) and E) is represented by a point with a species-specific symbol and colour according to its sex. The female symbol for *H. congener aquilionaris* is missing because there are no female specimens of that subspecies available.

Patterns of genetic differentiation

Genomic differentiation (F_{ST}) was strongest between *H. sapho* vs. *H. hewitsoni* followed by *H. congener* vs. *H. eleuchia,* and Andean vs. Amazonian *H. antiochus* (average F_{ST} = 0.33, 0.26, and 0.07 respectively; Fig 4). *H. sara* vs. *H. leucadia* were the least differentiated pair (average F_{ST} = 0.05, Fig 4). In contrast, absolute divergence (D_{XY}) was highest in *H. sara* vs. *H. leucadia* while all other species pairs exhibited a similar degree of divergence (Fig 4). We also observed elevated F_{ST} values on the Z chromosome compared to autosomes in all but one comparison (*H. eleuchia* vs. *H. congener*; Fig 4). Interestingly, Chr4 shows lower F_{ST} values and increased D_{XY} values (in all comparisons but *H. sara* vs. *H. leucadia*). Only in *H. sapho* vs. *H. hewitsoni* Chr9 showed lower F_{ST} but did not show higher D_{XY} (Fig 4). Chr14 behaves like Chr9 but only in *H. eleuchia* and *H. congener* (Fig 4). The same pattern was true when we compared F_{ST} and D_{XY} between subspecies (S30 – S33 Fig). Interestingly, F_{ST} values peaked at the end of Chr14 when comparing between subspecies of *H. congener*, in the same region that groups individuals by species and not sex in the local PCA (S30 Fig).

Fig 4. Genome-wide divergence (F_{ST}) between pairs of species in the *sara/sapho* clade. Each dot represents a chromosome, and chromosomes with evidence of SA fusions are colour coded. Analyses for pairs of subspecies are shown in S30 and S33 Fig.

Sex-specific differences in heterozygosity and coverage on multiple chromosome

The grouping by sex we observed in the local PCA and the trees exclusively of Chr4, Chr9 and Chr14 suggest possible fusions between these autosomes and the Z or W chromosome (females of *Heliconius* are ZW and males are ZZ). Because females of Lepidoptera lack crossing over, and their meiosis is achiasmatic, they do not recombine (37). This means that, if the Z chromosome is involved in the fusion (Fig 5A), only females would have unfused autosomes that would tend to accumulate mutations and/or structural variants leading to divergence from its (now fused) homologous. Alternatively, if the W

chromosome is involved in the fusion (Fig 5B), the SA fusion would be restricted to

females and the fused SA chromosome would tend to accumulate mutations and/or

structural variants leading to divergence from its unfused homologous. Both scenarios

would lead to high heterozygosity and low depth in females due to poor mapping (Fig 5).

Consistently, Chr4, Chr9 and Chr14 showed striking sex-specific differences in

heterozygosity in multiple species, supporting three sequential fusions of these

chromosomes with the Z or W chromosomes (Fig 6A and 6B). The strongest difference in

heterozygosity was observed on Chr4, where females of *H. eleuchia, H. congener, H.*

sapho, H. hewitsoni and H. antiochus showed much higher heterozygosity than males

(Fig 6B). In *H. eleuchia* and *H. congener*, females also showed very high heterozygosity

on Chr14, and in *H. hewitsoni* and *H. sapho,* on Chr9 (Fig 6B). Differences between males

and females were significant on these 3 chromosomes for all species of the sapho clade

(Wilcoxon test, p <0.01,   S34-38 Fig), except for *H. congener* and *H. hewitsoni* where

differences could not be tested due to low sample size. Heterozygosity of Chr4 for females

was also significantly higher than for the other chromosomes in *H. eleuchia*, *H. sapho*, *H.*

antiochus (Wilcoxon test, *p* <0.01, S36A Fig). The same was true for Chr14 in *H. eleuchia*

and Chr9 in *H. sapho* (Wilcoxon test, *p* <0.01, S36A Fig).

Fig 5. Scenarios of SA fusions involving either the W or Z chromosome. Scenario of **A)** ZA fusion and **B)** WA fusion for Chr4, Chr9 and Chr14 differentiating the expected pattern by sex. Yellow: Z chromosome. Pink: W chromosome. Green: autosome. Curved lines: fusions. Red dots: mutations and/or structural variants. Asterisks: chromosomes that do not recombine and may accumulate mutations and/or structural variants thus diverging from their homologous.

Females of *H. eleuchia*, *H. sapho*, *H. hewitsoni* and *H. antiochus* have reduced mean depth on chromosome 4, whereas the mean depth on that chromosome in males is normal (Fig 6B). These differences were significant in *H. eleuchia*, *H. sapho, H. antiochus* (Wilcoxon test, p ≤ 0.01; S34 Fig). However, this pattern was not true for Chr14 in *H. eleuchia* and *H. congener,* nor Chr9 for *H. sapho* and *H. hewitsoni* (Fig 6B and S35). The mean depth of Chr4 was also lower than that of all other chromosomes in females of *H. eleuchia*, *H. sapho* and *H. antiochus* (Wilcoxon test, p ≤ 0.01; S36A Fig). However, this was not true for Chr14 and Chr9 (S36A Fig).

Fig 6. Patterns of heterozygosity and depth across the genome. A) Orange squares highlight clades where SA fusions were detected, and the SA fusion is specified. **B)** Standardised proportion of heterozygosity sites and standardised mean depth per chromosome in each species (**p<0.01). Each line corresponds to one individual.

The sliding window analyses on Chr4, Chr9 and Chr14 revealed that the excess

heterozygosity in females is present in most windows along the entire chromosomes

(i.e., it is not concentrated in a single continuous region; Wilcoxon test, *p* <0.01, Fig 7

and S37-S39). Also, mean depth values were significantly lower for females than males

in most windows on Chr4 (*H. eleuchia, H. congener, H. sapho* and *H. antiochus*), Chr9

(*H. sapho*) and Chr14 (*H. eleuchia* and *H. congener*) (Wilcoxon test, *p* <0.01, Fig 7 and

S37-S39). Interestingly, one female of *H. congener* has a region of eight windows on

Chr4 with very high sequencing depth (Fig 7). There are also few windows on Chr4,

Chr9 and Chr14 where females exhibited both lower heterozygosity and mean depth

than males (Fig 7). *H. sara* and *H. leucadia* were the only species in the clade that did

not show sex-specific patterns in heterozygosity and mean depth in Chr4, Chr9 and

Chr14 (Fig 6B and 7). Interestingly, the heterozygosity in females of *H. congener*

dropped at the end of Chr14 to values similar to those of the males (S39 Fig).

Fig 7. Patterns of heterozygosity and depth across chromosome 4. Proportion of heterozygosity sites and mean

depth in 50Kb sliding windows in each species. Males are shown in blue and females in red, and their *n* is shown in

the top right corner of each panel. Shadowed regions are those where females show very low sequencing depth and

no heterozygous sites, suggesting that the female-limited haplotype diverged enough to become unmappable to the

H. sara reference genome.

Discussion

We find evidence for the fusion of three autosomes to sex chromosomes in the evolution

of the *sara/sapho* clade. The patterns of (i) sex-specific clustering in the phylogenies

and local PCA, (ii) low Fst on Chr4, Chr9 and Chr14, and (iii) high female heterozygosity

seen on these chromosomes suggest fusions of a sex chromosome with these

autosomes. As females are the heterogametic sex in butterflies and show no

recombination, SA fusions should generate female-specific haplotypes that do not

recombine with the unfused chromosomes. This results in high heterozygosity in

females and increases variation within populations and reduces variation between them,

as females share haplotypes with females from other populations, thus generating the

observed reduced F_{ST} pattern. Chr4 shows sex-specific clustering and high female

heterozygosity throughout the chromosome in all five species of the *sapho* group,

indicating that either the Z or W fused with Chr4 in the ancestor of the *sapho* group. In

line with an old fusion and increasing degeneration, females show a very high proportion

of heterozygous sites in most parts of the chromosome, whereas in other parts of this

chromosome, females show very low sequencing depth and no heterozygous sites,

indicating that the female-limited haplotype diverged enough to become difficult to map

to the *H. sara* reference genome. Similar, but less strong patterns of female-biased

heterozygosity and sex-specific clustering in phylogenies are found on Chr9 in *H. sapho*

and *H. hewitsoni* and on Chr14 in *H. eleuchia* and *H. congener.* In line with younger

fusions with only weak degeneration, the sequencing depth of females matches that of

the males on those chromosomes, indicating that the female-specific haplotypes still

map well to the *H. sara* genome. In summary, we find three independent SA fusions in

the *sapho* clade, each shared by two to five species (Fig 1). It is unknown if the

autosomes are fused with the Z or W chromosome, as both types of SA fusions would

lead to female-limited haplotypes that no longer recombine (Chr4/9/14 in the ZA fusion

and Chr W + Chr.4/9/14 in the WA fusion; Fig 5). Long-read sequencing data will be

required to distinguish between these possibilities. No SA fusions have been reported

previously for any *Heliconius* butterflies.

The finding of multiple SA fusions in the *sapho* group is particularly striking, since

this group is known for its high number of chromosomes compared to all other

Heliconius species. While most *Heliconius* species have 21 chromosomes, *H. eleuchia*

eleusinus have 56-57 chromosomes, and *H. eleuchia* and *H. congener* have 37 and 33

chromosomes, respectively (36). This clearly suggests many chromosomal fission

events, but surprisingly, our results show that SA fusions also occurred in this group. In

all species, the sex-linked chromosomes show even patterns of increased

heterozygosity and clustering by sex across, except for the end of Chr14 in *H. congener*.

This indicates that the many chromosome fission events in the *sapho* group appear to

have had little impact on the chromosomes that were fused with the sex chromosomes.

While we do not have evidence for an adaptive role of the SA fusions in the

sapho group, the fact that SA fusions occurred three times and remained fixed in

multiple species, suggests that they are at least not deleterious, or that any deleterious

effect is masked. In the latter scenario, low recombination around the SA fusions would

force their sex-specific transmission and result in a permanent heterozygosity that

protects against the expression of deleterious recessive mutations load and favours the

accumulation of adaptive mutations (4). Alternatively, selection for suppressed

recombination as a result of SA fusions could lead to sex-determining genes being

tightly linked to sexually antagonistic loci on Chr4/9/14, thus coupling sex-specific

beneficial alleles (1,7,9). Also, the SA fusions may have contributed to the particularly

high diversification rate in this clade if they linked together barrier loci in regions with

reduced recombination (12).

This is the first genomic study focused on the s*ara/sapho* clade. The inclusion of

multiple species and subspecies of this clade from a broad geographic range allowed us

to redefine some of the relations previously reported (34), and to identify the effect of

geography in shaping diversity. The phylogenetic position we found for *H. antiochus* and

H. hewitsoni contrasts with previous amplicon based phylogenies (32,34) but agrees

with a recent whole genome phylogeny based on *de novo* genome assemblies (38),

suggesting that the phylogenetic relations we describe for these two species are the

most plausible. We also identified *cis* and *trans*-Andean lineages for *H. sara* and *H.*

antiochus, as well as *H. congener* and *H. eleuchia* structured by the Andes. In addition,

H. sara was the only species in which we identified Andean and Amazonian lineages.

The driving forces behind the SA fusions identified here, as well as their role (if any) in

the speciation or adaptation of this clade, remain unknown. Our study adds to a growing

number of examples in butterflies that, in the future, may help to unravel the role of

these rearrangements in the evolution of Lepidoptera and other organisms.

Materials and Methods

Genome assembly of *Heliconius sara*

We used a female of *H. sara* (SAMEA5394385; S1 Table) to generate an

assembled genome for this species. We assembled the genome by combining PacBio,

HiC and 10X data, all generated by the Tree of Life Programme and the Wellcome

Sanger Institute (https://www.sanger.ac.uk/programme/tree-of-life/). The libraries were

produced following the manufacturer's guidelines. We sequenced four SMRT (single-

molecule real-time) cells with the PacBio Sequel II system. Dovetail Hi-C data was

sequenced on a HiSeq X Ten platform. A 10X Genomics Chromium linked-read

sequencing library was sequenced with 150-bp paired-end reads on four lanes on an

Illumina HiSeq X Ten platform.

An Initial contig assembly was generated from the PacBio CLR data using wtdbg2 v2.2

(39) . The PacBio data was then used to polish the contigs using Arrow

(https://github.com/PacificBiosciences/GenomicConsensus). The 10X Chromium linked-

reads were then mapped to the assembly using Longranger v2.2. (10X Genomics),

variants called using freebayes v1.1.0-3-g961e5f3 (40) . Next, the assembly was

polished using BCFtools consensus v1.9 (41) by applying homozygous non-reference

calls as edits. The 10X linked-reads were then used to scaffold contigs using Scaff10X

v2.3 (https://github.com/wtsi-hpag/Scaff10X). A round of manual curation was performed

on these polished scaffolds using gEVAL (42) . Lastly, Dovetail Genomics Hi-C data was

used to scaffold the assembly further using SALSA v2.2 (43), followed by another round

of manual curation with gEVAL (42). The chromosome-scale scaffolds were named by

synteny to the *Heliconius melpomene melpomene* assembly Hmel2.5 in LepBase. We

assessed the genome contiguity with gnx-tools (https://github.com/mh11/gnx-

tools/blob/master/README) and genome completeness with BUSCO v5.1.2 (44) using

the Lepidoptera gene set.

Sample collection for genome resequencing

We collected 114 individuals of *Heliconius* from 7 species and 18 subspecies in

the *sara/sapho* clade across their distribution range: 48 *H. sara*, 2 *H. leucadia*, 21 *H.*

antiochus, 13 *H. sapho*, 3 *H. hewitsoni*, 17 *H. eleuchia* and 10 *H. congener* (S1 Table).

The body of each was preserved in NaCl-saturated DMSO solution and stored at -80°C;

wings were kept for phenotype reference.

Whole-genome resequencing and genotype calling

Genomic DNA was extracted from thoracic tissue using a DNeasy Blood and

Tissue Kit (Qiagen). Library preparation and whole-genome Illumina resequencing (PE

reads) was carried out on Illumina's HiSeq X system by Novogene (Beijing, China), with

30X coverage per individual. We also downloaded two samples of *H. charithonia*

(SRR4032025 – SRR4032026) from SRA (https://www.ncbi.nlm.nih.gov/sra) to include

them as outgroups in phylogenetic analyses. The *H. sara* genome (HelSar1) was used

as a reference to map the reads of each individual using BWA mem v0.7.12 (45) with

default parameters. We then used samtools v1.12 to sort and index the alignment files

(46). PCR-duplicate reads were identified and removed using Picard tools v2.9.2 (47),

and variant calling was conducted with Haplotype Caller (GATK, v3.7.0) in BP-resolution

mode (48). Then, samples were jointly genotyped using GATK's GenotypeGVCFs (48).

We used vcftools v0.1.14 (49) and the final VCF to calculate: (i) mean depth per

individual and site, (ii) quality per site, (iii) the proportion of missing data per individual

and (iv) the proportion of missing data per site, and (v) percentage mapping per

individual. Based on these results, we kept sites with quality value (--minQ) ≥ 30 and

less than 5% missing data. We also excluded sites with a depth below 5 and mean

depth per individual more than 1.5 times the mean to exclude paralogous regions (50).

For this, we used the custom script *removeTooHighDepthSites.sh* from

((https://github.com/joanam/VictoriaRegionSuperflock/BashPipelines). We removed sites

with excess heterozygosity using the vcftools option *--hardy* and a p-value cut-off of <1e-

5. Finally, we checked our samples for contamination or other issues generating

erroneous heterozygotes using the python script *checkHetsIndvVCF.sh* by David

Marques, and modified by Jessica Rick (https://github.com/jessicarick/lates-

popgen/blob/master/scripts/).

Phylogenetic analysis

We generated a whole-genome ML tree using a vcf containing all sites as input in RAXML v8.2.9 **(51)**, with the GTRGAMMA model and 100 bootstrap replicates. We applied the same procedure to obtain ML trees for each chromosome. In all cases, *H. charitonia* was set as outgroup.

Population structure and shared ancestry

We performed a principal component analysis (PCA) to study the genetic structure of populations. We filtered out monomorphic or multiallelic sites, and sites with minor allele frequency (MAF) smaller than 0.1 with vcftools **(49)**. To reduce the linkage disequilibrium effect, we used the python script *ldPruning.sh* from (https://github.com/joanam/scripts), which removes sites with $r^2>0.2$ in windows of 50 Kbp sliding by 10 Kbp. This resulted in a vcf file with 3,685,916 high-quality SNPs sites. We conducted the PCA using Plink v2.0 with default parameters **(52,53)**.

We analysed patterns of population structure using the R package *lostruct* (54) on chromosomes revealing sex-clustering phylogenies. We performed this analysis between pairs of species (*H. eleuchia - H. congener* and *H. sapho - H. hewitsoni*), and between subspecies of *H. antiochus*. We first divided the chromosomes into non-overlapping windows of 100 SNPs and then performed a PCA for each window to identify regions that might indicate variation in the chromosome structure. We next used a two-dimension space multidimensional scaling (MDS) analysis to illustrate how dissimilar the patterns of relatedness are between each pair of windows. We then

manually selected sets of windows to calculate PCAs and test which regions do not

group samples by species. Finally, we used R (55) to visualise the PCA results and plot

the first two MDS coordinates against the midpoint of each window to visualise variation

across the genome.

The genetic ancestry of each sample was estimated using ADMIXTURE v. 1.3.0

(56). Input data were prepared using the same procedure for the genome wide PCA

analysis. We used K = 1-6 clusters and the cross-validation method to estimate the best

K value. All other parameters were set to default.

Patterns of genetic differentiation

We calculated F_{ST} and D_{XY} by pairs of sister species and by subspecies along

chromosomes in non-overlapping 50 Kbp windows. Because *H. antiochus* did not have

a sister species, we calculated these statistics between Andean and Amazonian

subspecies. Windows that contained less than 2,500 high-quality genotyped variable

sites were rejected. We used a dataset including SNPs and monomorphic sites and the

popgenWindows.py script from (https://github.com/simonhmartin/genomics_general).

Patterns of heterozygosity and mean depth by chromosome

To study chromosomes with F_{ST} and D_{XY} patterns different from the genome

average, we used the options --het and --depth of vcftools v. 0.1.14 (49), and calculated

heterozygosity and mean depth per chromosome for each individual of each species.

We also calculated these statistics in 50 Kbp non-overlapping sliding windows along the

'outlier' chromosomes identified. On these, we calculated π specifying each individual as

its own population, so π became a measure of heterozygosity. This was done with the

Python script *popgenWindows.py* from

(https://github.com/simonhmartin/genomics_general). We then averaged these values

across all individuals of the same sex and species. For depth, we first generated a file

containing the mean depth per site averaged across all individuals of the same sex and

species using the --site-mean-depth option of vcftools v. 0.1.14 (49). We then used the

R package windowscanr v. 0.1 from (https://github.com/tavareshugo/WindowScanR) to

calculate the mean of the mean depth per species, per sex, and per window. The few

individuals of *H. hewitsoni* and *H. leucadia* were not included in the sliding windows

analysis. Statistical tests were applied to assess significant differences in heterozygosity

and mean depth between sexes and between chromosomes. As the data were not

normally distributed, we performed a Wilcoxon signed rank test to compare sexes. To

assess differences between chromosomes, we applied a Kruskal-Wallis test and a *post*

hoc test (pairwise Wilcoxon test for Kruskal-Wallis).

Acknowledgments

We also thank "Autoridad Nacional de Licencias Ambientales—ANLA" in Colombia for

granting Universidad del Rosario the collecting permit 530 and the Instituto Chico

Mendes de Conservação da Biodiversidade in Brazil for granting SISBIO collection

licence 59194-1 under which we performed our collecting activities. The HPC Service of

Universidad del Rosario (CALDAS) for computing time. This work was funded by

Universidad del Rosario (BigGrant IV-FGD005, Fondos Concursables IV-FPD004) and

NRM was funded by the Colombian Ministry of Science and Technology (MinCiencias –

doctoral scholarship 727).

References

1. Ross JA, Urton JR, Boland J, Shapiro MD, Peichel CL. Turnover of Sex Chromosomes
in the Stickleback Fishes (Gasterosteidae). PLOS Genet. 2009 Feb 20;5(2):e1000391.

2. Yoshido A, Sahara K, Marec F, Matsuda Y. Step-by-step evolution of neo-sex
chromosomes in geographical populations of wild silkmoths, Samia cynthia ssp.
Heredity. 2011 Apr;106(4):614–24.

3. Šíchová J, Ohno M, Dincă V, Watanabe M, Sahara K, Marec F. Fissions, fusions, and
translocations shaped the karyotype and multiple sex chromosome constitution of the
northeast-Asian wood white butterfly, *Leptidea amurensis*. Biol J Linn Soc. 2016
Jul;118(3):457–71.

4. Jay P, Tezenas E, Véber A, Giraud T. Sheltering of deleterious mutations explains the
stepwise extension of recombination suppression on sex chromosomes and other
supergenes. PLOS Biol. 2022 Jul 19;20(7):e3001698.

5. Kitano J, Ross JA, Mori S, Kume M, Jones FC, Chan YF, et al. A role for a neo-sex
chromosome in stickleback speciation. Nature. 2009 Oct;461(7267):1079–83.

6. Pennell MW, Kirkpatrick M, Otto SP, Vamosi JC, Peichel CL, Valenzuela N, et al. Y
Fuse? Sex Chromosome Fusions in Fishes and Reptiles. PLOS Genet. 2015 May
20;11(5):e1005237.

7. Sigeman H, Ponnikas S, Chauhan P, Dierickx E, Brooke M de L, Hansson B.
Repeated sex chromosome evolution in vertebrates supported by expanded avian sex
chromosomes. Proc R Soc B Biol Sci. 2019 Dec 4;286(1916):20192051.

8. Charlesworth D, Charlesworth B. Sex differences in fitness and selection for centric
fusions between sex-chromosomes and autosomes. Genet Res. 1980 Apr;35(2):205–
14.

9. Pala I, Naurin S, Stervander M, Hasselquist D, Bensch S, Hansson B. Evidence of a
neo-sex chromosome in birds. Heredity. 2012 Mar;108(3):264–72.

10. Krátká M, Šmerda J, Lojdová K, Bureš P, Zedek F. Holocentric Chromosomes
Probably Do Not Prevent Centromere Drive in Cyperaceae. Front Plant Sci [Internet].
2021   [cited   2023   Mar   21];12.   Available   from:
https://www.frontiersin.org/articles/10.3389/fpls.2021.642661

11. Pardo-Manuel de Villena F, Sapienza C. Female meiosis drives karyotypic evolution
in mammals. Genetics. 2001 Nov;159(3):1179–89.

12. Guerrero RF, Kirkpatrick M. Local adaptation and the evolution of chromosome
fusions. Evolution. 2014 Oct;68(10):2747–56.

13. Yeaman S. Genomic rearrangements and the evolution of clusters of locally adaptive
loci. Proc Natl Acad Sci U S A. 2013 May 7;110(19):E1743-1751.

14. Nguyen P, Sýkorová M, Šíchová J, Kůta V, Dalíková M, Čapková Frydrychová R, et
al. Neo-sex chromosomes and adaptive potential in tortricid pests. Proc Natl Acad Sci
U S A. 2013 Apr 23;110(17):6931–6.

15. Matsumoto T, Kitano J. The intricate relationship between sexually antagonistic
selection and the evolution of sex chromosome fusions. J Theor Biol. 2016 Sep
7;404:97–108.

16. Schmid M, Feichtinger W, Steinlein C, Visbal García R, Fernández Badillo A.
Chromosome banding in Amphibia. XXVIII. Homomorphic XY sex chromosomes and
a derived Y-autosome translocation in Eleutherodactylus riveroi (Anura,
Leptodactylidae). Cytogenet Genome Res. 2003;101(1):62–73.

17. Král J, Kořínková T, Krkavcová L, Musilová J, Forman M, Herrera IMÁ, et al. Evolution
of karyotype, sex chromosomes, and meiosis in mygalomorph spiders (Araneae:
Mygalomorphae). Biol J Linn Soc. 2013 Jun 1;109(2):377–408.

18. Flores SV, Evans AL, McAllister BF. Independent Origins of New Sex-Linked
Chromosomes in the melanica and robusta Species Groups of Drosophila. BMC Evol
Biol. 2008 Jan 29;8:33.

19. Bressa MJ, Papeschi AG, Vítková M, Kubíčková S, Fuková I, Pigozzi MI, et al. Sex
chromosome evolution in cotton stainers of the genus Dysdercus (Heteroptera:
Pyrrhocoridae). Cytogenet Genome Res. 2009;125(4):292–305.

20. Mongue AJ, Nguyen P, Voleníková A, Walters JR. Neo-sex Chromosomes in the
Monarch Butterfly, *Danaus plexippus.* G3 GenesGenomesGenetics. 2017 Oct
1;7(10):3281–94.

21. Smith DAS, Gordon IJ, Traut W, Herren J, Collins S, Martins DJ, et al. A neo-W
chromosome in a tropical butterfly links colour pattern, male-killing, and speciation.
Proc R Soc B Biol Sci. 2016 Jul 27;283(1835):20160821.

22. Šíchová J, Voleníková A, Dincă V, Nguyen P, Vila R, Sahara K, et al. Dynamic
karyotype evolution and unique sex determination systems in Leptidea wood white
butterflies. BMC Evol Biol. 2015 Dec;15(1):89.

23. Carabajal Paladino LZ, Provazníková I, Berger M, Bass C, Aratchige NS, López SN,
et al. Sex Chromosome Turnover in Moths of the Diverse Superfamily Gelechioidea.
Barluenga M, editor. Genome Biol Evol. 2019 Apr 1;11(4):1307–19.

24. Picq S, Lumley L, Šíchová J, Laroche J, Pouliot E, Brunet BMT, et al. Insights into the
Structure of the Spruce Budworm (*Choristoneura fumiferana*) Genome, as Revealed
by Molecular Cytogenetic Analyses and a High-Density Linkage Map. G3
GenesGenomesGenetics. 2018 Aug 1;8(8):2539–49.

25. Yoshido A, Marec F, Sahara K. Resolution of sex chromosome constitution by
genomic in situ hybridization and fluorescence in situ hybridization with (TTAGG) n
telomeric probe in some species of Lepidoptera. Chromosoma. 2005 Aug;114(3):193–
202.

26. Lucek K, Augustijnen H, Escudero M. A holocentric twist to chromosomal speciation?
Trends Ecol Evol. 2022 Apr;S0169534722000854.

27. White MJD. Animal Cytology and Evolution [Internet]. 3rd ed. 1977 [cited 2022 Dec 6].
Available from: https://www.cambridge.org/co/academic/subjects/life-sciences/cell-
biology-and-developmental-biology/animal-cytology-and-evolution-3rd-edition,
https://www.cambridge.org/co/academic/subjects/life-sciences/cell-biology-and-
developmental-biology

28. Hill J, Rastas P, Hornett E, Neethiraj R, Clark N, Morehouse N, et al. Unprecedented
reorganization of holocentric chromosomes provides insights into the enigma of
lepidopteran chromosome evolution. Sci Adv. 2019 Jun 12;5:eaau3648.

29. Davey JW, Chouteau M, Barker SL, Maroja L, Baxter SW, Simpson F, et al. Major
Improvements to the Heliconius melpomene Genome Assembly Used to Confirm 10
Chromosome Fusion Events in 6 Million Years of Butterfly Evolution. G3
GenesGenomesGenetics. 2016 Mar 1;6(3):695–708.

30. Brown KS. The biology of Heliconius and related genera. Annu Rev Entomol.
1981;26(4):27–56.

31. Jiggins CD, Lamas G. The Ecology and Evolution of Heliconius Butterflies [Internet].
Oxford University Press; 2016 [cited 2022 Jul 18]. Available from:
https://oxford.universitypressscholarship.com/view/10.1093/acprof:oso/97801995665
70.001.0001/acprof-9780199566570

32. Beltrán M, Jiggins CD, Brower AVZ, Bermingham E, Mallet J. Do pollen feeding, pupal-
mating and larval gregariousness have a single origin in Heliconius butterflies?
Inferences from multilocus DNA sequence data. Biol J Linn Soc. 2007 Oct
1;92(2):221–39.

33. Engler-Chaouat HS, Gilbert LE. De novo synthesis vs. sequestration: negatively
correlated metabolic traits and the evolution of host plant specialization in cyanogenic
butterflies. J Chem Ecol. 2007 Jan;33(1):25–42.

34. Kozak KM, Wahlberg N, Neild AFE, Dasmahapatra KK, Mallet J, Jiggins CD.
Multilocus Species Trees Show the Recent Adaptive Radiation of the Mimetic
Heliconius Butterflies. Syst Biol. 2015 May 1;64(3):505–24.

35. Rueda-M N, Salgado-Roa FC, Gantiva-Q CH, Pardo-Díaz C, Salazar C.
Environmental Drivers of Diversification and Hybridization in Neotropical Butterflies.
Front Ecol Evol [Internet]. 2021 [cited 2022 Oct 31];9. Available from:
https://www.frontiersin.org/articles/10.3389/fevo.2021.750703

36. Brown KS, Emmel TC, Eliazar PJ, Suomalainen E. Evolutionary patterns in
chromosome numbers in neotropical Lepidoptera. I. Chromosomes of the Heliconiini
(family Nymphalidae: subfamily Nymphalinae). Hereditas. 1992;117(2):109–25.

37. Suomalainen E, Cook LM, Turner JRG. Achiasmatic oogenesis in the Heliconiine
butterflies. Hereditas. 1973;74(2):302–4.

38. Cicconardi F, Milanetti E, Castro ÉCP de, Mazo-Vargas A, Belleghem SMV, Ruggieri
AA, et al. Evolutionary dynamics of genome size and content during the adaptive
radiation of Heliconiini butterflies [Internet]. bioRxiv; 2022 [cited 2022 Oct 29]. p.
2022.08.12.503723. Available from:
https://www.biorxiv.org/content/10.1101/2022.08.12.503723v1

39. Ruan J, Li H. Fast and accurate long-read assembly with wtdbg2. Nat Methods. 2020
Feb;17(2):155–8.

40. Garrison E, Marth G. Haplotype-based variant detection from short-read sequencing
[Internet]. arXiv; 2012 [cited 2023 Mar 6]. Available from:
http://arxiv.org/abs/1207.3907

41. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years
of SAMtools and BCFtools. GigaScience. 2021 Feb 16;10(2):giab008.

42. Chow W, Brugger K, Caccamo M, Sealy I, Torrance J, Howe K. gEVAL — a web-
based browser for evaluating genome assemblies. Bioinformatics. 2016 Aug
15;32(16):2508–10.

43. Ghurye J, Rhie A, Walenz BP, Schmitt A, Selvaraj S, Pop M, et al. Integrating Hi-C
links with assembly graphs for chromosome-scale assembly. PLOS Comput Biol.
2019 Aug 21;15(8):e1007273.

44. Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. BUSCO Update: Novel
and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage
for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. Mol Biol Evol. 2021 Oct
1;38(10):4647–54.

45. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM [Internet]. ArXiv; 2013 [cited 2022 Jul 29]. Available from:
http://arxiv.org/abs/1303.3997

46. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009 Aug 15;25(16):2078–9.

47. Broad I. Picard Toolkit [Internet]. Broad Institute; 2019 [cited 2022 Jul 29]. Available
from: https://github.com/broadinstitute/picard

48. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, et al. A framework
for variation discovery and genotyping using next-generation DNA sequencing data.
Nat Genet. 2011 May;43(5):491–8.

49. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant
call format and VCFtools. Bioinformatics. 2011 Aug 1;27(15):2156–8.

50. Meier JI, Marques DA, Wagner CE, Excoffier L, Seehausen O. Genomics of Parallel
Ecological Speciation in Lake Victoria Cichlids. Mol Biol Evol. 2018 Jun 1;35(6):1489–
506.

51. Stamatakis A, Hoover P, Rougemont J. A Rapid Bootstrap Algorithm for the RAxML
Web Servers. Renner S, editor. Syst Biol. 2008 Oct 1;57(5):758–71.

52. Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. Second-generation
PLINK: rising to the challenge of larger and richer datasets. GigaScience. 2015 Dec
1;4(1):s13742-015-0047–8.

53. Purcell S, Chang CC. PLINK: Whole genome data analysis toolset [Internet]. 2007
[cited 2022 Jul 29]. Available from: www.cog-genomics.org/plink/2.0/

54. Li H, Ralph P. Local PCA Shows How the Effect of Population Structure Differs Along
the Genome. Genetics. 2019 Jan 1;211(1):289–304.

55. R Core Team. R: A language and environment for statistical computing. R Foundation
for Statistical Computing [Internet]. Viena, Austria; 2013. Available from: http://www.R-
project.org/

56. Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in
unrelated individuals. Genome Res. 2009 Sep;19(9):1655–64.

Supporting information captions

S1 Table. Sample information and genotyping statistics.

S1 Fig. Missing data and mean depth per individual.

S2 Fig. Maximum likelihood phylogeny of chromosome 1.

S3 Fig. Maximum likelihood phylogeny of chromosome 2.

S4 Fig. Maximum likelihood phylogeny of chromosome 3.

S5 Fig. Maximum likelihood phylogeny of chromosome 4.

S6 Fig. Maximum likelihood phylogeny of chromosome 5.

S7 Fig. Maximum likelihood phylogeny of chromosome 6.

S8 Fig. Maximum likelihood phylogeny of chromosome 7.

S9 Fig. Maximum likelihood phylogeny of chromosome 8.

S10 Fig. Maximum likelihood phylogeny of chromosome 9.

S11 Fig. Maximum likelihood phylogeny of chromosome 10.

S12 Fig. Maximum likelihood phylogeny of chromosome 11.

S13 Fig. Maximum likelihood phylogeny of chromosome 12.

S14 Fig. Maximum likelihood phylogeny of chromosome 13.

S15 Fig. Maximum likelihood phylogeny of chromosome 14.

S16 Fig. Maximum likelihood phylogeny of chromosome 15.

S17 Fig. Maximum likelihood phylogeny of chromosome 16.

S18 Fig. Maximum likelihood phylogeny of chromosome 17.

S19 Fig. Maximum likelihood phylogeny of chromosome 18.

S20 Fig. Maximum likelihood phylogeny of chromosome 19.

S21 Fig. Maximum likelihood phylogeny of chromosome 20.

S22 Fig. Maximum likelihood phylogeny of chromosome Z.

S23 Fig. Principal Component Analysis (PCA).

S24 Fig. Admixture plot.

S25 Fig. Local PCA in non-overlapping windows of 100 SNPs along Chr4 in *H.*
***eleuchia* and *H. congener*.**

699 **S37 Fig. Proportion of heterozygous sites and mean depth between sexes in sliding**
700 **windows along chromosome 4.**

701 **S38 Fig. Proportion of heterozygous sites and mean depth between sexes in sliding**
702 **windows along chromosome A) 14 and B) 9.**

703 **S39 Fig. Patterns of heterozygosity and depth across chromosome A) 14 and B) 9.**