Genomic evidence reveals three sequential W-autosome fusions in *Heliconius* butterflies

Nicol Rueda - M[1,7], Carolina Pardo-Diaz[1], Gabriela Montejo-Kovacevich[2], W. Owen McMillan[3], Krzysztof M. Kozak[3,4], Carlos F. Arias[3,5], Jonathan Ready[5,6], Shane McCarthy[7], Richard Durbin[2,7], Chris D. Jiggins[2], Joana I. Meier[2,7*], Camilo Salazar[1*+]

*These authors contributed equally to this work.

[1]Biology Program, Faculty of Natural Sciences, Universidad del Rosario, Bogotá, Colombia. 111221

[2]Department of Zoology, University of Cambridge, Cambridge, CB2 3EJ, United Kingdom

[3]Smithsonian Tropical Research Institute, Panama

[4]Museum of Vertebrate Zoology, UC Berkeley, USA

[5]Data Science Lab, Office of the Chief Information Officer, Smithsonian Institution, Washington, DC, United States

[5]Institute for Biological Sciences, Federal University of Pará - UFPA, R. Augusto Corrêa, 1 - Guamá, Belém - PA, 66075-110, Brazil

[6]Centre for Advanced Studies of Biodiversity - CEABIO, Guamá Science Park, Lot 11, Avenida Perimetral da Ciência km 01 - Guamá, Belém - PA, 66075-750, Brazil

[7]Tree of Life Programme, Wellcome Sanger Institute, United Kingdom

[+]camilo.salazar@urosario.edu.co (CS)

Abstract

Sex chromosomes are evolutionarily labile in many animals and sometimes fuse with autosomes, creating so-called neo-sex chromosomes. Fusions between sex chromosomes and autosomes have been proposed to reduce sexual conflict and to promote adaptation and reproductive isolation among species. Recently, advances in genomics have fuelled the discovery of such fusions across the tree of life. Here, we discovered multiple fusions leading to neo-sex chromosomes in the *sara/sapho* clade of the classical adaptive radiation of *Heliconius* butterflies. *Heliconius* butterflies generally have 21 chromosomes with very high synteny. However, the five species in the *sapho* subclade show large variation in chromosome number ranging from 21 to 60. We found that the W chromosome is fused with chromosome 4 in all of them. Two sister species pairs showed subsequent fusions between the W and chromosomes 9 or 14, respectively. These sequential fusions between autosomes and sex chromosomes make *Heliconius* butterflies an ideal system for studying the role of neo-sex chromosomes in adaptive radiations and the degeneration of sex chromosomes over time. Our findings emphasize the capability of WGS technology to detect neo-sex chromosomes even when sex chromosomes are not explicitly assembled.

Author Summary

Fusions between sex chromosomes and autosomes are thought to have the potential to resolve sexual conflict and enhance local adaptation or reproductive isolation between species. Here, we discovered such fusions in *Heliconius* butterflies. *Heliconius* butterflies mostly have a very stable karyotype with 21 chromosomes and high synteny across species. However, the five species in the *sapho* subclade have up to 60 chromosomes, suggesting that they have undergone many chromosomal fissions. We document that in addition to the fissions, the *sapho* subclade also showed multiple fusions between the female-specific sex chromosome (W) and autosomes. We found a W-4 fusion shared by all five species and additional W fusions with chromosomes 9 and 14 shared by two species each. Even though in many *sapho* subclade species the autosomes have undergone fissions, the chromosomes fused with the W represent the full-length autosomes, as seen in other *Heliconius* species, suggesting that fusing to the W protected them from fissions. Our study reveals the power of short-read sequencing to discover neo-sex chromosomes and showcases *Heliconius* butterflies as a promising system for studying the causes and consequences of sex chromosome evolution.

Introduction

Sex chromosome-autosome (Sex-A) fusions contribute to the evolution of neo-sex chromosomes (1,2), but it remains unclear what promotes them. Sexually antagonistic selection, direct selection, genetic drift, meiotic drive, and sheltering of deleterious mutations have all been suggested as possible drivers of Sex-A fusions (3–6). Sexually antagonistic selection is thought to favour the fusion of sex chromosomes with autosomes harbouring genes under sexually antagonistic selection (7). There is limited evidence for this hypothesis e.g. in sticklebacks (8), *Drosophila* flies (9), warblers (10) and butterflies (11). Sex-A fusions can also become fixed due to meiotic drive (including holocentric drive in holocentric organisms) (12), such as female meiotic drive elements on W/X-A fusions that preferentially end up in the egg instead of the polar bodies (13). An alternative hypothesis is deleterious mutation sheltering, which Sex-A fusions are favoured because they prevent the expression of recessive deleterious alleles in the heterogametic sex (3). As with other chromosomal rearrangements, Sex-A fusions can reduce recombination and potentially strengthen reproductive isolation (14,15). For instance, in the Japanese threespine stickleback *Gasterosteus aculeatus,* a Sex-A fusion resulted in a neo-X chromosome that linked loci underlying behavioural isolation traits and hybrid sterility (4). Sex-A fusions may also facilitate adaptation, such as the Sex-A fusion in *Cydia pomonella* (Tortricidae), which apparently linked two insecticide-resistance genes and genes involved in detoxifying plant metabolites (16).

Cytogenetic and genomic studies revealed that Sex-A fusions have occurred many

times across vertebrates (6,17,18) and invertebrates such as spiders (19,20), *Drosophila*

flies (21,22) or true bugs of the genus *Dysdercus* (23). In Lepidoptera (butterflies and

moths), examples of Sex-A fusions include *Danaus* (24,25) and *Leptidea* butterflies

(26,27), among others (16,28–31). Compared to taxa with a single centromere per

chromosome, the holocentric chromosomes of Lepidoptera may facilitate fusions, as they

are less likely to cause segregation problems during cytokinesis and thus reduce hybrid

fitness (32). Nonetheless, butterflies and moths have remarkably constrained

chromosome evolution (33), with most species having a ZW or Z0 sex determination

system and a haploid chromosome number ranging between 28 and 32, except for few

groups that have experienced extensive fission and fusion events (31,34).

Here, we focused on *Heliconius* butterflies, which have undergone 10 ancestral

fusions and thus display 20 autosomes, along with Z and W sex chromosomes with high

collinearity across species (35). Only a few species in the genus differ in this ancestral

chromosome number, especially species in the *sara/sapho* clade, with some having up to

60 chromosomes (36). The *sara/sapho* clade comprises 12 species (31) that are different

from other *Heliconius* due to their inability to synthesize cyanogens (compounds that deter

predators) thus forcing them to rely on sequestered plant toxins (37). A subclade of seven

species within the *sara/sapho* clade shows particularly high diversification rates (38) and

high number of chromosomes (36).

We assembled a reference genome of *Heliconius sara*, a species that shows the normal

karyotype of 21 chromosomes. We found that it was fully collinear with the other two

species with chromosome-level assemblies. Next, we generated whole-genome

resequencing data from 114 individuals of all seven species of the *sara/sapho* clade with

high diversification rates to completely resolve their phylogenomic relationships and

study genomic differences between the species. We discovered that the *sapho*

subclade, which shows the high number of chromosomes in some species, exhibits

fusions between the W chromosome and autosomes (W-A fusions). One W-A fusion is

shared by all five species, whereas two additional W-A fusions are shared by two

species each. The autosomes fused to the W represent the full-length chromosomes,

indicating that they have not undergone the same high rates of fissions as the unfused

autosomes. These sequential W-A fusions make the *Heliconius sapho* subclade a prime

study system for the evolution of neo-sex chromosomes.

Results

High-quality reference genome assembly for *Heliconius sara*

We assembled the genome of *Heliconius sara* using two laboratory-reared females from

a stock population originating from Panama (S1 Table). Using the first individual

(BioSample SAMEA8947140), we obtained 24 Gbp PacBio CLR data with a coverage of

50x and an N50 subread length of 16.5 kb 122 Gbp of 10X linked-read Illumina data with

a coverage of 294x. From the second individual (BioSample SAMEA8947139), we

obtained 137 Gbp Hi-C Illumina data with 111x coverage. The final assembled genome

consists of 348.8 Mbp in 384 scaffolds. The contig N50 was 8.2 Mbp and that of scaffold

N50 was 17.8 Mbp. Our genome showed the highest contiguity (S2 Table) and BUSCO

statistics (S3 Table) of some currently published *Heliconius* genomes. The BUSCO

completeness, using the Lepidoptera gene set, achieved 98.2% single-copy BUSCOs

and less duplicated, fragmented, and missing BUSCOs than in the genomes of *H. erato*

(39)*, H. melpomene* (40) and *H. charithonia* (41) (S3 Table). We assigned the largest

twenty-two scaffolds to 20 autosomes and one scaffold to the Z chromosome based on

synteny with the *Heliconius melpomene* genome. *H. sara* chromosomes are collinear

with this genome, as well as with *H. erato* and *H. charithonia* (S1 Fig). For more

information on the genome see

https://tolqc.cog.sanger.ac.uk/durbin/jiggins/Heliconius_sara/ and

https://www.ncbi.nlm.nih.gov/datasets/genome/GCA_917862395.2/.

The W chromosome did not assemble well, as is commonly seen in lepidopteran

genomes (31). We thus used whole genome resequencing data from 114 individuals

collected during this study (see below "Whole-genome resequencing dataset section") to

assign scaffolds to the W chromosome based on sequencing depth differences between

males and females. Among the 360 scaffolds not assigned to a specific chromosome,

32 exhibited a higher mean depth in females than in males in *Heliconius sara*. This

pattern suggests that these 32 scaffolds likely constitute a part of the W chromosome

(S2 Fig). Interestingly, the reads from the *sapho* subclade species (*H. antiochus*, *H.*

sapho, *H. hewitsoni*, *H. eleuchia*, and *H. congener*) did not align to these 32 scaffolds,

suggesting that W chromosome of these species is either too divergent from the *H. sara*

W chromosome (S2 Fig) or not present. The W scaffolds in *H. sara* correspond to a

single homolog in *H. charithonia* (S1C Fig).

Whole-genome resequencing dataset

A total of 114 individuals were successfully whole-genome resequenced. Our dataset

exhibits high taxonomic completeness covering all 7 species within the *sara/sapho*

subclade (*H. sara*, *H. leucadia*, *H. antiochus*, *H. sapho*, *H. hewitsoni*, *H. eleuchia*, and *H.*

congener) and 19 out of the 28 described subspecies (42) (S1 Table). The average

mapping percentage to the *H. sara* genome was 95.56% (range: 77.38% - 99.14%) (S1

Table). We observed a strong phylogenetic signal in the mapping proportion and,

consequently, in the proportion of missing data per individual (S1 Table and S3 Fig).

The average coverage was 97.7%, 97.6%, and 96.4% for *H. sara* specimens, its sister

species *H. leucadia*, and the *sapho* group, respectively. One *H. congener* and one *H.*

antiochus individual exhibited a particularly high proportion of missing data (19.5% and

39.5%, respectively) and low mean coverage (7.4X and 9.1X, respectively) (S1 Table

and S3 Fig), and were thus excluded from further analyses.

Phylogenetic analysis reveals two main subclades and uncovered incongruence across the genome

We reconstructed a Maximum Likelihood (ML) phylogenetic tree using 183,282,470 concatenated sites. This phylogeny separated individuals into two main subclades, consistent with the PCA analyses (S4 Fig): (i) *sara* and (ii) *sapho* (Fig 1). The *sara* subclade is composed of two species, namely *H. sara* and *H. leucadia*, where *H. sara* is subdivided into an Andean subgroup (*H. s. magdalena, H. s. sprucei,* and *H. s. elektra*) and an Amazonian subgroup (*H. s. sara*). The *sapho* subclade was split into two well-resolved lineages (*H. antiochus* and a subclade composed of two monophyletic groups: *H. eleuchia/H. congener* and *H. sapho/H. hewitsoni*). *H. antiochus* appeared as a monophyletic group split into an Andean group (*H. a. aranea* and *H. a. araneides*), and an Amazonian group (*H. a. antiochus* and *H. a. salvini*) (Fig 1). *H. antiochus* nested into the *sapho* subclade, whereas *H. hewitsoni* was found to be sister to *H. sapho,* thus resolving the previously undetermined position of these species (38).

To complement the concatenated phylogeny, we also reconstructed a species tree with Astral. The species phylogeny closely mirrored the genome wide phylogeny (S5A Fig). Although *H. hewitsoni* and *H. sapho* were recovered as sister species, their branch lengths (measured in coalescence units) (S5A Fig) were short. This was also the case for the subspecies of *H. eleuchia,* where *H. e. eleusinus* and *H. e. primularis* sometimes group with *H. congener* (S5A Fig). These phylogenetic discordances were also evident in the DensiTree analysis visualising 271 phylogenies together (S5 Fig).

We found a strong phylogenetic incongruence across chromosomes. The whole-

genome topology was only recovered on eight chromosomes (S6-S27 Fig), while nine

chromosomes showed *H. congener* appearing as sister either to *H. e. eleuchia* or to a

clade composed of *H. e. eleusinus + H. e. primularis* (S6-S27 Fig). Similarly, the position

of *H. hewitsoni* relative to *H. sapho* varied on eight chromosomes (S6–S27 Fig).

Interestingly, we observed sex-specific clustering on three chromosomes (S27B Fig). On

chromosome 4 (Chr4), all species in the *sapho* subclade showed females and males

forming separate clades within each species, whereas the males of *H. congener* and *H.*

eleuchia formed a shared clade and their females formed a shared clade (S27B and S9

Fig). Sex-specific clades were also observed on chromosome 9 in *H. sapho* and *H.*

hewitsoni (S27B and S14 Fig) and on chromosome 14 in *H. eleuchia* and *H. congener*

(S27B and S19 Fig)**.**

Fig 1. Phylogeny and distribution of the *sara/sapho* clade. Genome-wide ML phylogeny, highlighting the position

of the *Sara* and the *Sapho* subclades, with bootstrap support values on the branches. The distribution of all 17

subspecies in the clade from (43) is shown, where dots represent sampling locations in the present study. Each of the

17 subspecies is represented by a single colour, that is the same in the phylogeny and the distribution. The

chromosome number of each species is shown in grey rectangles (43). The species' distribution map was created

using the free and open source QGIS (44), with a base shapefile obtained from

https://www.naturalearthdata.com/downloads/10m-cultural-vectors/ and an elevation raster from

https://srtm.csi.cgiar.org/srtmdata/.

Haplotype-based phylogenetic analysis on chromosomes 4, 9 and 14

The grouping by sex we observed in the phylogenetic trees of Chr4, Chr9 and

Chr14 suggest possible fusions between these autosomes and either the Z or W

chromosome, or possibly both (females of *Heliconius* are ZW and males are ZZ) (42).

As females of Lepidoptera lack crossing over and their meiosis is achiasmatic, they do

not recombine (45). This means that if the W chromosome is involved in the fusion (Fig

2A), the Sex-A fusion would be restricted to females and the fused chromosome would

tend to accumulate mutations and/or structural variants leading to divergence from its

unfused homologous. The unfused chr4 would become a neo-Z2 chromosome in all

species of the *sapho* subclade, chr9 would become a neo-Z3 in *H. sapho,* and *H.*

hewitsoni and chr14 would become a neo-Z3 in *H. congener* and *H. eleuchia*.

Alternatively, if the Z chromosome is involved in the fusion (Fig 2B), females would

initially still have the unfused homologue (neo-W) that would start to accumulate

mutations and/or structural variants, leading to the divergence from the Z-fused

homologue. If the sex-autosome fusion was with the Z in the *sapho* subclade, chr4

would become neo-W2 in all five species, chr9 would become neo-W3 in *H. sapho* and

H. hewitsoni and chr14 would become neo-W3 in *H. congener* and *H. eleuchia*. Lastly, it

is possible that both the Z and the W fused to the autosomes, leading to neo-W and

neo-Z, reducing the number of chromosomes.

Under either fusion to the Z or the W, we predict i) site-specific genealogies

where each female has one haplotype that forms part of the same clade as the male

haplotypes, while the other haplotype forms a female-specific clade (hereafter called

female-specific haplotype clustering), ii) low genetic differentiation (F_{ST}) on Chr4, Chr9,

and Chr14, due to higher divergence between males and females within populations

and lower variation between populations if they share the same sex-autosome fusions

(see "Patterns of genetic differentiation"), iii) high proportion of sites where all females

are heterozygous due to the presence of two different haplotypes (see "Sex-specific

differences in heterozygosity and mean depth"), and iv) low sequencing depth in

females due to poor mapping of the female-specific haplotypes that have accumulated

mutations and structural variants (see "Sex-specific differences in heterozygosity and

mean depth") (Fig 2).

Fig 2. Scenarios of Sex-A fusions involving either the W or Z chromosomes. Scenario of (A) WA fusion and (B)

ZA fusion for Chr4, Chr9 and Chr14 differentiating the expected pattern by sex: (i) Female-specific haplotype

clustering, (ii) low female Fst, (iii) high female heterozygosity and (iv) low female mean depth. For more details see

main text. Yellow: Z chromosome. Pink: W chromosome. Green: autosome. Dark-grey semicircles: fusions. Red

dots: mutations and/or structural variants. Lastly, it is possible that both the W and Z chromosomes may have fused

with these autosomes, resulting in a neo-W and neo-Z sex chromosome constitution.

Consistent with our hypotheses, we identified 218,839 SNPs on Chr4, where all males

within the *sapho* subclade were homozygous, and most females were heterozygous (<1

female per species was homozygous). To study the phylogenetic relationships among

male and female haplotypes at these sites, we phased our dataset and inferred marginal

phylogenies from ancestral recombination graphs constructed using Relate. We

subsampled the dataset to every 1000th SNP with high female heterozygosity (214

SNPs). At 31% of these sites, we recovered the expected phylogeny with a female-specific

haplotype clade (Fig 3A). The sites displaying this pattern were distributed across the

entire chromosome and were not concentrated in a specific region (Fig 3B). Another set

of SNPs (53%) exhibited genealogies where female haplotypes from at least two species

within the *sapho* subclade clustered as expected for a Sex-A fusion (S28 Fig). The

remaining sites (16%) exhibited a mixed signal, similar to the unphased phylogenetic tree

obtained for the entire chromosome (S27B Fig). This is likely due to phasing errors where

the fused and unfused homologues of the females are mixed. On Chr9, we found 66,344

sites where *H. sapho* and *H. hewitsoni* exhibited the expected pattern, given a Sex-A

fusion. In this case, we found that the 23% of the 66 sampled sites displayed the expected

genealogy (S29A Fig). The remaining 77% of the SNPs showed a mixed signal as in Chr4

(Fig S27B). Finally, on Chr14 for *H. eleuchia* and *H. congener*, we identified 57,953 sites

with the described pattern. Among the 57 subsampled sites, 44% showed the expected

genealogical clustering (S30A Fig), while the remaining 56% displayed the mixed signal

described before for Chr4 and Chr9 (Fig S27B).

Fig 3. Marginal tree for one of the SNPs on Chr4. Genealogy of the SNP 1,900,983 in Chr4 showing a pattern fully consistent with a Sex-A fusion; a total of 66 subsampled SNPs across this chromosome showed the same pattern. Each vertical line represents an individual haplotype, and the haplotypes of all individuals are shown differentiating those of females (orange) from those of males (blue). Note that one allele of the females in the *sapho* subclade clustered with the alleles of males, while the other female allele formed a separate group (highlighted in orange). (B) Position of each of the 66 SNPs in Chr4 that showed genealogies consistent with a Sex-A fusion; they were not clustered in a specific region but rather distributed along the entire chromosome. The position of the SNP 1,900,983, whose genealogy shown in A, is indicated by an arrow.

Patterns of genetic differentiation

Genomic differentiation (F_{ST}) was strongest between *H. sapho* vs. *H. hewitsoni* followed

by *H. congener* vs. *H. eleuchia,* and Andean vs. Amazonian *H. antiochus* (average F_{ST} =

0.33, 0.26, and 0.07 respectively) (S31 Fig). *H. sara* vs. *H. leucadia* were the least

differentiated pair (average F_{ST} = 0.05) (S31 Fig). We also observed elevated F_{ST} values

on the Z chromosome compared to autosomes in all but one comparison (*H. eleuchia*

vs. *H. congener*) (S31 Fig). In line with predictions from sex-autosome fusions in the

sapho subclade, Chr4 showed lower than average F_{ST} values in the *sapho* subclade, but

not in *H. sara* vs. *H. leucadia* (Fig 4A and S31 Fig). This pattern is expected if the

females have a haplotype that is more similar to the females of the other population or

species compared than to their second haplotype or the males of their own population,

thus leading to high within-population diversity and low between-population differences.

Chr9 showed lower F_{ST} in *H. sapho* vs. *H. hewitsoni* Chr9 (Fig 4A and S31 Fig), and

Chr14 in *H. eleuchia* and *H. congener* (Fig 4A and S31 Fig). The same pattern was

observed when we compared F_{ST} between subspecies (S32 – S34 Fig). In line with

expectations (Fig. 2), the observed pattern of lower F_{ST} on Chr4, Chr9, and Chr14 was

exclusive to females and absent in males across all comparisons mentioned above

within the *sapho* subclade (Fig 4B). The observed differences in F_{ST} between males and

females were statistically significant for these three chromosomes (Wilcoxon test $p <$

0.01) (S35 Fig). The F_{ST} values of these chromosomes within females were also

significantly lower compared to those of the other chromosomes (Wilcoxon test $p < 0.01$)

(S35 Fig).

279
280

Fig 4. Genome-wide divergence (F_{ST}) in the *sara/sapho* clade. (A) F_{ST} between pairs of species. Each point
represents a 50Kb window, whereby the. the top 5% windows are shown in black. The numbers below correspond to
H. sara chromosomes. Subspecies comparisons are shown in S32-S34 Figs. (B) F_{ST} between pairs of species by sex.
Each circle represents a chromosome, and chromosomes with evidence of Sex-A fusions are colour coded (*
indicates outlier chromosomes, $p<0.01$). The observed reduction in F_{ST} in females is due to higher genetic diversity
within species due to the divergence between the sex chromosome-fused and unfused haplotypes in females.

Sex-specific differences in heterozygosity and mean depth

Consistent with our hypotheses (Fig 2), Chr4, Chr9, and Chr14 showed striking sex-

specific differences in the proportion of heterozygous sites and mean depth only in the

species of the *sapho* subclade, supporting three sequential fusions of these chromosomes

with the Z or W chromosomes or both (Fig 5). The strongest difference in the proportion

of heterozygous sites was observed on Chr4, where females of *H. eleuchia*, *H. congener*,

H. sapho, *H. hewitsoni,* and *H. antiochus* showed a higher proportion of heterozygous

sites than males and the other autosomal chromosomes in females (Fig 5B). Females of

H. eleuchia and *H. congener* also showed a high proportion of heterozygous sites on

Chr14 and in *H. hewitsoni* and *H. sapho* on Chr9 (Fig 5B). Differences between males

and females were significant on these three chromosomes for all species of the *sapho*

subclade (Wilcoxon test, p <0.01) (S36 Fig), except for *H. hewitsoni* where differences

could not be tested due to low sample size. The proportion of heterozygous sites of

females was also significantly higher for Chr4 than for the other chromosomes in *H.*

eleuchia, *H. sapho*, *H. antiochus* and *H. congener* (Wilcoxon test, p <0.01) (S36 Fig). The

same was true for Chr14 in *H. eleuchia* and *H. congener,* and Chr9 in *H. sapho* (Wilcoxon

test, p <0.01) (S36 Fig). A high proportion of heterozygous sites was not observed in the

females of the species *H. sara* and *H. leucadia* on any chromosome (Fig 5 and S36 Fig).

Chromosome Z exhibited lower proportion of heterozygous sites than autosomal

chromosomes in females across all species (Fig 5 and S36 Fig).

Fig 5. Genome-wide topology and patterns of heterozygosity and sequencing depth across the genome. (A) Genome-wide topology with grey squares highlighting nodes with putative Sex-A fusions. (B) The proportion of heterozygous sites vs. mean sequencing depth per chromosome, by sex and in each species. Each circle represents a chromosome, and chromosomes with evidence of Sex-A fusions are colour coded. The high heterozygosity in females is due to the presence of the fusion in only one of the haplotypes, which becomes divergent from its counterpart. The low mean depth is because the haplotype limited to females diverged enough to be difficult to map onto the reference genome of *H. sara*.

Females of *H. eleuchia*, *H. sapho*, *H. hewitsoni,* and *H. antiochus* also showed a reduced

mean depth on chromosome 4, whereas the mean depth on that chromosome in males

was normal (Fig 5B). However, these differences were only significant in *H. sapho*

(Wilcoxon test, $p \le 0.01$) (S37D Fig). This pattern was not true for Chr14 in *H. eleuchia*

and *H. congener*, nor Chr9 for *H. sapho* and *H. hewitsoni* (Figs 5). The mean depth of

Chr4 was also lower than that of all other autosomes in females of *H. sapho* (Wilcoxon

test, $p \le 0.01$) (S37D Fig). However, this was not true for Chr14 and Chr9 (S37 Fig).

The sliding window analyses on Chr4, Chr9, and Chr14 revealed that the excess

heterozygosity in females is present in most windows along the entire chromosomes

(i.e., it is not concentrated in a specific region on the chromosomes; Wilcoxon test, *p*

<0.01) (Fig 6 and S38-S40 Fig). Also, mean depth values were lower for females than

males in most windows on Chr4 for *H. eleuchia, H. congener, H. sapho* and *H.*

antiochus, Chr9 for *H. sapho* and Chr14 for *H. eleuchia* and *H. congener* (Wilcoxon test,

p <0.01) (Fig 6 and S38-S40 Fig). As expected, some peaks of high sequencing depth,

likely due to expansions of repeats or duplications were also visible in both females and

males (Fig 6). Interestingly, one female of *H. congener* has a region of eight windows on

Chr4 with very high sequencing depth (Fig 6), indicating that repeat expansions may still

be ongoing. There were also few windows on Chr4, Chr9, and Chr14 where females

exhibited both lower mean sequencing depth than males and (almost) no heterozygous

sites (highlighted with a grey rectangle) (Fig 6). In these regions, the female-specific

haplotype likely diverged too much from the reference genome to map well and thus the

heterozygosity is low in females because only one haplotype is represented in the data.

H. sara and *H. leucadia* were the only species in the clade that did not show sex-specific

patterns in heterozygosity and mean depth in Chr4, Chr9 and Chr14 (Fig 6 and S38-S40

Fig). Interestingly, the heterozygosity in females of *H. congener* and *H. sapho* dropped

at the end of Chr14 and Chr9, respectively, to values similar to those of the males (S38

Fig), indicating that the last part of these chromosomes may not be fused to a sex

chromosome in this species. However, the mean depth pattern did not change in these

regions (S38 Fig).

Fig 6. Patterns of heterozygosity and depth across Chr4. Proportion of heterozygous (Het) sites and mean sequencing depth in 50 kb sliding windows for each species. Males are shown in blue and females in orange. Grey rectangles highlight regions where females show very low sequencing depth and no heterozygous sites, suggesting that the female-limited haplotype diverged enough to become unmappable to the *H. sara* reference genome. The mean depth figures were trimmed to a maximum value of 20 in order to better visualize the differences between sexes.

Evidence for three W-autosome fusions

In order to elucidate if the W or Z chromosome or both are involved in the sex-autosome fusions, we produced Illumina Hi-C data for a female *H. congener* (0.71 Gbp) and a male *H. sapho* (0.73 Gbp). We mapped the Hi-C data of these two *sapho* subclade individuals and the *H. sara* Hi-C data produced for the reference genome to our *H. sara* genome and to a previously published *H. charithonia* genome which has the W assembled (41). In line with phylogenetic distances, the proportion of reads mapping to the *H. sara* reference genome were much higher (96%, 93% and 92%) than to the *H. charithonia* reference genome (69%, 70%, 71% for *H. sara*, *H. congener* and *H. sapho*, respectively. The mean mapping quality for all three species against both genomes exceeded a Phred quality score of 35.

We did not observe an excess of Hi-C contacts either between autosomes or between autosomes and sex chromosomes in the *H. sara* female (Fig 7A), as expected if the sex-autosome fusions were only in the *sapho* subclade. The Hi-C signal of the *H. sapho* male and the *H. congener* female showed that their genomes are split into 56 and 33 chromosomes, respectively (Fig 7B-C), consistent with findings by Brown et al (36)

and suggesting a high number of chromosome fissions. However, no fusion was

observed in the *H. sapho* male, suggesting that the Chr4 and Chr9 fusions are likely not

with the Z, but with the W (Fig 7B). In contrast, the female *H. congener* showed an

excess of contacts between Chr4 and Chr14, in line with a fusion (Fig 7C). There was

no excess of Hi-C contacts between Chr4 and Chr14 with the Z chromosome (Fig 7C

and 7D), indicating that the Chr4 and Chr14 are likely fused with the W instead. Even

though there is signal of Hi-C contact between these chromosomes and the W, the low

mapping rates of Illumina reads of the *sapho* subclade to the W of the *Heliconius sara*

genome (S2 Fig) likely explains the absence of a stronger signal.

As the Hi-C signal of chromosome fusions in the *H. congener* female represents

a mix of signals from the fused and unfused haplotypes (Fig 7C-D), we phased the Hi-C

data across the chromosomes of interest (Chr4, Chr14, W, Z) and used *chomper* (46) to

split the Hi-C data into two subsets of read pairs representing the two haplotypes (47.5%

and 52.2% of the read pairs in each subset). The excess of Hi-C contacts between Chr4

and Chr14 was completely absent on one haplotype (Fig 7E) and very strong on the

other haplotype (Fig 7F). The Hi-C contacts between Chr4 and Chr14 are unevenly

distributed (Fig 7E), suggesting not only a fusion between these chromosomes but also

the presence of other chromosomal rearrangements such as inversions or

translocations.

Fig. 7. Hi-C contact density map confirmed a high number of chromosome fissions and W-A fusions in the *sapho*

subclade. Hi-C contact heatmaps of the *H. sara* reference genome of (A) the *H. sara* female, (B) the *H. sapho* male,

and (C) the *H. congener* female. Note that the chromosomes of the *H. sara* reference genome are labelled and their

boundaries are denoted with vertical and horizontal black lines. (D) Zoom in (C) showing Hi-C contacts on Chr4,

Chr14, ChrW, and ChrZ. (E) Hi-C reads assigned to one of the two haplotypes in the *H. congener* female showing the

W-A fusion. (F) Hi-C reads assigned to the other haplotype in the *H. congener* female showing no evidence of a fusion.

Discussion

We found evidence of three W-A fusions involving Chr4, Chr9, and Chr14 in the

evolution of the *sara/sapho* clade in *Heliconius* butterflies. These autosomes seem to

have fused with the W chromosome as supported by: (i) females with one haplotype

each forming part of female-specific haplotype clade, (ii) low F_{ST} values in females, (iii)

high heterozygosity and low mean depth in females, and (iv) excess of Hi-C contacts in

the *H. congener* female but not in the *H. sapho* male. Because females are the

heterogametic sex in butterflies and show no recombination (45), the W-A fusion is

restricted to females and generates female-specific haplotypes that do not recombine

with the unfused chromosomes. This results in genealogies where one haplotype of the

females clusters with male haplotypes while the other haplotype of the females, the

fused one, forms a separate female-specific clade. Each female shows many

heterozygous sites on the ancestral Chr4 and depending on the species also on Chr9 or

Chr14, as it has inherited from the mother a fused W-4(-9/14) that never recombines

and accumulates mutations and structural variants and from the father unfused Z

chromosomes (Z1=ancestral Z, Z2=ancestral Chr4 (and Z3=ancestral Chr9 or Chr14).

This, in turn, increases variation within species and reduces variation between them, as

females share the fused haplotypes with females of other species, thus leading to

reduced $F_{ST.}$ Chr4 showed female-specific haplotype clustering and high female

heterozygosity throughout the chromosome in all five species of the *sapho* subclade,

indicating that the W chromosome fused with Chr4 in the ancestor of this group. In line

with an old fusion and increasing degeneration, females showed a high proportion of

heterozygous sites in most of the chromosome, and in some regions, the fused

haplotype had degenerated so much that mapping to the *H. sara* genome failed, leading

to low sequencing depth and the absence of heterozygous sites. Similar, but less strong

patterns of female-biased heterozygosity are found in Chr9 in *H. sapho* and *H.*

hewitsoni, and in Chr14 in *H. eleuchia* and *H. congener.* In line with younger fusions and

weak degeneration, the sequencing depth of females matches that of the males on

those chromosomes, indicating that the female-specific haplotypes still map well to the

H. sara genome.

Sex-A fusions in Lepidoptera usually involve the Z chromosome (31,47), but this

does not seem to be the case in *Heliconius*. Our *H. sapho* male (ZZ) did not have any

chromosome fusions, indicating that neither Chr4 nor Chr9 in that species is fused to the

Z, whereas Hi-C data from the female *H. congener* supports a fusion on one of the

haplotypes between Chr4 andChr14 with the W chromosome. Thus, in this species, the

unfused Chr4 and Chr14 become neo-Z2 and neo-Z3 chromosomes, respectively, and

the fused chromosome becomes the neo-W chromosome. Although the absence of a

fusion with the Z in the female *H. congener* clearly points to a W fusion, Hi-C contacts

between Chr4-Chr14 and the W are weak. This is likely due to the divergence of the W

in the *sapho* subclade from that of *H. sara*, making it difficult to map reads against this

reference (S2 Fig). Alternatively, the original W in the *sapho* subclade may have an

independent origin to that in *H. sara* or it may have been lost completely, as suggested

for other Lepidoptera (47).

The finding of multiple W-A fusions in the *sapho* subclade is particularly striking,

since this group is known for its high number of chromosomes compared to all other

Heliconius species that was also confirmed with our Hi-C data. While most species in

this genus have 21 chromosomes, *H. sapho* and *H. eleuchia eleusinus and H. e.*

primularis has 56-57 chromosomes, and *H. eleuchia eleuchia* and *H. congener* have 37

and 33 chromosomes, respectively (36), indicative of high rates of chromosomal fission

events in the group. However, even though in these species most autosomes are

broken up (Fig 7B-C), the chromosomes fused with the W are the full-length ancestral

chromosomes and thus likely did not undergo fissions. Interestingly, *H. antiochus* and *H.*

hewitsoni which also show W-A fusions have 21 chromosomes. These findings could be

explained by two alternative scenarios: a) The chromosomal fissions in *H. eleuchia*, *H.*

congener and *H. sapho* happened independently in each (sub)species after the W-A

fusions and fusing to the W protected Chr4 and Chr9/14 from fissions or b) the fissions

could be ancestral, and the W-fusions could have involved the largest chromosomes

that had not been broken up. In this second scenario, *H. hewitsoni* would have

undergone additional autosomal fusion events restoring chromosome number to 21 after

fission events. While we think the first scenario is more likely than the second one, full

genome assemblies will be required to distinguish these hypotheses.

While we cannot test for an adaptive role of the W-A fusions in the *sapho*

subclade, the fact that they occurred three times and remained fixed in multiple species

suggests they are at least not deleterious, or that any deleterious effect is masked. In

the latter scenario, low recombination around the W-A fusions would force their sex-

specific transmission and result in a permanent heterozygosity that protects against the

expression of deleterious recessive mutations load and favours the accumulation of

adaptive mutations. This is the case in inversions in *H. numata* (3,48). Other alternative

scenarios may have favoured the fixation of these fusions. First, a period of strong

genetic drift could have led to the fixation of these fusions even if they carry some

deleterious mutations. Second, positive natural selection acting on fusions is possible

(49), though it might be hard to imagine that such beneficial effects would be found on

all three chromosomes that fused to the W. Third, transmission bias such as meiotic

drive (50,51), or coincidental linkage with endosymbionts transmitted via females (e.g.

male-killing spiroplasma) (52) could explain how the W-A fusions might have fixed.

However, if chromosomal fusions occur through non-homologous recombination, meiotic

drive might counteract or facilitate the spread of the fusions as observed for *Leptidea*

butterflies (53). Fourth, the W-A fusions might have spread due to reduction of sexual

conflict if there are sexually antagonistic loci on Chr4/9/14. In Danaini butterflies, sex-

biased gene expression is consistent with this hypothesis (11). Finally, the W-A fusions

may have contributed to the particularly high diversification rate in this clade if they

linked together barrier loci in regions with reduced recombination (14).

This is the first genomic study focused on the *sara/sapho* clade. The inclusion of

multiple species and subspecies of this clade from a broad geographic range also

allowed us to redefine some of the relations previously reported (38), and to identify the

effect of geography in shaping diversity. The phylogenetic position we found for *H.*

antiochus and *H. hewitsoni* contrasts with previous amplicon based phylogenies (38,54)

but agrees with a recent whole genome phylogeny based on *de novo* genome

assemblies (55), suggesting that the phylogenetic relations we describe for these two

species are the most plausible. We also identified *cis* and *trans*-Andean lineages for *H.*

sara and *H. antiochus*, as well as *H. congener* and *H. eleuchia* structured by the Andes.

In addition, *H. sara* was the only species in which we identified Andean and Amazonian

lineages.

Further studies are needed to understand the evolutionary drivers of the W-A

fusions identified here, as well as their role (if any) in speciation or adaptation in this

clade. Our study highlights the importance of including both sexes in short-read

population WGS studies for identifying Sex-A fusions. Finally, we show what patterns to

expect if the fusions are recent enough that the previously autosomal chromosomes are

still diploid in both sexes. As W chromosomes are often not assembled in reference

genomes due to their high repeat content, genome assemblies might not necessarily

reveal W-A fusions. Our study thus demonstrates the power of short-read population

data to discover sex-A fusions, particularly for taxa where one sex is achiasmic.

Materials and Methods

Genome assembly of *Heliconius sara*

We used two laboratory-reared females from a stock population from Panama to

generate a reference genome for *H. sara* (BioSamples SAMEA8947140 and

SAMEA8947139; S1 Table). We assembled the genome by combining PacBio, 10X

data and Hi-C data, all generated by the Tree of Life Programme at the Wellcome

Sanger Institute (). The BioSample SAMEA8947140 was used to generate the PacBio

continuous long reads (CLR). Libraries were sequenced on four Single Molecule Real-

Time (SMRT) cells using the PacBio Sequel II system. The linked-reads from 10X

Genomics Chromium technology were generated with the same sample and sequenced

in four lanes on the Illumina HiSeq X Ten platform. The second BioSample

SAMEA8947139 was used to produce Dovetail Hi-C data and sequenced on a HiSeq X

Ten platform.

An initial contig assembly was generated from the PacBio (CLR) data using wtdbg2 v2.2

(56). The PacBio data was then used to polish the contigs using Arrow

(https://github.com/PacificBiosciences/GenomicConsensus). We then retained haplotig

identification with the Purge Haplotigs pipeline (57). The 10X data were mapped to this

assembly using Longranger v2.2 (10X Genomics) and variant calling was performed

using freebayes v1.1.0-3-g961e5f3 (58). Next, this first assembly was polished using

BCFtools consensus v1.9 (59) by applying homozygous non-reference calls as edits.

The 10X linked-reads were then used to scaffold contigs using Scaff10X v2.3

(https://github.com/wtsi-hpag/Scaff10X). A round of manual curation was performed on

these polished scaffolds using gEVAL (60). Lastly, Dovetail Genomics Hi-C data was

used to scaffold the assembly further using SALSA v2.2 (61), followed by another round

of manual curation with gEVAL (60). The chromosome-scale scaffolds were named by

synteny to the *Heliconius melpomene melpomene* assembly Hmel2.5 in LepBase. We

assessed the genome contiguity with gnx-tools (https://github.com/mh11/gnx-

tools/blob/master/README) and genome completeness with BUSCO v5.1.2 (62) using

the Lepidoptera gene set. To obtain synteny plots between *H. sara* vs. *H. melpomene*,

H. erato, and *H. charithonia*, we first performed pairwise alignments between these

genomes using minimap2 v. 2.24 (63). Subsequently, we plotted the minimap2 results

using custom scripts from (https://github.com/simonhmartin/asynt).

We used whole genome resequencing data from 114 individuals obtained in this

study (see sample collection for genome resequencing section) to identify the W

chromosome within the genome of *Heliconius sara*. We calculated the mean depth

across the scaffolds that were not yet assigned to a chromosome. We first generated a

file containing the mean depth per site averaged across all individuals of the same sex

and species using the --site-mean-depth option of vcftools v. 0.1.14 (64). Then, we used

the R package windowscanr v. 0.1 (https://github.com/tavareshugo/WindowScanR) to

calculate the average of the mean depth per species, per sex, and per 500 bp windows.

Scaffolds where we observed a higher mean depth in females compared to males were

assigned to the W chromosome.

Sample collection for genome resequencing

538   We collected 114 *Heliconius* individuals from 7 species and 18 subspecies in the

539   *sara/sapho* clade across their distribution range: 48 *H. sara*, 2 *H. leucadia*, 21 *H.*

540   *antiochus*, 13 *H. sapho*, 3 *H. hewitsoni*, 17 *H. eleuchia* and 10 *H. congener* (S1 Table).

541   The body of each individual was preserved in NaCl-saturated DMSO solution and stored

542   at -80°C; wings were kept for phenotype reference.

Whole-genome resequencing and genotype calling

Genomic DNA was extracted from thoracic tissue using a DNeasy Blood and

Tissue Kit (Qiagen). Library preparation and whole-genome Illumina resequencing (PE

reads) was carried out on Illumina's HiSeq X system by Novogene (Beijing, China), with

30X coverage per individual. We also downloaded two samples of *H. charithonia*

(SRR4032025 – SRR4032026) from SRA (https://www.ncbi.nlm.nih.gov/sra) to include

them as outgroups in phylogenetic analyses. Our *H. sara* genome (HelSar1) was used

as a reference to map the reads of each individual using BWA mem v0.7.12 (65) with

default parameters. We then used samtools v1.12 to sort and index the alignment files

(66). PCR-duplicate reads were identified and removed using Picard tools v2.9.2 (67),

and variant calling was conducted with Haplotype Caller (GATK, v3.7.0) in BP-resolution

mode (68). Then, samples were jointly genotyped using GATK's GenotypeGVCFs (68).

We used vcftools v0.1.14 (64) and the final VCF to calculate: (i) mean depth per

individual and site, (ii) quality per site, (iii) the proportion of missing data per individual

and (iv) the proportion of missing data per site, and (v) percentage mapping per

individual. Based on these results, we kept sites with quality value (--minQ) ≥ 30 and

less than 5% missing data. We also excluded sites with a sequencing depth below 5 and

mean depth per individual more than 1.5 times the mean to exclude paralogous regions.

For this, we used the custom script *removeTooHighDepthSites.sh* from

(https://github.com/joanam/VictoriaRegionSuperflock/BashPipelines). We additionally

removed sites with excess heterozygosity across all individuals using the vcftools option

--hardy and a p-value cut-off of <1e-5 to remove reads from paralogous regions that are

collapsed in the reference genome.

Analysis of population structure within the *sara/sapho* clade

We performed a principal component analysis (PCA) to study the genetic

structure of populations. We filtered out monomorphic or multiallelic sites, and sites with

minor allele frequency (MAF) smaller than 0.1 with vcftools (64). To reduce the linkage

disequilibrium effect, we used the python script *ldPruning.sh* from

(https://github.com/joanam/scripts), which removes sites with r^2>0.2 in windows of 50

Kbp sliding by 10 Kbp. This resulted in a vcf file with 3,685,916 high-quality SNPs sites.

We conducted the PCA using Plink v2.0 with default parameters (69,70).

Phylogenetic relationships among *sara/sapho* clade species

We generated a whole-genome Maximum Likelihood (ML) tree using a vcf

containing all sites as input in RAXML v8.2.9 (71), with the GTRGAMMA model and 100

bootstrap replicates. We applied the same procedure to obtain ML trees for each

chromosome to study the phylogenetic incongruence across the genome. We also

inferred the species tree using the coalescent-based method ASTRAL (72). For this, we

used vfctools v0.1.14 (64) to: (i) extract two males per subspecies so Sex-A fusions

present only in females do not alter the species tree, and (ii) extract 2 kbp loci spaced at

least 10 kbp apart to ensure no linkage disequilibrium between them (73). Then,

samtools (66) was used to generate 271 multilocus blocks, each resulting from

concatenating 100 loci. Each block was converted into PHYLIP format using our custom

script *vcf2phylip.py* from (https://github.com/joanam/scripts/blob/master/vcf2phylip.py)

and used to estimate a ML tree in IQ-tree (74,75) selecting the best model with

ModelFinder and assessing node support with 1000 ultrafast bootstraps. The resulting

271 topologies were used as input in ASTRAL. We also investigated these 271

topologies with DensiTree (76) to visualize discordance.

Haplotype-based phylogenetic analysis on chromosomes 4, 9 and 14

To further investigate sex clustering in the phylogenies of autosomes 4, 9 and 14,

we phased them to infer their haplotypes. Haplotype phasing was done by combining

two methods: WhatsHap, which is a haplotype assembly technique (77), and

SHAPEIT4, a statistical phasing method (78). To implement WhatsHap, we used the

BAM file of each individual as well as the reference genome to group nearby genetic

variants into fully resolved haplotype blocks or phase sets (77). Then, we used the

WhatsHap output file to run SHAPEIT4, which further phases haplotypes based on

population-level information using default parameters (78).

Next, we inferred SNP coalescent genealogies across each chromosome using

Relate v.1.1.2 (79). Initially, the phased vcf file was transformed into haplotype format

using RelateFileFormats with the "--mode ConvertFromVcf" flag. We performed this

analysis using an effective population size of 1×10^7 individuals, as estimated for *H. erato*

(80), and a mutation rate of 2.9×10^{-9} per site per generation from *H. melpomene* (81). As

Relate further requires a genetic map, we created one using the average recombination

rate of 6 cM/Mb calculated in *H. erato* (82), and our custom script

(https://github.com/joanam/scripts/blob/master/createuniformrecmap.r). Finally, the

ancestral allele was assigned as the one more common in the outgroup *H. charithonia*.

Because any possible Sex-A fusion would produce heterozygous females and

homozygous males, we identified sites with this pattern (see hypotheses in Fig 2). We

first calculated the number of heterozygous and homozygous individuals per species

and site using the *--hardy* option in vcftools version 0.1.14 (46). Then, we used custom

scripts to find sites where each species in the *sapho* subclade met the following criteria:

(i) no males were heterozygous, (ii) at least one female was heterozygous, and (iii) not

more than one female was homozygous (allowing for one female with one allele not

called). Finally, we selected one SNP for every 1000 SNPs from these filtered sites and

visualized their genealogies using the script *TreeView.sh* (58). This subsampling

approach enabled us to examine genealogies from various sites evenly distributed along

the chromosomes.

Patterns of genetic differentiation

We calculated F_{ST} by pairs of sister species along chromosomes in non-

overlapping 50 Kbp windows. Because *H. antiochus* did not have a sister species, we

calculated these statistics between Andean and Amazonian subspecies. Windows that

contained less than 2,500 high-quality genotyped variable sites were rejected. We used

a dataset including SNPs and monomorphic sites and the *popgenWindows.py* script

from (https://github.com/simonhmartin/genomics_general). We also calculated F_{ST} per

sex following the same methodology.

Patterns of heterozygosity and mean depth by chromosome

To study chromosomes with F_{ST} patterns different from the genome average, we

used the options --het and --depth of vcftools v. 0.1.14 (62) tocalculate heterozygosity

and mean depth per chromosome for each individual of each species. We also

calculated these statistics in 50 Kbp non-overlapping sliding windows along the 'outlier'

chromosomes identified. On these, we calculated π specifying each individual as its own

population, so π became a measure of proportion of heterozygous sites. This was done

with the Python script *popgenWindows.py* from

(https://github.com/simonhmartin/genomics_general). We then averaged these values

across all individuals of the same sex and species. For sequencing depth, we first

generated a file containing the mean depth per site averaged across all individuals of

the same sex and species using the --site-mean-depth option of vcftools v. 0.1.14 (64).

We then used the R package windowscanr v. 0.1 from

(https://github.com/tavareshugo/WindowScanR) to calculate the mean of the mean

depth per species, per sex, and per window. The few individuals of *H. hewitsoni* and *H.*

leucadia were not included in the sliding windows analysis. Statistical tests were applied

to assess significant differences in heterozygosity and mean depth between sexes and

between chromosomes. As the data were not normally distributed, we performed a

Wilcoxon signed rank test to compare the sexes. To assess differences between

chromosomes, we applied a Kruskal-Wallis test and a *post hoc* test (pairwise Wilcoxon

test for Kruskal-Wallis).

651   **Identification of fusions with Hi-C data**

To investigate whether autosomes 4, 9, and 14 are fused with the W or Z chromosomes,

we constructed Hi-C libraries from the thorax of one female of *Heliconius congener*

(BioSamples SAMEA112329098; S1 Table) and a male of *Heliconius sapho*

(BioSamples SAMEA112696452, S1 Table) using the Arima2 kit (Arima Genomics, Inc).

These libraries were then sequenced on an Illumina NovaSeq S4 platform with 150 bp

paired-end reads. We used BWA mem2 v 2.2.1 (83) to map the reads against the

genomes of *Heliconius sara* (this study) and *Heliconius charithonia* (41) using default

parameters. We also mapped the Hi-C data of *H. sara* to both references to confirm the

absence of the fusion in the *sara* subclade. Next, we removed PCR duplicates,

eliminated poorly aligned reads, and filtered out reads with a mapping quality <10 from

the resulting BAM files using samtools v1.12 (66). Additionally, we generated contact

maps with pretextview and pretextSnapshot (https://github.com/wtsi-hpag/PretextView

and https://github.com/wtsi-hpag/PretextSnapshot). All these steps were done using a

custom Perl pipeline developed by Shane McCarthy at the Wellcome Sanger Institute.

To further investigate the excess of Hi-C contacts between autosomes 4, 14 and

W in the female of *H. congener*, we generated haplotype-specific Hi-C maps for these

chromosomes. First, we created a version of the *H. sara* reference genome in which

chromosomes 4, 14, and W were concatenated together to allow for phasing across

chromosomes. Next, we mapped the Hi-C reads of *H. congener* to this modified

reference following the mapping pipeline by Arima Genomics, Inc. (). Then we called

heterozygous variants using freebayes v1.3.2-dirty (58). These variants were then

normalized with bcftools v1.8 (59), decomposed with vcfallelicprimitives (84) and filtered

for coverage (>21 and <141 reads) with vcftools v. 0.1.14 (64). Next, the remaining

SNPs were phased using HAPCUT2 v1.3.3, using both the bam and the vcf files as

input (85). We used the Python script *chomper.py* from () to separate haplotype aligned

HiC reads. Finally, these haplotype-specific sets of Hi-C reads were realigned to the

original *H. sara* assembly using a custom Perl pipeline developed by Shane McCarthy at

the Wellcome Sanger Institute. Hi-C contact map was generated using pretextview and

pretextSnapshot (and https://github.com/wtsi-hpag/PretextSnapshot).

681   Acknowledgments

We thank "Autoridad Nacional de Licencias Ambientales—ANLA" in Colombia for

granting Universidad del Rosario the collecting permit 530 and the Instituto Chico

Mendes de Conservação da Biodiversidade in Brazil for granting SISBIO collection

licence 59194-1 under which we performed our collecting activities. We also thank the

HPC Service of Universidad del Rosario (CALDAS) for computing time. We thank the

core lab of the Tree of Life Programme and the sequencing centre of the Sanger

Institute for support with the HiC sequencing.

References

1. Ma WJ, Rovatsos M. Sex chromosome evolution: The remarkable diversity in the
evolutionary rates and mechanisms. J Evol Biol. 2022;35(12):1581–8.

2. Abbott JK, Nordén AK, Hansson B. Sex chromosome evolution: historical insights and
future perspectives. Proc R Soc B Biol Sci. 2017 May 3;284(1854):20162806.

3. Jay P, Tezenas E, Véber A, Giraud T. Sheltering of deleterious mutations explains the
stepwise extension of recombination suppression on sex chromosomes and other
supergenes. PLOS Biol. 2022 Jul 19;20(7):e3001698.

4. Kitano J, Ross JA, Mori S, Kume M, Jones FC, Chan YF, et al. A role for a neo-sex
chromosome in stickleback speciation. Nature. 2009 Oct;461(7267):1079–83.

5. Pennell MW, Kirkpatrick M, Otto SP, Vamosi JC, Peichel CL, Valenzuela N, et al. Y
Fuse? Sex Chromosome Fusions in Fishes and Reptiles. PLOS Genet. 2015 May
20;11(5):e1005237.

6. Sigeman H, Ponnikas S, Chauhan P, Dierickx E, Brooke M de L, Hansson B.
Repeated sex chromosome evolution in vertebrates supported by expanded avian sex
chromosomes. Proc R Soc B Biol Sci. 2019 Dec 4;286(1916):20192051.

7. Charlesworth B. Model for evolution of Y chromosomes and dosage compensation.
Proc Natl Acad Sci U S A. 1978 Nov;75(11):5618–22.

8. Ross JA, Urton JR, Boland J, Shapiro MD, Peichel CL. Turnover of Sex Chromosomes
in the Stickleback Fishes (Gasterosteidae). PLOS Genet. 2009 Feb 20;5(2):e1000391.

9. Zhou Q, Bachtrog D. Sex-specific adaptation drives early sex chromosome evolution
in Drosophila. Science. 2012 Jul 20;337(6092):341–5.

10. Pala I, Naurin S, Stervander M, Hasselquist D, Bensch S, Hansson B. Evidence of a
neo-sex chromosome in birds. Heredity. 2012 Mar;108(3):264–72.

11. Mora P, Hospodářská M, Voleníková AC, Koutecký P, Štundlová J, Dalíková M, et al.
Sex-biased gene content is associated with sex chromosome turnover in Danaini
butterflies. Mol Ecol. n/a(n/a):e17256.

12. Krátká M, Šmerda J, Lojdová K, Bureš P, Zedek F. Holocentric Chromosomes
Probably Do Not Prevent Centromere Drive in Cyperaceae. Front Plant Sci [Internet].

2021 [cited 2023 Mar 21];12. Available from: https://www.frontiersin.org/articles/10.3389/fpls.2021.642661

13. Pardo-Manuel de Villena F, Sapienza C. Female meiosis drives karyotypic evolution in mammals. Genetics. 2001 Nov;159(3):1179–89.

14. Guerrero RF, Kirkpatrick M. Local adaptation and the evolution of chromosome fusions. Evolution. 2014 Oct;68(10):2747–56.

15. Yeaman S. Genomic rearrangements and the evolution of clusters of locally adaptive loci. Proc Natl Acad Sci U S A. 2013 May 7;110(19):E1743-1751.

16. Nguyen P, Sýkorová M, Šíchová J, Kůta V, Dalíková M, Čapková Frydrychová R, et al. Neo-sex chromosomes and adaptive potential in tortricid pests. Proc Natl Acad Sci U S A. 2013 Apr 23;110(17):6931–6.

17. Matsumoto T, Kitano J. The intricate relationship between sexually antagonistic selection and the evolution of sex chromosome fusions. J Theor Biol. 2016 Sep 7;404:97–108.

18. Schmid M, Feichtinger W, Steinlein C, Visbal García R, Fernández Badillo A. Chromosome banding in Amphibia. XXVIII. Homomorphic XY sex chromosomes and a derived Y-autosome translocation in Eleutherodactylus riveroi (Anura, Leptodactylidae). Cytogenet Genome Res. 2003;101(1):62–73.

19. Král J, Kořínková T, Krkavcová L, Musilová J, Forman M, Herrera IMÁ, et al. Evolution of karyotype, sex chromosomes, and meiosis in mygalomorph spiders (Araneae: Mygalomorphae). Biol J Linn Soc. 2013 Jun 1;109(2):377–408.

20. Maddison WP, Leduc-Robert G. Multiple Origins of Sex Chromosome Fusions Correlated with Chiasma Localization in Habronattus Jumping Spiders (araneae: Salticidae). Evolution. 2013;67(8):2258–72.

21. Flores SV, Evans AL, McAllister BF. Independent Origins of New Sex-Linked Chromosomes in the melanica and robusta Species Groups of Drosophila. BMC Evol Biol. 2008 Jan 29;8:33.

22. Charlesworth B, Coyne JA, Barton NH. The Relative Rates of Evolution of Sex Chromosomes and Autosomes. Am Nat. 1987;130(1):113–46.

23. Bressa MJ, Papeschi AG, Vítková M, Kubícková S, Fuková I, Pigozzi MI, et al. Sex chromosome evolution in cotton stainers of the genus Dysdercus (Heteroptera: Pyrrhocoridae). Cytogenet Genome Res. 2009;125(4):292–305.

24. Mongue AJ, Nguyen P, Voleníková A, Walters JR. Neo-sex Chromosomes in the
Monarch Butterfly, *Danaus plexippus*. G3 GenesGenomesGenetics. 2017 Oct
1;7(10):3281–94.

25. Smith DAS, Gordon IJ, Traut W, Herren J, Collins S, Martins DJ, et al. A neo-W
chromosome in a tropical butterfly links colour pattern, male-killing, and speciation.
Proc R Soc B Biol Sci. 2016 Jul 27;283(1835):20160821.

26. Šíchová J, Voleníková A, Dincă V, Nguyen P, Vila R, Sahara K, et al. Dynamic
karyotype evolution and unique sex determination systems in Leptidea wood white
butterflies. BMC Evol Biol. 2015 Dec;15(1):89.

27. Šíchová J, Ohno M, Dincă V, Watanabe M, Sahara K, Marec F. Fissions, fusions, and
translocations shaped the karyotype and multiple sex chromosome constitution of the
northeast-Asian wood white butterfly, *Leptidea amurensis*. Biol J Linn Soc. 2016
Jul;118(3):457–71.

28. Carabajal Paladino LZ, Provazníková I, Berger M, Bass C, Aratchige NS, López SN,
et al. Sex Chromosome Turnover in Moths of the Diverse Superfamily Gelechioidea.
Barluenga M, editor. Genome Biol Evol. 2019 Apr 1;11(4):1307–19.

29. Picq S, Lumley L, Šíchová J, Laroche J, Pouliot E, Brunet BMT, et al. Insights into the
Structure of the Spruce Budworm (*Choristoneura fumiferana*) Genome, as Revealed
by Molecular Cytogenetic Analyses and a High-Density Linkage Map. G3
GenesGenomesGenetics. 2018 Aug 1;8(8):2539–49.

30. Yoshido A, Marec F, Sahara K. Resolution of sex chromosome constitution by
genomic in situ hybridization and fluorescence in situ hybridization with (TTAGG) n
telomeric probe in some species of Lepidoptera. Chromosoma. 2005 Aug;114(3):193–
202.

31. Wright CJ, Stevens L, Mackintosh A, Lawniczak M, Blaxter M. Chromosome evolution
in Lepidoptera [Internet]. Evolutionary Biology; 2023 May [cited 2023 Dec 28].
Available from: http://biorxiv.org/lookup/doi/10.1101/2023.05.12.540473

32. Lucek K, Augustijnen H, Escudero M. A holocentric twist to chromosomal speciation?
Trends Ecol Evol. 2022 Apr;S0169534722000854.

33. White MJD. Animal Cytology and Evolution [Internet]. 3rd ed. 1977 [cited 2022 Dec 6].
Available from: https://www.cambridge.org/co/academic/subjects/life-sciences/cell-
biology-and-developmental-biology/animal-cytology-and-evolution-3rd-edition,
https://www.cambridge.org/co/academic/subjects/life-sciences/cell-biology-and-
developmental-biology

34. Hill J, Rastas P, Hornett E, Neethiraj R, Clark N, Morehouse N, et al. Unprecedented
reorganization of holocentric chromosomes provides insights into the enigma of
lepidopteran chromosome evolution. Sci Adv. 2019 Jun 12;5:eaau3648.

35. Cicconardi F, Lewis JJ, Martin SH, Reed RD, Danko CG, Montgomery SH.
Chromosome Fusion Affects Genetic Diversity and Evolutionary Turnover of
Functional Loci but Consistently Depends on Chromosome Size. Mol Biol Evol.
2021;38(10):4449–62.

36. Brown KS, Emmel TC, Eliazar PJ, Suomalainen E. Evolutionary patterns in
chromosome numbers in neotropical Lepidoptera. I. Chromosomes of the Heliconiini
(family Nymphalidae: subfamily Nymphalinae). Hereditas. 1992;117(2):109–25.

37. de Castro, Zagrobelny M, Zurano JP, Zikan Cardoso M, Feyereisen R, Bak S.
Sequestration and biosynthesis of cyanogenic glucosides in passion vine butterflies
and consequences for the diversification of their host plants. Ecol Evol.
2019;9(9):5079–93.

38. Kozak KM, Wahlberg N, Neild AFE, Dasmahapatra KK, Mallet J, Jiggins CD.
Multilocus Species Trees Show the Recent Adaptive Radiation of the Mimetic
Heliconius Butterflies. Syst Biol. 2015 May 1;64(3):505–24.

39. Belleghem SMV, Rastas P, Papanicolaou A, Martin SH, Arias CF, Supple MA, et al.
Complex modular architecture around a simple toolkit of wing pattern genes. Nat Ecol
Evol. 2017;1(52).

40. Davey JW, Chouteau M, Barker SL, Maroja L, Baxter SW, Simpson F, et al. Major
Improvements to the Heliconius melpomene Genome Assembly Used to Confirm 10
Chromosome Fusion Events in 6 Million Years of Butterfly Evolution. G3
GenesGenomesGenetics. 2016 Mar 1;6(3):695–708.

41. Chakraborty M, Lara AG, Dang A, McCulloch KJ, Rainbow D, Carter D, et al. Sex-
linked gene traffic underlies the acquisition of sexually dimorphic UV color vision in
Heliconius butterflies [Internet]. bioRxiv; 2022 [cited 2023 Jul 26]. p.
2022.07.04.498748. Available from:
https://www.biorxiv.org/content/10.1101/2022.07.04.498748v1

42. Jiggins CD, Lamas G. The Ecology and Evolution of Heliconius Butterflies [Internet].
Oxford University Press; 2016 [cited 2022 Jul 18]. Available from:
https://oxford.universitypressscholarship.com/view/10.1093/acprof:oso/97801995665
70.001.0001/acprof-9780199566570

43. Rueda-M N, Salgado-Roa FC, Gantiva-Q CH, Pardo-Díaz C, Salazar C.
Environmental Drivers of Diversification and Hybridization in Neotropical Butterflies.

Front Ecol Evol [Internet]. 2021 [cited 2022 Oct 31];9. Available from:
https://www.frontiersin.org/articles/10.3389/fevo.2021.750703

44. QGIS Development Team. QGIS Geographic Information System. Open Source
Geospatial Found Proj. 2018;URL:http://qgis.osgeo.org.

45. Suomalainen E, Cook LM, Turner JRG. Achiasmatic oogenesis in the Heliconiine
butterflies. Hereditas. 1973;74(2):302–4.

46. Mackintosh A, Laetsch DR, Baril T, Foster RG, Dincă V, Vila R, et al. The genome
sequence of the lesser marbled fritillary, Brenthis ino, and evidence for a segregating
neo-Z chromosome. G3 GenesGenomesGenetics. 2022 Jun 1;12(6):jkac069.

47. Chen X, Wang Z, Zhang C, Hu J, Lu Y, Zhou H, et al. Unraveling the complex
evolutionary history of lepidopteran chromosomes through ancestral chromosome
reconstruction and novel chromosome nomenclature. BMC Biol. 2023 Nov
20;21(1):265.

48. Jay P, Chouteau M, Whibley A, Bastide H, Llaurens V, Parrinello H, et al. Mutation
accumulation in chromosomal inversions maintains wing pattern polymorphism in a
butterfly [Internet]. bioRxiv; 2019 [cited 2023 Oct 18]. p. 736504. Available from:
https://www.biorxiv.org/content/10.1101/736504v1

49. Mackintosh A, Vila R, Martin SH, Setter D, Lohse K. Do chromosome rearrangements
fix by genetic drift or natural selection? A test in Brenthis butterflies [Internet]. bioRxiv;
2023 [cited 2023 Aug 13]. p. 2023.06.16.545248. Available from:
https://www.biorxiv.org/content/10.1101/2023.06.16.545248v1

50. Stewart NB, Ahmed-Braimah YH, Cerne DG, McAllister BF. Female meiotic drive
preferentially segregates derived metacentric chromosomes in *Drosophila* [Internet].
Genetics; 2019 May [cited 2024 Feb 12]. Available from:
http://biorxiv.org/lookup/doi/10.1101/638684

51. White MJD. Models of Speciation. Science. 1968 Mar 8;159(3819):1065–70.

52. Martin SH, Singh KS, Gordon IJ, Omufwoko KS, Collins S, Warren IA, et al. Whole-
chromosome hitchhiking driven by a male-killing endosymbiont. PLOS Biol. 2020 Feb
27;18(2):e3000610.

53. Boman J, Wiklund C, Vila R, Backström N. Meiotic drive against chromosome fusions
in butterfly hybrids [Internet]. bioRxiv; 2024 [cited 2024 Feb 9]. p. 2024.02.02.578558.
Available from: https://www.biorxiv.org/content/10.1101/2024.02.02.578558v1

54. Beltrán M, Jiggins CD, Brower AVZ, Bermingham E, Mallet J. Do pollen feeding, pupal-
mating and larval gregariousness have a single origin in Heliconius butterflies?
Inferences from multilocus DNA sequence data. Biol J Linn Soc. 2007 Oct
1;92(2):221–39.

55. Cicconardi F, Milanetti E, Castro ÉCP de, Mazo-Vargas A, Belleghem SMV, Ruggieri
AA, et al. Evolutionary dynamics of genome size and content during the adaptive
radiation of Heliconiini butterflies [Internet]. bioRxiv; 2022 [cited 2022 Oct 30]. p.
2022.08.12.503723.   Available   from:
https://www.biorxiv.org/content/10.1101/2022.08.12.503723v1

56. Ruan J, Li H. Fast and accurate long-read assembly with wtdbg2. Nat Methods. 2020
Feb;17(2):155–8.

57. Roach MJ, Schmidt SA, Borneman AR. Purge Haplotigs: allelic contig reassignment
for third-gen diploid genome assemblies. BMC Bioinformatics. 2018 Nov 29;19(1):460.

58. Garrison E, Marth G. Haplotype-based variant detection from short-read sequencing
[Internet]. arXiv; 2012 [cited 2023 Mar 6]. Available from:
http://arxiv.org/abs/1207.3907

59. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years
of SAMtools and BCFtools. GigaScience. 2021 Feb 16;10(2):giab008.

60. Chow W, Brugger K, Caccamo M, Sealy I, Torrance J, Howe K. gEVAL — a web-
based browser for evaluating genome assemblies. Bioinformatics. 2016 Aug
15;32(16):2508–10.

61. Ghurye J, Rhie A, Walenz BP, Schmitt A, Selvaraj S, Pop M, et al. Integrating Hi-C
links with assembly graphs for chromosome-scale assembly. PLOS Comput Biol.
2019 ago;15(8):e1007273.

62. Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. BUSCO Update: Novel
and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage
for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. Mol Biol Evol. 2021 Oct
1;38(10):4647–54.

63. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 2018
Sep 15;34(18):3094–100.

64. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant
call format and VCFtools. Bioinformatics. 2011 Aug 1;27(15):2156–8.

65. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM [Internet]. ArXiv; 2013 [cited 2022 Jul 29]. Available from:
http://arxiv.org/abs/1303.3997

66. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009 Aug 15;25(16):2078–9.

67. Broad I. Picard Toolkit [Internet]. Broad Institute; 2019 [cited 2022 Jul 29]. Available
from: https://github.com/broadinstitute/picard

68. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, et al. A framework
for variation discovery and genotyping using next-generation DNA sequencing data.
Nat Genet. 2011 May;43(5):491–8.

69. Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. Second-generation
PLINK: rising to the challenge of larger and richer datasets. GigaScience. 2015 Dec
1;4(1):s13742-015-0047–8.

70. Purcell S, Chang CC. PLINK: Whole genome data analysis toolset [Internet]. 2007
[cited 2022 Jul 29]. Available from: www.cog-genomics.org/plink/2.0/

71. Stamatakis A, Hoover P, Rougemont J. A Rapid Bootstrap Algorithm for the RAxML
Web Servers. Renner S, editor. Syst Biol. 2008 Oct 1;57(5):758–71.

72. Mirarab S, Reaz R, Bayzid MdS, Zimmermann T, Swenson MS, Warnow T. ASTRAL:
genome-scale coalescent-based species tree estimation. Bioinformatics. 2014 Sep
1;30(17):i541–8.

73. Dasmahapatra KK, Walters JR, Briscoe AD, Davey JW, Whibley A, Nadeau NJ, et al.
Butterfly genome reveals promiscuous exchange of mimicry adaptations among
species. Nature. 2012 Jul 5;487(7405):94–8.

74. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder:
fast model selection for accurate phylogenetic estimates. Nat Methods. 2017
Jun;14(6):587–9.

75. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and Effective
Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol Biol Evol.
2015 Jan 1;32(1):268–74.

76. Bouckaert RR. DensiTree: making sense of sets of phylogenetic trees. Bioinformatics.
2010 May 15;26(10):1372–3.

77. Patterson M, Marschall T, Pisanti N, Van Iersel L, Stougie L, Klau GW, et al. W HATS
H AP : Weighted Haplotype Assembly for Future-Generation Sequencing Reads. J
Comput Biol. 2015 Jun;22(6):498–509.

78. Delaneau O, Zagury JF, Robinson MR, Marchini JL, Dermitzakis ET. Accurate,
scalable and integrative haplotype estimation. Nat Commun. 2019 Nov 28;10(1):5436.

79. Speidel L, Forest M, Shi S, Myers SR. A method for genome-wide genealogy
estimation for thousands of samples. Nat Genet. 2019 Sep;51(9):1321–9.

80. Van Belleghem SM, Baquero M, Papa R, Salazar C, McMillan WO, Counterman BA,
et al. Patterns of Z chromosome divergence among Heliconius species highlight the
importance of historical demography. Mol Ecol. 2018;27(19):3852–72.

81. Keightley PD, Pinharanda A, Ness RW, Simpson F, Dasmahapatra KK, Mallet J, et al.
Estimation of the Spontaneous Mutation Rate in Heliconius melpomene. Mol Biol Evol.
2015 Jan 1;32(1):239–43.

82. Tobler A, Kapan D, Flanagan NS, Gonzalez C, Peterson E, Jiggins CD, et al. First-
generation linkage map of the warningly colored butterfly Heliconius erato. Heredity.
2005 Apr;94(4):408–17.

83. Vasimuddin Md, Misra S, Li H, Aluru S. Efficient Architecture-Aware Acceleration of
BWA-MEM for Multicore Systems. In: 2019 IEEE International Parallel and Distributed
Processing Symposium (IPDPS) [Internet]. 2019 [cited 2024 Jan 20]. p. 314–24.
Available from: https://ieeexplore.ieee.org/document/8820962

84. Garrison E, Kronenberg ZN, Dawson ET, Pedersen BS, Prins P. Vcflib and tools for
processing the VCF variant call format [Internet]. bioRxiv; 2021 [cited 2024 Feb 12].
p.   2021.05.21.445151.   Available   from:
https://www.biorxiv.org/content/10.1101/2021.05.21.445151v1

85. Bansal V. HapCUT2: A Method for Phasing Genomes Using Experimental Sequence
Data. In: Peters BA, Drmanac R, editors. Haplotyping: Methods and Protocols
[Internet]. New York, NY: Springer US; 2023 [cited 2023 Nov 3]. p. 139–47. (Methods
in Molecular Biology). Available from: https://doi.org/10.1007/978-1-0716-2819-5_9

Supporting information captions

S1 Table. Sample information and genotyping statistics.

S*2* Table. Lepidoptera genome assembly statistics.

S3 Table. BUSCO results statistics.

S1 Fig. Synteny plots showing high collinearity between *Heliconius* genomes*.*

S2 Fig. Identification of the W chromosome in the genome of *H. sara*.

S3 Fig. Missing data and mean depth per individual.

S4 Fig. Principal Component Analysis (PCA), performed with 3,685,916 SNPs.

S5 Fig. Species tree based on ASTRAL multi-species coalescence and Densitree.

S6 Fig. Maximum likelihood phylogeny of chromosome 1.

S7 Fig. Maximum likelihood phylogeny of chromosome 2.

S8 Fig. Maximum likelihood phylogeny of chromosome 3.

S9 Fig. Maximum likelihood phylogeny of chromosome 4.

S10 Fig. Maximum likelihood phylogeny of chromosome 5.

S11 Fig. Maximum likelihood phylogeny of chromosome 6.

S12 Fig. Maximum likelihood phylogeny of chromosome 7.

S13 Fig. Maximum likelihood phylogeny of chromosome 8.

S14 Fig. Maximum likelihood phylogeny of chromosome 9.

S15 Fig. Maximum likelihood phylogeny of chromosome 10.

S16 Fig. Maximum likelihood phylogeny of chromosome 11.

S17 Fig. Maximum likelihood phylogeny of chromosome 12.

S18 Fig. Maximum likelihood phylogeny of chromosome 13.

S19 Fig. Maximum likelihood phylogeny of chromosome 14.

S20 Fig. Maximum likelihood phylogeny of chromosome 15.

S21 Fig. Maximum likelihood phylogeny of chromosome 16.

S22 Fig. Maximum likelihood phylogeny of chromosome 17.

S23 Fig. Maximum likelihood phylogeny of chromosome 18.

S24 Fig. Maximum likelihood phylogeny of chromosome 19.

S25 Fig. Maximum likelihood phylogeny of chromosome 20.

S26 Fig. Maximum likelihood phylogeny of chromosome Z.

S27 Fig. Maximum Likelihood (ML) phylogenies inferred genome-wide and per
chromosome.

S28 Fig. Marginal tree for one SNPs in Chr4.

S29 Fig. Marginal tree for one SNPs in Chr9.

S30 Fig. Marginal tree for one SNPs in Chr14.