Promoting diversity in ecological systems through toxin production

Ga Ching Lui[1] and Sidhartha Goyal[1]

[1]Department of Physics,
University of Toronto, Ontario, Canada, M5S 1A7
[*]Corresponding author: goyal@physics.utoronto.ca

Abstract

Toxins are used by microbes to kill and compete, and hence, unlike cross-feeding from secondary metabolites, they are less often considered as a candidate to promote diversity in a microbial community. In contrast, some natural communities hint at the possibility that the presence of toxin genes and externally supplied antibiotics may support higher diversity. We address this gap with a model that shows how toxins in small and medium-sized communities can indeed promote diversity. The central idea of the model emerges from the combination of bacterial growth laws of resource partitioning and recent experiments suggesting a trade-off between toxin sensitivity and nutrient efficiency. The second puzzle in natural data is the presence of both negative and positive regulation of toxin production under similar environmental stress. Our model shows how this diversity in regulation of toxin production can promote toxin-producer fitness. In general, we provide a framework to integrate the role of toxins in microbial communities.

Significance statement

Structured organization of microbial communities into functional groups (consumers and producers) has been a framework to understand diverse populations. This framework leaves out the impact of toxin producers, which kill rather than provide the others with benefits, on diversification. Surprisingly, natural microbial communities show a correlation between the number of toxin genes in a community and its diversity. Our work shows how physiological constraints and trade-offs for individual bacteria can lead to a diverse, structured population where toxin-producers serve as a keystone species. This provides a pathway towards toxin production being established as a viable strategy in a diverse population: releasing toxins allows the survival of the less competitive strains, which would otherwise be driven towards extinction by competitive exclusion.

Introduction

Cross-feeding on secondary metabolites released in a microbial community can change the composition of the population. Since producers of these metabolites promote the fitness of other species, cross-feeding can mediate commensal interactions between different bacterial species and sustain diversity under nutrient-limited conditions [17, 34, 37]. This further points to a possible coarse-grained description of community compositions by functional groups [15]. The core idea of this line of work is to consider these secondary metabolites as additional degrees of freedom that describe the system, allowing greater diversity at steady state. In other words, high diversity requires the production of a diverse spectrum of secondary metabolites.

Another category of secondary metabolites is toxins. Unlike cross-feeding, toxin increases the competitive advantage of toxin producers through allelopathy, meaning that toxins in the environment negatively affect their competing species [29]. This advantage can be mediated by interfering with the cellular processes of the competing species, such as suppressing protein biosynthesis, modifying and damaging cellular DNA, or by increasing the permeability of the cytoplasmic membrane, which effectively kills these cells [29, 45, 46, 18, 55]. In this sense, the production of toxins has an opposite effect on diversity compared to cross-feeding.

Figure 1: (A) Environment defined by nutrient and toxin levels with a single species. In region (1), where the toxin level is low or the nutrient level is high, the growth rate of the species is larger than the removal rate, leading to large consumption of nutrients. In region (2), hostile environments lead to a small population size, and the overall nutrient level is replenished by the external supply. Due to this feedback, the nutrient and toxin levels at steady state are given by the ZNGI in Eq. (1). (B). The inset shows the ZNGIs of two species, which can coexist at the nutrient and toxin levels given by the red dot. For multiple species, it is difficult for all ZNGIs to intersect, and coexistence is not expected.

In models that describe interactions of species through toxins [22, 35, 19], it is common to observe multi-stability that leads to the dominance of a single species and lowers the diversity, which has been documented in early experiments such as [5]. In contrast, analysis of the Earth Microbiome Project dataset reports a correlation between diversity and antimicrobial toxin genes in rare taxa [33]. Furthermore, experimental work involving *Straphylococcus* spp. suggests that phenotypic diversity is promoted by intermediate antibi-otic levels, and decreases at both high and low levels of antibiotics such as ciprofloxacin, amikacin, and streptomycin [28]. A related work on mouse guts shows that the presence of bile salt, which solubilizes lipids and is bactericidal, can drive coexistence in small systems with few mutants of *Escherichia coli* due to trade-offs between bile resistance and growth rate [9]. On the theoretical side, the work in [8] suggests that an autoinhibitor-producing species can stabilize a two-species system, and allelopathy can be established through two different paths: either relying on high initial abundances of the producers due to bi-stability, or the producer has to actively change the environment to reduce its own fitness and coexist with its competitor. The latter is described as "problematic from an evolutionary perspective" (or evolutionarily unstable), and one of the ways proposed to resolve this is to consider detoxification by the toxin-resistant species instead of toxin production [30]. It has also been suggested that toxin-producing phytoplankton may be responsible for maintaining coexistence through the effective coupling between species [48, 4].

Unlike cross-feeding, where secretion of secondary metabolites is not considered a cost to the secreting bacterium, the production of toxins is costly in terms of bacterial growth. This suggests that toxins should only be produced when necessary, and tight regulations may be in place to balance the benefit-to-cost ratio. One of the regulation mechanisms is quorum sensing [1], where toxin production is dependent on species abundance, and it has been shown mathematically to exhibit both pure states (only one species survives) and persistence (multiple species survive in oscillatory or chaotic attractors) [3]. Another potential mechanism is competition sensing [7], where toxin producers respond to abiotic stressors [39] and use environmental cues to infer ecological competitions. For abiotic stressors, there has been evidence that unfavorable growth conditions, such as low nutrient availability in the environment, contribute to an increased level of toxin production, and a rich body of literature has been dedicated to studying the effects of toxin production under different stressors [20, 27, 13, 7]. These effects guide the ecological dynamics and dictate the community structures. The intuition is that while species compete for these growth-limiting nutrients, toxins can increase the competitive advantage of the producer, but at a cost. Modeling suggests that species with constitutive production of toxins tend to be out-competed by species with regulated production through quorum sensing or competitive sensing on abiotic stressors [39].

In this work, we present theoretical work that shows how toxins can counterintuitively play a role in supporting diversity. Suppose that the environment of a particular system can be described by some low-dimensional space such that for each species, the space can be partitioned into two different regions: (1)

bacterial growth rate greater than removal rate, and (2) negative net growth. The two regions are separated by a curve plotted in Figure 1A, which is called the zero net growth isocline (ZNGI). ZNGI denotes a set of environmental conditions that correspond to the steady states when that species is present in the system:

$$\text{Growth rate}(\text{Nutrient level}, \text{Toxin level}) - \text{Removal rate} = 0 \ . \tag{1}$$

If another species is added to the system, they can coexist only at the intersection of the ZNGIs (inset of Figure 1B). Now, if more species are added to the system as shown in Figure 1B, either through mutation or immigration, it is improbable that all of these ZNGIs intersect in this low-dimensional space. In this paper, we propose two different paths through which high diversity can be facilitated by toxins: (1) imposing symmetry to the entire population through trade-offs and (2) adjusting the ZNGIs by changing the rate of toxin production. We then address under what circumstances is the strategy of toxin production evolutionarily stable. Finally, we introduce regulation by considering the sensing of abiotic factors (nutrient or toxin levels) and show that persistence through regulation can also increase diversity.

1 The model

A. Growth rate functionals respecting proteome constraint

To investigate if toxins can play a role in sustaining diversity by trade-offs and by the increase in the number of abiotic factors, we need an approach that organically incorporates nutrient competition, toxin susceptibility, and metabolic constraints into the modeling of bacterial growth. We adopt recent models [57, 50] that describe metabolic constraints in terms of proteome partitioning, where the bacterial growth rate $g_i = \phi/(\nu_i^{-1} + \gamma_i^{-1})$ for species i depends on nutrient efficiency ν_i, translation efficiency γ_i in synthesizing proteins, and is subjected to the same proteome fraction devoted to growth (denoted by ϕ) for all species in the system. In SM Sect. 1.A., we show that in a system with one growth-limiting nutrient and one toxin, the bacterial growth rate for species i can be written as

$$g_i = \frac{g_m}{\nu_i^{-1}(n_N) + \gamma_i^{-1}(n_T)} = \frac{g_m}{K + \frac{k_N^{(i)}}{n_N} + \left(\frac{n_T}{k_T^{(i)}}\right)^q} \ , \tag{2}$$

where $k_N^{(i)}$ and $k_T^{(i)}$ are the Michaelis-Menten constants that determine the scales of nutrient level n_N and toxin level n_T in the environment, and $q > 0$ is the Hill coefficient for toxin susceptibility. The parameter K is a constant independent of nutrient and toxin levels, such that g_m/K is the maximal growth rate in an optimal environment with an abundance of nutrients at a low toxin level when $n_N >> k_N^{(i)}$ and $n_T << k_T^{(i)}$, and is proportional to ϕ.

Suppose that a certain fraction $f_i < 1$ of the nutrient flux is redirected from bacterial growth towards toxin production (see SM Sect. 1.B), there is an associated cost incurred in the nutrient efficiency ν_i, such that

$$\nu_i = \frac{1 - f_i}{\nu_m^{-1} + \tilde{k}_N^{(i)}/n_N} \approx \frac{1 - f_i}{\tilde{k}_N^{(i)}/n_N} \ , \tag{3}$$

in the limit of a nutrient-depleted environment $\tilde{k}_N^{(i)}/n_N >> \nu_m^{-1}$. Here, ν_m is the maximum efficiency when the nutrient is abundant, f_i is the production fraction and $k_N^{(i)} = \tilde{k}_N^{(i)}/(1 - f_i)$ depends on cost of production: $\tilde{k}_N^{(i)}$ is the Michaelis-Menten constant when no toxin is produced. By substitution of Eq. (3), the growth rate in Eq. (2) becomes

$$g_i = \frac{g_m}{K + \frac{\tilde{k}_N^{(i)}}{1 - f_i}\frac{1}{n_N} + \left(\frac{n_T}{k_T^{(i)}}\right)^q} \ . \tag{4}$$

As the toxin production fraction f_i increases, the bacterial growth rate g_i decreases due to a decrease in the nutrient efficiency ν_i.

B. Competition in a well-mixed continuous culture system

Populations in continuous culture systems are expected to be reasonably well-mixed with no appreciable spatial structures [56]. In laboratories, continuous long-term culturing is achievable with automated real-time monitoring of population compositions at an affordable cost [41]. As such, we focus on ecological diversity in a chemostat system, which is a continuous culturing device that pumps in fresh medium and removes stale medium at some constant rate. Chemostat experiments allow easy control over population sizes and environmental conditions, such as nutrient levels in the culture, with less fluctuation introduced externally. This provides a reasonable model for natural populations from gut microbiota to sewage [9, 31], although these have spatial structures that may also play a significant role [10, 54]. In general, a chemostat system is described by the following set of equations:

$$\dot{x}_i = (g_i - F)x_i \; ;$$
$$\dot{n}_\xi = (a_\xi - n_\xi)F + \sum_i (\theta_\xi^{+(i)} - \theta_\xi^{-(i)})x_i \; , \tag{5}$$

where x_i is the abundance of species i, g_i is the bacterial growth rate and n_ξ is the concentration of some abiotic factor ξ, with $\xi = T$ for the toxin and $\xi = N$ for the nutrient. A fresh medium, with nutrient and toxin concentrations a_ξ, is pumped into the chemostat at a flow rate of F. The production and consumption of abiotic factor ξ are denoted by $\theta_\xi^{+(i)}$ and $\theta_\xi^{-(i)}$, which describe the change in nutrient and toxin levels due to species i. By adjusting the flow rate F of the chemostat and the supply concentrations a_ξ of the abiotic factors, we can alter the selective pressure imposed on the species and modify the composition of the community.

For the remainder of the paper, we consider various systems defined by Eq. (5) with different forms of growth rates g_i and rates associated with abiotic factors $\theta_\xi^{\pm(i)}$. To survive in the chemostat at steady state, bacteria must change the levels of abiotic factors so that the bacterial growth rate is equal to the removal rate due to the flow of the medium in the chemostat, or $g_i(n_N, n_T) = F$. We first consider the constitutive production of toxins, before investigating the effect of regulation on diversity. Our system is summarized in Figure 2.

Figure 2: Diagram showing a system with a toxin non-producer (orange) and a toxin producer (blue), described by Eq. (4) and Eq. (5). Toxin production fraction f_1 is constant under constitutive production in Sect. 3.1. Regulation in Sect. 3.2 changes f_1 under nutrient sensing (green line) or toxin sensing (purple line), which are parametrized by ρ and σ in Eq. (9).

2 Results

A. Equalization by trade-offs and stabilization by toxin production

Can toxins, which are primarily produced to kill, increase diversity at steady states? We consider the simplest scenario of having only one growth-limiting nutrient N and one toxin T in the system, with concentrations n_N and n_T. The system is defined by Eq. (5) with bacterial growth rate given in Eq. (4). Nutrient N is externally supplied through the inflow of the fresh medium with concentration $a_N > 0$, and is consumed by

species i at a rate of $\theta_N^{-(i)} = g_i$ per cell to support either bacterial growth or toxin production. Toxin T may or may not be externally supplied with $a_T \geq 0$ and is produced by the species at a rate of $\theta_T^{+(i)} = g_i f_i/(1-f_i)$, which increases with production fraction f_i. In our model, the toxin is not consumed and the nutrient is not produced by the cells ($\theta_T^{-(i)} = \theta_N^{+(i)} = 0$). As f_i increases, the rate of toxin production $\theta_T^{+(i)}$ increases at the expense of a decreasing growth rate described in Eq. (4). We show the derivation in SM Sect. 1B.

Suppose that in this system there are five species ($i = 1, \ldots, 5$), each with the Hill coefficient $q = 1$ for toxin susceptibility in Eq. (4), what are the potential outcomes of the competition? According to the competitive exclusion principle [21], when there is only one externally supplied growth-limiting nutrient (with toxin supply concentration set to be $a_T = 0$, and that there is no cross-feeding), it is not possible to have more than one species survive at steady states. The fittest species would dominate the entire population and drive other species towards extinction. However, we notice that if there is at least one species that produces toxins, and that the following condition holds:

$$C = k_N^{(i)} + \frac{1}{k_T^{(i)}} , \qquad \forall i \tag{6}$$

where $C > 0$ is some constant, then the system can be brought to a state where all five species coexist. We will show in the next subsection and in SM Sect. 4 that Eq. (6) can be understood as a trade-off between the nutrient-driven growth rate and the toxin sensitivity observed in experiments. For the time being, we focus on the consequences of Eq. (6) and consider species $i = 1$ to be the only toxin producer in the system.

Given this trade-off, the amount of toxins released by the producer determines the competition outcome, and diversity is sustained only at intermediate values of production fraction f_1. Figure 3A shows the species abundances at steady state when f_1 changes, and illustrates three different regimes. At small $f_1 < f_{c1}$ in zone (1), the system shows a pure state where the toxin producer dominates the population. In zone (2) with $f_{c1} < f_1 < f_{c2}$, the producer only accounts for a portion of the total biomass, while other species successfully invade the population. Diversity increases while the total biomass sustained in the system decreases compared to that in zone (1). At large $f_1 > f_{c2}$ in zone (3), the system returns to a pure state that shows bistability: either the toxin producer or one of the non-producers survives.

A necessary condition for coexistence is fitness equalization. Consider the ZNGI to be $g_i(n_N, n_T) - F = 0$. We plotted the ZNGIs of the five species in Figure 3B-D for the three different zones. Notice that due to the symmetry imposed by the trade-off in Eq. (6), all ZNGIs intersect at $\vec{n}^\star \equiv (n_N = n_N^\star, n_T = n_T^\star)^T$: this denotes the environmental condition in which all species are equally fit and have a growth rate g_i equal to the flow rate F. This is in contrast to the case without trade-offs in Figure 1B, where $\vec{n}^\star$ does not exist. When there is no external supply of toxins ($a_T = 0$), the presence of a toxin producer is required to bring the toxin level up to $n_T^\star$ to achieve fitness equalization.

Under what circumstances then is the toxin producer able to create an environment in which all species are equally fit? To describe how species i changes the environment through both the consumption and production of toxin T and nutrient N, we define the impact factors $m_{T,N}^{(i)}$ as

$$m_{T,N}^{(i)} \equiv \left. \frac{\theta_T^{+(i)} - \theta_T^{-(i)}}{\theta_N^{+(i)} - \theta_N^{-(i)}} \right|_{g_i=F} = \frac{1}{1 - 1/f_i} \leq 0 , \tag{7}$$

which turns out to be uniquely determined by the production fraction f_i. We further define the supply vector $\vec{a} = (a_N/L_n, a_T/L_T)$ and the "impact lines" $n_T/L_T = m_i'(n_N/L_N) + (1-m_i')$ with slope $m_i' = (L_T/L_N)m_{T,N}^{(i)}$, where L_N and L_T are constants that provide proper scalings, and plotted them in Figure 3B-D. We show analytically in SM Sect. 2 that coexistence can only be sustained when the supply vector $\vec{a}$ lies within the shaded region bounded by the impact lines. For toxin producers, the impact lines always have a negative slope ($m_{T,N}^{(i)} < 0$) as nutrient N is always consumed and toxin T is always produced per unit cell growth; a species that does not produce toxins would have a horizontal impact line (or $m_{T,N}^{(i)} = 0$). Without any toxin producer, the bounded region has zero area and coexistence is not possible unless the supply vector $\vec{a}$ is fine-tuned: the supply concentration of toxin a_T has to match exactly with $n_T^\star$. The presence of a toxin producer provides an upward force to the flow in the abiotic factor space, so that fine-tuning is no longer

Figure 3: Diversity sustained by constitutive toxin production. (A) Species abundances and total biomass as production fraction f_1 changes. The three values $f_1 = 0.03, 0.55, 0.80$ are marked by solid black lines showing three phases, and the two transitions are at $f_1 = f_{c1}, f_{c2}$, separating the range of f_1 into zone (1), (2) and (3). (B-D) ZNGIs (solid) and impact lines (dashed) in abiotic space for the producer (blue) and the non-producers (orange). The nutrient level n_N and toxin level n_T are scaled by constants $(g_m/F - K)C^{-1} = L_T^q = L_N^{-1}$ (SM Sect. 2). If the supply vector $(a_T/L_T, a_N/L_N)$, notated by black dot, lies in the shaded region, the coexistence solution is physical, which can either be stable (green) or unstable (magenta). Black arrow shows directions of the dynamics, affecting the stability of coexistence. (E-G) Time series of species abundance (solid) and abiotic factors levels (dashed). In (F), the producer is removed at $t = 50$, leading to the collapse of the community.

required. As f_1 increases, the impact ratio $m_{T,N}^{(1)}$ of the producer increases, which widens the bounded area. Therefore, coexistence can be sustained at a lower nutrient supply concentration a_N. This gives the first transition at $f_1 = f_{c1}$ that we observe in Figure 3A, where $\vec{a}$ is on the impact line of the toxin producer, or $m_{T,N}^{(1)}(f_1 = f_{c1}) = (n_T^\star - a_T)/(n_N^\star - a_N)$. Figure 3E-F shows the time-series of both species abundances and abiotic levels before and after the transition. After the transition, we notice that an increase in diversity is accompanied by a matching of toxin level with nutrient level $n_T/L_T = n_N/L_N = 1$. We call this the shielding effect: the microbiome self-organizes to shield off the asymmetry in the supply concentrations $a_N \neq n_T$, such that the rescaled nutrient and toxin levels can be treated as equals. Therefore, strategies preferential towards nutrient competition are equally optimal to strategies that favor toxin resistance. This was previously observed in models of competition over alternate resources [52, 43, 32] in large randomized systems [53], but not in toxin models. Shielding effect is sustained only in the presence of the producer; removing the producer from the population leads to the collapse of diversity in the population, as shown in Figure 3F.

The implication is that the producer affects the stability of the coexistence state. As the production fraction f_1 increases, both $k_N^{(i)}$ and $k_T^{(i)}$ decrease due to Eq. (6) and the ZNGI is shifted. The ZNGI for the producer at large $f_1 > f_{c2} \equiv 1 - \tilde{k}_N^{(1)}/\max(\tilde{k}_N^{(i)})$ is shown in Figure 3D. Even though the supply vector $\vec{a}$ remains in the bounded region, the coexistence state is no longer stable, as shown in SM Sect. 3. The time-

series in Figure 3G shows the breakdown of the shielding effect as toxin level and nutrient level separate, resulting in the sole survival of a single species. As f_1 approaches 1, the toxin-producing species can no longer be sustained as its growth rate decreases to zero, as the nutrients consumed are funneled into the production of toxins.

Trade-offs

Do we see the above hypothesized trade-off between nutrient efficiency and toxin susceptibility in natural systems? Just to reiterate, the trade-off required for fitness equalization suggests that strains that are more efficient at consuming nutrients pay a price by being more susceptible to toxins. We went further and postulated a specific functional form for this trade-off. As potential evidence for this trade-off we first present the experimental data from [42] that utilized multiple *E. coli* strains in Figure 4A. The data plotted shows the growth rate of a strain relative to that of a reference strain in the absence of toxins against a strain's resistance to three different antibiotics. We show in SM Sect. 4 that the relation

$$k_N + \left(\frac{1}{k_T}\right)^q = C \,, \tag{8}$$

with k_N, k_T, C being the fitting constants, can recapture the different trends between the relative growth rate and toxin resistance shown in Figure 4A. Generally, the shapes of trade-offs affect the fitness of a species, and a convex (or concave) trade-off makes it more beneficial to be a specialist (or a generalist) [12]. In contrast, when the environmental condition $(n_T/L_T)^q = n_N/L_n = 1$ is satisfied, the shielding effect can be achieved in our model, such that all species (including both specialists and generalists) attain the same fitness, regardless of the values of q or the shapes of the trade-offs. It is clear that Eq. (6) in Sect. 2.A is a special case of Eq. (8) corresponding to $q = 1$. While Figure 4A shows that data generally follow the trend due to the trade-off, it also shows that data can depart from the exact relationship, and therefore from Eq. (8). This brings a natural question: how fine-tuned does the trade-off have to be for fitness equalization?

To answer this question, we considered a system with five species characterized by the Michaelis-Menten constants $\{(\tilde{k}_N^{(i)}, k_T^{(i)})\}_5^{i=1}$ which satisfy the trade-off in Eq. (6), but here we made these parameters noisy in time: $\tilde{k}_N^{(i)}(t) = \tilde{k}_N^{(i)} + \eta_N^{(i)}$ and $k_T^{(i)}(t) = k_T^{(i)} + \eta_T^{(i)}$. The deviation from the trade-offs may happen on long time scales due to mutations that lead to changes in the phenotype, or on short time scales due to fluctuations in gene expression. To test the time dependence, we sampled the Michaelis-Menten constants $\{(\tilde{k}_N^{(i)}, k_T^{(i)})\}_5^{i=1}$ across different timescales τ. The noise terms $\eta_\xi^{(i)}$ were sampled from normal distributions $\mathcal{N}(0, \sigma^2)$ for every τ units of time. The number of surviving species after transient dynamics is recorded over an ensemble.

For fast switching time, i.e. $\tau \ll \tau_D$, where τ_D is the cell division time, the diversity is most robust to noise; for example, even for $\tau = 0.3\tau_D$ the diversity remains unchanged until the noise gets beyond $\sim 20\%$. The observed diversity falls as the switching time τ increases. By $\tau \sim 300\tau_D$, a 10% noise is enough to destroy the shielding effect completely. Overall, this suggests that while mutations can break the equalization scheme, fitness equalization is quite robust to gene expression fluctuation.

B. Toxin regulation leads to new behavior - persistence, supersaturation, and producer's improved survival

We showed above how trade-offs sustain diversity through equalization under constitutive production. However, toxin production is costly for bacterial growth, and a more effective strategy is to avoid producing toxins unless necessary. Indeed, up-regulation of toxin synthesis in response to various stressors has been reported [7]. One way to model this is to assume that a species produces more toxins when the growth rate is slow [35], which is indicative of a stressed environment. Here, we explicitly model the dependence of toxin regulation on environmental cues by treating the toxin production fraction as a function of either the nutrient level $f_1 = f_1(n_N)$ or the toxin level $f_1 = f_1(n_T)$, as shown in Figure 2. Consequently, having a generic trade-off in the form of Eq. (8) is no longer possible. Without regulations, the expectation is that no more than two species can coexist in a system with only two growth-limiting factors [36], namely the externally supplied nutrient and the produced toxin.

To understand this better, we considered the competitions between a single toxin producer ($i = 1$) with one ($i = 2$) or more ($i \geq 2$) toxin non-producers, with abundances x_i. Specifically, the bacterial growth

Figure 4: (a) Data adapted from Ferenci et al. [42] to illustrate trade-offs, showing three different trends of nutrient competence (measured by selection coefficient) as toxin resistance increases. These trends (accelerating, linear and decelerating) are qualitatively captured by the model. (b) The number of surviving species as σ and switching time τ changes. Red, dotted line notates the upper limit of diversity given by the competitive exclusion principle. Parameters are given in SM Sect. 8.

rates in Eq. (4) and the dynamics for both the nutrient and the toxin levels in Eq. (5) for this system can be explicitly rewritten as

$$
g_{i=1} = \frac{g_m}{K + \frac{\tilde{k}_N^{(1)}}{n_N}\left(1 - \frac{f_m}{1+(\sigma/s)^p}\right)^{-1} + \frac{n_T}{k_T^{(1)}}} \; ; \quad g_{i\geq 2} = \frac{g_m}{K + \frac{\tilde{k}_N^{(i)}}{n_N} + \frac{n_T}{k_T^{(i)}}} \; ;
$$

$$
\dot{n}_N = (a_N - n_N)F - \sum_i g_i x_i \; ; \qquad \dot{n}_T = (a_T - n_T)F + \frac{\frac{f_m}{1+(\sigma/s)^p}}{1 - \frac{f_m}{1+(\sigma/s)^p}} g_1 x_1 \; ,
$$

(9)

where K is a constant unrelated to nutrient and toxin levels, such that g_m/K is the maximal growth rate in an optimal environment rich in nutrients and low on toxins. We model the dependence of the toxin production fraction from Eq. (4) as a Hill function, $f_1 = f_m/[1 + (\sigma/s)^p]$, where $s = n_N$ for nutrient sensing and $s = n_T$ for toxin sensing. The Hill coefficient p determines how acute the response is, and can take both positive and negative values to span both positive and negative regulations. The constant σ sets the concentration scale of s for sensing. And finally, the parameter f_m is the maximum value that the toxin production fraction f_1 can take; note that $f_m/2$ is the value of the production fraction without regulation.

In this section, we start with a two-species system in which the parameters are set such that the trade-off is intact under the constitutive production of toxins, but now toxin production is regulated. We saw that regulation changes the behavior of the system through altering the fitness of the toxin producer. As shielding is no longer in effect, only certain choices of p and σ would allow the toxin producer to thrive. To further understand the role of regulation for a community, we then considered cases where coexistence is infeasible without regulation; either because the species do not share any trade-offs or the state of fitness equalization is unstable. Lastly, we explored how regulation may enhance the diversity beyond what is supported by the environment - called supersaturation.

Regulation breaks shielding effect and can hurt the producer's ability to survive

How does regulation change the system that exhibits high diversity sustained by trade-offs under constitutive toxin production? Following Eq. (6), we set the Michaelis-Menten constants to respect the trade-off: $k_T^{(2)} = 1/(C - k_N^{(2)})$ and $k_T^{(1)} = 1/(C - k_N^{(1)}/(1 - f_m/2))$. As the toxin production fraction f_m increases, the toxin producer is more resistant to the toxin. Since here we are only considering 2 species, trade-off is not needed for the two ZNGIs to intersect, as shown in the subset of Figure 1B. But we use it to relate the behaviours with no regulations, or $p = 0$, studied in Sect. 2.A. Depending on f_m then, the two species can show coexistence or bistability under constitutive production ($p = 0$), which correspond to zones (2) and (3) in

Figure 3 respectively. We did not consider zone (1) as the change in regulation does not lead to transitions: the producer is always the fittest, with or without regulation.

We focus here on discussing the results for nutrient sensing, for it will become evident later in this section and in SM Sect. 6 that similar behaviours are also exhibited by toxin sensing. Figure 5A-B show the phase diagrams as we span the Hill coefficient p and scaling constant σ. For small $|p|$ the behavior is as expected, but as $|p|$ increases, the two species no longer share the trade-off between the effective nutrient efficiency $1/k_N^{(1)}$ and the toxin resistance $k_T^{(1)}$. This broken symmetry in competitiveness of the two species and can lead to pure states dominated by either the toxin producer (T) or the non-producer (NP), which signifies a change in fitness due to sensing. Note that in the cases shown in Figure 5A-B, there is some significant phase space where the toxin producer loses out due to regulation.

Figure 5: Regulation through nutrient sensing. The four regimes under consideration are: (A) small $f_1 = 0.55$ in zone (2); (B) large $f_1 = 0.9$ in zone (3); (C) no intersections of ZNGIs; and (D) the reverse case for nutrient efficiency. All except (D) has $k_N^{(1)} < k_N^{(2)}$, with ZNGIs similar to Figure 3D. Left figure shows ZNGIs of the producer (blue) and non-producer (orange) at $p = 0$. Right figure shows the phase diagram as Hill function p and sensing scale σ vary. L_N is a constant that sets the scales. Possible competition outcomes are coexistence (C), persistence (P) and pure states dominated by producer (T) or non-producing species (NP). Slash notates bistability of two outcomes (T/C,C/NP,T/NP). Parameters are given in SM Sect. 8.

Regulation rescues species with large toxin production fraction by increasing diversity

If toxin regulation can be detrimental to a toxin producer, as we have argued above, why do they choose to regulate the toxins then? To understand this better, we considered the situations where constitutive production was indeed too expensive and not beneficial to the producer. Here, we found that regulation provides a path for the producer to establish itself in the community.

For Figure 5C, we changed the Michaelis-Menten constants such that the toxin producer without sensing ($p = 0$) is always driven to extinction by the non-producer in the blue regions (NP) due to the cost of toxin production. Note that in the absence of the trade-off, the ZNGIs of the two species do not intersect under constitutive production: the non-producer can tolerate a higher toxin level and a lower nutrient level than the producer without regulation, and is therefore the fitter species. Here, nutrient sensing allows producers to survive in the regions of the phase diagrams that correspond to pure state (T), coexistence (C) or persistence (P). These regions in which the producer becomes more competitive either correspond to large σ with $p > 0$ or small σ with $p < 0$. We show in SM Sect. 5 that both positive and negative sensing shift the ZNGI of the producer by increasing its fitness, leading to coexistence of the two species.

In the presence of the trade-off, a producer with a large production fraction f_m will be able to invade under constitutive production only if its initial abundance is high. As shown in Figure 5B, regulation can move the system away from the brown region (T/NP), allowing the producer to gain fitness and remain within the population regardless of initial conditions.

Toxin down-regulation promotes toxin-producer fitness

When the toxin producer is more nutrient-efficient than the non-producer ($\tilde{k}_N^{(1)} < k_N^{(2)}$, ignoring the cost of toxin production), we observed a common driving force that acts on the competitiveness of the producer for both positive sensing ($p > 0$) and negative sensing ($p < 0$), as seen in regions where toxin producers are the sole survivors in top-right and bottom-left of Figure 5A-C. With positive sensing, the top-right corner corresponds to the limit where toxin production is down-regulated most of the time unless the toxin level is high. Interestingly, we also see a similar trend with negative sensing($p < 0$), where in the bottom-left corner the toxin production is again mostly down-regulated once the nutrient level exceeds some small threshold σ. Hence, for both sensing schemes: positive or negative, the producer thrives when it mostly down-regulates its toxin production over a wide range of values of nutrient levels n_N ($p > 0$, very large σ and $p < 0$, very small σ).

It might make sense for a species to dedicate some of the incoming nutrients to toxin production if it is one of the nutrient-competent species. But what if the producer is less nutrient-competent than the non-producer: ($\tilde{k}_N^{(1)} > k_N^{(2)}$)? Based on the observed trade-off, the producer would be more toxin resistant than the non-producer ($k_T^{(1)} > k_T^{(2)}$). This is the regime in which previous studies show that diversity is not sustainable under constitutive production [8]. We call this the reverse case, which is plotted in Figure 5D. At $p = 0$, either the producer or the non-producer, but not both, survives at the steady state. Regulation does not change the diversity of the system. Since the non-producer is more susceptible to the toxin, it is fitter when the toxin production is down-regulated, and dominates the population at either large σ for positive sensing and small σ for negative sensing.

Regulation allows for persistence and supersaturation

Beyond the steady-state behaviours we have discussed above, regulation also allows for persistence where the abundances of the different species oscillate. Notably, this is observed only for positive regulation $p > 0$. With positive sensing, a high level of nutrient or toxin (which corresponds to a fast growth rate) up-regulates toxin production, and thereby the production cost for the producer. This, in turn, lowers the producer's growth rate. Essentially, this provides negative feedback on the toxin producer and creates a possibility of oscillations. In contrast, during negative sensing ($p < 0$), a higher level of nutrient or toxin down-regulates toxin production and hence increases the growth rate, which serves as positive feedback. Not surprisingly, we see more bistable regions in this case.

Persistence then provides another pathway to increase diversity beyond what is supported by the environment, called supersaturation. It has been shown in models that supersaturation can be generated when the population competes for more than three nutrient resources [24], and that the system observes certain symmetries in the nutrient efficiencies [23]. To this end, we considered the system defined by Eq. (9) with one producer ($i = 1$) with positive sensing and multiple non-producers ($i = 2, 3...$). Without trade-offs, we should expect at most two coexisting species at steady states [36, 21], as discussed above. By comparison, Figure 6A shows that a producer growing along with two non-producers, even though the ZNGIs of the three species in Figure 6B share no intersections. However, this is not a steady state, and here regulation of toxin production leads to oscillating levels of the nutrient and the toxin, preventing any of the species from being consistently fit to dominate the population.

This pathway towards diversity, however, is limited in our model. First, small differences in the parameters drastically change how the system behaves and can break supersaturation. We kept all parameters constant while varying the toxin efficiency of the third species $k_T^{(3)}$ within a small range ($\sim 1\%$), with no discernible qualitative differences in the ZNGIs plot in Figure 6B, and plotted the species abundances in Figure 6A. As $k_T^{(3)}$ decreases within this range, which makes the species less competitive, the maximum abundance attained by the third species drops (from the green curve to the black and brown curves). Supersaturation is not sustained outside of this limited range. Whether there is a trade-off that may allow this "fine-tuning" of parameters for species to be achieved remains to be explored. Notice, however, that given the presence of the "fine-tuned" species, the setup of the system does not need to be fine-tuned, and supersaturation can be sustained over a range of flow rates and nutrient supply concentrations (see SM Sect. 7A).

Second, this mechanism is not scalable in our model. The parameter range that supports supersaturation quickly dropped from $\sim 1\%$ to $\sim 0.1\%$ when another species was added to the system (see SM Sect. 7B), and

Figure 6: Supersaturation of three species supported by a toxin and a nutrient source due to negative feedback under nutrient sensing. A third species with varying $k_T^{(3)}$ (Green, black and brown) is added to a two-species system (blue, orange). (A) Limit cycles of species abundances after transient dynamics. Supersaturation is not sustainable outside of this range. (B) ZNGIs of the three species. Slight changes in toxin resistance of the third species greatly affect the fitness and the competition outcome.

we have not been able to pack more than four species. If supersaturation is to be a significant driving force for diversity in nature, there is likely to be additional symmetries (e.g., trade-offs) at play. While oscillation is frequently observed in systems involving toxins, for example, in ecological suicide [6, 44] where a toxin producer releases autoinhibitors into the environment that lead to oscillating population sizes, its roles in supersaturation remain elusive.

Finally, qualitative behaviours of how regulation under nutrient sensing ($s = n_N$) are also observed in toxin sensing ($s = n_T$). In SM Sect. 6, we show that both sensing schemes share a common mechanism: they effectively regulate toxin production based on the growth rate of the toxin producer ($s = g_m/(k + \tilde{k}_N^{(1)}/n_N + n_T/k_T^{(1)})$). This is in support of a wide array of regulatory strategies observed in nature and will be discussed in more detail in the next section.

3 Discussion

Our first main result is that, contrary to intuition, toxin producers can promote diversity and act as the keystone species. This increase in diversity, however, comes at the cost of lower overall biomass and is a result of the trade-off between toxin resistance and nutrient competence [42, 26]. Second, we show that even when the toxins are not constitutively expressed but are regulated in response to the external environment, toxin producers can sustain diversity by creating a dynamic environment with persistence (oscillations and chaos) in species abundances.

Shielding, tradeoffs, and competitive exclusion

To our knowledge, the shielding effect due to trade-offs involving toxins has not been previously explored. But similar mechanisms were suggested as a pathway to diversity for competitions over alternate nutrients [32]. In a nutshell, the shielding effect for alternate resources results from a shared enzyme budget, which flattens the fitness landscape where different strains grow at the same rate but use different nutrients [52, 43, 32, 53]. As for the case of toxins, the mechanistic basis for this tradeoff in some examples has been linked to the permeability of the cell membrane to both nutrients and toxins [42].

This idea goes beyond ecology. In adaptive immunity, trade-offs have been evoked to explain the diversity observed in T cells [2]. More generally, this is related to multi-objective optimization under a set of constraints, leading to a set of solutions that are equally optimal [51]. In the context of toxins, the trade-off is reflected in the balance between the ability to self-preserve against toxins and the ability to compete for nutrients [26, 42]. Interestingly, experiments show plasticity exhibited by these trade-offs in different environments [42, 25], *i.e.* the exact quantitative trade-off is environment dependent. Despite the seeming differences of the measured shapes, such that a convex (concave) trade-off would generally promote the

fitness of the specialists (generalists) [12], our approach shows that both generalists and specialists share the same fitness under the shielding effect: for different values of q in Eq. (8) that determine these shapes, fitness equalization through the shielding effect remains valid for all species.

Competitive exclusion Generally, to reach steady-state coexistence in a flow driven system (for example, a chemostat) in Eq. (5), all surviving species reach the same growth rate that matches the flow rate $g_i(\vec{n}) = F$ with $i = 1, \cdots, M$, where M is the total number of possible species in the system. Therefore, the solution to this system involves solving M equations with $\dim(\vec{n})$ abiotic variables. This imposes an upper bound on the total number of species $M \leq \dim(\vec{n})$ that may survive in the system and is referred to as the competitive exclusion principle [21]. Previous work exploring how toxins can increase diversity [48, 4] included the role of toxins by treating growth rates as functions of the various species abundances. Consequently, these systems are no longer limited by the competitive exclusion principle due to the increase in dimensionality that scales with the number of species, *i.e.* now the dimensions are $M + \dim(\vec{n})$. Here, rather than postulating a (ad hoc) growth function in the presence of toxins, we built a growth function using recent models for bacterial growth [57, 50], where growth depends only on available nutrients and the toxins produced but not on abundances of other species, making the theoretical more challenging but also more generally applicable.

Previous work has shown that cross-feeding through secreted metabolites can enhance diversity in conditions limited by nutrients [34, 37]. These secreted metabolites, unlike toxins, create commensal and mutualistic interactions. We here suggest that perhaps a community with a large variety of secreted toxins can play a similar role, even if its effect on diversity is not as intuitive as cross-feeding. The impact of toxin-mediated interactions between strains on diversity was explored using random networks [38]: although it was suggested that self-inhibition of toxin producers can contribute to diversity, toxin-mediated interactions in general decrease the likelihood for a community to diversify, as trade-offs were not considered in these models.

Regulating toxins and persistence

Toxins are costly for producers and, therefore, are often regulated. However, the regulatory logic is not straightforward and shows a variety of strategies. For nutrient responses, although it is a common strategy that an increase in ppGpp, which is a signaling molecule in stringent responses such as nutrient limitation, up-regulates toxin production, there are also some cases where ppGpp suppresses toxin synthesis [7]. In response to toxins, autoinduction [49] senses toxins released through the lysis of sibling cells, which serves as an alarm cue of threats from its competitors. Hence, when the toxin level in the environment is elevated, an offensive approach is taken by producing more toxins. By responding to envelope stress, oxidative stress, and DNA damage through mechanisms such as the SOS response (which is important for DNA repairs) [7, 16], bacterial cells can, in effect, regulate toxin production through positive sensing. On the other hand, the same antibiotic stress can up-regulate or down-regulate the expressions of various type II toxin-antitoxin systems [40]. In our model of both nutrient sensing and toxin sensing, we see two different schemes that allow the toxin producer to survive: either large σ with positive sensing $p > 0$, or small σ with negative sensing $p < 0$. This seems to agree with the observed plethora of sensing schemes that act in opposite directions.

Our result shows that regulation of toxin production and its associated costs can change growth rates (fitness), allowing toxin producers to survive and potentially changing the overall diversity of the community. Here, however, the species can also be sustained in the system away from steady state, with species abundances showing persistence (oscillations and/or chaos). This is conceptually related to previously suggested resource competition models where oscillations are generated by competitions over at least three or more resources and is referred to as supersaturation [24, 23, 47, 14, 11]. Similarly, we show that persistence resulting from toxin regulation can also support a diversity higher than the number of abiotic factors and allow the system to tolerate the inexactness in trade-offs.

The rise and evolutionary stability of toxin producers

Given a population of multiple species, how does the strategy of toxin production arise? Taking into account the cost of toxin production, we used the toxin production fraction f_1 - the fraction of nutrients sacrificed for toxin production - to parameterize the ZNGI of the producer. With a large f_1, the producer can dominate the population provided that its initial abundance is high. But this is not evolutionarily stable: any new

producers introduced into the system through mutation would have a low initial abundance. With a small f_1, the newly introduced producer can consistently invade the population. However, in this regime, the toxin producer has a higher nutrient efficiency (small $k_N^{(1)}$) but lower toxin resistance (small $k_T^{(1)}$) compared to the competing species that does not produce toxins. Does it even make sense for a toxin-sensitive species to produce that toxin into the environment, thereby lowering its competitiveness? Both large and small f_1 align with the parameter space in [8] and [30], which consider only two-species competitions, and neither of the regimes reconciles with the rise of toxin production as a viable strategy. Building on their results, our work goes beyond two-species to multi-species competition, which provides a plausible explanation for this conundrum. In our model, it is still true that toxin production might not be a sensible strategy for specialists who invest in being the most nutrient-competent species. However, the toxin producer is now a keystone species, as we have shown in a high-diversity state: the community would collapse, and generalists that are neither the most nutrient-efficient nor the most toxin-resistant within the population would be driven to extinction if no toxin producers are present. By changing the production fraction such that $f_{c1} < f_1 < f_{c2}$ to shift the ZNGI, a generalist as a toxin producer can ensure its survival at a low abundance by sustaining a high diversity state through shielding. This is in support of the observed positive correlation between biodiversity and antimicrobial toxin genes abundances in rare taxa [33], and suggests that allelopathy might have a significant role in sustaining microbial diversity.

References

[1] Rhea G. Abisado, Saida Benomar, Jennifer R. Klaus, Ajai A. Dandekar, and Josephine R. Chandler. Bacterial quorum sensing and microbial community interactions. *mBio*, 9(3):e02331–17, July 2018.

[2] Clemente F. Arias, Miguel A. Herrero, Francisco J. Acosta, and Cristina Fernandez-Arias. Population mechanics: A mathematical framework to study T cell homeostasis. *Scientific Reports*, 7(1):9511, August 2017. Number: 1 Publisher: Nature Publishing Group.

[3] James P. Braselton and Paul Waltman. A competition model with dynamically allocated inhibitor production. *Mathematical Biosciences*, 173(2):55–84, October 2001.

[4] Subhendu Chakraborty, A Ramesh, and Partha Sharathi Dutta. Toxic phytoplankton as a keystone species in aquatic ecosystems: stable coexistence to biodiversity. *Oikos*, 125(5):735–746, May 2016.

[5] L Chao and B R Levin. Structured habitats and the evolution of anticompetitor toxins in bacteria. *Proceedings of the National Academy of Sciences*, 78(10):6324–6328, October 1981. Publisher: Proceedings of the National Academy of Sciences.

[6] Omar E Cornejo, Daniel E Rozen, Robert M May, and Bruce R Levin. Oscillations in continuous culture populations of *Streptococcus pneumoniae* : population dynamics and the evolution of clonal suicide. *Proceedings of the Royal Society B: Biological Sciences*, 276(1659):999–1008, March 2009.

[7] Daniel M. Cornforth and Kevin R. Foster. Competition sensing: the social side of bacterial stress responses. *Nature Reviews Microbiology*, 11(4):285–293, April 2013.

[8] M. J. De Freitas and A. G. Fredrickson. Inhibition as a factor in the maintenance of the diversity of microbial ecosystems. *Journal of General Microbiology*, 106(2):307–320, June 1978.

[9] Marianne De Paepe, Valérie Gaboriau-Routhiau, Dominique Rainteau, Sabine Rakotobe, François Taddei, and Nadine Cerf-Bensussan. Trade-off between bile resistance and nutritional competence drives Escherichia coli diversification in the mouse gut. *PLoS genetics*, 7(6):e1002107, June 2011.

[10] Gregory P. Donaldson, S. Melanie Lee, and Sarkis K. Mazmanian. Gut biogeography of the bacterial microbiota. *Nature Reviews Microbiology*, 14(1):20–32, January 2016. Publisher: Nature Publishing Group.

[11] David J. D. Earn, Pejman Rohani, and Bryan T. Grenfell. Persistence, chaos and synchrony in ecology and epidemiology. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 265(1390):7–10, January 1998. Publisher: Royal Society.

[12] Elias Ehrlich, Lutz Becks, and Ursula Gaedke. Trait–fitness relationships determine how trade-off shapes affect species coexistence. *Ecology*, 98(12):3188–3198, December 2017.

[13] Lusine Ghazaryan, Lilit Tonoyan, Ashraf Al Ashhab, M. Ines M. Soares, and Osnat Gillor. The role of stress in colicin regulation. *Archives of Microbiology*, 196(11):753–764, November 2014.

[14] Michael E. Gilpin. Limit cycles in competition communities. *The American Naturalist*, 109(965):51–60, January 1975. Publisher: The University of Chicago Press.

[15] Joshua E. Goldford, Nanxi Lu, Djordje Bajić, Sylvie Estrela, Mikhail Tikhonov, Alicia Sanchez-Gorostiaga, Daniel Segrè, Pankaj Mehta, and Alvaro Sanchez. Emergent simplicity in microbial community assembly. *Science (New York, N.Y.)*, 361(6401):469–474, August 2018.

[16] Diego Gonzalez and Despoina A. I. Mavridou. Making the best of aggression: the many dimensions of bacterial toxin regulation. *Trends in Microbiology*, 27(11):897–905, November 2019.

[17] Akshit Goyal, Tong Wang, Veronika Dubinkina, and Sergei Maslov. Ecology-guided prediction of cross-feeding interactions in the human gut microbiome. *Nature Communications*, 12(1):1335, February 2021. Number: 1 Publisher: Nature Publishing Group.

[18] Elisa T. Granato, Thomas A. Meiller-Legrand, and Kevin R. Foster. The evolution and ecology of bacterial warfare. *Current Biology*, 29(11):R521–R537, June 2019.

[19] James P. Grover and Feng-Bin Wang. Competition for one nutrient with internal storage and toxin mortality. *Mathematical Biosciences*, 244(2):82–90, August 2013.

[20] K. David Hambright, James D. Easton, Richard M. Zamor, Jessica Beyer, Anne C. Easton, and Brenda Allison. Regulation of growth and toxicity of a mixotrophic microbe: implications for understanding range expansion in *Prymnesium parvum*. *Freshwater Science*, 33(3):745–754, September 2014.

[21] Garrett Hardin. The competitive exclusion principle. *Science*, 131(3409):1292–1297, April 1960. Publisher: American Association for the Advancement of Science.

[22] S. B. Hsu and Paul Waltman. Competition in the chemostat when one competitor produces a toxin. *Japan Journal of Industrial and Applied Mathematics*, 15(3):471–490, October 1998.

[23] Jef Huisman, Anna M. Johansson, Eelke O. Folmer, and Franz J. Weissing. Towards a solution of the plankton paradox: the importance of physiology and life history. *Ecology Letters*, 4(5):408–411, September 2001.

[24] Jef Huisman and Franz J. Weissing. Biodiversity of plankton by species oscillations and chaos. *Nature*, 402(6760):407–410, November 1999.

[25] Christine M. Jessup and Brendan J. M. Bohannan. The shape of an ecological trade-off varies with environment. *Ecology Letters*, 11(9):947–959, 2008. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1461-0248.2008.01205.x.

[26] Thea King, Akira Ishihama, Ayako Kori, and Thomas Ferenci. A regulatory trade-off as a source of strain variation in the species Escherichia coli. *Journal of Bacteriology*, 186(17):5614–5620, September 2004. Publisher: American Society for Microbiology.

[27] Irena Kuhar and Darja Žgur Bertok. Transcription regulation of the colicin Kcka gene reveals induction of colicin synthesis by differential responses to environmental signals. *Journal of Bacteriology*, 181(23):7373–7380, December 1999. Publisher: American Society for Microbiology.

[28] Lewis Lee, Van M. Savage, and Pamela J. Yeh. Intermediate levels of antibiotics may increase diversity of colony size phenotype in bacteria. *Computational and Structural Biotechnology Journal*, 16:307–315, January 2018.

[29] Catherine Legrand, Karin Rengefors, Giovana O. Fistarol, and Edna Granéli. Allelopathy in phytoplankton - biochemical, ecological and evolutionary aspects. *Phycologia*, 42(4):406–419, July 2003.

[30] Richard E. Lenski and Susan E. Hattingh. Coexistence of two competitors on one resource and one inhibitor: A chemostat model based on bacteria and antibiotics. *Journal of Theoretical Biology*, 122(1):83–93, September 1986.

[31] J. N. Lester, R. Perry, and A. H. Dadd. Cultivation of a mixed bacterial population of sewage origin in the chemostat. *Water Research*, 13(6):545–551, 1979. Publisher: Elsevier.

[32] Zhiyuan Li, Bo Liu, Sophia Hsin-Jung Li, Christopher G. King, Zemer Gitai, and Ned S. Wingreen. Modeling microbial metabolic trade-offs in a chemostat. *PLOS Computational Biology*, 16(8):e1008156, August 2020. Publisher: Public Library of Science.

[33] Ya Liu, Yu Geng, Yiru Jiang, Peng Li, Yue-zhong Li, and Zheng Zhang. Global microbial community biodiversity increases with antimicrobial toxin abundance of rare taxa. *The ISME Journal*, 19(1):wraf012, January 2025.

[34] Robert Marsland III, Wenping Cui, Joshua Goldford, Alvaro Sanchez, Kirill Korolev, and Pankaj Mehta. Available energy fluxes drive a transition in the diversity, stability, and functional structure of microbial communities. *PLOS Computational Biology*, 15(2):e1006793, February 2019. Publisher: Public Library of Science.

[35] Ian P. Martines, Hristo V. Kojouharov, and James P. Grover. A chemostat model of resource competition and allelopathy. *Applied Mathematics and Computation*, 215(2):573–582, September 2009.

[36] Richard McGehee and Robert A Armstrong. Some mathematical problems concerning the ecological principle of competitive exclusion. *Journal of Differential Equations*, 23(1):30–52, January 1977.

[37] Pankaj Mehta and Robert Marsland III. Cross-feeding shapes both competition and cooperation in microbial ecosystems, October 2021. arXiv:2110.04965 [cond-mat, q-bio].

[38] Lori Niehaus, Ian Boland, Minghao Liu, Kevin Chen, David Fu, Catherine Henckel, Kaitlin Chaung, Suyen Espinoza Miranda, Samantha Dyckman, Matthew Crum, Sandra Dedrick, Wenying Shou, and Babak Momeni. Microbial coexistence through chemical-mediated interactions. *Nature Communications*, 10(1):2052, May 2019. Publisher: Nature Publishing Group.

[39] Rene Niehus, Nuno M Oliveira, Aming Li, Alexander G Fletcher, and Kevin R Foster. The evolution of strategy in bacterial warfare via the regulation of bacteriocins and antibiotics. *eLife*, 10:e69756, September 2021.

[40] Emeline Ostyn, Yoann Augagneur, and Marie-Laure Pinel-Marie. Insight into the environmental cues modulating the expression of bacterial toxin-antitoxin systems. *FEMS Microbiology Reviews*, page fuaf007, March 2025.

[41] Ue-Yu Pen, Christopher J. Nunn, and Sidhartha Goyal. An Automated Tabletop Continuous Culturing System with Multicolor Fluorescence Monitoring for Microbial Gene Expression and Long-Term Population Dynamics. *ACS Synthetic Biology*, 10(4):766–777, April 2021. Publisher: American Chemical Society.

[42] Katherine Phan and Thomas Ferenci. The fitness costs and trade-off shapes associated with the exclusion of nine antibiotics by OmpF porin channels. *The ISME journal*, 11(6):1472–1482, June 2017.

[43] Anna Posfai, Thibaud Taillefumier, and Ned S. Wingreen. Metabolic trade-offs promote diversity in a model ecosystem. *Physical Review Letters*, 118(2):028103, January 2017. Publisher: American Physical Society.

[44] Christoph Ratzke, Jonas Denk, and Jeff Gore. Ecological suicide in microbes. *Nature Ecology & Evolution*, 2(5):867–872, May 2018. Number: 5 Publisher: Nature Publishing Group.

[45] Margaret A. Riley and David M. Gordon. The ecological role of bacteriocins in bacterial competition. *Trends in Microbiology*, 7(3):129–133, March 1999.

[46] Margaret A. Riley and John E. Wertz. Bacteriocins: evolution, ecology, and application. *Annual Review of Microbiology*, 56(1):117–137, October 2002.

[47] Shovonlal Roy, S. Alam, and J. Chattopadhyay. Competing effects of toxin-producing phytoplankton on overall plankton populations in the Bay of Bengal. *Bulletin of Mathematical Biology*, 68(8):2303–2320, November 2006.

[48] Shovonlal Roy and Joydev Chattopadhyay. Toxin-allalopathy among phytoplankton species prevents competitive exclusion. *Journal of Biological Systems*, 15(01):73–93, March 2007.

[49] Stephanie Schmitz, Anja Hoffmann, Christiane Szekat, Brian Rudd, and Gabriele Bierbaum. The lantibiotic mersacidin is an autoinducing peptide. *Applied and Environmental Microbiology*, 72(11):7270–7277, November 2006.

[50] Matthew Scott, Stefan Klumpp, Eduard M. Mateescu, and Terence Hwa. Emergence of robust growth laws from optimal regulation of ribosome synthesis. *Molecular Systems Biology*, 10(8):747, August 2014.

[51] O. Shoval, H. Sheftel, G. Shinar, Y. Hart, O. Ramote, A. Mayo, E. Dekel, K. Kavanagh, and U. Alon. Evolutionary Trade-Offs, Pareto Optimality, and the Geometry of Phenotype Space. *Science*, 336(6085):1157–1160, June 2012. Publisher: American Association for the Advancement of Science.

[52] Thibaud Taillefumier, Anna Posfai, Yigal Meir, and Ned S Wingreen. Microbial consortia at steady supply. *eLife*, 6:e22644, May 2017. Publisher: eLife Sciences Publications, Ltd.

[53] Mikhail Tikhonov and Remi Monasson. Collective Phase in Resource Competition in a Highly Diverse Ecosystem. *Physical Review Letters*, 118(4):048103, 2017.

[54] Carolina Tropini, Kristen A. Earle, Kerwyn Casey Huang, and Justin L. Sonnenburg. The Gut Microbiome: Connecting Spatial Organization to Function. *Cell Host & Microbe*, 21(4):433–442, April 2017.

[55] Lauren R. Walling and J. Scott Butler. Toxins targeting transfer RNAs: Translation inhibition by bacterial toxin-antitoxin systems. *Wiley interdisciplinary reviews. RNA*, 10(1):e1506, January 2019.

[56] Marvin Whiteley, Erin Brown, and Robert J. C. McLean. An inexpensive chemostat apparatus for the study of microbial biofilms. *Journal of Microbiological Methods*, 30(2):125–132, August 1997.

[57] Conghui You, Hiroyuki Okano, Sheng Hui, Zhongge Zhang, Minsu Kim, Carl W. Gunderson, Yi-Ping Wang, Peter Lenz, Dalai Yan, and Terence Hwa. Coordination of bacterial proteome with metabolism by cyclic AMP signalling. *Nature*, 500(7462):301–306, August 2013. Number: 7462 Publisher: Nature Publishing Group.