Detection and differentiation of *Xanthomonas translucens* pv. *translucens* and pv. *undulosa*

from wheat and barley by duplex quantitative PCR

Heting Fu[1,†], María Constanza Fleitas[2,†], Alian Sarkes[1], Lipu Wang[2], Yalong Yang[1],

Michael W. Harding[3], David Feindel[1], Randy Kutcher[2,*], and Jie Feng[1,*]

[1]Alberta Plant Health Lab, Crop Diversification Centre North, Alberta Agriculture and Irrigation

(AGI), Edmonton, AB, T5Y 6H3, Canada;

[2]Department of Plant Sciences, University of Saskatchewan, Saskatoon, SK, S7N 5A8, Canada;

[3]Crop Diversification Centre South, AGI, Brooks, AB, T1R 1E6, Canada.

[†]Equal contributions.

[*]Corresponding authors:

Randy Kutcher, email: randy.kutcher@usask.ca

Jie Feng, email: jie.feng@gov.ab.ca

Abstract

Two probe-based quantitative PCR (qPCR) systems, namely P-Xtt and P-Xtu, were developed

for diagnosis of cereal bacterial leaf streak pathogens *Xanthomonas translucens* pv. *translucens*

and pv. *undulosa*, respectively. P-Xtt is specific to pv. *translucens*. P-Xtu is specific to pv.

undulosa, pv. *cerealis*, pv. *secalis* and pv. *pistacia*. P-Xtt and P-Xtu worked on all accessible

strains of pv. *translucens* and pv. *undulosa*, respectively. Both systems could detect 100 copies

of the target gBlock DNA. The two systems could be used in both singleplex qPCR and duplex

qPCR with similar efficiencies. On genomic DNA from strains of various *X. translucens*

pathovars, both singleplex qPCR and duplex qPCR could specifically detect and differentiate pv.

translucens and pv. *undulosa*. On infected barley and wheat grain samples, the duplex qPCR

showed similar efficiency compared to a previously published qPCR system but with the

additional capability of pathovar differentiation. The duplex qPCR system developed in this

study will be useful in studies on bacterial leaf streak and detection/differentiation of the

pathogens.

Keywords: bacterial leaf streak, Xtt, Xtu, pathovar *translucens*, diagnostics, qPCR

Introduction

Xanthomonas translucens is a Gram-negative bacterial pathogen causing bacterial leaf streak (BLS) on a range of cereal crops, including wheat, barley, and rye (Jones et al. 1917). The disease was firstly reported on barley (Jones et al. 1917) and wheat (Smith et al. 1919) in the United States and the pathogens were named *Bacterium translucens* and *B. translucens var. undulosum*, respectively. Once Dowson (1939) created the genus *Xanthomonas*, *B. translucens* was reclassified as *X. translucens*. This pathogen is known to cause significant yield losses and reduce crop quality (Bamberg 1936; Duveiller 1994), which can cause serious economic implications for growers and food producers. According to Forster (1982), Idaho in the United States has experienced yield losses of up to 40 percent. In Canada, Drayton (1923) firstly reported that BLS on wheat and barley in Manitoba as early as 1922.

The taxonomy and nomenclature of *Xanthomonas* have undergone multiple changes due to new studies on bacterial biology, phylogenetics, and genomics (Sapkota et al. 2020). The translucens group was created to classify strains of *X. translucens* causing diseases on small grains (Bragard et al. 1995; Vauterin et al. 1995). This group was believed to consist of three different biological entities, pathovar (pv.) *translucens* pathogenic to barley, pv. *undulosa* pathogenic to both wheat and barley, and pv. *cerealis* pathogenic to wheat, barley, oat and bromegrass (Bragard et al. 1995; 1997). Another pathovar, *X. translucens* pv. *secalis*, has been reported to be pathogenic to rye (Reddy et al. 1924). However, phylogenetic studies indicated that strains of pv. *secalis* were clustered with pv. *undulosa* (Bragard et al. 1997; Curland et al. 2018).

Timely and accurate detection and diagnosis of *X. translucens* are critical for the effective

management of this pathogen. The subspecies classification of *X. translucens* has presented

challenges in distinguishing between pathovars. While whole genome sequencing (Langlois et al.

2017) and multilocus sequence analysis (Curland et al. 2018; Ledman et al. 2021; Tambong et al.

2021) have been effective in identifying pathovars, they are time-consuming and impractical on

plant samples. One loop-mediated isothermal amplification (LAMP) protocol has been

developed (Langlois et al. 2017) for the specific detection of three *X. translucens* pathovars, but

cannot differentiate between them. A BLS-specific qPCR has been developed, but also cannot

differentiate between pv. *undulosa* and pv. *translucens* (Sarkes et al. 2022).

In recent years, there has been a rise in cases of BLS in wheat crops in the Midwestern United

States (Adhikari et al., 2012) and western Canada (Kuropatwa, 2014; Arnason, 2021; Tambong

et al., 2021), leading to an increased frequency of grain samples with BLS symptoms being

submitted to the Alberta Plant Health Lab (APHL) since 2019. The objective of this study was to

develop a qPCR-based diagnostic protocol that can detect, quantify and differentiate between

Xanthomonas translucens pathovars, specifically pv. *translucens* and pv. *undolusa*.

Materials and Methods

Chemicals and standard techniques

All chemicals and instrument were purchased from Fisher Scientific Canada (Ottawa, ON)

unless otherwise specified. All primers, probes and gBlocks were synthesized by Integrated

DNA Technologies (Coralville, IA). PCR was conducted in Promega PCR master mix with a

ProFlex PCR system. SYBR Green-based qPCR and probe-based qPCR were conducted in

SsoAdvanced universal SYBR Green supermix (Bio-Rad Canada) and PrimeTime gene

expression master mix (Integrated DNA Technologies), respectively, in a CFX96 touch real-time

PCR detection system (Bio-Rad Canada, Mississauga, ON).

Bacterial isolates

Isolation of X. *translucens* pv. *translucens* and pv. *undulosa* from wheat and/or barley grains

was conducted according to Sarkes et al. (2022). Two pv. *translucens* stains, namely CDCN-

B053 and B054, six pv. *undulosa* strains, namely CDCN-B045, B046, B047, B055, B057 and

B058, were isolated from barley and wheat leaf samples. The isolation and pathovar

identification were described in Sarkes et al (2022). Among these strains, CDCN-B046 and

CDCN-B053 were whole-genome sequenced (GenBank accession numbers JANVBQ010000026

and JANVBP010000007, respectively). Other bacterial species, if not otherwise specified, were

retrieved from the plant pathogen culture collection of APHL and their identities were confirmed

by sequencing of the 16S ribosomal RNA or the *cpn60* gene. PCR amplification of the 16S and

cpn60 followed Klindworth et al. (2013) with the primer pair S-17/A-21 and Sahin et al. (2010)

with the primer pair H1594/H1595, respectively. Sequencing of the PCR products was done with

a SeqStudio genetic analyzer. All bacterial isolates were maintained on 2.3% (w/v) nutrient agar

in darkness at 28°C.

DNA preparation

Two 394-bp double-strand DNA fragments, with a sequence of GenBank accession number

LT604072 (nt: 4206041 to 4206434) and CP043500 (nt: 4106698 to 4107091), were synthesized

as gBlocks and named as G-Xtt and G-Xtu, respectively. Upon receiving, each gBlock was

dissolved in water and a stock solution at 8.3 pM was prepared, which roughly equals to $0.5 \times$

10^7 molecules/µL.

All DNA extraction was conducted using the DNeasy Plant Pro kits (Qiagen Canada, Toronto,

ON) with a Qiacube (Qiagen Canada). DNA extracted from each sample was eluted in 50 µL

elution buffer. For DNA extraction from bacteria, 1 mL cell suspension containing 1×10^7 cells

was centrifuged at 5,000 rcf for 5 min and then the supernatant was removed to leave 100 µL

remaining to avoid disturbing the cell pellet. DNA was extracted from the resultant 100-µL

samples.

Genomic DNA samples of other *X. translucens* strains were kindly provided by researchers in

the United States. These included two pv. *translucens* strains from Dr. Zhaohui Liu, North

Dakota State University, Fargo, ND, and four pv. *translucens* strains, four pv. *undulosa* strains,

one pv. *graminis* strain, one pv. *arrhenatheri* strain, one pv. *poae* strain, one pv. *phlei* strain, one

pv. *phleipratensis* strain and two pv. *secalis* strains from Rebecca Curland, University of

Minnesota, St. Paul, MN. Upon receiving, a 20-30 µL subsample was transferred from each

DNA sample into a new tube and diluted with water to a final volume of 100 µL. The DNA

concentration in each subsample was measured using the NanoDrop 1000 with water as the

blanking sample.

Primer design

In a previous study (Sarkes et al. 2022), 27 genes from *X. translucens* pv. *translucens* strain

DSM18974 were selected as candidates for the development of the qPCR system specific to both

pv. *translucens* and pv. *undulosa*. Among these 27 genes, one gene (GenBank accession number:

SCB06379) was found present in pv. *translucens* only, i.e. not present in any other organisms

based on the current National Center for Biotechnology Information (NCBI) database

(https://www.ncbi.nlm.nih.gov). The genomic sequence of SCB06379 was retrieved and its

location on the whole genome sequence of DSM18974 (GenBank accession number: LT604072)

was identified. The sequence was extended to its 5' end and 3' end, both for 1,000 base pairs at

once, and then the extended sequence was used as the query to BLAST the *X. translucens* pv.

undulosa (taxid:487909) whole-genome shotgun contigs (wgs) database. The sequence extension

and BLAST were repeated until homologous sequences in the two pathovars were found. By this

approach, a heterologous region was identified (Fig. 1). Based on the sequence of SCB06379 and

the sequence of a pv. *undulosa* gene located in the heterologous region, two qPCR primer sets

were designed, respectively, using Primer-Blast (https://www.ncbi.nlm.nih.gov/tools/primer-

blast). For each of the two primer sets, a probe was designed using Primer 3

(https://primer3.ut.ee). The probes in the two primer sets were labelled with different fluorescent

dyes (6-FAM for pv. *translucens* and HEX for pv. *undulosa*). The two primer/probe sets were

named P-Xtt and P-Xtu, respectively (Table 1).

Primer sequence analysis

All primers and probes were further analyzed by Blastn against the NCBI nr/nt database. For

each primer or probe, the presence of identical sequence (only one alignment, with 100%

coverage and 100% identity) in organisms other than *X. translucens* pathovars was checked.

The sequences of the primer target, i.e., the sequence from the 5' end of the forward primer site

to the 3' end of the reverse primer site, of the two qPCR systems were used as queries to Blast

two databases: nr/nt and wgs (Xanthomonas taxid: 338). The Blast results were manually

checked for the absence of sequences identical to the queries in species other than *X. translucens*

pathovars. In addition, the presence of P-Xtt and P-Xtu in all whole-genome sequenced *X.*

translucens pv. *translucens* and pv. *undulosa* was confirmed by checking the presence of all

whole-genome sequenced strains in the Blast results. The lists of whole-genome sequenced

strains of the two pathovars were obtained by searching "Xanthomonas translucens" in the NCBI

genome database (www.ncbi.nlm.nih.gov/genome) and manually checking the resultant Genome

Assembly and Annotation reports.

qPCR reaction

Each qPCR reaction was 20 µL containing 2 µL template and 0.25 µM of each primer. The

probe-based qPCR reaction also contained 0.15 µM probe. The qPCR program consisted of an

initial denaturation step at 95°C for 2 min, followed by 40 cycles of 5 s at 95°C and 30 s at 60°C.

Each qPCR reaction was conducted with three technical repeats.

Specificity test of P-Xtt and P-Xtu on the species level

The specificity of the primer sets P-Xtt and P-Xtu was tested on templates from the following

organisms: *X. translucens* pv. *undulosa* strain CDCN-B046, pv. *translucens* strain CDCN-B053,

healthy wheat grains and healthy barley grains, pellets of 20 mL of YPG media (1% yeast extract,

1% peptone, 2% glucose, w/v) after incubating five healthy wheat grains and five healthy barley

grains at 150 rpm 30°C for 48 hours (the pellets were prepared by centrifuging the culture after

the grains being removed at 4,000 rcf for 5 min) and eight bacterial cultures on nutrient agar

including *Pseudomonas syringae* pv. *atrofaciens*, *P. syringae* pv. *syringae*, *Erwinia rhapontici*,

Clavibacter michiganensis, *Xanthomonas campestris* pv. *pelargonii*, *Pantoea agglomerans*, *P.*

allii and *P. ananatis*. Template DNA in each 20-µL reaction was 2 ng for bacterial samples or 10

ng for plant samples and the grain culture sample. All tests were conducted twice with similar

results.

Specificity test of P-Xtt and P-Xtu on the pathovar level

Genomic DNA from ten strains of *X. translucens* pv. *undulosa*, eight strains of pv. *translucens*,

and one strain of pv. *cerealis*, pv. *secalis,* pv. *graminis*, pv. *arrhenatheri*, pv. *poae*, pv. *phlei* and

pv. *phleipratensis* were tested by P-Xtu and P-Xtt in both singleplex and duplex qPCR. All

templates were also tested with the primer pair specific to *X. translucens* 16S region (F16S/R16S;

Sarkes et al. 2022) by SYBR Green qPCR, which served as DNA quantity control.

Sensitivity test of P-Xtt and P-Xtu on gBlocks

From the stock solution of each of the two gBlocks (G-Xtt and G-Xtu), a set of 10× serial

dilutions were prepared from 0.5×10^7 to 5 molecules/µL (seven solutions for each gBlock).

Using the serial dilutions as templates, sensitivity of P-Xtt and P-Xtu was evaluated in both

singleplex and singleplex qPCR reactions.

Detection of *X. translucens* pv. *translucens* and pv. *undulosa* from plant samples

Three barley grain samples and two wheat grain samples were tested by duplex qPCR using P-

Xtt and P-Xtu in the same reactions. The primers/probe set developed by Sarkes et al (2022) was

used as control. Preparation of samples for DNA extraction followed Sarkes et al. (2022). Briefly,

100 grains from each sample were soaked in 4 mL water in a 50-mL tube. The tube was vortexed

for 30 sec, from which a 1-mL sample of supernatant was transferred to a 1.5-mL tube. The tube

was centrifuged at $11,600 \times g$ for 2 min and then 900 µL of the supernatant was removed. DNA

was extracted from the remaining 100 µL pellet/supernatant mixture, which represented all

extractible DNA from the surface of 25 grains.

Data analysis

In all qPCR, means of the quantification cycle (Cq) values from the three technical repeats were

calculated and treated as one data point. The qPCR standard curves were constructed by

regression analysis using the SAS software (version 9.4; SAS Institute, Cary, NC). qPCR primer

efficiencies were calculated as $E = -1+10^{(-1/\text{slope})}$ (Svec et al. 2015). For the same template DNA

at the same concentration, the Cq values of singleplex qPCR and duplex qPCR were compared

by Student's t-test using the SAS software.

Results and Discussion

Sequence analysis of the specificity of P-Xtt and P-Xtu

Blastn of the primer and probe sequence against NCBI nr/nt database indicated that the forward

primer and the reverse primer of P-Xtt are present in one non-*X. translucens* organism,

respectively, and that the forward primer and the probe of P-Xtt are present in two and four non-

X. translucens organisms, respectively. However, all of these non-*X. translucens* are different

species. Aside from the above exceptions, all the primers and probes are present exclusively in *X.*

translucens pathovars. The results also indicated the specificity of P-Xtt and P-Xtu at the

pathovar levels, which are identical with the results from Blastn of the primer targets of P-Xtt

and P-Xtu as described below.

The primer targets of P-Xtt and P-Xtu are 103 and 93 bp, respectively (Table 1). Using the two

fragments as queries, Blastn against NCBI nr/nt database and wgs (Xanthomonas taxid: 338)

database was conducted (the results were verified on March 2, 2023; Table 2). For both

fragments, Blastn against the two databases covered all whole-genome sequenced *X. translucens*

pv. *translucens* and pv. *undulosa*, and did not produce any hit from a non-*X. translucens*

organism. Orthologs of the P-Xtt target was found in 23 out of the 27 whole-genome sequenced

pv. *translucens*, all the whole-genome sequenced pv. *graminis* (9), pv. *arrhenatheri* (3) and pv.

hordei (1), as well as seven unknown *X. translucens* pathovars. The P-Xtu target was found in all

the whole-genome sequenced pv. *undulosa* (18), pv. *cerealis* (3), pv. *secalis* (2) and pv. *pistacia*,

as well as 17 unknown *X. translucens* pathovars. Orthologs of the P-Xtu target are also present in

the four pv. *translucens* strains that lack the P-Xtt target: strains XtKm33, BLSB3, SIMT-07 and

SLV-2. On the target of the qPCR system F11/R11/P11 development by Sarkes et al. (2022),

these four strains had the same genotype as pv. *undulosa* rather than pv. *translucens* (Table S3 in

Sarkes et al. 2022). Phylogenetic studies (Khojasteh et al. 2019; Shah et al. 2021) indicated that

these four strains are indeed pv. *undulosa* but mislabeled as pv. *translucens*. Manually checking

the primer and probe sequences indicated that P-Xtt and P-Xtu are present in strains where an

ortholog of their target was present except for the forward primer of P-Xtt that was partially

present in the nine pv. *graminis* strains, the three pv. *arrhenatheri* strains and one unknown

pathovar. P-Xtt and P-Xtu, as well as their targets, are not present in pv. *poae*, pv. *phlei* and pv.

phleipratensis. These results indicated that P-Xtt is specific to pv. *translucens* and pv. *hordei*, P-

Xtu is specific to pv. *undulosa*, pv. *cerealis*, pv. *secalis* and pv. *pistacia*, and both P-Xtt and P-

Xtu are ubiquitous among strains of the specific pathovars. It has not escaped our notice that the

whole genome sequence of pv. *hordei* strain UPB458 was relabeled as pv. *translucens* (Table 2).

Thus, based on the database available genomic sequences, we can conclude that P-Xtt is specific

to pv. *translucens* only.

Specificity of P-Xtt and P-Xtu on species level

When P-Xtt and P-Xtu were tested on DNA from healthy (non-symptomatic) wheat and barley,

microorganisms associated with wheat/barley grain surface, and eight other bacterial species, no

positive signal was observed (Table 3). This result indicated that both P-Xtt and P-Xtu are

specific to *X. translucens* and DNA from host plants or other bacteria commonly associated with

wheat/barley grains will not interfere with the qPCR reactions.

Specificity of P-Xtt and P-Xtu on bacterial DNA

DNA from strains of *X. translucens* pathovars were tested by P-Xtt and P-Xtu by singleplex and

duplex qPCR (Table 4). Positive signals (mean Cq value less than 30) were produced by P-Xtt

from all pv. *translucens* strains but not from other pathovars. Cq values at 30 were produced

from the pv. *graminis* and pv. *arrhenatheri* strains, which in agreement with the sequence

analysis that the forward primer of P-Xtt is partially present in these two pathovars. Cq values

higher than 37 were produced from some strains of other pathovars, however, in most cases only

from one or two of the three technical repeats. Similarly, positive signals (mean Cq value less

than 30) were produced by P-Xtu from all pv. *undulosa*, pv. *cerealis* and pv. *secalis* strains but

not from other pathovars. Cq values higher than 37 were produced from some strains of other

pathovars, however, in most cases only from one or two of the three technical repeats. Both P-

Xtt and P-Xtu did not produce positive signals from pv. *poae*, pv. *phlei* and pv. *phleipratensis*.

These results confirmed that P-Xtt is specific to pv. *translucens*, P-Xtu is specific to pv.

undulosa, pv. *cerealis* and pv. *secalis*, and both P-Xtt and P-Xtu are ubiquitous among strains of

the specific pathovars.

Cq values generated by singleplex qPCR and duplex qPCR were compared by Student's t-test.

Among all the applicable samples, p values between 0.0813 and 0.9118 were obtained. These

results indicated that duplex qPCR with P-Xtt and P-Xtu in the same reaction had the same

robustness as singleplex qPCR on detection and differentiation of *X. translucens* pv. *translucens*

and pv. *undulosa*.

Sensitivity of P-Xtt and P-Xtu on gBlocks

The serial dilutions of the gBlock G-Xtt were tested in singleplex qPCR with P-Xtt and in duplex

qPCR with P-Xtt/P-Xtu. Similarly, G-Xtu was tested with P-Xtu and P-Xtt/P-Xtu. Based on the

Cq values, a standard curve was constructed for each primer set (Fig. 2). The efficiencies of P-

Xtt in singleplex and duplex qPCR are 0.98, respectively (Fig. 2a and 2b). The efficiencies of P-

Xtu in singleplex and duplex qPCR are 1.04 (Fig. 2c) and 1.03 (Fig. 2d), respectively. For P-Xtt,

on each concentration of the gBlock serial dilutions, the Cq value from singleplex qPCR (Fig. 2a)

and that from duplex qPCR (Fig. 2b) were compared by two-tailed Student's t-test. Of all gBlock

concentrations, the p values of the t-test were in the range of 0.1233 to 0.9095, indicating no

statistically significant difference in Cq between singleplex and multiplex reactions. Similar

results were observed for P-Xtu (Fig. 2c and d), in which the p values were in the range of

0.2416 to 0.9974. The results confirmed that P-Xtt and P-Xtu can be used in duplex qPCR and

singleplex qPCR with same robustness. In addition, all four standard curves (Fig. 2) indicated

that the low limit of the two qPCR systems for a positive detection in both singleplex and duplex

qPCR is 10 double stranded DNA copies per reaction. However, with consideration of the qPCR

results from bacterial genomic DNA in which a Cq value at 37 could be produced from non-

targeted strains (Table 4), and the influence of DNA extraction efficiency on diagnostic tests

(Yang et al. 2021), we suggest that the duplex qPCR system can reliably detect DNA from 100

bacterial cells.

Detection of *X. translucens* pv. *translucens* and pv. *undulosa* from plant samples

Both duplex qPCR using P-Xtt and P-Xtu and singleplex qPCR using F11/P11/R11 detected *X.*

translucens pv. *translucens* from the three barley samples and pv. *undulosa* from one of the two

wheat samples (Table 5). From the other wheat sample, no signal was detected, indicating the

grains in this sample were not infected by ether of the two pathovars. From the barley sample B2,

pv. *undulosa* was also detected by the duplex qPCR. This result was supported by bacterial

isolation: both pv. *translucens* and pv. *undulosa* were isolated from this sample and the identities

of the isolates were confirmed by sequencing of the 16S ribosomal RNA or the *cpn60* gene.

Our results indicated that P-Xtt and P-Xtu are specific to *Xanthomonas translucens* pv.

translucens and pv. *undulosa*, respectively. The two systems work for all the whole-genome

sequenced strains of these two pathovars and strains/DNA accessible to our lab. The two

primers/probe sets can be used in duplex qPCR to detect and differentiate the two pathovars.

Acknowledgements

This study was supported by a grant from SaskBarley Development Commission to RK.

References

Adhikari, T. B., Gurung, S., Hansen, J. M., and Bonman, J. M. 2012. Pathogenic and genetic

diversity of *Xanthomonas translucens* pv. *undulosa* in North Dakota. Phytopathology

102:390-402.

Arnason, R. 2021. Bacterial leaf streak persists in Alberta.

https://www.producer.com/crops/bacterial-leaf-streak-persists-in-alberta.

Bamberg, R. H. 1936. Black chaff disease of wheat. J. Agri. Res. 52:397-417.

Bragard, C., Verdier, V., and Maraite, H. 1995. Genetic diversity among *Xanthomonas*

campestris strains pathogenic for small grains. Appl. Environ. Microbiol. 61:1020-1026.

Bragard, C., Singer, E., Alizadeh, A., Vauterin, L., Maraite, H., and Swings, J. 1997.

Xanthomonas translucens from small grains: diversity and phytopathological relevance.

Phytopathology 87:1111-1117.

Curland, R. D., Gao, L., Bull, C. T., Vinatzer, B. A., Dill-Macky, R., Van Eck, L., and Ishimaru,

C. A. 2018. Genetic diversity and virulence of wheat and barley strains of *Xanthomonas*

translucens from the Upper Midwestern United States. Phytopathology 108:443-453.

Dowson, W. J. 1939. On the systematic position and generic names of the Gram-negative

bacterial plant pathogens. Zentralblatt fur Bakteriologie, Parasitenkunde,

Infektionskrankheiten und Hygiene. 100:177-193.

Drayton, F. L. 1923. Survey of the prevalence of plant diseases in the Dominion of Canada. p11

and p24. https://phytopath.ca/wp-content/uploads/2021/02/CPDS_1923_4.pdf.

Duveiller, E. 1994. Bacterial leaf streak or black chaff of cereals. EPPO Bull. 1994:24.

Forster, R. L. 1982. The status of black chaff disease in Idaho. Idaho Wheat 12:3-4.

Jones, L. R., Johnson, A. G., and Reddy, C. S. 1917. Bacterial-blight of barley. J. Agricul. Res.,

11:625-643.

Khojasteh, M., Taghavi, S. M., Khodaygan, P., Hamzehzarghani, H., Chen, G., Bragard, C.,

Koebnik, R., and Osdaghi, E. 2019. Molecular typing reveals high genetic diversity of

Xanthomonas translucens strains infecting small-grain cereals in Iran. Appl Environ

Microbiol 85: e01518-19.

Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., and Glöckner, F. O.

2013. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-

generation sequencing-based diversity studies. Nucl. Acids Res. 41:e1.

Kuropatwa, R. 2014. Detect and avoid bacterial leaf streak.

https://www.grainews.ca/features/detect-and-avoid-bacterial-leaf-streak.

Langlois, P. A., Snelling, J., Hamilton, J. P., Bragard, C., Koebnik, R., Verdier, V., Triplett, L.

R., Blom, J., Tisserat, N. A., and Leach, J. E. 2017. Characterization of the *Xanthomonas*

translucens complex using draft genomes, comparative genomics, phylogenetic analysis, and

diagnostic LAMP assays. Phytopathology 107:519-527.

Ledman, K. E., Curland, R. D., Ishimaru, C. A., and Dill-Macky, R. 2021. *Xanthomonas*

translucens pv. *undulosa* identified on common weedy grasses in naturally infected wheat

fields in Minnesota. Phytopathology 111:1114-1121.

Reddy, C. S., Godkin, J., and Johnson, A. G. 1924. Bacterial blight of rye. J. Agr. Res. 28:1039-

1040.

Sahin, N., Gonzalez, J. M., Iizuka, T., and Hill, J. E. 2010. Characterization of two aerobic

ultramicrobacteria isolated from urban soil and a description of *Oxalicibacterium solurbis* sp.

nov. FEMS Microbiol. Lett. 307:25-29.

Sapkota, S., Mergoum, M., and Liu, Z. 2020. The translucens group of *Xanthomonas translucens*:

complicated and important pathogens causing bacterial leaf streak on cereals. Mol. Plant

Pathol. 21:291-302.

Sarkes, A., Yang, Y., Dijanovic, S., Fu, H., Zahr, K., Harding, M. W., Feindel, D., and Feng, J.

2022. Detection of *Xanthomonas translucens* pv. *undulosa*, pv. *translucens*, and pv. *secalis* by

quantitative PCR. Plant Dis. 106:2876-2883.

Shah, S. M. A., Khojasteh, M., Wang, Q., Taghavi, S. M., Xu, Z., Khodaygan, P., Zou, L.,

Mohammadikhah, S., Chen, G., and Osdaghi, E. 2021. Genomics-enabled novel insight into

the pathovar-specific population structure of the bacterial leaf streak pathogen *Xanthomonas*

translucens in small grain cereals. Front. Microbiol. 12: 674952.

Smith, E. F., Jones, L. R., and Reddy, C. S. 1919. The black chaff of wheat. Science 50:48.

Stadhouders, R., Pas, S. D., Anber, J., Voermans, J., Mes, T. H., and Schutten, M. 2010. The

effect of primer-template mismatches on the detection and quantification of nucleic acids

using the 5′ nuclease assay. J. Mol. Diagn. 12:109-117.

Svec, D., Tichopad, A., Novosadova, V., Pfaffl, M. W., and Kubista, M. 2015. How good is a

PCR efficiency estimate: recommendations for precise and robust qPCR efficiency

assessments. Biomol. Detect. Quantif. 3:9-16.

Tambong, J. T., Xu, R., Gerdis, S., Daniels, G. C., Chabot, D., Hubbard, K., and Harding, M. W.

2021. Molecular analysis of bacterial isolates from necrotic wheat leaf lesions caused by

Xanthomonas translucens, and description of three putative novel species, *Sphingomonas*

albertensis sp. nov., *Pseudomonas triticumensis* sp. nov. and *Pseudomonas foliumensis* sp.

nov. Front. Microbiol. 12:838.

Vauterin, L., Hoste, B., Kersters, K., and Swings, J. 1995. Reclassification of Xanthomonas. Int.

J. Syst. Evol. Microbiol. 45:472-489.

Yang, Y., Zhou, Q., Zahr, K., Harding, M. W., Feindel, D., and Feng, J. 2021. Impact of DNA

extraction efficiency on the sensitivity of PCR-based plant disease diagnosis and pathogen

quantification. Eur. J. Plant Pathol. 159:583-591.

Table 1. Primers and probes designed in this study
381

Primer set[a]		Sequence	GenBank accession number
	F	CGGAAGCCTGACGAGATGG	LT604072
P-Xtt	P	/56-FAM/AGAAATCCAGGGCCATCGTC/3IABkFQ/	
	R	TGCCTACGCCGGAATACCG	Nucleotide: 4206294 to 4206396 (103 bp)[b]
	F	ACCACCTCGCCGCTCACAG	CP043500
P-Xtu	P	/5HEX/AAACGTCAT/ZEN/CGGCAGCATG /3IABkFQ/	
	R	CATGATGCTCGCCGATATGC	Nucleotide: 4106793 to 4106885 (93 bp)[b]

382 [a]Primer sets for *Xanthomonas translucens* pv. *translucens* (P-Xtt) and pv. *undulosa* (P-Xtu), each consisting of a forward primer (F), a
383 probe (P) and a reverse primer (R).
384 [b]The fragment corresponding to the sequence from the 5' end of the forward primer to the 3' end of the reverse primer.

385   **Table 2.** Orthologs of the qPCR targets in the whole-genome sequenced strains of *Xanthomonas translucens* pathovars

Pathovar	Strain	GenBank number	Blast LT604072 (4206294-206396)[a]	Blast CP043500 (4106793-41068858)[b]	Database[c]
translucens	AB16-8-8a	JANVBV010000003	27384-27486	No Match	wgs
translucens	AB20-11-12a	JANVBS010000004	29978-30080	No Match	wgs
translucens	AB20-12-12b	JANVBR010000006	143557-143659	No Match	wgs
translucens	ATCC19319	JAJITG010000083	5099-5201	No Match	wgs
translucens	BC10-1-2a	JANUQZ010000003	143578-143680	No Match	wgs
translucens	BC10-1-2b	JANUQY010000006	143578-143680	No Match	wgs
translucens	BC11-3-3a	JANVCA010000012	89200-89302	No Match	wgs
translucens	BC13-4-5a	JANVBZ010000002	143578-143680	No Match	wgs
translucens	BC14-5-5b	JANVBY010000003	27384-27486	No Match	wgs
translucens	BC14-6-6a	JANVBX010000012	89200-89302	No Match	wgs
translucens	BC14-7-6b	JANVBW010000003	27384-27486	No Match	wgs
translucens	CDCN-B053	JANVBP010000007	143564-143666	No Match	wgs
translucens	CIX43	CP072988	3104098-3104200	No Match	nr/nt
translucens	CIX95	CP072990	3691826-3691928	No Match	nr/nt
translucens	DSM18974	CAPJ01000170	13120-13222	No Match	wgs
translucens	DSM18974	LT604072	4206294-4206396	No Match	nr/nt
translucens	Km7	CP064005	1169811-1169913	No Match	nr/nt
translucens	Km8	CP064004	3191872-3191974	No Match	nr/nt
translucens	Km9	CP064003	3740658-3740760	No Match	nr/nt
translucens	UPB458[d]	LWSV01000094	3822-3924	No Match	wgs
translucens	UPB787	JTEM01000253	6566-6668	No Match	wgs
translucens	UPB886	VCIH01000001	3836944-3837046	No Match	wgs
translucens	XtKm34	CP064001	980127-980229	No Match	nr/nt

386   [a,b]The targets of P-Xtt and P-Xtu were used as queries in Blast.
387   [c]Blastn was conducted against two database: the nr/nt database and the wgs database (Xanthomonas taxid: 338).
388   [d]Previously labeled as pv. *hordei*.

389 **Table 2.** Orthologs of the qPCR targets in the whole-genome sequenced strains of *Xanthomonas translucens* pathovars (Cont.)

Pathovar	Strain	GenBank number	Blast LT604072 (4206294-206396)[a]	Blast CP043500 (4106793-41068858)[b]	Database[c]
arrhenatheri	LMG727	CXOI01000063	1773-1875	No Match	wgs
arrhenatheri	LMG727	CP086333	4303903-4304005	No Match	nr/nt
arrhenatheri	UPB455	LWSX01000031	7756-7858	No Match	wgs
graminis	ART-Xtg10	FLTY01000287	2417-2519	No Match	wgs
graminis	ART-Xtg2	FLTV01000281	2190-2292	No Match	wgs
graminis	ART-Xtg29	FLTW01000291	4434-4536	No Match	wgs
graminis	ART-Xtg29	ANGG01000735	952-1054	No Match	wgs
graminis	ART-Xtg9	FLTU01000273	2401-2503	No Match	wgs
graminis	CFBP2053	LHSI01000001	3932637-3932739	No Match	wgs
graminis	ICMP6431	FLTZ01000271	2887-2989	No Match	wgs
graminis	LMG726	CP076254	4384134-4384236	No Match	nr/nt
graminis	UPB437	LWSW01000249	6664-6766	No Match	wgs
hordei	UPB458[e]	CP076249	451351-451453	No Match	nr/nt
unknown	B1	LNTA01000081	28705-28807	No Match	wgs
unknown	B1FA	CP090000	451895-451997	No Match	nr/nt
unknown	B2	LNSZ01000033	4132-4234	No Match	wgs
unknown	B8GF	CP089999	452595-452697	No Match	nr/nt
unknown	CFBP8304	CP074365	4126320-4126422	No Match	nr/nt
unknown	LB5[f]	LNSS01000139	50-152	No Match	wgs
unknown	XT8	LNVW01000003	4130-4232 (range 1)	No Match	wgs
unknown	XT8	LNVW01000003	292242-292344 (range 2)	No Match	wgs

390 [a,b]The targets of P-Xtt and P-Xtu were used as queries in Blast.
391 [c]Blastn was conducted against two database: the nr/nt database and the wgs database (Xanthomonas taxid: 338).
392 [e]Now labeled as pv. *translucens*.
393 [f]Carries the targets of both P-Xtt and P-Xtu.
394

395   **Table 2.** Orthologs of the qPCR targets in the whole-genome sequenced strains of *Xanthomonas translucens* pathovars (Cont.)

Pathovar	Strain	GenBank number	Blast LT604072 (4206294-206396)[a]	Blast CP043500 (4106793-41068858)[b]	Database[c]
translucens	BLSB3[g]	MADS01000080	No Match	14004-14096	wgs
translucens	SIMT-07[g]	MADQ01000160	No Match	3616-3708	wgs
translucens	SLV-2[g]	MADR01000243	No Match	2086-2178	wgs
translucens	XtKm33[g]	CP064002	No Match	1024529-1024621	nr/nt
undulosa	AB17-9-9a	JANVBU010000038	No Match	8493-8585	wgs
undulosa	AB17-10-9b	JANVBT010000037	No Match	4100-4192	wgs
undulosa	BLSW16	MADU01000070	No Match	13990-14082	wgs
undulosa	CDCN-B046	JANVBQ010000026	No Match	15106-15198	wgs
undulosa	CFBP2055	CP074361	No Match	4095104-4095196	nr/nt
undulosa	DOAB1058	JACONR010000071	No Match	8492-8584	wgs
undulosa	ICMP11055	CP009750	No Match	4242324-4242416	nr/nt
undulosa	LW16	CP043540	No Match	4207406-4207498	nr/nt
undulosa	MAI5034	CP089584	No Match	4114745-4114837	nr/nt
undulosa	NARK-1	MADT01000084	No Match	14051-14143	wgs
undulosa	P3	CP043500	No Match	4106793-4106885	nr/nt
undulosa	UPB513	MADV01000055	No Match	5857-5949	wgs
undulosa	XtFa1	CP063996	No Match	3656115-3656207	nr/nt
undulosa	XtKm12	CP064000	No Match	788227-788319	nr/nt
undulosa	XtKm15	CP063997	No Match	3621919-3622011	nr/nt
undulosa	XtLr8	CP063993	No Match	2897415-2897507	nr/nt
undulosa	Xtu4699	CP008714	No Match	4040196-4040288	nr/nt
undulosa	Xtu-UPB513	CP096570	No Match	1158486-1158578	nr/nt

396   [a,b]The targets of P-Xtt and P-Xtu were used as queries in Blast.
397   [c]Blastn was conducted against two database: the nr/nt database and the wgs database (Xanthomonas taxid: 338).
398   [e]Now labeled as pv. *translucens*.
399   [g]These four pv. *undulosa* strains were mislabeled as pv. *translucens* in NCBI database

400 **Table 2.** Orthologs of the qPCR targets in the whole-genome sequenced strains of *Xanthomonas translucens* pathovars (Cont.)

Pathovar	Strain	GenBank number	Blast LT604072 (4206294-206396)[a]	Blast CP043500 (4106793-41068858)[b]	Database[c]
cerealis	01	CP038228	No Match	3385846-3385938	nr/nt
cerealis	CFBP2541	JWHD01000016	No Match	1251724-1251816	wgs
cerealis	CFBP2541	CP074364	No Match	3999642-3999734	nr/nt
pistacia	ICMP16317	CP083804	No Match	3925920-3926012	nr/nt
secalis	CFBP2539	CP074363	No Match	4055150-4055242	nr/nt
secalis	NCPPB1943	MADM01000023	No Match	30140-30232	wgs
unknown	CR31	LNVJ01000003	No Match	131316-131408	wgs
unknown	CS2	LNVK01000002	No Match	8639-8731	wgs
unknown	CS22	LNVL01000002	No Match	119817-119909	wgs
unknown	CS4	LNVM01000050	No Match	15054-15146	wgs
unknown	DAR61454	AMXY01000020	No Match	31611-31703	wgs
unknown	LB10	LNVN01000002	No Match	8639-8731	wgs
unknown	LB5[h]	LNSS01000096	No Match	8639-8731	wgs
unknown	LG48	LNVO01000018	No Match	15298-15390	wgs
unknown	LG54	LNVP01000001	No Match	8639-8731	wgs
unknown	LW16	LNVQ01000045	No Match	9794-9886	wgs
unknown	P3	LNVR01000051	No Match	15300-15392	wgs
unknown	XT123	LNSU01000080	No Match	1138-1230	wgs
unknown	XT130	LNVS01000016	No Match	9273-9365	wgs
unknown	XT5523	LNVT01000017	No Match	90627-90719	wgs
unknown	XT5770	LNVU01000044	No Match	14016-14108	wgs
unknown	XT5791	LNVV01000001	No Match	5833-5925	wgs
unknown	XT-Rocky	LNST01000199	No Match	5872-5964	wgs

401 [a,b]The targets of P-Xtt and P-Xtu were used as queries in Blast.
402 [c]Blastn was conducted against two database: the nr/nt database and the wgs database (Xanthomonas taxid: 338).
403 [h]Carries the targets of both P-Xtt and P-Xtu.

404　**Table 3.** Specificity test of P-Xtt and P-Xtu on DNA of *Xanthomonas translucens* pv.
405　*translucens* and pv. *undulosa* and other organisms
406

DNA template[a]	Cq value (mean±SD; n = 3)	
	P-Xtt	P-Xtu
CDCN-B046 (*X. translucens* pv. *undulosa*)	ns	24.9±0.1
CDCN-B053 (*X. translucens* pv. *translucens*)	23.6±0.1	ns
Healthy wheat leaves	ns	ns
Healthy barley leaves	ns	ns
Healthy wheat grains	ns	ns
Health barley grains	ns	ns
Grain culture in YPG medium[b]	ns	ns
Pseudomonas syringae pv. *atrofaciens*	ns	ns
P. syringae pv. *syringae*	38.7[c]	ns
Erwinia rhapontici	ns	ns
Clavibacter michiganensis	ns	ns
Xanthomonas campestris pv. *pelargonii*	39.3[c]	ns
Pantoea agglomerans	ns	ns
P. allii	ns	ns
P. ananatis	ns	ns

407　[a]DNA template in a 20-µL qPCR reaction: bacteria, 2 ng; plants and the grain culture, 10 ng.
408　[b]DNA from microorganisms grown in YPG media (1% yeast extract, 1% peptone, 2% glucose,
409　w/v) after incubating healthy wheat and barley grains at 150 rpm 30°C for 48 hours
410　[c]Only one of the three technical repeats produced a Cq value
411

412   **Table 4.** Specificity test of P-Xtt and P-Xtu on DNA of *Xanthomonas translucens* pathovars by singleplex and duplex qPCR

Strain Name	Pathovar	Origin[a]	P-16S[b]	P-Xtt Singleplex	P-Xtt Duplex	t-test[e]	P-Xtu Singleplex	P-Xtu Duplex	t-test[e]
CDCN-B045	*undulosa*	APHL	17.1 ± 0.2^c	ns	ns	na	25.3±0.1	25.7±0.1	0.1005
CDCN-B046	*undulosa*	APHL	17.5±0.1	ns	ns	na	25.3±0.3	25.7±0.1	0.1480
CDCN-B047	*undulosa*	APHL	17.2±0.1	ns	ns	na	25.7±0.4	25.4±0.2	0.4324
CDCN-B055	*undulosa*	APHL	18.9±0.2	ns	39.4	na	27.1±0.1	27.2±0.1	0.5415
CDCN-B057	*undulosa*	APHL	19.1±0.1	38.1	38.4 ± 1.0^d	na	27.4±0.3	27.4±0.2	0.8439
CDCN-B058	*undulosa*	APHL	18.8±0.1	38.9	ns	na	26.8±0.4	26.6±0.1	0.6108
CIX23	*undulosa*	UM	17.8±0.2	ns	ns	na	26.4±0.2	26.6±0.2	0.4235
CIX40	*undulosa*	UM	17.9±0.2	38.4 ± 1.3^d	38.5 ± 1.4^d	0.9118	26.9±0.1	26.9±0.1	0.6842
CIX102	*undulosa*	UM	19.0±0.1	38.2 ± 1.1^d	38.4	na	27.7±0.2	27.8±0.2	0.4307
LMG892	*undulosa*	UM	18.8±0.1	38.9	ns	na	27.6±0.2	27.6±0.1	0.8463
CDCN-B053	*translucens*	APHL	18.0±0.1	23.4±0.4	23.3±0.3	0.8348	ns	ns	na
CDCN-B054	*translucens*	APHL	19.3±0.1	24.3±0.2	24.6±0.2	0.0977	ns	ns	na
XttB1FA	*translucens*	NDSU	15.9±0.1	21.3±0.2	21.3±0.3	0.8151	ns	ns	na
XttB8GF	*translucens*	NDSU	15.5±0.2	20.6±0.4	20.2±0.2	0.2720	ns	ns	na
CIX86	*translucens*	UM	16.1±0.1	22.2±0.1	22.2±0.2	0.6104	ns	37.9±0.2	na
CIX87	*translucens*	UM	18.2±0.1	24.2±0.2	24.2±0.2	0.8833	ns	ns	na
CIX94-95	*translucens*	UM	18.4±0.2	23.9±0.3	24.3±0.3	0.2170	ns	ns	na
LMG876	*translucens*	UM	16.4±0.1	22.0±0.3	22.5±0.3	0.1133	39.1	38.9 ± 0.8^d	na
LMG679	*cerealis*	UM	17.5±0.2	ns	38.6 ± 1.7^d	na	26.1±0.1	25.9±0.1	0.0813
LMG883	*secalis*	UM	16.7±0.1	39.1	38.4 ± 0.6^d	na	25.2±0.4	25.0±0.2	0.4993
LMG726	*graminis*	UM	18.6±0.2	30.3±0.2	30.4±0.5	0.6458	ns	39.1 ± 0.4^d	na
LMG727	*arrhenatheri*	UM	18.0±0.1	30.3±0.2	30.9±0.4	0.0939	39.2	38.8 ± 0.6^d	na
LMG728	*poae*	UM	17.6±0.1	37.0±0.2	37.4±0.9	0.4954	ns	ns	na
LMG730	*phlei*	UM	19.4±0.2	ns	38.7	na	ns	39.4	na
LMG843	*phleipratensis*	UM	17.1±0.1	37.3	ns	na	ns	38.6 ± 0.3^d	na

[a]Origin of strains or DNA. APHL, strains isolated by this study; UM, DNA provided by Rebecca Curland, University of Minnesota;
NDSU, DNA provided by Dr. Zhaohui Liu, North Dakota State University.
[b]Primer pair for the 16S region used in SYBR Green qPCR as DNA controls.
[c]Cq value shown as mean ± SD (n =3); ns = no signal from all of the three technical repeats; Data without a SD: only one of the three
technical repeat produced Cq.
[d]Two of the three technical repeat produced Cq.
[e]p value for two-tailed Student's t-test between the Cq value of the singleplex qPCR and that of the duplex qPCR; na = not applicable.
420
421

422   **Table 5.** Test of three barley grain samples (B1-B3) and two wheat grain samples (W1 and W2)

423   by duplex qPCR

Primers[a]	B1	B2	B3	W1	W2	CDCN-B046[b]	CDCN-B053[b]
P-Xtt	23.7±0.1[c]	25.6±0.1	21.6±0.2	38.1[d]	ns[e]	ns	22.8±0.1
P-Xtu	39.2[d]	25.9±0.1	ns	27.6±0.1	ns	25.3±0.1	ns
F11/P11/R11	24.5±0.1	25.0±0.1	22.5±0.1	26.8±0.2	ns	24.4±0.1	23.3±0.1

[a]P-Xtt and P-Xtu were used in duplex qPCR; the primers/probe set F11/P11/R11 (Sarkes et al.
2023) was used as control.
[b]The *X. translucens* pv. *undulosa* strain CDCN-B046 and the pv. *translucens* strain CDCN-B053
were used as control (2 ng DNA per qPCR reaction).
[c]Cq values shown as mean ± SD (n =3).
[d]Only one of the three technical repeat produced Cq.
[e]ns = no signal from all of the three technical repeats.

Figures

Fig. 1. Diagrammatic sketch illustrating the genomic positions of P-Xtt and P-Xtu. The

heterologous region, framed by homologous regions, on GenBank accession number LT604072

(*Xanthomonas translucens* pv. *translucens* strain DSM18974) and on GenBank accession

number CP043500 (*X. translucens* pv. *undulosa* strain P3) is indicated. The brackets indicate the

locations of P-Xtt and P-Xtu. The double-headed arrows indicate the location of the two gBlocks

(G-Xtt and G-Xtu).

Fig. 2. Sensitivity of the qPCR systems P-Xtt for *Xanthomonas translucens* pv. *translucens* and
P-Xtu for pv. *undulosa* on gBlock DNA. a and b, P-Xtt singleplex qPCR and P-Xtt/P-Xtu duplex
qPCR on the pv. *translucens* gBlock, respectively. c and d, P-Xtu singleplex qPCR and P-Xtt/P-
Xtu duplex qPCR on the pv. *undulosa* gBlock, respectively. The qPCR standard curves were
generated from the mean of quantification cycle (Cq) values against log10 of gBlock DNA
copies in one reaction. The R^2 score of the equation and the efficiency of the primers (E) are
indicated over the curve. Efficiency was calculated as $E = -1+10^{(-1/slope)}$. Each data point is shown
as mean $\pm$ SD (n=3).