Vaccination with mycobacterial lipid loaded nanoparticle leads to lipid antigen persistence and memory differentiation of antigen-specific T cells

Eva Morgun,[1] Jennifer Zhu,[2] Sultan Almunif,[2] Sharan Bobbala,[2, †], Melissa S. Aguilar,[3] Junzhong Wang, [1, ‡] Kathleen Conner, [1] Yongyong Cui, [1] Liang Cao,[1] Chetan Seshadri,[3] Evan A. Scott,[1,2]* Chyung-Ru Wang[1]*

[1]Department of Microbiology and Immunology, Feinberg School of Medicine, Northwestern University, Chicago, IL, USA

[2]Department of Biomedical Engineering, Northwestern University, Evanston, IL, USA

[3]Department of Medicine, University of Washington School of Medicine, Seattle, WA, USA

Present addresses

[†]Department of Pharmaceutical Sciences, School of Pharmacy, West Virginia University, Morgantown, WV, USA

[‡]Department of Infectious Disease, Union Hospital of Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China

***Corresponding authors:** Dr. Chyung-Ru Wang (chyung-ru-wang@northwestern.edu)

Dr. Evan A Scott (evan.scott@northwestern.edu)

Running title: Nanoparticle encapsulated lipid elicits antigen persistence

Keywords: nanoparticle, antigen persistence, CD1, mycolic acid, T cell, vaccination, *Mycobacterium tuberculosis*

Abstract

Mycobacterium tuberculosis (Mtb) infection elicits both protein and lipid antigen-specific T cell responses. However, the incorporation of lipid antigens into subunit vaccine strategies and formulations has been underexplored, and the characteristics of vaccine-induced Mtb lipid-specific memory T cells have remained elusive. Mycolic acid (MA), a major lipid component of the Mtb cell wall, is presented by human CD1b molecules to unconventional T cell subsets. These MA-specific CD1b-restricted T cells have been detected in the blood and disease sites of Mtb-infected individuals, suggesting that MA is a promising lipid antigen for incorporation into multicomponent subunit vaccines. In this study, we utilized the enhanced stability of bicontinuous nanospheres (BCN) to efficiently encapsulate MA for *in vivo* delivery to MA-specific T cells, both alone and in combination with an immunodominant Mtb protein antigen (Ag85B). Pulmonary administration of MA-loaded BCN (MA-BCN) elicited MA-specific T cell responses in humanized CD1 transgenic mice. Simultaneous delivery of MA and Ag85B within BCN activated both MA- and Ag85B-specific T cells. Notably, pulmonary vaccination with MA-Ag85B-BCN resulted in the persistence of MA, but not Ag85B, within alveolar macrophages in the lung. Vaccination of MA-BCN through intravenous or subcutaneous route, or with attenuated Mtb likewise reproduced MA persistence. Moreover, MA-specific T cells in MA-BCN-vaccinated mice differentiated into a T follicular helper-like phenotype. Overall, the BCN platform allows for the dual encapsulation and *in vivo* activation of lipid and protein antigen-specific T cells and leads to persistent lipid depots that could offer long-lasting immune responses.

Introduction

Vaccination is currently the most effective method for the prevention and eradication of infectious diseases. These efforts have been enhanced by the development of subunit vaccine formulations that include only immunodominant antigens from pathogens paired with select adjuvants. In contrast to attenuated and inactivated vaccines that contain diverse molecular components of pathogens, subunit vaccines are often preferred due to their simplicity, safety, stability, and scalability of production [1]. Yet the development of effective vaccines for several key endemic parasitic and bacterial pathogens has remained elusive. Antigen selection for subunit vaccines has thus far focused on protein molecules, overlooking a vast family of non-protein antigens such as lipids/glycolipids, metabolites, and phosphoantigens, which can be found on and within pathogens and can be recognized by various subsets of unconventional T cells [2].

Persistent antigen exposure is known to occur naturally after infection through antigen archiving by follicular dendritic cells (FDCs), which periodically present antigen to cognate B cells and aid in affinity maturation and somatic hypermutation [3, 4]. Antigen persistence can also support the maintenance of T cell response through the periodic priming of circulating T cells [5-10]. Subcutaneous vaccination has also been shown to lead to archiving of protein antigens within lymphatic endothelial cells (LECs), which induce the development of protective T cell immunity [11, 12]. However, little is known about how other vaccination routes and vaccine formulations may affect antigen archiving and persistence.

Lipid and glycolipid antigens can be presented by group 1 (CD1a, b, and c) and group 2 (CD1d) CD1 molecules to CD1-restricted T cells. Unlike MHC molecules, CD1 molecules exhibit limited polymorphism [13]. Therefore, CD1-restricted microbial lipid antigens are likely to be recognized by most individuals, making them attractive targets for vaccine development [14]. Group 1 CD1-restricted T cells have diverse T cell receptors and can recognize a variety of lipid and glycolipid antigens [15]. Microbial lipid-specific group 1 CD1-restricted T cells have been identified from several pathogens, including Mtb and other mycobacterial species: *Staphylococcus aureus* (SA), *Borrelia*

burgdorferi, *Salmonella typhimurium*, and *Brucella melitensis* [15]. CD1b-restricted T cells specific to

Mtb lipids including mycolic acid (MA), an immunodominant lipid antigen[16] and component of the

mycobacterial cell wall, have been identified in patients with tuberculosis (TB) [17, 18]. As group 1

CD1 molecules are not expressed in mice, we have previously developed a human CD1 transgenic

mouse model (hCD1Tg). [19] Using hCD1Tg mice, we have shown that transgenic CD1b-restricted

DN1 T cells specific for MA could protect against Mtb infection [20], making these potential antigens

of interest in a subunit vaccine against TB.

The primary goal of a vaccine is to induce the differentiation of the adaptive immune system

such that upon re-exposure to the antigen, a faster and larger response occurs, halting pathogen spread.

While conventional T cells and B cells are known to undergo this differentiation, unconventional T

cells such as NKT cells do not possess a comparable memory phenotype. The presence of Mtb lipid-

specific group 1 CD1-restricted memory T cells has been described in both human and mice [17-19].

However, the properties of memory group 1 CD1-restricted T cells, which share similar developmental

pathway as NKT cells, are not known.

Lipid antigens are difficult to formulate and administer due to their hydrophobic nature. To

assess the utility of MA as a component of a subunit vaccine, we previously encapsulated MA in

poly(ethylene glycol)-block-poly(propylene sulfide) (PEG-*b*-PPS) micellar nanocarriers (MA-MC)

[21]. Nanocarriers composed of PEG-*b*-PPS possess several unique benefits compared to other

delivery platforms, including high stability in aqueous solution [22], minimal background

immunostimulation [23], and triggered release of cargo when exposed to physiological levels of

reactive oxygen species [24-26]. We showed MA-MC could effectively activate adoptively transferred

DN1 T cells and elicit polyclonal group 1 CD1-restricted T cell response in hCD1Tg mice after

intranasal delivery [21]. While effective for stable loading of lipophilic and amphiphilic payloads,

micellar nanocarriers, do not allow for facile encapsulation of hydrophilic payloads like protein

antigens.

Here, we build upon our prior lipid antigen vaccine formulation by employing a nanocarrier

platform engineered for the dual loading of both lipid and protein antigens. Bicontinuous nanospheres

(BCN) are polymeric analogs to lipid cubosomes and thus possess a highly organized internal cubic

organization containing both intertwined hydrophobic bilayers for lipid loading as well as hydrophilic

aqueous channels for protein loading [27]. Using the flash nanoprecipitation (FNP) technique [28, 29],

we fabricated MA-loaded BCN (MA-BCN) and found them to have superior stimulatory capacity *in*

vivo compared to MA loaded poly(D,L-lactide-co-glycolide) (PLGA) nanocarriers (MA-PLGA),

which is a widely used nanocarrier platform for Mtb vaccine development [30, 31]. Furthermore, we

found that after MA-BCN vaccination, MA could persist for 6 weeks post-vaccination within alveolar

macrophages, a phenomenon which was dependent on the presence of an encapsulating vector,

nanocarrier or bacterial, but not the route of vaccination. Due to the enhanced stability of the BCN

architecture that can support the loading of diverse and multiple payloads [22, 32], we were able to

efficiently co-encapsulate within BCN both MA and Ag85B, an immunodominant protein antigen of

Mtb [33]. Interestingly, while both antigens activated their corresponding antigen-specific T cells,

only MA resulted in antigen persistence, suggesting a potentially unique characteristic of subunit

vaccines containing lipid antigens.

                                                    Results

BCN morphology preserved after MA loading

We scalably assembled spherical BCN with and without loaded MA using flash nanoprecipitation (FNP) as previously described [29]. Dynamic light scattering (DLS) showed that both MA-BCN and BCN were consistent in size (364 ±19 nm and 354 ± 14 nm, respectively), and were monodisperse based on their polydispersity indices (PDIs) of 0.24 ± 0.04 and 0.21 ±0.05, respectively (**Fig 1A**). We next verified that the BCN formulation maintained its characteristic interconnected aqueous channels using cryogenic transmission electronic microscopy (cryo-TEM) (**Fig 1B**) and negative staining transmission electron microscopy (TEM) (**Fig 1C**). Using small angle X-ray scattering (SAXS) studies, we confirmed the internal cubic organization of BCN. Bragg peaks at the √2, √4, and √6 ratios show that the primitive type of cubic internal organization was preserved between MA-BCNs and BCNs (**Fig 1D**). Thus, MA encapsulation did not disturb the BCN architecture. We also manufactured a PLGA nanocarrier formulation (PLGA-NP) with and without MA encapsulation. PLGA-NP morphology were confirmed using cryoTEM (**Suppl Fig 1A**) and SAXS (**Suppl Fig 1B**) and the size of blank PLGA-NP and MA-loaded PLGA-NP (MA-PLGA) were found to be 169 ± 4 and 163 ± 5 nm, respectively, with PDIs of 0.18 ± 0.04 and 0.19 ± 0.12, respectively (**Fig 1E**). Using the coumarin derivatization method [21], we found that BCNs could more efficiently encapsulate MA than PLGA-NPs (**Fig 1F**). We previously noted that the highly stable cubic architecture of BCN can lead to the retention of cargo within the endolysosomal compartment [22]. Indeed, Texas Red-Dextran loaded BCN were mostly found in lysosomes while Texas Red-Dextran loaded PLGA-NP could be found both within the lysosome and cytosol of cells (**Supp Fig 2A, B**).

MA-BCN activate MA-specific T cells *in vitro* and *in vivo*

As PLGA has served as a component of a wide range of vaccine formulations, including Mtb vaccines [30, 31], we benchmarked PEG-*b*-PPS BCN against PLGA-NP as a nanocarrier platform for lipid antigens. Both MA-BCN and MA-PLGA were highly effective in activating both mouse (DN1)

and human (M11) MA-specific T cells *in vitro* and inducing IFN-γ production (**Fig 2A-D**). In particular, MA-BCN were significantly better at activating MA-specific T cells compared to free MA at equivalent concentrations (**Fig 2A, B and D**). In comparison to MA-BCN, MA-PLGA was significantly more effective at stimulating mouse MA-specific T cells *in vitro* at lower concentrations. Interestingly, even in the absence of MA, PLGA-NP showed strong stimulation particularly for M11 T cells. These results demonstrate that MA encapsulation within BCN enhances its ability to stimulate antigen-specific T cells and reveal a considerable background, and thus difficult to control, stimulatory effect of blank PLGA-NP. In contrast, blank BCN showed little background immunostimulation [23], allowing a more controlled dose-dependent increase in T cell responses as the loaded MA concentration increased.

We next tested the ability of MA-loaded and unloaded BCN and PLGA to stimulate DN1 T cells *in vivo*. Intranasal (IN) vaccination of hCD1Tg mice was followed by the adoptive transfer of DN1 T cells. Surprisingly, we found that vaccination with MA-BCN induced a significantly higher percent of proliferation and activation in DN1 T cells than MA-PLGA in the draining lymph nodes (LN) and lungs at 1 week post-vaccination (**Fig 2E-G**). The extent of cell proliferation induced by MA-PLGA was not significantly different from that of blank PLGA (**Fig 2F**). Since MA-BCN effectively stimulated MA-specific T cells *in vivo* while MA-PLGA did not, we focused on further characterizing and assessing vaccination via solely MA-BCN formulations.

MA persists while Ag85B does not after vaccination with Ag85B-MA-BCN

Given the lack of knowledge regarding BCN and lipid antigen kinetics, it would be of interest to test whether delivered antigen could persist in the mouse weeks post-vaccination (**Fig 3A**). We switched from intranasal vaccination to intratracheal (IT) vaccination, as it allowed for the reliable delivery of a larger volume of the vaccine. We found that a significant proportion of the adoptively transferred DN1 T cells could proliferate and become activated in the LNs, lung, and spleen of MA-BCN-vaccinated mice 6 weeks post vaccination (**Fig 3B-D**). This activation and proliferation were

MA-specific, as no activation and proliferation of Ag85B-specific p25 Tg (p25) T cells were

observed[34].

Since BCN allow for the co-loading of lipid and protein antigens, we assessed the ability of

dual loaded Ag85B-MA-BCN to activate antigen-specific T cells as well as lead to antigen persistence.

We co-loaded Ag85B and MA in BCN and found Ag85B-MA-BCN to have a diameter of 380 nm and

PDI of 0.223, comparable to MA-BCN (**Fig 4A**). Encapsulation efficiency was also relatively high at

70%.Vaccination of hCD1Tg mice with Ag85B-MA-BCN activated and induced proliferation of both

p25 and DN1 T cells in the LN, lung, and spleen (**Fig 4B-D**) at one-week post-vaccination. In addition,

DN1 T cells were still able to be activated 6 weeks post-vaccination in the context of an Ag85B-MA-

BCN vaccination, while p25 T cells did not show increased activation or proliferation compared to

blank BCN control at this time point (**Fig 4E-H**). Therefore, MA, but not Ag85B, appears to persist in

the context of an Ag85B-MA-BCN vaccination.

MA persistence was dependent on encapsulation but not route of delivery or delivery vector

To determine whether BCN encapsulation contributed to MA persistence, we designed an

experiment that would allow direct comparison between MA-BCN and free MA. To account for their

differential efficiency in stimulating DN1 T cells, we pulsed bone marrow-derived dendritic cells

(BMDCs) from hCD1Tg mice with MA or MA-BCN at 10 μg/mL and 5 μg/mL MA concentration,

respectively (**Fig 5A**). hCD1Tg mice were immunized with MA or MA-BCN-pulsed BMDCs at either

1 week or 6 weeks before the adoptive transfer of DN1 T cells, allowing comparison of short-term *vs*.

long-term vaccination conditions. We found that while there were no significant differences in DN1 T

cell activation and proliferation between mice vaccinated with free MA and MA-BCN pulsed BMDCs

at 1-week post-vaccination, after 6 weeks, DN1 T cell response could only be detected in the MA-BCN

vaccination condition (**Fig 5B, C**). Thus, BCN encapsulation contributed to the persistence of MA. We

then tested whether the BCN structure could contribute to antigen persistence by comparing PEG-*b*-

PPS BCNs to PEG-*b*-PPS MCs using a similar experimental set up (**Suppl Fig 3**). We found that MA-

BCN and MA-MC overall had similar ability to activate DN1 T cells in both short-term and long-term

vaccination time points (**Suppl Fig 3B, C**). Thus, the structure of the encapsulating NP did not play a significant role in antigen persistence of MA.

To investigate if the IT route of vaccination played a role in the development of antigen persistence, we tested intravenous (IV) and subcutaneous (SC) vaccination routes with MA-BCN. DN1 T cell activation and proliferation were detected at 6-weeks post-vaccination through both routes of vaccination (**Fig 5D-F**), suggesting that the antigen persistence is not unique to IT vaccination. To assess whether Mtb lipid antigen persistence could also be observed in mice vaccinated with an attenuated strain of Mtb, we vaccinated hCD1Tg mice SC with Mtb mc^{2}6206 strain and tested for the persistence of both Ag85B and MA after 6 weeks. The Mtb mc^{2}6206 strain displays less virulence in mouse studies than BCG but still confers long-term protection against virulent Mtb challenge equivalent to BCG[35]. We confirmed that no bacterial burden remained at the 6 weeks time point (**Suppl Fig 4A**). Similar to MA-BCN-vaccinated mice, DN1 T cells were activated and proliferated in the draining lymph nodes and spleen of mice vaccinated with the attenuated Mtb strain (**Fig 5D-F**). In line with our findings with BCN, no proliferation was seen in p25 T cells (**Suppl Fig 4B-D**). These results suggest that incorporation of MA in a nanocarrier vaccine allows for the mimicry of the lipid antigen persistence induced by vaccination with an attenuated Mtb vaccine.

MA persists within alveolar macrophages in the lung

To identify the location of antigen persistence, we determined the biodistribution of BCN after IT vaccination using BCN loaded with the hydrophobic dye DiD (DiD-BCN). Since DiD is a lipophilic dye which, like MA, stably loads into the BCN bilayers, we could track the localization of BCN without having to perform additional chemical modifications. We found that IT-administered DiD-BCN were located almost exclusively within the lung at all time points tested (4 h, 24 h, 48 h, 6 days, and 6 weeks) (**Fig 6A**). We were able to detect fluorescence through both *in vivo* imaging system (IVIS) and flow cytometry at 6-weeks post-vaccination that was significantly above unvaccinated control (**Fig 6A, B**), despite DiD quickly losing fluorescence upon mixture into aqueous environments [36], suggesting that some DiD-BCN structure remained intact at this time point. Flow cytometric

analysis (**Suppl Fig 5**) revealed that DiD BCN was primarily found within alveolar macrophages (AMs) at both early and late time points with neutrophils and DCs showing some fluorescence at early time points (4 and 24 h) (**Fig 6C**).

We next tested whether MA likewise persisted primarily within AMs. 6-weeks after vaccination, we enriched for AMs using a column-based magnetic cell isolation system containing SiglecF antibody, which is highly expressed on AMs and a small population of eosinophils in the lung (**Fig 6D**). We co-cultured enriched AMs or the flow through with or without hCD1Tg-expressing BMDCs and DN1 T cells (**Suppl Fig 6**). While both enriched AM and flow through fractions from MA-BCN vaccinated mice co-culture with BMDCs could activate DN1 T cells (**Fig 6E**), co-culture with enriched AMs led to significantly higher DN1 T cell activation, suggesting AMs are the primary location of MA persistence. The ability of flow through to likewise lead to DN1 T cell activation suggests MA may also persist in other cell types or could be attributed to residual AMs in flow through. Furthermore, AMs could not themselves present the MA to DN1 T cells, noted by the lack of T cell activation in the absence of BMDCs, suggesting that MA may be transferred from AMs to DCs for presentation. In fact, CD1b is not expressed on AMs [19, 37] and therefore the presence of CD1b-expressing DCs may be necessary both *in vitro* and *in vivo* for T cell activation to occur. These data indicate that MA-BCN mainly persists within AMs in the lung after IT vaccination and CD1b-expressing DCs is required for antigen presentation and activation of MA-specific T cells.

Vaccination leads to DN1 T cell differentiating into T follicular helper-like T cells

To study the memory phenotype of DN1 T cells after MA-BCN vaccination, we constructed a mixed DN1 bone marrow (BM) chimera mouse model (DN1-hCD1Tg), since adoptively transferred DN1 T cells in unvaccinated mice cannot survive long term (**Fig 7A**). After vaccination, the percent of $CD44^+$ DN1 T cells in the blood increased quickly and remained high until the last time point of 40 days (**Fig 7B**). Various classical memory T cell subsets have been characterized 1 month after initial antigen-encounter, we therefore determined the DN1 T cell phenotype 6-weeks post vaccination. We found that the most prevalent memory population within DN1 T cells were $CD44^+CD62L^+$, markers

used to define central memory T cells, particularly in the LN (**Fig 7C** and **Supp Fig 7A**). To characterize the memory DN1 T cells, we performed RNA-seq analysis on sorted CD44$^+$CD62L$^+$ (memory) and CD44$^-$CD62L$^+$ (naïve) DN1 T cells from LNs of MA-BCN vaccinated DN1-hCD1Tg BM chimeric mice. We found that memory and naïve DN1 T cells clustered separately after principal component analysis (PCA), despite these samples coming from the same animals (**Suppl Fig 7B**). A total of 995 differentially expressed genes (DEGs) were identified of which 542 upregulated and 453 downregulated in the memory subset (**Suppl Fig 7C**). Next, we determined which T cell population memory DN1 T cells most resembled. Toward this end, we obtained data from ImmGen database (including terminally differentiated effector (Te), memory precursor (Tmp), central memory (Tcm), effector memory (Tem), regulatory (Treg))[42], and two additional publications (follicular helper (Tfh)[43], exhausted (Texh)[44]) and compared the respective DEG lists. Using PCA, we found that memory DN1 T cells clustered most closely to Tfh, Treg, and Texh (**Fig 7D**) and when this subset is isolated, most closely to Tfh cells (**Fig 7E**).

Within the DEGs, we noted upregulation of key Tfh cell transcription factors BCL6 and TCF1 (*TCF7*), chemokines CXCR5 and CXCR4 and surface receptor ICOS, important for Tfh cell migration into the germinal center, and adaptor protein SAP (*SH2D1A*), important for T cell-dependent B cell immunity (**Fig 7F**)[45]. Inhibitory receptors PD-1 (*PDCD1*), TIGIT, CTLA4, and transcription factor EOMES, which are known to be expressed on Tfh cells were also upregulated. In addition, we observed downregulation of transcription factor KLF2, which can inhibit Tfh cell differentiation, and CCR7 and IL-7R, receptors known to have decreased expression in T$_{FH}$ cells. Consistent with our transcriptome analysis, we found increased percentage of CXCR5$^+$PD-1$^+$ DN1 T cells in MA-BCN vaccinated mice 6 weeks post-immunization (**Fig 7G, H**). Both our transcriptomic and experimental data highlighted that memory, but not naïve DN1 T cells, were proliferating, given the enrichment for the mitotic cell cycle process (**Fig 7I**) and increased KI67$^+$ percentage within this population (**Fig 7J**). MA-BCN vaccination thus led to the differentiation of DN1 T cells into a proliferating Tfh-like T cell population.

Discussion

Research on the development of subunit vaccines containing lipid antigens is limited. This is in part due to a lack of research into appropriate lipid vaccine formulations and their associated properties. Some early evidence suggests that lipid vaccines could contribute to protection in Mtb challenge in guinea pig model [38, 39], which naturally express group 1 CD1 molecules [40]. Vaccination with a formulation composed of total Mtb lipid encapsulated within liposomes with either dimethyl-dioctadecyl-ammonium (DDA) and/or QS-21 adjuvants showed lower bacterial burden and pathology in lung and spleen compared to unvaccinated controls [38]. In another study, two Mtb lipids (diacylated sulfoglycolipids and the phosphatidylinositol dimannosides) encapsulated into liposomes formulated with DDA and trehalose 6,6'-dibehenateled (TDM) to a decreased bacterial burden in the spleen and overall improved pathology in lung and spleen [39]. This work motivated our current study, where we used hCD1Tg mice that express human group 1 CD1 molecules, including CD1b and MA-specific (DN1), as well as p25-specific T cells to investigate the use of nanocarriers coloaded with both Mtb lipid and protein antigens during vaccination (**Fig 8**).

We found that PEG-*b*-PPS BCN could effectively encapsulate MA as well as the protein antigen Ag85B to elicit activation of antigen-specific T cells in vaccinated mice (**Fig 4**). To our knowledge, this is the first time a dual protein and lipid loaded nanocarrier has been shown to effectively stimulate both lipid and peptide antigen-specific T cells simultaneously *in vivo*. PEG-*b*-PPS BCN can encapsulate both hydrophobic and hydrophilic molecules, making it an excellent platform for subunit vaccine delivery as both antigens and adjuvants can be co-loaded and delivered to the same cellular targets. Prior work on TB vaccines has shown that the inclusion of several protein antigens spanning the Mtb life cycle along with adjuvants with multiple boosts is paramount in the development of a vaccine that can provide effective protection against Mtb [30, 41]. As we have established that PEG-*b*-PPS BCN can easily incorporate a variety of payloads, future work will involve determining the optimal antigen and adjuvant combination for providing protection in a Mtb challenge experiment.

As several formulations of PLGA-NP have been approved for clinical uses [42], we compared MA encapsulated by either BCN or PLGA-NP and found that while PLGA appeared to offer greater, yet less controllable stimulation *in vitro* (**Fig 2**), BCN exhibited significantly greater efficacy *in vivo* (**Fig 2**). The result is not completely unexpected as a more fragile PLGA-NP could allow for fast intracellular release and antigen presentation *in vitro* while also leading to significant antigen loss prior to APC interaction *in vivo* (**Fig 3**) [43]. Previous studies have documented PLGA's innate pro-inflammatory effects, while other studies have highlighted the absence and suppression of inflammatory response [44-46]. As PLGA has not been previously used for lipid antigen delivery via IT vaccination, our findings suggest that PLGA may have limited effectiveness in such a vaccine system, although alternative formulations or polymer compositions could potentially yield improved results. In contrast and as previously observed for PEG-*b*-PPS nanocarriers [23], we showed that unloaded BCN have little inflammatory effect (**Fig 2**), allowing their immunomodulation to be dictated by the properties of the loaded antigen and adjuvant. Thus, PEG-*b*-PPS is a promising polymer for the formulation of subunit vaccines.

We proceeded with a pulmonary (IN or IT) vaccination approach in this study based on emerging evidence that mucosal vaccination can induce a stronger immune response in the lung against Mtb in humans [47] and improve protection following vaccination with nanocarriers in mice [48]. We found that in the pulmonary route of vaccination, BCN primarily were taken up and remained within alveolar macrophages in the lung (**Fig 6**). In fact, the ability of nanocarriers to target alveolar macrophages or other APCs may contribute to their distinct capacity for inducing antigen persistence, which was not seen with cellular carriers such as MA-pulsed BMDCs (**Fig 5**). Alveolar macrophages are the first line of defense of the lungs, responsible for keeping alveoli sterile by taking up inhaled particles and pathogens through phagocytosis [49-51]. Targeting these cells may be particularly beneficial in a TB vaccine design since they may also be the main site of Mtb latency [52]. While MA remained within alveolar macrophages in the lung, T cell activation appeared to occur prominently within the draining lymph nodes (**Fig 2**), suggesting that either MA was transported from the lungs to lymph nodes where the presentation took place or T cells activated in the lung migrated to the lymph

nodes. The first option seems most likely, as CD1b-expressing DCs are more prevalent in the lymph

nodes [53]. Furthermore, in Mtb infection models, it has been previously shown that apoptotic

macrophage and vesicles could deliver mycobacterial antigens to DCs that then activate cognate T

cells [54, 55]. Similarly, we showed that alveolar macrophages from MA-BCN vaccinated mice could

only lead to T cell activation in the presence of BMDCs from hCD1Tg mice (**Fig 6**).

We demonstrated MA persistence in the lung 6 weeks post-vaccination using an *in vivo* antigen

presentation assay with adoptive transfer of naïve MA-specific T cells. Sustained release of antigen

through vaccination is thought to be key in the development of a robust adaptive immune response [56,

57]. Recently, spike protein was found within the draining lymph nodes of vaccinated human subjects

60 days after COVID-19 vaccination with mRNA-1273 or BNT162b2 [58]. In mice, pulmonary

vaccination with adenovirus expressing influenza nucleoprotein supported long term maintenance of

CD8 T_{RM} (resident memory) cells [59]. Unlike in this work, antigen persistence did not occur after

subcutaneous vaccination, suggesting a distinct mechanism of antigen archiving. In a different vaccine

model, subcutaneous administration of ovalbumin paired with adjuvants polyI:C and anti-CD40 led to

antigen archiving by LECs [11, 12]. Similar to our findings, antigen-specific T cell activation required

cross-presentation by DCs [11, 60]. Both LECs and FDCs retain antigens within a non-degradative

compartment [11, 61], suggesting that residual antigens in our model may similarly remain in the

endosome within undigested BCN. We have previously demonstrated the enhanced stability of BCN

within cell endosomes *in vitro* for extensive periods of time [22]. This possibility is corroborated by

the finding that DiD, which is easily quenched in aqueous solution [36], remained fluorescent 6 weeks

post-vaccination (**Fig 6**). The lack of persistence for Ag85B could be due to its leakage out from the

aqueous pores of BCN, which we have observed for other proteins [29]. Peptide antigens may be

modified to include a hydrophobic tail for stable retention within BCN and therefore allow for peptide

antigen persistence as well. MA's extreme hydrophobic surface, which correlates with its low

permeability as a part of Mtb cell wall, may also contribute to its prolonged persistence *in vivo* [62].

After infection and vaccination, peptide antigen persistence has been shown to occur through

archiving by binding to CD21 receptors on follicular dendritic cells (FDCs) in lymph nodes [3, 4]. MA

antigen persistence may likewise be a natural phenomenon that can occur after infections, although its

detection may be difficult given the lack of available assays to detect very small amounts of lipid

antigen. T cells may, in fact, be more sensitive than any commercially available tool. While we

determined that MA persists within AMs following pulmonary vaccination, this localization is likely

dependent on the vaccination route. In the case of SC and IV routes, significant exposure of AMs to

MA-BCN is unlikely to occur. Consequently, the exact localization of MA after vaccination via SC,

IV, or attenuated Mtb remains to be determined, and it is plausible that other types of macrophages

may be involved in this process. Additionally, we found that different forms of encapsulation (such as

BCN, MC, attenuate Mtb) can induce antigen persistence, DC internalization alone was not sufficient

to replicate this phenomenon (**Fig 5**). Thus, the question still remains regarding the necessary

properties of the delivery vector for the persistence of antigens to occur. Overall, MA antigen

persistence is a robust phenomenon which occurs independently of the vector and vaccine route of

administration, and thus may play a role in the efficacy of a variety of vaccine strategies. Additional

work will focus on whether lipid antigen persistence is found across the lipid family or specific to

lipids with certain properties.

Antigen persistence can also occur in chronic infections in which it is known to induce T cell

exhaustion [63]. Despite differentiating in an environment of persistent antigen exposure, memory

DN1 T cells exhibited a gene expression profile more similar to that of Tfh cells rather than exhausted

T cells (**Fig 7**). In particular, memory DN1 T cells upregulated the expression of TCF7 and BCL6

which are known to sustain a non-exhausted phenotype in CD8 T cells during chronic viral infections

[64, 65]. Previous studies on antigen archiving have suggested that the mere persistence of antigen is

insufficient for exhaustion to develop, and the inflammatory context likely plays a role in T cell

differentiation [11]. In many subunit vaccines, virulence factors are used as antigens to target and

suppress the pathogen's capacity to infect the host. The persistence of lipid virulence factors could

raise concerns regarding the safety of lipid vaccines. However, it is worth noting that otherwise toxic

compounds such as the lipid TDM have been successfully incorporated into numerous vaccines

without any known safety concerns to date [66-71]. The absence of reported safety issues may be

attributed to limited research or the administration of low doses of TDM. It is important to highlight that while MA play a role in immune evasion, MA without additional functional groups has not been shown to elicit similar toxic effects as TDM [62].

The use of memory T cells in the setting of antigen persistence may be debatable, given that memory response has historically referred to the adaptive immune response in the setting of antigen clearance [72]. However, given that the persistence of MA occurs even in the setting of attenuated Mtb vaccination, Mtb lipid-specific long-term adaptive immunity will likely naturally exist in an antigen persistent environment. We found that DN1 T cells in this setting expressed key markers of Tfh cells such as CXCR5, PD-1, BCL6, and TCF1 (**Fig 7**). T cells differentiate into Tfh cells through the expression of master transcription factor regulator BCL6, which directs the upregulation of chemokine receptors CXCR5 and CXCR4 and thus migration into the germinal center[73, 74]. In the germinal center, Tfh cells provide help to B cells for affinity maturation and class switching. Other unconventional T cells are known to behave similarly such as CD1d-restricted NKT cells, which can acquire a Tfh phenotype and provide B cell help through both cognate and non-cognate interactions[75]. Additional work will look at whether DN1 or other group 1 CD1-restricted T cells can functionally act as Tfh cells to provide B cell help, as may be suggested by the existence of IgG antibodies against MA and glycolipid lipoarabinomannan in TB patients [76-78].

While persistence of protein antigen has been shown to improve the quality of protective immunity [7, 11, 79, 80], it is not yet known how lipid antigen persistence may affect the protective efficacy of lipid-specific T cells. At the very least, antigen persistence increases the probability of T cell/DC encounters by increasing the time that the antigen is available. Additional work will need to address the function of lipid-specific T cells differentiated in antigen persistent environments. As the ultimate goal of this work is to improve the efficacy of existing subunit vaccines, future work will assess the protective capacity of lipid antigen vaccines paired with additional protein antigens and adjuvants.

Materials and Methods

Ethics statement

This study was carried out in accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. The protocol was approved by the Institutional Animal Care and Use Committee of Northwestern University (Protocol number: IS000011717).

Poly (ethylene glycol)-block-poly (propylene sulfide) polymer synthesis

Poly (ethylene glycol)$_{17}$-block-poly-(propylene sulfide)$_{75}$ (PEG$_{17}$-*b*-PPS$_{75}$) copolymer was synthesized as previously reported [81, 82]. Briefly, monomethoxy PEG (molecular weight=750 Da) was functionalized with mesyl chloride and substituted with thioacetate to produce a PEG acetate initiator for the polymerization for PPS.

Bicontinuous nanosphere fabrication via flash nano precipitation

BCN were produced using FNP technique as previously described[81]. In brief, PEG$_{17}$-*b*-PPS$_{75}$ BCN polymer and any hydrophobic cargo was dissolved in THF and loaded onto a syringe. The hydrophilic cargo was loaded onto a separate syringe, and both impinged against one another through the confined impingement mixer into scintillation vial. The vial was placed in a vacuum desiccator overnight to remove the organic solvent. NPs were filtered through LH-20 column with PBS as an eluent and suspended in PBS for later use. A natural mixture of Mtb MA consists of α-, keto-, and methoxy forms of MA with chain lengths of 72-86 carbon atom was obtained from Sigma (M4537, Sigma-Aldrich, St. Louis, MO) [83]. MA encapsulation efficiency was analyzed using coumarin derivatization as previously described [21]. MA was loaded at a concentration of 100 µg/mL of total BCN formulation and DiD was loaded at 5 µg/mL total BCN formulation. MA loading capacity in BCNs was 2%. To measure the encapsulation efficiency of Ag85B, Ag85B-MA-BCN were centrifuged at 10,000 g for 10 minutes. The supernatant was collected and concentrated using Amicon® Ultra-4 10K centrifugal filter

devices. To determine the unencapsulated protein concentration, the supernatant was incubated with

Pierce™ BCA Protein assay reagent following the manufacturer protocol and the absorbance was

measured at 660 nm using a SpectraMax M3 multi-mode microplate reader (Molecular Devices, LLC).

The percentage encapsulation efficiency was calculated as a percent of encapsulated protein to the total

amount of protein added. Ag85B was encapsulated at a concentration of 140 μg/mL and loading

capacity was found to be 2.88%.

Poly (lactic-co-glycolic acid nanoparticle (NP) fabrication

PLGA-NP were prepared by the Oil-in-Water single emulsion method. Briefly, PLGA (20 mg in 800

μL dichloromethane) organic phase was emulsified using an ultrasonic processor with aqueous phase

containing 2 mL of polyvinyl alcohol (PVA) solution (2.5% w/v) to form an emulsion. The emulsion

was then added dropwise into 4 mL of stirring 0.5% w/v PVA solution at room temperature to

evaporate the organic solvent. NPs were collected after 4 hours of stirring followed by centrifugation at

10,000 x g for 10 minutes. After centrifugation, NPs were washed twice with cold water to remove

residual PVA and redispersed in PBS. Hydrophilic (Texas red-Dextran) or hydrophobic (MA) loading

was performed by adding the payloads to the aqueous and organic phases, respectively.

Nanocarrier characterization

The nanocarrier size (z-average diameter) and polydispersity was measured using dynamic light

scattering (DLS) using a Nano 300 ZS Zetasizer (Malvern Panalytical, Malvern, U.K.). BCN and

PLGA morphology was visualized with cryogenic transmission electron microscopy (cryoTEM) as

previously detailed [21]. Transmission electron microscopy (TEM) was performed to image BCN

using uranyl acetate as a negative stain as previously reported [22]. BCN aggregate structure and

internal morphology was characterized with small angle X-ray scattering (SAXS). These studies were

performed at the DuPont-Northwestern-Dow Collaborative Access Team beamline at Argonne

National Laboratory's Advanced Photon Source with 10 keV (wavelength $\lambda = 1.24$ Å) collimated X-

rays, as described previously[81].

Confocal imaging

Confocal images of RAW 264.7 macrophages stained with lysosomal dye Lysotracker™ (green) and NucBlue™ stain (blue) following incubation with Texas Red-labeled PLGA-NP and BCN (red) for 8 hr. The cells were then imaged within a humidified chamber using a 63X oil-immersion objective on a SP5 Leica Confocal Microscope using HyD detectors and lasers. Data analysis was performed using ImageJ software.

Mouse strains

Human CD1 transgenic mice (hCD1Tg) in C57BL/6 background or MHC II-deficient background [19] and CD1b-restricted MA-specific TCR transgenic mice on Rag$^{-/-}$ background (DN1Tg/hCD1Tg/Rag$^{-/-}$) [20] were generated in our lab as previously described. P25-specific TCR transgenic mice (Jackson lab) were crossed onto Rag$^{-/-}$ background. Both males and females were used.

Antibodies and flow cytometry

For cell surface staining, cells were pre-incubated with 2.4G2 Fcγ RII/RIII blocking mAb for 15 minutes and then stained with the appropriate combinations of mAbs diluted in HBSS + 2% FBS for 30 minutes at 4°C. Cells were analyzed on a BD FACS CantoII, or Cytek Aurora Spectral Cytometer and data were processed with FlowJo software (TreeStar). The complete list of antibodies use is as follows: anti-human Vβ5.1 (LC4, Invitrogen), anti-TCRβ (H57-597, BioLegend), anti-CD44 (IM7, BioLegend), anti-Vβ11 (KT11, BioLegend), anti-CD45.2 (104, BioLegend), anti-CD45.1 (A20, BioLegend), anti-SiglecF (E50-2440, BD Bioscience), anti-CD45 (30-F11, BioLegend), anti-CD11c (N418, BioLegend), anti-CD11b (M1/70, BioLegend), anti-Ly6G (1A8, BioLegend), anti-CD19 (6D5, BioLegend), anti-NK1.1 (PK136, BioLegend), anti-CD3 (17A2, BioLegend), Zombie Red (BioLegend), anti-mouse CD185/CXCR5 (L138D7, Biolegend), anti-mouse CD279/PD-1 (29F.1A12, Biolegend), and anti-Ki67 (SolA15, Invitrogen).

***In vitro* nanoparticle and lipid titration with MA-specific DN1 T cells**

Nanoparticle and lipid titrations were performed as previously described [21]. In brief, hCD1Tg MHC

II$^{-/-}$ bone marrow-derived dendritic cells (BMDCs) were pulsed with varying quantities of nanoparticle

of sonicated MA and then co-cultured for 48 hours with DN1 T cells isolated from lymph nodes (axial,

brachial, submandibular, inguinal) of DN1Tg/hCD1Tg/Rag$^{-/-}$ mice. The supernatant was used for IFN-

γ enzyme-linked immunosorbent assay (ELISA) performed as described previously [21] and cells were

stained for activation.

***In vitro* nanoparticle and lipid titration with human cells via ELISPOT assay**

Hydrophobic polyvinylidene fluoride 96-well plates (Millipore, Bedford, MA) were coated with anti-

human IFN-γ mAb 1-D1K (Mabtech, Cincinnati, OH) diluted 1:400 in PBS (Gibco, Waltham, MA)

and incubated at 4°C overnight. The following day, mAb was removed and each plate was washed and

incubated at room temperature for 2 hours in the presence of media. Human monocyte-derived

dendritic cells and MA-specific M11 T cells were seeded 50,000 cells and 2,000 cells per well,

respectively, together with serially diluted MA and NPs. After 16-hour incubation, plates were washed

and incubated with biotinylated anti-human IFN-g (7-B6-1), diluted 1:3000 in PBS + 0.5% FBS, for 4

hours. Plates were then washed with PBS 5 times and incubated for 2 hours with ExtrAvidin-Alkaline

phosphatase diluted 1:1000 in PBS + 0.5% FBS. Spots were visualized after incubation with the

BCIP/NBT substrate for up to 20 minutes and then imaged with ImmunoSpot reader v.2.0.

Mouse vaccination

For both intranasal (IN) and intratracheal (IT) vaccination, mice were anesthetized using inhaled

isoflurane. For IN vaccination, 25 μL of liquid was administered at the nostrils of the mice for a total

of 1 μg of MA per mouse. For IT vaccination, mice were administered 25 or 50 μL of liquid for a total

of 2.5 μg of MA with or without 2.0 μg of Ag85B. For BMDC vaccinations, hCD1Tg BMDCs were

cultured in GM-CSF and IL-4 for 5 days[84] and pulsed with MA-BCN (5 μg/mL of MA) or free MA

(10 μg/mL of MA) for 24 hours. Cells were washed and resuspended in PBS and IT vaccinated at a

dose of 1x10^6 BMDCs/mouse. Intravenous vaccination was administered by tail vein injection.

Subcutaneous vaccination was administered between the shoulders over the neck portion of the mouse.

Vaccination with attenuated Mtb

 Attenuated Mtb strain (H37Rv Δ*panCD* Δ*leuCD* referred to as mc^2 6206) was kindly provided by

Jacobs' Laboratory at Albert Einstein College of Medicine. Bacteria were grown as previously

described in supplemented Middlebrook 7H9 media [35, 85]. Mice were vaccinated subcutaneously

with 1x10^6 CFU in 100 μL of PBS. Mycobacterial burden quantification was performed by plating

serial dilutions of lung or spleen homogenate on Middlebrook 7H11 agar plates [35, 85].

T cell Adoptive transfer and cell preparation

DN1 or p25 T cells were isolated from lymph nodes of DN1Tg/hCD1Tg/Rag$^{-/-}$ or P25Tg/Rag$^{-/-}$ mice,

respectively. Cells were labeled with CellTrace™ Violet reagent (Invitrogen™) as per manufacturer's

instructions and 3-5x10^6 cells were injected into hCD1Tg mice either 1 day before vaccination (short

term) or 6 weeks post-vaccination (long term). On day 7 post-adoptive transfer, draining lymph nodes

(axial, brachial, mediastinal), lung, and spleen were obtained. Single-cell suspensions were prepared

by mechanical disruption in HBSS/2% FBS. The lung was digested with collagenase IV (1mg/ml;

Sigma-Aldrich) and DNase I (30μg/ml; Sigma- Aldrich) for 30 minutes at 37°C before the disruption.

Biodistribution imaging

BCN were loaded with DiD dye (Thermo Fisher) and administered to mice IT at 4 hours, 24 hours, 48

hours, 6 days, and 6 weeks prior to the final time point, at which all mice were sacrificed and analyzed

together. Lung, spleen, and draining lymph nodes were harvested and imaged using a near-IR *in vivo*

Imaging System (IVIS; Center for Advanced Molecular Imaging, Northwestern University). The

single-cell suspension was then prepared from indicated organs and cells were stained for flow

cytometric analysis.

Alveolar macrophage enrichment and T cell co-culture

hCD1Tg mice were immunized IT with either BCN or MA-BCN (2.5 μg of MA/mouse). 6 weeks

later, lungs were isolated and single-cell suspension was stained with anti-SiglecF-PE (E50-2440, BD

547   Bioscience). SiglecF-positive cells were enriched using anti-PE MultiSort Kit (Miltenyi Biotec). $1x10^5$

548   flow through and AM enriched fractions were co-cultured with $1x10^5$ hCD1Tg BMDCs and $3x10^5$

549   DN1 T cells for 48 hours and cells were stained for the expression of activation markers.

Generation and vaccination of BM chimeras

Bone marrow cells from DN1Tg/hCD1Tg/Rag$^{-/-}$ (CD45.2) and hCD1Tg mice (CD45.1) were depleted of

mature T cells using anti-Thy-1.2 (AT83.A-6) plus rabbit complement (Cedarlane Laboratories). Equal

numbers of cells ($5x10^6$) were adoptively transferred into hCD1Tg (CD45.1) mice irradiated with 900

rads. After 6 weeks, BM chimeric mice were intratracheally vaccinated with either 2.5 μg of MA in

MA-BCN or equivalent volume of PBS and the activation status of DN1 T cells were monitored in the

blood 1-2 times per week until experiment completion.

RNA-Seq analysis

CD44$^+$CD62L$^+$ and CD44$^-$CD62L$^+$ DN1 T cells were sorted from LNs of DN1-hCD1Tg BM chimera

mice vaccinated with MA-BCN at 6 weeks post vaccination by BD FACS Aria with purity >98%.

RNA was extracted with RNAeasy® mini kit (Qiagen). Libraries were generated by using the Illumina

Truseq preparation kit and sequenced on HiSeq4000. Reads were analyzed with Ceto pipeline

(https://github.com/ebartom/NGSbartom) using STAR and HTseq for alignment on mm10 mouse

genome and reading counting, as described previously [86]. Paired differential expression analysis was

performed using DESeq2 to account for original mouse sourcing[87]. All downstream analysis was

performed in R and figured generated using ggplot. Gene enrichment analysis was performed using

Metascape [88]. Comparisons to other T cell profiles were performed using raw transcriptome data

from ImmGen database[89], and two additional publications for follicular helper (T$_{FH}$)[90], exhausted

(Texh)[91]. Analysis was performed as described above to obtain gene expression fold change values

relative to internal naïve condition.

Data accessibility

The data discussed in this publication have been deposited in NCBI's Gene Expression Omnibus and are accessible through GEO Series accession number GSE226075 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE226075).

Statistical analysis

Statistical analyses were performed using Prism software 9 (GraphPad software, Inc.). Multi-group comparisons were done using two-way analysis of variance (ANOVA) with significance determined by Tukey's multiple comparison. Where appropriate, outliers were identified through Grubbs method with alpha=0.05. Statistically significance denoted as ns = not significant, *$p<0.05$, **$p<0.01$, ***$p<0.001$, and ****$p< 0.0001$.

<div align="center">**Acknowledgments**</div>

We thank Ying He for her support through maintaining laboratory reagents and equipment, Elizabeth Bartom for use of her RNASeq analysis pipeline, and Michael Kleyman for his advice on comparing gene expression among multiple T cell populations. We would like to thank the Center for Advanced Molecular Imaging (CAMI) for their help in performing IVIS and The Northwestern University's Atomic and Nanoscale Characterization Experimental Center (NUANCE) Keck-II facility for their help with NP characterization. Ag85B from *Mycobacterium tuberculosis* H37Rv was obtained through BEI Resources, NIAID, NIH. SAXS experiments were performed at the DuPont-Northwestern-Dow Collaborative Access Team (DND-CAT) located at Sector 5 of the Advanced Photon Source. DND-CAT is supported by Northwestern University, The Dow Chemical Company, and DuPont de Nemours, Inc. This research used resources of the Advanced Photon Source; a U.S. Department of Energy (DOE) Office of Science User Facility operated for the DOE Office of Science by Argonne National Laboratory under Contract No. DE-AC02-06CH11357. Data was collected using an instrument funded by the National Science Foundation under Award No. 0960140.

Funding:
This work was supported by the following grans:

National Institutes of Health grant R01AI057460 (CRW), R01AI145345 (CRW, EAS)

National Institutes of Health grant F30AI157314 (EM)

National Institutes of Health grant T32GM008152 (EM)

National Institutes of Health grant R01AI146072 (CS)

Competing interests: Authors declare that they have no competing financial interests.

Data and materials availability:

All data are available in the main text or the supplementary materials. Derived data supporting the

findings of this study are available from the corresponding author CRW and EAS on request.

Generated transgenic mice and nanoparticles are available upon request upon signing of Material

Transfer Agreement (MTA).

Nonstandard Abbreviations:

Mtb, *Mycobacterium tuberculosis*; MA, mycolic acid; BCN, bicontinuous nanospheres; MA-BCN,

mycolic acid loaded bicontinuous nanospheres; FDCs, follicular dendritic cells; LECs, lymphocytic

endothelial cells; hCD1Tg, human CD1 transgenic mouse model; APCs, antigen-presenting cells;

PEG-b-PPS, poly(ethylene glycol)-block-poly(propylene sulfide); MA-MC, mycolic acid loaded

micellar nanocarrier; FNP, flash nanoprecipitation; PLGA, poly(D,L-lactide-co-glycolide); DLS,

dynamic light scattering; PDI, polydispersity index; cryo-TEM, cryogenic transmission electron

microscopy; SAXS, small angle X-ray scattering; PLGA-NP, poly(D,L-lactide-co-glycolide)

nanoparticle; MA-PLGA, mycolic acid loaded poly(D,L-lactide-co-glycolide); IN, intranasal; IT,

intratracheal; Ag85B-MA-BCN, Ag85B and mycolic acid loaded bicontinuous nanosphere; BMDCs,

bone marrow-derived dendritic cells; DiD-BCN, DiD loaded bicontinuous nanosphere; AMs, alveolar

macrophages.

References

1.   Karch CP, Burkhard P. Vaccine technologies: From whole organisms to rationally designed

protein assemblies. Biochem Pharmacol. 2016;120:1-14. Epub 20160506. doi:

10.1016/j.bcp.2016.05.001. PubMed PMID: 27157411; PubMed Central PMCID: PMCPMC5079805.

2.   Legoux F, Salou M, Lantz O. Unconventional or Preset alphabeta T Cells: Evolutionarily

Conserved Tissue-Resident T Cells Recognizing Nonpeptidic Ligands. Annu Rev Cell Dev Biol.

2017;33:511-35. Epub 20170629. doi: 10.1146/annurev-cellbio-100616-060725. PubMed PMID:

28661722.

3.   Heesters BA, Myers RC, Carroll MC. Follicular dendritic cells: Dynamic antigen libraries.

Nature Publishing Group; 2014. p. 495-504.

4.   Hirosue S, Dubrot J. Modes of antigen presentation by lymph node stromal cells and their

immunological implications. Frontiers in Immunology. 2015;6. doi: ARTN 446

10.3389/fimmu.2015.00446. PubMed PMID: WOS:000361632700001.

5.   Takamura S, Roberts AD, Jelley-Gibbs DM, Wittmer ST, Kohlmeier JE, Woodland DL. The

route of priming influences the ability of respiratory virus-specific memory CD8+ T cells to be

activated by residual antigen. Journal of Experimental Medicine. 2010;207(6):1153-60. doi:

10.1084/jem.20090283.

6.   Kim TS, Hufford MM, Sun J, Fu YX, Braciale TJ. Antigen persistence and the control of local

T cell memory by migrant respiratory dendritic cells after acute virus infection. Journal of

Experimental Medicine. 2010;207(6):1161-72. doi: 10.1084/jem.20092017.

7.   Demento SL, Cui W, Criscione JM, Stern E, Tulipan J, Kaech SM, et al. Role of sustained

antigen release from nanoparticle vaccines in shaping the T cell memory phenotype. Biomaterials.

2012;33(19):4957-64. Epub 20120406. doi: 10.1016/j.biomaterials.2012.03.041. PubMed PMID:

22484047; PubMed Central PMCID: PMCPMC5724530.

8.   Zammit DJ, Turner DL, Klonowski KD, Lefrançois L, Cauley LS. Residual Antigen

Presentation after Influenza Virus Infection Affects CD8 T Cell Activation and Migration. Immunity.

2006;24(4):439-49. doi: 10.1016/j.immuni.2006.01.015.

9.   Woodland DL, Kohlmeier JE. Migration, maintenance and recall of memory T cells in

peripheral tissues. Nat Rev Immunol. 2009;9(3):153-61. doi: 10.1038/nri2496. PubMed PMID:

19240755.

10.   Vokali E, Yu SS, Hirosue S, Rincon-Restrepo M, F VD, Scherer S, et al. Lymphatic endothelial

cells prime naive CD8(+) T cells into memory cells under steady-state conditions. Nat Commun.

2020;11(1):538. Epub 20200127. doi: 10.1038/s41467-019-14127-9. PubMed PMID: 31988323;

PubMed Central PMCID: PMCPMC6985113.

11.   Tamburini BA, Burchill MA, Kedl RM. Antigen capture and archiving by lymphatic

endothelial cells following vaccination or viral infection. Nature Communications. 2014;5(1):3989-.

doi: 10.1038/ncomms4989.

12.   Walsh SM, Sheridan RM, Lucas ED, Doan TA, Ware BC, Schafer J, et al. Molecular tracking

devices quantify antigen distribution and archiving in the murine lymph node. Elife. 2021;10. Epub

20210412. doi: 10.7554/eLife.62781. PubMed PMID: 33843587; PubMed Central PMCID:

PMCPMC8116055.

13.   Reinink P, Van Rhijn I. Mammalian CD1 and MR1 genes. Immunogenetics. 2016;68(8):515-

23. Epub 20160728. doi: 10.1007/s00251-016-0926-x. PubMed PMID: 27470004; PubMed Central

PMCID: PMCPMC5002277.

14.   Joosten SA, Ottenhoff THM, Lewinsohn DM, Hoft DF, Moody DB, Seshadri C. Harnessing

donor unrestricted T-cells for new vaccines against tuberculosis. 2019. p. 3022-30.

15.   Morgun E, Cao LC, Wang C-R. Role of Group 1 CD1-restricted T Cells in Host Defense and

Inflammatory Diseases. Critical Reviews in Immunology. 2021. doi:

10.1615/critrevimmunol.2021040089.

16.   Busch M, Herzmann C, Kallert S, Zimmermann A, Hofer C, Mayer D, et al.

Lipoarabinomannan-responsive polycytotoxic T cells are associated with protection in human

tuberculosis. American Journal of Respiratory and Critical Care Medicine. 2016;194(3):345-55. doi:

10.1164/rccm.201509-1746OC.

17. Lopez K, Iwany SK, Suliman S, Reijneveld JF, Ocampo TA, Jimenez J, et al. CD1b Tetramers

Broadly Detect T Cells That Correlate With Mycobacterial Exposure but Not Tuberculosis Disease

State. Frontiers in Immunology. 2020;11:199-. doi: 10.3389/fimmu.2020.00199.

18. Montamat-Sicotte DJ, Millington KA, Willcox CR, Hingley-Wilson S, Hackforth S, Innes J, et

al. A mycolic acid-specific CD1-restricted T cell population contributes to acute and memory immune

responses in human tuberculosis infection. Journal of Clinical Investigation. 2011;121(6):2493-503.

doi: 10.1172/JCI46216.

19. Felio K, Nguyen H, Dascher CC, Choi H-J, Li S, Zimmer MI, et al. CD1-restricted adaptive

immune responses to Mycobacteria in human group 1 CD1 transgenic mice. The Journal of

experimental medicine. 2009;206(11):2497-509. doi: 10.1084/jem.20090898.

20. Zhao J, Siddiqui S, Shang S, Bian Y, Bagchi S, He Y, et al. Mycolic acid-specific T cells

protect against Mycobacterium tuberculosis infection in a humanized transgenic mouse model. eLife.

2015;4. doi: 10.7554/eLife.08525.

21. Shang S, Kats D, Cao L, Morgun E, Velluto D, He Y, et al. Induction of Mycobacterium

Tuberculosis Lipid-Specific T Cell Responses by Pulmonary Delivery of Mycolic Acid-Loaded

Polymeric Micellar Nanocarriers. Frontiers in Immunology. 2018;9:2709-. doi:

10.3389/fimmu.2018.02709.

22. Bobbala S, Allen SD, Yi S, Vincent M, Frey M, Karabin NB, et al. Employing bicontinuous-to-

micellar transitions in nanostructure morphology for on-demand photo-oxidation responsive cytosolic

delivery and off-on cytotoxicity. Nanoscale. 2020;12(9):5332-40. doi: 10.1039/c9nr10921h.

23. Burke JA, Zhang X, Bobbala S, Frey MA, Bohorquez Fuentes C, Freire Haddad H, et al.

Subcutaneous nanotherapy repurposes the immunosuppressive mechanism of rapamycin to enhance

allogeneic islet graft viability. Nat Nanotechnol. 2022. Epub 20220117. doi: 10.1038/s41565-021-

01048-2. PubMed PMID: 35039683.

24. Du F, Liu YG, Scott EA. Immunotheranostic Polymersomes Modularly Assembled from

Tetrablock and Diblock Copolymers with Oxidation-Responsive Fluorescence. Cell Mol Bioeng.

2017;10(5):357-70. Epub 20170410. doi: 10.1007/s12195-017-0486-7. PubMed PMID: 28989540;

PubMed Central PMCID: PMCPMC5628512.

25.   Scott EA, Stano A, Gillard M, Maio-Liu AC, Swartz MA, Hubbell JA. Dendritic cell activation

and T cell priming with adjuvant- and antigen-loaded oxidation-sensitive polymersomes. Biomaterials.

2012;33(26):6211-9. doi: 10.1016/j.biomaterials.2012.04.060.

26.   Napoli A, Valentini M, Tirelli N, Müller M, Hubbell JA. Oxidation-responsive polymeric

vesicles. Nature Materials. 2004;3(3):183-9. doi: 10.1038/nmat1081.

27.   Allen SD, Bobbala S, Karabin NB, Scott EA. On the advancement of polymeric bicontinuous

nanospheres toward biomedical applications. Nanoscale Horiz. 2019;4(2):258-72. Epub 20181120.

doi: 10.1039/c8nh00300a. PubMed PMID: 32254084.

28.   Saad WS, Prud'Homme RK. Principles of nanoparticle formation by flash nanoprecipitation.

Nano Today. 2016;11(2):212-27. doi: 10.1016/j.nantod.2016.04.006.

29.   Bobbala S, Allen SD, Scott EA. Flash nanoprecipitation permits versatile assembly and loading

of polymeric bicontinuous cubic nanospheres. Nanoscale. 2018;10(11):5078-88. doi:

10.1039/C7NR06779H.

30.   Lin LCW, Chattopadhyay S, Lin JC, Hu CMJ. Advances and Opportunities in Nanoparticle-

and Nanomaterial-Based Vaccines against Bacterial Infections. Wiley-Blackwell; 2018. p. 1701395-.

31.   Khademi F, Derakhshan M, Yousefi-Avarvand A, Tafaghodi M. Potential of polymeric

particles as future vaccine delivery systems/adjuvants for parenteral and non-parenteral immunization

against tuberculosis: A systematic review. Mashhad University of Medical Sciences; 2018. p. 116-23.

32.   Modak M, Bobbala S, Lescott C, Liu YG, Nandwana V, Dravid VP, et al. Magnetic

Nanostructure-Loaded Bicontinuous Nanospheres Support Multicargo Intracellular Delivery and

Oxidation-Responsive Morphological Transitions. ACS Appl Mater Interfaces. 2020;12(50):55584-95.

Epub 20201201. doi: 10.1021/acsami.0c15920. PubMed PMID: 33259182.

33.   Huygen K. The Immunodominant T-Cell Epitopes of the Mycolyl-Transferases of the Antigen

85 Complex of M. tuberculosis. Front Immunol. 2014;5:321. Epub 20140709. doi:

10.3389/fimmu.2014.00321. PubMed PMID: 25071781; PubMed Central PMCID: PMCPMC4089088.

34.   Tamura T, Ariga H, Kinashi T, Uehara S, Kikuchi T, Nakada M, et al. The role of antigenic

peptide in CD4+ T helper phenotype development in a T cell receptor transgenic model. International

Immunology. 2004;16(12):1691-9. doi: 10.1093/intimm/dxh170.

35.   Sampson SL, Dascher CC, Sambandamurthy VK, Russell RG, Jacobs WR, Jr., Bloom BR, et

al. Protection elicited by a double leucine and pantothenate auxotroph of Mycobacterium tuberculosis

in guinea pigs. Infect Immun. 2004;72(5):3031-7. doi: 10.1128/IAI.72.5.3031-3037.2004. PubMed

PMID: 15102816; PubMed Central PMCID: PMCPMC387862.

36.   Invitrogen™. DiD' solid; DiIC18(5) solid (1,1'-Dioctadecyl-3,3,3',3'-

Tetramethylindodicarbocyanine, 4-Chlorobenzenesulfonate Salt).

37.   Barral DC, Brenner MB. CD1 antigen presentation: how it works. Nature Reviews

Immunology. 2007;7(12):929-41. doi: 10.1038/nri2191.

38.   Dascher CC, Hiromatsu K, Xiong X, Morehouse C, Watts G, Liu G, et al. Immunization with a

mycobacterial lipid vaccine improves pulmonary pathology in the guinea pig model of tuberculosis.

Oxford University Press; 2003. p. 915-25.

39.   Larrouy-Maumus G, Layre E, Clark S, Prandi J, Rayner E, Lepore M, et al. Protective efficacy

of a lipid antigen vaccine in a guinea pig model of tuberculosis. Vaccine. 2017;35(10):1395-402. doi:

10.1016/J.VACCINE.2017.01.079.

40.   Eckhardt E, Bastian M. Animal models for human group 1 CD1 protein function. Molecular

Immunology. 2021;130:159-63. doi: 10.1016/j.molimm.2020.12.018.

41.   Schrager LK, Vekemens J, Drager N, Lewinsohn DM, Olesen OF. The status of tuberculosis

vaccine development. Lancet Infect Dis. 2020;20(3):e28-e37. Epub 20200131. doi: 10.1016/S1473-

3099(19)30625-5. PubMed PMID: 32014117.

42.   Wang Y, Qin B, Xia G, Choi SH. FDA's Poly (Lactic-Co-Glycolic Acid) Research Program

and Regulatory Outcomes. Aaps J. 2021;23(4):92. Epub 20210629. doi: 10.1208/s12248-021-00611-y.

PubMed PMID: 34189655.

43.   Ding D, Zhu Q. Recent advances of PLGA micro/nanoparticles for the delivery of

biomacromolecular therapeutics. Materials Science & ENgineering C. 2018. doi:

10.1016/j.msec.2017.12.036.

44.   Barillet S, Fattal E, Mura S, Tsapis N, Pallardy M, Hillaireau H, et al. Immunotoxicity of poly

(lactic-co-glycolic acid) nanoparticles: influence of surface properties on dendritic cell activation.

Nanotoxicology. 2019;13(5):606-22. Epub 20190214. doi: 10.1080/17435390.2018.1564078. PubMed PMID: 30760076.

45.   Ma S, Feng X, Liu F, Wang B, Zhang H, Niu X. The pro-inflammatory response of macrophages regulated by acid degradation products of poly(lactide-co-glycolide) nanoparticles. Eng Life Sci. 2021;21(10):709-20. Epub 20210512. doi: 10.1002/elsc.202100040. PubMed PMID: 34690640; PubMed Central PMCID: PMCPMC8518582.

46.   Nicolete R, dos Santos DF, Faccioli LH. The uptake of PLGA micro or nanoparticles by macrophages provokes distinct in vitro inflammatory response. Int Immunopharmacol. 2011;11(10):1557-63. Epub 20110527. doi: 10.1016/j.intimp.2011.05.014. PubMed PMID: 21621649.

47.   Stylianou E, Paul MJ, Reljic R, McShane H. Mucosal delivery of tuberculosis vaccines: a review of current approaches and challenges. Expert Rev Vaccines. 2019;18(12):1271-84. Epub 20191226. doi: 10.1080/14760584.2019.1692657. PubMed PMID: 31876199; PubMed Central PMCID: PMCPMC6961305.

48.   Ballester M, Nembrini C, Dhar N, de Titta A, de Piano C, Pasquier M, et al. Nanoparticle conjugation and pulmonary delivery enhance the protective efficacy of Ag85B and CpG against tuberculosis. Vaccine. 2011;29(40):6959-66. doi: 10.1016/j.vaccine.2011.07.039. PubMed PMID: WOS:000295300500018.

49.   Fels AO, Cohn ZA. The alveolar macrophage. J Appl Physiol (1985). 1986;60(2):353-69. doi: 10.1152/jappl.1986.60.2.353. PubMed PMID: 3005225.

50.   Bowden DH. The alveolar macrophage. Environ Health Perspect. 1984;55:327-41. doi: 10.1289/ehp.8455327. PubMed PMID: 6376105; PubMed Central PMCID: PMCPMC1568378.

51.   Geiser M. Update on macrophage clearance of inhaled micro- and nanoparticles. J Aerosol Med Pulm Drug Deliv. 2010;23(4):207-17. doi: 10.1089/jamp.2009.0797. PubMed PMID: 20109124.

52.   Pai M, Behr MA, Dowdy D, Dheda K, Divangahi M, Boehme CC, et al. Tuberculosis. Nat Rev Dis Primers. 2016;2:16076. Epub 20161027. doi: 10.1038/nrdp.2016.76. PubMed PMID: 27784885.

53.   Dougan SK, Kaser A, Blumberg RS. CD1 Expression on Antigen-Presenting Cells. 314. Berlin, Heidelberg: Springer Berlin Heidelberg; 2007. p. 113-41.

54.   Espinosa-Cueto P, Magallanes-Puebla A, Castellanos C, Mancilla R. Dendritic cells that

phagocytose apoptotic macrophages loaded with mycobacterial antigens activate CD8 T cells via

cross-presentation. PLoS One. 2017;12(8):e0182126. Epub 20170802. doi:

10.1371/journal.pone.0182126. PubMed PMID: 28767693; PubMed Central PMCID:

PMCPMC5540487.

55.   Schaible UE, Winau F, Sieling PA, Fischer K, Collins HL, Hagens K, et al. Apoptosis

facilitates antigen presentation to T lymphocytes through MHC-I and CD1 in tuberculosis. Nat Med.

2003;9(8):1039-46. Epub 20030720. doi: 10.1038/nm906. PubMed PMID: 12872166.

56.   Silva AL, Soema PC, Slutter B, Ossendorp F, Jiskoot W. PLGA particulate delivery systems

for subunit vaccines: Linking particle properties to immunogenicity. Hum Vaccin Immunother.

2016;12(4):1056-69. Epub 20160111. doi: 10.1080/21645515.2015.1117714. PubMed PMID:

26752261; PubMed Central PMCID: PMCPMC4962933.

57.   Correia-Pinto JF, Csaba N, Alonso MJ. Vaccine delivery carriers: insights and future

perspectives. Int J Pharm. 2013;440(1):27-38. Epub 20120423. doi: 10.1016/j.ijpharm.2012.04.047.

PubMed PMID: 22561794.

58.   Roltgen K, Nielsen SCA, Silva O, Younes SF, Zaslavsky M, Costales C, et al. Immune

imprinting, breadth of variant recognition, and germinal center response in human SARS-CoV-2

infection and vaccination. Cell. 2022;185(6):1025-40 e14. Epub 20220125. doi:

10.1016/j.cell.2022.01.018. PubMed PMID: 35148837; PubMed Central PMCID: PMCPMC8786601.

59.   Uddback I, Cartwright EK, Scholler AS, Wein AN, Hayward SL, Lobby J, et al. Long-term

maintenance of lung resident memory T cells is mediated by persistent antigen. Mucosal Immunol.

2021;14(1):92-9. Epub 20200609. doi: 10.1038/s41385-020-0309-3. PubMed PMID: 32518368;

PubMed Central PMCID: PMCPMC7726002.

60.   Kedl RM, Lindsay RS, Finlon JM, Lucas ED, Friedman RS, Tamburini BAJ. Migratory

dendritic cells acquire and present lymphatic endothelial cell-archived antigens during lymph node

contraction. Nat Commun. 2017;8(1):2034. Epub 20171211. doi: 10.1038/s41467-017-02247-z.

PubMed PMID: 29229919; PubMed Central PMCID: PMCPMC5725486.

61.   Heesters BA, Chatterjee P, Kim YA, Gonzalez SF, Kuligowski MP, Kirchhausen T, et al.

Endocytosis and recycling of immune complexes by follicular dendritic cells enhances B cell antigen

binding and activation. Immunity. 2013;38(6):1164-75. Epub 20130613. doi:

10.1016/j.immuni.2013.02.023. PubMed PMID: 23770227; PubMed Central PMCID:

PMCPMC3773956.

62.   Daffé M, Quémard A, Marrakchi H. Mycolic Acids: From Chemistry to Biology. 2019. p. 181-

216.

63.   Barber DL, Wherry EJ, Masopust D, Zhu B, Allison JP, Sharpe AH, et al. Restoring function in

exhausted CD8 T cells during chronic viral infection. Nature. 2006;439(7077):682-7. doi:

10.1038/nature04444.

64.   Wu T, Ji Y, Ashley Moseman E, Xu HC, Manglani M, Kirby M, et al. The TCF1-Bcl6 axis

counteracts type I interferon to repress exhaustion and maintain T cell stemness. Science Immunology.

2016;1(6). doi: 10.1126/sciimmunol.aai8593.

65.   Wang Y, Hu J, Li Y, Xiao M, Wang H, Tian Q, et al. The transcription factor TCF1 preserves

the effector function of exhausted CD8 T cells during chronic viral infection. Frontiers in Immunology.

2019;10(FEB):169-. doi: 10.3389/fimmu.2019.00169.

66.   Kwon KW, Kang TG, Lee A, Jin SM, Lim YT, Shin SJ, et al. Protective Efficacy and

Immunogenicity of Rv0351/Rv3628 Subunit Vaccine Formulated in Different Adjuvants Against

Mycobacterium tuberculosis Infection. Immune Netw. 2023;23(2):e16. Epub 20230216. doi:

10.4110/in.2023.23.e16. PubMed PMID: 37179749; PubMed Central PMCID: PMCPMC10166659.

67.   Lima KM, Santos SA, Lima VM, Coelho-Castelo AA, Rodrigues JM, Jr., Silva CL. Single dose

of a vaccine based on DNA encoding mycobacterial hsp65 protein plus TDM-loaded PLGA

microspheres protects mice against a virulent strain of Mycobacterium tuberculosis. Gene Ther.

2003;10(8):678-85. doi: 10.1038/sj.gt.3301908. PubMed PMID: 12692596.

68.   Niu H, Hu L, Li Q, Da Z, Wang B, Tang K, et al. Construction and evaluation of a multistage

Mycobacterium tuberculosis subunit vaccine candidate Mtb10.4-HspX. Vaccine. 2011;29(51):9451-8.

Epub 20111021. doi: 10.1016/j.vaccine.2011.10.032. PubMed PMID: 22024175.

69.   Sarkar S, Mishra A, Periasamy S, Dyett B, Dogra P, Ball AS, et al. Prospective Subunit

Nanovaccine against Mycobacterium tuberculosis Infection horizontal line Cubosome Lipid

Nanocarriers of Cord Factor, Trehalose 6,6' Dimycolate. ACS Appl Mater Interfaces.

2023;15(23):27670-86. Epub 20230601. doi: 10.1021/acsami.3c04063. PubMed PMID: 37262346.

70.   Welsh KJ, Hunter RL, Actor JK. Trehalose 6,6'-dimycolate--a coat to regulate tuberculosis

immunopathogenesis. Tuberculosis (Edinb). 2013;93 Suppl:S3-9. doi: 10.1016/S1472-9792(13)70003-

9. PubMed PMID: 24388646.

71.   Xin Q, Niu H, Li Z, Zhang G, Hu L, Wang B, et al. Subunit vaccine consisting of multi-stage

antigens has high protective efficacy against Mycobacterium tuberculosis infection in mice. PLoS One.

2013;8(8):e72745. Epub 20130815. doi: 10.1371/journal.pone.0072745. PubMed PMID: 23967337;

PubMed Central PMCID: PMCPMC3744459.

72.   Martin MD, Badovinac VP. Defining memory CD8 T cell. Frontiers Media SA; 2018.

73.   Johnston RJ, Poholek AC, DiToro D, Yusuf I, Eto D, Barnett B, et al. Bcl6 and Blimp-1 are

reciprocal and antagonistic regulators of T follicular helper cell differentiation. Science.

2009;325(5943):1006-10. doi: 10.1126/science.1175870.

74.   Nurieva RI, Chung Y, Martinez GJ, Yang XO, Tanaka S, Matskevitch TD, et al. Bcl6 mediates

the development of T follicular helper cells. Science. 2009;325(5943):1001-5. doi:

10.1126/science.1176676.

75.   Clerici L, Casorati G, Dellabona P. B Cell Help by CD1d-Rectricted NKT Cells. Antibodies.

2015;4(4):279-94. doi: 10.3390/antib4040279.

76.   Correia-Neves M, Sundling C, Cooper A, Kallenius G. Lipoarabinomannan in Active and

Passive Protection Against Tuberculosis. Front Immunol. 2019;10:1968. Epub 20190911. doi:

10.3389/fimmu.2019.01968. PubMed PMID: 31572351; PubMed Central PMCID: PMCPMC6749014.

77.   Schleicher GK, Feldman C, Vermaak Y, Verschoor JA. Prevalence of anti-mycolic acid

antibodies in patients with pulmonary tuberculosis co-infected with HIV. Clinical Chemistry and

Laboratory Medicine. 2002;40(9):882-7. doi: 10.1515/CCLM.2002.156.

78.   Pan J, Fujiwara N, Oka S, Maekura R, Ogura T, Yano I. Anti-cord factor (trehalose

6,6'dimycolate) IgG antibody in tuberculosis patients recognizes mycolic acid subclasses. Microbiol

Immunol. 1999;43(9):863-9. doi: 10.1111/j.1348-0421.1999.tb01221.x. PubMed PMID: 10553679.

79.   Bioley G, Lassus A, Terrettaz J, Tranquart F, Corthésy B. Long-term persistence of immunity

induced by OVA-coupled gas-filled microbubble vaccination partially protects mice against infection

by OVA-expressing Listeria. Biomaterials. 2015;57:153-60. doi: 10.1016/j.biomaterials.2015.04.008.

80.   Silva AL, Rosalia RA, Sazak A, Carstens MG, Ossendorp F, Oostendorp J, et al. Optimization

of encapsulation of a synthetic long peptide in PLGA nanoparticles: low-burst release is crucial for

efficient CD8(+) T cell activation. Eur J Pharm Biopharm. 2013;83(3):338-45. Epub 20121129. doi:

10.1016/j.ejpb.2012.11.006. PubMed PMID: 23201055.

81.   Allen SD, Bobbala S, Karabin NB, Modak M, Scott EA. Benchmarking Bicontinuous

Nanospheres against Polymersomes for in Vivo Biodistribution and Dual Intracellular Delivery of

Lipophilic and Water-Soluble Payloads. ACS Applied Materials & Interfaces. 2018;10(40):33857-66.

doi: 10.1021/acsami.8b09906.

82.   Frey M. Development and Optimization of Functionalized Self-Assembling Polymeric

Nanobiomaterials: Northwestern University; 2021.

83.   Van Rhijn I, Iwany SK, Fodran P, Cheng TY, Gapin L, Minnaard AJ, et al. CD1b-mycolic acid

tetramers demonstrate T-cell fine specificity for mycobacterial lipid tails. European Journal of

Immunology. 2017;47(9):1525-34. doi: 10.1002/eji.201747062.

84.   Li S, Choi HJ, Felio K, Wang CR. Autoreactive CD1b-restricted T cells: A new innate-like T-

cell population that contributes to immunity against infection. Blood. 2011;118(14):3870-8. doi:

10.1182/blood-2011-03-341941.

85.   Larsen MH, Biermann K, Chen B, Hsu T, Sambandamurthy VK, Lackner AA, et al. Efficacy

and safety of live attenuated persistent and rapidly cleared Mycobacterium tuberculosis vaccine

candidates in non-human primates. Vaccine. 2009;27(34):4709-17. Epub 20090602. doi:

10.1016/j.vaccine.2009.05.050. PubMed PMID: 19500524; PubMed Central PMCID:

PMCPMC3512200.

86.   Weng X, Kumar A, Cao L, He Y, Morgun E, Visvabharathy L, et al. Mitochondrial metabolism

is essential for invariant natural killer T cell development and function. Proc Natl Acad Sci U S A.

2021;118(13). Epub 2021/03/24. doi: 10.1073/pnas.2021385118. PubMed PMID: 33753493.

87.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-

seq data with DESeq2. Genome Biol. 2014;15(12):550. Epub 2014/12/18. doi: 10.1186/s13059-014-

0550-8. PubMed PMID: 25516281; PubMed Central PMCID: PMCPMC4302049.

88.   Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, et al. Metascape

provides a biologist-oriented resource for the analysis of systems-level datasets. Nature

Communications. 2019;10(1). doi: 10.1038/s41467-019-09234-6.

89.   Yoshida H, Lareau CA, Ramirez RN, Rose SA, Maier B, Wroblewska A, et al. The cis-

Regulatory Atlas of the Mouse Immune System. Cell. 2019;176(4):897-912.e20. doi:

10.1016/j.cell.2018.12.036.

90.   Lahmann A, Kuhrau J, Fuhrmann F, Heinrich F, Bauer L, Durek P, et al. Bach2 Controls T

Follicular Helper Cells by Direct Repression of Bcl-6. The Journal of Immunology. 2019;202(8):2229-

39. doi: 10.4049/jimmunol.1801400.

91.   Man K, Gabriel SS, Liao Y, Gloury R, Preston S, Henstridge DC, et al. Transcription Factor

IRF4 Promotes CD8+ T Cell Exhaustion and Limits the Development of Memory-like T Cells during

Chronic Infection. Immunity. 2017;47(6):1129-41.e5. doi: 10.1016/j.immuni.2017.11.021.

Fig. 1. Physicochemical characterization of BCN and PLGA-NP. (A) Dynamic light scattering

(DLS) analysis of blank BCN and MA-BCN. **(B)** CryoTEM of MA-BCN (scale = 500 nm). **(C)**

Negative staining TEM images of MA-BCN. **(D)** SAXS of blank BCN and MA-BCN. **(E)** DLS

analysis of blank PLGA and MA PLGA. **(F)** MA encapsulation efficiency for MA-BCN and MA

PLGA. N = 3 per condition. Data represented as mean ± SD.

Fig. 2. MA-BCN effectively activated MA-specific T cells *in vitro* and *in vivo*. (A, B) BMDCs were

pulsed for 18 hrs. with selected nanoparticles at various concentrations, co-cultured for 48 hrs. with

DN1 T cells, and T cell activation was measured. **(A)** Percentage of CD69 and CD25-expressing DN1

T cells. **(B)** IFN-g production measured by ELISA. (N=3). **(C, D)** ELISPOT of IFN-g from MA-

specific human M11 T cells cultured for 16 hours with monocyte-derived dendritic cells with selected

nanoparticles at high **(C)** and low **(D)** concentrations. Data are representative of two independent

experiments and displayed as mean ± SEM. Statistical analysis: 2-way ANOVA, significance

designated for MA-BCN vs free MA. **(E-G)** hCD1Tg mice were IN vaccinated with selected

nanoparticles and CellTrace Violet-labeled DN1 T cells were adoptively transferred the next day. After

1 week, DN1 T cell activation and proliferation was measured. **(E)** Representative FACS plots of DN1

T cells in the LN. Percentage of proliferating **(F)** and CD44-expressing **(G)** DN1 T cells in the LN and

lung. Data represented as mean ± SEM. Statistical analysis: 2-way ANOVA. *p<0.05, **p<0.01,

***p<0.001, ****p< 0.0001.

Fig. 3. Vaccination with MA-BCN leads to antigen persistence 6 weeks post-vaccination. hCD1Tg

mice were IT vaccinated with MA-BCN or BCN. 6 weeks later, CellTrace-labeled p25 and DN1 T

cells were adoptively transferred, and T cell activation was measured after 1 week. **(A)** Experimental

design. **(B)** Representative FACS plots of p25 and DN1 T cells in the LN. Percentage of proliferating

(C) and CD44-expressing **(D)** p25 and DN1 T cells in the LN, lung, and spleen. N = 3 or 6 per

condition. Outliers were identified through Grubbs method with alpha=0.05 with one sample removed.

Data represented as mean ± SEM. Statistical analysis: 2-way ANOVA. *$p < 0.05$, **$p < 0.01$, ***$p <$

0.001, ****$p < 0.0001$.

Fig. 4. Ag85B and MA dual loaded BCNs activate antigen-specific T cells without Ag85B

persistence. (A) DLS analysis of blank BCN and Ag85B-MA-BCN. (**B-D**) hCD1Tg mice were IT

vaccinated with Ag85B-MA-BCN or BCN and CellTrace-labeled p25 and DN1 T cells were

adoptively transferred the next day. After 1 week, T cell activation and proliferation was measured. (**B**)

Representative FACS plots of p25 and DN1 T cells in the LN. Percentage of proliferating (**C**) and

CD44-expressing (**D**) p25 and DN1 T cells in the LN, lung, and spleen (N = 4 or 5). (**E-G**) hCD1Tg

mice were IT vaccinated with blank BCN or Ag85B-MA-BCN. 6 weeks later, CellTrace-labeled p25

and DN1 T cells were adoptively transferred, and T cell activation was measured after 1 week. (**E**)

Representative FACS plots of p25 and DN1 T cells in the LN. (**F**) Percentage of proliferating p25 and

DN1 T cells**. (G)** Percentage of CD44-expressing p25 and DN1 T cells in the LN and lung. (N = 4).

Data represented as mean ± SEM. Statistical analysis: 2-way ANOVA. ns = not significant, *$p<0.05$,

**$p<0.01$, ***$p<0.001$, ****$p< 0.0001$.

Fig. 5. Route of vaccination or vector type do not affect antigen persistence. hCD1Tg BMDCs

were pulsed with MA-BCN or MA at concentrations of 5 μg/mL and 10 μg/mL, respectively. hCD1Tg

mice were IT vaccinated with MA or MA-BCN pulsed BMDCs at 6 weeks or 1 week prior to the

adoptive transfer of CellTrace-labeled DN1 T cells. T cell activation and proliferation was measured 1

week after adoptive transfer. (**A**) Experimental design. (**B, C**) Percentage of proliferating (**B**) and

CD44-expressing (**C**) DN1 T cells in the LN, lung, and spleen of vaccinated mice. N = 4 per condition.

N = 4 per condition. hCD1Tg mice were vaccinated SC (subcutaneously) or IV (intravenously) with

blank BCN, MA-BCN, or attenuated Mtb strain and 6 weeks later CellTrace-labeled DN1 T cells were

adoptively transferred. (**D**) Representative FACS plots of DN1 T cells in the LN of mice vaccinated

with indicated conditions. (**E**) Percentage of proliferating DN1 T cells and (**F**) Percentage of CD44-

expressing DN1 T cells in the LN and spleen of mice vaccinated with Blank-BCN (SC), MA-BCN (IV

and SC), and attenuated Mtb. N = 3-5 per condition. Data represented as mean ± SEM. Statistical

analysis: 2-way ANOVA. ns = not significant, $*p<0.05$, $**p<0.01$, $***p<0.001$, $****p<0.0001$.

Fig. 6. Persistent MA remains in part due to encapsulation inside alveolar macrophages and

activates T cells through DCs. B6 or hCD1Tg mice were IT vaccinated with BCN loaded with a

hydrophobic fluorescent dye (DiD) at 6 weeks, 6 days, 48 hours, 24 hours, or 4 hours prior to the

experiment. LN, lung, and spleen were analyzed using **(A)** In Vivo Imaging System (IVIS). Single cell

suspension was then obtained and presence of DiD BCN was quantified in **(B)** total lung, and **(C)**

CD45⁻ cells, neutrophils (Ly6G⁺), alveolar macrophages (CD11c⁺SiglecF⁺), DCs (CD11c⁺), monocytes

(CD11b⁺CD11c⁻), B cells (CD19⁺), T cells (CD3⁺), NK cells (NK1.1⁺), eosinophils (CD11c⁻SiglecF⁺)

by flow cytometry. Gating strategy exemplified in **Suppl Fig 4**. hCD1Tg mice were IT vaccinated with

either BCN or MA-BCN. After 6 weeks, alveolar macrophages were enriched from lungs using anti-

SiglecF. Enriched or flow through cells were co-cultured with DN1 T cells in the presence or absence

of hCD1Tg-expressing BMDCs for 48 hours, and DN1 T cells activation was measured. **(D)**

Experimental design. **(E)** Percentage of CD25-expressing DN1 T cells. Data represented as mean ±

SEM. Statistical analysis: 2-way ANOVA. ns = not significant, $*p<0.05$, $**p<0.01$, $***p<0.001$.

Fig. 7. **MA-BCN vaccination leads to DN1 T cell differentiation into T follicular helper-like T**

cells. Mixed bone marrow chimeric mice were created by adoptive transfer of DN1 bone marrow and

congenic CD45.1 hCD1Tg bone marrow to irradiated hCD1Tg mice. After 5 weeks, mice were

vaccinated with either MA-BCN or PBS, and DN1 surface marker expression was monitored in the

blood and in organs at 6 wks. **(A)** Experimental design. **(B)** Percent CD44⁺ DN1 T cells in the blood at

various time points post vaccination. **(C)** Percent CD44⁺CD62L⁺ DN1 T cells in indicated organs at 6

weeks post vaccination. Memory (CD44⁺CD62L⁺) and naïve (CD44⁻CD62L⁺) DN1 T cells were sorted

from MA-BCN vaccinated hCD1Tg-DN1 BM chimeras at 6 weeks post vaccination and subjected to

RNAseq analysis. N=3 per condition **(D, E)** PCA of log fold change of gene expression for each T cell

subset was performed relative to internal naïve T cell control. **(F)** Relative expression of key T_FH cell

and DEGs in memory subset. **(G)** Gene enrichment analysis of upregulated DEGs was performed

using Metascape. **(H)** Representative FACS plots of DN1 T cells in the LN of mice. Percentage of **(I)** CXCR5$^+$PD1$^+$ or **(J)** KI67$^+$ DN1 T cells DN1 T cells in LN, lung, and spleen. N = 3-5 per condition. Data represented as mean ± SEM. Statistical analysis: 2-way ANOVA. *p<0.05, **p<0.01, ****p < 0.0001.

Fig. 8. **Summary schematic illustrates MA antigen persistence and memory differentiation of antigen-specific T cells induced by vaccination with MA-loaded nanoparticle vaccine.** *Mycobacterium tuberculosis* (Mtb) derived mycolic acid (MA) lipid and Ag85B protein are encapsulated into bicontinuous nanosphere (BCNs) composed of PEG-*b*-PPS polymer. These MA-BCN or Ag85B-MA-BCN are used for intratracheal (IT) vaccination of human CD1 transgenic (hCD1Tg) mice. Following vaccination, dendritic cells (DCs) present the antigens, leading to the activation of CD1b-restricted MA-specific or MHCII-restricted Ag85B-specific T cells (DN1 or p25-specific T cells, respectively). Notably, MA can persist within alveolar macrophages and continue to activate MA-specific T cells even at 6 weeks post-vaccination. These MA-specific T cells have the capacity to differentiate into memory T cells with T follicular-like phenotype.

Figure 1

A

Number (%)

Size (d.nm)

— Blank BCN
— MA BCN

B

C

D

Intensity (AU)

√2
√4
√6
√2
√4
√6

Q(1/Å)

● MA BCN
● Blank BCN

E

Number (%)

Size (d.nm)

— Blank PLGA
— MA PLGA

F

Encapsulation efficiency (%)

MA
PLGA

MA
BCN

Figure 2

Figure 3

A Vaccinate IT with MA-BCN or Blank-BCN   DN1 T cells   P25 T cells   Test activation proliferation

hCD1Tg   6 wks post vaccination   1 wk post AT

B

P25   DN1

4.3   7.6   **Blank BCN**

6.0   82   **MA BCN**

Count

Cell trace

□ Blank-BCN
■ MA-BCN

C

% Dividing cells

LN   Lung   Spleen

****   ***   *

P25 DN1   P25 DN1   P25 DN1

D

% CD44+ T cells

LN   Lung   Spleen

****   ***   *

P25 DN1   P25 DN1   P25 DN1

Figure 4

Figure 5

A

hCD1Tg — MA-BCN or MA-pulsed DCs — hCD1Tg — MA-BCN or MA-pulsed DCs — DN1 T cells — Test activation proliferation

6 wks pre-AT | 1 wk pre-AT | AT | 1 wk post AT

B

LN   Lung   Spleen

% Dividing T cells

ns ** ns *** ns **

Vaccination week: 1 6

C

LN   Lung   Spleen

% CD44⁺ T cells

ns *** ns ** ns **

Vaccination week: 1 6

▲ MA pulsed DCs
■ MA-BCN pulsed DCs

□ SC Blank-BCN
■ IV MA-BCN
■ SC MA-BCN
● Attenuated Mtb

D

Blank BCN: 20 | IV MA-BCN: 95 | SC MA-BCN: 79 | SC Mtb: 85

Count

Cell trace

E

% Dividing DN1 T cells

**** *** **** *** *** *

LN   Spleen

F

% CD44⁺ DN1 T cells

**** *** **** **** ** *

LN   Spleen

Figure 6

A

B

C

D

E

Figure 7

Figure 8

Fig. S1. Physicochemical characterization of PLGA-NP. (A) CryoTEM image and DLS. **(B)** SAXS

scattering curve fitted with a sphere model.

Fig. S2. Intracellular localization of PLGA-NP and BCN. (A) Representative confocal image of

intracellular localization of PLGA or BCN (Texas red Dextran dye), lysosome (lysotracker), nucleus

(NucBlue). **(B)** Quantification of co-localization of lysotracker with Texas red dye by Pearson

correlation. (8 cells were analyzed per experiment, 3 independent experiments), ****$p < 0.0001$.

Fig. S3. MA encapsulated by either BCN or MC formulation leads to lipid antigen persistence.

hCD1Tg mice were IT vaccinated with MA-BCN or MA-MC at 6 weeks or 1 week prior to adoptive

transfer of cell trace stained DN1 T cells. T cell activation and proliferation was measured 1 week after

adoptive transfer. **(A)** Experimental diagram. **(B)** Percentage of proliferating DN1 T cells in LN, lung,

and spleen. **(C)** Percentage of CD44-expressing DN1 T cells in the LN, lung, and spleen. N = 3 or 4

per condition. Data represented as mean ± SEM. ns = not significant, *$p < 0.05$.

Fig. S4. P25-specific T cell activation and proliferation and bacterial burden in attenuated Mtb

vaccination. hCD1Tg were SC vaccinated with attenuated Mtb. **(A)** Bacterial burden in the spleen and

lung were determined at 2- or 6-weeks post vaccination. hCD1Tg mice were vaccinated with

attenuated Mtb and after 6 weeks, CellTrace-labeled p25 T cells were adoptively transferred into the

mice. After 1 week, T cell activation and proliferation was measured. **(B)** Representative FACS plots

of in LN. **(C)** Percentage of proliferating P25 T cells in the LN and spleen. **(D)** Percentage of CD44-

expressing P25 T cells in the LN and spleen (N = 5).

Fig. S5. Gating strategy for analyzing biodistribution of BCN in the lung. Lung single cell

suspension was gated as follows: CD45$^-$ cells, neutrophils (Ly6G$^+$), alveolar macrophages

(CD11c$^+$SiglecF$^+$), DCs (CD11c$^+$), monocytes (CD11b$^+$CD11c$^-$), B cells (CD19$^+$), T cells (CD3$^+$), NK

cells (NK1.1$^+$), and eosinophils (CD11c$^-$SiglecF$^+$).

Fig. S6. Representative FACS plot of alveolar macrophages within enriched and flow through

fractions.

Fig. S7. Memory and naïve DN1 T cells display distinct gene expression profiles. Memory

(CD44$^+$CD62L$^+$) and naïve (CD44$^-$CD62L$^+$) DN1 T cells were sorted from LNs of MA-BCN

vaccinated hCD1Tg-DN1 BM chimeras at 6 weeks post vaccination and subjected to RNAseq analysis

(A) Representative figure of CD44 and CD62L expression in DN1 T cells in LN, lung, and spleen. **(B)**

MA plot representing memory vs naïve analysis. **(C)** PCA of normalized counts. N=3 per condition.

39
40

Supplementary Figure 1

A

B

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

Supplementary Figure 5

Supplementary Figure 6

Supplementary Figure 7

A

PBS

MA-BCN

CD44

CD62L

B

C

Memory
542

Naïve
453