# Novel axonemal protein ZMYND12 is required for flagellum function and male fertility

Denis Dacheux[1,2,18], Guillaume Martinez[3,4,18], Christine E. Broster Reix[1], Julie Beurois[4], Patrick Lorès[5], Magamba Tounkara[1], Jean-William Dupuy[6], Derrick R. Robinson[1,2], Corinne Loeuillet[4], Emeline Lambert[4], Zeina Wehbe[4], Amir Amiri-Yekta[7], Abbas Daneshipour[7], Seyedeh-Hanieh Hosseini[8], Raoudha Zouari[9], Sélima Fourati Ben Mustapha[9], Lazhar Halouani[9], Xiaohui Jiang[10,11,12], Ying Shen[11,12], Chunyu Liu[13], Nicolas Thierry-Mieg[14], Amandine Septier[14], Marie Bidart[4,15], Véronique Satre[3,4], Caroline Cazin[4,16], Zine-Eddine Kherraf[4,16], Christophe Arnoult[4], Pierre F. Ray[4,16], Aminata Touré[17,19], Mélanie Bonhivers[1,19] and Charles Coutton[3,4,19,*]

[1] Univ. Bordeaux, CNRS, MFP, UMR 5234, F-33000 Bordeaux, France.

[2] Bordeaux INP, Microbiologie Fondamentale et Pathogénicité, UMR 5234, F-33000 Bordeaux, France.

[3] CHU Grenoble Alpes, UM de Génétique Chromosomique, 38000 Grenoble, France.

[4] Institute for Advanced Biosciences, INSERM U1209, CNRS UMR 5309, Univ. Grenoble Alpes, Team Genetics Epigenetics and Therapies of Infertility, 38000 Grenoble, France.

[5] Institut Cochin, INSERM U1016, CNRS UMR 8104, Université Paris Cité, 75014 Paris, France.

[6] Univ. Bordeaux, Plateforme Protéome, F-33000, Bordeaux, France.

[7] Department of Genetics, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran-Iran.

[8] Department of Andrology, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran-Iran.

[9] Polyclinique les Jasmins, Centre d'Aide Médicale à la Procréation, Centre Urbain Nord, Tunis, Tunisia.

[10] Human Sperm Bank, West China Second University Hospital of Sichuan University, Chengdu, Sichuan, China.

[11] NHC Key Laboratory of Chronobiology, Sichuan University, Chengdu, Sichuan, China.

[12] Key Laboratory of Birth Defects and Related Diseases of Women and Children (Sichuan University), Ministry of Education, China.

[13] Obstetrics and Gynecology Hospital, Fudan University, Shanghai, China.

[14] Univ. Grenoble Alpes, CNRS, UMR5525, TIMC, 38000 Grenoble, France.

[15] CHU Grenoble Alpes, Laboratoire de Génétique Moléculaire: Maladies Héréditaires et Oncologie, Grenoble, France.

[16] CHU de Grenoble, UM GI-DPI, Grenoble, 38000, France.

[17] Institute for Advanced Biosciences, INSERM U 1209, CNRS UMR 5309, Université Grenoble Alpes, Team Physiology and Pathophysiology of Sperm cells 38000 Grenoble, France.

[18] These authors contributed equally to this work.

[19] These authors contributed equally to this work.

*Correspondence: CCoutton@chu-grenoble.fr

Running title: Role of ZMYND12 in flagellum function

# ABSTRACT

Male infertility is common and complex, presenting a wide range of heterogeneous phenotypes. Although about 50% of cases are estimated to have a genetic component, the underlying cause often remains undetermined. Here, from whole-exome sequencing on samples from 168 infertile men with severe sperm flagellum defects, we identified homozygous *ZMYND12* variants in four unrelated patients. In sperm cells from these individuals, immunofluorescence revealed altered localization of DNAH1, DNALI1, WDR66 and TTC29. Axonemal localization of ZMYND12 ortholog TbTAX-1 was confirmed using the *Trypanosoma brucei* model. RNAi knock-down of TbTAX-1 dramatically affected flagellar motility, with a phenotype similar to ZMYND12-variant-bearing human sperm. Co-immunoprecipitation and ultrastructure expansion microscopy in *T. brucei* revealed TbTAX-1 to form a complex with TTC29. Comparative proteomics with samples from *Trypanosoma* and *Ttc29* KO mice identified a third member of this complex: DNAH1. The data presented revealed the axonemal ZMYND12/TTC29/DNAH1 trio to critically contribute to flagellum function and assembly in humans and *Trypanosoma*. *ZMYND12* is thus a new asthenoteratozoospermia-associated gene, bi-allelic variants of which cause severe flagellum malformations and primary male infertility.

**Keywords:** asthenoteratozoospermia / male infertility / MMAF / sperm flagellum / ZMYND12

# INTRODUCTION

Male infertility is one of the most challenging health issues that will face us in the near future. Recent decades have been marked by a significant and constant deterioration of quantitative and qualitative sperm parameters in both humans and animals, demonstrating the detrimental impact of recent environmental changes (Rolland *et al*, 2013). However, environmental factors are not the sole cause of male infertility, which is often multifactorial. Among the various possible causes, it is estimated that a genetic component, alone or in association, can be found in about half of cases (Oud *et al*, 2022).

Spermatozoa are the most highly differentiated and specialized human cells, as reflected by the 2,300 genes necessary for normal spermatogenesis (Lu *et al*, 2019). The complex process of spermatogenesis is divided into three phases: proliferation and differentiation of spermatogonia, meiosis, and the final critical step, spermiogenesis. During spermiogenesis, haploid round spermatids undergo an extraordinary series of transformations to acquire the specific morphological features necessary for them to competently fertilize an egg. The stages in spermiogenesis include nuclear compaction, acrosome formation and flagellum biogenesis (O'Donnell, 2014). Any defects in genes driving these differentiation steps can lead to severe functional or morphological sperm defects, and thus male infertility.

Interestingly, the prevalence of genetic defects is higher in conditions with severe sperm phenotypes (Coutton *et al*, 2015). Multiple Morphological Abnormalities of the sperm Flagellum (MMAF) is a condition associated with extreme morphological sperm defects characterized by a mosaic of phenotypes. Thus, MMAF sperm cells may lack a flagellum or bear a short, irregular, or coiled flagellum. Because of these characteristics, MMAF causes total asthenozoospermia with almost no progressive sperm (Touré *et al*, 2020). At the ultrastructural level, MMAF is associated with severe disorganization of both axonemal and peri-axonemal

structures (Nsota Mbango *et al*, 2019). Approximately 40 genes have been strongly or definitively associated with the MMAF phenotype so far (Wang *et al*, 2020). However, this high genetic heterogeneity means the prevalence of individual variants of each gene remains low, and the underlying genetic component remains unknown for about half of affected individuals. It is therefore necessary to pursue the identification of new candidate MMAF genes to increase the diagnostic yield and improve our understanding of the pathophysiological mechanisms inducing this sperm phenotype. In addition, the identification of MMAF genes provides invaluable clues to the molecular mechanisms underlying sperm flagellum biogenesis. Therefore, efforts are continuously being deployed to decipher the precise organization, localization and interactions of each protein encoded by these genes within the flagellum.

Here, following analysis of an initial cohort of 167 MMAF individuals by whole-exome sequencing (WES), we identified truncating homozygous variants in *ZMYND12* in three unrelated individuals. A fourth individual with a similar mutation was subsequently identified in a Chinese cohort. In this paper, we present a thorough investigation of the role of this new gene by studying its ortholog, TbTAX-1, in *Trypanosoma*, an evolutionarily-distant model sharing a highly conserved flagellar structure. We first showed the importance of ZMYND12 for the axonemal structure and function of the flagellum. Subsequently, we shed light on ZMYND12 function in the flagellum using the TbTAX-1 protist model and *Tttc29* KO mice. Results formally demonstrated that ZMYND12 interacts directly with DNAH1 and TTC29 (two known MMAF-related proteins) within the same axonemal complex.

# RESULTS

**Whole-exome sequencing (WES) identified homozygous truncating variants in *ZMYND12* in MMAF individuals.**

In our cohort of 167 MMAF individuals, we previously identified 83 individuals (49.7%) with harmful gene variants in known MMAF-related genes. Reanalysis of the remaining negative exomes identified three unrelated MMAF individuals (ZMYND12$_{1-3}$) (Fig. 1A, Table 1) carrying homozygous variants in *ZMYND12* (Zinc Finger MYND-Type Containing 12, NM_032257.5). This gene was not previously associated with any pathology. Two individuals (ZMYND12$_1$ and ZMYND12$_3$) originated from Tunisia, and one individual (ZMYND12$_2$) was of Iranian origin.

*ZMYND12* is located on chromosome 1 and contains eight exons encoding the Zinc finger MYND domain-containing protein 12, a predicted 365-amino acid protein (Q9H0C1). According to data from GTEx, *ZMYND12* is strongly expressed in the testis and is associated with cilia and flagella (Yamamoto *et al*, 2006, 2021). Moreover, quantitative single-cell RNA sequencing datasets from human adult testis (ReproGenomics Viewer) (Darde *et al*, 2019) indicate abundant expression in germ cells from zygotene spermatocyte to late spermatid stages. This expression pattern suggests a role in sperm cell differentiation and/or function. ZMYND12 protein was also detected at significant levels in the human sperm proteome (Wang *et al*, 2013), but at very low levels in human airway cilia (Blackburn *et al*, 2017). We performed RT-qPCR experiments on human tissue panels (Fig. S3), and results confirmed that expression of *ZMYND12* mRNA is predominant in testis as compared with the other tissues tested.

A stop-gain variant c.433C>T; p.Arg145Ter (NM_032257, GRCh38 g.chr1:42440017G>A, rs766588568), located in *ZMYND12* exon 4 (Fig. 1B), was identified in individuals ZMYND12$_1$ and ZMYND12$_3$. Subsequently, a Chinese MMAF subject harboring

the same homozygous stop-gain variant in *ZMYND12* (ZMYND12_4) was identified in the cohort collected by the Human Sperm Bank of West China Second University Hospital of Sichuan University hospital (Fig. 1, Table 1). This c.433C>T variant is present in the Genome Aggregation Database (gnomAD v2.1.1, http://gnomad.broadinstitute.org/variant/1-42905688-G-A) database with a minor allele frequency (MAF) of 3.391e-05. This nonsense variant, identified by exome sequencing, was confirmed by Sanger sequencing in all four individuals (Fig. 1C).

Using the ExomeDepth software package (Plagnol *et al*, 2012), individual ZMYND12_2, was identified based on a homozygous deletion removing *ZMYND12* exons 6, 7 and 8 (Fig. S1). We confirmed the presence of the homozygous deletion in the affected individual by custom MLPA (Fig. 1D). The deletion identified was not listed in public databases, including DECIPHER (https://www.deciphergenomics.org/) and the Database of Genomic Variants (http://dgv.tcag.ca/dgv/app/home).

For the individuals carrying *ZMYND12* mutations, analysis of control databases revealed no low-frequency variants in other genes reported to be associated with cilia, flagella or male fertility. We therefore focused our study on *ZMYND12*, which appeared as the best MMAF candidate gene. All of the *ZMYND12* variants identified here have been deposited in ClinVar under reference SUB12540308.

## ZMYND12 is located in the sperm flagellum, its variants cause severe axonemal disorganization

To further investigate the pathogenicity of the *ZMYND12* variants identified, we examined the distribution of the ZMYND12 protein in control and mutated sperm cells by immunofluorescence (IF). Due to sample availability, these analyses were only carried out for

individual ZMYND12_3 – carrier of the recurrent stop-gain variant c.433C>T. In sperm from control individuals, ZMYND12 antisera decorated the full length of the sperm flagellum (Fig. 1E). Conversely, and in line with a truncation or degradation of the mutated protein, ZMYND12 was totally absent from all sperm cells from individual ZMYND12_3 (Fig. 1E). Importantly, ZMYND12 was detected in sperm cells from MMAF individuals bearing mutations in other genes (*ARMC2* and *WDR66*) or with unknown genetic causes (Fig. S2). Thus, the absence of ZMYND12 is not a common feature of the MMAF phenotype, and is specifically associated with *ZMYND12* variants.

To explore the ultrastructural defects induced by *ZMYND12* variants in human sperm cells, we next studied the presence of proteins present in various axonemal and peri-axonemal substructures by IF. The following proteins were investigated: AKAP4 (fibrous sheath, FS), CFAP70 (axoneme-binding protein with unknown precise location), DNAI1 (outer arm intermediate chain), DNALI1 (inner arm light intermediate chain), DNAH1 (inner arm heavy chain), DNAH8 (outer arm heavy chain), DNAH17 (outer arm heavy chain), GAS8 (Nexin-Dynein Regulatory Complex, N-DRC), RSPH1 (radial spoke, RS), SPAG6 (central pair complex, CPC), WDR66/CFAP251 (calmodulin-associated and spoke-associated complex, CSC), and TTC29 (axoneme-binding protein with unknown precise location). In sperm cells from ZMYND12_3, we first observed a total lack of SPAG6 staining (Fig. 2A), suggesting severe CPC defects. This type of defect is a common feature of MMAF spermatozoa (Coutton *et al*, 2015). In addition, no immunostaining for DNAH1, DNALI1, or WDR66 was detected in sperm from individual ZMYND12_3 (Fig. 2B, 2C and 2D). These results suggested a strong disorganization of the IDAs and the CSC. ZMYND12 could therefore be a key axonemal component responsible for stability of these structures.

We subsequently examined the presence of TTC29 and CFAP70, two proteins with as yet incompletely characterized function and localization. In control sperm, TTC29 and CFAP70

immunostaining decorated the full length of the flagellum, but neither protein was detected in cells from individual ZMYND12_3 (Fig. 2E and 2F). Intriguingly, ZMYND12 was undetectable in the sperm flagellum from an MMAF individual carrying a previously-reported TTC29 splicing variant c.1761+1G>A (Lorès *et al*, 2019b) (Fig. S4). Unfortunately, no sample from an individual carrying a bi-allelic *CFAP70* variant could be analyzed. Nevertheless, the available data suggest that TTC29, CFAP70, and ZMYND12 may be physically or functionally associated within the same axonemal complex.

We also observed heterogeneous RSPH1 staining with all flagellar morphologies (Fig. S5), indicating that radial spokes are inconsistently present in sperm cells from individual ZMYND12_3. In contrast, the immunostaining patterns for AKAP4, DNAH8, DNAH17, DNAI1, and GAS8 were comparable to those observed in control sperm cells. This result indicates that the fibrous sheath (FS), the outer dynein arms (ODAs), and the N-DRC were not directly affected by *ZMYND12* variants (Fig. S6). Transmission Electron Microscopy (TEM), which could provide direct evidence of ultrastructural defects, could not be performed due to the very low number of sperm cells available from *ZMYND12* individuals.

**TbTAX-1, the *Trypanosoma brucei* ortholog of ZMYND12, is an axoneme-associated protein involved in flagellar motility**

To validate our candidate gene and better characterize ZMYND12 localization and function, we used the Trypanosoma model. Forward and reverse genetic approaches are well established for this model, and have largely contributed to the characterization of the molecular pathogenesis of a number of human diseases caused by defective cilia and/or flagella (Vincensini *et al*, 2011). BlastP analysis of the *Trypanosoma brucei* genome database (Aslett *et al*, 2010) identified the putative ortholog of ZMYND12 as Tb927.9.10370. The *T. brucei* gene encodes TbTAX-1, a 363 amino acid protein previously identified by proteomics analyses

as a flagellar protein, annotated as the inner dynein arm protein/p38 (FAP146) (Broadhead *et al*, 2006a; Ralston *et al*, 2009). Human ZMYND12 (Q9H0C1) and TbTAX-1 share 29% protein identity and 48% similarity. Both proteins contain a TPR motif domain potentially involved in protein-protein interaction (D'Andrea & Regan, 2003) (Fig. S7A and B).

RNAi knock-down of TbTAX-1 in the bloodstream form (BSF) of *T. brucei* is reported to lead to multi-flagellated cells, suggesting incomplete cell division (Broadhead *et al*, 2006a). This phenotype is described when proteins directly or indirectly involved in flagellar motility are knocked down in the BSF (Coutton *et al*, 2018b; Ralston *et al*, 2011). In contrast, in the procyclic form, knock-down induces cell sedimentation due to motility defects (Baron *et al*, 2007).

Here, to analyze the protein's localization by immunofluorescence (IF) on detergent-extracted cells (cytoskeletons, CSK), we generated a procyclic *T. brucei* cell line endogenously expressing myc-tagged TbTAX-1 ($_{myc}$TbTAX-1). $_{Myc}$TbTAX-1 specifically localized to the flagellum, as shown by co-staining with the Paraflagellar Rod-2 protein (PFR2), a marker of the para-axonemal and axoneme-associated structure (PFR) (Fig. 3A). Importantly, $_{Myc}$TbTAX-1 was found to associate with the axoneme fraction of the flagellum, with staining observed in the distal part of the flagellum, beyond the limit of PFR labeling (Fig. 3A). Furthermore, mycTbTAX-1 labeling was observed in both the old flagellum and the new flagellum, indicating that the protein is present at all stages of the cell cycle, in line with proteomics data reported for the Trypanosoma cell cycle (Crozier *et al*, 2018).

### *Trypanosoma brucei* TAX-1 is involved in flagellum motility

To assess the functional role of TbTAX-1 in the procyclic parasite model, we used a tetracycline-inducible RNA interference (RNAi) system (Wirtz *et al*, 1999) to knock-down TbTAX-1 expression. Using the $_{myc}$TbTAX-1 / TbTAX-1$^{RNAi}$ *T. brucei* cell line (Fig. 3), we

observed the effects of TbTAX-1 knock-down following the addition of tetracycline. Western blot analysis confirmed successful RNA interference, leading to a significant reduction in TbTAX-1 protein levels (Fig. 3B). These results were confirmed by IF, where expression of $_{Myc}$TbTAX-1 was strongly reduced in the new flagellum (NF) compared to the old flagellum (OF) (Fig. 3C). Comparison of TEM images of thin sections from wild-type (WT) and RNAi-induced cells after 7 days revealed no obvious differences in flagellum ultrastructure (Fig. S8), and TbTAX-1 knock-down had no effect on cell growth (Fig. 3D). However, video microscopy evidence pointed to a strong mobility defect (Movies S1 and S2). Tracking analysis (Fig. 3E) indicated that reduced mobility was linked to problems with flagellar beating. In line with these observations, sedimentation assays showed that after induction for 48 h, 70% of RNAi-induced cells had sedimented (Fig. 3F).

**The truncated TbTAX-1 protein is not associated with the flagellum and cannot rescue the motility defect induced by TbTAX-1 RNAi**

The p.Arg145Ter mutation of *ZYMND12* identified in MMAF individuals could theoretically produce a truncated form of the protein (aa 1-144). To assess where a similar truncated protein – TbTAX-1 (aa 1-128) – would localize, we generated a TAX-1$_{HA}$ *Trypanosoma* cell line expressing the truncated form TbTAX-1$^{Trunc}_{Ty1}$ alongside the tagged full-length form. The relative sizes of the mutated, truncated ZMYND12 protein potentially encoded in human patients and the equivalent modified, truncated protein – TbTAX1$^{Trunc}$ – are presented in Fig. S7B.

In this cell line, expression of *TbTAX-1$^{RNAi}$* – specifically targeting the full-length protein and not the truncated form – was inducible (Fig. 4A). Western blot analysis of fractionated samples (whole cells (WC), detergent-extracted cells or cytoskeletons (CSK), and the flagellum fraction (FG)), we localized the truncated TbTAX-1$^{Trunc}_{Ty1}$ protein (Fig. 4B). The

full-length TbTAX-1 protein was strongly associated with the axoneme, and was detected in the salt-extracted flagellum fraction. In contrast, the truncated TbTAX-1$^{Trunc}$$_{Ty1}$ form was detected neither in the flagellum nor in the cytoskeleton-associated fractions (Fig. 4B). These results were confirmed by IF on CSK (Fig. 4C). As expected, induction of *TbTAX-1*$^{RNAi}$ led to decreased TbTAX-1 levels, but had no effect on TbTAX-1$^{Trunc}$$_{Ty1}$ (Fig. 4D). Cell growth was not affected (Fig. 4E), and the sedimentation phenotype was still observed (Fig. 4F). Thus, it appears that the N-terminal domain of TbTAX-1 is not involved in protein targeting and function, and cannot complement TbTAX-1 RNAi knock-down.

### TbTTC29 and TAX-1 belong to the same protein complex; TbTTC29 expression depends on TbTAX-1

As indicated above, in IF analyses of spermatozoa from individuals bearing a mutated ZYMND12, proteins CFAP70, WDR66, TTC29, and SPAG6 were undetected. To further study associations between these proteins, we generated *T. brucei* TbTAX-1$_{HA}$/ *TbTAX-1*$^{RNAi}$ background cell lines expressing their respective orthologs i.e., TbCFAP70$_{Ty1}$ (Tb927.10.2380), TbWDR66$_{Ty1}$ (Tb927.3.1670), TbTTC29$_{Ty1}$ (Tb927.3.1990), and $_{Ty1}$TbPF16/SPAG6 (Tb927.1.2670).

Western blotting revealed that CFAP70, WDR66 and SPAG6 were tightly associated with the flagellum fraction (Fig. S9A). However, none of these proteins was pulled down with Tb-TAX1 upon co-IP with the anti-Ty1 antibody (Fig. S9B). Furthermore, RNAi knock-down of TbTAX-1 expression did not alter their overall levels (Fig. S9C). The lack of direct interaction between TbTAX-1 and these three proteins was further confirmed by ultrastructure expansion microscopy IF (Gambarotto *et al*, 2019; Kalichava & Ochsenreiter), which provided no evidence of colocalization (Fig. S9D).

In contrast, decreased TbTAX-1 levels were associated with decreased TbTTC29 expression, and *vice versa* according to western blot quantification (Fig. 5A and 6B). By IF, TbTAX-1 was also observed to co-localize with TbTTC29 on the axoneme (Fig. 5C). However, analysis of ultrastructure expansion microscopy (U-ExM) and confocal microscopy intensity peaks revealed the colocalization to be incomplete (Fig. 5D). Nevertheless, IP of TbTTC29$_{Ty1}$ from a TbTTC29$_{Ty1}$/TbTAX-1$_{HA}$ cell line pulled down TbTAX-1$_{HA}$ and *vice versa* (Fig. 5E). These results suggest that TbTAX-1 and TbTTC29 are part of the same protein complex.

The co-immunoprecipitated proteins were further analyzed by mass spectrometry. Comparative analysis of the significant hits in each co-IP (anti-Ty1 and anti-HA) (enrichment fold ≥2) identified two additional proteins enriched in both IPs: Tb927.11.3880 – a flagellar actin-like protein (Ersfeld & Gull, 2001a) – and Tb927.11.8160 – a putative inner arm dynein heavy chain protein belonging to the IAD-4 protein family – (data summarized in Table S1). Tb927.11.8160 is potentially involved in flagellum motility (Wickstead & Gull, 2007), it localizes along the *T. brucei* axoneme according to the TrypTag genome-wide protein localization resource (http://tryptag.org/) (Dean *et al*, 2017). Significantly, BlastP searches of mammalian genomes for Tb927.11.8160 orthologs identified DNAH1 (Protein ID: XP_828890.1).

**DNAH1 and DNALI1 both interact with TTC29 in mouse**

In a recent human study, we showed that the presence of bi-allelic *TTC29* variants induces a MMAF phenotype and male infertility (Lorès *et al*, 2019b). By CRISPR/Cas9 mutagenesis and RNAi silencing, we demonstrated that the orthologous TTC29 proteins in mouse and in *Trypanosoma brucei*, respectively, also constitute flagellar proteins that are

required for proper flagellum beating, indicating an evolutionarily-conserved function for TTC29 (Lorès *et al*, 2019b).

Here, we took advantage of the availability of the mutant mouse model to further decipher the protein complexes associated with TTC29 function within the flagella. We performed comparative analyses of co-immunoprecipitations of testis protein extracts from WT mice using TTC29-specific antibodies, and from two previously generated TTC29 KO mouse lines lacking the TTC29 protein (*Ttc29*$^{-/-}$ L5 and L7) (Lorès *et al*, 2019b, 29). Mass spectrometry analyses of four independent experimental replicates for each mouse line identified a set of 124 proteins specifically immunoprecipitated in WT testes.

These proteins were further examined to identify TTC29 binding proteins (Table S2). Filtering of immunoprecipitated proteins based on peptide number and replicate constancy identified DNAH1 and DNALI1, in addition to TTC29, as the only proteins that were co-immunoprecipitated with a significant number of peptides in all four experimental replicates. In our previous study, DNAH1 and DNALI1 were detected at equal levels in flagella purified from WT and Ttc29$^{-/-}$ KO mice (Lorès *et al*, 2019b). Therefore, we considered DNAH1 and DNALI1 to be authentic protein partners of TTC29 in sperm flagella, in line with data obtained with *Trypanosoma brucei*. Thus, our data indicate that ZMYND12 interacts with DNAH1 and DNALI1, two established components of the axonemal IDA.

## DISCUSSION

The results presented here show that the presence of bi-allelic *ZMYND12* variants induces a typical MMAF phenotype in humans, suggesting that this gene is necessary for sperm flagellum structure and function. ZMYND12 is a previously uncharacterized MYND-domain-containing zinc finger protein, we therefore explored its localization and function using a recognized model: the flagellated protozoan *Trypanosoma*. This model is widely used to

investigate flagellum biogenesis and the function of flagellar proteins. In addition to having a very similar axonemal organization and composition to human flagellum, *Trypanosoma* has numerous advantages for functional studies, and is very useful for reverse genetics and biochemical or proteomics studies (Vincensini *et al*, 2011).

We first confirmed that ZMYND12 is an axonemal protein, and then used RNAi experiments to confirm its critical role in flagellum function. ZMYND12 knock-down led to dramatic flagellar motility defects like those observed in human sperm (Fig. 3). In addition, a truncated form of TAX-1 protein lacking the TPR domain – mimicking the expected truncated ZMYND12 protein in MMAF individuals bearing the p.Arg145Ter variant – was not associated with the flagellum, and failed to rescue TAX-1 RNAi-induced motility defects (Fig. 4). These results clearly indicate that the TPR motif of ZMYND12 is critical for protein targeting and function in both human sperm and *Trypanosoma*. Other groups also previously reported a significant role for TPR proteins in sperm flagellar biogenesis and male infertility (Liu *et al*, 2019b; Lorès *et al*, 2019a). Indeed, TPR domains are known to be involved in protein-protein interaction and the assembly of multiprotein complexes (D'Andrea & Regan, 2003). We therefore decided to further explore the potential interactions between ZMYND12 and other axonemal proteins.

IF experiments in sperm cells from ZMYND12 individuals revealed severe ultrastructural defects dramatically affecting several components of the axoneme. More specifically, IDAs and the CSC were severely disorganized, as shown by the total absence of DNAH1, DNALI1 and WDR66 immunostaining (Fig. 2B, 2C and 2D). Intriguingly, TTC29 and CFAP70 were also absent from sperm cells from individuals bearing ZMYND12 mutations (Fig. 2E and 2F). These observations suggest some functional or physical interaction between these proteins. However, the exact function and location of TTC29 and CFAP70 remain unclear. In *Chlamydomonas*, TTC29 has been suggested to be a component of IDAs

(Yamamoto *et al*, 2008), but this protein was also reported to possibly contribute to the IFT machinery (Lorès *et al*, 2019a; Brooks, 2014). For CFAP70, functional studies in *Chlamydomonas* demonstrated that the protein localized at the base of the ODA, where it could regulate ODA activity (Shamoto *et al*, 2018). In *T. brucei*, we screened the axonemal proteins WDR66, CFAP70, SPAG6, and TTC29 to test their interaction with the ZMYND12 homolog. Our results indicated that TbTAX-1 only interacts with TbTTC29, and that the two proteins are part of the same axonemal complex (Fig. 5A and 5B). This interaction was confirmed by ultrastructure expansion microscopy and confocal microscopy (Fig. 5C and 5D). A third member of the complex – Tb927.11.8160, the ortholog of human DNAH1 – was subsequently identified by comparative proteomics analysis of co-immunoprecipitated proteins (Fig. 5E and Table S1). These results were further confirmed using the *Ttc29*-null mouse model previously generated in our laboratory (Table S2). All of these results are consistent with previous studies in *Chlamydomonas* suggesting that p38 (ZMYND12 ortholog) and p44 (TTC29 ortholog) interact and play a role in IDAd docking on the axoneme (Yamamoto *et al*, 2008). Interestingly, the axonemal IDAd in *Chlamydomona*s is composed of the DHC2 heavy chain and the p28 light chain, orthologs of human DNAH1 and DNALI1, respectively. Moreover, the recent human reference interactome map obtained by high-throughput two-hybrid screening (Luck *et al*, 2020) also supports direct protein-protein interaction between ZMYND12 and TTC29. Our results thus formally demonstrate that ZMYND12 interacts with TTC29 within the axonemal complex, in association with DNAH1 and inner dynein arms. DNAH1 is directly connected through an arc-like structure (King, 2013) to the base of radial spoke 3 (RS3), which is anchored to the axoneme by the CSC (Dymek *et al*, 2011). In humans and *Trypanosoma*, the absence of ZMYND12 in the flagellum probably destabilizes this axonemal complex, and consequently alters the stability of the RS3 base-docked IDAs including DNAH1, the CSC and RS. Each of these defects leads to severe axonemal disorganization, and a MMAF sperm phenotype. Our

team previously reported deleterious variants of *TTC29* and *DNAH1* associated with a MMAF phenotype. The flagellar defects caused by these variants are similar to those observed in sperm from individuals with *ZMYND12* defects (Ben Khelifa *et al*, 2014; Lorès *et al*, 2019a; Liu *et al*, 2019a). The results presented here thus contribute to elucidating the physical and functional associations between ZMYND12, TTC29 and DNAH1 in the axoneme, and emphasize the key role played by this triumvirate in maintaining the structure and function of the sperm flagellum.

Adding the three ZMYND12-mutated individuals to the 83 individuals previously identified with variants in known MMAF-related genes in our cohort of 167 individuals, we now have a diagnostic efficiency of 51.4% (86/167). This figure demonstrates the efficiency and the clinical utility of WES for the investigation of the genetic causes and pathophysiology of MMAF syndrome. Among the variants identified, we found the same *ZMYND12* nonsense variant c.433C>T; p.(Arg145Ter) in three unrelated individuals from diverse geographical origins (*i.e*., Tunisia, China), suggesting that the Arg145 residue may constitute a mutational hotspot. Importantly, as previously reported (Kherraf *et al*, 2018a), we confirmed that WES can detect small pathogenic exon deletions. These events may constitute a recurrent disease mechanism in MMAF, and should therefore be systematically explored. Despite these significant advances, the genetic cause of MMAF remains unknown for about half the individuals affected, due to the high genetic heterogeneity of the disorder. To improve the diagnostic rate, more powerful techniques such as whole-genome sequencing may now be envisaged for MMAF patients for whom WES fails to provide results (Meienberg *et al*, 2016).

The results presented here demonstrate that the combined use of exome sequencing and the *Trypanosoma* model can efficiently reveal new genes responsible for a MMAF phenotype in human and decipher how they affect sperm flagellum architecture and function. Overall, our strategy identified bi-allelic variants in *ZMYND12* associated with severe flagellum malformations resulting in asthenoteratozoospermia and primary male infertility. In conclusion,

our work with human sperm and the flagellated protist *T. brucei* demonstrated that ZMYND12 is part of the same axonemal complex as TTC29 and DNAH1, and plays a critical role in flagellum function and assembly.

# MATERIALS AND METHODS

## Study participants

WES data from a previously-established cohort of 167 MMAF individuals were analyzed (Coutton *et al*, 2019). All individuals presented a typical MMAF phenotype characterized by severe asthenozoospermia (total sperm motility below 10%), with >5% of the spermatozoa displaying at least three of the following flagellar abnormalities: short, absent, coiled, bent or irregular flagella (Coutton *et al*, 2019). All individuals had a normal somatic karyotype (46,XY) with normal bilateral testicular size, hormone levels, and secondary sexual characteristics. Sperm were analyzed in the source laboratories during routine biological examination of the individuals according to World Health Organization (WHO) guidelines (Wang *et al*, 2014). An additional Chinese individual was included in the study from a familial case. This individual was enrolled from the Human Sperm Bank of West China Second University Hospital of Sichuan University; he presented with severe asthenoteratozoospermia and a typical MMAF phenotype. WES analysis for this subject was performed as described in the previous report (Liu *et al*, 2021). Sperm morphology was assessed following Papanicolaou staining (Fig. 1A). Detailed semen parameters of the four individuals carrying *ZMYND12* mutations (*ZMYND12*_1-4) are presented in Table 1. Sperm samples for additional phenotypic characterization were obtained from *ZMYND12*_3. Written informed consent was obtained from all individuals participating in the study, and institutional approval was given by the local medical ethics committee (CHU Grenoble Alpes institutional review board). Samples were stored in the Fertithèque collection registered with the French Ministry of health (DC-2015-2580) and the French Data Protection Authority (DR-2016-392).

**Sanger sequencing**

*ZMYND12* single nucleotide variants identified by exome sequencing were validated by Sanger sequencing as previously described (Coutton *et al*, 2019). PCR primers and protocols used for each individual are listed in the Table S3.

**MLPA analysis**

Deletion of exons 6-8 identified in individual ZMYND12_2 by exome sequencing was confirmed by custom MLPA analysis, according to our previously described method (Coutton *et al*, 2012). The MLPA probes were designed, and the MLPA reaction was performed according to the recommendations set out in the MRC-Holland synthetic protocol ([www.mlpa.com](www.mlpa.com)). Data analysis was also performed in line with this protocol. For this study, three synthetic MLPA probes specific for exons 5, 6 and 8 of *ZMYND12* were designed. MLPA probes are listed in the Table S4.

**Quantitative real-time RT-PCR (RT-qPCR) analysis**

RT-qPCR experiments were performed as previously described (Coutton *et al*, 2019) with cDNAs extracted from various human tissues purchased from Life Technologies®. A panel of 10 organs was used for experiments: testis, brain, trachea, lung, kidney, liver, skeletal muscle, pancreas, placenta, and heart. Human RT-qPCR data were normalized relative to the two reference housekeeping genes *RPL6 and RPL27* by the $-\Delta\Delta C_t$ method (Livak & Schmittgen, 2001). Primer sequences and RT-qPCR conditions are indicated in Table S5. Relative expression of *ZMYND12* transcripts was compared in several organs, and statistical analysis involved a two-tailed *t*-test (Prism 4.0 software; GraphPad, San Diego, CA). A P-value $\leq 0.05$ was considered significant.

**Immunostaining in human sperm cells**

Immunofluorescence (IF) experiments were performed on sperm cells from control individuals, from individual ZMYND12_3 – carrier of the nonsense variant c.103C>T – and from a previously-reported MMAF individual – carrier of the *TTC29* splicing variant c.1761+1G>A (Lorès *et al*, 2019b). For each MMAF individual, 200 sperm cells were analyzed by two experienced operators, and the IF staining intensity and patterns were compared to those obtained from fertile control individuals. Sperm cells were fixed in phosphate-buffered saline (PBS)/4% paraformaldehyde for 1 min at room temperature. After washing in 1 mL PBS, the sperm suspension was spotted onto 0.1% poly L-lysine pre-coated slides (Thermo Scientific). After attachment, sperm were permeabilized with 0.1% (v/v) Triton X-100–DPBS (Triton X-100; Sigma-Aldrich) for 5 min at RT. Slides were then blocked by applying 5% normal serum–DPBS (normal goat or donkey serum; GIBCO, Invitrogen) before incubating overnight at 4 °C with the primary antibody. The following primary antibodies were used: ZMYND12, DNAI1, DNALI1, DNAH1, DNAH8, DNAH17, RSPH1, SPAG6, GAS8, CFAP70, WDR66 (also known as CFAP251), TTC29, AKAP4, and acetylated-α-tubulin. The primary antibody references and dilutions are fully detailed in Table S6. Slides were washed with 0.1% (v/v) Tween 20–DPBS before incubating for 1 h at room temperature with secondary antibodies. Highly cross-adsorbed secondary antibodies (Dylight 488 and Dylight 549, 1:1000) were purchased from Jackson Immunoresearch®. Experimental controls were treated similarly, but omitting the primary antibodies. Samples were counterstained with DAPI and mounted with DAKO mounting media (Life Technology). Fluorescence images were captured with a confocal microscope (Zeiss LSM 710).

***Trypanosoma brucei* cell lines, culture and transfection**

The trypanosome cell lines used in this study are derived from the procyclic parental form, *T. brucei* SmOxP427(Sunter, 2016). These cells co-express the T7 RNA polymerase and

the tetracycline repressor and were grown and transfected as previously described (Kherraf *et al*, 2018b). When required, the culture medium was supplemented with puromycin (1 µg/mL), neomycin (10 µg/mL), phleomycin (5 µg/mL), or hygromycin (25 µg/mL). Cells were transfected using an AMAXA electroporator (Lonza) as previously described (Schumann Burkard *et al*, 2011). *TbTAX-1* RNA interference was induced with tetracycline (10 µg/mL).

Tagged proteins were obtained from SmOxP427 cells, by inserting epitope-tagged transgenes at the endogenous locus. Transgenes were N-terminally or C-terminally tagged with 10myc, 10Ty1, or 10HA tags using the pPOTv7 vector series, as previously described (Dean *et al*, 2015; Kherraf *et al*, 2018b). For RNAi$^{TbTAX-1}$ knock-down, bp 219-667 were cloned between the *XhoI* and *XbaI* sites of p2T7tiB (LaCount *et al*, 2002) and the linearized vector was transfected into endogenously-tagged TbTAX-1-background cell lines. To target the full-length *TbTAX-1* and not the shorter form, similar cloning was performed with bp 493-917. The TbTTC29$_{Ty1}$ RNAi cell line is described elsewhere (Lorès *et al*, 2019b).

**Co-immunoprecipitation and Mass spectrometry on *Trypanosoma* cells**

In a 15-mL polycarbonate tube, $2.10^8$ cells expressing the different tagged proteins were washed in PBS and incubated for 5 min at RT in 1 mL of immunoprecipitation buffer for cell lysis (IP buffer; 25 mM Tris-HCl pH 7.4, 100 mM NaCl, 0.25% Nonidet P-40 (Igepal), 1 mM DTT, protease inhibitor cocktail, and 1 unit of nuclease) (Pierce). Lysis was completed by sonication (5x 30 s ON/30 s OFF) in a Bioruptor sonication system. Cell lysates were clarified by centrifugation (10 min, 16,000 x g, 4 °C) and split into two samples each. One was incubated with 3 µL of anti-Ty1 and the other with 2 µL of mouse anti-HA for 1 h at 4 °C on a rotating wheel. Protein-G Dynabead (Invitrogen 10003D) slurry (40 µL) was washed in IP buffer and incubated with the samples for 1 h at 4 °C. After washing three times in IP buffer, beads were resuspended in 40 µL sample buffer and boiled before loading 20 µL on SDS-PAGE.

For mass spectrometry analyses, $1.8 \times 10^9$ cells were washed in PBS, resuspended in 2 mL of immunoprecipitation buffer without DTT and processed as described above. Antibodies used were: mouse anti-HA (10 µL) and anti-TY1 (15 µL). Complex precipitation was achieved by adding 200 µL of Protein-G Dynabead slurry. After washing three times in IP buffer, beads were transferred to new tubes and washed 3x in PBS before storing dry at -20 °C.

Protein-G Dynabeads were resuspended in Laemmli buffer and boiled. Supernatants were loaded on a 10% acrylamide SDS-PAGE gel, and proteins were revealed by Colloidal Blue staining. Sample preparation and protein digestion by trypsin were performed as previously described (Allmann *et al*, 2014). NanoLC-tandem mass spectrometry (MS/MS) analyses were performed using an Ultimate 3000 RSLC Nano-UPHLC system (Thermo Scientific, USA) coupled to a nanospray Orbitrap Fusion™ Lumos™ Tribrid™ Mass Spectrometer (Thermo Fisher Scientific, California, USA). Each peptide extract was loaded on a 300 µm ID x 5-mm PepMap C18 pre-column (Thermo Scientific, USA) at a flow rate of 10 µL/min. After a 3-min desalting step, peptides were separated on a 50-cm EasySpray column (75 µm ID, 2 µm C18 beads, 100 Å pore size; ES903, Thermo Fisher Scientific) by applying a 4-40% linear gradient of solvent B (0.1% formic acid in 80% ACN) over 48 min. The flow rate for separation was set to 300 nL/min. The mass spectrometer was operated in positive ion mode at a 2.0 kV needle voltage. Data were acquired using Xcalibur 4.4 software in data-dependent mode. MS scans (m/z 375-1500) were recorded at a resolution of R = 120,000 (@ m/z 200) and an Automated Gain Control (AGC) target of $4 \times 10^5$ ions collected within 50 ms was applied along with a top speed duty cycle of up to 3 seconds for MS/MS acquisition. Precursor ions (2 to 7 charge states) were isolated in the quadrupole within a mass window of 1.6 Th, and fragmented by Higher energy Collisional Dissociation (HCD), applying 28% normalized collision energy. MS/MS data were acquired in the Orbitrap cell at a resolution of R=30000 (m/z 200), a standard AGC target and a maximum injection time in automatic mode. A 60-s

exclusion window was applied to previously selected precursors. Proteome Discoverer 2.5 was used to identify proteins and perform Label-Free Quantification (LFQ). Proteins were identified in batch mode using MS Amanda 2.0, Sequest HT and Mascot 2.5 algorithms, with searches performed against a *Trypanosoma brucei brucei* TREU927 protein database (9,788 entries, release 57, https://tritrypdb.org/ website). Up to two missed trypsin cleavages were allowed. Mass tolerances in MS and MS/MS were set to 10 ppm and 0.02 Da, respectively. Oxidation (M) and acetylation (K) were included as dynamic modifications, and carbamidomethylation (C) as a static modification. Peptides were validated using the Percolator algorithm (Käll *et al*, 2007), retaining only "high confidence" peptides, corresponding to a 1% false discovery rate at the peptide level. The minora feature-detector node (LFQ) was used along with the feature-mapper and precursor ion quantifier. The following quantification parameters were applied: (1) Unique peptides (2) Precursor abundance based on intensity (3) No normalization (4) Protein abundance calculation: summed abundances (5) Protein ratio calculation: pairwise ratios (6) Imputation mode: Low abundance resampling and (7) Hypothesis test: t-test (background-based). For master proteins, quantitative data were considered based on a minimum of two unique peptides, a fold-change greater than 2 and a Benjamini-Hochberg-corrected p-value for the FDR of less than 0.05. Mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium *via* the PRIDE partner repository (Deutsch *et al*, 2020), under dataset identifier PXD039470.

**Immunofluorescence on *Trypanosoma* cells**

Cells were harvested, washed, and processed for immunolabelling on methanol-fixed detergent-extracted cells (cytoskeleton, CSK) as described in (Albisetti *et al*, 2017). The antibodies used and their dilutions are listed in Table S7. Images were acquired on a Zeiss Imager Z1 microscope, using a Photometrics Coolsnap HQ2 camera, with a 100x Zeiss objective (NA 1.4) using Metamorph software (Molecular Devices), and processed with ImageJ

(NIH).

**Ultrastructure expansion microscopy on *Trypanosoma* cells**

The U-ExM protocol has been adapted and optimized for use on trypanosomes, as described (Gambarotto *et al*, 2021). Protocol details can be found online at https://doi.org/10.17504/protocols.io.bvwqn7dw (Casas *et al*, 2022). Images were acquired on a Zeiss AxioImager Z1 using a 63x (1.4 NA) objective, or at the Bordeaux Imaging Center with a Leica TCS SP5 on an upright stand DM6000 (Leica Microsystems, Mannheim, Germany), fitted with a HCX Plan Apo CS 63X oil NA 1.40 objective. Images were processed and analyzed using ImageJ and Fiji software (Schindelin *et al*, 2012; Schneider *et al*, 2012).

***Trypanosoma* cells: sedimentation assays and video microscopy**

Sedimentation assays were performed as described (Dacheux *et al*, 2012a). Briefly, cells were placed in cuvettes and incubated for 24 h without shaking. Optical density ($OD_{600nm}$) was measured before mixing (ODb, to detect "swimming" cells) and after mixing (ODa, to detect "swimming" plus "sedimenting" cells). The percentage of sedimented cells was calculated as 100-(ODb/ODa) x100 for the WT and RNAi cells. Sedimentation of RNAi cells was normalized relative to the tagged TbTAX-1 parental cell line. Video microscopy was carried out as described (Lorès *et al*, 2019b). Briefly, PCF cells (WT cell and RNAi$^{TbTAX-}$1 cells 72 h after induction) were washed in PBS, and cell mobility was recorded by phase-contrast microscopy on a Zeiss AxioImager Z1, 40x lens (NA 1.3). A 28-s sequence of digital video was captured from separate regions and analyzed using Metamorph software (Molecular Devices) (speed x8). The positions of individual cells were plotted over 0.28-s intervals. The start and end positions of each cell were marked.

**Transmission Electron Microscopy (TEM) on *Trypanosoma* cells**

TEM blocks of Trypanosoma cells and thin sections were prepared as previously described (Albisetti *et al*, 2017). Thin sections were visualized on an FEI Tecnai 12 electron microscope. Images were captured on an ORIUS 1000 11M Pixel camera (resolution 3±5 nm) controlled by Digitalmicrograph, and processed using ImageJ (NIH).

**Western blot analyses of *Trypanosoma* cells**

Proteins from whole cells, detergent-extracted cytoskeletons and flagella, or co-IPs were separated on SDS-PAGE gels. Separated proteins were transferred by semi-dry transfer to PVDF and low-fluorescence PVDF membranes for further processing, as described (Albisetti *et al*, 2017). The antibodies used and their dilutions are listed in Table S7. For IP analysis, primary antibodies – anti-TY1 and rabbit anti-HA (GTX115044) – and fluorescent secondary antibodies – anti-mouse Starbright Blue 520 (Bio-Rad 12005867) and anti-rabbit Starbright Blue 700 (Bio-Rad 12004162) – were used.

**Co-immunoprecipitation and mass spectrometry analysis on *Ttc29*[-/-] mouse testes**

Testes from wild type and Ttc29-/- L5 and L7 mice (Lorès *et al*, 2019b) were dissected in cold PBS; the albuginea was removed and the seminiferous tubules were dissociated and pelleted by brief centrifugation. Proteins from the pellets were extracted in 1 mL lysis buffer: 50 mM Tris-HCl, pH 8, 150 mM NaCl, 10 mM MgCl2, 0.5% NP40, and Complete protease inhibitor cocktail (Roche Applied science). Immunoprecipitation was performed on total-protein lysate using 1 µg of rabbit polyclonal TTC29 antibody (HPA061473, Sigma-Aldrich) and 15 µl of Protein A/G magnetics beads (Bio-Ademtec) according to the manufacturer's instructions. The experiment was replicated with four mice for each genotype.

Total immunoprecipitated proteins were recovered in 25 µL 100 mM Tris/HCl pH 8.5 containing 10 mM TCEP, 40 mM chloroacetamide and 1% sodium deoxycholate, by sonicating three times, and incubating for 5 min at 95 °C. An aliquot (20 µg) of protein lysate was diluted (1:1) in Tris 25 mM pH 8.5 in 10% Acetonitrile (ACN) and subjected to overnight trypsin

digestion with 0.4 µg of sequencing-grade bovine trypsin (Promega) at 37 °C. Deoxycholate was removed using liquid-liquid phase extraction after adding 50 µL of 1% TriFluoroacetic Acid (TFA) in ethyl-acetate. A six-layer home-made Stagetip with Empore Styrene- divinylbenzene - Reversed Phase Sulfonate (SDB RPS; 3M) disks was used. Eluted peptides were dried and re-solubilized in 2% TFA before preparing five fractions on hand-made Strong Cation eXchange (SCX) StageTips (Kulak *et al*, 2014). Fractions were analyzed on a system combining an U3000 RSLC nanochromatograph eluting into an Orbitrap Fusion mass spectrometer (both Thermo Scientific). Briefly: peptides from each SCX fraction were separated on a C18 reverse phase column (2 µm particle size, 100 Å pore size, 75 µm inner diameter, 25 cm length) with a 145-min linear gradient starting from 99% solvent A (0.1% formic acid in $H_2O$) and ending with 40% of solvent B (80% ACN and 0.085% formic acid in $H_2O$). The MS1 scans spanned 350-1500 m/z with an AGC target of $1.10^6$ within a maximum ion injection time (MIIT) of 60 ms, at a resolution of 60,000. The precursor selection window was set to 1.6 m/z with quadrupole filtering. HCD Normalized Collision Energy was set to 30%, and the ion trap scan rate was set to "rapid" mode with AGC target $1.10^5$ and 60 ms MIIT. Mass spectrometry data were analyzed using Maxquant (v.1.6.1.0)(Cox *et al*, 2014). The database used was a concatenation of human sequences from the Uniprot-Swissprot database (release 2017-05) and the Maxquant list of contaminant sequences. Cysteine carbamidomethylation was set as fixed modification, and acetylation of protein N-terminus and oxidation of methionine were set as variable modifications. Second peptide search and "match between runs" (MBR) options were allowed. Label-free protein quantification (LFQ) was performed using both unique and razor peptides, requiring at least two peptides per protein. The quality of raw quantitative data was evaluated using PTXQC software v. 0.92.3 (Bielow *et al*, 2016). Perseus software, version 1.6.2.3 (Tyanova *et al*, 2016) was used for statistical analysis and data comparison. Proteomics data obtained by mass spectrometry analysis were deposited to the

ProteomeXchange Consortium *via* the PRIDE partner repository (Deutsch *et al*, 2020) under

dataset identifier PXD039854.

**A**

**B**

**ZMYND12** *(ENST00000372565.8)*

c.443C>T

Del.ex6_ex8

1    100    200    300    366

N    C

p.Arg145Ter

■ Zinc-finger domain
■ Tetratricopeptide-like helical domain

**C**                          **D**

*ZMYND12* Exon 4

Ref.Seq.

Control

ZMYND12_1

ZMYND12_3

ZMYND12_4

ZMYND12_2

ZMYND12_ex8    c1    ZMYND12_ex6    c2    ZMYND12_ex5    c3

Control

**E**

MERGE    DAPI    ZMYND12    TUBULIN

Control

ZMYND12_3

**Fig. 1. Morphology of normal and *ZMYND12* mutant spermatozoa and variants identified in *ZMYND12* individuals.** (**A**) Light microscopy analysis of spermatozoa from fertile control individuals and from individuals ZMYND12_2 and ZMYND12_3. Most spermatozoa from ZMYND12 individuals had short (#), coiled (@), or irregular caliber (&) flagella. Head malformations were also observed (δ). (**B**) Location of the variants identified within the *ZMYND12* gene (ENST00000372565.8) and protein according to UniprotKB (Q9H0C1). The blue box represents a Zinc Finger domain and the orange boxes represent TPR domains (tetratricopeptide repeat). (**C**) Sanger sequencing electropherograms indicating the homozygous state of the *c.433C>T* variant located in exon 4 in the three individuals ZMYND12_1,3 and 4. The substituted nucleotide is highlighted in red. Variants are annotated in line with HGVS recommendations. (**D**) Confirmation of *ZMYND12* exon deletion by MLPA. MLPA profiles obtained for control and ZMYND12_2 individuals. Each set contains three control probes for normalization purposes (c1-c3) and three *ZMYND1*2-specific probes. Blue arrows indicate homozygous deletion of exons 6 and 8 in the *ZMYND12* gene. *ZMYND12* exon 5 is not included in the deletion. (**E**) Sperm cells from a fertile control individual and from individual ZMYND12_3 – carrier of the *c.433C>T* variant – were stained with anti-ZMYND12 (green) and anti-acetylated tubulin (red) antibodies. Sperm nuclear DNA was counterstained with DAPI (blue). In control sperm, ZMYND12 immunostaining was present along the whole length of the flagellum; no staining was observed in sperm cells from individual ZMYND12_3. Scale bars: 10 μm.

**Fig. 2. Altered immunostaining for SPAG6, DNAH1, DNALI1, WDR66, CFAP70, and TTC29 in the presence of *ZMYND12* variants.** Immunofluorescence experiments were performed using sperm cells from control individuals and from individual ZMYND12_3 carrying the nonsense variant *c.433C>T*. Although tubulin staining remained detectable, sperm cells from individual ZMYND12_3 showed no immunostaining for (**A**) SPAG6, (**B**) DNAH1, (**C**) DNALI1, (**D**) WDR66, (**E**) TTC29, and (**F**) CFAP70. The CPC, IDAs, and the CSC were therefore strongly disorganized in these cells. Scale bars: 10 µm.

**Fig. 3**. **TbTAX-1 is an axoneme-associated protein; its knock-down leads to flagellar motility defects.** (**A**) Representative image of a post-mitotic detergent-extracted cell immunolabelled with anti-PFR2 (magenta) staining the paraflagellar rod (a para-axonemal structure) and anti-myc (yellow) to reveal $_{myc}$TbTAX-1. The protein was detected on the axoneme in both the old flagellum (OF) and the new flagellum (NF). Immunolabelling indicated that the axonemal localization of $_{myc}$TbTAX-1 extended distally along the flagellum beyond the region labelled by PFR2 (see zoom). The mitochondrial genome (kinetoplast, k) and nuclei (N) were stained with DAPI. The outline of the cell body is indicated by the dashed white line. Scale bars: 5 μm in whole-cell images; 1 μm in zoomed images. (**B**) Anti-myc western blot analysis of RNAi$^{TbTAX-1}$ knock-down 24, 48, or 72 h post-induction. $_{myc}$TbTAX-1 protein levels were reduced in induced cells (I) compared to non-induced cells (NI); TbSAXO, a flagellum-specific protein, was used as loading control. (**C**) Representative image of a detergent-extracted post-mitotic cell after RNAi$^{TbTAX-1}$ induction, showing a strong decrease in $_{myc}$TbTAX-1 labeling in the NF compared to the OF. (**D**) Comparative *T. brucei* growth curves for wild-type (WT), non-induced (NI), and tetracycline-induced RNAi$^{TbTAX-1}$ cell lines at 0-, 24-, 48- and 72-hours post-induction. (**E**) Mobility tracking from video microscopy recordings of live cells: WT and RNAi$^{TbTAX-1}$ after induction for 72 h. The positions of individual cells are plotted at 0.28-s intervals; circles indicate start positions, arrowheads indicate end positions. A dramatic loss of progressive mobility was observed after TbTAX1 protein knock-down. Scale bars: 20 μm. (**F**) Sedimentation assays for *T. brucei* WT, non-induced (NI), and tetracycline-induced RNAi$^{TbTAX-1}$ at 24, 48, and 72 h post-induction. The percentage of cells sedimenting increased from 20 to 70% after depletion of the TbTAX-1 protein.

**A**

TbTAX-1

RNAi targeted sequence in
TbTAX-1 ORF

1 | TPR-like domain | HA | 363 aa

TbTAX-1$^{Trunc}$

1 | Ty1 | 128 aa

**B**

| kDa | WC | CSK | FG | |
|---|---|---|---|---|
| 70- | | | | TbTAX-1$_{HA}$ |
| 40- | | | | TbTAX-1$^{Trunc}_{Ty1}$ |
| 40- | | | | Enolase |
| 55- | | | | Tubulin |

**C**

TbTAX-1$_{HA}$     TbTAX-1$^{Trunc}_{Ty1}$

**D**

| kDa | NI | 24H | 48H | 72H | 96H | |
|---|---|---|---|---|---|---|
| 70- | | | | | | TbTAX-1$_{HA}$ |
| 35- | | | | | | TbTAX-1$^{Trunc}_{Ty1}$ |
| 40- | | | | | | Enolase |

**E**

LOG number of cells

— WT
— NI
---- RNAi$^{TbTAX-1}$

Hours of induction

**F**

% sedimentation

WT   NI   24H   48H   72H

**Fig. 4. The truncated form of TbTAX-1 is not functional.** (**A**) Schematic representation of full-length TbTAX-1 and its truncated form. The TPR-like domain is illustrated in light blue, and the targeted RNA interference sequence in dark blue. (**B**) Western blot analysis of subcellular fractions to determine localization of TbTAX-1$_{HA}$ and its truncated form TbTAX-1$^{Trunc}_{Ty1}$. Enolase (cytoplasm) and tubulin (cytoskeleton) were used as loading controls. (**C**) Immunofluorescence-labeling of detergent-extracted cytoskeletons to detect TbTAX-1$_{HA}$ (yellow) and TbTAX-1$^{Trunc}_{Ty1}$ (stained green, but not visible as the protein was eliminated during detergent extraction). Scale bars: 5 μm. (**D**) Western blot analysis of the impact of RNAi$^{TbTAX-1}$ expression on TbTAX-1$^{Trunc}_{Ty1}$ levels. Enolase was used as a loading control. (**E**) Growth curves of WT, non-induced (NI), and RNAi$^{TbTAX-1}$ induced cells expressing TbTAX-1$_{HA}$ and TbTAX-1$^{Trunc}_{Ty1}$. (**F**) Sedimentation assays for *T. brucei* WT, non-induced (NI), and RNAi$^{TbTAX-1}$ induced cells at 24, 48, and 72 h.

**A** TbTTC29$_{10Ty1}$/TbTAX-1$_{10HA}$/***TbTAX-1$^{RNAi}$***

NI  24  48  72

kDa

70 — TbTAX-1$_{10HA}$

100 —

70 — TbTTC29$_{10Ty1}$

55 — Enolase

■ TbTAX-1$_{10HA}$
▨ TbTTC29$_{10Ty1}$

Change in protein level

100% 80% 60% 40% 20% 0%

0  24  48  72

Time of induction (Hr)

**B** TbTTC29$_{10Ty1}$/TbTAX-1$_{10HA}$/***TbTTC29$^{RNAi}$***

NI  24  48  72

kDa

70 — TbTAX-1$_{10HA}$

100 —

70 — TbTTC29$_{10Ty1}$

55 — Enolase

▨ TbTTC29$_{10Ty1}$
■ TbTAX-1$_{10HA}$

change in protein level

100% 80% 60% 40% 20% 0%

0  24  48  72

Time of induction (Hr)

**C**

NF
N  N
k  k
OF

TbTAX-1$_{10HA}$   TbTTC29$_{10TY1}$

**D** TbTAX-1$_{10HA}$  TbTTC29$_{10TY1}$

10 µm

Intensity (A.U.)

120 100 80 60 40 20 0

TbTAX 1$_{10HA}$
TbTTC29$_{10TY1}$

0  5  10  15

Distance (um)

**E** TbTTC29$_{TY1}$+TbTAX-1$_{10HA}$   TbTTC29$_{TY1}$   TbTAX-1$_{10HA}$

Input  IP$_{TY1}$  IP$_{HA}$   Input  IP$_{TY1}$  IP$_{HA}$   Input  IP$_{TY1}$  IP$_{HA}$

TbTTC29$_{10TY1}$ (anti-Ty1)

TbTAX-1$_{10HA}$ (anti-HA)

**Fig. 5. TbTAX-1 and TbTTC29 are part of the same protein complex.** (**A**) Analysis of the impact of *TbTAX-1* RNAi on TbTTC29$_{Ty1}$ protein levels. Western blot (upper panel) and quantification (lower panel). (**B**) Analysis of the impact of *TbTTC29* RNAi induction on TbTAX-1$_{HA}$ protein levels. Western blot (upper panel) and quantification (lower panel). (**C**) U-ExM analysis of TbTAX-1$_{HA}$ (yellow) and TbTTC29$_{TY1}$ (purple) colocalization in a dividing cell with an old flagellum (OF) and a new flagellum (NF). Kinetoplast (k) and nuclei (N) were counterstained with Hoechst (blue). (**D**) Confocal image of U-ExM labelling of TbTAX-1$_{HA}$ (yellow) and TbTTC29$_{TY1}$ (magenta) (upper panel). Lower panel: profile plot along the white line showing overlapping intensity peaks for TbTAX-1 and TbTTC29, confirming true colocalization. (**E**) Anti-Ty1 (IP$_{Ty1}$) or anti-HA (IP$_{HA}$) co-immunoprecipitation assays in cell lines expressing both TbTAX-1$_{HA}$ and TbTTC29$_{Ty1}$, or TbTTC29$_{Ty1}$ alone or TbTAX-1$_{HA}$ alone to verify IP specificity. Proteins were detected simultaneously by western blotting using anti-Ty1 and anti-HA antibodies and fluorescent anti-mouse (detecting mouse anti-Ty1) and anti-rabbit (detecting rabbit anti-HA) secondary antibodies. IP assays on cell lines expressing only one or the other tagged protein did not pull down the other tagged protein.

**Table 1. Detailed semen parameters in the four MMAF individuals harboring a *ZMYND12* variant.**

| ZMYND12 mutated individuals | | Semen parameters | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Individuals | *ZMYND12* variant | Sperm volume (mL) | Sperm concentration ($10^6$/mL) | Total motility 1 h | Vitality | Normal spermatozoa | Absent flagella | Short Flagella | Coiled Flagella | Bent Flagella | Flagella of irregular caliber | Tapered head | Thin head | Micro-cephalic | Macro-cephalic | Multiple heads | Abnormal base | Abnormal acrosomal region |
| ZMYND12_1 | c.433C>T | 2 | 6.5 | 4 | 70 | 0 | 35 | 48 | 10 | 0 | 34 | 22 | 2 | 2 | 4 | 6 | 20 | 80 |
| ZMYND12_2 | Del ex6-8 | 2.5 | 4.25 | 2 | NA | 0 | NA | 72 | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA |
| ZMYND12_3 | c.433C>T | 3 | 34 | 2 | 50 | 0 | 20 | 20 | 28 | 0 | 46 | 22 | 6 | 8 | 2 | 2 | 24 | 88 |
| ZMYND12_4 | c.433C>T | 2.6 | 0.1 | 0 | NA | NA | NA | 70 | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA |
| **Reference limits**[a] | | 1.5 (1.4-1.7) | 15 (12-16) | 40 (38-42) | 58 (55-63) | 23 (20-26) | 5 (4-6) | 1 (0-2) | 17 (15-19) | 13 (11-15) | 2 (1-3) | 3 (2-4) | 14 (12-16) | 7 (5-9) | 1 (0-2) | 2 (1-3) | 42 (39-45) | 60 (57-63) |

Values are percentages unless specified otherwise. NA: not available

[a] Reference limits (5th centiles and their 95% confidence intervals) according to World Health Organization (WHO) standards (Cooper *et al*, 2010) and the distribution range of morphologically normal spermatozoa observed in 926 fertile individuals (Auger *et al*, 2016).

# REFERENCES AND NOTES

Albisetti A, Florimond C, Landrein N, Vidilaseris K, Eggenspieler M, Lesigang J, Dong G, Robinson DR & Bonhivers M (2017) Interaction Between the Flagellar Pocket Collar and the Hook Complex via a Novel Microtubule-binding Protein in Trypanosoma brucei. *PLOS Pathog* 13: e1006710

Allmann S, Mazet M, Ziebart N, Bouyssou G, Fouillen L, Dupuy J-W, Bonneu M, Moreau P, Bringaud F & Boshart M (2014) Triacylglycerol Storage in Lipid Droplets in Procyclic Trypanosoma brucei. *PLOS ONE* 9: e114628

Arafah K, Lopez F, Cazin C, Kherraf Z-E, Tassistro V, Loundou A, Arnoult C, Thierry-Mieg N, Bulet P, Guichaoua M-R, *et al* (2021) Defect in the nuclear pore membrane glycoprotein 210-like gene is associated with extreme uncondensed sperm nuclear chromatin and male infertility: a case report. *Hum Reprod Oxf Engl* 36: 693–701

Aslett M, Aurrecoechea C, Berriman M, Brestelli J, Brunk BP, Carrington M, Depledge DP, Fischer S, Gajria B, Gao X, *et al* (2010) TriTrypDB: a functional genomic resource for the Trypanosomatidae. *Nucleic Acids Res* 38: D457-462

Auger J, Jouannet P & Eustache F (2016) Another look at human sperm morphology. *Hum Reprod Oxf Engl* 31: 10–23

Baron DM, Ralston KS, Kabututu ZP & Hill KL (2007) Functional genomics in Trypanosoma brucei identifies evolutionarily conserved components of motile flagella. *J Cell Sci* 120: 478–91

Ben Khelifa M, Coutton C, Zouari R, Karaouzène T, Rendu J, Bidart M, Yassine S, Pierre V, Delaroche J, Hennebicq S, *et al* (2014) Mutations in DNAH1, which encodes an inner arm heavy chain dynein, lead to male infertility from multiple morphological abnormalities of the sperm flagella. *Am J Hum Genet* 94: 95–104

Bielow C, Mastrobuoni G & Kempa S (2016) Proteomics Quality Control: Quality Control Software for MaxQuant Results. *J Proteome Res* 15: 777–787

Blackburn K, Bustamante-Marin X, Yin W, Goshe MB & Ostrowski LE (2017) Quantitative Proteomic Analysis of Human Airway Cilia Identifies Previously Uncharacterized Proteins of High Abundance. *J Proteome Res* 16: 1579–1592

Broadhead R, Dawe HR, Farr H, Griffiths S, Hart SR, Portman N, Shaw MK, Ginger ML, Gaskell SJ, McKean PG, *et al* (2006a) Flagellar motility is required for the viability of the bloodstream trypanosome. *Nature* 440: 224–7

Broadhead R, Dawe HR, Farr H, Griffiths S, Hart SR, Portman N, Shaw MK, Ginger ML, Gaskell SJ, McKean PG, *et al* (2006b) Flagellar motility is required for the viability of the bloodstream trypanosome. *Nature* 440: 224–227

Brookman JL, Stott AJ, Cheeseman PJ, Burns NR, Adams SE, Kingsman AJ & Gull K (1995) An immunological analysis of Ty1 virus-like particle structure. *Virology* 207: 59–67

Brooks ER (2014) Control of intraflagellar transport : studies of the planar cell polarity effector Fuz, the small GTPase Rsg1, and the novel protein TTC29.

Casas E, Landrein N & Bonhivers M (2022) Ultra Expansion microscopy protocol with improved setup for upright and inverted microscopes. *protocols.io*

Cooper TG, Noonan E, von Eckardstein S, Auger J, Baker HWG, Behre HM, Haugen TB, Kruger T, Wang C, Mbizvo MT, *et al* (2010) World Health Organization reference values for human semen characteristics. *Hum Reprod Update* 16: 231–245

Coutton C, Escoffier J, Martinez G, Arnoult C & Ray PF (2015) Teratozoospermia: spotlight on the main genetic actors in the human. *Hum Reprod Update* 21: 455–485

Coutton C, Martinez G, Kherraf Z-E, Amiri-Yekta A, Boguenet M, Saut A, He X, Zhang F, Cristou-Kent M, Escoffier J, *et al* (2019) Bi-allelic Mutations in ARMC2 Lead to Severe Astheno-Teratozoospermia Due to Sperm Flagellum Malformations in Humans and Mice. *Am J Hum Genet* 104: 331–340

Coutton C, Vargas AS, Amiri-Yekta A, Kherraf Z-E, Ben Mustapha SF, Le Tanno P, Wambergue-Legrand C, Karaouzène T, Martinez G, Crouzy S, *et al* (2018a) Mutations in CFAP43 and CFAP44 cause male infertility and flagellum defects in Trypanosoma and human. *Nat Commun* 9: 686

Coutton C, Vargas AS, Amiri-Yekta A, Kherraf Z-E, Mustapha SF, Tanno P, Wambergue-Legrand C, Karaouzène T, Martinez G, Crouzy S, *et al* (2018b) Mutations in CFAP43 and CFAP44 cause male infertility and flagellum defects in Trypanosoma and human. *Nat Commun* 9: 686

Coutton C, Zouari R, Abada F, Ben Khelifa M, Merdassi G, Triki C, Escalier D, Hesters L, Mitchell V, Levy R, *et al* (2012) MLPA and sequence analysis of DPY19L2 reveals point mutations causing globozoospermia. *Hum Reprod Oxf Engl* 27: 2549–2558

Cox J, Hein MY, Luber CA, Paron I, Nagaraj N & Mann M (2014) Accurate proteome-wide label-free quantification by delayed normalization and maximal peptide ratio extraction, termed MaxLFQ. *Mol Cell Proteomics MCP* 13: 2513–2526

Crozier TWM, Tinti M, Wheeler RJ, Ly T, Ferguson MAJ & Lamond AI (2018) Proteomic Analysis of the Cell Cycle of Procylic Form Trypanosoma brucei. *Mol Cell Proteomics MCP* 17: 1184–1195

Dacheux D, Landrein N, Thonnus M, Gilbert G, Sahin A, Wodrich H, Robinson DR & Bonhivers M (2012a) A MAP6-Related Protein Is Present in Protozoa and Is Involved in Flagellum Motility. *PLoS One* 7: e31344

Dacheux D, Landrein N, Thonnus M, Gilbert G, Sahin A, Wodrich H, Robinson DR & Bonhivers M (2012b) A MAP6-related protein is present in protozoa and is involved in flagellum motility. *PloS One* 7: e31344

D'Andrea LD & Regan L (2003) TPR proteins: the versatile helix. *Trends Biochem Sci* 28: 655–662

Darde TA, Lecluze E, Lardenois A, Stévant I, Alary N, Tüttelmann F, Collin O, Nef S, Jégou B, Rolland AD, *et al* (2019) The ReproGenomics Viewer: a multi-omics and cross-species resource compatible with single-cell studies for the reproductive science community. *Bioinforma Oxf Engl*

Dean S, Moreira-Leite F, Varga V & Gull K (2016) Cilium transition zone proteome reveals compartmentalization and differential dynamics of ciliopathy complexes. *Proc Natl Acad Sci U S A* 113: E5135-5143

Dean S, Sunter J, Wheeler RJ, Hodkinson I, Gluenz E & Gull K (2015) A toolkit enabling efficient, scalable and reproducible gene tagging in trypanosomatids. *Open Biol* 5: 140197

Dean S, Sunter JD & Wheeler RJ (2017) TrypTag.org: A Trypanosome Genome-wide Protein Localisation Resource. *Trends Parasitol* 33: 80–82

Deutsch EW, Bandeira N, Sharma V, Perez-Riverol Y, Carver JJ, Kundu DJ, García-Seisdedos D, Jarnuczak AF, Hewapathirana S, Pullman BS, *et al* (2020) The ProteomeXchange consortium in 2020: enabling 'big data' approaches in proteomics. *Nucleic Acids Res* 48: D1145–D1152

Dymek EE, Heuser T, Nicastro D & Smith EF (2011) The CSC is required for complete radial spoke assembly and wild-type ciliary motility. *Mol Biol Cell* 22: 2520–2531

Ersfeld K & Gull K (2001a) Targeting of cytoskeletal proteins to the flagellum of Trypanosoma brucei. *J Cell Sci* 114: 141–148

Ersfeld K & Gull K (2001b) Targeting of cytoskeletal proteins to the flagellum of Trypanosoma brucei. *J Cell Sci* 114: 141–148

Gambarotto D, Hamel V & Guichard P (2021) Chapter 4 - Ultrastructure expansion microscopy (U-ExM). In *Methods in Cell Biology*, Guichard P & Hamel V (eds) pp 57–81. Academic Press

Gambarotto D, Zwettler FU, Guennec ML, Schmidt-Cernohorska M, Fortun D, Borgers S, Heine J, Schloetel J-G, Reuss M, Unser M, *et al* (2019) Imaging cellular ultrastructures using expansion microscopy (U-ExM). *Nat Methods* 16: 71–74

Hannaert V, Albert M-A, Rigden DJ, da Silva Giotto MT, Thiemann O, Garratt RC, Van Roy J, Opperdoes FR & Michels PAM (2003) Kinetic characterization, structure modelling studies and crystallization of Trypanosoma brucei enolase. *Eur J Biochem* 270: 3205–3213

Kalichava A & Ochsenreiter T Ultrastructure expansion microscopy in Trypanosoma brucei. *Open Biol* 11: 210132

Käll L, Canterbury JD, Weston J, Noble WS & MacCoss MJ (2007) Semi-supervised learning for peptide identification from shotgun proteomics datasets. *Nat Methods* 4: 923–925

Kherraf Z-E, Amiri-Yekta A, Dacheux D, Karaouzène T, Coutton C, Christou-Kent M, Martinez G, Landrein N, Le Tanno P, Fourati Ben Mustapha S, *et al* (2018a) A Homozygous Ancestral SVA-Insertion-Mediated Deletion in WDR66 Induces Multiple Morphological Abnormalities of the Sperm Flagellum and Male Infertility. *Am J Hum Genet* 103: 400–412

Kherraf Z-E, Amiri-Yekta A, Dacheux D, Karaouzène T, Coutton C, Christou-Kent M, Martinez G, Landrein N, Le Tanno P, Fourati Ben Mustapha S, *et al* (2018b) A Homozygous Ancestral SVA-Insertion-Mediated Deletion in WDR66 Induces Multiple Morphological Abnormalities of the Sperm Flagellum and Male Infertility. *Am J Hum Genet* 103: 400–412

Kim S, Scheffler K, Halpern AL, Bekritsky MA, Noh E, Källberg M, Chen X, Kim Y, Beyter D, Krusche P, *et al* (2018) Strelka2: fast and accurate calling of germline and somatic variants. *Nat Methods* 15: 591–594

King SM (2013) A solid-state control system for dynein-based ciliary/flagellar motility. *J Cell Biol* 201: 173–175

Kulak NA, Pichler G, Paron I, Nagaraj N & Mann M (2014) Minimal, encapsulated proteomic-sample processing applied to copy-number estimation in eukaryotic cells. *Nat Methods* 11: 319–324

LaCount DJ, Barrett B & Donelson JE (2002) Trypanosoma brucei FLA1 is required for flagellum attachment and cytokinesis. *J Biol Chem* 277: 17580–8

Liu C, He X, Liu W, Yang S, Wang L, Li W, Wu H, Tang S, Ni X, Wang J, *et al* (2019a) Bi-allelic Mutations in TTC29 Cause Male Subfertility with Asthenoteratospermia in Humans and Mice. *Am J Hum Genet* 105: 1168–1181

Liu C, Tu C, Wang L, Wu H, Houston BJ, Mastrorosa FK, Zhang W, Shen Y, Wang J, Tian S, *et al* (2021) Deleterious variants in X-linked CFAP47 induce asthenoteratozoospermia and primary male infertility. *Am J Hum Genet* 108: 309–323

Liu W, He X, Yang S, Zouari R, Wang J, Wu H, Kherraf Z-E, Liu C, Coutton C, Zhao R, *et al* (2019b) Bi-allelic Mutations in TTC21A Induce Asthenoteratospermia in Humans and Mice. *Am J Hum Genet* 104: 738–748

Livak KJ & Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods San Diego Calif* 25: 402–408

Lorès P, Dacheux D, Kherraf Z-E, Nsota Mbango J-F, Coutton C, Stouvenel L, Ialy-Radio C, Amiri-Yekta A, Whitfield M, Schmitt A, *et al* (2019a) Mutations in TTC29, Encoding an Evolutionarily Conserved Axonemal Protein, Result in Asthenozoospermia and Male Infertility. *Am J Hum Genet* 105: 1148–1167

Lorès P, Dacheux D, Kherraf Z-E, Nsota Mbango J-F, Coutton C, Stouvenel L, Ialy-Radio C, Amiri-Yekta A, Whitfield M, Schmitt A, *et al* (2019b) Mutations in TTC29, Encoding an Evolutionarily Conserved Axonemal Protein, Result in Asthenozoospermia and Male Infertility. *Am J Hum Genet* 105: 1148–1167

Lu Y, Oura S, Matsumura T, Oji A, Sakurai N, Fujihara Y, Shimada K, Miyata H, Tobita T, Noda T, *et al* (2019) CRISPR/Cas9-mediated genome editing reveals 30 testis-enriched genes dispensable for male fertility in mice†. *Biol Reprod* 101: 501–511

Luck K, Kim D-K, Lambourne L, Spirohn K, Begg BE, Bian W, Brignall R, Cafarelli T, Campos-Laborie FJ, Charloteaux B, *et al* (2020) A reference map of the human binary protein interactome. *Nature* 580: 402–408

McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GRS, Thormann A, Flicek P & Cunningham F (2016) The Ensembl Variant Effect Predictor. *Genome Biol* 17: 122

Meienberg J, Bruggmann R, Oexle K & Matyas G (2016) Clinical sequencing: is WGS the better WES? *Hum Genet* 135: 359–362

Nsota Mbango J-F, Coutton C, Arnoult C, Ray PF & Touré A (2019) Genetic causes of male infertility: snapshot on morphological abnormalities of the sperm flagellum. *Basic Clin Androl* 29: 2

O'Donnell L (2014) Mechanisms of spermiogenesis and spermiation and how they are disturbed. *Spermatogenesis* 4: e979623

Oud MS, Smits RM, Smith HE, Mastrorosa FK, Holt GS, Houston BJ, de Vries PF, Alobaidi BKS, Batty LE, Ismail H, *et al* (2022) A de novo paradigm for male infertility. *Nat Commun* 13: 154

Plagnol V, Curtis J, Epstein M, Mok KY, Stebbings E, Grigoriadou S, Wood NW, Hambleton S, Burns SO, Thrasher AJ, *et al* (2012) A robust model for read count data in exome sequencing

experiments and implications for copy number variant calling. *Bioinforma Oxf Engl* 28: 2747–2754

Ralston KS, Kabututu ZP, Melehani JH, Oberholzer M & Hill KL (2009) The Trypanosoma brucei flagellum: moving parasites in new directions. *Annu Rev Microbiol* 63: 335–62

Ralston KS, Kisalu NK & Hill KL (2011) Structure-function analysis of dynein light chain 1 identifies viable motility mutants in bloodstream-form Trypanosoma brucei. *Eukaryot Cell* 10: 884–94

Rolland M, Le Moal J, Wagner V, Royère D & De Mouzon J (2013) Decline in semen concentration and morphology in a sample of 26,609 men close to general population between 1989 and 2005 in France. *Hum Reprod Oxf Engl* 28: 462–470

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, *et al* (2012) Fiji - an Open Source platform for biological image analysis. *Nat Methods* 9: 10.1038/nmeth.2019

Schneider CA, Rasband WS & Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* 9: 671–675

Schumann Burkard G, Jutzi P & Roditi I (2011) Genome-wide RNAi screens in bloodstream form trypanosomes identify drug transporters. *Mol Biochem Parasitol* 175: 91–94

Shamoto N, Narita K, Kubo T, Oda T & Takeda S (2018) CFAP70 Is a Novel Axoneme-Binding Protein That Localizes at the Base of the Outer Dynein Arm and Regulates Ciliary Motility. *Cells* 7

Sunter JD (2016) A vanillic acid inducible expression system for Trypanosoma brucei. *Mol Biochem Parasitol* 207: 45–48

Touré A, Martinez G, Kherraf Z-E, Cazin C, Beurois J, Arnoult C, Ray PF & Coutton C (2020) The genetic architecture of morphological abnormalities of the sperm tail. *Hum Genet*

Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, Mann M & Cox J (2016) The Perseus computational platform for comprehensive analysis of (prote)omics data. *Nat Methods* 13: 731–740

Vincensini L, Blisnick T & Bastin P (2011) 1001 model organisms to study cilia and flagella. *Biol Cell* 103: 109–130

Wang G, Guo Y, Zhou T, Shi X, Yu J, Yang Y, Wu Y, Wang J, Liu M, Chen X, *et al* (2013) In-depth proteomic analysis of the human sperm reveals complex protein compositions. *J Proteomics* 79: 114–122

Wang W-L, Tu C-F & Tan Y-Q (2020) Insight on multiple morphological abnormalities of sperm flagella in male infertility: what is new? *Asian J Androl* 22: 236–245

Wang Y, Yang J, Jia Y, Xiong C, Meng T, Guan H, Xia W, Ding M & Yuchi M (2014) Variability in the morphologic assessment of human sperm: use of the strict criteria recommended by the World Health Organization in 2010. *Fertil Steril*

Wickstead B & Gull K (2007) Dyneins Across Eukaryotes: A Comparative Genomic Analysis. *Traffic* 8: 1708–1721

Wirtz E, Leal S, Ochatt C & Cross GeorgeAM (1999) A tightly regulated inducible expression system for conditional gene knock-outs and dominant-negative genetics in Trypanosoma brucei. *Mol Biochem Parasitol* 99: 89–101

Woods A, Sherwin T, Sasse R, MacRae TH, Baines AJ & Gull K (1989) Definition of individual components within the cytoskeleton of Trypanosoma brucei by a library of monoclonal antibodies. *J Cell Sci* 93 ( Pt 3): 491–500

Yamamoto R, Hwang J, Ishikawa T, Kon T & Sale WS (2021) Composition and function of ciliary inner-dynein-arm subunits studied in Chlamydomonas reinhardtii. *Cytoskelet Hoboken NJ* 78: 77–96

Yamamoto R, Yagi T & Kamiya R (2006) Functional binding of inner-arm dyneins with demembranated flagella of Chlamydomonas mutants. *Cell Motil Cytoskeleton* 63: 258–265

Yamamoto R, Yanagisawa H, Yagi T & Kamiya R (2008) Novel 44-Kilodalton Subunit of Axonemal Dynein Conserved from Chlamydomonas to Mammals. *Eukaryot Cell* 7: 154–161

**Acknowledgments**: We thank all individuals for their participation. We thank Samuel Dean (Warwick Medical School) and Jack Sunter (Oxford Brookes University) for the pPOT plasmids and SmOx T. brucei cell lines, F. Bringaud (University of Bordeaux) for the anti-enolase antibody, P. Bastin (Institut Pasteur) for the anti-Ty1 antibody. We thank G. Cougnet-Houlery, S. Guit, and J. Marcos for ongoing access to the MFP lab infrastructure. This work was supported by the Institut National de la Santé et de la Recherche Médicale (INSERM), the Centre National de la Recherche Scientifique (CNRS), University Grenoble Alpes, and French National Research Agency funding for specific projects: MASFLAGELLA (ANR-14-CE15-0002), FLAGEL-OME (ANR-19-CE17-0014), OLIGO-SPERM (ANR-21-CE17-0007). Mass spectrometry experiments were performed at the Centre de Génomique Fonctionnelle facility (CGFB, Bordeaux) for Trypanosoma studies and at the Proteom'IC core-facility (Institut Cochin, Paris - François Guillonneau and Johanna Bruce) for mouse studies. The Orbitrap Fusion mass spectrometer at the Proteom'IC core-facility was acquired with funds from the FEDER through the « Operational Program for Competitiveness Factors and employment 2007-2013 » and from the « Cancéropôle Ile-de-France ». The Bordeaux Imaging Center is a service unit of the CNRS INSERM and Bordeaux University, a member of the French BioImaging national infrastructure funded through the French National Research Agency [ANR-10-INBS-04]. **Author contributions:** D.D., M.Bo., G.M., P.F.R., Z-E.K., C.Ca., C.L., E.L., M.Bi., C.A., A.T. and C.Co. analyzed the data and wrote the manuscript; C.Co., Z.-E.K., C.Ca., G.M., M.Bi., A.S., and N.T.-M. performed and analyzed the genetic data; D.D., C.E.B.R., D.R.R., M.T., J-W.D., and M.Bo. performed the Trypanosoma studies. J.B. and G.M. performed the immunofluorescence assays. P.L. and A.T. performed studies on the $Ttc29^{-/-}$ mouse model. A.A.-Y., A.D., S.-H.H., X.J., Y.S., C.L., R.H., L.H., and V.S. provided clinical samples and data; G.M., P.F.R., A.T., D.D., M.Bo., and C.Co. designed the study, supervised all molecular laboratory work, had full access to all study data, and took responsibility for data integrity and accuracy. All authors have read and agreed to the published version of the manuscript. **Competing interests:** The authors declare no competing interests. **Data and materials availability:** All data needed to evaluate the conclusions presented in the paper are provided in the paper and/or the Supplementary Materials.

# Supplementary Materials for

## Novel axonemal protein ZMYND12 is required for flagellum function and male fertility

Denis Dacheux *et al.*

*Corresponding author. Email: CCoutton@chu-grenoble.fr

**This PDF file includes:**

**Supplementary text**

<u>Whole-Exome Sequencing (WES) and bioinformatics analysis</u>

Genomic DNA was isolated from EDTA blood using DNeasy Blood & Tissue Kits (QIAGEN SA). Genetic data were obtained through several sequencing centers, in particular Novogene, Genoscope, and Integragen. Coding regions and intron/exon boundaries were sequenced after enrichment using Agilent SureSelect Human All Exon V5, V6, or Clinical Research capture kits.

An alignment-ready GRCh38 reference genome (including ALT, decoy, and HLA) was produced using "run-gen-ref hs38DH" from Heng Li's bwakit package (https://github.com/lh3/bwa). Exomes were analyzed using a bioinformatics pipeline developed in-house. The pipeline consists of two modules, both distributed under the GNU General Public License v3.0 and available on github (grexome-TIMC-Primary and grexome-TIMC-Secondary, URLs indicated in the web resources).

The first module takes FASTQ files as input and produces a single merged GVCF file per variant-caller, as follows: trim adaptors and filter low-quality reads with fastp 0.23.2, align reads with BWA-MEM 0.7.17, mark duplicates using samblaster 0.1.26, and sort and index BAM files with samtools 1.15.1. SNVs and short indels are called from each BAM file using strelka 2.9.10 (Kim *et al*, 2018) and GATK 4.1.8.1 to produce individual GVCF files from each variant-caller. These files are finally merged using mergeGVCFs.pl to obtain a single multi-sample GVCF per caller. Using several variant-callers helps to compensate for each caller's flaws.

The second module takes each merged GVCF as input and produces annotated analysis-ready TSV files. Annotation involved 17 streamlined tasks: Low-quality variant calls (DP < 10, GQ < 20, or less than 15% of reads supporting the ALT allele) were discarded. Variant Effect

Predictor v108 (McLaren *et al*, 2016) was used to annotate variants and predict their impact, in order to filter out low-impact (MODIFIER) variants and/or prioritize high-impact ones (e.g., stop-gain or frameshift variants). Variants with a minor allele frequency greater than 1% in gnomAD v.2.1 or 3% in 1000 Genomes Project phase 3 were excluded. The pipeline performs an integrative analysis of all available exomes, both at the individual variant level and at the gene level. Thus, recurring genomic variants are readily identified alongside genes that are severely impacted in several patients but not in control subjects. Additional information can be found in Arafah et al. (Arafah *et al*, 2021). Copy number variants (CNVs) were searched using the ExomeDepth software package, as previously reported (Kherraf *et al*, 2018a; Plagnol *et al*, 2012).

**Fig. S1. Depth of coverage at the *ZMYND12* locus for the ZMYND12_2 individual, carrier of a homozygous deletion of exons 6-8, and four other unrelated MMAF individuals.** Compared with the control individuals (blue), affected individual ZMYND12_2 (red) showed a complete absence of sequence coverage for exons 6, 7 and 8. ZMYND12 is on the reverse genomic strand. The five samples shown were processed by exome-seq in a single batch, using the same Agilent Clinical Research capture kit, Illumina HiSeq 4000 sequencing technology, and bioinformatics pipeline (see Supplemental Methods).

**Fig. S2. ZMYND12 immunostaining in human spermatozoa from MMAF individuals carriers of pathogenic variants in *ARMC2* and *WDR66* genes, or with unknwon genetic cause.** Sperm cells from a fertile control individual, two MMAF patients, carriers of mutations in *ARMC2* and *WDR66*, and one MMAF patient with unknown genetic cause were stained with anti-ZMYND12 (green) and anti-acetylated tubulin (red) antibodies. Sperm nuclear DNA was counterstained with DAPI (blue). ZMYND12 immunostaining was present throughout the flagellum in all samples, regardless of phenotype. Variants are specified for each patient. Scale bars: 10 μm.

**Fig. S3. Relative mRNA Expression of human *ZMYND12* transcripts.** RT-qPCR was performed with cDNAs from various human tissues purchased from Life Technologies®. A panel of 10 organs was used for experiments: placenta, lung, pancreas, testis, trachea, kidney, whole brain, skeletal muscle, liver and heart. Each sample was assayed in triplicate for each gene on a StepOnePlus (Life Technologies®) with Power SYBR®Green PCR Master Mix (Life Technologies®). The PCR cycle was as follows: 10 min at 95 °C, 1 cycle for enzyme activation; 15 s at 95 °C, 60 s at 60 °C with fluorescence acquisition, 40 cycles for the PCR. RT-qPCR data were normalized using the two reference housekeeping genes *RPL6* and *RPL27* for human samples, applying the -$\Delta\Delta C_t$ method(Livak & Schmittgen, 2001). The $2^{-\Delta\Delta Ct}$ value was set to 0 in brain cells, resulting in an arbitrary maximum expression of 1. The efficacy of primers was checked using a standard curve. Melting curve analysis was used to verify the presence of a single PCR product. Statistical significance of differences in expression of *ZMYND12* transcripts in several organs was determined by applying a two-tailed *t*-test using Prism 4.0 software (GraphPad, San Diego, CA). ***P< 0.001.

**Fig. S4. ZMYND12 is absent from sperm cells from *TTC29* individuals.** Sperm cells from a fertile control individual and a TTC29 individual carrying the previously-reported c.1761+1G>A splice-variant (Lorès *et al*, 2019b) were stained with anti-ZMYND12 (green) and anti-acetylated tubulin (red) antibodies. DAPI counterstaining was used to reveal sperm nuclear DNA (blue). In control sperm, ZMYND12 immunostaining decorated the full length of the flagellum, but in sperm from TTC29 individuals, a total absence of ZMYND12 was recorded. Scale bars: 10 µm.

**Fig. S5. Radial spokes are inconstantly affected in *ZMYND12* individuals.** Sperm cells from

a fertile control individual and from individual ZMYND12_3 stained with anti-RSPH1 (green)

– located in the radial spokes – and anti-acetylated tubulin (red) antibodies. RSPH1

immunostaining is uniformly present along the length of the flagellum in control sperm cells.

Sperm cells from individual ZMYND12_3 show heterogeneous staining regardless of flagellar

morphology, suggesting that the radial spokes are inconsistently present in sperm cells from

this individual. Scale bars: 10 µm. Lower panel: Histogram showing the percentage of analyzed spermatozoa of various morphologies displaying or lacking RSPH1 immunostaining (number of sperm cells analyzed = 200).

**Fig. S6. Immunostaining for AKAP4, DNAH2, DNAH8, DNAH17, DNAI1, and GAS8 is not affected by *ZMYND12* mutation.** Immunofluorescence (IF) experiments were performed using sperm cells from control individuals and from individual ZMYND12_3, carrier of the nonsense variant c.433C>T. Immunostaining patterns for AKAP4, DNAH2, DNAH8, DNAH17, DNAI1, and GAS8 in sperm from individual ZMYND12_3 were comparable to those observed in control sperm cells, demonstrating that the fibrous sheath, outer dynein arms (ODAs) and the N-DRC were not directly affected by *ZMYND12* variants. Scale bars: 10 µm.

**A**

```
ZYMND12  MNVIYPLAVPKGRRLCCEVCEAPAERVCAACTVTYYCGVVHQKADWDSIHEKICQLLIPL
TbTAX-1  ---------MQGANL----KVLPSTRAAAK---ELVDPLDITNVGWSTLEPKFDELMQLM
                   :*  .*  *:  *..*  :  :..*.::.  *:  :*:  :

ZYMND12  RTSMPFYNSEEERQHGLQQLQQRQKYLIEFCYTIAQKYLFEGKHEDAVPAALQSLRFRVK
TbTAX-1  DAPSSGINALAARSAHREKVGAVIEQLLTRAQDESRRLLVEGNGEAAAEAGVKTLRLKER
          :  *:  *.  :::  :  *:  .  :::  *.**:  *  *.  *.:::**::  :

ZYMND12  LYGLSSVELVPAYLLLAEASLGLGRIVQAEEYLFQAQWTVLKSTDCSN-ATHSLLHRNLG
TbTAX-1  FYGKGSVKLVPAHFHLARTNQFLKRYGNAEEILSLAHFIILQNPDEADATIKAELHQTFG
          :**  .**:****::  **.:.  *  *  :*** *  *::  :*:.  *  ::  :  ::  **:.:*

ZYMND12  LLYIAKKNYEEARYHLANDIYFASCAFGTEDIRTSGGYFHLANIFYDLKKLDLADTLYTK
TbTAX-1  LLYAADNKLDVSVKHLTCATYYLSVMNGPEHVLTTFAYFDLANVFATKACMEAAMALYDT
          ***  *.::  :  :  **:  *:  *  *  *.:  *:  .**.***:*  ::  *  :**  .

ZYMND12  VSEIWHAYLNNHYQVLSQAHIQQMDLLGKLFENDTGLDEA-----------QEAEAIRIL
TbTAX-1  VKNIWLKHLRRVLKDIVDETMAA--KLVKRYDDDEVTHEVGHASARAFGKENLADVSKML
          *.:**  :*..  :  :  :  :  *  *  :::*  .*.  :  *:.  ::*

ZYMND12  TSILNIRES---TSDKAPQKTIFVLKILVMFYYLMMNSSKAQEYGMRALSLAKEQQLDVH
TbTAX-1  FGIFSIQKERLTISHPTTARAQFLLG----LYLLWVNKNDEAA--EHLLSAR-------
          .*:.*::.  *.  :  ::  *:*  :*  *  :*...  :  **

ZYMND12  EQSTIQELLSLISTEDHPIT-------------------
TbTAX-1  --TT----SQKFYGERHPIVQDIEDWCIWFEIPFRGVAAEQ
          :*  .  :  *  ***.
```

**B**

**Fig. S7. TbTAX-1 is the *T. brucei* ortholog of ZMYND12.** (**A**) Clustal Omega alignment of human ZYMND12 and TbTAX-1. (**B**) Schematic representation of wild-type and truncated forms of ZYMND12 and TbTAX-1.

**Fig. S8. TbTAX-1 RNAi knock-down.** Micrograph of WT cells and TbTAX-1 RNAi-induced cells (48 h) in thin section. The characteristic 9+2 structure of the axoneme and the paraflagellar structure (PFR) are visible, and no obvious structural defects are observed in RNAi-induced cells.

**Fig. S9. TbTAX-1 knock-down does not affect TbWDR66, TbCFAP70, or TbSPAG6.** (**A**) TbWDR66, TbCFAP70, and TbSPAG6 are axoneme-associated proteins according to western blot analysis of whole cell lysate (WC), detergent-extracted cytoskeletons (CSK), salt-extracted flagella (Fg). Enolase was used as extraction control for cytoskeleton and flagella samples. (**B**) Co-immunoprecipitation assays using anti-Ty1 ($IP_{Ty1}$) in cell lines expressing TbTAX-1$_{HA}$ alongside TbWDR66$_{Ty1}$ or TbCFAP70 or TbSPAG6$_{Ty1}$. Proteins were detected simultaneously by western blotting using anti-Ty1 and anti-HA antibodies and their corresponding fluorescent secondary antibodies.

(**C**) Western blot analysis of the fate of TbWDR66$_{Ty1}$, TbCFAP70$_{Ty1}$, or TbPF16/SPAG6$_{Ty1}$ in TbTAX-1 RNAi cells; non-induced (NI) cells, and cells induced for 24, 48, or 72 h. Enolase served as loading control. (**D**) U-ExM co-immunolabelling of TbTAX-1$_{HA}$ (cyan) and TbWDR66$_{Ty1}$, TbCFAP70$_{Ty1}$, or TbSPAG6$_{Ty1}$ (magenta).

**Table S1.** Proteins co-immunoprecipitated with TbTAX-1 and TbTTC29 identified by mass spectrometry analyses.

| TbTAX-1$_{HA}$ immunoprecipitation | | |
|---|---|---|
| **Accession number** | **Protein** | **Fold-enrichment** |
| Tb927.9.10370 | TbTAX-1 (Broadhead *et al*, 2006b) | 20.6 |
| Tb927.3.1990 | TbTTC29 (Lorès *et al*, 2019a) | 20.3 |
| Tb927.11.8160 | Dynein Heavy Chain putative. DNAH1 ortholog putative (Dean *et al*, 2016) | 6.66 |
| Tb927.10.15350 | Histone H3 variant V | 6.25 |
| Tb927.10.14680 | Ribosome biogenesis, putative | 5.88 |
| Tb927.11.3880 | TrypARP, Actin putative(Ersfeld & Gull, 2001b) | 5.7 |
| Tb927.11.3830 | Hypothetical protein, conserved | 4.42 |
| Tb927.10.8940 | Flagellum targeting protein kharon1 | 4.24 |
| Tb927.11.970 | Hypothetical protein, conserved | 4.1 |
| Tb927.10.170 | Pseudouridine synthase, Cbf5p | 3.7 |
| Tb927.9.12610 | Glycerol kinase, glycosomal | 3.7 |
| Tb927.2.2950 | Nop14-like family, putative | 3.5 |
| Tb11.v5.0732.1 | Hexokinase, putative | 3.1 |
| Tb927.11.16720 | Mitotic cyclin 6, CYC6 | 2.4 |
| Tb927.11.8870 | DEAD-box ATP-dependent RNA helicase, MHEL61 | 2.3 |

| TbTTC29$_{TY1}$ immunoprecipitation | | |
|---|---|---|
| **Accession number** | **Protein** | **Fold-enrichment** |
| Tb927.3.1990 | TbTTC29 (Lorès *et al*, 2019a) | 27.3 |
| Tb927.11.8160 | Dynein Heavy Chain putative. DNAH1 ortholog putative (Dean *et al*, 2016) | 9.2 |
| Tb927.9.10370 | TbTAX-1 (Broadhead *et al*, 2006b) | 8.6 |
| Tb927.11.11330 | HSP70 | 6. 5 |
| Tb927.11.3880 | TrypARP, Actin putative(Ersfeld & Gull, 2001b) | 5.88 |
| Tb927.3.1790 | Pyruvate dehydrogenase E1 beta subunit, putative | 2.4 |

# Table S2. Proteins identification by tandem mass spectrometry (MS/MS) analysis of proteins co-immunoprecipitated with TTC29 from wild-type and Ttc29$^{-/-}$ L5 and L7 mouse testes.

Analysis of the 124 proteins specifically identified in wild-type testis

| | |
|---|---|
| In blue, proteins identified in all 4 experimental replicates : n=3 | |
| In yellow, proteins identified in 3 out of 4 experimental replicates : n= 4 | |
| In grey, proteins identified in 2 out of 4 experimental replicates : n= 24 | |
| In white , proteins identified in 1 out of 4 experimental replicates : n= 93 | |

| Selected proteins | Peptides number per genotype and distribution per replicate | | | | | MW (kDa) | Protein Description | Species |
|---|---|---|---|---|---|---|---|---|
| | WT | | | L5 KO | L7 KO | | | |
| DYH1_MOUSE | 153 | 4 | 55-47-30-21 | - | - | 487.1 | Dynein heavy chain 1, axonemal | Mus musculus OX=10090 |
| IDLC_MOUSE | 33 | 4 | 5-13-9-6 | - | - | 29.7 | Axonemal dynein light intermediate polypeptide 1 | Mus musculus OX=10090 |
| TTC29_MOUSE | 31 | 4 | 4-10-10-7 | - | - | 54.3 | Tetratricopeptide repeat protein 29 | Mus musculus OX=10090 |
| PRS8_MOUSE | 8 | 3 | 4-2-2-0 | - | - | 35.7 | Aldo-keto reductase family 1 member B1 | Mus musculus OX=10090 |
| BASI_MOUSE | 5 | 3 | 3-1-1-0 | - | - | 59.8 | ATP synthase subunit alpha, mitochondrial | Mus musculus OX=10090 |
| MPCP_MOUSE | 3 | 3 | 1-1-1-0 | - | - | 45.6 | 26S proteasome regulatory subunit 8 | Mus musculus OX=10090 |
| RL14_MOUSE | 3 | 3 | 1-1-1-0 | - | - | 44.9 | Phosphoglycerate kinase 2 | Mus musculus OX=10090 |
| ALDR_MOUSE | 8 | 2 | 6-2-0-0 | - | - | 50.2 | RuvB-like 1 | Mus musculus OX=10090 |
| ATPA_MOUSE | 8 | 2 | 6-0-2-0 | - | - | 114.9 | Sarcoplasmic/endoplasmic reticulum calcium ATPase 2 | Mus musculus OX=10090 |
| RUVB1_MOUSE | 7 | 2 | 6-1-0-0 | - | - | 26.7 | 40S ribosomal protein S3 | Mus musculus OX=10090 |
| RUVB2_MOUSE | 6 | 2 | 4-2-0-0 | - | - | 51.1 | RuvB-like 2 | Mus musculus OX=10090 |
| ABCF2_MOUSE | 4 | 2 | 1-3-0-0 | - | - | 58.0 | T-complex protein 1 subunit zeta | Mus musculus OX=10090 |
| ADT1_MOUSE | 4 | 2 | 3-1-0-0 | - | - | 42.4 | Basigin | Mus musculus OX=10090 |
| ARL3_MOUSE | 4 | 2 | 2-0-2-0 | - | - | 136.3 | Cullin-associated NEDD8-dissociated protein 1 | Mus musculus OX=10090 |
| DHB4_MOUSE | 4 | 2 | 3-0-1-0 | - | - | 111.2 | Hypoxia up-regulated protein 1 | Mus musculus OX=10090 |
| RAB6A_MOUSE | 4 | 2 | 3-0-1-0 | - | - | 57.1 | Protein disulfide-isomerase | Mus musculus OX=10090 |
| RB11A_MOUSE | 4 | 2 | 0-2-2-0 | - | - | 68.5 | Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit 1 | Mus musculus OX=10090 |
| SRSF2_MOUSE | 4 | 2 | 2-0-2-0 | - | - | 95.9 | Protein 4.1 | Mus musculus OX=10090 |
| CKAP4_MOUSE | 3 | 2 | 2-0-1-0 | - | - | 71.8 | ATP-binding cassette sub-family F member 2 | Mus musculus OX=10090 |
| HVM03_MOUSE | 3 | 2 | 0-1-2-0 | - | - | 32.9 | ADP/ATP translocase 1 | Mus musculus OX=10090 |
| RS14_MOUSE | 3 | 2 | 1-2-0-0 | - | - | 20.5 | ADP-ribosylation factor-like protein 3 | Mus musculus OX=10090 |
| RSPH1_MOUSE | 3 | 2 | 0-2-1-0 | - | - | 16.8 | Calmodulin-1 | Mus musculus OX=10090 |
| ACYP1_MOUSE | 2 | 2 | 1-1-0-0 | - | - | 79.5 | Peroxisomal multifunctional enzyme type 2 | Mus musculus OX=10090 |
| DOPD_MOUSE | 2 | 2 | 1-1-0-0 | - | - | 123.6 | Importin-5 | Mus musculus OX=10090 |
| HMGA1_MOUSE | 2 | 2 | 1-0-1-0 | - | - | 72.0 | Protein disulfide-isomerase A4 | Mus musculus OX=10090 |
| HNRPL_MOUSE | 2 | 2 | 0-1-1-0 | - | - | 24.9 | Peroxiredoxin-6 | Mus musculus OX=10090 |
| KAP2_MOUSE | 2 | 2 | 1-1-0-0 | - | - | 47.4 | 26S proteasome non-ATPase regulatory subunit 11 | Mus musculus OX=10090 |
| PPT1_MOUSE | 2 | 2 | 1-0-1-0 | - | - | 23.6 | Ras-related protein Rab-6A | Mus musculus OX=10090 |
| SMIM4_MOUSE | 2 | 2 | 1-0-0-1 | - | - | 24.4 | Ras-related protein Rab-11A | Mus musculus OX=10090 |
| STX12_MOUSE | 2 | 2 | 0-1-1-0 | - | - | 25.5 | Serine/arginine-rich splicing factor 2 | Mus musculus OX=10090 |
| U5S1_MOUSE | 2 | 2 | 2-1*-0-0 | - | - | 139.9 | Tripeptidyl-peptidase 2 | Mus musculus OX=10090 |
| PGK2_MOUSE | 7 | 1 | 7-0-0-0 | - | - | 69.6 | Xaa-Pro aminopeptidase 1 | Mus musculus OX=10090 |
| AT2A2_MOUSE | 6 | 1 | 6-0-0-0 | - | - | 36.1 | Aldose reductase-related protein 2 | Mus musculus OX=10090 |
| RS3_MOUSE | 6 | 1 | 6-0-0-0 | - | - | 42.3 | Calcium-binding and spermatid-specific protein 1 | Mus musculus OX=10090 |
| TCPZ_MOUSE | 6 | 1 | 6-0-0-0 | - | - | 63.7 | Cytoskeleton-associated protein 4 | Mus musculus OX=10090 |
| CAND1_MOUSE | 5 | 1 | 5-0-0-0 | - | - | 191.6 | Clathrin heavy chain 1 | Mus musculus OX=10090 |
| HYOU1_MOUSE | 5 | 1 | 5-0-0-0 | - | - | 69.7 | Dynein assembly factor 1, axonemal | Mus musculus OX=10090 |
| PDIA1_MOUSE | 5 | 1 | 5-0-0-0 | - | - | 49.2 | Dynein regulatory complex protein 9 | Mus musculus OX=10090 |
| RPN1_MOUSE | 5 | 1 | 5-0-0-0 | - | - | 26.8 | EF-hand domain-containing protein D2 | Mus musculus OX=10090 |
| 41_MOUSE | 4 | 1 | 4-0-0-0 | - | - | 36.9 | Fructose-1,6-bisphosphatase 1 | Mus musculus OX=10090 |

| | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| CALM1_MOUSE | 4 | 1 | 4-0-0-0 | - | | | - | | 272.4 | Fatty acid synthase | Mus musculus OX=10090 |
| IPO5_MOUSE | 4 | 1 | 4-0-0-0 | - | | | - | | 13.3 | Ig heavy chain V region 36-65 | Mus musculus OX=10090 |
| PDIA4_MOUSE | 4 | 1 | 4-0-0-0 | - | | | - | | 46.7 | Platelet-activating factor acetylhydrolase IB subunit alpha | Mus musculus OX=10090 |
| PRDX6_MOUSE | 4 | 1 | 4-0-0-0 | - | | | - | | 36.5 | Malate dehydrogenase, cytoplasmic | Mus musculus OX=10090 |
| PSD11_MOUSE | 4 | 1 | 4-0-0-0 | - | | | - | | 35.6 | Malate dehydrogenase, mitochondrial | Mus musculus OX=10090 |
| TPP2_MOUSE | 4 | 1 | 4-0-0-0 | - | | | - | | 39.6 | Phosphate carrier protein, mitochondrial | Mus musculus OX=10090 |
| XPP1_MOUSE | 4 | 1 | 4-0-0-0 | - | | | - | | 28.1 | Thioredoxin-dependent peroxide reductase, mitochondrial | Mus musculus OX=10090 |
| ALD2_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 44.2 | 26S proteasome regulatory subunit 10B | Mus musculus OX=10090 |
| CABS1_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 29.5 | Proteasome subunit alpha type-4 | Mus musculus OX=10090 |
| CLH1_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 29.1 | Proteasome subunit beta type-4 | Mus musculus OX=10090 |
| DAAF1_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 105.7 | 26S proteasome non-ATPase regulatory subunit 1 | Mus musculus OX=10090 |
| DRC9_MOUSE | 3 | 1 | 0-0-3-0 | - | | | - | | 60.7 | 26S proteasome non-ATPase regulatory subunit 3 | Mus musculus OX=10090 |
| EFHD2_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 22.2 | Ras-related protein Rab-1B | Mus musculus OX=10090 |
| F16P1_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 23.6 | 60S ribosomal protein L14 | Mus musculus OX=10090 |
| FAS_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 16.3 | 40S ribosomal protein S14 | Mus musculus OX=10090 |
| LIS1_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 34.2 | Radial spoke head 1 homolog | Mus musculus OX=10090 |
| MDHC_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 170.1 | Bifunctional glutamate/proline--tRNA ligase | Mus musculus OX=10090 |
| MDHM_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 75.7 | Arginine--tRNA ligase, cytoplasmic | Mus musculus OX=10090 |
| PRDX3_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 71.2 | Thioredoxin reductase 3 | Mus musculus OX=10090 |
| PRS10_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 85.5 | Aconitate hydratase, mitochondrial | Mus musculus OX=10090 |
| PSA4_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 11.2 | Acylphosphatase-1 | Mus musculus OX=10090 |
| PSB4_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 36.6 | Aldo-keto reductase family 1 member A1 | Mus musculus OX=10090 |
| PSMD1_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 93.8 | A-kinase anchor protein 4 | Mus musculus OX=10090 |
| PSMD3_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 54.0 | Aldehyde dehydrogenase family 3 member A2 | Mus musculus OX=10090 |
| RAB1B_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 42.4 | Protein arginine N-methyltransferase 1 | Mus musculus OX=10090 |
| SYEP_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 50.5 | Damage-control phosphatase ARMT1 | Mus musculus OX=10090 |
| SYRC_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 64.3 | Asparagine synthetase [glutamine-hydrolyzing] | Mus musculus OX=10090 |
| TRXR3_MOUSE | 3 | 1 | 3-0-0-0 | - | | | - | | 42.5 | Cytosolic acyl coenzyme A thioester hydrolase | Mus musculus OX=10090 |
| ACON_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 70.8 | Carnitine O-acetyltransferase | Mus musculus OX=10090 |
| AK1A1_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 78.2 | Caprin-1 | Mus musculus OX=10090 |
| AKAP4_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 38.2 | Sorbitol dehydrogenase | Mus musculus OX=10090 |
| AL3A2_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 40.6 | DnaJ homolog subfamily B member 11 | Mus musculus OX=10090 |
| ANM1_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 13.1 | D-dopachrome decarboxylase | Mus musculus OX=10090 |
| ARMT1_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 53.4 | Dynein regulatory complex protein 10 | Mus musculus OX=10090 |
| ASNS_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 49.0 | Spliceosome RNA helicase Ddx39b | Mus musculus OX=10090 |
| BACH_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 24.7 | Elongation factor 1-beta | Mus musculus OX=10090 |
| CACP_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 28.8 | Endoplasmic reticulum resident protein 29 | Mus musculus OX=10090 |
| CAPR1_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 32.5 | Glucosamine-6-phosphate isomerase 1 | Mus musculus OX=10090 |
| DHSO_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 31.1 | Glucosamine-6-phosphate isomerase 2 | Mus musculus OX=10090 |
| DJB11_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 23.6 | Glutathione S-transferase P 1 | Mus musculus OX=10090 |
| DRC10_MOUSE | 2 | 1 | 0-0-2-0 | - | | | - | | 54.0 | Solute carrier family 2, facilitated glucose transporter member 1 | Mus musculus OX=10090 |
| DX39B_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 11.6 | High mobility group protein HMG-I/HMG-Y | Mus musculus OX=10090 |
| EF1B_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 35.7 | Heme oxygenase 2 | Mus musculus OX=10090 |
| ERP29_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 64.0 | Heterogeneous nuclear ribonucleoprotein L | Mus musculus OX=10090 |
| GNPI1_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 16.8 | Eukaryotic translation initiation factor 5A-1 | Mus musculus OX=10090 |
| GNPI2_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 45.4 | cAMP-dependent protein kinase type II-alpha regulatory subunit | Mus musculus OX=10090 |
| GSTP1_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 50.8 | LanC-like protein 2 | Mus musculus OX=10090 |
| GTR1_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 34.8 | Leucine zipper transcription factor-like protein 1 | Mus musculus OX=10090 |
| HMOX2_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 34.1 | NADH-cytochrome b5 reductase 3 | Mus musculus OX=10090 |
| IF5A1_MOUSE | 2 | 1 | 2-0-0-0 | - | | | - | | 113.1 | Poly [ADP-ribose] polymerase 1 | Mus musculus OX=10090 |

| Protein | | | | | | | | | Mass | Description | Species |
|---|---|---|---|---|---|---|---|---|---|---|---|
| LANC2_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 67.8 | Protein disulfide-isomerase-like protein of the testis | Mus musculus OX=10090 |
| LZTL1_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 534.2 | Plectin | Mus musculus OX=10090 |
| NB5R3_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 34.5 | Palmitoyl-protein thioesterase 1 | Mus musculus OX=10090 |
| PARP1_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 108.5 | Pre-mRNA-processing factor 40 homolog A | Mus musculus OX=10090 |
| PDILT_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 49.5 | 26S proteasome regulatory subunit 6A | Mus musculus OX=10090 |
| PLEC_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 28.4 | Proteasome subunit alpha type-3 | Mus musculus OX=10090 |
| PR40A_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 34.6 | 26S proteasome non-ATPase regulatory subunit 14 | Mus musculus OX=10090 |
| PRS6A_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 23.0 | Ras-related protein Rab-18 | Mus musculus OX=10090 |
| PSA3_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 20.5 | Ras-related protein Rap-2b | Mus musculus OX=10090 |
| PSDE_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 24.9 | 60S ribosomal protein L10a | Mus musculus OX=10090 |
| RAB18_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 17.8 | 60S ribosomal protein L12 | Mus musculus OX=10090 |
| RAP2B_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 17.7 | 60S ribosomal protein L23a | Mus musculus OX=10090 |
| RL10A_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 69.1 | Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit 2 | Mus musculus OX=10090 |
| RL12_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 23.7 | Small nuclear ribonucleoprotein-associated protein B | Mus musculus OX=10090 |
| RL23A_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 32.8 | 40S ribosomal protein SA | Mus musculus OX=10090 |
| RPN2_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 56.5 | Succinyl-CoA:3-ketoacid coenzyme A transferase 2A, mitochondrial | Mus musculus OX=10090 |
| RSMB_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 31.8 | Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial | Mus musculus OX=10090 |
| RSSA_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 9.7 | Small integral membrane protein 4 | Mus musculus OX=10090 |
| SCO2A_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 24.6 | Superoxide dismutase [Mn], mitochondrial | Mus musculus OX=10090 |
| SDHB_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 100.7 | Sperm-associated antigen 1 | Mus musculus OX=10090 |
| SODM_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 62.6 | Stress-induced-phosphoprotein 1 | Mus musculus OX=10090 |
| SPAG1_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 38.4 | Serine-threonine kinase receptor-associated protein | Mus musculus OX=10090 |
| STIP1_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 31.2 | Syntaxin-12 | Mus musculus OX=10090 |
| STRAP_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 57.1 | Aspartate--tRNA ligase, cytoplasmic | Mus musculus OX=10090 |
| SYDC_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 140.2 | Valine--tRNA ligase | Mus musculus OX=10090 |
| SYVC_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 37.4 | Transaldolase | Mus musculus OX=10090 |
| TALDO_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 44.8 | Acetyl-CoA acetyltransferase, mitochondrial | Mus musculus OX=10090 |
| THIL_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 41.8 | 3-ketoacyl-CoA thiolase, mitochondrial | Mus musculus OX=10090 |
| THIM_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 67.6 | Transketolase | Mus musculus OX=10090 |
| TKT_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 24.9 | Transmembrane emp24 domain-containing protein 10 | Mus musculus OX=10090 |
| TMEDA_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 181.9 | DNA topoisomerase 2-beta | Mus musculus OX=10090 |
| TOP2B_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 109.4 | 116 kDa U5 small nuclear ribonucleoprotein component | Mus musculus OX=10090 |
| UBP7_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 128.5 | Ubiquitin carboxyl-terminal hydrolase 7 | Mus musculus OX=10090 |
| WDR92_MOUSE | 2 | 1 | 2-0-0-0 | - | | | | - | 39.8 | WD repeat-containing protein 92 | Mus musculus OX=10090 |

Tandem mass spectrometry (MS/MS) analysis was performed on TTC29-immunoprecipitated pellets obtained from wild-type and Ttc29$^{-/-}$ L5 and L7 mouse testes. N=4 for controls and N=4 for *TTC29$^{-/-}$* mutants. 124 proteins were specifically identified in wild-type and absent from samples from both Ttc29$^{-/-}$ L5 and L7 mutants. Proteins are ranked by their total peptides count. Only three proteins were systematically identified in the four experimental replicates: DNAH1, DNALI1 and TTC29. For each protein, the total number of peptides is indicated alongside the distribution of peptides in each experimental replicate.

**Table S3.** Primer sequences for Sanger sequencing verification of *ZMYND12* variants.

| Primers | Primer sequence (5'-3') | Tm (°C) | Product length (bp) |
|---|---|---|---|
| *ZMYND12*-F-Ex4 | AGGCTGCTGTGAACATTCCT | 57.3 | 250 |
| *ZMYND12*-R-Ex4 | AATGCTTGAACTCGGGTCGG | 59.4 | |

**Table S4**. *ZMYND12* MLPA probes used in this work.

| MLPA probes | Upstream hybridizing sequence (5'-3') | Downstream hybridizing sequence (5'-3') | Total probe length (nt) |
|---|---|---|---|
| *ZMYND12* specific probes | | | |
| exons 5 | GCTACTTCCACCTGGCTAATATATTCTATGACCTT | AAAAAGTTGGACCTGGCAGACACATTGTACACCAA | 112 |
| exon 6 | GGTCTCTGAGATCTGGCATGCATATTTGAAC | AATCACTATCAAGTCCTCTCACAGGCTCACATCCA | 108 |
| exon 8 | CTCAGTCTAGCCAAAGAACAACAGCTTGA | TGTCCATGAGCAAAGCACCATTCAAGAGT | 98 |

For each probe, the unique hybridizing sequence of the two half-probes is given. Total probe length represents the total length of amplification product (nt=nucleotides) = forward primers + upstream hybridizing sequence + downstream hybridizing sequence + reverse primers.

**Table S5. Primers used for RT-qPCR detection of *ZMYND12* in human tissue extracts.**

| Primer names | Primer sequences (5'-3') | Tm |
|---|---|---|
| ZMYND12_RT_F | TACTGCATCGGAATCTGGGAC | 59 °C |
| ZMYND12_RT_R | GCCAGATGATAACGGGCCTC | |
| RPL6_RTqPCR_F | TCCATTCGTCAGAGCAAACA | 56 °C |
| RPL6_RTqPCR_R | TACGGAGCAGCGCAAGAT | |
| RPL27_RTqPCR_F | AGAGTACCTTGTGGGCAT | 56 °C |
| RPL27_RTqPCR_R | TGATGGCACCTCAGATCGC | |

Tm: melting temperature

**Table S6.** Primary antibodies used in immunofluorescence experiments with human samples

| Primary antibodies | Reference | Species | Localization | Dilution |
|---|---|---|---|---|
| **Human experiments** | | | | |
| **AKAP4** | Sigma-Aldrich® HPA020046 | Rabbit | Fibrous sheath | 1/100 |
| **CFAP70** | Sigma-Aldrich® HPA062453 | Rabbit | Unknown (base of ODA?) | 1/50 |
| **DNAH1** | Sigma-Aldrich® HPA036806 | Rabbit | IDA | 1/25 |
| **DNAH17** | Sigma-Aldrich® HPA024354 | Rabbit | ODA | 1/200 |
| **DNAH8** | Sigma-Aldrich® HPA028447 | Rabbit | ODA | 1/200 |
| **DNAI1** | Sigma-Aldrich® HPA021649 | Rabbit | ODA | 1/200 |
| **DNALI1** | Sigma-Aldrich® HPA028305 | Rabbit | IDA | 1/100 |
| **GAS8** | Sigma-Aldrich® HPA041311 | Rabbit | N-DRC | 1/100 |
| **RSPH1** | Sigma-Aldrich® HPA017382 | Rabbit | RS | 1/100 |
| **SPAG6** | Sigma-Aldrich® HPA020046 | Rabbit | CPC | 1/500 |
| **TTC29** | Sigma-Aldrich® HPA061473 | Rabbit | Unknown (IDA? IFT?) | 1/100 |
| **WDR66** | Sigma-Aldrich® HPA040005 | Rabbit | CSC | 1/20 |
| **ZMYND12(PA5-69715)** | Sigma-Aldrich® HPA062453 | Rabbit | Unknown | 1/100 |
| **β-tubulin** | Cell signal 2128 | Rabbit | microtubules | 1/500 |
| **β-tubulin** | Sigma-Aldrich® T4026 | Mouse | microtubules | 1/500 |

Abbreviations: ODA, Outer Dynein Arms; IDA, Inner Dynein Arms; RS, Radial Spokes; FS, Fibrous sheath; CPC, Central Pair Complex; N-DRC: Nexin-Dynein Regulatory Complex, CSC: calmodulin-associated and spoke-associated complex

**Table S7.** Primary antibodies used in immunofluorescence experiments with *Trypanosoma* cells

| Antibody | Provider & Reference | Dilution IF | Dilution U-ExM | WB Dilution | IP Dilution |
|---|---|---|---|---|---|
| **Primary antibodies** | | | | | |
| Anti-PFR2 | (Coutton *et al*, 2018a) | 1:500 | - | - | - |
| Anti-TbSAXO | (Dacheux *et al*, 2012b) | 1:1000 | - | 1:1000 | |
| Anti-Myc | Mouse monoclonal 9E10 | 1:1000 | - | - | - |
| Anti-Myc | Sigma c-3956 | - | - | 1:1000 | - |
| Anti-Ty1 (purified supernatant) | (Brookman *et al*, 1995) | 1:2000 | 1:500 | 1:10,000 | 1:33 |
| Anti-HA | Genetex GTX628902 | - | - | - | 1:50 |
| Anti-Enolase | (Hannaert *et al*, 2003) | - | - | 1:25,000 | |
| Anti-Tubulin | (Woods *et al*, 1989) | - | - | 1:1000 | - |
| **Secondary antibodies** | | | | | |
| Anti-mouse FITC | Sigma F2012 | 1:100 | - | - | - |
| Anti-rabbit Alexa fluor 594 | Molecular Probes A21201 | 1:200 | - | - | - |
| Anti-rabbit Alexa fluor A647 | Molecular Probes A31573 | - | 1:500 | - | - |
| Anti-mouse Alexa fluor A488 | Molecular Probes A11001 | - | 1:500 | - | - |
| Anti-mouse HRP | Jackson 115-035-062 | - | - | 1:10,000 | - |
| Anti-rabbit HRP | Sigma A-9169 | - | - | 1:10,000 | - |
| Anti-rabbit StarBright Blue 700 | Bio-Rad 12004162 | - | - | 1:2500 | - |
| Anti-mouse StarBright Blue 520 | Bio-Rad 12005867 | - | - | 1:2500 | - |

**Movie S1.** Video microscopy of WT cells.

**Movie S2.** Video microscopy of RNAi$^{TbTAX-1}$ cells 7 days post-induction.