Effects of selfing on the evolution of sexual reproduction

Abstract

Plants exhibit diverse breeding systems, with populations capable of outcrossing, selfing, and/or

asexual reproduction. However, interactions between the three reproductive pathways remain not fully

clear. Sexual reproduction introduces segregation and recombination, but incurs several costs. Selfing can

affect the relative costs and benefits of sexual vs. asexual reproduction. Building population genetic

models, I explore how selfing affects the evolution of a sexual reproduction rate modifier via (1) indirect

selection due to segregation, (2) indirect selection from changes in recombination rates, and (3) selection

from the cost of meiosis and mate limitation. The dominant selective force mediating the evolution of sex

is found to vary with the rate of sexual reproduction and selfing, but selective force (1) and (3) are

generally stronger than selective force (2). A modifier enhancing sexual reproduction tends to be favored

by indirect selection generated by partially recessive, small-effect deleterious mutations, while hindered

by highly recessive lethal mutations. Overall, evolution towards higher sexual reproduction is hindered at

low sexual reproduction rates and intermediate selfing rates, but favored under high selfing rates. The

results suggest that asexual reproduction may precede the evolution of selfing, and offer insights into the

evolution of mechanisms reducing geitonogamy in partially clonal populations.

Introduction

Plants can reproduce through outcrossing, self-fertilization, and asexual reproduction. Many plant

populations reproduce through a combination of the three modes. As most plants are hermaphroditic,

selfing occurs in more than 50% of plant species (Jarne and Charlesworth 1993, Goodwillie et al. 2005).

Asexual reproduction is also prevalent in plants and occurs mainly in two ways: vegetative reproduction

and apomixis. Vegetative reproduction is very common in plants (Gustafsson, 1947, Abrahamson 1980,

Silvertown 1987), which can be realized through budding from organs such as roots, stems, leaves, and

inflorescences (Eckert 1999, Silvertown 2014, Barrett 2015). Apomixis refers to asexual seed production

(Asker and Jerling 1992, Richards 2003, Whitton et al. 2008, Hojsgaard and Hörandl 2019), but it is

much rarer compared to vegetative reproduction (Whitton et al. 2008). Also, unlike vegetative

reproduction, apomixis seems to be unlikely to coexist with sexual reproduction within the same

population (Bengtsson and Ceplitis 2000, Nakayama et al. 2002, Whitton et al. 2008).

To understand the diversity of breeding systems in plants, it is important to know how asexual

reproduction, outcrossing, and selfing interact to affect the evolution of each other. Empirical studies

show that apomictic populations often derived from self-incompatible lineages (Hörandl 2010, Hojsgaard

and Hörandl 2019), and many facultatively asexual reproducing species are outcrossing (Ellstrand and

Roose 1987). Nevertheless, partially clonal populations can still exhibit a wide range of selfing rates

(Honnay and Jacquemyn 2008, Vallejo-Marín et al. 2010), with some being highly self-fertilizing (e.g.,

Iris versicolor, Kron et al. 1993). Therefore, how self-fertilization and asexual reproduction interact with

each other is still not well understood. This is partly because studies on the evolution of selfing often

assume complete sexual reproduction, while studies on the evolution of sex usually consider random

mating.

The evolution of sexual reproduction has attracted long-lasting interest in the field of

evolutionary biology. Although sexual reproduction is prevalent, it bears several costs compared to

asexual reproduction (reviewed in Lehtonen et al. 2012). Table 1 summarizes several key selective forces

on sex and their causes. One of the major costs of sex is the "cost of males". In a sexually reproducing

population, females allocate resources to produce males in separate-sex populations (or the male function

in hermaphrodites), but male gametes contribute few resources to the offspring zygote. Also, mate search

and choice require extra resources and energy of females (Smith and Maynard-Smith 1978, Lehtonen et al.

2012, Meirmans et al. 2012). Such costs have been observed in various taxa (Michaels and Bazzaz 1986,

Innes et al. 2000, Kumpulainen et al. 2004, Wolinska and Lively 2008). However, the cost of males can

be compensated by benefits provided by males such as parental care, or increased access to resources for

offspring production (Lehtonen et al. 2012). Another cost of sex is the "cost of meiosis" (Williams 1975)

or the cost of "genome dilution" (Lewis 1987), which occurs because a sexually reproducing individual

contributes only half of its genomes to the offspring, while an asexual individual transmits the entire

genome. However, this cost is weaker if increased allocation to asexual reproduction reduces gamete

production. The cost of males can apply to selection both within and between populations, and may occur

in both hermaphroditic and separate-sex populations. In contrast, the cost of meiosis only applies to

selection within a population and only occurs in hermaphrodites, because in separate-sex populations,

asexually reproducing females do not bear the male function to fertilize sexually reproducing females.

Despite the costs reviewed above, sexual reproduction is common in eukaryotes (Bell 1982), so

numerous studies have tried to identify the selective advantages of sex. Sexual reproduction differs from

asexual reproduction by two genetic characteristics. First, sexual reproduction allows the possibility of

recombination. A sex modifier can thus be indirectly selected due to a change in the effective

recombination rate under interactions between selected loci, hereafter referred to as selection caused by

inter-locus interactions. Previous studies show that an allele increasing the recombination rate can be

indirectly selected for by breaking negative linkage disequilibria between loci, thereby enhancing the

efficiency of selection. This outcome may arise from epistatic interactions between loci (Charlesworth et

al. 1979, Barton 1995, Roze and Lenormand 2005, Roze and Stetsenko, Roze 2022) or the interplay

between selection and drift in finite population size, known as the Hill-Robertson effect (Hill and

Robertson 1966, Otto and Barton 2001, Keightley and Otto 2006, Roze and Barton 2006, Roze 2021).

Secondly, in a diploid population, sexual reproduction entails segregation, and consequently, it

can be favored under interactions between alleles within a locus, referred to as selection caused by intra-

locus interactions. Specifically, consider a fitness locus with allele A and a, where a is deleterious. When

allele a is recessive enough so that the heterozygote Aa is on average fitter than homozygotes (i.e., $w_{Aa}^2 >$

$w_{AA}w_{aa}$), or when the population size is finite, asexual reproduction will generate an excess of

heterozygotes than Hardy-Weinberg equilibrium in a random mating population (Chasnov 2000, Agrawal

and Chasnov 2001, Roze and Michod 2010). This intra-locus association reduces genetic variance and

hinders selection against the deleterious allele a. A modifier allele that increases the rate of sexual

reproduction can gain a long-term advantage by breaking this association through segregation. This is

because it augments genetic variance and thus enhancing mean fitness at equilibrium, as selection

becomes more efficient in eliminating the deleterious allele (Otto 2003). However, the modifier will face

a short-term disadvantage since it immediately produces more homozygous offspring, which has a lower

average fitness than heterozygotes.

Regarding the multiple selective forces on the evolution of sex reviewed above, selfing can have

various effects on the relative costs and advantages of sexual vs. asexual reproduction, as summarized in

Table 1. First, selfing is a mode of uniparental reproduction, thus reducing the cost of meiosis during

sexual reproduction (Charlesworth 1980). Additionally, selfing serves as an alternative method to asexual

reproduction for providing reproductive assurance under mate limitation, where outcrossing is disfavored

(Busch and Delph 2012, Eckert et al. 2016).

Also, selfing influences indirect selection on sex through changes in the recombination rate

caused by inter-locus interactions. Selfing generates positive associations among homozygotes at different

loci, referred to as identity disequilibrium (Vitalis and Couvet 2001). This effect can lead to selection

against a modifier increasing the recombination rate in the absence of epistasis. However, it can generate

indirect selection for a higher recombination rate under negative epistatic interactions between deleterious

mutations (Charlesworth et al. 1979, Roze and Lenormand 2005, Stetsenko and Roze 2022). Indirect

selection for a higher recombination rate can also be caused by the Hill-Robertson effect, which tends to

be stronger under selfing for two reasons (Roze 2021, Stetsenko and Roze 2022). First, selfing reduces

the effective recombination rate across the genome due to increased homozygosity. Second, selfing

reduces the effective population size by increasing the variance in allele frequency changes (Caballero

and Hill 1992) and strengthening background selection (selection against linked deleterious mutations;

Stephan 2010, Charlesworth 2013). Nevertheless, previous studies on the evolution of the recombination

rate often assume complete sexual reproduction, and the effects of selfing under the presence of asexual

reproduction are not fully clear.

Furthermore, selfing will affect indirect selection on sex through segregation caused by intra-

locus interactions. One reason is that selfing can alter genetic associations between the sex modifier locus

and fitness locus (Otto 2003). Another factor is that selfing may change the relative fitness of sexually

versus asexually reproduced offspring due to increased homozygosity (Muirhead and Lande 1997),

thereby affecting the short-term (dis)advantage of a sex modifier. This effect is akin to inbreeding

depression, where selfed offspring tends to have lower fitness than outcrossed offspring because selfing

exposes partially recessive deleterious mutations to homozygotes (Charlesworth and Willis 2009). A

higher selfing rate can lower inbreeding depression by purging deleterious mutations (Roze 2015).

However, it is not clear whether selfing will decrease or increase the relative fitness of sexually

reproduced compared to asexually reproduced offspring.

As for the overall effects of selfing on indirect selection caused by intra-locus interactions, Otto

(2003) showed that gametophytic selfing (a mode of selfing in some mosses and ferns that involves the

union of two gametes from the same haploid gametophyte; Soltis and Soltis 1992) can qualitatively alter

the genetic conditions for a sex modifier to invade compared to those under outcrossing. However, in

most plants and some animal groups, self-fertilization occurs through the union of two gametes from the

same diploid sporophyte (referred to as sporophytic selfing, Jarne and Charlesworth 1993). Since gametes

from the same haploid parent are identical, gametophytic selfing always produces homozygotes. In

contrast, sporophytic selfing can still generate heterozygous offspring if the parent is heterozygous.

Therefore, the effects of sporophytic selfing on indirect selection on sex through segregation are not fully

clear. Uyenoyama and Bengtsson (1989) showed that sporophytic selfing can promote the evolution of

sex, assuming the deleterious mutation is completely lethal. However, since the lethal allele only exists in

the heterozygous state in their recursion equations, whether the results obtained in this special case hold

for more general genetic conditions remains to be investigated.

Due to the complicated effects of selfing on the evolution of sex, we still lack have a

comprehensive understanding of the relative importance of these forces, as well as the overall effects of

selfing on the evolution of sex. Answering these questions requires quantifying the strength of each

selective force. Nevertheless, most previous studies that compare the strength of different selective forces

assume random mating (e.g., Roze and Michod 2010, Roze 2014). Therefore, in this paper, I use

population genetic models and individual-based simulations to investigate the effects of selfing on the

evolution of a sex modifier through: (1) indirect selection through segregation caused by intra-locus

interactions (2) indirect selection caused by inter-locus interactions due to changes in the effective

recombination rate, and (3) selection due to the cost of meiosis and mate limitation.

I find that for indirect selection caused by intra-locus interactions, under sporophytic selfing, a

modifier increasing the rate of sexual reproduction is favored either when the deleterious allele is

recessive and selection is weak, or when the deleterious mutation is dominant enough and selection is

strong. This is qualitatively similar to results found under gametophytic selfing (Otto 2003). Moreover,

since deleterious are often recessive, a higher selfing rate lowers the fitness of sexually reproduced

offspring relative to asexually reproduced offspring, generating a short-term disadvantage to a modifier

increasing the rate of sex due to segregation. Using empirically estimated genetic parameters of

deleterious mutations, I show that indirect selection caused by intra-locus interactions has comparable

strength to selection due to the cost of meiosis and mate limitation. Both selective forces are stronger than

indirect selection caused by inter-locus interactions, unless the selfing rate is high. Overall, the evolution

towards a higher sexual reproduction rate tends to be disfavored under an intermediate selfing rate and a

low sexual reproduction rate, but is favored when the selfing rate is high. Given that clonality is often

associated with outcrossing in plant populations, the current results suggest that asexual reproduction may

evolve prior to the evolution of selfing in plant populations.

Methods

This section consists of three subsections. The first and second subsections present a population

genetic model for selection on a sex modifier caused by intra-locus interactions and inter-locus

interactions, respectively. The third subsection describes the algorithm of individual-based simulations.

The biological meanings of key symbols used in the model are summarized in Table 2. The Mathematica

Notebook for all population genetic models and related derivations, as well as the C++ code for

individual-based simulation, will be available on Zenodo [uploaded upon acceptance].

Indirect selection through segregation caused by intra-locus interactions

I consider a diploid population with an infinitely large population size. Each generation begins

with viability selection, followed by mutations and reproduction. During reproduction, there may be

selection due to the cost of meiosis and mate limitation. I consider a sex modifier locus **M** controlling the

proportion of sexually reproduced offspring, and a fitness locus **A** governing individual viability, with a

recombination rate r between them.

The sex modifier locus **M** has two alleles M and m. The rate of sexual reproduction of genotype

MM, Mm and mm is σ_0, $\sigma_1 = \sigma_0 + h_\sigma \Delta\sigma$, $\sigma_2 = \sigma_0 + \Delta\sigma$, respectively. The model focuses on the

invasion of allele m in a population previously fixed with allele M, so I refer to σ_0 as the background

sexual reproduction rate. During sexual reproduction, the sporophytic selfing rate is denoted by α. The

fitness locus **A** has two alleles A and a, where allele a is deterious. The fitness of AA, Aa and aa are 1,

$1 - hs$, and $1 - s$, respectively, where s and h are the selection coefficient and dominance of allele a.

Allele A mutates to a with a rate μ, and I assume there is no backward mutation from a to A. I denote the

genotype frequency at the beginning of each generation by x_{ij}, where $i \leq j$ and $i, j = 1,2,3,4$ (1 for MA,

2 for Ma, 3 for mA, and 4 for ma).

Denote the genotype frequency after viability selection as $x'_{ij} = x_{ij}/\overline{W}$, where $\overline{W} =$

$\sum_{i \leq j \in \{1,2,3,4\}} x_{ij} w_{ij}$ is the mean fitness, and w_{ij} is the fitness of genotype x_{ij}. The genotype frequencies

after mutation and selection, denoted by x''_{ij}, are given by

$$x''_{11} = (1 - \mu^2)x'_{11}, \tag{1a}$$

$$x''_{12} = 2\mu(1 - \mu)x'_{11} + (1 - \mu)x'_{12}, \tag{1b}$$

174 etc., and the frequency of other genotypes can be expressed in a similar way.

175 After selection and mutation, reproduction occurs. I denote the genotype frequency within

176 asexually reproduced, selfed and outcrossed offspring by $x_{ij}^{(asex)}$, $x_{ij}^{(self)}$ and $x_{ij}^{(out)}$, respectively. The

177 genotype frequency within asexually reproduced offspring is $x_{ij}^{(asex)} = x''_{ij}$. Under sporophytic selfing,

178 the genotype frequency within selfed offspring can be calculated as

$$\overline{\sigma} x_{11}^{(self)} = \sigma_0 \left(x''_{11} + \frac{x''_{12}}{4} \right) + \sigma_1 \left(\frac{x''_{13}}{4} + \frac{x''_{14}(1 - r)^2}{4} + \frac{x''_{23}r^2}{4} \right), \tag{2a}$$

$$\bar{\sigma} x_{12}^{(self)} = \sigma_0 \frac{x_{12}''}{2} + \sigma_1 \left(\frac{x_{14}'' r(1-r)}{2} + \frac{x_{23}'' r(1-r)}{2} \right), \tag{2b}$$

181   etc. Here, $\bar{\sigma}$ is the average sexual reproduction rate, given by

182    $$\bar{\sigma} = \sigma_0(x_{11}'' + x_{12}'' + x_{22}'') + \sigma_1(x_{13}'' + x_{14}'' + x_{23}'' + x_{24}'') + \sigma_2(x_{33}'' + x_{34}'' + x_{44}''). \tag{3}$$

183   The frequency of other genotypes $x_{ij}^{(self)}$ can be obtained in a similar way.

184   For outcrossed offspring, I denote the haplotype frequency within female gametes (i.e., ovules)

and male gametes (i.e., pollen) by y_i and z_i, respectively, which are calculated as

$$y_1 = \frac{1}{\bar{\sigma}} \left[\sigma_0 \left(x_{11}'' + \frac{x_{12}''}{2} \right) + \sigma_1 \left(\frac{x_{13}''}{2} + \frac{x_{14}''(1-r)}{2} + \frac{x_{23}'' r}{2} \right) \right], \tag{4a}$$

$$z_1 = \frac{1}{1 - c(1-\bar{\sigma})} \left[\left(1 - c(1-\sigma_0)\right) \left(x_{11}'' + \frac{x_{12}''}{2} \right) + \left(1 - c(1-\sigma_1)\right) \left(\frac{x_{13}''}{2} + \frac{x_{14}''(1-r)}{2} + \frac{x_{23}'' r}{2} \right) \right], \tag{4b}$$

respectively. The frequency of other haplotypes can be calculated in a similar way. In equation (4a), I

assume an increased rate of asexual reproduction causes a proportional reduction in female gamete

production. In equation (4b), the parameter c measures the reduction in male gamete production due to an

increased rate of asexual reproduction. When $c = 1$, an increased rate of asexual reproduction causes a

proportional reduction in male gamete production (thus $y_i = z_i$), so the sex modifier is neutral when there

is no fitness locus. When $c < 1$, sex suffers the cost of meiosis since asexual reproduction can transmit

extra male gametes. I assume pollination is random, so the genotype frequency within outcrossed

offspring is

$$x_{ij}^{(out)} = \begin{cases} y_i z_j, \text{for } i = j \\ y_i z_j + y_j z_i, \text{for } i < j \end{cases}. \tag{5}$$

197   The reproductive success of outcrossing, selfing and asexual reproduction, as well as the fitness

198   of offspring produced from the three reproductive modes may differ due to factors other than the focal

199   fitness locus (e.g., mate limitation, different seed provisioning). To account for these effects, I use w_{asex}

200   and w_{self} to represent the reproductive success and the offspring fitness of asexual reproduction and

201   selfing relative to outcrossing. The genotype frequency in the next generation is thus given by

202
$$\tilde{x}_{ij} = \frac{w_{asex}(1-\overline{\sigma})x_{ij}^{(asex)} + w_{self}\alpha\overline{\sigma}x_{ij}^{(self)} + (1-\alpha)\overline{\sigma}x_{ij}^{(out)}}{w_{asex}(1-\overline{\sigma}) + w_{self}\alpha\overline{\sigma} + (1-\alpha)\overline{\sigma}}. \tag{6}$$

To investigate whether a sex modifier allele m can invade into a population previously fixed with

allele M, I numerically calculate the leading eigenvalue λ of the Jacobian matrix given by equation (6) (a

complicated analytical expression of λ is available in Section 3.3 of the Mathematica Notebook). m can

invade when $\lambda > 1$. I also track the genetic associations within loci and between the modifier and fitness

locus after the introduction of a modifier allele m in the population, to see how genetic associations affect

the capacity for a sex modifier to invade (the calculation of genetic associations is available in Section 1.3

of the Mathematica Notebook).

To estimate the short-term (dis)advantage of a sex modifier due to fitness differences between

sexually and asexually reproduced offspring caused by the focal fitness locus, I calculate the fitness of

asexually reproduced offspring and selfed offspring relative to outcrossed offspring contributed by the

focal locus. To do this, I first solve the genotypic frequencies at the fitness locus under mutation-selection

balance based on equation (6). The mean fitness of asexually reproduced offspring is merely equal to the

mean fitness of the population $\overline{W}$. The mean fitness of selfed and outcrossed offspring can be calculated

as

$$x_{11}^{(k)} + x_{13}^{(k)} + x_{33}^{(k)} + (1-hs)\left(x_{12}^{(k)} + x_{14}^{(k)} + x_{23}^{(k)} + x_{34}^{(k)}\right) + (1-s)\left(x_{22}^{(k)} + x_{44}^{(k)}\right), \tag{7}$$

where $k = self$ or out, and $x_{ij}^{(self)}$ and $x_{ij}^{(out)}$ $(i,j \in \{1,2,3,4\})$ is given by equations (2) and (5),

respectively.

Indirect selection caused by inter-locus interactions

221   Indirect selection on sex through changes in the recombination rate caused by inter-locus

222   interactions can occur due to two causes: deterministic interactions between loci and the Hill-Robertson

223   effect in a finite population. I present models for estimating the strength of selection caused by deleterious

224   mutations in the genome under each cause, respectively.

<u>*Deterministic interactions between loci*</u>

To estimate the strength of indirect selection caused by deterministic interactions between

deleterious mutations in an infinitely large population, I modify the model described in Stetsenko and

Roze (2022). Briefly, I consider two chromosomes with map length R, and assume the loci where

deleterious mutations occur are uniformly distributed along the chromosome with a genomic mutation

rate U. A sex modifier locus i is located at the midpoint of the chromosome. I assume the recombination

rate between locus j and k is $r_{jk} = (1 - e^{-2D_{jk}})/2$ (Haldane 1919), where D_{jk} is the genetic distance

between locus j and k. For a sex modifier that changes the rate of sexual reproduction by $\Delta\sigma$, it

effectively changes the map length by $\Delta R = R\Delta\sigma$. The modifier thus effectively changes the

recombination rate between locus j and k by $\Delta r_{jk} = r_{jk}\Delta R/R$.

I denote the strength of selection on the modifier locus i contributed by fitness locus j and k as

$s_{jk}(D_{ij}, D_{jk}, D_{ik})$, where D_{ij}, D_{jk} and D_{ik} are genetic distance between pairwise loci. A complicated

expression of s_{jk} is available in Section 4.1 of the Mathematica Notebook. Briefly, s_{ij} is obtained using

equation (6) in Stetsenko and Roze (2022), by replacing the probability of identity by descent at one locus

F (inbreeding coefficient) and at two loci ϕ with

$$F = \frac{\alpha}{(2 - \alpha) + s\left(\alpha + \frac{2}{\sigma_0} - 2\right)} + o(\mu), \tag{8a}$$

$$\phi \approx \frac{\alpha\sigma_0\left[(1 - s)\left(2 - \alpha - 2(1 - r)r(3\alpha - 2)\right)\sigma_0 + 2(1 - 2(1 - r)r)s\right]}{[2s + (1 - s)(2 - \alpha)\sigma_0][(1 - s)^2(2 - \alpha + 2\alpha(1 - r)r)\sigma_0 + 2(2 - s)s]} + o(\mu). \tag{8b}$$

Equations (8a) and (8b) account for the coexistence of outcrossing, selfing and asexual reproduction in a

population, and the derivation is available in Section 1.3 of the Mathematica Notebook. F converges to

the classical expression $F = \alpha/(2 - \alpha)$ as $s \to 0$. Note that a high asexual reproduction rate (i.e., large

s/σ_0) can greatly reduce F and ϕ, and makes F less than 1 even under complete selfing $\alpha = 1$.

Ignoring interactions between deleterious alleles at more than two loci, the genome-wide strength

of selection caused by pairwise interactions of deleterious mutations is

$$248 \quad \left(\frac{U}{\mu R}\right)^2 \left[2\int_0^{\frac{R}{2}} \left(\int_0^{\frac{R}{2}-D_{ij}} s_{jk}(D_{ij}, D_{jk}, D_{ij}+D_{jk})dD_{jk} \right) dD_{ij} + \int_0^{\frac{R}{2}}\int_0^{\frac{R}{2}} s_{jk}(D_{ij}, D_{ij}+D_{ik}, D_{ik})dD_{ij}dD_{ik} \right]. (9)$$

In equation (9), the term $U/\mu R$ is the density of fitness loci on the chromosome. The first term in the

square bracket corresponds to the contribution from fitness loci located on the same side of the modifier

locus, while the second term represents the contribution from fitness loci located on the opposite side of

the modifier locus on the chromosome.

Hill-Robertson effect

To estimate the strength of selection due to the Hill-Robertson effect, I modify equation (1) in

Roze (2021). In a population with a background sexual reproduction rate σ_0 and inbreeding coefficient F,

the effective recombination rate is $R\sigma_0(1-F)$. A modifier that alters the rate of sex by $\Delta\sigma$ changes the

effective recombination rate by $R(1-F)\Delta\sigma$. Given that the effective recombination rate is large

compared to selection (i.e., $R\sigma_0(1-F) \gg (h+(1-h)F)s$), which will not hold when $\sigma_0 \to 0$ or $F \to$

1), the strength of selection caused by the Hill-Robertson effect is

$$260 \qquad\qquad \frac{1.8U^2}{4[\sigma_0 R(1-F)]^3 N_e} R(1-F)\Delta\sigma. \qquad\qquad (10)$$

261 The effective population size N_e is approximately (Glémin 2007)

$$262 \qquad\qquad \frac{N_e}{N} \approx \frac{1}{1+F} \times \exp\left[-\frac{U}{2(h+(1-h)F)s + R\sigma_0(1-F)} \right]. \qquad (11)$$

In equation (11), the first term captures the reduction of N_e due to increased variance in allele frequency

change under inbreeding (Caballero and Hill 1992). The second term accounts for an increased strength of

background selection due to a reduced effective recombination rate under selfing (Nordborg et al. 1996).

Individual-based simulations

I use individual-based simulations to investigate how selfing influences the overall selection on a

sex modifier caused by deleterious mutations in the genome. The simulation is similar to that described in

Roze (2015). Briefly, the simulation considers a population with N diploid individuals, each carrying two

chromosomes with the physical length scaled to be 1. I assume deleterious mutations occur at an infinite

number of loci (Kondrashov 1985), and the sex modifier locus is located at the midpoint of the

chromosome. At each generation, the number of new deleterious mutations per chromosome is drawn

from a Poisson distribution with parameter $U/2$, with their positions on the chromosome being drawn

from a uniform distribution $\sim \mathrm{U}(0,1)$. The fitness of an individual is $w = (1 - hs)^{n_{het}}(1 - s)^{n_{hom}}$,

where n_{het} and n_{hom} are the number of fitness loci in heterozygous and homozygous state that the

individual carries.

To generate the offspring population, for each offspring, I first sample a parent i from the

parental population, with the probability being proportional to individual fitness w_i. I then obtain its rate

of sexual reproduction σ_i, and generate a random number $\epsilon_1 \sim \mathrm{U}(0,1)$. If $\epsilon_1 > \sigma_i$, the offspring is

asexually reproduced, and the two chromosomes of the offspring are the same as its parent's. If $\epsilon_1 \leq \sigma_i$,

the offspring is sexually reproduced, and I draw a second number $\epsilon_2 \sim \mathrm{U}(0,1)$ to determine whether the

offspring is reproduced from outcrossing or selfing. If $\epsilon_2 < \alpha$, the offspring is selfed. If $\epsilon_2 > \alpha$, the

offspring is outcrossed, and a second parent j is sampled with the probability being proportional to

$w_j\left(1 - c(1 - \sigma_j)\right)$, where the term in the parenthesis captures the assumption that asexual reproduction

may reduce male gamete production. To obtain gametes generated by meiosis, the number of crossovers

between two chromosomes is drawn from a Poisson distribution with parameter L, and the position of

each crossover is drawn from $\mathrm{U}(0,1)$.

For each parameter combination, I heat the population for $6N$ generations to reach selection-

mutation-drift balance, with the sex modifier locus fixed with allele M. I then introduce a mutant allele m

at the sex modifier locus in a randomly chosen chromosome, and run the simulation until m becomes

fixed or lost. The fixation probability of m is calculated based on 2×10^5 replications.

Results

The result section contains five subsections. The first subsection investigates the effects of selfing

on indirect selection caused by intra-locus interactions, assuming that there is no cost of meiosis (set $c =$

1). The second and third subsections present the strength of indirect selection on a sex modifier generated

by genome-wide deleterious mutations due to intra-locus interactions and inter-locus interactions,

respectively. The fourth subsection presents results from individual-based simulations on how selfing

affects the overall selection on a sex modifier caused by deleterious mutations in the genome. The last

subsection investigates the effects of selfing on selection due to the cost of meiosis and mate limitation. In

this paper, I mainly present results under sporophytic selfing, since results under gametophytic selfing are

qualitatively similar. Also, I only present results for a modifier that increases the rate of sex, since when a

modifier that increases the rate of sex is favored, a modifier that reduces the rate sex is selected against,

and *vice versa*.

Indirect selection through segregation caused by intra-locus interactions

For indirect selection through segregation, I consider two cases. In the first case, the deleterious

allele a at the fitness locus is at mutation-selection balance, and I investigate whether a sex modifier allele

m can invade into a population in which the modifier locus is fixed with allele M. In the second case, the

fitter allele A sweeps from a low initial frequency $p_{A,0}$ to a high frequency $p_{A,T}$, and I investigate changes

in the frequency of an initially rare sex modifier allele m.

Mutation-selection balance

Under complete outcrossing or random mating ($\alpha = 0$), a modifier that increases the rate of sex

invades when the dominance coefficient of the deleterious mutation h is intermediate (Fig. 1(a)).

Specifically, when h is low ($h < \frac{1}{1+\sqrt{1-s}}$), selection causes an excess of Aa when there is asexual

reproduction (Otto 2003). A modifier that increases the rate of sex enjoys a long-term benefit by breaking

this negative association. However, the modifier initially suffers a short-term cost, since the offspring

contain more homozygotes, and thus is on average less fit than asexually reproduced offspring. As a result,

invasion of the modifier requires that the dominance h is not too high, so that the long-term benefit exists,

but also not too low so that the short-term cost is not too strong.

When selfing is present, given that the selfing rate is much larger than the mutation rate ($\alpha \gg \mu$),

inbreeding generates an excess of homozygotes at a locus (see equation (8a)). Consequently, the invasion

conditions of a modifier increasing the rate of sexual reproduction are altered compared to the case of

outcrossing. There exists a narrow range of selection coefficient s where the modifier cannot invade for

all levels of dominance h. This range can be represented by a critical value θ under the first-order

approximation of the mutation rate μ (the expression of θ is in Section 3.3 of the Mathematica Notebook).

A modifier that increases the rate of sex can invade either when $s < \theta$ and dominance h is low enough, or

$s > \theta$ and h is high enough (bottom-left and upper-right regions in Fig. 1(b)).

The invasion region is mainly determined by the value of the critical selection coefficient θ. As θ

increases, the bottom-left region expands, while the top-right region shrinks. Generally, θ becomes

smaller when the selfing rate α is higher (Fig. 1(b)), the background sexual reproduction rate σ_0 is lower

(Figs. 2(a), 2(b)), and when the sex modifier allele m is more dominant (compare Figs. 2(a)-(c)). These

results are qualitatively similar to those found under gametophytic selfing in Otto (2003). The only

difference is that under sporophytic selfing, the critical selection coefficient θ will slightly increase when

the recombination rate between the modifier and fitness locus r is lower (Fig. S2(a)). In contrast, under

gametophytic selfing, the value of θ is independent of r (see equation (13) in Otto (2003)).

We can check the short-term selective (dis)advantage of a sex modifier due to segregation by

computing the fitness of offspring from sexual and asexual reproduction. Section 1.4 of the Mathematica

Notebook Materials shows that the relative fitness reduction of asexually reproduced and selfed offspring

compared to outcrossed offspring is respectively,

339
$$d_a = 1 - \frac{w_{asex}}{w_{out}} = \frac{F(1-2h)}{h+(1-h)F}\mu + o(\mu), \tag{12a}$$

340
$$d_s = 1 - \frac{w_{self}}{w_{out}} = \frac{(1+F)(1-2h)}{2(h+(1-h)F)}\mu + o(\mu), \tag{12b}$$

where F is given by equation (8a), and d_s is just inbreeding depression. Equations (12a) and (12b) imply

that when the deleterious allele is partially recessive ($h < 0.5$), both asexually reproduced and selfed

offspring are less fit than outcrossed offspring. Since $\frac{d_s}{d_a} \approx \frac{1}{\alpha} + s\left(\frac{1-\sigma_0}{\alpha\sigma_0} + \frac{1}{2}\right) \geq 1$, selfed offspring are less

fit than asexually reproduced offspring. Moreover, Section I in Supplementary Materials shows that a

lower background sexual reproduction rate σ_0 or selfing rate α will increase the relative fitness of

asexually reproduced offspring d_a, but reduce the relative fitness of selfed offspring d_s (i.e., stronger

inbreeding depression).

Since sexually reproduced offspring contain both selfed and outcrossed offspring, their mean

fitness is $w_{sex} = \alpha(1 - d_s) + 1 - \alpha = 1 - \alpha d_s$. When the fitness locus is under selection-mutation

balance the fitness reduction of sexually versus asexually reproduced offspring is

351
$$\frac{1 - w_{sex}}{1 - w_{asex}} = \frac{d_s}{d_a}\alpha = \begin{cases} 1, & \alpha = 0 \\ 1 + s\left(\dfrac{\alpha}{2} + \dfrac{1}{\sigma_0} - 1\right) \geq 1, & \alpha > 0 \end{cases}. \tag{13}$$

Equation (13) shows that when the deleterious allele is recessive ($h < 0.5$), a higher selfing rate α and a

lower background sexual reproduction rate σ_0 lowers the relative fitness of sexually reproduced offspring,

thus increasing the short-term disadvantage of a modifier that increases the rate of sexual reproduction.

The above results qualitatively hold under gametophytic selfing. However, unlike sporophytic

selfing, the fitness of sexually reproduced offspring relative to asexually reproduced offspring is

independent of the gametophytic selfing rate (see section II in Supplementary Materials).

Selective Sweep

When allele A sweeps from a low to high frequency at the fitness locus, selfing generally inhibits

the evolution of sexual reproduction. Under outcrossing, similar to the results under mutation-selection

balance, a modifier increasing the sexual reproduction rate increases in frequency when the dominance h

is intermediate (Fig. 1(c)). When selfing is present, this region continuously shrinks as the selfing rate α

increases (region Fig. 1(d)). However, there appears a new parameter region that the modifier will

increase in frequency, which occurs when s is small and dominance h is low (bottom-left corner in Fig.

1(d)). This region expands when the selfing rate becomes higher (Fig. 1(d)) and the sex modifier m is

more recessive (compare Figs. 2(d)-(e)). Moreover, a higher recombination rate r slightly reduces the

parameter regions that favor a higher sexual reproduction rate (Fig. S2(b)).

Genetic association analysis

Selfing generates positive association between homozygotes at different loci (see equation (8a)).

Therefore, genotype mm will be associated with homozygotes at the fitness locus (AA and aa), while

Mm will be associated with heterozygote Aa. When h is low so that Aa is on average fitter than

homozygotes ($w_{Aa} > \sqrt{w_{AA}w_{aa}}$), the modifier m found in Mm individuals will enjoy a short-term benefit

by associating with Aa, but suffers long-term disadvantages due to a reduction in genetic variance. In

contrast, m found in mm individuals is associated more with homozygotes AA and aa, thereby suffering

a short-term cost but enjoying a long-term benefit. The invasion of m found in Mm individuals requires a

low enough h so that the short-term benefit is strong enough, while the invasion of m found in mm

individuals requires a high enough h.

Under mutation-selection balance, consistent with Otto (2003), when selection is weak ($s < \theta$),

the evolution of the modifier is mainly determined by those found in Mm individuals, so allele m is

associated more with Aa (i.e., $corr(m, Aa) > 0$). Therefore, invasion occurs when h is low enough.

When selection is strong ($s > \theta$), the evolution is mediated by m often found in mm individuals (thus

$corr(m, Aa) < 0$), so invasion occurs when h is high enough.

Under selective sweep, when selection is weak, the modifier m is associated with Aa, so a sex

modifier m increases in frequency when h is low enough. When selection is strong, whether the

correlation between the modifier m and the heterozygote Aa will be positive or negative depends on the

dominance h. When h is low, the modifier m is associated with homozygotes, so the modifier m is

favored when h is high enough. As h increases, the modifier m becomes associated with Aa, so the

modifier m is favored when h is low enough. As a result, the modifier will increase in frequency only

when h is intermediate, and this intermediate range may not exist for some values of selection coefficient.

Strength of indirect selection through segregation caused by intra-locus interactions

The previous section investigates selection contributed by a single fitness locus. This result can

be extended to estimate the strength of indirect selection on a modifier of sex through segregation

generated by deleterious mutations in the genome. When a modifier allele is rare and has small effects,

the selective strength $\Delta p_m / p_m (1 - p_m)$ can be approximated by the leading eigenvalue as $\lambda - 1$ (Otto

(2003)). For simplicity, I assume loci are identical and ignore genetic associations between loci and

epistasis. When the genomic mutation rate is U, the number of identical fitness loci is $L = U/\mu$, and the

genome-wide strength of selection is approximately $L(\lambda - 1)$. In addition, I estimate the strength of the

short-term (dis)advantage of a sex modifier due to fitness differences between sexually and asexually

reproduced offspring caused by deleterious mutations in the genome. When there are L loci, the fitness of

asexually reproduced and selfed offspring relative to outcrossed offspring is $w_{asex} = e^{-d_a L}$ and $w_{self} =$

$e^{-d_s L}$, where d_a and d_s are given by equation (12). The strength of this short-term selective force can

thus been estimated by setting $s = 0$, $w_{asex} = e^{-d_a L}$ and $w_{self} = e^{-d_s L}$ in equation (6), and solving for

the leading eigenvalue.

Previous analyses show that there are two major categories of deleterious mutations: small-effect,

partially dominant mutations and large-effect, highly recessive lethal mutations (Mukai et al. 1972, Eyre-

Walker and Keightley 2007, Charlesworth and Willis 2009). Based on previous estimations (Simmons

and Crow 1979), I compare the strength of indirect and direct selection caused by deleterious mutations

by setting $s = 0.05$, $h = 0.2$ for small-effect deleterious mutations, and $s = 0.8$, $h = 0.02$ for large-effect

lethal mutations.

For selection caused by small-effect deleterious mutations, a modifier that increases the rate of

sex is indirectly selected for either when the rate of sexual reproduction σ_0 is not low and the selfing rate

is not high (region I in Fig. 3(a)), or when σ_0 is low and α is high (region II in Fig. 3(a)). In region I,

sexually reproduced offspring is fitter than asexually reproduced offspring, so the modifier enjoys a weak

short-term advantage (upper-left corner in Fig. 3(b)). In contrast, in region I, fitness of sexually

reproduced offspring is much lower than asexually reproduced offspring, indicating a strong short-term

cost to the modifier (bottom-right corner in Fig. 3(b)). However, the modifier is actually strongly favored

by indirect selection (region II in Fig. 3(a)), which suggests that the modifier has a strong long-term

benefit by increasing the equilibrium mean fitness in the population.

I also investigated the influences of the genetic property of deleterious mutations. Consistent with

the results in the previous subsection, a modifier increasing the rate of sexual reproduction is more likely

to invade when the selection coefficient of small-effect deleterious mutations is smaller (Figs. S4(a)-(b)).

Also, a lower h reduces the relative fitness of sexually reproduced offspring relative to asexually

reproduced offspring, and thus reduces the parameter space for the modifier to invade (Figs. S4(c)-(d)).

For selection generated by large-effect, highly recessive mutations, a modifier increasing the rate

of sexual reproduction is indirectly selected against in a large parameter space (Fig. 3(c)). This is partly

contributed by the short-term disadvantage of the modifier, as sexually reproduced offspring tend to have

lower fitness than asexually reproduced offspring when deleterious alleles are very recessive (see Fig.

3(d)). However, similar to selection generated by small-effect deleterious mutations in Fig. 3(a), the

modifier is favored when the background sexual reproduction rate σ_0 is low and the selfing rate α is high

(bottom-right corner in Fig. 3(c)).

Regarding selection on a sex modifier when both categories of deleterious mutations are present

in the genome (compare Figs. 3(a) and 3(c)), the only parameter space where the modifier will always be

favored is when the background sexual reproduction rate σ_0 is low, and the selfing rate α is high (bottom-

left corner in panels of Fig. 3(a), 3(c)). On the other hand, in the parameter space represented by region III

in Fig. 3(a), the modifier should always be selected against. When σ_0 is relatively high, and α is relatively

low (region I in Fig. 3(a)), small-effect deleterious mutations and highly recessive, large-effect deleterious

mutations have opposing effects on the evolution of sex with comparable selective strength. Therefore, in

this parameter region, whether a modifier that increases the rate of sexual reproduction is favored or not

should depend on the relative genomic mutation rate of the two categories of mutations.

Strength of indirect selection caused by inter-locus interactions

Here I estimate how selfing impacts the strength of indirect selection generated by deleterious

mutations in the genome due to inter-locus interactions. This indirect selection can be caused by two

forces: deterministic interactions between deleterious mutations, and the Hill-Robertson effect in a finite

poupulation (Stetkenko and Roze 2022). Generally, unless the selfing rate is high, indirect selection on a

sex modifier caused by inter-locus interactions is weak compared to indirect selection caused by intra-

locus interactions.

Deterministic interactions between deleterious mutations

448   As Fig. 4(a) illustrates, even with a small map length ($R = 1$), indirect selection caused by

deterministic interactions between deleterious mutations is quite weak 1. Selfing causes selection against

sex when there is no epistasis (solid lines in Fig. 4(a)), but can favor sex when there is negative epistasis

between deleterious mutations (dashed lines in Fig. 4(a)). Generally, the background sexual reproduction

rate σ_0 only slightly influences how the strength of selection changes with the selfing rate, except when

the selfing rate is high. Under complete selfing, the strength of selection will reduce to 0 in a completely

sexually reproducing population since the inbreeding coefficient becomes $F = 1$ (Stetsenko and Roze

2022). In contrast, when there is asexual reproduction, the strength of selection will not reduce to 0 under

complete selfing (see results at $\alpha = 1$ in Fig. 4(a)), because F remains smaller than 1 (see equation (8a)).

The Hill-Robertson effect

458   Fig. 4(b) shows that selection on a sex modifier due to the Hill-Robertson is weak compared to

selection caused by intra-locus interactions ($\sim 10^{-4} - 10^{-3}$), unless the population size is small ($N \sim$

10^2), the background sexual reproduction rate σ_0 is low and the selfing rate α is high (black solid line in

Fig. 4(b)). As equations (13) and (14) indicate, selfing increases the strength of selection due to the Hill-

Robertson effect by reducing both the effective recombination rate and effective population size, while a

higher background sexual reproduction rate σ_0 reduces the selective strength, as illustrated in Fig. 4(b).

Overall effects of selfing on indirect selection caused by deleterious mutations

465   Results from individual-based simulations in Fig. 5 confirm that the evolution of a sex modifier is

mainly mediated by selection due to intra-locus interactions rather than inter-locus interactions.

Considering selection generated by small-effect deleterious mutations, a modifier that increases the rate of

sexual reproduction is generally disfavored when the background sexual reproduction rate σ_0 is low,

unless the selfing rate is high (line with $\sigma_0 = 0.2$ in Fig. 5(a)). Under a higher level of σ_0, as the selfing

rate increases, the fixation probability of the sex modifier is reduced within a specific range of the selfing

rate (see the range of α=0.6-0.9 at $\sigma_0 = 0.5, 0.9$ in Fig. 5(a)). For selection caused by highly recessive,

large-effect alleles, the modifier tends to be strongly disfavored under an intermediate selfing rate, unless

the selfing rate is high (Fig. 5(b)). Referring to the results at $\sigma_0 = 0.2, 0.5, 0.9$ in Fig. 3(a) and 3(c), the

above findings are qualitatively align with results for indirect selection through segregation caused by

intra-locus interactions.

Moreover, when the genomic recombination rate is low or the population size is small, selection

driven by intra-locus interactions still tends to be the dominant force, despite the increased strength of

selection caused by inter-locus interactions (Figs. 5(c), 5(d)). When the background sexual reproduction

rate σ_0 is low, a small map length or a small population size weakens selection against the sex modifier

(compare results at $\sigma_0 = 0.2$ in Figs. 5(c), 5(d) with 5(a)). Nevertheless, this effect may primarily stem

from a reduced short-term cost of sex caused by intra-locus interactions. Specifically, under tight linkage

or a small population size, there will be fewer segregating deleterious mutations, which can increase the

fitness of sexually reproduced offspring relative to asexually reproduced offspring (Kirkpatrick and

Batillon 2000). Moreover, under a higher level of σ_0 ($\sigma_0 = 0.5, 0.9$), the pattern of how the fixation

probability changes with the selfing rate α in Fig. 5(c) and 5(d) similar to Fig. 5(a).

Moreover, a high selfing rate can increase the strength of selection on a sex modifier by lowering

the effective recombination rate, it also reduces the effective population size (see equation (10)). As a

result, under a high selfing rate, the fixation probability is close to the neutral case when deleterious

mutations are partially dominant and have small effects (Figs. 5(a), (c), (d)).

Selection caused by the cost of meiosis and mate limitation

Previous subsections focus on indirect selection on a sex modifier caused by selection at fitness

loci. To quantify how selfing can reduce the strength of selection against sex by lowering the cost of

meiosis and providing reproductive assurance under mate limitation, I define the strength of mate

limitation τ as the reduced proportion of outcrossed ovules (Xu 2022). Therefore, the relative fertility via

outcrossing is $1 - \tau$ compared to selfing and asexual reproduction. The strength of selection on a sex

modifier can be obtained by setting $s = 0$ and $w_{asex} = w_{self} = 1/(1 - \tau)$ in the recursion equation (6).

For a rare modifier allele with small effect ($|\Delta\sigma| \ll 1$), the selective strength is approximately $s_e(h_\sigma +$

$(1 - h_\sigma)F)$, where the effective selection coefficient s_e is

499
$$s_e = -\frac{(1 - \alpha)(1 - c + \tau - \tau c(1 - 2\sigma_0))}{2(1 - c(1 - \sigma_0))(1 - \tau(1 - \alpha)\sigma_0)}\Delta\sigma. \tag{14}$$

In the denominator, the term $2(1 - c(1 - \sigma_0))$ is the amount of male gamete production, and the term

$1 - \tau(1 - \alpha)\sigma_0$ accounts for fitness reduction due to mate limitation. In the numerator, the term $1 - c$

and τ capture selection due to the cost of meiosis and mate limitation, respectively, while the $-\tau c(1 -$

$2\sigma_0)$ captures interactions between the two forces.

Fig. 6 shows that generally, the strength of selection caused by the cost of meiosis and mate

limitation is often comparable to indirect selection generated by deleterious mutations in the genome.

Interestingly, the strength of selection against the modifier is stronger when the background sexual

reproduction rate σ_0 is higher (compare solid and dashed lines in Fig. 6). This is because when there is

mate limitation, a higher σ_0 reduces the mean fitness of the population (the term $1 - \tau(1 - \alpha)\sigma_0$ in

equation (14)). Generally, a higher selfing rate tends to reduce the strength of selection. However, under

strong mate limitation ($\tau = 0.8$ in Fig. 6), selection against sexual reproduction is strongest under a low

selfing rate rather than under complete outcrossing. This is because although selfing reduces the effective

selection coefficient s_e, it strengthens selection by increasing homozygosity (the term $h_\sigma + (1 - h_\sigma)F$).

Discussion

This study investigates the effects of selfing on the evolution of an allele through three major

selective forces: (1) indirect selection caused by an altered rate of segregation at selected loci, referred to

as selection caused by intra-locus interactions, (2) indirect selection due to changes in the effective

recombination rate between selected loci, referred to as selection caused by inter-locus interactions, and

(3) selection due to the cost of meiosis and mate limitation. The effects of selfing on the three selective

forces are summarized in Table 1. Briefly, selfing affects indirect selection caused by both intra-locus and

inter-locus interactions by primarily increasing homozygosity across the genome, and generating

correlation between homozygotes at different loci, as indicated by equations (8a) and (8b). Also, similar

to asexual reproduction, selfing is also a mode of uniparental reproduction. Therefore, selfing can weaken,

selection against sexual reproduction due to the cost of meiosis and mate limitation, although may not

revert the direction.

The mechanisms by which selfing influences indirect selection caused by intra-locus selection are

complicated. One important mechanism is that selfing will induce higher homozygosity in sexually

reproduced offspring compared to asexually reproduced offspring. Therefore, when deleterious alleles are

recessive, similar to inbreeding depression, sexually reproduced offspring will be less fitness than

asexually reproduced offspring, resulting in a short-term disadvantage of a modifier that increases the rate

of sexual reproduction. Selfing also affects indirect selection by altering the genetic associations between

a modifier locus and fitness loci. Generally, under sporophytic selfing, a modifier that increases the rate

of sexual reproduction can invade in two scenarios: either when the deleterious allele is recessive enough

under weak selection, or when the allele is dominant enough under stronger selection. These results are

qualitatively similar to the findings under gametophytic selfing (Otto 2003).

Consequently, unless the selfing rate is high, highly recessive, large-effect deleterious mutations

in the genome often exert indirect selection against a modifier that increases the rate of sexual

reproduction through intra-locus interactions. On the other hand, partially recessive, small-effect

deleterious mutations can favor the evolution of sexual reproduction in a larger parameter space. In

general, a modifier that enhances the rate of sexual reproduction will be indirectly selected against by

intro-locus interactions at an intermediate selfing rate and when the sexual reproduction rate in the

population is low. Given that the map length in most organisms lies between 1 ~ 10 (Otto and Payseur

2019), indirect selection caused by inter-locus interactions should be much weaker than indirect selection

caused by intra-locus interactions. When the selfing rate is very high, both intra-locus and inter-locus

interactions will generate strong indirect selection for a modifier that increases the rate of sexual

reproduction.

By comparing the strength of the three selective forces on a sex modifier, I find that the dominant

selective force will depend on the rates of selfing and sexual reproduction. Under outcrossing, selection

on a sex modifier is primarily driven by the cost of meiosis and mate limitation. Deleterious mutations in

the genome result in weak indirect selection against a higher sexual reproduction rate. As the selfing rate

increases, selection due to the cost of meiosis and mate limitation gets weaker, while indirect selection

through intra-locus interactions becomes stronger and may become the dominant force. In highly selfing

populations, indirect selection caused by both intra-locus and inter-locus interactions is significant.

Furthermore, under a higher rate of sexual reproduction, selection on a sex modifier due to the cost of

meiosis and mate limitation intensifies, while indirect selection generated by deleterious mutations

weakens. This suggests that evolutionary transition from complete sexual reproduction to partially

asexual reproduction may initially be driven by the cost of meiosis and mate limitation.

The current results suggest that high selfing rate should favor the evolution of sexual reproduction

is consistent with the evidence that populations with high selfing rates are rarely found in partially clonal

herbs (Vallejo-Marín et al. 2010). Empirical evidence show that apomictic populations often derive from

self-incompatible populations (Hörandl 2010, Hojsgaard and Hörandl 2019). However, our results

suggest that intermediate selfing rates tend to favor the evolution of asexual reproduction, suggesting that

asexual reproduction tends to evolve before, instead of after, the evolution of selfing. In fact, equation

(12b) shows that asexual reproduction will inhibit the evolution of selfing by increasing inbreeding

depression. This hypothesis may be tested based on ancestral state reconstructions.

The offers a novel interpretation for the observed association between vegetative reproducing

populations and self-incompatibility/outcrossing (Ellstrand and Roose 1987, Vallejo-Marín et al. 2010).

Traditionally, it has been proposed that in partially clonal populations, outcross-pollination among cloned

individuals (geitonogamy) will produce offspring equivalent to those produced through selfing, thus

suffering the cost of inbreeding depression (Vallejo-Marín et al. 2010). Consequently, mechanisms

avoiding geitonogamy may be favored, such as self-incompatibility and dichogamy (temporal separation

of female and male function). However, I find that assuming no spatial structure, even with geitonogamy,

the fitness of outcrossed offspring is always higher than of asexually reproduced offspring (equation

(12a)). Moreover, equation (12b) reveals that a higher level of asexual reproduction can lower the fitness

of selfed offspring, potentially selecting against alleles that increases the rate of selfing. Thus, in partially

clonal populations, mechanisms promoting obligate outcrossing may be primarily selected for by

avoiding intra-individual self-fertilization, instead of geitonogamy.

The current study did not explore the impact of selfing on the evolution of sex due to the cost of

males. However, empirical evidence suggests that selfing may generally weaken the cost of males by

reducing resource allocation to the male function in plants. Higher selfing rates are correlated with lower

pollen-ovule ratios (Cruden 1977, Cruden and Lyon 1985, de Cunha and Aizen 2023) and smaller flower

display size (Sicard et al. 2011). While a corresponding increase in allocation to the female function

might be anticipated, the evidence is less conclusive (Cruden 1977, Cruden and Lyon 1985). In some

groups, selfing species produce more ovules per flower than outcrossing species (Moine and Anderson

1992, Delesalle et al. 2008). In other groups, selfing reduces the ovule number per flower (Ritland and

Ritland 1989). This may be because in highly selfing populations, male and female gamete production is

expected to be perfectly correlated (Mazer et al. 2007). Also, selfing populations may experience smaller

stochastic variation in mating success, which selects for fewer ovules (Burd et al. 2009). Ovule

production may not fully reflect female function allocation. In fact, self-compatible species tend to have a

high fruit-flower ratio than self-incompatible species, indicating a greater allocation to the female

function (Sutherland 1986).

The current study investigates the evolution of sexual reproduction within a population by

considering a modifier locus. Another category of studies on the evolution of sex looks at selection

between populations, by investigating how the sexual reproduction rate affects the mean fitness of a

population (reviewed in Otto (2009)). Analyses in Section I of Supplementary Materials suggests that

selfing tends to increase the strength of sexual reproduction caused by selection between populations.

In conclusion, this study indicates that an intermediate selfing rate tends to impede the evolution

of sexual reproduction, especially when the baseline sexual reproduction rate is low. A high selfing rate

strongly favors the evolution of sexual reproduction. Regarding empirical evidence, these findings

suggest that the evolution of asexual reproduction may often precede the evolution of selfing.

References

Agrawal, A. F., & Chasnov, J. R. (2001). Recessive mutations and the maintenance of sex in structured populations. *Genetics*, 158(2), 913-917.

Asker, S. E., & Jerling, L. (2017). Apomixis in plants. Routledge.

Barrett, S. C. (2008). Major evolutionary transitions in flowering plant reproduction: an overview. *International Journal of Plant Sciences*, 169(1), 1-5.

Barrett, S. C. (2015). Influences of clonality on plant sexual reproduction. *Proceedings of the National Academy of Sciences*, 112(29), 8859-8866.

Barton N. H. (1995). A general model for the evolution of recombination. *Genetic Research*, 65, 123-144.

Bell, G. (1982). The paradox of sexuality. The masterpiece of nature: The evolution and genetics of sexuality, 19-78.

Bengtsson, & Ceplitis. (2000). The balance between sexual and asexual reproduction in plants living in variable environments. *Journal of Evolutionary Biology*, 13(3), 415-422.

Burd, M., Ashman, T. L., Campbell, D. R., Dudash, M. R., Johnston, M. O., Knight, T. M., ... & Vamosi, J. C. (2009). Ovule number per flower in a world of unpredictable pollination. *American Journal of Botany*, 96(6), 1159-1167.

Busch, J. W., & Delph, L. F. (2012). The relative importance of reproductive assurance and automatic selection as hypotheses for the evolution of self-fertilization. *Annals of botany*, 109(3), 553-562.

Caballero, A., & Hill, W. G. (1992). Effective size of nonrandom mating populations. *Genetics*, 130(4), 909-916.

Charlesworth, B. (1980). The cost of sex in relation to mating system. Journal of Theoretical Biology, 84(4), 655-671.

Charlesworth, B. (2013). Background selection 20 years on: the Wilhelmine E. Key 2012 invitational lecture. *Journal of Heredity*, 104(2), 161-171.

Charlesworth, B., Charlesworth, D., & Morgan, M. T. (1990). Genetic loads and estimates of mutation rates in highly inbred plant populations. *Nature*, 347(6291), 380-382.

Charlesworth, D., & Willis, J. H. (2009). The genetics of inbreeding depression. *Nature reviews genetics*, 10(11), 783-796.

Charlesworth, D., Charlesworth, B., & Strobeck, C. (1979). Selection for recombination in partially self-fertilizing populations. *Genetics*, 93(1), 237-244.

Chasnov, J. R. (2000). Mutation-selection balance, dominance and the maintenance of sex. *Genetics*, 156(3), 1419-1425.

Cruden, R. W. (1977). Pollen-ovule ratios: a conservative indicator of breeding systems in flowering plants. *Evolution*, 32-46.

Cruden, R. W., & Lyon, D. L. (1985). Patterns of biomass allocation to male and female functions in plants with different mating systems. *Oecologia*, *66*, 299-306.

da Cunha, N. L., & Aizen, M. A. (2023). Pollen production per flower increases with floral display size across animal‑pollinated flowering plants. *American Journal of Botany*, e16180.

Damgaard, C., & Abbott, R. J. (1995). Positive correlations between selfing rate and pollen-ovule ratio
within plant populations. *Evolution*, 214-217.

Damgaard, C., & Loeschcke, V. (1994). Genotypic variation for reproductive characters, and the
influence of pollen-ovule ratio on selfing rate in rape seed (Brassica napus). Journal of Evolutionary
Biology, 7(5), 599-607.

Delesalle, V. A., Mazer, S. J., & Paz, H. (2008). Temporal variation in the pollen: ovule ratios of Clarkia
(Onagraceae) taxa with contrasting mating systems: field populations. *Journal of Evolutionary Biology*,
21(1), 310-323.

Eckert, C. G. (1999). Clonal plant research: proliferation, integration, but not much evolution. *American
Journal of Botany*. 86(11), 1649-1654.

Eckert, C. G., Samis, K. E., Dart, S., Harder, L. D., & Barrett, S. C. H. (2006). Reproductive assurance
and the evolution of uniparental reproduction in flowering plants. Ecology and evolution of flowers, 183,
203.

Ellstrand, N. C., & Roose, M. L. (1987). Patterns of genotypic diversity in clonal plant species. *American
Journal of Botany*, 74(1), 123-131.

Eyre-Walker, A., & Keightley, P. D. (2007). The distribution of fitness effects of new mutations. *Nature
Reviews Genetics*, 8(8), 610-618.

Glémin, S. (2007). Mating systems and the efficacy of selection at the molecular level. *Genetics*, 177(2),
905-916.

Goldberg, E. E., Kohn, J. R., Lande, R., Robertson, K. A., Smith, S. A., & Igić, B. (2010). Species
selection maintains self-incompatibility. *Science*, 330(6003), 493-495.

Goodwillie, C., Kalisz, S., & Eckert, C. G. (2005). The evolutionary enigma of mixed mating systems in
plants: occurrence, theoretical explanations, and empirical evidence. *Annu. Rev. Ecol. Evol. Syst.*, 36, 47-
79.

Haldane, J. B. (1919). The combination of linkage values and the calculation of distances between the loci
of linked factors. *J Genet*, 8(29), 299-309.

Hartfield, M., Bataillon, T., & Glémin, S. (2017). The evolutionary interplay between adaptation and self-
fertilization. *Trends in Genetics*, 33(6), 420-431.

Hill W. G., & Robertson. A. (1966) The effect of linkage on limits to artificial selection. *Genetic
Research*, 8, 269-294.

Hojsgaard, D., & Hörandl, E. (2019). The rise of apomixis in natural plant populations. *Frontiers in Plant
Science*, 10, 358.

Honnay, O., & Jacquemyn, H. (2008). A meta-analysis of the relation between mating system, growth
form and genotypic diversity in clonal plant species. *Evolutionary Ecology*, 22, 299-312.

Hörandl, E. (2010). The evolution of self-fertility in apomictic plants. Sexual plant reproduction, 23(1),
73-86.

Innes, D. J., Fox, C. J., & Winsor, G. L. (2000). Avoiding the cost of males in obligately asexual Daphnia
pulex (Leydig). *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 267(1447),
991-997.

Jarne, P., & Charlesworth, D. (1993). The evolution of the selfing rate in functionally hermaphrodite
plants and animals. *Annual Review of Ecology and Systematics*, 24(1), 441-466.

Keightley, P. D., & Otto, S. P. (2006). Interference among deleterious mutations favours sex and
recombination in finite populations. *Nature*, 443(7107), 89-92.

Klekowski, E. J., & Godfrey, P. J. (1989). Ageing and mutation in plants. *Nature*, 340(6232), 389-391.

Kondrashov, A. S. (1985). Deleterious mutations as an evolutionary factor. II. Facultative apomixis and
selfing. Genetics, 111(3), 635-653.

Kron, P., Stewart, S. C., & Back, A. (1993). Self-compatibility, autonomous self-pollination, and insect-
mediated pollination in the clonal species Iris versicolor. *Canadian Journal of Botany*, 71(11), 1503-1509.

Kumpulainen, T., Grapputo, A., & Mappes, J. (2004). Parasites and sexual reproduction in psychid moths.
Evolution, 58(7), 1511-1520.

Lande, R., & Schemske, D. W. (1985). The evolution of self-fertilization and inbreeding depression in
plants. I. Genetic models. *Evolution*, 39(1), 24-40.

Lehtonen, J., Jennions, M. D., & Kokko, H. (2012). The many costs of sex. *Trends in ecology & evolution*,
27(3), 172-178.

Lewis W. M. Jr. 1987. The cost of sex. Pages 33–57 in The Evolution of Sex and its Consequences,
edited by S. C. Stearns. Basel (Switzerland): Birkhäuser Verlag.

Marriage, T. N., & Orive, M. E. (2012). Mutation-selection balance and mixed mating with asexual
reproduction. *Journal of theoretical biology*, 308, 25-35.

Mazer SJ, Delesalle VA, Paz H. 2007. Evolution of mating system and the genetic covariance between
male and female investment in Clarkia (onagraceae): selfing opposes the evolution of trade-offs.
Evolution 61: 83 – 98.

Meirmans, S., Meirmans, P. G., & Kirkendall, L. R. (2012). The costs of sex: facing real-world
complexities. *The Quarterly Review of Biology*, 87(1), 19-40.

Michaels, H. J., & Bazzaz, F. A. (1986). Resource allocation and demography of sexual and apomictic
Antennaria parlinii. *Ecology*, 67(1), 27-36.

Mione, T., & Anderson, G. J. (1992). Pollen-ovule ratios and breeding system evolution in Solanum
section Basarthrum (Solanaceae). *American Journal of Botany*, 79(3), 279-287.

Muirhead, C. A., & Lande, R. (1997). Inbreeding depression under joint selfing, outcrossing, and
asexuality. *Evolution*, 51(5), 1409-1415.

Mukai, T., Chigusa, S. I., Mettler, L. E., & Crow, J. F. (1972). Mutation rate and dominance of genes
affecting viability in Drosophila melanogaster. *Genetics*, 72(2), 335-355.

Nakayama, Y., Seno, H., & Matsuda, H. (2002). A population dynamic model for facultative
agamosperms. *Journal of theoretical biology*, 215(2), 253-262.

Nordborg, M., Charlesworth, B., & Charlesworth, D. (1996). The effect of recombination on background
selection. *Genetics Research*, 67(2), 159-174.

Otto, S. P. (2003). The advantages of segregation and the evolution of sex. *Genetics*, 164(3), 1099-1118.

Otto, S. P. (2009). The evolutionary enigma of sex. *The American Naturalist*, 174(S1), S1-S14.

Otto, S. P., & Barton, N. H. (2001). Selection for recombination in small populations. *Evolution*, 55(10),
1921-1931.

Otto, S. P., & Payseur, B. A. (2019). Crossover interference: shedding light on the evolution of
recombination. *Annual review of genetics*, 53, 19-44.

Richards, A. J. (2003). Apomixis in flowering plants: an overview. *Philosophical Transactions of the
Royal Society of London. Series B: Biological Sciences*, 358(1434), 1085-1093.

Ritland, C., & Ritland, K. (1989). Variation of sex allocation among eight taxa of the Mimulus guttatus
species complex (Scrophulariaceae). *American Journal of Botany*, 76(12), 1731-1739.

Roze, D. (2014). Selection for sex in finite populations. *Journal of evolutionary biology*, 27(7), 1304-
1322.

Roze, D. (2021). A simple expression for the strength of selection on recombination generated by
interference among mutations. *Proceedings of the National Academy of Sciences*, 118(19).

Roze, D., & Barton, N. H. (2006). The Hill–Robertson effect and the evolution of recombination.
Genetics, 173(3), 1793-1811.

Roze, D., & Lenormand, T. (2005). Self-fertilization and the evolution of recombination. *Genetics,* 170(2),
841-857.

Sicard, A., & Lenhard, M. (2011). The selfing syndrome: a model for studying the genetic and
evolutionary basis of morphological adaptation in plants. Annals of botany, 107(9), 1433-1443.

Silvertown, J. (2008). The evolutionary maintenance of sexual reproduction: evidence from the ecological
distribution of asexual reproduction in clonal plants. *International journal of plant sciences*, 169(1), 157-
168.

Simmons, M. J., & Crow, J. F. (1977). Mutations affecting fitness in Drosophila populations. *Annual
review of genetics*, 11(1), 49-78.

Smith, J. M., & Maynard-Smith, J. (1978). The evolution of sex (Vol. 4). Cambridge: Cambridge
University Press.

Soltis, D. E., & Soltis, P. S. (1992). The distribution of selfing rates in homosporous ferns. *American
Journal of Botany*, 79(1), 97-100.

Stephan, W. (2010). Genetic hitchhiking versus background selection: the controversy and its
implications. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365(1544), 1245-
1253.

Stetsenko, R., & Roze, D. (2022). The evolution of recombination in self-fertilizing
organisms. *Genetics*, *222*(1), iyac114.

Sutherland, S. (1986). Patterns of fruit‐set: What controls fruit‐flower ratios in plants?. *Evolution*, 40(1),
117-128.

Uyenoyama, M. K., & Bengtsson, B. O. (1989). On the origin of meiotic reproduction: a genetic modifier
model. *Genetics*, 123(4), 873-885.

Vallejo-Marín, M., Dorken, M. E., & Barrett, S. C. (2010). The ecological and evolutionary consequences
of clonality for plant mating. *Annual Review of Ecology, Evolution, and Systematics*, 41, 193-213.

Vitalis, R., & Couvet, D. (2001). Two-locus identity probabilities and identity disequilibrium in a
partially selfing subdivided population. *Genetics Research*, 77(1), 67-81.

Whitton, J., Sears, C. J., Baack, E. J., & Otto, S. P. (2008). The dynamic nature of apomixis in the
angiosperms. *International Journal of Plant Sciences*, 169(1), 169-182.

Williams, G. C. (1975). Sex and evolution. Princeton Univ. Press, Princeton, NJ.

Wolinska, J., & Lively, C. M. (2008). The cost of males in Daphnia pulex. *Oikos*, 117(11), 1637-1646.

Xu, K. (2022). The genetic basis of selfing rate evolution. *Evolution*, 76(5), 883-898.

Tables

Table 1: A summary of selective forces on a sex modifier and the effects of selfing on each force.

Selective forces on a sex modifier	Causes of the selective force	How selfing affects the selective force		Related figures in this paper
Indirect selection due to interactions between alleles within a selected locus (intra-locus interaction)	Changes in the rate of segregation	Alter the fitness of sexually reproduced offspring relative to asexually reproduced offspring	1. An excess of homozygosity at a locus 2. Generate positive association between homozygotes at different loci.	Fig. 1, 2, 3
Indirect selection due to interactions between selected loci (inter-locus interactions)	Changes of the effective recombination rate across the genome	Reduce the effective recombination rate in the genome	3. Alter genetic association between the sex modifier locus and fitness loci	Fig. 4
Cost of males	Resources cost to produce males or the male function	Reduce resource allocation to the male function		N/A
Cost of meiosis	Sexual individuals transmit only half of the genome to the offspring	Reduce the cost of meiosis as a mode of uniparental reproduction		Fig. 6(b)
Mate limitation	Outcrossing requires availability of gametes from other individuals	Reduce the strength of mate limitation by providing reproductive assurance		Figs. 6(a), 6(b)

763   **Table 2.** Biological meaning of key symbols used in the model.

Symbol	Biological meaning
α	Sporophytic selfing rate within sexual reproduction
σ_0	Background sexual reproduction rate in the population when the sex modifier locus is fixed with allele M.
$\Delta\sigma$	Effects of a sex modifier on changing the rate of sexual reproduction
h_σ	Dominance of a sex modifier
s, h	Selection coefficient and dominance of allele a at the fitness locus
μ	Mutation rate from allele A to a deleterious allele a at a fitness locus
r	Recombination rate between the sex modifier and fitness loci
U	Genomic mutation rate of deleterious mutations
R	Chromosome map length/Mean number of crossovers in the genome
F	Inbreeding coefficient, which is the probability of identity by descent at a locus
ϕ	The probability of identity by descent at two loci
c	Reduction in the male gamete output due to asexual reproduction, measuring the strength of cost of meiosis. There is no cost of meiosis when $c = 1$.
τ	Strength of mate limitation
d_a	Fitness reduction of asexually reproduced offspring relative to outcrossed offspring caused by a single locus
d_s	Fitness reduction of selfed offspring relative to outcrossed offspring caused by a single locus

Figure 1. Invasibility of a modifier that increases sexual reproduction when allele a at the fitness locus is under selection-mutation balance (panels (a), (b)), and when the fitness locus experiences selective sweep (panels (c), (d)). The symbols "+" and "-" mean that the modifier can invade (or increase in frequency) and cannot invade, respectively. Panels (a) and (c) show results under outcrossing ($\alpha = 0$). Panels (b) and (d) show results when there is selfing. In panel (b), the modifier can invade either at the bottom left or upper right region of the parameter space. In panel (d), the modifier increases in frequency in the top left region or the region between the two lines. For panels (c) and (d), the initial and final frequency of allele A is $p_{A,0} = 0.001, p_{A,T} = 0.999$, and the initial frequency of the modifier allele m is $p_{m,0} = 0.001$. $\sigma_0 = 0.5$ for panels (a) and (b), $\sigma_0 = 0.2$ for panels (c) and (d). Other parameters are $\Delta\sigma = 0.01, h_\sigma = 0.5, r = 0.5, \mu = 10^{-5}, c = 1, \tau = 0$. The magnitude of the modifier effect $\Delta\sigma$ only slightly affects the invasion conditions (Fig. S1).

Figure 2. Effects of the dominance coefficient of the modifier (h_σ) and background sexual reproduction
rate (σ_0) on the invasibility of a modifier that increases sexual reproduction under sporophytic selfing.
Results are based on numerical solution of the eigenvalue. Left and right columns show the case when the
fitness locus contains deleterious mutations at selection-mutation balance (panels (a)-(c)) and when the
fitness locus experiences selective sweep (panels (d)-(f)), respectively. The meanings of "+" and "-" are
the same as in Figure 1. For panels (d)-(e), the initial and final frequency of allele A is $p_{A,0} =$
$0.001, p_{A,T} = 0.999$, and the initial frequency of m is $p_{m,0} = 0.001$. Other parameters are $\Delta\sigma =$
$0.01, \alpha = 0.1, r = 0.5, \mu = 10^{-5}, c = 1, \tau = 0$.

Figure 3. Genome-wide strength of indirect selection (in the order of 10^{-4}) on a modifier increasing the sexual reproduction rate caused by within-locus interactions. The upper and bottom panels show selection caused by small-effect and large-effect deleterious mutations. Panels (a) and (c) show the overall strength of indirect selection. Panels (b) and (d) show the strength of short-term (dis)advantage of the modifier due to fitness differences between sexually and asexually reproduced offspring. Similar results are found under gametophytic selfing (Fig. S3). I use $s = 0.05, h = 0.2, U = L\mu = 0.2$ for small-effect deleterious mutations, and $s = 0.8, h = 0.02, U = L\mu = 0.025$ for large-effect deleterious mutations (Mukai et al. 1972, Simmons and Crow 1977, Klekowski and Godfrey 1989). Other parameters are $\Delta\sigma = 0.01, h_\sigma = 0.5, c = 1, \tau = 0, r = 0.5$.

Figure 4. Strength of indirect selection ($\Delta p_m / p_m (1 - p_m)$) on a rare sex modifier due to effective changes in the recombination rate. Panels (a) and (b) show the strength of selection caused by deterministic interactions between deleterious mutations, and the Hill-Robertson effect in a finite population, respectively. In panel (a), solid lines show results when fitness loci have multiplicative interactions on individual fitness (no epistasis). Dashed lines show results under additive-additive epistasis, so that for two loci j and k, the fitness of genotype $A_j a_j A_k a_k$, $A_j a_j a_k a_k$, and $a_j a_j a_k a_k$ is further changed from the multiplication by $e_{a \times a}$, $2e_{a \times a}$, and $4e_{a \times a}$, respectively. Other parameters are $s = 0.05, h = 0.2, U = 0.2, \Delta\sigma = 0.01, h_\sigma = 0.5, R = 1$.

Figure 5. Effects of selfing on the fixation probability of a sex modifier allele with deleterious mutations
from individual-based simulations. The fixation probability is divided by that under the neutral case $1/2N$.
The genomic mutation rate is $U = 0.2$ for small-effect deleterious mutations (panels (a), (c), (d)), and
$U = 0.05$ for highly recessive, large-effect mutations (panel (b)). Other parameters are $\Delta\sigma = 0.05, h_\sigma = 0.5, c = 1, \tau = 0$.

815 **Figure 6.** Effects of the selfing rate on the strength of selection against a modifier that increases sex
816 ($\Delta\sigma = 0.01$, $h_\sigma = 0.5$) caused by the cost of meiosis and mate limitation. Panels (a) and (b) show results
817 when there is no and strong cost of meiosis, respectively. Lines with different colors show results under
818 different strength of mate limitation. The strength of selection is calculated based on equation (14).