The ability to sense the environment is heterogeneously distributed in cell populations

Andrew Goetz[1*], Hoda Akl[2*], and Purushottam Dixit[1,2,3,4]

[1]Department of Biomedical Engineering, Yale University, New Haven CT
[2]Department of Physics, University of Florida, Gainesville FL
[3]Systems Biology Institute, Yale University, West Haven, CT
[4]Corresponding author: purushottam.dixit@yale.edu
[*]Contributed equally

Abstract

Channel capacity of signaling networks quantifies their fidelity in sensing extracellular inputs. Low estimates of channel capacities for several mammalian signaling networks suggest that cells can barely detect the presence/absence of environmental signals. However, given the extensive heterogeneity and temporal stability of cell state variables, we hypothesize that the sensing ability itself may depend on the state of the cells. In this work, we present an information theoretic framework to quantify the distribution of sensing abilities from single cell data. Using data on two mammalian pathways, we show that sensing abilities are widely distributed in the population and most cells achieve better resolution of inputs compared to an "*average cell*". We verify these predictions using live cell imaging data on the IGFR/FoxO pathway. Importantly, we identify cell state variables that correlate with cells' sensing abilities. This information theoretic framework will significantly improve our understanding of how cells sense in their environment.

Introduction

In cell populations, there is a significant overlap in responses to environmental stimuli of differing

strengths. This raises a fundamental question[1]: do signaling networks in cells relay accurate information

about their environment to take appropriate action? And if not, where along the signal transduction

pathway is the information lost[2, 3]? Mutual information (MI) quantifies the information content in an

intracellular output about extracellular inputs. For an input u (e.g., concentration of a ligand) and an

output x (e.g., intracellular species concentration[4-6] or a cellular phenotype[7, 8]), the MI is defined as[9]

$$I = \sum_{x,u} p(x|u)p(u) \log_2 \frac{p(x|u)}{\sum_{u'} p(x|u')p(u')}. \tag{1}$$

Experimental single cell methods such as flow cytometry[10], immunofluorescence[10], mass spectrometry[11],

or live cell imaging[12] allow us to estimate response histograms $p(x|u)$ across several inputs. Using these

distributions, we can estimate the maximum of the MI (the channel capacity, CC) by optimizing Eq. 1

over input distributions $p(u)$. The CC quantifies fidelity of signal transduction[1, 2]. For example, a CC of 1

bit implies that that the cells can *at best* distinguish between two input levels (e.g. presence versus

absence), with higher CCs indicating that cells can resolve multiple input states. Importantly, CC can be

used to identify bottlenecks in signaling[2, 13].

The CC has been estimated for input-output relationships in several mammalian signaling networks[4-6, 14-

16]. When the output is defined as levels of a single protein a fixed time, the CC was found to be

surprisingly low, $\sim 1 - 1.5$ bits. These estimates have been improved by considering multidimensional

outputs[5] or time varying inputs[17]. While these modifications led to somewhat higher CC estimates, the

overall conclusion that cells know little about their environment remains well established. In contrast,

significantly higher CC estimates have been found when the output at the level of cell population

averages are considered[6], suggesting that the only way to overcome low sensing fidelity is population

average response.

These previous calculations estimated one channel capacity for all cells in a population, implicitly

assuming that individual cells have similar sensing abilities. However, we now know that cells exhibit

extensive heterogeneity in cell state variables[18, 19] such as abundances of key signaling proteins[20, 21],

mRNA abundances[22], and chromatin conformation[23] and accessibility[24]. Notably, the time scale of

fluctuations in these variables can be significantly slower than relevant signaling time scales[25, 26],

sometimes extending across multiple generations[27]. Heterogeneity in cell state variables leads to a

heterogeneity in response to extracellular cues, including chemotherapeutic drugs[25, 26, 28], mitogens[29],

hormones[30], chemotactic signals[7, 27, 31, 32], and other electrical and chemical stimuli[33, 34]. Therefore, we

hypothesize that that the ability to sense the environment varies from cell-to-cell in populations, in a cell

state dependent manner.

There is no conceptual framework to estimate the variation in sensing abilities in cell populations and its

dependence on cell state variables. To that end, we introduce an information theoretic quantity CeeMI:

Cell stat**e** dependent **M**utual **I**nformation. We show that using typically collected single cell data and

computational models of signaling networks, we can estimate the distribution $p_{\text{CeeMI}}(I)$ of single cell

signaling fidelities (single cell mutual information values). We also show that we can identify cell state

variables that make some cells better and others worse at sensing their environment.

Using an illustrative example, we show that in heterogeneous cell populations, estimates of mutual

information that average over cell states can be significantly lower than the mutual information of

signaling networks in typical cells. Next, using previously collected experimental data, we estimate

$p_{\text{CeeMI}}(I)$ for two important mammalian signaling pathways[35]; the Epidermal growth factor (EGF)/EGF

receptor pathway and the Insulin like growth factor (IGF)/Forkhead Box protein O (FoxO) pathway, we

show that while the cell state agnostic CC estimates for both pathways are $\sim$ 1 bit, most individual cells

are predicted to be significantly better at resolving different inputs. Using live cell imaging data for the

IGF/FoxO pathway, we show that our estimate of variability in sensing abilities matches closely with

experimental estimates. Importantly, for this pathway, we also verify our prediction that specific cell

state variables dictate cells' sensing abilities. Finally, using a simple receptor/ligand model, we show

how the time scales cell state dynamics affects cells' individuality and therefore sensing abilities. We

believe that CeeMI will be an important tool to explore variability in cellular sensing abilities and in

identifying what makes some cells better and others worse in sensing their environment.

Results

Conditional mutual information models single cells as cell state dependent channels

Consider a cell population where cells are characterized by state variables $\boldsymbol{\theta}$. These include abundances

of signaling proteins and enzymes, epigenetic states, etc. We assume that cell states are temporally

stable, that is, $\boldsymbol{\theta}$ remains constant over a time scale that is longer than typical fluctuations in

environmental inputs. Later, using a toy simple model, we explicitly study the effect of cell state

dynamics on cells' ability to sense their environment.

We assume that cell state variables are distributed in the population according to a distribution $p(\boldsymbol{\theta})$. If

x denotes an output of choice (e.g., phosphorylation levels of one or more protein(s) at one or more

time point(s)) and u denotes the input (e.g., ligand concentration), the experimentally measured

response distribution $p(x|u)$ can be decomposed as:

$$p(x|u) = \int p(x|u,\boldsymbol{\theta})p(\boldsymbol{\theta})d\boldsymbol{\theta}. \qquad (2)$$

117   where $p(x|u,\boldsymbol{\theta})$ is the distribution of the output x conditioned on the input u *and* cell state variables $\boldsymbol{\theta}$.

118   We note that in most cases, the same cell cannot be probed multiple times. Consequently, $p(x|u,\boldsymbol{\theta})$

119   may not be experimentally accessible. However, it is conceptually well defined and mathematically

120   accessible when interactions amongst molecules are specified[36].

121
Figure 1. A schematic of our computational approach. (top) single cell data across different input conditions and
time points are integrated with a stochastic model of a signaling network using a previous developed maximum
entropy approach leading to a distribution over signaling network parameters $p(\boldsymbol{\theta})$ (middle). (bottom) In silico
cells are generated using the inferred parameter distribution and cell-state specific mutual information $I(\boldsymbol{\theta})$ and

population distribution of cell performances $p_{\text{CeeMI}}(I)$ is estimated. The model also evaluates the correlation
between cells' performance and biochemical parameters.
128

129 Using $p(x|u, \boldsymbol{\theta})$, we can define the cell state dependent mutual information $I(\boldsymbol{\theta})$ for a fixed input
130 distribution $p(u)$ analogously to Eq. 1:

$$I(\boldsymbol{\theta}) = \sum_{x,u} p(x|u, \boldsymbol{\theta})p(u) \log_2 \frac{p(x|u, \boldsymbol{\theta})}{\sum_{u'} p(x|u', \boldsymbol{\theta})p(u')}. \tag{3}$$

131 $I(\boldsymbol{\theta})$ quantifies individual cells' ability to sense their environment as a function of the cell state
132 parameters $\boldsymbol{\theta}$. The distribution $p_{\text{CeeMI}}(I)$ of single cell sensing abilities is

$$p_{\text{CeeMI}}(I) = \int p(\boldsymbol{\theta}) \delta(I - I(\boldsymbol{\theta})) d\boldsymbol{\theta} \tag{4}$$

133 where $\delta(\cdot)$ is the Dirac delta function. We can also compute the joint distribution between the single
134 cell mutual information and any cell state variable of interest χ (e.g., abundance of cell surface
135 receptors):

$$p_{\text{CeeMI}}(I, \chi) = \int p(\boldsymbol{\theta}) \delta(I - I(\boldsymbol{\theta})) \delta(\chi - \chi(\boldsymbol{\theta})) d\boldsymbol{\theta} \tag{5}$$

136 where $\chi(\boldsymbol{\theta})$ is the value of the biochemical parameter when cell state variables are fixed at $\boldsymbol{\theta}$. The
137 distributions in Eq. 5 quantify the interdependency between a cell's signaling fidelity $I(\boldsymbol{\theta})$ and cell
138 specific biochemical parameters $\chi(\boldsymbol{\theta})$. As we will see below, the distributions in Eq. 4 and Eq. 5 can be
139 experimentally verified when appropriate measurements are available. Finally, we define the population
140 average of the cell state dependent mutual information:

$$I_{\text{Cee}} = \int I(\boldsymbol{\theta}) \, p(\boldsymbol{\theta}) d\boldsymbol{\theta}. \tag{6}$$

In information theory, I_{Cee} is known as the conditional mutual information[9] between the input u and
the output x conditioned on $\boldsymbol{\theta}$. If cell state variables remain constant and if distribution over cell states
is independent of the input distribution, it can be shown that $I_{\text{Cee}} \geq I$ (SI Section 1). It then follows that
at least some cells in the population have better sensing abilities compared to the cell state agnostic
mutual information (Eq. 1).

Since I_{Cee} depends on the input distribution $p(u)$, we can find an optimal input distribution $p(u)$ that
maximizes I_{Cee} (SI Section 1). Going forward, unless an input distribution is specified, the distributions
$p_{\text{CeeMI}}(I)$ and $p_{\text{CeeMI}}(I, \chi)$ are discussed in the context of this optimal input distribution.

Maximum entropy inference can estimate $p_{\text{CeeMI}}(I)$

Estimation of $p_{\text{CeeMI}}(I)$ requires decomposing the experimentally observed response $p(x|u)$ into cell

specific output distributions $p(x|u, \boldsymbol{\theta})$ and the distribution of cell state variables $p(\boldsymbol{\theta})$ (Eq. 3 and 4). This

problem is difficult to solve given that neither $p(x|u, \boldsymbol{\theta})$ nor $p(\boldsymbol{\theta})$ are accessible in typical experiments.

However, for many signaling networks, stochastic biochemical models can be constructed to obtain the

cell specific output distribution $p(x|u, \boldsymbol{\theta})$. Here, $\boldsymbol{\theta}$ comprise protein concentrations and effective rates

of enzymatic reactions and serve as a proxy for cell state variables. Given the experimentally measured

cell state averaged response $p(x|u)$ and the model predicted cell specific output distribution $p(x|u, \boldsymbol{\theta})$,

we need a computational method to infer $p(\boldsymbol{\theta})$ (see Eq. 2). The problem of inference of parameter

heterogeneity is a non-trivial inverse problem[37]. While there are several proposed methods to solve this

problem (reviewed in[38]), most cannot efficiently infer $p(\boldsymbol{\theta})$ for signaling networks with even a moderate

($n \sim 10$) number of parameters. Here, we use our previously develop maximum entropy-based

approach to infer $p(\boldsymbol{\theta})$[37]. This way, we can estimate $p_{\text{CeeMI}}(I)$ using experimentally obtained cell state

agnostic response $p(x|u)$ and a stochastic biochemical model $p(x|u, \boldsymbol{\theta})$ of the underlying signaling

network.

An "average cell" can discern much less than a typical cell about the environment

To illustrate the effect of heterogeneity of cell state variables on the cell state agnostic estimate of

mutual information (which we call I_{CSA} from now onwards), we consider a simple stochastic biochemical

network of a receptor-ligand system. Details of the model and the calculations presented below can be

found in SI Section 2. Briefly, extracellular ligand L binds to cell surface receptors R. Steady state levels

of the ligand bound receptor B is considered the output. The signaling network is parametrized by

several cell state variables $\boldsymbol{\theta}$ such as receptor levels, rates of binding/unbinding, etc. For simplicity, we

assume that only one variable, steady state receptor level R_0 in the absence of the ligand, varies in the

population. Calculations with variability in other parameters are presented in SI Section 2.

In this population, cells' response $p(B|L, R_0)$ is distributed as a Poisson distribution whose mean is

proportional to the cell state variable R_0 (SI Section 2). That is, when all other parameters are fixed, a

higher R_0 leads to lower noise (coefficient of variation). To calculate cell state dependent mutual

information (Eq. 3), we assume that $p(L)$ is a gamma distribution. As expected, $I(R_0)$ (Eq. 3) between

the output B and the input L increases monotonically with R_0 (inset in Fig. 2A). Moreover, given that R_0

varies in the population (also assumed to be a gamma distribution), the sensing ability varies in the

population as well (Fig. 2A). Notably, the average I_{Cee} of $I(R_0)$ remains relatively robust to variation in

. At the same time, the traditional estimate , which is the mutual information between the input

and the cell state agnostic population response (response of the

"average" cell, Eq. 1 and Eq. 2), decreases as the population heterogeneity in increases. Importantly,

is significantly lower than the sensing ability of most cells in the population (Fig. 2A). This is because

the overlap in the population response distributions is significantly larger than that in single cell

response distributions (Fig. 2B). This simple example illustrates that the traditional mutual information

estimates may severely underestimate cells' ability to resolve inputs, especially when cell state variables

are heterogeneously distributed. Moreover, it is crucial to explicitly account for heterogeneity in cell

state variables when estimating fidelity of cellular communication channels.

Figure 2. A. The distribution of single cell sensing abilities (horizontal blue histograms) and its average

plotted as a function of the coefficient of variation of the distribution of one cell state variable, the cell surface

receptor number The dashed blue lines show the traditional cell state averaged mutual information (Eq.

1). The inset shows the dependence between cell state specific mutual information and cell state variable

The input distribution is assumed to be a gamma distribution. **B.** A schematic showing the effect of

heterogeneity in cell states on population level response. Even when individual cells have little overlap in their responses to extracellular signal (bottom), the population level responses could have significant overlap (top), leading to a low mutual information between cell state averaged response and the input. **C.** A combined schematic of the two growth factor pathways. Extracellular growth factor ligand (red circle) binds to cell surface receptors which are shuttled to and from the plasma membrane continuously. Ligand bound receptors are phosphorylated and activate Akt. Phosphorylated Akt leads to phosphorylation of FoxO which bars it from entering the nucleus. EGF/EGFR model is limited to the reactions on the plasma membrane. The corresponding cell state variables are given by: $\theta \equiv \{k_{prod}, k_{bind}, k_{unbind}, k_p, k_{dp}, k_{deg}, k_{deg}^*\}$. The cell state variables for the IGF/FoxO model are given by: $\theta \equiv \{k_{prod}, k_{bind}, k_{unbind}, k_p, k_{dp}, k_{deg}, k_{deg}^*, [Akt], k_{ap}, k_{adp}, k_{in}, k_{ef}, k_{fp}, k_{fdp}, [FoxO]\}$. **D.** The estimated distribution of single cell mutual information values $p_{\mathrm{CeeMI}}(I)$ for the EGF/EGFR pathway using a maximum entropy estimation of $p(\theta)$. The inset shows the input distribution $p(u)$ corresponding to the maximum of the average I_{Cee} of $p_{\mathrm{CeeMI}}(I)$ (blue), along with the input distribution corresponding to the channel capacity of I_{CSA} (green). **E.** Same as **D** for the IGF/FoxO pathway. We additionally show the experimentally estimated $p_{\mathrm{CeeMI}}(I)$ (pink).

Experimental verification of $p_{\mathrm{CeeMI}}(I)$ using growth factor signaling networks

In cell populations, state variables that govern signaling dynamics such as protein levels (receptors, kinases, dephosphatases, etc.)[20, 21] as well as effective rate constants such as endocytosis rates[39], ligand binding rates[40], etc. vary from cell to cell. Therefore, we expect the downstream phenotype of environmental sensing to be widely distributed as well. To experimentally verify the computational prediction of the distribution $p_{\mathrm{CeeMI}}(I)$ of sensing abilities, we need a system that allows us to approximate the cell state specific response distribution $p(x|u, \theta)$. The IGF/FoxO pathway[35] is an ideal system for these explorations for several reasons. First, following IGF stimulation, the transcription factor FoxO is pulled out of the nucleus. GFP-tagged variant of FoxO can be used to detect the dynamics of nuclear levels of FoxO at the single cell level[41]. Second, nuclear FoxO levels reach an approximate steady state within 30-45 minutes of constant IGF stimulation with FoxO levels decreasing with increasing IGF dose[41]. As a result, an approximate cell state specific distribution $p(x|u, \theta)$ of steady state levels of FoxO can be obtained by stimulating the same cell with successive IGF levels. Finally, the biochemical interactions in the IGF/FoxO are well studied (Fig. 2C), allowing us to a build stochastic biochemical model based on previous computational efforts[42, 43] that fits the single cell data accurately. Another system where $p_{\mathrm{CeeMI}}(I)$ can in principle be verified is the EGF/EGFR pathway (Fig. 2C). Here too, abundance of cell surface EGFR levels can be tracked in live cells following EGF stimulation[44], allowing us to obtain cell-state specific response distribution $p(x|u, \theta)$. Below, we show estimates of

$p_{\mathrm{CeeMI}}(I)$ for both pathways and an experimental verification of our estimates for the IGF/FoxO

pathway where live-cell imaging data was previously collected[41].

The details of the calculations presented below can be found in SI Sections 3 and 4. Briefly, we first

constructed stochastic biochemical models for the two pathways based on known interactions (Fig. 2C)

and previous models[42, 43]. The output for the EGFR pathway was defined as the steady state levels of cell

surface EGF receptors and the output for the IGF/FoxO pathway was defined as the steady state nuclear

levels of the transcription factor FoxO. Using previously collected single cell data on the two pathways[37,

41, 43] and our maximum entropy-based framework[37], we estimated the distribution over parameters $p(\boldsymbol{\theta})$

for the two pathways (SI Section 3). Using these inferred distributions, and the model-predicted cell

state specific response distribution $p(x|u, \boldsymbol{\theta})$, we could compute $p_{\mathrm{CeeMI}}(I)$ for any specified input

distribution $p(u)$. We choose the support of the input distribution as the ligand concentrations used in

the experimental setup. The estimates shown in Fig. 2D and Fig. 2E show $p_{\mathrm{CeeMI}}(I)$ corresponding to the

input distribution $p(u)$ that maximizes the conditional mutual information I_{Cee} (see Eq. 6). This input

distribution is shown in the inset of Fig. 2D and 2E.

Similar to the illustrative example (Fig. 2A), there is a wide distribution of single cell sensing fidelities in

real populations (Fig. 2D and Fig. 2E). Moreover, most cells are better sensors compared to the "average

cell", a cell whose response $p(x|u)$ is averaged over cell state variability which was estimated to be ~ 1

bit for both pathways. The cellular signaling fidelity skews towards the upper limit of 2 bits which

corresponds to the logarithm of the number of inputs used in the experiment. Indeed, as seen in the

insets of Fig. 2D and Fig. 2E, the input distribution corresponding to the maximum of the cell state

agnostic mutual information I_{CSA} is concentrated on the lowest and the highest input for both

pathways, indicating that cells may be able to detect only two input levels. In contrast, the input

distribution corresponding to the maximum of I_{Cee} is close to uniform, suggesting that individual cells

can in fact resolve different ligand levels.

To verify our computational estimate of $p_{\mathrm{CeeMI}}(I)$ for the IGF/FoxO pathway, we reanalyzed previously

collected data wherein several cells were stimulated using successive IGF levels[41]. The details of our

calculations can be found in SI Section 4. Briefly, the cells reached an approximate steady state within 60

minutes of each stimulation and nuclear FoxO levels measured between 60 to 90 minutes were used to

approximate an experimental cell state specific response distribution $p(x|u, \boldsymbol{\theta})$. The distribution

$p_{\mathrm{CeeMI}}(I)$ was then obtained by maximizing the average mutual information I_{Cee} (averaged over all

cells) with respect to the input distribution. As seen in Fig. 2D and Fig. 2E, the experimentally evaluated
single cell signaling fidelities match closely with our computational estimates. Moreover, as predicted
using our computational analysis, individual cells in the population were significantly better at sensing
than what is implied by the maximum of I_{CSA}. Indeed, the distribution of steady state FoxO levels were
found to be well-resolved at the single cell level as well (Fig. 2E). Live cell imaging data for the EGFR
pathway was not available and we leave it to future studies to validate our predictions.
272
Our calculations show that real cell populations comprise cells that have differing sensing fidelities,
individual cells are significantly better at sensing their environment than what traditional estimates
would indicate, and importantly, the CeeMI approach can accurately estimate the variation in signaling
performances using readily collected experimental data and stochastic biochemical models. Notably, the
variability in cell states and therefore the heterogeneity in sensing abilities is likely to be stable over
time; the same cell's FoxO responses to the same input were found to have significantly less variation
compared to the variation within the population (SI Section 4).
280
281 **CeeMI identifies biochemical parameters that determine cells' ability to sense their**
282 **environment**

283
Figure 3. Dependence on cell state dependent mutual information on biochemical parameters. (left) The joint
distribution $p_{CeeMI}(I, \chi)$ of cell state specific mutual information and biochemical parameter χ chosen to be the
single cell response range of nuclear FoxO levels (x-axis, see inset for a cartoon). The shaded blue regions are
model predictions, and the green line is the model average. The darker shades represent higher probabilities. The
red dots represent experimental cells. The cyan line represents experimental averages. (right) same as (left) with
biochemical parameter χ chosen to be steady state nuclear Foxo levels in the absence of stimulation. The contours
represent 1% to 10%, 10% to 50%, and 50% to 100% of the total probability mass (from faint to dark shading).
291
292 Like other phenotypes[21] and signaling dynamics[20, 30], we expect that cells' ability to sense their
293 environment depends on their state variables. For example, cells with faster endocytosis rates[39] may
294 integrate environmental fluctuations with higher accuracy[43]. Or cells with higher receptor numbers[21] will
295 lead to a lowered relative noise in bound ligand concentration.

To systematically identify the variables that differentiate between cells' ability to sense the environment, we quantify the joint distribution $p_{\text{CeeMI}}(I, \chi)$ of single cell signaling fidelity and biochemical state variables that take part in the signaling network. To test whether we can identify variables that are predictive of cellular fidelity, we estimated the joint distribution $p_{\text{CeeMI}}(I, \chi)$ for two variables that were experimentally accessible, response range of nuclear FoxO (Fig. 3 left, see inset) and total nuclear FoxO levels prior to IGF stimulation (Fig. 3 right, see inset). In both figures, the shaded regions show computational estimates of the joint probability distributions, and the red circles represent real cells. The green and the cyan trend lines represent computational and experimental binned averages.

One may expect that higher total nuclear FoxO levels could result in lower noise (coefficient of variation) and therefore better sensing abilities. However, we find that total nuclear FoxO levels only weakly correlate with cell state dependent mutual information (correlation coefficient $r = 0.16$ for computational estimates and $r = 0.04$ for experimental data). In contrast, cell state dependent mutual information depended strongly on the dynamic range of the response (correlation coefficient $r = 0.53$ for computational estimates and $r = 0.29$ for experimental data). Importantly, the model captured the observation that cells with a small response range had a variable sensing abilities while cells with a large response range all performed well in resolving extracellular IGF levels. Surprisingly, the total nuclear FoxO levels only weakly correlated with the cell specific mutual information. In SI Section 5, we show the predicted joint distributions $p_{\text{CeeMI}}(I, \chi)$ for several other biochemical variables that can potentially govern single cells' response to extracellular IGF stimuli. This way, CeeMI can be used to systematically identify cell state variables that differentiate between good sensors and bad sensors of the environmental stimuli.

Time scale of stochastic dynamics of cell states dictates the divergence between state-specific and state-agnostic sensing abilities

A key limitation of the presented analysis is the assumption that cell state variables remain approximately constant over the time scale of typical environmental fluctuations. While many cell state variables change slowly, remaining roughly constant over the life spans of cells and beyond[25, 26], state changes may occur within the lifespan of a cell as well[45]. These dynamical changes can be

accommodated easily in our calculations. Here, instead of fixing the cell state variables $\boldsymbol{\theta}$, we can treat

them as initial conditions and propagate them stochastically with their own dynamics. In the limit of

very fast dynamics where individual cells rapidly transition cell states, we expect that the individuality of

cells in a population vanishes and consequently cell state specific mutual information for each cell will

agree with traditional cell state agnostic estimates of the channel capacity. In contrast, if the cell state

dynamics are slow, our framework highlights differences between cells in the population.

Figure 4. Cell state dynamics governs cell state conditioned mutual information. A. In a simple stochastic model,
receptor mRNA is produced at a constant rate from the DNA and the translated into ligand-free receptors. The
number of ligand-bound receptors after a short exposure to ligands is considered the output. **B.** A schematic
showing dynamics of receptor numbers when mRNA dynamics are slower compared to signaling time scales. **C.**
Conditioning on receptor numbers leads to differing abilities in sensing the environment when the time scale of
mRNA dynamics is slow. In contrast, when the mRNA dynamics are fast (large τ^{-1}), conditioning on cell state
variables does not lead to difference in sensing abilities.

To elucidate the role of cell state dynamics, we built a simple model of ligand-receptor system (SI

Section 6, Fig. 4). Briefly, the model included production and degradation of the receptor mRNA,

translation of the receptor protein from the mRNA, and ligand binding to the receptor (Fig.4A). The

number of ligand-free receptors and the number mRNA molecules were together considered the state

variables. We tuned the mRNA dynamics by keeping the average mRNA number constant while

simultaneously changing mRNA production and degradation rates. The time scale of mRNA dynamics is

denoted by τ.

When the mRNA dynamics are slower than translation, ligand binding, and receptor degradation,

individual cells are effectively frozen around a particular receptor number (Fig. 4B). The distribution of

the number of ligand-bound receptors after a short-term ligand exposure reflects this frozen state. As a

result, the cell-state conditioned mutual information depends strongly the cell state, with higher mutual

information for cells with higher receptor numbers. In contrast, as the mRNA dynamics become faster,

cells quickly lose the memory of cell state conditioning. As a result, the distribution of the number of

ligand-bound receptors after a short-term ligand exposure reflects an ergodic averaging over the

underlying mRNA dynamics. This averaging effectively washes out differences between cells (Fig. 4C).

This example shows that when cells states change at time scales slower than signaling dynamics, cells in

a population can be identified by their cell state and their ability to sense the environment can differ

from one another.

Discussion

Cell populations are characterized by heterogeneity in cell state variables[18, 19] that is responsible for

important phenotypic differences and selective advantage for example in sensitivity to drugs[25, 26, 28],

response to chemotactic signals[46], and following proliferation cues[29]. Therefore, it is reasonable to

expect that cells' ability to sense their environment depends on their state and is therefore variable

across cells in a population. To quantify this heterogeneity, here, we developed a novel information

theoretic perspective that allowed us to quantify the distribution $p_{\mathrm{CeeMI}}(I)$ of single cell sensing abilities

using easily measurable single cell data and stochastic models of signaling networks. We also quantified

the joint distribution $p_{\mathrm{CeeMI}}(I, \chi)$ of cell specific sensing ability and biochemical cell state variable.

Importantly, using two growth factor pathways, we showed that individual cells in real cell populations

were much better at sensing their environment compared to what is implied by the traditional estimate

of channel capacities of signaling networks. Typical single cell data are time stamped and do not give

information about dynamics in the single cell. Moreover, these data do not provide information about

dependence of cell-to-cell variability on cell state variables. Therefore, computational modeling of

cellular trajectories from single cell data and stochastic models is essential in deciphering mechanistic

origins of heterogeneity in cellular phenotypes.

The approach presented here will be useful in identifying bottlenecks in signal transduction. Many

cellular phenotypes such as chemotaxis[7, 8] and cell proliferation[47] exhibit a weak correlation between

cellular outputs (e.g., directional alignment with chemical gradients) with the input (e.g., gradient

strength), resulting in a low channel capacity even for individual cells. In such cases, it is important to

understand where exactly along the information transduction pathway is the information about the

gradient is lost. If traditional calculations are pursued, for example, for movement of mammalian cells

under growth factor gradients[7, 31], one may conclude that the information loss likely happens right at the

receptor level (Fig. 2D). In contrast, CeeMI will allow us to disentangle the effect of cell state
heterogeneity and noisy cellular response to precisely pinpoint intracellular signaling nodes that are
responsible for signal corruption.

How do we contrast our results with previous low estimates of channel capacities? There is extensive
population heterogeneity in cell state variables and this heterogeneity often is stable over time.
Nonetheless, this heterogeneity arises from stochasticity in underlying processes including dynamics of
epigenetic transitions and gene expression. When one specifies state variables[19] $\boldsymbol{\theta}$, one effectively
conditions on the stochastic trajectory of the underlying dynamics to a specific narrow window (Fig. 4).
This extra conditioning narrows the cells' response distributions, revealing the heterogeneous nature of
cell populations[19].

In summary, we showed that like other phenotypes, the ability to sense the environment is itself
heterogeneously distributed in a cell population. Moreover, we also showed that when conditioned on
cell state variables, mammalian cells appear to be significantly better at sensing their environment than
what traditional mutual information calculation suggests. Finally, we showed that we could identify cell
state variables that made some cells better sensors compared to others. We believe that CeeMI will be
an important framework in quantifying fidelity of input/output relationships in heterogeneous cell
populations.

Acknowledgments

AG, HA, and PD are supported by NIGMS grant R35GM142547. The authors would like to thank Andre
Levchenko for useful discussions.

430 **References**
431

1. Levchenko A, Nemenman I. Cellular noise and information transmission. Curr Opin
Biotechnol. 2014;28:156-64.
2. Rhee A, Cheong R, Levchenko A. The application of information theory to biochemical
signaling systems. Phys Biol. 2012;9(4):045011.
3. ten Wolde PR, Becker NB, Ouldridge TE, Mugler A. Fundamental Limits to Cellular
Sensing. J Stat Phys. 2016;162(5):1395-424.
4. Cheong R, Rhee A, Wang CJ, Nemenman I, Levchenko A. Information transduction
capacity of noisy biochemical signaling networks. Science. 2011;334(6054):354-8.
5. Selimkhanov J, Taylor B, Yao J, Pilko A, Albeck J, Hoffmann A, Tsimring L, Wollman R.
Accurate information transmission through dynamic biochemical signaling networks. Science.
2014;346(6215):1370-3.
6. Suderman R, Bachman JA, Smith A, Sorger PK, Deeds EJ. Fundamental trade-offs
between information flow in single cells and cellular populations. P Natl Acad Sci USA.
2017;114(22):5755-60.
7. Varennes J, Moon HR, Saha S, Mugler A, Han B. Physical constraints on accuracy and
persistence during breast cancer cell chemotaxis. Plos Comput Biol. 2019;15(4). PubMed PMID:
WOS:000467530600065.
8. Mattingly HH, Kamino K, Machta BB, Emonet T. Escherichia coli chemotaxis is
information limited. Nat Phys. 2021;17(12):1426-+.
9. Cover TM, Thomas JA. Elements of information theory. 2nd ed. Hoboken, N.J.: Wiley-
Interscience; 2006. xxiii, 748 p. p.
10. Wu M, Singh AK. Single-cell protein analysis. Curr Opin Biotechnol. 2012;23(1):83-8.
11. Slavov N. Single-cell protein analysis by mass spectrometry. Curr Opin Chem Biol.
2021;60:1-9.
12. Lemon WC, McDole K. Live-cell imaging in the era of too many microscopes. Curr Opin
Cell Biol. 2020;66:34-42.
13. Hansen AS, O'Shea EK. Limits on information transduction through amplitude and
frequency regulation of transcription factor activity. Elife. 2015;4.
14. Billing U, Jetka T, Nortmann L, Wundrack N, Komorowski M, Waldherr S, Schaper F,
Dittrich A. Robustness and Information Transfer within IL-6-induced JAK/STAT Signalling.
Commun Biol. 2019;2:27.
15. Garner KL, Perrett RM, Voliotis M, Bowsher C, Pope GR, Pham T, Caunt CJ, Tsaneva-
Atanasova K, McArdle CA. Information Transfer in Gonadotropin-releasing Hormone (GnRH)
Signaling: EXTRACELLULAR SIGNAL-REGULATED KINASE (ERK)-MEDIATED FEEDBACK LOOPS
CONTROL HORMONE SENSING. J Biol Chem. 2016;291(5):2246-59.

16. Frick CL, Yarka C, Nunns H, Goentoro L. Sensing relative signal in the Tgf-beta/Smad
pathway. Proc Natl Acad Sci U S A. 2017;114(14):E2975-E82.
17. Lee JB, Caywood LM, Lo JY, Levering N, Keung AJ. Mapping the dynamic transfer
functions of eukaryotic gene regulation. Cell Syst. 2021;12(11):1079-93 e6.
18. Trapnell C. Defining cell types and states with single-cell genomics. Genome Res.
2015;25(10):1491-8.
19. Wollman R. Robustness, Accuracy, and Cell State Heterogeneity in Biological Systems.
Curr Opin Syst Biol. 2018;8:46-50.
20. Meyer R, D'Alessandro LA, Kar S, Kramer B, She B, Kaschek D, Hahn B, Wrangborg D,
Karlsson J, Kvarnstrom M, Jirstrand M, Lehmann WD, Timmer J, Hofer T, Klingmuller U.
Heterogeneous kinetics of AKT signaling in individual cells are accounted for by variable protein
concentration. Front Physiol. 2012;3:451.
21. Frankel NW, Pontius W, Dufour YS, Long J, Hernandez-Nunez L, Emonet T. Adaptability
of non-genetic diversity in bacterial chemotaxis. Elife. 2014;3.
22. Chen KH, Boettiger AN, Moffitt JR, Wang S, Zhuang X. RNA imaging. Spatially resolved,
highly multiplexed RNA profiling in single cells. Science. 2015;348(6233):aaa6090. Epub
20150409.
23. Nagano T, Lubling Y, Stevens TJ, Schoenfelder S, Yaffe E, Dean W, Laue ED, Tanay A,
Fraser P. Single-cell Hi-C reveals cell-to-cell variability in chromosome structure. Nature.
2013;502(7469):59-64.
24. Buenrostro JD, Wu B, Litzenburger UM, Ruff D, Gonzales ML, Snyder MP, Chang HY,
Greenleaf WJ. Single-cell chromatin accessibility reveals principles of regulatory variation.
Nature. 2015;523(7561):486-90. PubMed PMID: 26083756.
25. Flusberg DA, Roux J, Spencer SL, Sorger PK. Cells surviving fractional killing by TRAIL
exhibit transient but sustainable resistance and inflammatory phenotypes. Mol Biol Cell.
2013;24(14):2186-200.
26. Spencer SL, Gaudet S, Albeck JG, Burke JM, Sorger PK. Non-genetic origins of cell-to-cell
variability in TRAIL-induced apoptosis. Nature. 2009;459(7245):428-32.
27. Pleska M, Jordan D, Frentz Z, Xue B, Leibler S. Nongenetic individuality, changeability,
and inheritance in bacterial behavior. Proc Natl Acad Sci U S A. 2021;118(13).
28. Emert BL, Cote CJ, Torre EA, Dardani IP, Jiang CL, Jain N, Shaffer SM, Raj A. Variability
within rare cell states enables multiple paths toward drug resistance. Nat Biotechnol.
2021;39(7):865-76.
29. Matson JP, Cook JG. Cell cycle proliferation decisions: the impact of single cell analyses.
FEBS J. 2017;284(3):362-75.
30. Norris D, Yang P, Shin SY, Kearney AL, Kim HJ, Geddes T, Senior AM, Fazakerley DJ,
Nguyen LK, James DE, Burchfield JG. Signaling Heterogeneity is Defined by Pathway
Architecture and Intercellular Variability in Protein Expression. Iscience. 2021;24(2):102118.
31. Moon HR, Saha S, Mugler A, Han B. Signal processing capacity of the cellular sensory
machinery regulates the accuracy of chemotaxis under complex cues. Iscience. 2021;24(11).
32. Waite AJ, Frankel NW, Dufour YS, Johnston JF, Long J, Emonet T. Non-genetic diversity
modulates population performance. Mol Syst Biol. 2016;12(12):895.
33. Clark JA, Campbell SG. Diverse relaxation rates exist among rat cardiomyocytes isolated
from a single myocardial region. J Physiol. 2019;597(3):711-22.

34.   Yao J, Pilko A, Wollman R. Distinct cellular states determine calcium signaling response.
Mol Syst Biol. 2016;12(12):894.
35.   Lewis Cantley TH, Richard Sever, Jeremy Thorner Signal Transduction: Principles,
Pathways, and Processes 1st ed: Cold Spring Harbor Laboratory Press; 2013.
36.   Feinberg AP, Levchenko A. Epigenetics as a mediator of plasticity in cancer. Science.
2023;379(6632):eaaw3835. Epub 20230210. doi: 10.1126/science.aaw3835. PubMed PMID:
36758093; PMCID: PMC10249049.
37.   Dixit PD, Lyashenko E, Niepel M, Vitkup D. Maximum Entropy Framework for Predictive
Inference of Cell Population Heterogeneity and Responses in Signaling Networks. Cell Syst.
2020;10(2):204-12 e8.
38.   Loos C H, J. Mathematical modeling of variability in intracellular signaling. Current
Opinion in Systems Biology. 2019;16:17-24.
39.   Kallenberger SM, Unger AL, Legewie S, Lymperopoulos K, Klingmuller U, Eils R, Herten
DP. Correlated receptor transport processes buffer single-cell heterogeneity. Plos Comput Biol.
2017;13(9):e1005779.
40.   Chung JC, Sciaky N, Gross DJ. Heterogeneity of epidermal growth factor binding kinetics
on individual cells. Biophys J. 1997;73(2):1089-102.
41.   Gross SM, Dane MA, Bucher E, Heiser LM. Individual Cells Can Resolve Variations in
Stimulus Intensity along the IGF-PI3K-AKT Signaling Axis. Cell Syst. 2019;9(6):580-8 e4.
42.   Wimmer RJ, Liu Y, Schachter TN, Stonko DP, Peercy BE, Schneider MF. Mathematical
modeling reveals modulation of both nuclear influx and efflux of Foxo1 by the IGF-I/PI3K/Akt
pathway in skeletal muscle fibers. Am J Physiol Cell Physiol. 2014;306(6):C570-84.
43.   Lyashenko E, Niepel M, Dixit PD, Lim SK, Sorger PK, Vitkup D. Receptor-based
mechanism of relative sensing and cell memory in mammalian signaling networks. Elife. 2020;9.
44.   Fortian A, Sorkin A. Live-cell fluorescence imaging reveals high stoichiometry of Grb2
binding to the EGF receptor sustained during endocytosis. J Cell Sci. 2014;127(Pt 2):432-44.
45.   MacLean AL, Hong T, Nie Q. Exploring intermediate cell states through the lens of single
cells. Curr Opin Syst Biol. 2018;9:32-41.
46.   Mattingly HH, Emonet T. Collective behavior and nongenetic inheritance allow bacterial
populations to adapt to changing environments. Proc Natl Acad Sci U S A.
2022;119(26):e2117377119.
47.   Gross SM, Rotwein P. Unraveling Growth Factor Signaling and Cell Cycle Progression in
Individual Fibroblasts. J Biol Chem. 2016;291(28):14628-38.
544