Original Title: β-carotene accelerates the resolution of atherosclerosis in mice

Authors: Ivan Pinos[1]*, Johana Coronel[2]*, Asma'a Albakri[1]*, Amparo Blanco[1], Patrick
McQueen[1], Donald Molina[2], JaeYoung Sim[2], Edward A Fisher[3], Jaume Amengual[1,2]

Affiliations:

1: Division of Nutritional Sciences, University of Illinois Urbana Champaign, Urbana, IL
2: Department of Food Science and Human Nutrition, University of Illinois Urbana Champaign,
Urbana, IL
3: The Leon H. Charney Division of Cardiology, Department of Medicine, The Marc and Ruti
Bell Program in Vascular Biology, New York University Grossman School of Medicine, NYU
Langone Medical Center, NY

* These authors contributed equally to the work.

Corresponding author: jaume6@illinois.edu (J. Amengual)

Funding: This work was supported by the National Institutes of Health (R01HL147252 to JA)
and the United States Department of Agriculture (W4002 to JA).

ABSTRACT

β-carotene oxygenase 1 (BCO1) catalyzes the cleavage of β-carotene to form vitamin A. Besides

its role in vision, vitamin A regulates the expression of genes involved in lipid metabolism and

immune cell differentiation. BCO1 activity is associated with the reduction of plasma cholesterol

in humans and mice, while dietary β-carotene reduces hepatic lipid secretion and delays

atherosclerosis progression in various experimental models. Here we show that β-carotene also

accelerates atherosclerosis resolution in two independent murine models, independently of

changes in body weight gain or plasma lipid profile. Experiments in $Bco1^{-/-}$ mice implicate

vitamin A production in the effects of β-carotene on atherosclerosis resolution. To explore the

direct implication of dietary β-carotene on regulatory T cells (Tregs) differentiation, we utilized

anti-CD25 monoclonal antibody infusions. Our data show that β-carotene favors Treg expansion

in the plaque, and that the partial inhibition of Tregs mitigates the effect of β-carotene on

atherosclerosis resolution. Our data highlight the potential of β-carotene and BCO1 activity in

the resolution of atherosclerotic cardiovascular disease.

Keywords: Retinoic acid, forkhead box P3 (FoxP3), inflammation

1. INTRODUCTION

Atherosclerotic cardiovascular disease is a progressive pathological process initiated by the accumulation of cholesterol-rich lipoproteins within the intima layer of the arterial wall. These particles ultimately lead to the production of chemoattractant cues for circulating monocytes that transmigrate across the endothelial layer to reach the intima and then differentiate to macrophages to become cholesterol-laden foam cells. During lesion development, macrophages promote local inflammation and plaque weakening by degrading extracellular matrix components such as collagen fibers, which can evolve in the rupture of the lesion and thrombus formation [1].

Conventional therapies aim to lower plasma cholesterol to mitigate the progression of atherosclerosis. Novel strategies are currently under development to stimulate the resolution of plaque inflammation more directly, and eventually a reduction in lesion size, in a process named atherosclerosis regression [2]. Among these strategies, the modulation of CD4$^+$ regulatory T cells (Tregs) number is gaining interest over the past years since the discovery that regressing lesions are enriched in Tregs, where they promote plaque stabilization and repair [3, 4]. Tregs typically express the surface marker CD25, which has been utilized to target and eliminate Tregs [5, 6]. However, strategies to deplete CD25$^+$ Tregs do not affect CD25$^-$ Tregs, which possess similar immunomodulatory properties as CD25$^+$ Tregs [7]. Among the different Tregs markers, the forkhead box P3 (FoxP3) acts as lineage specification factor regulating gene expression of proteins implicated in the immunosuppressive activity of these cells [8]. Cell culture studies show that retinoic acid, the transcriptionally active form of vitamin A, promotes Treg differentiation by upregulating FoxP3 expression [9], however, whether dietary vitamin A affects Tregs during atherosclerosis development and resolution remains unanswered.

Humans obtain vitamin A primarily from β-carotene and other provitamin A carotenoids present

in most fruits and vegetables. Upon absorption, provitamin A carotenoids are cleaved by the

action of β-carotene oxygenase 1 (BCO1), the limiting enzyme in vitamin A formation [10]. Our

data show that the enzymatic activity of BCO1 mediates the bioactive actions of β-carotene in

various preclinical models of obesity and atherosclerosis [11-16]. We showed that the dietary

supplementation with β-carotene delays atherosclerosis progression by reducing cholesterol

hepatic secretion in low-density lipoprotein receptor (LDLR)-deficient ($Ldlr^{-/-}$) mice [15]. We

also reported that subjects harboring a genetic variant linked to greater BCO1 activity was

associated with a reduction in plasma cholesterol [17].

In this study, we tested the effect of dietary β-carotene on atherosclerosis resolution in two

independent experimental models. We characterized plaque composition by probing for

macrophage and collagen contents, two parameters utilized to characterize atherosclerotic lesions

undergoing resolution [18-21]. Lastly, we utilized $Bco1^{-/-}$ mice and $CD25^{+}$ Treg depletion

experiments in $Foxp3^{EGFP}$ mice to tease out the direct implications of vitamin A formation and

Tregs on atherosclerosis resolution.

2. METHODS

2.1. Animal husbandry and diets

All procedures were approved by the Institutional Animal Care and Use Committees of the

University of Illinois at Urbana Champaign. For all our studies, we utilized comparable number

of male and female wild-type, $Ldlr^{-/-}$ (#002207, Jackson Labs, Bar Harbor, ME), $Bco1^{-/-}$ [11], and

$Foxp3^{EGFP}$ mice (#006772, Jackson Labs). All mice were in C57BL/6J background. Mice were

kept under controlled temperature and humidity conditions with a 12-hours light/dark cycle and

free access to food and water. Mice were weaned at three weeks of age onto a breeder diet

(Teklad global 18% protein diet: Envigo, Indianapolis, IN) in groups of three to four mice.

Control and β-carotene diets contained either placebo beadlets or β-carotene beadlets at a final

concentration of 50 mg β-carotene/kg diet. For reference, the content of β-carotene in carrots is

80 mg/kg (USDA Database). Beadlets were a generous gift from DSM Nutritional Products

(Sisseln, Switzerland). All diets were prepared by Research Diets (New Brunswick, NJ) by cold

extrusion. The exact composition of all the diets used for this study are provided in

Supplementary Table 1. For all the experiments, we stimulated the development of

atherosclerosis with a Western diet deficient in vitamin A (WD-VAD), as done in the past [15,

17].

2.2. Blood sampling and tissue collection

Before tissue harvesting, mice were deeply anesthetized by intraperitoneal injection with a

mixture of ketamine and xylazine at 80 and 8 mg/kg body weight, respectively. We collected

blood by cardiac puncture using ethylenediaminetetraacetic acid (EDTA)-coated syringes. We

then perfused the mice with 10% sucrose in 0.9% sodium chloride (NaCl)-saline solution prior to

tissue harvesting. Tissues were immediately snap-frozen in liquid nitrogen and kept at -80 ℃.

Aortic roots were collected after removing fat under a binocular microscope, embedded in

optimum cutting temperature compound (OCT, Tissue-Tek, Sakura, Torrance, CA) and kept at -

80°C.

2.3. Antisense oligonucleotide (ASO) targeting LDLR expression (ASO-LDLR)

To transiently deplete LDLR expression in wild-type, $Bco1^{-/-}$, and $Foxp3^{EGFP}$ mice, we

administered weekly an intraperitoneal injection containing 5 mg/kg body weight of ASO-LDLR

for a period of 16 weeks. At the end of the treatment, we injected a single dose of 20 mg/kg of

the sense oligonucleotide (SO-LDLR) antidote, which binds and deactivates the ASO-LDLR

[22]. All oligonucleotide treatments were generously provided by Ionis Pharmaceuticals

(Carlsbad, CA).

2.4. HPLC analyses of carotenoids and retinoids

Nonpolar compounds were extracted from 100 μl of plasma or 30 mg of liver under a dim yellow

safety light using methanol, acetone, and hexanes. The extracted organic layers were then pooled

and dried in a SpeedVac (Eppendorf, Hamburg, Germany). We performed the HPLC with a

normal phase Zobax Sil (5 μm, 4.6 × 150 mm) column (Agilent, Santa Clara, CA). Isocratic

chromatographic separation was achieved with 10% ethyl acetate/hexane at a flow rate of 1.4

ml/min. For molar quantification of β-carotene and retinoids, we scaled the HPLC with a

standard curve using the parent compound.

2.5. Plasma lipid analyses

We measured plasma total cholesterol, cholesterol in the high-density lipoprotein fraction (HDL-

C), and triglyceride levels using commercially available kits (FUJIFILM Wako Diagnostic,

Mountain View, CA), according to the manufacturer's instructions.

Plasma lipoproteins were fractionated using fast performance liquid chromatography (FPLC)

with two Superose 6 10/300 GL columns (G.E. Healthcare, Boston, MA) on a Shimadzu HPLC

system (Columbia, MD), as done in the past [15, 18]. Pooled samples (five mice/group) were

used for analysis, and lipoprotein fractions were identified by quantifying cholesterol and

triglyceride levels in each fraction.

2.6. RNA isolation and RT-PCR

We isolated the total RNA using TRIzol reagent (Thermo Fisher Scientific, Waltham, MA) and

Direct-zol RNA Miniprep kit (Zymo Research, Irvine, CA) according to the manufacturer's

instructions. RNA was reverse-transcribed to cDNA using high-capacity cDNA reverse

transcription kit (Applied Biosystems, Foster City, CA). Then we used Taqman Master Mix

(Applied Biosystems) to perform RT-PCR using the StepOnePlus RT-PCR system (ABI 7700,

Applied Biosystems). We calculated the relative gene expression using the ΔCt method and

normalized the data to β-actin (*Actb)*. Probes (Applied Biosystems) include mouse *Ldlr*

(Mm00440169_m1), mouse cytochrome P450 26a1 (*Cyp26a1,* Mm00514486_m1) and mouse

Actb (Mm02619580_g1).

2.7. Western blot analysis

For the determination of hepatic levels of LDLR, proteins were extracted from liver lysates in

RIPA buffer (50 mM Tris pH 7.4, 150 mM NaCl, 0.25% sodium deoxycholate, 1% Nonidet P-

40) in the presence of protease inhibitors. Total protein amounts were quantified using the

Pierce® BCA Protein Assay Kit (ThermoFisher Scientific, Waltham, MA). A total of 80 µg of

protein homogenate were separated by SDS-PAGE and transferred onto PVDF membranes (Bio-

Rad, Hercules, CA). Membranes were blocked with fat-free milk powder (5% w/v) dissolved in

Tris-buffered saline (15 mM NaCl and 10 mM Tris/HCl, pH 7.5) containing 0.01% Tween 100

(TBS-T), washed, and incubated overnight at 4 °C with mouse anti-LDLR (Santa Cruz

Biotechnologies, Dallas, TX) and mouse anti-GAPDH (ThermoFisher Scientific) as a

housekeeping control. Infrared fluorescent-labeled secondary antibodies were prepared at

1:15,000 dilution in TBS-T with 5% fat-free milk powder and incubated for 1 h at room

temperature.

2.8. Treg depletion in $Foxp3^{EGFP}$ mice

To deplete Tregs, mice were injected twice with 250 µg of either the isotype control IgG (Ultra-

LEAFPurified Rat IgG1, λ Isotype Ctrl Antibody no. 401916, Biolegend, CA) or PC61 anti-

CD25 monoclonal antibody (Ultra-LEAF Purified anti-mouse CD25 Antibody no. 102040,

Biolegend, CA). The first treatment took place a day after the administration of SO-LDLR, and

the second injection two weeks after, following established protocols [6].

2.9. Monocyte/macrophage trafficking studies

One week before harvesting the baseline group, we injected the mice intraperitoneally with 4

mg/ml of 5-ethynyl-2′-deoxyuridine (EdU) (Invitrogen, Waltham, MA) with the goal of labeling

circulating monocytes to assess macrophage retention among groups. To compare monocyte

recruitment among groups, we labeled circulating monocytes by injecting the mice retro-orbitally

with 1 µm diameter Flouresbrite flash red plain microspheres beads (Polysciences Inc,

Warrington, PA) diluted in sterile PBS 24 hours before harvesting, regardless of the

experimental group. We assessed the efficiency of EdU and bead labeling by flow cytometry 48

hours and 24 hours after injection by tail bleeding, respectively [23].

2.9. Flow cytometry

To assess the number of monocytes labeled with either EdU or fluorescent beads, we incubated

blood samples with red blood cell lysis buffer (Thermo Fisher Scientific, Waltham, MA) and

blocked unspecific bindings with using anti-mouse CD16/CD32 (Mouse BD Fc Block, BD

Biosciences, Franklin Lakes, NJ). Cells were then stained with FITC-conjugated anti-mouse

CD45 (BioLegend), PE-conjugated anti-mouse CD115 (BioLegend), and PerCP-conjugated anti-

mouse Ly6C/G (BioLegend, San Diego, CA) for 30 minutes on ice. Cells were fixed,

permeabilized, and stained for EdU using Click-iT EdU Pacific Blue Flow Cytometry Assay Kit

(Invitrogen) following manufacturer's instructions. Beads were detected with the 640-nm red

laser using a 660/20 bandpass filter. Monocyte recruitment and egress was estimated based on

monocyte labeling efficiency, following established protocols [24].

For Treg quantifications, spleens were mashed and filtered through a 100 µm sterile cell strainer

(Thermo Fisher Scientific). Blood and spleen homogenates were incubated with red blood cell

lysis buffer (Thermo Fisher Scientific) and subsequently blocked with anti-mouse CD16/CD32

(BD Biosciences). We stained the cells with Fixable Viability Dye eFluor 780 (eBioscience, San

Diego, CA) and eFluor 506-conjugated anti-mouse CD45 (BioLegend), PE-conjugated anti-

mouse CD3e (BioLegend), FITC-conjugated anti-mouse CD4 (BioLegend), and BV421-

conjugated anti-mouse CD25 (BioLegend) for 30 minutes on ice. We washed the cells twice

before fixing and permeabilizing cell membranes using the eBioscience FoxP3/Transcription

Factor Staining Buffer Set (Invitrogen) for 1 hour at room temperature, followed by incubation

with Alexa Fluor 647-conjugated anti-mouse FoxP3 antibody (BioLegend) for 30 minutes at

room temperature. All samples were measured on a BD LSR II analyzer (BD Biosciences) and

results were analyzed with the FCS express 5 software (De Novo Software, Pasadena, CA).

2.10. Atherosclerotic lesion analysis

Six μm-thick sections were fixed and permeabilized with ice-cold acetone, blocked, and stained

with rat anti-mouse CD68 primary antibody (clone FA-11; Bio-Rad, Hercules, CA) followed by

biotinylated rabbit anti-rat IgG secondary antibody (Vector Laboratories, Burlingame, CA).

CD68$^+$ area was visualized using a Vectastain ABC kit (Vector Laboratories). Sections were then

counterstained with hematoxylin, dehydrated in an ethanol gradient, xylene, and mounted with

Permount medium (Thermo Fisher Scientific). Images were acquired using Axioskop 40

microscope (Carl Zeiss, Jena, Germany). For collagen content, frozen sections were fixed and

stained using picrosirius red (Polysciences). Sections were scanned using the Axioscan.Z1

microscope (Carl Zeiss) using both bright field and polarized light. Total lesion area, CD68$^+$, and

collagen-positive areas were quantified using ImageJ software (NIH).

Aortic root sections were fixed in 10% formalin, permeabilized with 1% Triton X-100 in

phosphate buffer, and stained using anti-FoxP3 (clone MF-14; BioLegend), anti-GFP (cat

#ab290; Abcam), anti-CD25 (clone IL2R.1; Thermo Fisher Scientific), anti-arginase 1 (cat

#16001-1-AP; Thermo Fisher Scientific), anti-Cyp26b1b (cat #21555-1-AP; Thermo Fisher

Scientific), and anti-Ki67 (clone SP6; Abcam). EdU+ cells in the lesion were visualized using

Click-IT EdU Imaging Kit (Invitrogen). Secondary antibodies conjugated to Alexa fluorophores

were purchased from Molecular Probes (Life Technologies). Images were taken using a Zeiss

Axioscan.Z1 slide scanner (Carl Zeiss). Quantifications were performed using the ImageJ

software (NIH).

2.11. Statistical Analysis

Data represented as bar charts are expressed as mean ± standard error of the mean (SEM). Data

were analyzed using GraphPad Prism software (GraphPad Software Inc., San Diego, CA) by

one-way ANOVA, followed by Tukey's multiple comparisons test. Differences between groups

were considered significant with an adjusted P value < 0.05. The relationship between the two

dependent factors, CD68 and collagen contents, and group differences were evaluated by

descriptive discriminant analysis. Briefly, models were fitted under null and alternative

hypotheses and goodness of fit was evaluated using the Likelihood Ratio Test with randomized

inference (bootstrap; n = 1000 simulations). Simultaneous hypothesis testing was corrected using

the Benjamini-Hochberg procedure [24]. The resulting False Discovery Rate (FDR) was

considered significant with a cut-off < 0.05.

3. Results

3.1. β-carotene supplementation accelerates atherosclerosis resolution

We recently showed that dietary β-carotene delays atherosclerosis progression in *Ldlr*[-/-] mice [15], which prompted us to examine whether β-carotene also impacts the resolution of inflammation in complex atherosclerotic lesions. To achieve these lesions, we utilized two distinct mouse models of atherosclerosis in combination with WD-VAD. In our first model, we established LDLR deficiency in wild-type mice by injecting ASO-LDLR weekly for a period of 16 weeks. The characteristics of the plaques before undergoing resolution were established by sacrificing a subset of mice after 16 weeks on diet (Baseline). The remaining mice were injected once with SO-LDLR to promote atherosclerosis resolution and divided into two groups fed either WD-VAD (Resolution - Control) or WD-β-carotene (Resolution - β-carotene). Mice were harvested three weeks after SO-LDLR injections (Figure 1A).

Male mice gained more weight than female mice, although we did not observe differences between the three experimental groups (Supplementary Figure 1A). For the remaining outcomes (plasma lipid profile, retinoid levels, and plaque composition), we did not observe sex differences in any of our experiments, and therefore, we combined results from both sexes. Both Resolution groups, independently of β-carotene, showed drastic reduction in total plasma cholesterol and triglyceride levels in comparison to the Baseline group. HDL-C levels remained constant between groups (Figure 1B). Using pooled samples, we compared plasma cholesterol and triglyceride partitioning between groups by FPLC. Very low-density lipoprotein (VLDL) and LDL fractions in both Resolution groups showed a comparable reduction in total cholesterol and triglyceride content in comparison to Baseline mice (Figure 1C, D). Lipid normalization was accompanied by the upregulation of hepatic LDLR expression at the mRNA and protein levels (Figure 1E, F).

Despite mice being fed a WD-VAD for several weeks, HPLC measurements of circulating

vitamin A and hepatic retinoid stores ruled out vitamin A deficiency in any of the experimental

groups (Figure 1G, H). However, mice fed WD-β-carotene showed an increase in hepatic

vitamin A content in comparison to those fed WD-VAD, which was mediated by the conversion

of β-carotene to vitamin A (Figure 1H). Hepatic *Cyp26a1* expression, which is commonly

utilized as a surrogate marker of vitamin A and retinoic acid production [25, 26], appeared

upregulated in response to WD-β-carotene (Supplementary Figure 1B).

To determine the effect of β-carotene supplementation on atherosclerosis resolution, we

examined lesion size and composition at the level of the aortic root (Figure 1I). Lesion area

remained constant among the three experimental groups, as previously reported in mice treated

with ASO-LDLR under similar experimental conditions (Figure 1J) [27]. We next examined

plaque composition by focusing on two parameters commonly utilized as a surrogate indicators

of atherosclerosis resolution: CD68$^+$ area, a myeloid (macrophage) marker that represents plaque

inflammation, and collagen area, a marker of plaque stability in humans [28, 29]. In comparison

to Baseline mice, the lesions of Resolution – Control and Resolution - β-carotene groups

presented a reduction in CD68 content of approximately 36% and 45%, respectively (Figure 1K).

The collagen content in lesions increased in both Resolution groups in comparison to Baseline,

reaching statistical significance only in the β-carotene-fed mice. In this group, we observed

215% and 60% increase in collagen content compared to the Baseline and the Resolution -

Control groups, respectively (Figure 1L). Lastly, we plotted the relative CD68 and collagen

contents in the lesion to perform a descriptive discriminant analysis. Individual samples from

three distinct experimental groups clustered together, highlighting significant differences

between the three experimental groups for all the comparisons (Figure 1M).

To validate the results obtained in our ASO-LDLR reversible model of atherosclerosis, we

utilized *Ldlr*$^{-/-}$ mice fed WD-VAD subjected to a dietary switch strategy to lower cholesterol in

plasma and promote atherosclerosis resolution [18, 30]. Baseline *Ldlr*$^{-/-}$ mice were harvested after

12 weeks on WD-VAD, while the remaining animals were switched to either a Standard diet

without vitamin A (Resolution - Control) or the same diet supplemented with β-carotene

(Resolution - β-carotene) for four more weeks. To prevent changes in food intake due to

consistency and hardness of the feed, we provided Standard diets as powder. This approach

prevented a reduction in body weight, which could have resulted in confounding alterations in

atherosclerosis resolution (Supplementary Figure 2A).

Ldlr$^{-/-}$ mice in both Resolution groups presented a comparable reduction in plasma cholesterol

and triglyceride levels in comparison to the Baseline group. These changes were not

accompanied by alterations in HDL-C levels between groups (Figure 2A). Systemic and hepatic

vitamin A levels failed to show indications of vitamin A deficiency, and hepatic vitamin A stores

increased in β-carotene-fed mice (Supplementary Figure 2B).

We next characterized atherosclerotic lesions at the level of the aortic root (Figure 2B). Lesion

size area was comparable between experimental groups (Figure 2C), although CD68 content

decreased to the same extent in both Resolution groups in comparison to Baseline mice (Figure

2D). In comparison to the Baseline group, collagen accumulation in the lesion increased 100% in

the Resolution - Control and over 200% in the Resolution - β-carotene group, respectively

(Figure 2E). Descriptive discriminant analysis showed comparable results to those observed for

our ASO-LDLR model, where differences between the three experimental groups reached

statistical significance (Figure 2F).

Together, these data show that β-carotene supplementation during atherosclerosis resolution

results in the improvement of lesion composition in two independent mouse models.

3.2. BCO1 drives the effect of β-carotene on atherosclerosis resolution

To establish the contribution of BCO1 on the effect of β-carotene on atherosclerosis resolution,

we utilized LDLR-ASO to promote atherogenesis in $Bco1^{-/-}$ mice following the same

experimental approach described for wild-type mice (Figure 1A). Consistent with our results in

our two Resolution models (Figure 1B and 2A), plasma cholesterol levels decreased in both

Resolution groups in comparison to Baseline mice (Supplementary Figure 3A). Similarly, HPLC

measurements of plasma and hepatic vitamin A ruled out vitamin A deficiency in $Bco1^{-/-}$ mice

Baseline and Resolution mice (Supplementary Figure 3B, C). The accumulation of β-carotene in

tissues and plasma is characteristic in $Bco1^{-/-}$ mice [11]. Indeed, HPLC quantification of β-

carotene in plasma and liver of $Bco1^{-/-}$ Resolution - β-carotene mice presented four-fold and 400-

fold greater β-carotene levels found in wild-type Resolution - β-carotene, respectively (Figure

3A, B).

The characterization of the atherosclerotic lesions showed no alterations in lesion size between

the three experimental groups (Figure 3C, D). When compared to Baseline mice, both Resolution

groups displayed lower CD68 and higher collagen contents, although we didn't observe

differences in CD68 and collagen contents between both Resolution groups (Figure 3E, F).

Descriptive discriminant analysis highlighted differences between Baseline $Bco1^{-/-}$ mice and their

Resolution littermates, independently of the presence of β-carotene in the diet (Figure 3G).

3.3. Effect of β-carotene and anti-CD25 depletion on Treg cell number

Our data show that β-carotene favors atherosclerosis resolution in two independent mouse

models (Figure 1M, 2F), and results in $Bco1^{-/-}$ mice directly implicate vitamin A formation in this

process (Figure 3G). Retinoic acid is the transcriptionally active form of vitamin A and promotes

Treg differentiation by upregulating FoxP3 expression in various experimental models [31-36].

Treg number decreases in developing lesions and increases during atherosclerosis resolution [6,

37, 38], but whether the dietary manipulation of β-carotene or vitamin A affects Treg number

during atherosclerosis remains unanswered. To investigate whether the Tregs are responsible for

the effect of dietary β-carotene on atherosclerosis resolution, we induced atherosclerosis in

$Foxp3^{EGFP}$ mice by injecting ASO-LDLR as described above (Figure 1 and 3). $Foxp3^{EGFP}$ mice

co-express EGFP and FoxP3 under the regulation of the endogenous FoxP3 promoter [39]. After

16 weeks on diet, mice undergoing resolution were treated twice with either the PC61 anti-CD25

monoclonal antibody (anti-CD25) or an isotype control (IgG) (Figure 4A) [6]. We did not

observe changes in body weight among groups, and the three groups undergoing resolution

showed a normalization in plasma lipids in comparison to Baseline controls (data not shown).

Flow cytometry analyses demonstrated the effectiveness of the anti-CD25 treatment by reducing

the ratio of circulating and splenic $CD25^{+}FoxP3^{+}$ ($CD25^{+}$ Tregs) in comparison to the other

experimental groups (Figure 4B-D). We also observed a depletion of $CD25^{+}$ Tregs and

$CD25^{+}FoxP3^{-}$ T cells in the lesion and lymph nodes (Figure 4E, F), in agreement with previous

reports showing that anti-CD25 fails to completely deplete $CD25^{-}FoxP3^{+}$ Tregs ($CD25^{-}$ Tregs),

which retain strong anti-inflammatory properties [7]. Hence, we quantified the number of total

Tregs independently of the presence of CD25 by counting the number of GFP/FoxP3/DAPI triple

positive cells (Figure 4G). All the groups undergoing Resolution presented a greater total Treg

number than Baseline mice, although the results did not reach statistical significance between

Baseline and Resolution - Control group (P = 0.10). Resolution - β-carotene injected with IgG

presented the highest Treg cell number among all the other experimental groups, suggesting that

dietary β-carotene favors Treg expansion in the plaque (Figure 4H).

Circulating total Treg number remained constant, but we observed a slight decrease in the

number of splenic total Tregs in mice undergoing Resolution in comparison to Baseline mice.

This reduction was more pronounced in the Resolution – β-carotene mice + anti-CD25

(Supplementary Figure 4A, B). The number of circulating CD25$^-$ Tregs increased in the

Resolution – β-carotene mice + anti-CD25 group in comparison to Baseline mice, remaining

constant in the spleen among all groups (Supplementary Figure 4C, D).

3.4. Anti-CD25 treatment partially abrogates the effect of β-carotene on atherosclerosis

resolution

Tregs possess strong anti-inflammatory properties independently of the presence of CD25 [7,

40], and play an important role on plaque remodeling during atherosclerosis regression [6].

Hence, we evaluated plaque composition in our $Foxp3^{EGFP}$ mice (Figure 5A). As expected, all

experimental groups showed comparable lesion size at the level of the aortic root (Figure 5B).

Mice in the Resolution – β-carotene + IgG group displayed a reduction in CD68 content in the

lesion of approximately 50% in comparison to Baseline mice. Resolution – Control + IgG and

Resolution – β-carotene + anti-CD25 groups showed a 30% reduction in comparison to Baseline

mice, although only the former reached statistical significance (Figure 5C). Collagen content in

the lesion of Resolution – β-carotene + IgG mice was significantly higher in comparison to any

other experimental group. Resolution Control + IgG and Resolution - β-carotene + Anti-CD25

showed comparable results, while Baseline mice had the lowest collagen content among all the

groups (Figure 5D).

We next performed a descriptive discriminant analysis to examine pairwise comparisons between

our four experimental groups. The Baseline group was significantly different in comparison to all

the Resolution counterparts. Resolution - β-carotene + IgG mice were different from their

Resolution – Control + IgG, as we observed in our two previous resolution experiments (Figure

1M and 2F).   We did not observe statistical differences when we compared Resolution - β-

carotene + anti-CD25 to Resolution - Control + IgG (FDR = 0.38) or Resolution - β-carotene +

IgG to (FDR = 0.10) (Figure 5E).

The presence of anti-inflammatory macrophages in the lesions, together with a net egress of pro-

inflammatory macrophages, are key features of atherosclerosis resolution [41]. We quantified

arginase 1 content in the lesion, a key anti-inflammatory marker in mice that contributes to

collagen synthesis [42], and is synergistically upregulated in anti-inflammatory macrophages

exposed to retinoic acid [43, 44] (Figure 5F). Only those mice fed β-carotene, independent of the

injection with IgG or anti-CD25, showed an upregulation of arginase 1 in the lesion (Figure 5G).

We next evaluated retinoic acid signaling in the atherosclerotic lesion by quantifying the levels

of the retinoic acid-sensitive marker Cyp26b1 in the plaque (Figure 5H) [45]. We observed an

upregulation in Cyp26b1 expression in mice fed β-carotene, independently of receiving IgG or

anti-CD25 (Figure 5I). We also evaluated macrophage egress by injecting a single dose of EdU a

week before sacrificing the Baseline mice (see methods for details) (Figure 5J). Only those mice

fed β-carotene, independent of the antibody treatment, displayed a greater egress in comparison

to Baseline mice (Figure 5K). We did not observe changes in monocyte recruitment evaluated by

injecting fluorescently labeled beads 24 h before the sacrifice (Figure 5L, M), nor changes in

Ki67$^+$CD68$^+$ cells, as an indicator of macrophage proliferation in the lesion (Supplementary

Figure 5A, B).

4. Discussion

Seminal studies showed that β-carotene delays atherosclerosis progression in various

experimental models by reducing plasma cholesterol levels [46-48], while greater levels of

circulating β-carotene and retinoic acid are associated with lower risk of cardiovascular disease

in people ([49, 50]. In 2020, we demonstrated that these effects depend on BCO1 activity in

mice, and that a genetic variant in the *BCO1* gene is associated with a reduction in plasma

cholesterol levels in people [15, 17]. Our study shows that β-carotene promotes atherosclerosis

resolution in two independent experimental models in a BCO1-dependent manner. Treg

depletion studies revealed that dietary β-carotene favors Treg expansion in resolving lesions.

Together, these findings identify dietary β-carotene and its conversion to vitamin A as a

promising strategy to ameliorate plaque burden not only by delaying atherosclerosis progression,

but also by reversing atherosclerosis inflammation.

Carotenoids are the main source of vitamin A in human diet, and the only source of this vitamin

in strict vegetarians [51]. Among those carotenoids with provitamin A activity, β-carotene is the

most abundant in our diet and the only compound capable of producing two vitamin A

molecules. Vitamin A is required for vision, embryo development, and immune cell maturation,

making β-carotene a crucial nutrient for human health [52]. Unlike preformed vitamin A, the

intestinal uptake of carotenoids is a protein-mediated process that depends on vitamin A status.

This regulatory mechanism prevents the excessive uptake of provitamin A carotenoids to avoid

vitamin A toxicity [53, 54].

BCO1 is the only enzyme in mammals capable of synthetizing vitamin A from carotenoids [10,

12]. Therefore, it is not surprising that SNPs in the *BCO1* gene are associated to alterations in

circulating carotenoids including β-carotene [55-57]. We recently described that subjects

harboring at least a copy of rs6564851-T allele, which increases BCO1 activity [58], show a

reduction in total cholesterol and non-HDL-C in comparison to those with two copies of the

rs6564851-G variant [17]. A recent study revealed that this variant is also associated with

triglyceride levels in middle-aged individuals [59], implicating BCO1 activity as a novel

regulator of plasma lipid profile. These studies provide a clinical relevance to studies performed

in rodents and rabbits in which β-carotene-rich diets reduce plasma cholesterol and mitigate the

development of atherosclerosis [15, 46-48].

Carotenoids, including β-carotene, possess antioxidant properties in lipid-rich environments [60].

The development of *Bco1*$^{-/-}$ mice of contributed to solving a long-lasting controversy in the

carotenoid field by dissecting the biological effects of intact β-carotene and its role in vitamin A

formation. When *Bco1*$^{-/-}$ mice are fed β-carotene, these mice accumulate large amounts of this

compound in tissues and plasma [11]. In 2011, we reported that wild-type mice fed β-carotene

exhibited a reduction in adipose tissue size. *Bco1*$^{-/-}$ mice subjected to the same experimental

conditions did not chow differences in adipose tissue size in comparison to control-fed mice

despite accumulating large amounts of β-carotene in this tissue [16]. We recently over-expressed

BCO1 in the adipose tissue of *Bco1*$^{-/-}$ mice fed β-carotene. Under these conditions, we observed

an increase in vitamin A levels including retinoic acid, and a reduction of adipose tissue size

[16]. These effects are in line with those reported by Palou's group utilizing both dietary vitamin

A and retinoic acid supplementation, which promotes fatty acid oxidation and thermogenesis in

various experimental models [61-71].

In 2020, we confirmed the implication of BCO1 activity and vitamin A in the development of

atherosclerosis. We compared the effect of β-carotene on *Ldlr*$^{-/-}$ and *Ldlr*$^{-/-}$*Bco1*$^{-/-}$ mice using the

same experimental diets utilized in this study (Supplementary Table 1). In alignment with our

clinical data, β-carotene reduced total cholesterol and non-HDL-C in *Ldlr*$^{-/-}$ mice, but not in *Ldlr*$^{-}$

$^{/-}$*Bco1*$^{-/-}$ mice [15, 17]. β-carotene supplementation during atherosclerosis resolution, either in

Western or Standard diets, failed to improve plasma lipids in comparison to Resolution – Control

groups (Figure 1B, 2A). This discrepancy could be attributed to the duration of the β-carotene

feedings: 12 weeks in our Progression study vs. three to four weeks in our Resolution studies

[15]. Another possibility is that the effect of β-carotene is only evident in mice displaying

elevated plasma lipid levels such as reported in studies using $Ldlr^{-/-}$ and apolipoprotein E-

deficient mice fed atherogenic diets [15, 46, 47].

In our atherosclerosis progression study, we attributed the effects of dietary β-carotene on plasma

lipids to the formation of retinoic acid in the liver. We showed that retinoic acid treatment

decreased cholesterol and triglyceride hepatic secretion rates in cultured hepatocytes and wild-

type mice, suggesting that the effects of β-carotene on plasma lipid profile depend on the

lipidation of hepatic lipoprotein particles [15]. Whether BCO1 activity is associated with the

development of atherosclerotic cardiovascular disease in humans, however, remains unexplored.

Besides its effects on lipid metabolism, retinoic acid modulates immune cell function [72].

Exogenous retinoic acid promotes alternative macrophage activation in various cell culture

models, and skews naïve T cells to anti-inflammatory Tregs by directly upregulating the

transcription factor FoxP3 [73, 74]. The expression of FoxP3 is necessary and sufficient for the

anti-inflammatory phenotype of Tregs [8, 75]. Studies carried out by Loke's group highlighted

the interplay between alternative macrophage activation and Treg differentiation, a process that

was proposed to be mediated by the production and release of retinoic acid by the macrophage

[33, 36]. This hypothesis has not been demonstrated to date, partially due to the technical

difficulty to quantify retinoic acid production in biological samples [76]. We recently overcame

this limitation and demonstrated for the first time that alternatively-activated macrophages

produce and release retinoic acid in a STAT6-dependent manner [44]. We also demonstrated that

exogenous retinoic synergizes with interleukin 4 (IL4) to stimulate the expression of arginase 1,

which is considered an anti-inflammatory marker in murine macrophages [44, 77]. Arginase 1

catalyzes the production of ornithine, which is a precursor of proline [42]. Together with its

derivative hydroxyproline, proline is a key limiting amino acid in the synthesis of collagen.

These results indicate that the combination of retinoic acid with anti-inflammatory signals

typically present during regressing lesions could favor atherosclerosis resolution by stimulating

collagen deposition by the macrophage.

Our PCR data in liver homogenates show β-carotene supplementation upregulates Cyp26a1

expression in the liver, which together with Cyp26b1 and Cyp26c1, are widely considered as

surrogate markers for retinoic signaling in tissues (Supplementary Figure 1B, [26, 45]). Cyp26

family members are responsible for the degradation of intracellular retinoic acid, and their

expression is regulated by retinoic acid response elements in their promoter [26, 45]. To examine

whether β-carotene also stimulates retinoic acid signaling in the atherosclerotic lesion, we

stained the lesions of $Foxp3^{EGFP}$ mice with Cyp26b1 (Figure 5H, I). Among the three Cyp26s,

we selected Cyp26b1 based on our recent RNA seq data in murine bone marrow-derived

macrophages exposed to retinoic acid [44]. We observed that Cyp26b1 was the only Cyp26

family member responsive to retinoic acid in macrophages. Our immunostaining showed that

Cyp26b1 co-localizes with the macrophage marker CD68, in agreement with results obtained in

human lesions showing that CYP26B1 are present in macrophage-rich areas [78]. This study also

reported an association between aggravated atherosclerosis with a SNP in the *CYP26B1* gene

with a greater retinoic acid catabolic activity [78]. These data suggest that greater retinoic acid

levels in the lesion could decrease atherogenesis in humans, in agreement with our preclinical

findings.

Despite the upregulation of Cyp26b1 expression, Treg expansion in regressing lesions in

response to β-carotene was partially suppressed in mice injected with anti-CD25 (Figure 4H). To

rule out intact β-carotene as the driving factor responsible for these effects, we quantified the

number of Tregs in our experiment using $Bco1^{-/-}$ mice. As expected, both Resolution groups

displayed a greater number of Tregs in comparison to Baseline mice, although we did not

observe any effect in response to β-carotene (Supplementary Figure 3D, E). These data suggest

that vitamin A formation is required for Treg expansion in the atherosclerotic lesion of mice. To

our knowledge, this is the first report showing that a dietary intervention with a nutrient

administered at physiological concentrations can modulate Treg levels in the atherosclerotic

lesion. As a reference, our diets were supplemented with 50 mg of β-carotene/kg in comparison

to 80 mg of β-carotene/kg in carrots (USDA Database).

Our HPLC analyses in liver and plasma revealed that our experimental approach did not result in

vitamin A deficiency despite feeding our mice vitamin A-deficient diets for several weeks to

develop atherosclerosis. This was not surprising, since mice are notably resistant to vitamin A

deficiency. Protocols to develop vitamin A deficiency in wild-type mice require feeding the

dams a vitamin A deficient diet during pregnancy through weaning [31], although the use of

genetic models lacking key proteins in vitamin A transport, uptake, or storage are more adequate

and time-efficient [79-81]. Future experiments utilizing these genetic models could shed light on

the role of vitamin A deficiency on Treg expansion and its consequence on atherosclerosis

resolution.

To establish a direct link between the effect of β-carotene supplementation and Tregs in our

experimental model, we decided to deplete Tregs in $Foxp3^{EGFP}$ mice by infusing them with anti-

CD25, following Sharma and colleagues' approach [6]. However, our data show that this

strategy failed to completely deplete Tregs, as previously reported by other investigators [7, 40].

CD25$^-$ Tregs remained in circulation and tissues including aortic lesions (Figure 4E-H and

Supplementary Figure 4). It is possible that the conversion of β-carotene to retinoic acid favored

Treg expansion by specifically favoring CD25⁻ Treg proliferation, which are insensitive to anti-

CD25.

CD25 is a crucial component of the IL2 receptor and it is expressed in $CD4^+$ T cell subsets other

than Tregs ($CD4^+CD25^+FoxP3^-$) [82]. Therefore, strategies targeting $FoxP3^+$ cells directly are an

attractive approach to specifically dissect the role of Tregs in our experimental model. For

example, the use of *Foxp3^{DTR}* mice (#016958, Jackson Labs), which express the diphtheria toxin

receptor, would allow us to specifically deplete $FoxP3^+$ cells independently of the expression of

CD25 [83]. This approach would also eliminate $CD8^+CD25^+FoxP3^+$ Tregs, a relatively new Treg

subpopulation [84]. However, Treg depletion in *Foxp3^{DTR}* mice requires injections with

diphtheria toxin, which can result in potential adverse effects under certain experimental

conditions [85]. Future studies dissecting the contribution of T cell subpopulations and the role

vitamin A and other nutrients could play in their differentiation are needed to fully understand

their role during atherosclerosis resolution.

Whether β-carotene supplementation during atherosclerosis resolution alters Treg content in

other organs and disease models were outside of the scope of this study. For example, an

increase in Treg number in developing tumors could result in the reprograming of pro-

inflammatory macrophages towards resolving macrophages that could contribute to tumor

expansion [86]. In summary, partial Treg depletion was sufficient to mitigate the effects of β-

carotene on plaque composition during the resolution of atherosclerosis, providing a direct link

between β-carotene and Treg number (Figure 5A-I).

Another question remains unanswered: What is the source of vitamin A that favors Treg

expansion in the lesion? Our HPLC analyses show that circulating β-carotene in wild-type mice

is marginal, although hepatic vitamin A stores highlight an increase in vitamin A formation

(Figure 1H and Figure 3A, B). It is not clear whether naïve T cells express BCO1, which would

enable them to locally produce vitamin A and its derivative retinoic acid. Previous work has

related tissue macrophages as the responsible for retinoic acid production and suggested that

these cells would release it to signal nearby naïve T cells that in turn, could differentiate into

Tregs. Retinoic acid could be originated from β-carotene or retinol/retinyl esters stored in the

cell, although our RNA sequencing analysis revealed that BCO1 expression in plaque

macrophages is relatively limited [15]. Plaque macrophages, and T cells in a lesser extent,

express various scavenger receptors that have been linked to the uptake of retinoids and

carotenoids such as the scavenger receptor class BI, lipoprotein lipase and the cluster of

differentiation 36 [87]. Determining whether naïve T cells rely on plaque macrophages to obtain

retinoids and activate FoxP3 to differentiate into Treg is not clear to this day.

In summary, we report for the first time that vitamin A production from β-carotene favors Treg

expansion in the atherosclerotic lesion, a process that contributes to atherosclerosis resolution.

Together with the cholesterol-lowering effects of β-carotene described in the past, unveils a dual

role of dietary β-carotene and the enzyme BCO1: (1) BCO1 delays atherosclerosis progression

by reducing plasma cholesterol, and (2) favor atherosclerosis resolution by modulating Treg

levels and macrophage polarization status.

Figures

Figure 1. β-carotene accelerates atherosclerosis resolution in wild-type mice infused with anti-sense oligonucleotide targeting the low-density lipoprotein receptor (ASO-LDLR). (A) Four-week-old male and female wild-type mice were fed a purified Western diet deficient in vitamin A (WD-VAD) and injected with antisense oligonucleotide targeting the low-density lipoprotein receptor (ASO-LDLR) once a week for 16 weeks to induce atherosclerosis. After 16 weeks, a group of mice was harvested (Baseline) and the rest of the mice were injected once with sense oligonucleotide (SO-LDLR) to inactivate ASO-LDLR and promote atherosclerosis

resolution. Mice undergoing resolution were either kept on the same diet (Resolution - Control)

or switched to a Western diet supplemented with 50 mg/kg of β-carotene (Resolution - β-

carotene) for three more weeks. **(B)** Plasma lipid levels at the moment of the sacrifice. **(C)**

Cholesterol and **(D)** triglyceride distribution in FPLC-fractioned plasma (data pooled from five

mice/group). **(E)** Relative LDLR mRNA, and **(F)** protein expressions in the liver. **(G)** Circulating

vitamin A (all-*trans* retinol), and **(H)** hepatic retinyl ester stores determined by HPLC. **(I)**

Representative images for macrophage (CD68$^+$, top panels), and picrosirius staining to identify

collagen using the bright-field (middle panels) or polarized light (bottom panels). **(J)** Plaque size,

(K) relative CD68 content, and **(L)** collagen content in the lesion**. (M)** Descriptive discriminant

analysis employing the relative CD68 and collagen contents in the lesion as variables

highlighting the FDR for each comparison. Each dot in the plot represents an individual mouse

(n = 10 to 12/group). Values are represented as means ± SEM. Statistical differences were

evaluated using one-way ANOVA with Tukey's multiple comparisons test. Differences between

groups were considered significant with a p-value < 0.05. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.005$;

**** $p < 0.001$. Size bar = 200 µm.

578
Figure 2. β-carotene accelerates atherosclerosis resolution in low-density lipoprotein

deficient (*Ldlr⁻/⁻*) mice subjected to dietary switch. Four-week-old male and female *Ldlr⁻/⁻* mice

were fed a purified Western diet deficient in vitamin A (WD-VAD) for 12 weeks to induce

atherosclerosis. After 12 weeks, a group of mice was harvested (Baseline) and the rest of the

mice were switched to a Standard diet (Resolution-Control) or the same diet supplemented with

50 mg/kg of β-carotene (Resolution-β-carotene) for four more weeks. **(A)** Total cholesterol

plasma levels at the moment of the sacrifice. **(B)** Representative images for macrophage (CD68⁺,

top panels), and picrosirius staining to identify collagen using the bright-field (middle panels) or

polarized light (bottom panels). **(C)** Plaque size, **(D)** relative CD68 content, and **(E)** collagen

content in the lesion**. (F)** Descriptive discriminant analysis employing the relative CD68 and

collagen contents in the lesion as variables highlighting the FDR for each comparison. Each dot

in the plot represents an individual mouse (n = 9 to 12/group). **(A-E)** Values are represented as

means ± SEM. Statistical differences were evaluated using one-way ANOVA with Tukey's

multiple comparisons test. Differences between groups were considered significant with a p-

value < 0.05. * p < 0.05; *** p < 0.005; **** p < 0.001. Size bar = 200 µm.

Figure 3. β-carotene supplementation does alter atherosclerosis resolution in β-carotene oxygenase 1-deficient (*Bco1⁻/⁻*) mice infused with ASO-LDLR. Four-week-old male and female *Bco1⁻/⁻* mice were fed a purified Western diet deficient in vitamin A (WD-VAD) and injected with antisense oligonucleotide targeting the low-density lipoprotein receptor (ASO-LDLR) once a week for 16 weeks to induce the development of atherosclerosis. After 16 weeks, a group of mice was harvested (Baseline) and the rest of the mice were injected once with sense oligonucleotide (SO-LDLR) to block ASO-LDLR (Resolution). Mice undergoing resolution were either kept on the same diet (Resolution-Control) or switched to a Western diet supplemented with 50 mg/kg of β-carotene (Resolution- β-carotene) for three more weeks. **(A)** β-carotene levels in plasma and **(B)** liver at the sacrifice determined by HPLC. **(C)** Representative images for macrophage (CD68⁺, top panels), and picrosirius staining to identify collagen using the bright-field (middle panels) or polarized light (bottom panels). **(D)** Plaque size, **(E)** relative CD68 content, and **(F)** collagen content in the lesion. **(G)** Descriptive discriminant analysis employing the relative CD68 and collagen contents in the lesion as variables highlighting the FDR for each comparison. Each dot in the plot represents an individual mouse (n = 5 to 11/group). **(A-F)** Values are represented as means ± SEM. Statistical differences were evaluated using one-way ANOVA with Tukey's multiple comparisons test. Differences between groups were considered significant with a p-value < 0.05. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.005$; **** $p < 0.001$. Size bar = 200 µm.

Figure 4. Effect of anti-CD25 treatment on Treg number. (A) Four-week-old male and female

mice expressing enhanced green fluorescence protein (EGFP) under the control of the forkhead

box P3 (*Foxp3*) promoter (*Foxp3^{EGFP}* mice) were fed a purified Western diet deficient in vitamin

A (WD-VAD) and injected with antisense oligonucleotide targeting the low-density lipoprotein

receptor (ASO-LDLR) once a week for 16 weeks to induce the development of atherosclerosis.

After 16 weeks, a group of mice was harvested (Baseline) and the rest of the mice were injected

once with sense oligonucleotide (SO-LDLR) to block ASO-LDLR (Resolution). Mice

undergoing resolution were either kept on the same diet (Resolution-Control) or switched to a

Western diet supplemented with 50 mg/kg of β-carotene (Resolution- β-carotene) for three more

weeks. An additional group of mice fed with β-carotene was injected twice before sacrifice with

anti-CD25 monoclonal antibody to deplete Treg (Resolution-β-carotene+anti-CD25). The rest of

the resolution groups were injected with IgG isotype control antibody. **(B)** Representative flow

cytometry panels showing splenic $CD25^+FoxP3^+$ ($CD25^+$ Treg) cells in mice injected with IgG or

anti-CD25. **(C)** Quantification of the splenic and **(D)** circulating blood levels of $CD25^+$ Treg

cells determined by flow cytometry. **(E)** Representative confocal images show the presence of

total Tregs ($CD25^-FoxP3^+$ + $CD25^+$ Tregs) and $CD25^+FoxP3^-$ T cells in the lesion of mice

injected with IgG (left panel) or anti-CD25 (right panel). **(F)** Representative confocal images

show the presence of total Tregs and $CD25^+FoxP3^-$ T cells in lymph nodes of mice injected with

IgG (left panel) or anti-CD25 (right panel). quantification of $CD25^+$ Tregs in the lesions. **(G)**

Representative confocal image and **(H)** quantification of total Tregs in the lesion. Each dot in the

plot represents an individual mouse (n = 5 to 11 mice/group). Values are represented as means ±

SEM. Statistical differences were evaluated using one-way ANOVA with Tukey's multiple

comparisons test. Differences between groups were considered significant with a p-value < 0.05.

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.005$; **** $p < 0.001$.

Figure 5. Effect of anti-CD25 treatment on lesion composition and monocyte/macrophage

trafficking. Four-week-old male and female expressing enhanced green fluorescence protein

(EGFP) under the control of the forkhead box P3 (*Foxp3*) promoter (*Foxp3^{EGFP}* mice) were fed a

purified Western diet deficient in vitamin A (WD-VAD) and injected with antisense

oligonucleotide targeting the low-density lipoprotein receptor (ASO-LDLR) once a week for 16

weeks to induce the development of atherosclerosis. After 16 weeks, a group of mice was

harvested (Baseline) and the rest of the mice were injected once with sense oligonucleotide (SO-

LDLR) to block ASO-LDLR (Resolution). Mice undergoing resolution were either kept on the

same diet (Resolution-Control) or switched to a Western diet supplemented with 50 mg/kg of β-

carotene (Resolution- β-carotene) for three more weeks. An additional group of mice fed with β-

carotene was injected twice before sacrifice with anti-CD25 monoclonal antibody to deplete Treg

(Resolution-β-carotene+anti-CD25). The rest of the resolution groups were injected with IgG

isotype control antibody. To quantify macrophage egress and monocyte recruitment, we injected

a dose of EdU at week 15 and fluorescently labeled beads two days before harvesting the mice,

respectively (see methods for details). **(A)** Representative images for macrophage (CD68^{+}, top

panels), and picrosirius staining to identify collagen using the bright-field (middle panels) or

polarized light (bottom panels). Size bar = 200 µm. **(B)** Plaque size, **(C)** relative CD68 content,

and **(D)** collagen content in the lesion. **(E)** Descriptive discriminant analysis employing the

relative CD68 and collagen contents in the lesion as variables highlighting the FDR for each

comparison. **(F)** Representative confocal image showing arginase 1 (green), CD68 (red), and

DAPI (blue) in the lesion. **(G)** relative arginase 1 area in the lesion. **(H)** Representative confocal

image showing Cyp26b1 (green), CD68 (red), and DAPI (blue) in the lesion. **(I)** relative

Cyp26b1 area in the lesion. **(J)** EdU+ macrophages were identified by the colocalization of EdU

(green) and DAPI (blue) in CD68^{+} (red) cells. **(K)** Number of EdU+ macrophages in the lesion.

(L) Newly recruited monocytes were identified, and **(M)** quantified by the presence of beads

(green) on the lesion. Size bars = 50 µm. Each dot in the plot represents an individual mouse (n =

5 to 11 mice/group). Values are represented as means ± SEM. Statistical differences were

evaluated using one-way ANOVA with Tukey's multiple comparisons test. Differences between

groups were considered significant with a p-value < 0.05. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.005$;

**** $p < 0.001$.

672
673
674

REFERENCES

1. Newby, A.C., et al., *Vulnerable atherosclerotic plaque metalloproteinases and foam cell*
phenotypes. Thrombosis and haemostasis, 2009. **101**(6): p. 1006-1011.
2. Goldberg, I.J., G. Sharma, and E.A. Fisher, *Atherosclerosis: Making a U Turn.* Annu Rev
Med, 2020. **71**: p. 191-201.
3. Andersson, J., P. Libby, and G.K. Hansson, *Adaptive immunity and atherosclerosis.*
Clinical Immunology, 2010. **134**(1): p. 33-46.
4. Witztum, J.L. and A.H. Lichtman, *The influence of innate and adaptive immune*
responses on atherosclerosis. Annual review of pathology, 2014. **9**: p. 73-102.
5. Onda, M., K. Kobayashi, and I. Pastan, *Depletion of regulatory T cells in tumors with an*
anti-CD25 immunotoxin induces CD8 T cell-mediated systemic antitumor immunity. Proc
Natl Acad Sci U S A, 2019. **116**(10): p. 4575-4582.
6. Sharma, M., et al., *Regulatory T Cells License Macrophage Pro-Resolving Functions*
During Atherosclerosis Regression. Circ Res, 2020. **127**(3): p. 335-353.
7. Couper, K.N., et al., *Incomplete depletion and rapid regeneration of Foxp3+ regulatory*
T cells following anti-CD25 treatment in malaria-infected mice. J Immunol, 2007.
178(7): p. 4136-46.
8. Fontenot, J.D., et al., *Regulatory T cell lineage specification by the forkhead*
transcription factor foxp3. Immunity, 2005. **22**(3): p. 329-41.
9. Elias, K.M., et al., *Retinoic acid inhibits Th17 polarization and enhances FoxP3*
expression through a Stat-3/Stat-5 independent signaling pathway. Blood, 2008. **111**(3):
p. 1013-20.
10. von Lintig, J. and K. Vogt, *Filling the gap in vitamin A research. Molecular*
identification of an enzyme cleaving beta-carotene to retinal. J Biol Chem, 2000.
275(16): p. 11915-20.
11. Hessel, S., et al., *CMO1 deficiency abolishes vitamin A production from beta-carotene*
and alters lipid metabolism in mice. J Biol Chem, 2007. **282**(46): p. 33553-61.
12. Amengual, J., et al., *Two carotenoid oxygenases contribute to mammalian provitamin A*
metabolism. J Biol Chem, 2013. **288**(47): p. 34081-96.
13. Amengual, J., et al., *Beta-carotene reduces body adiposity of mice via BCMO1.* PLoS
One, 2011. **6**(6): p. e20644.
14. Lobo, G.P., et al., *Beta,beta-carotene decreases peroxisome proliferator receptor gamma*
activity and reduces lipid storage capacity of adipocytes in a beta,beta-carotene
oxygenase 1-dependent manner. J Biol Chem, 2010. **285**(36): p. 27891-9.
15. Zhou, F., et al., *beta-carotene conversion to vitamin A delays atherosclerosis progression*
by decreasing hepatic lipid secretion in mice. J Lipid Res, 2020.
16. Coronel, J., et al., *The conversion of beta-carotene to vitamin A in adipocytes drives the*
anti-obesogenic effects of beta-carotene in mice. Mol Metab, 2022: p. 101640.
17. Amengual, J., et al., *beta-Carotene Oxygenase 1 Activity Modulates Circulating*
Cholesterol Concentrations in Mice and Humans. J Nutr, 2020.
18. Amengual, J., et al., *Short-Term Acyl-CoA:Cholesterol Acyltransferase Inhibition,*
Combined with Apoprotein A1 Overexpression, Promotes Atherosclerosis Inflammation
Resolution in Mice. Mol Pharmacol, 2021. **99**(3): p. 175-183.
19. Josefs, T., et al., *Atherosclerosis Regression and Cholesterol Efflux in*
Hypertriglyceridemic Mice. Circ Res, 2021. **128**(6): p. 690-705.

20. Josefs, T., et al., *Neutrophil extracellular traps promote macrophage inflammation and*
impair atherosclerosis resolution in diabetic mice. JCI Insight, 2020. **5**(7).
21. Barrett, T.J., et al., *Apolipoprotein AI) Promotes Atherosclerosis Regression in Diabetic*
Mice by Suppressing Myelopoiesis and Plaque Inflammation. Circulation, 2019. **140**(14):
p. 1170-1184.
22. Basu, D., et al., *Novel Reversible Model of Atherosclerosis and Regression Using*
Oligonucleotide Regulation of the LDL Receptor. Circulation Research, 2018. **122**(4): p.
560-567.
23. Weinstock, A. and E.A. Fisher, *Methods to Study Monocyte and Macrophage Trafficking*
in Atherosclerosis Progression and Resolution. Methods Mol Biol, 2019. **1951**: p. 153-
165.
24. Benjamini, Y., et al., *Controlling the false discovery rate in behavior genetics research.*
Behav Brain Res, 2001. **125**(1-2): p. 279-84.
25. Emoto, Y., et al., *Retinoic acid-metabolizing enzyme Cyp26a1 is essential for*
determining territories of hindbrain and spinal cord in zebrafish. Dev Biol, 2005. **278**(2):
p. 415-27.
26. Abu-Abed, S., et al., *Differential expression of the retinoic acid-metabolizing enzymes*
CYP26A1 and CYP26B1 during murine organogenesis. Mech Dev, 2002. **110**(1-2): p.
173-7.
27. Basu, D., et al., *Novel Reversible Model of Atherosclerosis and Regression Using*
Oligonucleotide Regulation of the LDL Receptor. Circ Res, 2018. **122**(4): p. 560-567.
28. Libby, P. and M. Aikawa, *Stabilization of atherosclerotic plaques: New mechanisms and*
clinical targets. Nature Medicine, 2002. **8**(11): p. 1257-1262.
29. Soehnlein, O. and P. Libby, *Targeting inflammation in atherosclerosis - from*
experimental insights to the clinic. Nat Rev Drug Discov, 2021. **20**(8): p. 589-610.
30. Yu, M., et al., *Targeted Nanotherapeutics Encapsulating Liver X Receptor Agonist*
GW3965 Enhance Antiatherogenic Effects without Adverse Effects on Hepatic Lipid
Metabolism in Ldlr(-/-) Mice. Adv Healthc Mater, 2017. **6**(20).
31. Gundra, U.M., et al., *Vitamin A mediates conversion of monocyte-derived macrophages*
into tissue-resident macrophages during alternative activation. Nat Immunol, 2017.
18(6): p. 642-653.
32. Ouimet, M., et al., *MicroRNA-33-dependent regulation of macrophage metabolism*
directs immune cell polarization in atherosclerosis. J Clin Invest, 2015. **125**(12): p. 4334-
48.
33. Gundra, U.M., et al., *Alternatively activated macrophages derived from monocytes and*
tissue macrophages are phenotypically and functionally distinct. Blood, 2014. **123**(20): p.
e110-22.
34. Girgis, N.M., et al., *Ly6Chigh monocytes become alternatively activated macrophages in*
schistosome granulomas with help from CD4+ cells. PLoS Pathog, 2014. **10**(6): p.
e1004080.
35. Girgis, N.M., U.M. Gundra, and P. Loke, *Immune regulation during helminth infections.*
PLoS Pathog, 2013. **9**(4): p. e1003250.
36. Broadhurst, M.J., et al., *Upregulation of retinal dehydrogenase 2 in alternatively*
activated macrophages during retinoid-dependent type-2 immunity to helminth infection
in mice. PLoS Pathog, 2012. **8**(8): p. e1002883.

37. Foks, A.C., A.H. Lichtman, and J. Kuiper, *Treating atherosclerosis with regulatory T*
cells. Arterioscler Thromb Vasc Biol, 2015. **35**(2): p. 280-7.
38. Ait-Oufella, H., et al., *Natural regulatory T cells control the development of*
atherosclerosis in mice. Nat Med, 2006. **12**(2): p. 178-80.
39. Haribhai, D., et al., *Regulatory T cells dynamically control the primary immune response*
to foreign antigen. J Immunol, 2007. **178**(5): p. 2961-72.
40. Hayes, E.T., et al., *Regulatory T Cells Maintain Selective Access to IL-2 and Immune*
Homeostasis despite Substantially Reduced CD25 Function. J Immunol, 2020. **205**(10):
p. 2667-2678.
41. Moore, K.J., et al., *Macrophage Trafficking, Inflammatory Resolution, and Genomics in*
Atherosclerosis: JACC Macrophage in CVD Series (Part 2). J Am Coll Cardiol, 2018.
72(18): p. 2181-2197.
42. Szondi, D.C., et al., *Arginase Signalling as a Key Player in Chronic Wound*
Pathophysiology and Healing. Front Mol Biosci, 2021. **8**: p. 773866.
43. Yurdagul, A., Jr., et al., *Macrophage Metabolism of Apoptotic Cell-Derived Arginine*
Promotes Continual Efferocytosis and Resolution of Injury. Cell Metab, 2020. **31**(3): p.
518-533 e10.
44. Pinos, I., et al., *Functional characterization of interleukin 4 and retinoic acid signaling*
crosstalk during alternative macrophage activation. Biochim Biophys Acta Mol Cell
Biol Lipids, 2023: p. 159291.
45. Ross, A.C. and R. Zolfaghari, *Cytochrome P450s in the regulation of cellular retinoic*
acid metabolism. Annu Rev Nutr, 2011. **31**: p. 65-87.
46. Relevy, N.Z., et al., *Vitamin A-deficient diet accelerated atherogenesis in apolipoprotein*
E(-/-) mice and dietary beta-carotene prevents this consequence. Biomed Res Int, 2015.
2015: p. 758723.
47. Harari, A., et al., *A 9-cis beta-carotene-enriched diet inhibits atherogenesis and fatty*
liver formation in LDL receptor knockout mice. J Nutr, 2008. **138**(10): p. 1923-30.
48. Shaish, A., et al., *Beta-carotene inhibits atherosclerosis in hypercholesterolemic rabbits.*
J Clin Invest, 1995. **96**(4): p. 2075-82.
49. Wang, Y., et al., *Dietary carotenoids are associated with cardiovascular disease risk*
biomarkers mediated by serum carotenoid concentrations. J Nutr, 2014. **144**(7): p. 1067-
74.
50. Huang, J., et al., *Serum Beta Carotene and Overall and Cause-Specific Mortality.* Circ
Res, 2018. **123**(12): p. 1339-1349.
51. Grune, T., et al., *Beta-carotene is an important vitamin A source for humans.* J Nutr,
2010. **140**(12): p. 2268S-2285S.
52. Coronel, J., I. Pinos, and J. Amengual, *beta-carotene in Obesity Research: Technical*
Considerations and Current Status of the Field. Nutrients, 2019. **11**(4).
53. Lobo, G.P., et al., *Genetics and diet regulate vitamin A production via the homeobox*
transcription factor ISX. J Biol Chem, 2013. **288**(13): p. 9017-27.
54. Seino, Y., et al., *Isx participates in the maintenance of vitamin A metabolism by*
regulation of beta-carotene 15,15'-monooxygenase (Bcmo1) expression. J Biol Chem,
2008. **283**(8): p. 4905-11.
55. Clifford, A.J., et al., *Single nucleotide polymorphisms in CETP, SLC46A1, SLC19A1,*
CD36, BCMO1, APOA5, and ABCA1 are significant predictors of plasma HDL in
healthy adults. Lipids Health Dis, 2013. **12**: p. 66.

56.   Yabuta, S., et al., *Common SNP rs6564851 in the BCO1 Gene Affects the Circulating*
Levels of beta-Carotene and the Daily Intake of Carotenoids in Healthy Japanese
Women. PLoS One, 2016. **11**(12): p. e0168857.
57.   Moran, N.E., et al., *Single Nucleotide Polymorphisms in beta-Carotene Oxygenase 1 are*
Associated with Plasma Lycopene Responses to a Tomato-Soy Juice Intervention in Men
with Prostate Cancer. J Nutr, 2019. **149**(3): p. 381-397.
58.   Lietz, G., et al., *Single nucleotide polymorphisms upstream from the beta-carotene*
15,15'-monoxygenase gene influence provitamin A conversion efficiency in female
volunteers. J Nutr, 2012. **142**(1): p. 161S-5S.
59.   Leon-Reyes, G., et al., *Common variant rs6564851 near the beta-carotene oxygenase 1*
gene is associated with plasma triglycerides levels in middle-aged Mexican men adults.
Nutr Res, 2022. **103**: p. 30-39.
60.   Rodriguez-Concepcion, M., et al., *A global perspective on carotenoids: Metabolism,*
biotechnology, and benefits for nutrition and health. Prog Lipid Res, 2018. **70**: p. 62-93.
61.   Bonet, M.L., et al., *Carotenoids and carotenoid conversion products in adipose tissue*
biology and obesity: Pre-clinical and human studies. Biochim Biophys Acta Mol Cell
Biol Lipids, 2020: p. 158676.
62.   Amengual, J., et al., *Retinoic Acid Increases Fatty Acid Oxidation and Irisin Expression*
in Skeletal Muscle Cells and Impacts Irisin In Vivo. Cell Physiol Biochem, 2018. **46**(1):
p. 187-202.
63.   Granados, N., et al., *Vitamin A supplementation in early life affects later response to an*
obesogenic diet in rats. Int J Obes (Lond), 2013. **37**(9): p. 1169-76.
64.   Amengual, J., et al., *Induction of carnitine palmitoyl transferase 1 and fatty acid*
oxidation by retinoic acid in HepG2 cells. Int J Biochem Cell Biol, 2012. **44**(11): p.
2019-27.
65.   Amengual, J., et al., *Retinoic acid treatment enhances lipid oxidation and inhibits lipid*
biosynthesis capacities in the liver of mice. Cell Physiol Biochem, 2010. **25**(6): p. 657-66.
66.   Amengual, J., et al., *Retinoic acid treatment increases lipid oxidation capacity in skeletal*
muscle of mice. Obesity (Silver Spring), 2008. **16**(3): p. 585-91.
67.   Mercader, J., et al., *Remodeling of white adipose tissue after retinoic acid administration*
in mice. Endocrinology, 2006. **147**(11): p. 5325-32.
68.   Felipe, F., et al., *Effects of retinoic acid administration and dietary vitamin A*
supplementation on leptin expression in mice: lack of correlation with changes of adipose
tissue mass and food intake. Biochim Biophys Acta, 2005. **1740**(2): p. 258-65.
69.   Felipe, F., et al., *Modulation of resistin expression by retinoic acid and vitamin A status.*
Diabetes, 2004. **53**(4): p. 882-9.
70.   Puigserver, P., et al., *In vitro and in vivo induction of brown adipocyte uncoupling*
protein (thermogenin) by retinoic acid. Biochem J, 1996. **317 (Pt 3)**: p. 827-33.
71.   Berry, D.C. and N. Noy, *All-trans-retinoic acid represses obesity and insulin resistance*
by activating both PPAR{beta}/{delta} and RAR. Mol Cell Biol, 2009.
72.   Miller, A.P., J. Coronel, and J. Amengual, *The role of beta-carotene and vitamin A in*
atherogenesis: Evidences from preclinical and clinical studies. Biochim Biophys Acta
Mol Cell Biol Lipids, 2020: p. 158635.
73.   He, L., et al., *Global characterization of macrophage polarization mechanisms and*
identification of M2-type polarization inhibitors. Cell Rep, 2021. **37**(5): p. 109955.

74.   Mucida, D., et al., *Reciprocal TH17 and regulatory T cell differentiation mediated by*
retinoic acid. Science, 2007. **317**(5835): p. 256-60.

75.   Hori, S., T. Nomura, and S. Sakaguchi, *Control of regulatory T cell development by the*
transcription factor Foxp3. Science, 2003. **299**(5609): p. 1057-61.

76.   Kane, M.A., et al., *Quantitative profiling of endogenous retinoic acid in vivo and in vitro*
by tandem mass spectrometry. Anal Chem, 2008. **80**(5): p. 1702-8.

77.   Murray, P.J., et al., *Macrophage activation and polarization: nomenclature and*
experimental guidelines. Immunity, 2014. **41**(1): p. 14-20.

78.   Krivospitskaya, O., et al., *A CYP26B1 polymorphism enhances retinoic acid catabolism*
and may aggravate atherosclerosis. Mol Med, 2012. **18**(1): p. 712-8.

79.   Quadro, L., et al., *Impaired retinal function and vitamin A availability in mice lacking*
retinol-binding protein. Embo J, 1999. **18**(17): p. 4633-44.

80.   O'Byrne, S.M., et al., *Retinoid absorption and storage is impaired in mice lacking*
lecithin:retinol acyltransferase (LRAT). J Biol Chem, 2005. **280**(42): p. 35647-57.

81.   Amengual, J., et al., *STRA6 is critical for cellular vitamin A uptake and homeostasis.*
Hum Mol Genet, 2014. **23**(20): p. 5402-17.

82.   Wing, K., et al., *Characterization of human CD25+ CD4+ T cells in thymus, cord and*
adult blood. Immunology, 2002. **106**(2): p. 190-9.

83.   Kim, J.M., J.P. Rasmussen, and A.Y. Rudensky, *Regulatory T cells prevent catastrophic*
autoimmunity throughout the lifespan of mice. Nat Immunol, 2007. **8**(2): p. 191-7.

84.   Churlaud, G., et al., *Human and Mouse CD8(+)CD25(+)FOXP3(+) Regulatory T Cells*
at Steady State and during Interleukin-2 Therapy. Front Immunol, 2015. **6**: p. 171.

85.   Christiaansen, A.F., P.M. Boggiatto, and S.M. Varga, *Limitations of Foxp3(+) Treg*
depletion following viral infection in DEREG mice. J Immunol Methods, 2014. **406**: p.
58-65.

86.   Ohue, Y. and H. Nishikawa, *Regulatory T (Treg) cells in cancer: Can Treg cells be a new*
therapeutic target? Cancer Sci, 2019. **110**(7): p. 2080-2089.

87.   Reboul, E., *Mechanisms of Carotenoid Intestinal Absorption: Where Do We Stand?*
Nutrients, 2019. **11**(4).