Title:

The CRL plastid outer envelope protein supports TOC75-V / OEP80 complex formation

in *Arabidopsis*

Authors:

Ryo Yoshimura[1], Syun Minamikawa[1], Takamasa Suzuki[2], David Latrasse[3,4], Sanchari

Sicar[3,4], Cécile Raynaud[3,4], Moussa Benhamed[3,4,5], Yasushi Yoshioka[1]

Authors address:

[1]Graduate School of Science, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, 464-

8602, Japan. [2]Department of Biological Chemistry, College of Bioscience and

Biotechnology, Chubu University, 1200 Matsumoto-cho, Kasugai, Aichi 487-8501, Japan.

[3]Université Paris-Saclay, CNRS, INRAE, Univ Evry, Institute of Plant Sciences Paris-

Saclay (IPS2), 91405, Orsay, France. [4]Université de Paris, CNRS, INRAE, Institute of

Plant Sciences Paris-Saclay (IPS2), 91405, Orsay, France. [5]Institut Universitaire de

France (IUF).

Corresponding author:

Yasushi Yoshioka

Graduate School of Science, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, 464-

8602, Japan.

e-mail: yoshioka@bio.nagoya-u.ac.jp

Associate Professor, Grad Sch Sci, Nagoya Univ.

Abstract

Embedded β-barrel proteins in the outer envelope membrane mediate most cellular

traffic between the cytoplasm and the plastids. Although TOC75-V/OEP80 has been

implicated in the insertion and assembly of β-barrel proteins in the outer envelope

membrane of *Arabidopsis thaliana*, relatively little is known about this process.

CRUMPLED LEAF (*CRL*) encodes a protein localizing in the outer envelope membrane,

and its loss of function results in pleiotropic defects, including altered plant

morphogenesis, growth retardation, suppression of plastid division, and spontaneous

light intensity-dependent localized cell death. A suppressor screen conducted on

mutagenized *crl* mutants with ethyl methanesulfonate revealed that a missense mutation

in *OEP80* suppresses *crl*'s pleiotropic defects. Furthermore, we found that the complex

formation of OEP80 was compromised in *crl*. Furthermore, we demonstrated that CRL

interacts with OEP80 *in vivo* and that a portion of CRL is present in protein complexes

with the same molecular weight as the OEP80-associated complex. Our results suggest

that CRL interacts with OEP80 to regulate its complex formation. CRL has been shown

to be involved in plastid protein import; therefore, pleiotropic defects in *crl* are likely due

to the combined effects of decreased plastid protein import and altered membrane

integration of β-barrel proteins in the outer envelope membrane. This study sheds light

on the mechanisms that allow the integration of β-barrel proteins into the outer envelope

membrane of plastids and the significance of this finding for plant cellular processes.

Introduction

Plastids are double membrane-bound organelles that are isolated from the

cytoplasm by an inner envelope membrane (IEM) and an outer envelope membrane

(OEM). Since numerous essential metabolic pathways reside in plastids, the normal

cellular activity requires dynamic exchanges of substances between the cytosol and

plastids via OEM and IEM. Multiple membrane-embedded β-barrel proteins localized in

the OEM are essential for the exchange of substances between the cytosol and plastid.

Porin-like β-barrel proteins, OUTER ENVELOPE PROTEIN 21, 24, 37, 40 kDa (OEP21,

24, 37, 40), were discovered in the pea OEM. These proteins form channels and

exchange ions and metabolites with solute-specific selectivity (Pohlmeyer et al., 1998;

Bölter et al., 1999; Goetze et al., 2006; Harsman et al., 2016).

TRIGALACTOSYLDIACYLGLYCEROL4 (TGD4) is proposed to be a β-barrel protein

localized in OEM in *Arabidopsis thaliana*. TGD4 is involved in the transfer of lipids from

the ER to chloroplasts (Wang et al., 2012). β-barrel proteins also contribute to protein

import within plastids. Most plastid-localized proteins are encoded in the nucleus and

translated into the cytosol as precursors with an N-terminal transit peptide (Leister, 2003).

These are imported from OEM and IEM through the transport complex (TOC) (translocon

on the outer chloroplast membrane) and TIC (translocon on the inner chloroplast

membrane) protein import apparatus (Bölter and Soll, 2016). TOC75, one of the core

components of the TOC, is a β-barrel protein that is a member of the Omp85 family (Day

et al., 2014).

Although numerous β-barrel proteins have been shown to contribute to the

transport of various substances across the OEM, much less is known about the

mechanisms that permit the incorporation of these β-barrel proteins into the OEM.

However, a paralog of TOC75, TOC75-V/OEP80 (hereafter OEP80), which is a member

of the Omp85 family, was recently proposed to play a role in the integration of β-barrel

proteins into OEM (Gross et al., 2021). OEM β-barrel proteins can be translocated into

the intermembrane space (IMS) via the transmembrane TOC and then integrated into

the OEM by interacting with OEP80 (Gross et al., 2021). It has been hypothesized that additional factors are involved in this process, but none have been identified (Gross et al., 2021). In addition to OEP80, it has been suggested that the TamB homolog in DEFECTIVE KERNEL5 maize has bacterial TamB-like function (Zhang et al., 2019), while its *Arabidopsis* ortholog TIC236 is involved in the import of protein into the plastid as a component of TIC (Chen et al., 2018).

Although little is known about the molecular mechanisms of integration of β-barrel proteins into the OEM of plastids, it is well understood how β-barrel proteins are inserted into the outer membranes (OMs) of Gram-negative bacteria and mitochondria. This process involves β-barrel proteins belonging to the Omp85 family, such as BamA, and TamA in gram-negative bacteria and Sam50 in mitochondria, respectively (Walther et al., 2009; Hagan et al., 2011; Selkrig et al., 2012). For example, BamA in *Escherichia coli* forms the β-barrel assembly machinery with BamB-E and facilitates the assembly of β-barrel proteins in the OM (Walther et al., 2009; Hagan et al., 2011). TamA and TamB form the translocation and assembly module in *E. coli* to facilitate the assembly and secretion of β-barrel proteins, including autotransporters (Selkrig et al., 2012). Sam50 is the central component of the sorting and assembly machinery (SAM) in *Saccharomyces cerevisiae* (Kozjak et al., 2003; Paschen et al., 2003). Sam50, Sam35, and Sam37 are accessory subunits of SAM, which insert β-barrel proteins into the mitochondrial OM (Walther et al., 2009).

This study provides evidence that the CRUMPLED LEAF (CRL) protein associates with OEP80 to regulate the formation of its complex. *CRL* encodes a conserved intrinsic membrane protein with a putative transmembrane domain that is localized in the OEM (Asano et al., 2004). The *crl* mutant exhibits multiple defects in plastid division, leaf shape and color, cell division and differentiation, suppression of spontaneous light-intensity-dependent localized cell death (LCD), and regulation of nuclear-encoded stress-related genes (Asano et al., 2004; Hudik et al., 2014; Šimková et al., 2012). The mechanism of the *crl* mutation's effect on plant morphology, plant

growth, nuclear gene regulation, and plastid division is largely unknown. However, recent

reports indicate that gain-of-function mutations of *TIC236* suppress multiple *crl* defects

(Fang et al., 2022). Since CRL interacts with translocon components and TIC236 is

involved in the import of plastid proteins as the physical link between TOC and TIC, it

has been proposed that CRL is a component of TOC (Chen et al., 2018; Fang et al.,

2022). These findings suggest that multiple *crl* defects may be due to plastid protein

deficiency (Fang et al., 2022).

Here, we demonstrate that amino acid substitutions in OEP80 suppressed crl

multiple defects of *crl* and that CRL interacts with OEP80 to aid in the formation of its

complex. Since OEP80 is a paralog of TOC75 (Hsu and Inoue, 2009), CRL may aid in

the formation of both the OEP80 and TOC complexes by interacting with TOC75. In

addition to the suppression of plastid protein import (Fang et al., 2022), our findings

suggest that the decreased efficiency of OEM integration of β-barrel proteins may be a

cause of multiple *crl* defects.

Results

Identification of suppressors of the *crl-2* mutant

Through the use of an ethyl methanesulfonate (EMS) mutagenesis screen for suppressors of the *crl-2* mutant encoding the CRL protein with amino acid substitution, we isolated two semidominant suppressor mutants, designated S1-9 and S2-6. S2-6 plants with expanded green leaves and smooth margins were indistinguishable from wild-type (WT) plants, thus rescuing the growth retardation and pale green phenotype of *crl-2* (Figure 1A). The fresh weight of the S2-6 seedlings was consistently greater than that of *crl-2* and equal to that of the WT (Figure 1B). Concerning chloroplasts in mesophyll cells, both chloroplast chlorophyll content and plastid division were fully restored in S2-6 (Figure 1A, C, D). Finally, spontaneous cell death observed in *crl-2* was completely eliminated in S2-6 (Figure 1A, E). We performed RNA sequencing to examine the effect of the level of suppressor mutation on gene expression in *crl-2*. Cluster analysis revealed that the WT and S2-6 samples clustered and were significantly different from the *crl-2* samples (Figure 1F). These results demonstrate that S2-6 suppresses pleiotropic phenotypic defects in *crl-2* nearly completely.

The other suppressor mutant, S1-9, exhibited a phenotype similar to S2-6. It inhibited the pale green color and growth retardation of *crl-2* (Supplemental Figures S1A, B). However, the size and number of chloroplasts in the mesophyll cells of S1-9 were intermediate between those of *crl-2* and WT (Supplemental Figure S1A, C), indicating that the suppressor mutation in S1-9 partially suppresses the *crl-2* phenotype. Therefore, we conclude that the suppressor mutation in S1-9 is less potent than in S2-6.

Amino acid substitutions in the β-barrel domain of OEP80 are responsible for the rescue of phenotypic defects in crl-2

The causal mutations of S1-9 and S2-6 were identified in the upper arm of chromosome 5, where both S1-9 and S2-6 harbored a missense mutation in the OEP80 coding region: a C-to-T mutation at position 1271 (Ala-to-Val substitution at position 424, OEP80 ^{A424V}) in S1-9 and a G-to-A mutation at position 1547 (Gly-to-Glu substitution at position 516, OEP80 ^{G516E}) in S2-6. OEP80 is predicted to have a transit peptide, 16-stranded β-barrel domain, and three-repeated polypeptide transport-associated (POTRA) domain facing the IMS (Day et al., 2014; Day et al., 2019; Gross et al., 2020). Amino acid substitutions A424V in S1-9 and G516E in S2-6 occurred on the putative second and seventh β-strands of the β-barrel domain, respectively (Supplemental Figure S2).

In order to determine whether *OEP80* was the causal gene for *crl* suppression, the genomic sequences of *OEP80* S1-9, S2-6, and WT were amplified by PCR, and each was introduced into *crl-2* by *Agrobacterium*-mediated transformation. The *OEP80* of S2-6 suppressed growth retardation and chloroplast division defects of *crl-2* (*genomic oep80^{G516E}*), while WT did not (Figure 2). (*genomic OEP80^{WT}* in Figure 2). When comparing *crl-2* with the *oep80^{G516E}* transgene to S2-6, the restoration of chloroplast division in *crl-2* was less pronounced (Figure 2B). This is consistent with the observation that suppressor mutations in S2-6 are semidominant. In fact, chloroplast division was fully rescued in a double mutant with an *OEP80* null allele and the *oep80^{G516E}* transgene but not in a double mutant with the *OEP80^{WT}* transgene (Figure 2B). In the absence of WT *OEP80*, *oep80^{G516E}* fully suppresses the *crl-2* mutation. Furthermore, the genomic fragments of *OEP80* from S1-9 inhibited both *crl-2* growth retardation and chloroplast division defects (*genomic OEP80^{A424V}* in Supplemental Figure S3). These complementation analyses confirm that *OEP80* is the causal gene for the suppression of *crl-2* in S1-9 and S2-6.

The *oep80^{G516E}* single mutant exhibits no apparent phenotypic defects

Since the degree of suppression of S2-6 was greater than that of S1-9, *oep80^{G516E}* was chosen for further examination. To examine the

physiological and morphological properties of *oep80^{G516E}*, the phenotype of a single *oep80^{G516E}* mutant was examined. Plant growth, leaf shape, leaf color, and chloroplast size of *oep80^{G516E}* single mutants were visually indistinguishable from those of WT plants under normal growth conditions (Supplemental Figure S4A, B). Quantitative analysis revealed that the fresh weight and chlorophyll concentration of the *oep80^{G516E}* single mutant were comparable to those of the WT (Supplemental Figure S4C, D). According to these findings, *oep80^{G516E}* has no discernible effect on plant growth, development, or chloroplast division.

Immunoblot analysis revealed that the amount of OEP80 protein in the *oep80^{G516E}* single mutant was lower than that of the WT, while the amounts of the OEM proteins TOC75-III (TOC75 of *A. thaliana*, hereafter TOC75) and TOC33 were unchanged (Supplemental Figure S5A, B). This reduction in OEP80 accumulation is likely not responsible for *crl-2* suppression, as *oep80-2/OEP80*, which led to a greater reduction in OEP80 accumulation than the *oep80^{G516E}* mutation (Supplemental Figure S5A, B), did not rescue the above-ground appearance and chloroplast size of *crl-2* (Supplemental Figure S5C, D).

oep80^{G516E} suppressed pleiotropic defects of the *CRL* null mutant

The *oep80^{G516E}* mutation resulted in slightly elevated CRL protein levels in both WT and *crl-2*, as demonstrated by immunoblot analysis with an anti-CRL antibody (Supplemental Figure S6). To determine whether this increase in CRL accumulation accounts for the phenotypic rescue of *crl-2*, we examined whether the *oep80^{G516E}* mutation rescues *crl-1*, a null *CRL* allele. Thus, by genetic crossing, we introduced *oep80^{G516E}* into *crl-1* and observed the phenotype (i.e., growth, leaf shape, color, chloroplast division, and spontaneous LCD) of the double mutant (Figure 3). Our analyses revealed that *oep80^{G516E}* completely suppressed the observed *crl-1* defects, including plant size, fresh weight, color, and spontaneous LCD (Figure 3A, D, E, G), while the size and number in mesophyll cells were only partially restored (Figure 3B, F). Thus, increased CRL accumulation due to the *oep80^{G516E}* mutation is not essential for

crl suppression, indicating that the OEP80[G516E] mutation must act through a different

mechanism.

A424V and G516E substitutions likely affect β-barrel conformation of OEP80

Recent advances in AlphaFold2 have made precise structural modeling

possible (Jumper et al., 2021). Therefore, we examined the 3D structure of *A. thaliana*

OEP80 obtained from the AlphaFold Protein Structure Database. The confidence score

(pLDDT) was quite high in the POTRA and β-barrel domains of OEP80, indicating that

these regions were accurately predicted (blue color in Supplemental Figure S7A). In

contrast, the regions preceding the POTRA domain appeared disorganized because

their score was low (red color in Supplemental Figure S7A). Consistent with the C-

terminal homology modeling of OEP80 (Day et al., 2014), AlphaFold2 predicted a 16-

stranded β-barrel, with A424 and G516 located on the second and seventh β-strands,

respectively (Supplemental Figure S2). According to the Alphafold2 description, when

the main chain of a protein is predicted with high confidence, so is the rotamer of amino

acid side chains (Jumper et al., 2021). Consequently, we hypothesized that the side

chain rotamer in the β-barrel domain of OEP80 was also highly accurate. To evaluate

the conformational effects of amino acid substitutions (A424V and G516E) on OEP80,

we performed *in silico* mutagenesis. All possible rotamers of A424V and G516E

exhibited unfavorable atomic interactions with their adjacent residues. In a

representative rotamer, the substitution of A424 in V results in two clashes, whereas

the substitution of G516 into E results in seven clashes on the inner side of the β-barrel

(Supplemental Figure S7B, B', C, C'). Therefore, it is probable that substitutions of

A424V and G516E alter the β-barrel structure of OEP80.

The complex formation of OEP80 to form complexes is impaired in *crl-2*

In silico analysis suggests that A424V and G516E mutations alter the

conformation of OEP80. Subsequently, using chloroplast proteins and two-dimensional

gel electrophoresis with Blue Native-PAGE and SDS-PAGE (2D-BN/SDS-PAGE), we compared to OEP80's capacity to form complexes between WT, *crl-2*, and S2-6. In the WT, OEP80 is most abundant in a complex of approximately 200 kDa (Figure 4A; *** in Col-0), while it is less abundant in complexes of ca. 70 kDa to ca.150 kDa and ca. 242 to ca.480 kDa (Figure 4A; * and ** in Col-0). This result is similar to that of Gross et al. (2021). The 200 kDa complex was significantly reduced in *crl-2* (Figure 4A; *crl-2*), and no other complexes were detected (Figure 4A; *crl-2*). Compared to *crl-2*, the number of OEP80-containing complexes in S2-6 increased but did not reach WT levels (Figure 4A; S2-6). These findings indicate that the ability of OEP80 to form complexes is perturbed in *crl-2* and partially restored in S2-6. Since OEP80 accumulation is not altered in *crl-2* (Figure 4B), the decrease in OEP80-containing complexes may indicate that the majority of OEP80 becomes digitonin-insoluble in *crl-2*.

CRL likely interacts with OEP80 complexes *in vivo*

We sought to examine the possibility that CRL was one of the complex components. First, we analyzed the interaction between CRL and OEP80 by co-immunoprecipitation using chloroplasts from plants expressing functional CRL-GFP or plastid-targeted GFP (pt-GFP). OEP80 was co-immunoprecipitated with CRL-GFP but not with pt-GFP when solubilized chloroplast proteins were immunoprecipitated with an anti-GFP antibody (Figure 5A). Neither CRL-GFP nor pt-GFP co-immunoprecipitated with the IEM protein TIC40 (Figure 5A). These results suggest that CRL and OEP80 interact *in vivo*.

Next, the molecular weight of the CRL-containing complexes was analyzed by 2D-BN/SDS-PAGE using chloroplasts of WT plants. Most of the CRL-containing complexes were detected at molecular weights below 146 kDa (Figure 5B), including the molecular weight of one of the less abundant complexes (* in Figure 4A), while others were detected between 146 and 242 kDa (Figure 5B, solid line), where the most abundant complex was detected (*** in Figure 4A). Additionally, a weak CRL signal

was detected between ca. 242 kDa and ca. 480 kDa (Figure 5B, dotted line), where a

weak OEP80 signal was observed (Figure 4A, **). These results suggest that a portion

of the CRL pool is associated with complexes, not the entire pool. Our data suggest

that CRL is an OEP80 associating factor and is involved in the formation of OEP80

complexes. By altering the conformation of the β-barrel domain, the two amino acid

substitutions in OEP80^{G516E} and OEP80^{A424V} can facilitate the formation of the OEP80

complex in the absence of CRL.

Discussion

We have identified two allelic mutations of *OEP80*, *oep80^A424V,* and *oep80^G516E* (Figure 2, Supplemental Figure S3), which suppress pleiotropic defects of the hypomorphic mutant (Figure 1, Supplemental Figure S1) and of the null mutant, *crl-1* (Figure 3). Moreover*, crl-2* impaired the ability of OEP80 to form complexes, indicating that CRL is involved in the formation of complexes (Figure 4A). Furthermore, CRL-GFP interacted with OEP80 *in vivo,* and a portion of CRL comigrated with OEP80 (Figure 5). From these findings, CRL interacts with OEP80-containing complexes and promotes their formation. Recently, CRL has been shown to interact with TOC75 and other TOC components *in vivo* using plants expressing functional CRL-GFP under the 35S promoter (Fang et al., 2022). Since TOC75 and OEP80 are members of the Omp85 protein family (Hsu and Inoue, 2009), CRL may facilitate not only the formation of OEP80-containing complexes but also the formation of the TOC complex by interacting with OEP80 and TOC75. Because endogenous CRL was not detected in the molecular mass range of 880 kDa to 1.3 MDa (Figure 5B, Chen, and Li, 2017), the interaction between CRL and TOC75-III may be transient or very weak.

OEP80 is involved in the incorporation of plastidic β-barrel proteins into OEM (Gross et al., 2021). Therefore, CRL may participate in this process in conjunction with OEP80. However, mutants of loss of function of *OEP80* exhibit embryonic lethality (Patel et al., 2008), while the *crl-1* mutant is not (Asano et al., 2004). By preventing the formation of OEP80-containing complexes, the efficiency of membrane integration of β-barrel proteins into OEM may be partially but not completely compromised in *crl*. According to *in silico* mutagenesis of the modeled structure of OEP80, the substitutions for OEP80^A424V and OEP80^G516E slightly alter the β-barrel structure (Supplemental Figure S7B, B', C, C'). Therefore, the structural change in the β-barrel domain of OEP80 in the S1-9 and S2-6 mutants probably restores the ability to form complexes in the absence of CRL, resulting in the restoration of β-barrel protein to OEM. Compared to Gram-negative bacteria and mitochondria, the mechanism of assembly

of plastidic β-barrel proteins in OEM is not well understood. Furthermore, to date, no components of OEP80-containing complexes have been identified. To our knowledge, CRL is the first factor that has been experimentally implicated in the integration of the OEP80-dependent β-barrel membrane. Additional analysis of OEP80 and the other β-barrel proteins in *crl* would aid in understanding the OEM integration mechanism.

CRL has been implicated in the importation of plastid protein (Fang et al., 2022). Consequently, pleiotropic defects of *crl* may be a result of the additive or synergistic effect of the suppression of plastid protein import and the partial dysfunction of the integration of plastidic β-barrel proteins in the OEM. DEK5, a TamB ortholog in maize, has been implicated in the integration of OEM-localized β-barrel proteins such as OEP80 and TOC75 (Zhang et al., 2019). A *dek5* mutant with a weak allele displays *crl*-like phenotypes, including stunted growth, enlarged chloroplasts with well-developed thylakoid membranes, and decreased accumulation of OEM proteins, including OEP80, TOC75, and TOC159 (Zhang et al., 2019). These analyses of the *dek5* mutant confirm the importance of β-barrel protein integration in OEM and plastid protein import for plant development and chloroplast division.

The mutant *tic236-2* shows a *crl*-like phenotype and decreased levels of plastid proteins (Fang et al., 2022), whereas the *Arabidopsis oep80* RNAi line does not (Huang et al., 2011). This may indicate that *crl* pleiotropic defects are mainly attributable to a decrease in the import of plastid proteins and that OEP80 dysfunction of OEP80 has a minor effect on these defects. This is supported by the suppression of *crl* by gain-of-function mutations in *TIC236* (Fang et al., 2022). However, no plastid protein import mutant other than *tic236-2* exhibits a *crl*-like phenotype, and the phenotype of *tic236-2* is not identical to that of *crl*. In *tic236-2*, not all mesophyll cells exhibit chloroplast division defects, unlike *crl* (Asano et al., 2004; Fang et al., 2022). The leaf lamina of *tic236-2* is similar but not identical to that of *crl* (Asano et al., 2004; Chen et al., 2018). Additionally, the import of plastid proteins would not be severely compromised in *crl* due to the fact that the amounts of TOC75 and OEP80 were not altered in *crl-2* (Figure 4B), and a

stroma-targeting YFP protein is normally imported into chloroplasts in *crl-1* (Chen et al.,

2009). Therefore, we propose that in addition to suppression of plastid protein import

(Fang et al., 2022), partial dysfunction of the integration of plastidic β-barrel proteins in

OEM, as described in this study, is responsible for pleiotropic defects in *crl*. Quantitative

analysis of plastid protein import and examination of membrane localization of β-barrel

proteins in OEM in *crl* will be necessary to determine the contribution of OEP80 and

plastid protein import to *crl* pleiotropic defects.

Materials and methods

Plant materials and growth conditions

Previously, techniques for seed sterilization and growth in soil and Murashige and Skoog (MS) medium were outlined (Chen et al., 2009). All experiments were carried out in a Columbia (Col) background using *A. thaliana*. Previously, *crl-2* (Šimková et al., 2012), *crl-1* (Asano et al., 2004), *crl-1* expressing *35S::CRL-sGFP(S65T)* (*CRL-GFP*) (Asano et al., 2004) and Col expressing *35S::pt-sGFP(S65T)* (*pt-GFP*) (Niwa et al., 1999) described o*ep80-2* (GK 429H12) was described in Patel et al., (2008) and obtained from the Arabidopsis Biological Resource Center. Before analysis, the suppressor lines S1-9 and S2-6 were isolated from *crl-2* that had been mutagenized with EMS and backcrossed three times with *crl-2*. From the original S2-6, *oep80*^{G516E} *crl-1* was generated by crossing with the *crl-1* heterozygote. The *oep80*^{G516E} single mutant was produced by crossing the original S2-6 with Col-0. Additionally, *crl-2 oep80-2* was produced by crossing *crl-2* with the *oep80-2* hemizygous line. The genotyping primers are listed in Supplemental Table S1.

Through *Agrobacterium*-mediated transformation, stable transgenic plants were produced (Clough and Bent, 1998)*.* Briefly, genomic fragments that encompass OEP80 and its putative promoter region (1 kb upstream of the translation initiation site to the end of the annotated 3'UTR) were amplified from genomic DNA extracted from S2-6, S1-9, or Col-0. The amplification primers are listed in Supplemental Table S1 of the supplementary materials. Subsequently, using the TA cloning kit, amplified PCR products were cloned into the pCR8 vector (Thermo Fisher Scientific K.K. Tokyo, Japan). After confirming the nucleotide sequences, genomic fragments were transferred to the binary vector by the LR recombination reaction (Nakagawa et al., 2007). The resulting binary vectors were electroporated in *Agrobacterium tumefaciens* (GV3101) and then introduced to *crl-2* plants using the floral dip technique (Clough and Bent, 1998). The transgenic *crl-2* plants were evaluated in an MS medium containing 15 µg ml^{-1} of hygromycin B. Several lines derived from independent T1 individuals were obtained, and two or three lines were used for subsequent analyses.

EMS mutagenesis and screening

The *crl-2* seeds (200 mg equivalent to ~10,000 seeds) were incubated in 100 ml of 0.3% EMS for 9 h at room temperature (M1 seeds). After being washed six times for 1 min with 100 ml of deionized water and three times for 30 min with 500 ml of deionized water, the seeds were separated into 10 pools and planted in the soil. From the 10 pools, a total of 7,050 M2 lines (self-pollinated M1 offspring) were obtained. In MS medium, M2 and M3 seeds were planted, and approximately 700 individuals were evaluated based on their appearance and mesophyll chloroplasts.

Microscopy

AxioCam MRm (Carl Zeiss AG, Jena, Thüringen, German) was used with an epifluorescent microscope (AxioImager M1; Carl Zeiss AG) or a confocal microscope to capture images of chlorophyll autofluorescence (FV10i, Olympus, Tokyo, Japan). For quantification, 14-day seedling cotyledons were fixed overnight at 4°C in 4% (w/v) paraformaldehyde in phosphate-buffered saline (PBS; 8.1 mM Na_2HPO_4, 1.5 mM KH_2PO_4, 137 mM NaCl, 2.7 mM KCl). More than 2 days were spent immersing the fixed tissues in a solution of 0.1 M EDTA (pH 9.0) at 4°C. A confocal laser microscope was used to acquire bright-field autofluorescent images of mesophyll cells (FV10i, Olympus). The images were reconstructed using Fiji (https://fiji.sc), and the number of mesophyll chloroplasts per 5000 μm^2 mesophyll cell surface was determined.

Trypan blue staining

Trypan blue staining was used to detect cell death in accordance with Weigel and Glazebrook (2002), with modifications. Briefly, the cotyledons were submerged in the staining solution (0.1 g trypan blue, 10 ml of lactic acid, 10 ml of glycerol, 10 ml of phenol, 10 ml of water, 80 ml of ethanol) and boiled at 90°C for 3 min. The samples were then allowed to stand at room temperature for 6 h to ensure full penetration of the stain into the

cotyledons. The cotyledons were next covered with a chloral hydrate solution (2.5 g ml^{-1} chloral hydrate) at room temperature and incubated for 18 h by shaking. Finally, cotyledons were transferred to a 70% glycerol solution and photographed using a stereomicroscope. The LCD area was measured with Fiji. Briefly, raw RGB color images were converted to 8-bit grayscale images. The brightness threshold was set between the LCD and the vasculature regions.

Measurement of chlorophyll content

The 14-day seedlings were submerged in N,N-dimethylformamide for 18 h at 4°C and kept in the dark to extract chlorophyll. Chlorophyll content was calculated according to the formula (Chl a + b (nmol/ml) = 19.43A_{648} + 8.05A_{663}) (Porra et al., 1989) and normalized per fresh weight.

RNA sequencing

Total RNA was extracted from 12-day-old plantlets grown *in vitro* using the RNA Plus Kit (Macherey-Nagel GmbH & Co. KG, Düren, NRW, Germany) following the manufacturer's instructions. Libraries were prepared with 2 µg of RNA using the NEBNext® Ultra™ II Directional RNA Library Prep Kit for Illumina, following the manufacturer's instructions. The sequencing of libraries was performed on an Illumina NextSeq® 500 (Illumina, San Diego, CA, USA). Single-end sequencing of RNA sequence samples was trimmed using Trimmomatic-0.38 (Bolger et al., 2014) with the parameters: Minimum length 30 bp; Mean Phred quality score greater than 30; Removal of leading and trailing bases with base quality of <5. Bowtie2 aligner (Langmead and Salzberg, 2012) was used to map the assembly of the TAIR10 genome. The raw read counts were then extracted using feature counts (Liao et al., 2014) based on the gene annotations in the Araport11 annotations. Finally, we use DESeq2 (Love et al., 2014) to identify differentially expressed genes. Genes with read counts ≥ 20 were considered for differential analysis. Clustering and heatmap generation were performed using Pheatmap v1.0.12 (KOLDE,

R. pheatmap v1. 0.12. 2019) with parameter cutree_rows = 8 and scale = 'row.' The

RNAseq data reported in this article have been deposited in the NCBI GEO database

under accession code GSE193615.

Whole-genome sequencing

Genomic DNAs were extracted from 20 homozygotes of the F_2 population after S2-6

were crossed once with *crl-2,* and from 20, homozygotes of the F_3 population after S1-9

were backcrossed once with *crl-2*. DNA extracted DNA was fragmented by sonication

(S220 from Covaris) and then converted to libraries using TruSeq DNA Sample Prep Kits

(Illumina) according to the manufacturer's protocol. The libraries were sequenced using

NextSeq500 (Illumina). The bcl files were converted to fastq files using bcl2fastq

(Illumina). Additionally, fastq files were analyzed by Mitsucal (Suzuki et al., 2018).

Structure modeling and analysis

The modeled structure of OEP80 (AlphaFold identifier AF-Q9C5J8-F1) was obtained from

the AlphaFold Protein Structure Database (https://alphafold.ebi.ac.uk), which was updated

on 1 July 2021 with AlphaFold v2.0 (Jumper et al., 2021). Visualization, mutagenesis, and

clash analysis of the predicted OEP80 structure were performed using UCSF ChimeraX

according to the user guide (https://www.rbvi.ucsf.edu/chimerax/, Goddard et al., 2018;

Pettersen et al., 2021). *In silico* mutagenesis was based on the Dunbrack rotamer library

(Shapovalov and Dunbrack, 2011). A clash was defined as an interaction between atoms

with van der Waals overlap $\geqq 0.6$ Å.

Protein extraction and immunoblotting

Fourteen-day-old seedlings homogenized in liquid nitrogen were suspended in a protein

extraction buffer (2% SDS, 10% sucrose, 56 mM Na_2CO_3, 2 mM EDTA, pH 8.0) and

boiled at 95°C for 5 min. The supernatant was collected after two centrifugations (14,000

g, 10 min). Protein concentration was measured using a BCA assay kit (Takara Bio Inc.,

Kusatsu, Siga, Japan). Twenty micrograms (μg) of proteins were combined with 2-

mercaptoethanol and bromophenol blue (to final concentrations of 5% and 0.0093%, respectively) and boiled at 95°C for 5 min. The samples were separated by SDS-PAGE and immunoblotted for analysis. Next, using the semi-dry method, the proteins on the gel were transferred to a PVDF membrane. The membrane was then immersed for 1 h in the blocking solution (1% skim milk, 0.1% Tween 20 / PBS). Overnight incubation with the primary antibody was performed at 4°C. The membrane was washed with PBS, and then the secondary antibody was incubated for 90 min at room temperature. Subsequently, the membrane was washed with PBS, and ECL prime solution (Cytiva, Marlborough, MA, USA) was added. A CCD imager (LAS4000 mini, Cytiva) was used to record chemiluminescence. For the detection of OEP80, an anti-OEP80 antibody (PHY0814A, PhytoAB Inc., San Jose, CA, USA) was used. The antibody against TOC75-III was a gift from Dr. Inaba. Antibody against TOC33 was a gift from Dr. Nakai. The secondary antibody was Donkey anti-rabbit IgG-HRP (NA934, GE Healthcare, Chicago, IL, USA), which was detected using ECL select (Cytiva). Each antibody was diluted with a blocking buffer.

Isolation of intact chloroplasts

Three-week-old plants grown on MS medium were cut into pieces, treated with an enzyme solution (1% Cellulase "Onozuka" R-10, 0.25% Macerozyme "Onozuka" R-10, 400 mM Mannitol, 20 mM MES-KOH pH5.7, 20 mM KCl, 10 mM $CaCl_2$, 0.1% BSA) and degassed. After a 2 h incubation, the protoplasts were pelleted by centrifugation (100 g, 4°C, 5 min). After the elimination of plant tissues through Miracloth filtration, protoplasts were suspended in a buffer (300 mM Sorbitol, 20 mM Tricine-KOH, pH8.4, 5 mM EDTA, 5 mM EGTA, 10 mM $NaHCO_3$, 0.1% BSA) and disrupted by filtrating through a 20 µm nylon mesh using a syringe, and the solution was placed on 40% / 85% step gradients. The lower layer was collected after centrifugation (2,500 g, 4°C, 10 min). After washing with two volumes of suspension buffer (330 mM Sorbitol, 50 mM HEPES-KOH pH 8.0), intact chloroplasts were pelleted by centrifugation (700 g, 4°C, 5 min). The pellet was

resuspended in a suspension buffer and used for further analysis. The chlorophyll

content was calculated using the formula: Chl (mg ml^{-1}) = 2.899A_{652}.

In vivo immunoprecipitation assay

The GFP polyclonal antibody (Anti-GFP pAb(Medical & Biological Laboratories Co., Ltd.,

Tokyo, Japan) was conjugated with M280-tosyl activated Dynabeads (Thermo Fisher

Scientific K.K.) according to the manufacturer's instructions. Chloroplasts (approximately

500–700 µg chlorophyll content) were resuspended in 1 ml of solubilization buffer (50 mM

HEPES-KOH pH7.3, 1% n-dodecyl-β-D-maltoside) and solubilized on ice for 5 min. A small

amount of supernatant was collected as input after centrifugation (14,000 _g_, 4°C, 5 min).

Combining the remaining supernatant with antibody-coupled Dynabeads. The binding was

achieved by rotating the mixture at room temperature for 1 h. Immunoprecipitants were

eluted with 16 µl of 0.1 M glycine-HCl pH 2.8 and neutralized with 4 µl of 1.0 M Tris-HCl

pH 8.0 after four washes of the beads with 1 ml of solubilization buffer. After adding 4 ×

sample buffer (250 mM Tris-HCl pH 6.8, 8% SDS, 20% sucrose) to the elution and input

samples, they were boiled at 95°C for 5 min. As described previously, the samples were

separated by SDS-PAGE and analyzed by immunoblotting. For TIC40 detection, the anti-

TIC40 antibody (AS10709, Agrisera, Vännäs, Västerbotten, Sweden) was used.

Two-dimensional gel analysis (2D-BN/SDS-PAGE)

Chloroplasts containing 30 µl of 1 mg ml^{-1} chlorophyll were resuspended in 2x

solubilization buffer (100 mM Bis-Tris, 20% glycerol, 100 mM NaCl, 2% digitonin) and

stored at 4°C for 30 min. After ultracentrifugation (100,000 _g_, 4°C, 30 min), 30 µl of the

supernatant was added to 4 × native sample buffer (200 mM Bis-Tris, 40% glycerol, 200

mM NaCl, 0.5% CBB-G250, 50 mM aminocaproic acid). The samples were separated

using a 3%–12% concentration linear gradient acrylamide gel (Blue Native-PAGE). The

lanes were cut and incubated at 37°C for 30 min in 50 ml of denaturing buffer (3.3% SDS,

4% 2-mercaptoethanol, and 65 mM Tris-HCl, pH 6.8) with rotation. As described previously,

each gel fragment was separated by SDS-PAGE and analyzed by immunoblotting.

Antibody against OEP80 was able to detect OEP80 (PHY2423A, PhytoAB Inc.). Earlier,

the anti-CRL antibody was described (Asano et al., 2004). The secondary antibody was

Donkey anti-rabbit IgG-HRP (NA934, GE HealthCare), and it was detected using ECL

select (Cytiva).

Acknowledgments

We are grateful to Dr. Takehito Inaba for providing the anti-TOC75-III antibody, Dr. Masato Nakai for providing the anti-TOC33 antibody, and Dr. Niwa for providing *35S::pt-sGFP(S65T)*. We thank Dr. Tetsuya Higashiyama, Dr. Wataru Sakamoto, Dr. Akira Kanamori, Dr. Toshinori Kinoshita, Dr. Shin Sugiyama, Dr. Shin Takagi, and Dr. Yoshimasa Yagi for the helpful discussion. The authors also thank Tomotaka Itaya, Ayami Furuta, Tomomi Shinagawa, Sae Miyazaki, Aya Murata, Kimika Takeuchi, and Ryohei Seta for technical support. This collaboration was supported by a CNRS PICS-ChloroCycle grant.

Authors' contributions

Y.Y. designed the research. Y.Y. and R.Y wrote the manuscript. R.Y and S.M. isolated the suppressor lines and performed the experiments. C.R did the RNA extraction and library construction for RNAseq. D.L. did the nucleotide sequencing of the RNAseq. T.S., D.V., S.S., C.R. and M.B. performed bioinformatic analysis. All authors have read and approved the final manuscript.

Funding

This work was supported by the Japan Society for the Promotion of Science (No. 26440137 to Y.Y.) and the Joint Usage/Research Center, Institute of Plant Science and Resources, Okayama University.

Disclosures

The authors have no conflicts of interest to declare.

References

Asano T, Yoshioka Y, Kurei S, Sakamoto W, Sodmergen, Machida Y (2004) A mutation of the *CRUMPLED LEAF* gene that encodes a protein localized in the outer envelope membrane of plastids affects the pattern of cell division, cell differentiation, and plastid division in *Arabidopsis*. Plant J **38:** 448–459

Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics **30:** 2114–2120

Bölter B, Soll J, Hill K, Hemmler R, Wagner R (1999) A rectifying ATP-regulated solute channel in the chloroplastic outer envelope from pea. EMBO J **18:** 5505–5516

Bölter B, Soll J (2016) Once upon a Time – Chloroplast Protein Import Research from Infancy to Future Challenges. Molecular Plant **9:** 798–812

Chen Y, Asano T, Fujiwara MT, Yoshida S, Machida Y, Yoshioka Y (2009) Plant Cells Without Detectable Plastids are Generated in the *crumpled leaf* Mutant of *Arabidopsis thaliana*. Plant Cell Physiol **50:** 956–969

Chen YL, Chen LJ, Chu CC, Huang PK, Wen JR, Li H (2018) TIC236 links the outer and inner membrane translocons of the chloroplast. Nature **564:** 125–129

Chen L, Li H. (2017) Stable megadalton TOC-TIC supercomplexes as major mediators of protein import into chloroplasts. Plant J **92:** 178-188

Clough SJ, Bent AF (1998) Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. Plant J **16:** 735–743

Day PM, Potter D, Inoue K (2014) Evolution and targeting of Omp85 homologs in the

chloroplast outer envelope membrane. Front Plant Sci **5:** 535

Day PM, Inoue K, Theg SM (2019) Chloroplast Outer Membrane β-Barrel Proteins

Use Components of the General Import Apparatus. Plant Cell **31:** 1845–1855

Fang J, Li B, Chen Lj, Dogra V, Luo S, Wu W, Wang P, Hwang I, Li Hm, Kim C

(2022) *TIC236* gain-of-function mutations unveil the link between plastid division and

plastid protein import. Proc Natl Acad Sci USA **119:** e2123353119

Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, Ferrin TE

(2018) UCSF ChimeraX: Meeting modern challenges in visualization and analysis. Protein

Sci **27:** 14–25

Goetze TA, Philippar K, Ilkavets I, Soll J, Wagner R (2006) OEP37 Is a New Member

of the Chloroplast Outer Membrane Ion Channels. J Biol Chem **281:** 17989–17998

Gross LE, Spies N, Simm S, Schleiff E (2020) Toc75-V/OEP80 is processed during

translocation into chloroplasts, and the membrane-embedded form exposes its POTRA

domain to the intermembrane space. FEBS Open Bio **10:** 444–454

Gross LE, Klinger A, Spies N, Ernst T, Flinner N, Simm S, Ladig R, Bodensohn U,

Schleiff E (2021) Insertion of plastidic β-barrel proteins into the outer envelopes of

plastids involves an intermembrane space intermediate formed with Toc75-V/OEP80.

Plant Cell **33:** 1657–1681

Hagan CL, Silhavy TJ, Kahne D (2011) β-Barrel Membrane Protein Assembly by the

Bam Complex. Annu Rev Biochem **80:** 189-210

Harsman A, Schock A, Hemmis B, Wahl V, Jeshen I, Bartsch P, Schlereth A, Pertl-Obermeyer H, Goetze TA, Soll J, et al. (2016) OEP40, a Regulated Glucose-permeable β-Barrel Solute Channel in the Chloroplast Outer Envelope Membrane. J Biol Chem **291:** 17848–17860

Hsu SC, Inoue K (2009) Two evolutionarily conserved essential β-barrel proteins in the chloroplast outer envelope membrane. Biosci Trends **3:** 168–178

Huang W, Ling Q, Bédard J, Lilley K, Jarvis P (2011) In Vivo Analyses of the Roles of Essential Omp85-Related Proteins in the Chloroplast Outer Envelope Membrane. Plant Physiol **157:** 147–159

Hudik E, Yoshioka Y, Domenichini S, Bourge M, Soubigout-Taconnat L, Mazubert, C, Yi D, Bujaldon S, Hayashi H, De Veylder L, et al. (2014) Chloroplast dysfunction causes multiple defects in cell cycle progression in the Arabidopsis *crumpled leaf* mutant. Plant Physiol **166:** 152–167

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Zidek A, Potapenko A, et al. (2021) Highly accurate protein structure prediction with AlphaFold. Nature **596:** 583–589

Kozjak V, Wiedemann N, Milenkovic D, Lohaus C, Meyer HE, Guiard B, Meisinger C, Pfanner N (2003) An essential role of Sam50 in the protein sorting and assembly machinery of the mitochondrial outer membrane. J Biol Chem **278:** 48520-48523

Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. Nat Methods **9:** 357–359

Leister D (2003) Chloroplast research in the genome age. Trend Genet **19:** 47-56

Liao Y, Smyth GK, Shi W (2014) featureCounts: an efficient general purpose program

for assigning sequence reads to genomic features. Bioinformatics **30:** 923–930

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol **15:** 550

Nakagawa T, Suzuki T, Murata S, Nakamura S, Hino T, Maeo K, Tabata R, Kawai

T, Tanaka K, Niwa Y, et al. (2007) Improved Gateway binary vectors: high-

performance vectors for creation of fusion constructs in transgenic analysis of plants.

Biosci Biotechnol Biochem **71:** 2095–2100

Niwa Y, Hirano T, Yoshimoto K, Shimizu M, Kobayashi, H. (1999) Non-invasive

quantitative detection and applications of non-toxic, S65T-type green fluorescent

protein in living plants. Plant J **18:** 455–463

Paschen CA, Waizenegger T, Stan T, Preuss M, Cyrklaff M, Hell K, Rapaport D,

Neupert W (2003) Evolutionary conservation of biogenesis of β-barrel membrane proteins.

Nature **426:** 862-866

Patel R, Hsu SC, Bédard J, Inoue K, Jarvis P (2008) The Omp85-Related Chloroplast

Outer Envelope Protein OEP80 Is Essential for Viability in Arabidopsis. Plant Physiol **148:**

235–245

Pettersen EF, Goddard TD, Huang CC, Meng EC, Couch GS, Croll TI, Morris JH, Ferrin TE (2021) UCSF ChimeraX: Structure visualization for researchers, educators, and developers. Protein Sci **30:** 70–82

Pohlmeyer K, Soll J, Grimm R, Hill K, Wagner R (1998) A high-conductance solute channel in the chloroplastic outer envelope from Pea. Plant Cell **10:** 1207–1216

Porra RJ, Thompson WA, Kriedemann PE (1989) Determination of accurate extinction coefficients and simultaneous equations for assaying chlorophylls a and b extracted with four different solvents: verification of the concentration of chlorophyll standards by atomic absorption spectroscopy. Biochimica Biophysica Acta **975:** 384–394

Selkrig J, Mosbahi K, Webb CT, Belousoff MJ, Perry AJ, Wells TJ, Morris F, Leyton DL, Totsika M, Phan MD, et al. (2012) Discovery of an archetypal protein transport system in bacterial outer membranes. Nat Struct Mol Biol **19:** 506–510

Shapovalov MV, Dunbrack RL (2011) A smoothed backbone-dependent rotamer library for proteins derived from adaptive kernel density estimates and regressions. Structure **19:** 844–858

Šimková K, Kim C, Gacek K, Baruah A, Laloi C, Apel K (2012) The chloroplast division mutant *caa33* of *Arabidopsis thaliana* reveals the crucial impact of chloroplast homeostasis on stress acclimation and retrograde plastid-to-nucleus signaling. Plant J **69:** 701–712

Suzuki T, Kawai T, Takemura S, Nishiwaki M, Suzuki T, Nakamura K, Ishiguro S, Higashiyama T (2018) Development of the Mitsucal computer system to identify causal mutation with a high-throughput sequencer. Plant Reprod **31:** 117–128

Walther DM, Rapaport D, Tommassen J (2009) Biogenesis of β-barrel membrane

proteins in bacteria and eukaryotes: evolutionary conservation and divergence. Cell Mol

Life Sci **66:** 2789–2804

Wang Z, Xu C, Benning C. (2012) TGD4 involved in endoplasmic reticulum-to-

chloroplast lipid trafficking is a phosphatidic acid binding protein. Plant J **70:** 614–623

Weigel D, Glazebrook J (2002) Trypan Blue Stain for Fungi, Oomycetes, and Dead

Plant Cells. *In* Arabidopsis: A Laboratory Manual. Chapter 4. Cold Spring Harbor

Laboratory Press, Cold Spring Harbor, New York, pp 86-87

Zhang J, Wu S, Boehlein SK, McCarty DR, Song G, Walley JW, Myers A, Settles

AM (2019) Maize *defective kernel5* is a bacterial TamB homologue required for

chloroplast envelope biogenesis. J Cell Biol **218:** 2638–2658

Legends to figures

Figure 1 S2-6 suppresses *crl-2*

(A) Phenotypic analysis of S2-6. The overall morphology of 3-week-old plants grown

on soil. Bar = 1 cm. Autofluorescent images of mesophyll chloroplasts in cotyledons of

2-week-old seedlings. Bar = 20 μm. Cell death in cotyledons of 10-day-old seedlings

visualized by trypan blue staining. Bar = 1 mm. (B) Fresh weight of 2-week-old

seedlings (n = 12). (C) Number of chloroplasts per cell (n = 10). Mesophyll cells were

used in the cotyledons of 2-week-old seedlings. (D) Chlorophyll content of 2-week-old

seedlings (n = 12). (E) Percentage of cotyledon area affected by localized cell death in

10-day seedlings (n = 10). (F) Similarity analysis of RNAseq samples. Metagene data

from biological replicates of the three biological replicates were used to calculate the

correlation coefficient based on Euclidean distances, and the samples were clustered

based on their similarity. Asterisks indicate statistical significance (Tukey multiple

comparison test, ****P < 0.001, *P < 0.05, NS, not significant).

Figure 2 *oep80^{G516E}* complements *crl-2*

(A) Two-week-old seedlings grown on MS plates. Bar = 5 mm. (B) Autofluorescent

images of mesophyll chloroplasts in cotyledons of 2-week-old seedlings. Bar = 50 μm.

Figure 3 *oep80^{G516E}* suppresses *crl-1*

(A) Two-week-old seedlings grown on MS plates. Bar = 5 mm. (B) Autofluorescent

images of mesophyll chloroplasts in cotyledons of 2-week-old seedlings. Bar = 50 μm.

(C) Cell death in the cotyledons of 10-day-old seedlings visualized by trypan blue staining.

Bar = 1 mm. (D) Fresh weight of 2-week-old seedlings (n = 15). (E) Chlorophyll content

of 2-week-old seedlings (n = 15). (F) The number of chloroplasts per cell ($n \geqq$ 5).

Mesophyll cells were used in the cotyledons of 2-week-old seedlings. (G) Percentage of

the area of cotyledons affected by localized cell death in 10-day-old seedlings (n = 10).

Asterisks indicate statistical significance (Tukey multiple comparison test, ****P < 0.001,

NS, not significant.

Figure 4 The formation of the OEP80 complex is disturbed in *crl-2* and restored

in S2-6

(A) Immunoblotting of the OEP80 complex. Intact chloroplasts from 3-week-old seedlings

grown on MS plates were solubilized with 1% digitonin, and the supernatants were

separated by 2D-BN/SDS-PAGE. OEP80 was detected by an anti-OEP80 antibody. The

molecular weight of ca. 200 kDa (***), from ca. 70 kDa to 150 kDa (*), and from ca. 242–

480 kDa (**) are indicated. IB, immunoblotting (B) Immunoblot analysis of OEP80 and

TOC75-III. Twenty micrograms (μg) of the total proteins extracted from 2-week-old

seedlings were separated by SDS-PAGE. OEP80 and TOC75-III were detected by anti-

OEP80 and anti-TOC75-III antibodies, respectively.

Figure 5 CRL is suggested to be contained in the OEP80 complex

(A) Immunoblotting of immunoprecipitants with anti-GFP antibody. Intact chloroplasts

from 3-week-old seedlings grown in MS plates were solubilized with 1% n-Dodecyl-β-D-

maltoside (DDM). Immunoprecipitants were separated by SDS-PAGE and detected by

anti-OEP80, anti-TIC40, and anti-GFP antibodies, respectively. *pt-GFP*, *35S::pt-*

sGFP(S65T); *CRL-GFP, 35S::CRL-sGFP(S65T)*; IgG(H), IgG heavy chain; IgG(L), IgG

light chain; IP, immunoprecipitation. (B) Immunoblotting of the CRL complex. Intact

chloroplasts from 3-week-old seedlings grown on MS plates were solubilized with 1%

digitonin, and the supernatants were separated by 2D-BN/SDS-PAGE. CRL was

detected by an anti-CRL antibody. The molecular weight is about 146–242 kDa (solid

line), and from ca. 242–480 kDa (dotted line) are indicated. IB, immunoblotting.

Supplemental Figure S1 S1-9 suppresses *crl-2*

(A) Phenotypic analysis of S1-9. Two-week-old seedlings grown on MS plates. Bar = 5

mm. Autofluorescent images of mesophyll chloroplasts in the mesophyll cell of 2-week-

old seedlings. The margin of the cell was indicated by a dashed line. Bar = 20 μm. (B)

Fresh weight of 2-week-old seedlings (n = 10). (C) A number of chloroplasts per cell (n

$\geqq$ 7). Mesophyll cells were used in the cotyledons of 2-week-old seedlings. The asterisks

represent statistical significance (Tukey multiple comparison test, ***P < 0.001, NS, not

significant).

Supplemental Figure S2 Schematic structure of OEP80

The rectangles in the β-barrel domain represent β-strands. Amino acid substitutions in

S1-9 and S2-6 are indicated.

Supplemental Figure S3 *oep80^{A424V}* complements *crl-2*

(A) Fresh weight of 2-week-old seedlings (n = 10). (B) Number of mesophyll chloroplasts

per cell (n 3). Mesophyll cells were used in the cotyledons of 2-week-old seedlings. The

different letters above the plots represent statistical significance (Tukey multiple

comparison test, P < 0.05).

Supplemental Figure S4 The phenotypes of the *oep80^{G516E}* single mutant

(A) Two-week-old seedlings grown on MS plates. Bar = 5 mm. (B) Autofluorescent

images of mesophyll chloroplasts in cotyledons of 2-week-old seedlings. Bar = 50 μm.

(C) Fresh weight of 2-week-old seedlings (n = 12). (D) Chlorophyll content of 2-week-

old seedlings (n = 12). NS not significant (Student's t-test, P > 0.05).

Supplemental Figure S5 *oep80^{G516E}* reduces its own protein level, but the reduction

of the protein level is insufficient for the suppression of *crl-2*

(A) Immunoblot analysis of OEP80, TOC75-III, and TOC33. Twenty micrograms (μg) of

total proteins extracted from 2-week-old seedlings were separated by SDS-PAGE.

OEP80, TOC75-III and TOC33 were detected by anti-OEP80, anti-TOC75-III and anti-

TOC33 antibodies, respectively. *oep80-2*, mutant null mutant; +/*oep80-2*, heterozygous

for *oep80-2*; *oep80^{G516E}*, single mutant for *oep80^{G516E}*. (B) Protein abundance of OEP80,

TOC75-III, aminocapronic and TOC33. Protein immunoblot intensities were normalized

to those of Col-0. The mean of three biological replicates ± SD was indicated. (C) Two-

week-old seedlings grown in MS plates. Bar = 5 mm. (D) Autofluorescent images of

mesophyll chloroplasts in cotyledons of 2-week-old seedlings. Bar = 50 μm.

Supplemental Figure S6 *oep80^{G516E}* slightly enhances the protein level of CRL^{G31D}

An immunoblot analysis of CRL and CRL^{G31D} encoded by *crl-2* in *oep80^{G516E}*. Twenty

micrograms (µg) of total proteins extracted from 2-week-old seedlings were separated

by SDS-PAGE. CRL or CRL^{G31D} was detected by anti-CRL antibody. CRL^{G31D} migrated

slightly faster and was detected weaker than that of WT. High exp., high exposure time.

Supplemental Figure S7 Predicted structure and *in silico* mutagenesis of OEP80

(A) Predicted structure of OEP80 of *A. thaliana* (AlphaFold identifier AF-Q9C5J8-F1).

Each residue is colored according to the pLDDT (per-residue estimate of its confidence).

The following rules provide guidance on the expected reliability of a given region

(AlphaFold Protein Structure Database: https://alphafold.ebi.ac.uk). Regions with

pLDDT > 90 (cornflower blue) are expected to be modeled to high accuracy. Regions

with pLDDT between 70 (yellow) and 90 (cornflower blue) are expected to be well

modeled (a generally good backbone prediction). Note that regions with pLDDT between

50 (orange) and 70 (yellow) have low confidence. The 3D coordinates of regions with

pLDDT < 50 (orange) often have a ribbon-like appearance. (B and C) Magnified images

around A424 or G516 in (A), which were captured from the interior side of β-barrel

domain. The residues within 5 Å from A424 or G516 were visualized. Gray, red, and blue

balls represent carbon, oxygen, and nitrogen, respectively. (B' and C') Representative

results of *in silico* mutagenesis of A424V or G516E. The purple dashed lines indicate

unfavorable atomic interactions.

Figure 1 S2-6 suppresses *crl-2*

(A) Phenotypic analysis of S2-6. The overall morphology of 3-week-old plants grown on soil. Bar = 1 cm. Autofluorescent images of mesophyll chloroplasts in cotyledons of 2-week-old seedlings. Bar = 20 μm. Cell death in cotyledons of 10-day-old seedlings visualized by trypan blue staining. Bar = 1 mm. (B) Fresh weight of 2-week-old seedlings (n = 12). (C) Number of chloroplasts per cell (n = 10). Mesophyll cells were used in the cotyledons of 2-week-old seedlings. (D) Chlorophyll content of 2-week-old seedlings (n = 12). (E) Percentage of cotyledon area affected by localized cell death in 10-day seedlings (n = 10). (F) Similarity analysis of RNAseq samples. Metagene data from biological replicates of the three biological replicates were used to calculate the correlation coefficient based on Euclidean distances, and the samples were clustered based on their similarity. Asterisks indicate statistical significance (Tukey multiple comparison test, ***P < 0.001, *P < 0.05, NS, not significant).

Figure 2 *oep80*^{G516E} complements *crl-2*

(A) Two-week-old seedlings grown on MS plates. Bar = 5 mm. (B) Autofluorescent images of mesophyll chloroplasts in cotyledons of 2-week-old seedlings. Bar = 50 μm.

Figure 3 *oep80^G516E* suppresses *crl-1*

(A) Two-week-old seedlings grown on MS plates. Bar = 5 mm. (B) Autofluorescent

images of mesophyll chloroplasts in cotyledons of 2-week-old seedlings. Bar = 50 μm.

(C) Cell death in the cotyledons of 10-day-old seedlings visualized by trypan blue staining.

Bar = 1 mm. (D) Fresh weight of 2-week-old seedlings (*n* = 15). (E) Chlorophyll content

of 2-week-old seedlings (*n* = 15). (F) The number of chloroplasts per cell (*n* $\geqq$ 5).

Mesophyll cells were used in the cotyledons of 2-week-old seedlings. (G) Percentage of

the area of cotyledons affected by localized cell death in 10-day-old seedlings (*n* = 10).

Asterisks indicate statistical significance (Tukey multiple comparison test, ****P* < 0.001,

NS, not significant.

Figure 4 The formation of the OEP80 complex is disturbed in *crl-2* and restored

in S2-6

(A) Immunoblotting of the OEP80 complex. Intact chloroplasts from 3-week-old seedlings

grown on MS plates were solubilized with 1% digitonin, and the supernatants were

separated by 2D-BN/SDS-PAGE. OEP80 was detected by an anti-OEP80 antibody. The

molecular weight of ca. 200 kDa (***), from ca. 70 kDa to 150 kDa (*), and from ca. 242–

480 kDa (**) are indicated. IB, immunoblotting (B) Immunoblot analysis of OEP80 and

TOC75-III. Twenty micrograms (µg) of the total proteins extracted from 2-week-old

seedlings were separated by SDS-PAGE. OEP80 and TOC75-III were detected by anti-

OEP80 and anti-TOC75-III antibodies, respectively.

Figure 5 CRL is suggested to be contained in the OEP80 complex

(A) Immunoblotting of immunoprecipitants with anti-GFP antibody. Intact chloroplasts

from 3-week-old seedlings grown in MS plates were solubilized with 1% n-Dodecyl-β-D-

maltoside (DDM). Immunoprecipitants were separated by SDS-PAGE and detected by

anti-OEP80, anti-TIC40, and anti-GFP antibodies, respectively. *pt-GFP*, *35S::pt-*

sGFP(S65T); *CRL-GFP, 35S::CRL-sGFP(S65T)*; IgG(H), IgG heavy chain; IgG(L), IgG

light chain; IP, immunoprecipitation. (B) Immunoblotting of the CRL complex. Intact

chloroplasts from 3-week-old seedlings grown on MS plates were solubilized with 1%

digitonin, and the supernatants were separated by 2D-BN/SDS-PAGE. CRL was

detected by an anti-CRL antibody. The molecular weight is about 146–242 kDa (solid

line), and from ca. 242–480 kDa (dotted line) are indicated. IB, immunoblotting.

Supplemental Figure S1 S1-9 suppresses *crl-2*

(A) Phenotypic analysis of S1-9. Two-week-old seedlings grown on MS plates. Bar = 5

mm. Autofluorescent images of mesophyll chloroplasts in the mesophyll cell of 2-week-

old seedlings. The margin of the cell was indicated by a dashed line. Bar = 20 μm. (B)

Fresh weight of 2-week-old seedlings (*n* = 10). (C) A number of chloroplasts per cell (*n*

≧ 7). Mesophyll cells were used in the cotyledons of 2-week-old seedlings. The asterisks

represent statistical significance (Tukey multiple comparison test, ***P < 0.001, NS, not

significant).

2 **Supplemental Figure S2 Schematic structure of OEP80**

3 The rectangles in the β-barrel domain represent β-strands. Amino acid substitutions in

4 S1-9 and S2-6 are indicated.

Supplemental Figure S3 *oep80^{A424V}* complements *crl-2*

(A) Fresh weight of 2-week-old seedlings (*n* = 10). (B) Number of mesophyll chloroplasts
per cell (*n* 3). Mesophyll cells were used in the cotyledons of 2-week-old seedlings. The
different letters above the plots represent statistical significance (Tukey multiple
comparison test, *P* < 0.05).

Supplemental Figure S4 The phenotypes of the *oep80^{G516E}* single mutant

(A) Two-week-old seedlings grown on MS plates. Bar = 5 mm. (B) Autofluorescent

images of mesophyll chloroplasts in cotyledons of 2-week-old seedlings. Bar = 50 μm.

(C) Fresh weight of 2-week-old seedlings (*n* = 12). (D) Chlorophyll content of 2-week-

old seedlings (*n* = 12). NS not significant (Student's t-test, *P* > 0.05).

Supplemental Figure S5 *oep80^{G516E}* reduces its own protein level, but the reduction

of the protein level is insufficient for the suppression of *crl-2*

(A) Immunoblot analysis of OEP80, TOC75-III, and TOC33. Twenty micrograms (µg) of

total proteins extracted from 2-week-old seedlings were separated by SDS-PAGE.

OEP80, TOC75-III and TOC33 were detected by anti-OEP80, anti-TOC75-III and anti-

TOC33 antibodies, respectively. *oep80-2*, mutant null mutant; +/*oep80-2*, heterozygous

for *oep80-2*; *oep80^{G516E}*, single mutant for *oep80^{G516E}*. (B) Protein abundance of OEP80,

TOC75-III, aminocapronic and TOC33. Protein immunoblot intensities were normalized

to those of Col-0. The mean of three biological replicates ± SD was indicated. (C) Two-

week-old seedlings grown in MS plates. Bar = 5 mm. (D) Autofluorescent images of

mesophyll chloroplasts in cotyledons of 2-week-old seedlings. Bar = 50 µm.

Supplemental Figure S6 *oep80^{G516E}* slightly enhances the protein level of CRL^{G31D}

An immunoblot analysis of CRL and CRL^{G31D} encoded by *crl-2* in *oep80^{G516E}*. Twenty micrograms (µg) of total proteins extracted from 2-week-old seedlings were separated by SDS-PAGE. CRL or CRL^{G31D} was detected by anti-CRL antibody. CRL^{G31D} migrated slightly faster and was detected weaker than that of WT. High exp., high exposure time.

Supplemental Figure S7 Predicted structure and *in silico* mutagenesis of OEP80

(A) Predicted structure of OEP80 of *A. thaliana* (AlphaFold identifier AF-Q9C5J8-F1).

Each residue is colored according to the pLDDT (per-residue estimate of its confidence).

The following rules provide guidance on the expected reliability of a given region

(AlphaFold Protein Structure Database: https://alphafold.ebi.ac.uk). Regions with

pLDDT > 90 (cornflower blue) are expected to be modeled to high accuracy. Regions

with pLDDT between 70 (yellow) and 90 (cornflower blue) are expected to be well

modeled (a generally good backbone prediction). Note that regions with pLDDT between

50 (orange) and 70 (yellow) have low confidence. The 3D coordinates of regions with

pLDDT < 50 (orange) often have a ribbon-like appearance. (B and C) Magnified images

around A424 or G516 in (A), which were captured from the interior side of β-barrel

domain. The residues within 5 Å from A424 or G516 were visualized. Gray, red, and blue

balls represent carbon, oxygen, and nitrogen, respectively. (B' and C') Representative

results of *in silico* mutagenesis of A424V or G516E. The purple dashed lines indicate

unfavorable atomic interactions.

1 **Supplemental Table S1**

Primers	Sequence (5' to 3')	Purpose
OEP80_Hph1_F	AAAGTGGAATGGTACTGGTG	$oep80^{G516E}$ genotyping. PCR amplified fragments from $oep80^{G516E}$ have a *Hph* I site
OEP80_Hph1_R	GGCATCTATCAAATTCATATG	
OEP80_Dde1_F	GATTTGATCATGAACTGTGTCGAG	$oep80^{A424V}$ genotyping. PCR amplified fragments from *OEP80* have a *Dde* I site
OEP80_Dde1_R	CCAAACAGATTGCGGTGGGACTAA	
OEP80_T-DNA_F	AGTAAGAACGAAAGATGGTGAGGA	*oep80-2* genotyping
OEP80_T-DNA_R	CCCATTTGGACGTGAATGTAGACAC	
OEP80_WT_F	CGCTTTGCTCTTACGGCGATACG	*OEP80* genotyping
OEP80_WT_R	TTTACACACCTCCTGCAGGTTGAGG	
crl-2u	GAGGCGAAAACGTCGATCGTCACAACCTTAC	*crl-2* genotyping. PCR amplified fragments from *CRL* have *Hae* II site
crl-2d	TTTAGTGAGACGCTTGATGAGAAGAGCAGCG	
BAR_d1	AGAAACCCACGTCATGCCAGT	*crl-1* genotyping
1primer_U1	TACACGCCGAAATAAACGACC	
CRL1_F	ATGGGTACCGAGTCG	*CRL* genotyping

CRL1_R	CTAGTCTTGCAAGATGAGGGACC	
gOEP80_F	GGTGTTGCCTTGCCTTGGTTTGATCCT	Amplification of *OEP80* containing the regions from 1 kb upstream of the translation initiation site to 3'UTR
gOEP80_R	GTTGAGTTATTATCACTCTATCGGAC	