1

High light can alleviate chilling stress in maize

Authors

Lee Cackett*, Angela C. Burnett*, Jessica Royles, Julian M. Hibberd, Johannes
Kromdijk

Address for all authors

Department of Plant Sciences, University of Cambridge, Cambridge, CB2 3EA, UK

* Contributed equally

Author for correspondence

Email: jk417@cam.ac.uk

Running head (max. 50 characters)

High light can alleviate chilling stress in maize

Abstract

Chilling stress has the potential to significantly decrease growth and yield of sensitive crop plants such as maize. Based on previous work, high light during chilling may exacerbate stress via enhanced photoinhibition but may also aid acclimation responses to chilling. To further understand molecular processes behind responses to chilling with and without high light, two maize accessions with contrasting tolerance (B73 and F7) were exposed to three treatments: chilling, chilling combined with high light and high light alone. Transcriptome data indicated that the chilling treatment resulted in the largest stress response. Addition of high light to chilling stress had a mitigating, rather than additive effect on stress, as evident from alleviated repression of photosynthesis-related genes and less induction of stress-related pathways such as abscisic acid signalling and senescence compared with the response to chilling alone. Five transcription factors belonging to well-known stress-related transcription factor families were identified as candidates for driving the transcriptional changes behind the high-light induced mitigation of chilling stress. Physiological measurements of non-photochemical quenching and the maximum quantum efficiency of photosystem II corroborated the transcriptome results, showing that the addition of high light alleviated photoinhibition and membrane damage caused by chilling. High light alone had little effect on the plant transcriptome or physiological response. Overall, this study overturns previous reports, offers a new outlook on the impact of high light during chilling stress and has the potential to provide clearer targets for crop engineering.

Introduction

The world's population is rising dramatically and with it the need for increased global food production (FAO, 2018). Adding to this challenge is the increasing frequency and severity of extreme weather events (IPCC, 2018) meaning that crop production must also become more climate-resilient if we are to meet the food security challenges of the future. Specifically, temperature is an important determinant of photosynthesis and crop yield (Yoshida et al., 1981; Geange et al., 2021). Therefore, in order to maintain crop yield stability in future environments it is imperative that we continue to develop crop plants which are increasingly resilient to temperature extremes and are adapted to growing and thriving in different climatic conditions. To meet this challenge, a more extensive understanding of the physiological and molecular foundations of temperature stress responses is required.

Maize is the most-produced cereal crop in the world, with a production mass of 1.1 billion tonnes in 2019 (FAO, 2021). It has diverse uses including food, feed, energy, and raw materials for industry (Ma et al., 2022) and makes a major contribution to the caloric intake of the global population (Erenstein et al., 2022). Maize is highly diversified at a genomic level and has dispersed into a wide range of environments distinct from its environment of domestication (Tenaillon and Charcosset, 2011; Sowinski et al., 2020). Despite this diversification, the tropical origins of maize mean that it is highly susceptible to chilling stress when grown in temperate locations. Here, chilling is defined as suboptimal cool temperatures below 15°C and above 0°C (i.e. not freezing stress) with severe chilling stress occurring below 8°C (Frascaroli and Revilla, 2019; Burnett and Kromdijk, 2022). Chilling stress significantly limits seed germination, seedling establishment, and subsequent plant growth and development including yield (Jompuk et al., 2005; Revilla et al., 2005; Sanghera et al., 2011). For example, cold snaps – short periods of chilling temperatures – result in significant yield losses (Sanghera et al., 2011). In addition, poor early-season establishment can elevate farm costs associated with suppression of weeds via herbicide applications or mechanical measures.

The morphological and physiological impacts of chilling stress on maize have been extensively studied and recently reviewed (Burnett and Kromdijk 2022; Ma et al.,

2022). Chilling stress in maize results in excessive wilting, chlorosis and necrosis
and decreased germination rate, root growth, leaf area and leaf appearance rate
(Revilla et al., 2005; Frascaroli and Revilla 2019). Physiologically, chilling stress in
maize results in increased cell membrane disruption (Miedema, 1982), decreased
chlorophyll content and decreased operating efficiency of photosystem II (Φ_{PSII})
(reviewed by Burnett and Kromdijk, 2022). Further, chilling stress in maize
decreases the maximum quantum efficiency of photosystem II (F_v/F_m) (Dolstra et al.,
1994, Fracheboud et al., 1999, Ying et al., 2002) and increases non-photochemical
quenching (NPQ) (Fracheboud et al., 1999, Rodríguez et al., 2013, Riva-Roveda et
al., 2016). NPQ comprises rapidly-reversible photoprotective responses as well as
damage-related photoinhibitive processes from which the plant requires more time to
recover (Baker, 2008, Malnoë, 2018).

These physiological changes resulting from chilling stress contribute to lowered rates
of photosynthesis, ultimately impacting growth, morphology, and productivity, making
chilling tolerance of maize an important target for breeding (Burnett and Kromdijk,
2022). Many transcription factors (TFs), alleles and quantitative trait loci (QTL)
related to chilling stress in maize have been identified (Fracheboud et al., 2004;
Waters et al., 2017; Zeng et al., 2021; Ma et al., 2022). For example, genes such as
CYTOKININ RESPONSE REGULATOR 1 (CRR1), MAP KINASE 8 (*MPK8)* and the
DEHYDRATION-RESPONSIVE ELEMENT-BINDING proteins (DREBs) have been
shown to influence chilling tolerance of maize varieties (Qin et al., 2004, Wang and
Dong, 2009, Wang, Yang and Yang, 2011, Zeng et al., 2021). Despite this progress,
the factors underpinning chilling tolerance in maize, and determining whether a
genotype will display chilling tolerance or chilling sensitivity, are not completely
understood (Frascaroli and Revilla, 2019). Notably, significant gaps remain in
knowledge of the molecular foundations which govern the well-known physiological
responses of maize to low temperatures, specifically key genes controlling signalling
pathways and responses (reviewed in Sowinski et al., 2020).

Photosynthesis is particularly light- and temperature-sensitive due to the tight
coupling between the thylakoid reactions and downstream C_4 acid shuttle and
Calvin-Benson-Bassham cycle (Hüner et al., 1998). Chilling stress and high light
often coincide during clear-sky cold snaps in temperate regions. High light levels can

aid maize growth but can also cause photoinhibition. In the context of maize chilling responses, the role of high light is currently unclear, with two competing mechanisms reported across literature. On the one hand, combined chilling and high light stress has long been considered as additive, with high light exacerbating chilling stress and vice versa (Ortiz-Lopez, 1990). Indeed, it has been demonstrated that high light combined with chilling stress increases photoinhibition in field-grown maize (Farage and Long, 1987). Conversely, other studies have reported that high light can aid acclimation to chilling stress and thereby mitigate the negative effects of chilling. For example, increasing light intensity during a mild chilling treatment enhanced tolerance to a subsequent severe chilling event (Szalai et al., 2018), a phenomenon known as stress priming (Liu et al., 2022). In addition, several cold-tolerant maize lines evaluated by Grzybowski et al. (2019) displayed more photo-sensitivity than cold-sensitive lines in the same study, suggesting that exposure to high light does not have to be additive with chilling stress.

In this work we explored the relationship between chilling and high light in greater depth, in order to elucidate the possible mitigation of chilling stress by high light. In particular, we used a combined genetic and physiological approach to determine whether high light plays a role in exacerbation or mitigation of stress induced by a short cold spell (a cold night and subsequent cool morning at two contrasting light levels, 400 and 1000 µmol photons m^{-2} s^{-1}). We selected two contrasting maize inbred lines: a chilling tolerant accession, F7 (a European Flint line) and a mildly chilling-sensitive accession, B73 (an American Dent line); these accessions have previously been observed to have different chilling tolerances in a two to four day chilling stress period (Enders et al., 2018). Specifically, we set out to test the following hypotheses: 1. High light will exacerbate chilling stress in maize leading to greater physiological damage and reduced photosynthesis. 2. The chilling sensitive accession (B73) will be more susceptible to chilling stress than the chilling tolerant accession (F7). 3. Physiological responses to high light and chilling stress are underpinned by transcriptomic responses, and differences in physiological and photosynthetic changes will reflect differences in gene expression.

The transcriptomic responses confirmed the expected difference in chilling tolerance

between F7 and B73, based on both the overall magnitude of the transcriptomic

response, as well as changes in transcript abundance of previously characterised

cold-responsive genes. However, surprisingly transcriptomic and physiological data

in both accessions indicated a mitigating effect of high light on the chilling stress.

Five candidate transcription factors were identified as control factors for the

mitigating effect of high light on chilling, all of which belong to transcription factor

families which have been previously implicated in abiotic stress responses.

Responses to chilling in combination with 1000 µmol m^{-2} s^{-1} consistently displayed

lower values for physiological stress proxies as well as for transcriptomic responses,

compared to chilling in combination with 400 µmol m^{-2} s^{-1}. Overall these results

clearly demonstrate that high light levels can alleviate chilling stress in maize.

Results

High light impacted the transcriptome signature of chilling stress

Plants of two contrasting maize accessions, B73 and F7, were exposed to one of three treatments. Each treatment was composed of twelve hours night and three hours photoperiod, following which the plants were sampled. The first treatment (referred to as chilling) exposed plants to chilling temperature of 5 $^{\circ}$C during the night and 8 $^{\circ}$C during the photoperiod, while keeping light intensity during the three hour photoperiod the same as during growth prior to the experiment (400 µmol m^{-2} s^{-1}). The second treatment (referred to as chilling plus high light), exposed plants to the same chilling temperatures of 5/8 $^{\circ}$C N/D, but also raised the light intensity during the three morning hours to 2.5 times the growth intensity (1000 µmol m^{-2} s^{-1}). Finally, the third treatment (referred to as high light) kept temperature at 20/28 $^{\circ}$C N/D, while raising the light intensity during the three morning hours to 1000 µmol m^{-2} s^{-1}. Plants from each treatment were compared to matching control plants (0/400 µmol m^{-2} s^{-1}, 20/28 $^{\circ}$C N/D) sampled at the same time of day, one day prior to the start of each treatment. To explore the molecular basis of the responses to chilling and high light, leaf tissue from all untreated and treated plants was sampled for 3'UTR RNA sequencing (RNA-seq). Principal component analysis (PCA) on the transcripts per million (TPM) data (Fig. 1A) assessed the quality and structure of the data and demonstrated that for both B73 and F7 accessions, the chilling treated samples (blue) differed most from the control samples (grey) across the first principal component (explaining 37.3% of transcriptomic variation) whilst high light treated samples (red) appeared similar to the control samples. The chilling plus high light treated samples (yellow) were positioned in between the chilling only and high light only samples, suggesting that the addition of high light to chilled plants had a mitigating effect. The second principal component, explaining 22.8% of the transcriptomic variation, mostly depicted the difference between B73 and F7 accessions (circles and triangles respectively). When TPM data from the two accessions was separated and independent PCAs performed, the differences in treatments described in Figure 1A were conserved (Fig. 1B and C), with the chilling treated samples (blue) differing the most from the control samples, followed by the chilling and high light treated samples (yellow) and the high light treated samples (red) differing very little from the control samples. This explained 43.7% and 61.7% of the transcriptomic variation for B73 and F7 accessions respectively.

Differentially expressed genes (DEGs) for each treatment were identified by comparing the four treated samples with their four matching control samples. Only genes with statistically significant changes in expression (i.e. adjusted p-value < 0.05) in each treatment were retained for subsequent analyses. The chilling treatment led to differential expression of 6065 and 9545 genes for B73 and F7 respectively (Fig. 1D, E and F). Addition of high light to the chilling treatment (chilling + HL) decreased the number of DEGs in both maize lines substantially, resulting in 3405 and 8964 genes for B73 and F7 respectively. Of these genes, the majority were also differentially expressed in the chilling treatment (68% for B73 and 81% for F7, Fig. 1E and F). The transcriptome profile of the high light treatment without chilling was considerably less responsive compared with the chilling treatments, with only 729 and 662 DEGs for B73 and F7 respectively (Fig. 1D, E and F). Results of a Gene Ontology (GO) enrichment analysis on DEGs after high light treatment included the "response to high light intensity" GO term, indicating that the plants did still respond to the high light treatment, but only to a limited degree. When comparing between accessions, 73% of the chilling responsive DEGs in B73 were also differentially expressed in F7 and 70% of the chilling plus high light responsive genes in B73 were also present in the corresponding F7 list (Supp. figure 1). Only 20% of the high light responsive genes in B73 were also present in the corresponding F7. Overall, the RNA-seq data indicated that the chilling treatment alone had the greatest impact on transcription, whilst the addition of high light to chilling treatment decreased the response and the plants treated with high light showed much fewer transcriptional change.

F7 showed transcriptional indicators of improved chilling tolerance

When comparing the transcriptome data from the two accessions, evidence for enhanced tolerance of F7 to chilling stress emerged and provided indications of the mechanisms driving such tolerance. Notably, the transcriptional response of F7 to chilling and chilling plus high light was much broader (i.e. more DEGs) compared to B73 despite using the B73 genome for alignment (Fig. 1D). This is consistent with previously observed differences in the number of cold responsive genes between tolerant and sensitive genotypes (Zhou et al., 2022a). Additionally, several individual genes which have previously been linked with chilling tolerance showed opposite patterns of expression in F7 and B73 (Fig. 2). Zeaxanthin accumulation is negatively

associated with chilling tolerance in maize (Fracheboud et al., 2002). *ZEAXANTHIN*
EPOXIDASE 2 (*ZEP2*), which catalyses the conversion of zeaxanthin to violaxanthin,
showed down-regulation in B73 in response to chilling and chilling plus high light,
whereas expression was up-regulated in F7 in the treatments (Fig. 2A). Conversely,
VIOLAXANTHIN DE-EPOXIDASE 3b (*VDE3b*) which catalyses the conversion of
violaxanthin to zeaxanthin was up-regulated in B73 but down-regulated in F7 in
response to chilling and chilling plus high light (Fig. 2B). Both DEGs point towards a
greater accumulation of zeaxanthin in B73. Additionally, a positive regulator of
chilling stress, *CYTOKININ RESPONSE REGULATOR 1* (*CRR1*) (Zeng et al., 2021),
was up-regulated in F7 and down regulated in B73 whereas the negative regulator of
chilling stress, *MAP KINASE 8* (*MPK8*) (Zeng et al., 2021) showed the opposite
expression (Fig. 2C and D). DESeq2 and TPM values for each of these genes can
be found in Supp. dataset 1 and 2.

A recent study by Zhou et al. (2022a) carried out a transcriptome study on
B73 and Mo17 accessions after a similar chilling treatment to that reported here.
Mo17 is, like F7, considered more chilling tolerant than B73. To explore if
mechanisms of chilling tolerance may be conserved between these two resilient
accessions, genes differentially expressed in response to chilling in both F7 and
Mo17, but not in B73, were determined. This resulted in 233 common genes, 133 up-
regulated and 100 down-regulated (Supp. dataset 3). A GO enrichment analysis on
these 233 genes showed "brassinosteroid metabolic process" as one of the most
significantly enriched terms (FDR = 0.027), suggesting that a brassinosteroid-derived
component of chilling tolerance (Ramirez and Poppenberger, 2020) may be common
across these two accessions.

High light mitigated the repressive effect of chilling on the transcription of
photosynthetic genes
Rather than the hypothesized additive effect, the global transcriptome data
suggested that high light had a mitigating effect on chilling responses (Fig. 1). To find
groups of genes and subsequent individual candidate genes which may be involved
in this response, a weighted gene co-expression network analysis (WGCNA) was
performed. This analysis identified groups of genes (modules) which have similar
patterns of expression. The eigengene values for the resulting 11 modules are

shown in Supp. figure 2 (values provided in Supp. dataset 4) and provide a visual
representation of the gene expression profiles in each module.

Modules one, two, three and four were of specific interest for further analyses The
expression of the genes in modules one and two (Fig. 3A and B) showed large
decreases in expression in response to the chilling treatment while this decrease in
expression was not as severe (i.e. was mitigated) in the chilling plus high light
treatment and there was minimal change in expression in response to high light
alone. Thus, genes in these modules could be involved in the mitigating effect of
high light on the severity of chilling stress. Based on the eigengene values, the
expression of the genes in module one was similar across the two accessions (Fig.
3A) whilst genes in module two had slightly higher expression in B73 compared to
F7 (Fig. 3B), although the pattern of expression was the same. To gain further
biological insight into the role of the genes in these two modules, a GO enrichment
analysis was performed on the gene lists from each module. This gave rise to
multiple enriched GO terms in modules one and two (Supp. dataset 5). GO terms
enriched in both modules were determined to identify possible physiological
responses being most severely impacted by the chilling treatment and mitigated
upon addition of high light (Supp. dataset 6). Several GO terms relating to
photosynthesis were significantly enriched in modules one and two (Fig 3E.),
including "photosystem II assembly", "chlorophyll metabolic process", "thylakoid
membrane organization", and "photosynthetic electron transport chain". Some other
commonly enriched GO terms of interest here include "starch biosynthetic process",
"response to cold", "membrane lipid biosynthetic process" and "response to oxygen-
containing compound" (Fig. 3E). A heatmap of the Log2 Fold Change in expression
of the genes with the "photosynthesis" GO terms found in modules one and two
showed the progressive increase in expression from chilling to high light, with
intermediate expression in the chilling plus high light treatment (Fig. 4A). Selected
individual genes in Figure 4B exemplify the mitigating influence of high light on the
chilling treatment, with all genes showing significant down-regulation in response to
chilling and a notable increase in expression in the chilling plus high light treatment
versus the chilling treatment. None of the genes showed a significant change in
expression in response to high light alone. The pattern of expression of all genes is
consistent between B73 and F7. These representative genes are involved in
photosynthesis (*LCHB1, LHCB2*), NPQ (*PSBS*), carbon fixation (*PPDK1, PEPC1,*

RBCS), xanthophyll metabolism (*VDE1, VP7*), chlorophyll metabolism (*CHAO2*),

membrane integrity (*FAD6*), ROS scavenging (*SOD14*) and sucrose metabolism

(*GAPA*) and give an example of the downregulated processes in the chilling

treatment, for which high light addition seems to allow more sustained activity under

chilling conditions. The DESeq2 and TPM values for each gene can be found in

Supp. dataset 1 and 2. Overall, these results suggest that high light is able to

mitigate the repressive effect of chilling on the transcription of genes primarily

involved in photosynthesis and thus could improve growth in chilled environments.

High light mitigated the inducing effect of chilling on the transcription of ABA

signalling, plant growth and transcription related genes

As with modules one and two, modules three and four (Fig. 3C and D) were of

specific interest for further analysis as they also showed a mitigating effect of high

light when combined with a chilling treatment, but in the opposite direction to

modules one and two. The expression of the genes in these modules showed large

increases in expression in response to the chilling treatment but this was weaker in

the chilling plus high light treatment and there was minimal change in expression in

response to high light alone. Based on eigengene values, expression of genes in

module three was similar across the two accessions whilst genes in module four had

higher expression in F7 compared to B73, although the pattern of expression was

the same. GO enrichment analysis of the module gene lists showed that the "gene

expression" GO term as one of the most significantly enriched term in both gene lists

(Fig. 3F, Supp. dataset 5) several GO terms related to gene expression, including

"regulation of transcription, DNA templated", "RNA splicing", "ribosome biogenesis"

and "chromatin organization". Other GO terms of interest enriched in both module

gene lists included "response to abscisic acid", "leaf development" and "flower

development". Figure 5A shows a heatmap of the Log2 Fold Change in expression

of the genes with the "regulation of transcription, DNA-templated" GO term.

Expression of these genes is most up-regulated in the chilling treated samples,

whilst this up-regulation is decreased in the chilling plus high light samples. As with

modules one and two above (Fig. 4B), several genes representative of transcript

abundance and overall enriched processes in modules three and four were graphed

for clearer visualisation (Fig. 5B). Each gene was significantly up-regulated in

response to chilling and a notable decrease in the expression in response to chilling

plus high light was evident. Again, there were no statistically significant changes in
expression in response to high light alone. Genes in figure 5B are involved in
processes including ABA signalling (*SnRK2.9, CDPK4, CDPK7, PYL8, PYL13*),
stress responsive transcription (*DREB1.2, DREB1.7*), regulation of translation
(*AGO1A, AGO1B*) and leaf senescence (*S40-1, S40-2, S40-4*). The DESeq2 and
TPM values for each gene can be found in Supp. dataset 1 and 2. These results
suggest that the addition of high light during chilling dampens the chilling-enhanced
transcription of genes modulating ABA signalling, growth and transcription and which
could improve growth in chilled environments.
330

Identification of candidate transcription factors driving the high light
mitigation of chilling stress
Transcription factors (TFs) play a vital role in the response to all abiotic stresses,
including chilling and high light (reviewed by Abdullah, Azzeme and Yousefi, 2022,
Sharma et al., 2020, Wang et al., 2016). With this in mind, the TFs and enriched TF
binding motifs present in the WGCNA module gene lists were investigated to find
candidates that could be responsible for the mitigating effect of high light treatment
on the transcriptome signature and stress response of chilling.
Firstly, the TFs present in each module gene list were identified using the
iTAK tool (Zheng et al., 2016). A total of 39 TFs were identified in the gene list from
module one and 88 TFs identified in module two (Table 1). In both modules, the
most represented types of TF were homeobox (HB) TFs (9 and 8 present in modules
one and two respectively) and MYB-related TFs (4 and 10 present in modules one
and two respectively). Both modules also contained TFs from bZIP, C3H and GATA
TF families. Only module two contained TFs from the heat shock factor (HSF) family.
A total of 213 TFs were identified in the gene list from module three and 78
TFs identified in module four (Table 2). APETALA2/Ethylene Responsive Element
Binding Factor domain protein (AP2/EREB) TFs were the most represented type of
TF in both module three and four (24 and 9 present in module three and four
respectively). Other TFs identified in both modules included WRKY, bZIP and NAC
TF families. As expected, most of the highly represented TF families from modules
three and four were different from those in modules one and two. All of the highly
represented TF families in both gene lists are well known for modulating abiotic

stress responses (Abdullah, Azzeme and Yousefi, 2022, Sharma et al., 2020, Wang et al., 2016).

Following the identification of TFs in each module, a Simple Enrichment Analysis (SEA) using the MEME Suite of Motif-based sequence analysis tools was performed to identify enriched TF binding motifs present in the promoters of genes in each module, using 1000 bp of genomic sequence located upstream of the ATG start codon of each gene. Motif enrichment was assessed in comparison to a control set of 5036 genes which did not show significant differential expression in any of the treatments. The promoters of the genes in modules one and two had 28 and 40 enriched TF binding motifs respectively (Table 3). The majority of these were from the bHLH TF family (10 in module one and 17 in module two) and the bZIP TF family (10 in module one and 13 in module 2). Motifs for the HSF TF family were also enriched in modules one and two (1 on module one and 6 in module two). The promoters of the genes in modules three and four had 52 and 18 enriched TF binding motifs respectively (Table 4). Unlike modules one and two, the types of TF families with enriched promoter binding sites in modules three and four differed, with the AP2/EREB TF family being most represented in module three (24 enriched motifs) and the bHLH TF family being most represented in module four (7 enriched motifs).

To identify candidate transcription factors that may underpin the mitigation of chilling stress by high light, the results from the iTAK analysis (Tables 1 and 2) and SEA (Tables 3 and 4) were combined. Here, the gene IDs of the top 20 enriched TF binding motifs from the Simple Enrichment Analysis (SEA) results were scanned (except for module four where only 18 were identified) to determine whether they were present in the list of TFs in the iTAK output (i.e. a transcription factor present in module one which also had an enriched binding motif in the promoters of the other genes in module one). The subsets of transcription factors were then filtered for a module membership (kME) value greater than 0.9. The kME values indicate the centrality of a gene within a WGCNA module, with the most central 'hub' genes within a module having a kME value approaching 1. This resulted in the identification of five TFs, present only in module two or three (Table 5 and Figure 6). Heatmaps in Figure 6 represent the 18-20 most enriched TF binding motifs in the four modules of interest, the TF family of each motif indicated by the text to the right of the corresponding heatmap box. The five TF candidates identified are highlighted by

black boxes within the heatmap and the DESeq2 expression results included
adjacent. *HEAT STRESS TRANSCRIPTION FACTOR 4 (HSF4)* and (*BASIC HELIX-*
LOOP-HELIX TRANSCRIPTION FACTOR 99 (BHLH99) were identified from module
two. And *AP2-EREBP-TRANSCRIPTION FACTOR 52 (EREB52), TEOSINTE-*
BRANCHED1/CYCLOIDEA/PCF TRANSCRIPTION FACTOR 21 (TCP21) and
TEOSINTE-BRANCHED1/CYCLOIDEA/PCF TRANSCRIPTION FACTOR 20
(TCP20) were identified from module three. Assuming that expression changes of
these TF encoding genes are indicative of expression changes of the target genes
which they regulate (i.e. the genes present in each module), the five TFs identified
above serve as candidate transcription factors that modulate the mitigating effect
which high light has on chilling treatment and may have potential to improve maize
growth in chilled conditions.

Decrease in F_v/F_m due to chilling stress was alleviated by high light
A number of common physiological parameters were measured in response to each
treatment to find out if these would be consistent with the aforementioned
transcriptomics-based observations. No detectable effect of chilling, chilling plus high
light or high light alone was found on the concentration of malondialdehyde (MDA)
which is a measure of the lipid peroxidation state (Supplemental Fig. 3), or on the
chlorophyll a/b ratio (Supplemental Fig. 4). There was an increase in total chlorophyll
content compared to the control under the high light treatment but there was no
effect of chilling or chilling plus high light on chlorophyll content (Supplemental Fig.
4). Chlorophyll fluorescence imaging was used to assess the impact of chilling and
high light treatments on the photosynthetic apparatus. Leaf discs taken from plants
exposed to chilling and/or high light stress were dark adapted for 1 hour after which
F_v/F_m was measured; this provided an estimate of the maximum quantum efficiency
of photosystem II. F_v/F_m was significantly decreased by 9-12% in the chilling-
exposed plants (*p*-value < 0.001, Fig. 7A) indicating damage to photosystem II. F_v/F_m
decreased much less in leaves exposed to chilling stress in combination with high
light (2-3% decrease, *p*-value < 0.05, Fig. 7B) and in leaves exposed to high light
alone (3% decrease, *p*-value < 0.05, Fig. 7C).

Chilling stress impaired photoprotective responses

Following measurement of F_v/F_m, leaf discs were immediately exposed to a 10
minute light treatment and a 12 minute dark relaxation period to assess how the
ability to dissipate absorbed energy via photochemical and nonphotochemical
quenching processes was affected by the treatments. Non-photochemical quenching
(NPQ) is the photoprotective downregulation of light-harvesting efficiency, allowing
dissipation of excess light energy not used in photochemistry. Significantly less NPQ
induction occurred in leaves that had been exposed to the chilling treatment
compared to the relevant control: amplitude of NPQ at the end of the light phase was
20-26% lower (p-value < 0.001, Fig. 7D). This was also the case for chilling stress in
combination with high light (19-23% decrease, p-value < 0.001, Fig. 7E). There was
a marginal decrease in NPQ at the end of the light phase (5-9%, p-value < 0.05, Fig.
7F) in leaves that had been subjected to the high light treatment compared to the
relevant control. During the 12 minute dark relaxation phase, significantly less NPQ
relaxation occurred in the leaves that had been exposed to the chilling treatment,
leading to higher residual NPQ at the end of the dark phase (29-64% increase, p-
value < 0.001, Fig. 7G). However, in leaves that had been exposed to combined
chilling and high light, there was no significant difference in residual NPQ between
treatment and control (Fig. 7H) suggesting that the addition of high light allowed a
more complete relaxation of NPQ in the 12 minute dark relaxation. In leaves that had
been exposed to high light only, residual NPQ was 31-37% lower than in the control
(p-value < 0.001, Fig. 7I). Again, these results confirm the findings in the
transcriptome data indicating that high light is able to mitigate chilling stress.

Relative injury was increased by chilling stress but mitigated by high light
In plants exposed to chilling, the relative injury of leaf tissue, a measure of stress-
induced electrolyte leakage due to membrane breakdown, was 57-61% greater than
in control plants (p-value < 0.01, Fig. 7J). When plants were exposed to chilling
stress in combination with high light, relative injury was not different than the control
(Fig. 7K) suggesting that high light mitigated the chilling-induced damage, in line with
the transcriptome data above. Relative injury in response to high light without chilling
also did not differ from the control (Fig. 7L).

Discussion

Contrasting results have been reported on the effect of adding high light to chilling-
induced stress in cold-sensitive plants like maize. Some work suggests that high light
would add to chilling-induced stress, primarily via photoinhibition and photodamage
(Farage and Long, 1987, Ortiz-Lopez, 1990), whereas others suggest that high light
can aid acclimation to chilling stress and thereby mitigate the negative effects of
chilling (Szalai et al., 2018, Grzybowski et al., 2019). These discrepancies could be
due to different chilling and high light treatments and growth environments, thus
making direct comparisons of the results difficult. The results presented here show
that the addition of three hours of high light in the morning as part of a 12 h night and
3 hr morning chilling treatment had a surprisingly consistent mitigating impact on the
effect of chilling on both transcriptomic profiles and physiological status of the leaves
(Fig. 1, 3, 4, 5, and 7). These responses were found in both B73 and F7 accessions,
although the transcriptomic responses of F7 indicated that it is more tolerant to
chilling than B73, consistent with previous reports (Enders et al., 2018).

The role of high light in mitigating chilling stress

The transcriptome and physiological data suggest that three hours of
increased light in the morning had very little impact as an individual treatment but
was significantly more impactful when plants had already had 12 hours of cold, dark
treatment. Modules one, two, three and four of the WGCNA contained genes with
patterns of expression which depicted the mitigating effect of high light on the chilling
treatment (Fig. 3A, B, C and D). GO enrichment analyses and analyses of individual
genes in modules one and two showed that processes such as photosynthesis,
xanthophyll metabolism, chlorophyll metabolism, response to ROS and sucrose
responses were all decreased in response to chilling but mitigated by high light
treatment (Fig. 3E, Fig. 4A). These physiological responses have previously been
reported as repressed in chilling environments and result in damaging physiological
changes and reduced plant growth (reviewed by Burnett and Kromdijk, 2022, Savitch
et al., 2011). Based on the expression patterns of many genes with these GO terms,
it is plausible that the addition of high light was able to reduce the severity of chilling-
induced down-regulation of these processes, thus improving physiological integrity.
F_v/F_m and NPQ relaxation results (Fig. 7) substantiate that the addition of high light
alleviates chilling-induced strain on photosynthesis and photoinhibition. GO
enrichment analyses and analyses of individual genes in modules three and four

indicated that processes such as ABA signalling, leaf senescence, transcription, and translation were all enhanced during chilling stress, but less so after the addition of high light (Fig. 3F, Fig. 5A). The up-regulation of ABA and senescence related processes in response to chilling is expected, as they have been widely reported previously (Capell and Dörffling, 1993, Janowiak, Maas and Dörffling, 2002, Janowiak, Luck and Dörffling, 2003, Masclaux-Daubresse et al., 2007, Wingler and Roitsch, 2008). A reduction in ABA responses upon the addition of high light to the chilling treatment indicates a less stressed state and could result in the re-opening of stomata to allow for continued transpiration and uptake of CO_2. The reduced induction of senescence related genes upon the addition of high light could also be a significant mechanism behind the mitigating effect of high light during chilling. Delayed leaf senescence and aging allows for enhanced photosynthesis and decreased ROS and has been linked with improved abiotic stress resistance (reviewed by Zhao, Zhao and Wang, 2022). The significant increase in the expression of genes associated with gene expression was somewhat unexpected and counterintuitive. Generally, stress conditions limit energetically consumptive processes such transcription and translation (Muñoz and Castellano, 2012) and translation rates in plants are proportional to temperature (Guillaume-Schöpfer et al., 2020). However, the stress-induced decrease of global translation is often accompanied by a switch to selective transcription and translation of proteins needed for survival during the stress (Holcik and Sonenberg, 2005, Muñoz and Castellano, 2012), in the case of chilling stress, these could be genes involved in processes such as NPQ, chlorophyll metabolism or ABA signalling.

The mechanisms for how high light might mitigate chilling stress suggested here corroborate those reported by Szalai et al. (2018) who found that exposing maize plants to high light (relative to intensity during growth) prior to a severe chilling stress enhanced acclimation to the chilling by modulating genes involved in photosynthesis, chlorophyll biosynthesis, ribosome biogenesis and translation. Thus, it appears that high light has a fundamental impact on plant photosynthesis, metabolism and signalling during chilling stress responses.

Transcription factors which modulate chilling stress responses and mitigation by high light have been identified

The global transcriptome analyses above identified groups of genes and
physiological responses that are significantly impacted during chilling and show a
diminished response in the combination with high light. TFs are master regulators of
chilling stress responses and can be manipulated to alter tolerance (reviewed:
Abdullah, Azzeme and Yousefi, 2022, Sharma et al., 2020, Wang et al., 2016). To
identify TFs with the potential to be key drivers of the transcriptional changes
observed in modules one, two, three and four, we filtered for TFs which were present
in the modules of interest (Table 1 and 2), had enriched binding motifs in the
promoters of genes in the same module (Table 3 and 4) and had high kME values
for the module in which they belonged (indicating a high level of centrality or
connectivity with the other genes in the module). This filtering was based on the
premise that TFs and their target genes are co-expressed, such that a change in
abundance of a TF alters the interaction of that TF with cis-acting elements in target
gene promoters to control stress-responsive gene clusters. Five individual TFs stood
out from this analysis: *bHLH99* and *HSF4* from module two and *TCP20, TCP21* and
EREB52 from module three (Fig. 6 and Table 5). All of these are part of TF families
often reported in association with stress tolerance and may have promise as
candidates for improving chilling tolerance, potentially more so in high light
environments. *bHLH99* belongs in the large group of bHLH TFs which have been
linked with modulating cold stress responses in several plant species, including
maize (reviewed by Guo et al., 2021, Sun, Wang and Sui, 2018). The expression of
several bHLH genes was reported as differentially expressed in response to cold in
the high-vigour maize variety, Zhongdi175 (Li et al., 2021). Overexpression of the
maize bHLH TF, *ZmPTF1*, improved the drought tolerance of transgenic maize,
indicating a role for this group of TFs in manipulating abiotic stress responses (Li et
al., 2019). As the name indicates, TFs of the heat shock factor (HSF) family often
respond strongly to temperature. A study which surveyed several independent maize
cold response transcriptomes to detect key genes involved in the response identified
HSF4 as consistently differentially expressed in response to chilling in multiple
datasets (Sowiński et al., 2020), illustrating a robust and conserved role for this
gene. *TCP20* and *TCP21* form part of the TEOSINTE BRANCHED 1, CYCLOIDEA,
PCF1 (TCP) transcription factor family. TCP transcription factors are key modulators
of plant growth processes such as branching, floral symmetry and cell cycle and
control these processes in response to diverse environmental cues, including abiotic

stress (Danisman, 2016). In rice, overexpression of *OsTCP19* improved tolerance to

NaCl and mannitol stresses (Mukhopadhyay and Tyagi, 2015), whilst down-

regulation of *OsTCP21* increased plant tolerance to cold stress (Wang et al., 2014).

Overexpression of maize *TCP42* in Arabidopsis resulted in enhanced drought

tolerance (Ding et al., 2019). *EREB52* belongs to the large AP2-EREB transcription

factor family which is extensively associated with plant cold stress responses

(Ritonga et al., 2021, Zhou et al., 2022b). The most well-known members of this

family are the DREBs which form part of the C-repeat binding factor (CBF)-regulon

(Liu et al., 2019) which increase the expression of cold-regulated genes and

enhance chilling tolerance in maize (Qin et al., 2004, Wang and Dong, 2009, Wang,

Yang and Yang, 2011). Thus, the five TFs that arose from analyses presented here

are clearly implicated to impact maize chilling responses and tolerance and warrant

further exploration.

F7 is more chilling tolerant than B73

Whilst significant differences were observed between F7 and B73 for some of the

physiological parameters measured here (Fig. 7), there were no significant

interactions between accession and treatment, indicating a lack of accession-specific

differential response to the stress that would indicate variation in chilling tolerance.

However, this may be due to the relatively short (15 hour) chilling treatment period

imposed here rather than a lack of difference in chilling tolerance between the

accessions. Enders et al. (2018) reported that maize grown under a 24 hour period

of chilling had few changes in growth inhibition (plant height, area, and width)

compared with plants under control conditions. Thus, observing physiological

differences in the stress response between accessions may require a longer

treatment period. There were, however, transcriptomic indicators which suggested

greater chilling tolerance in F7. Firstly, F7 had substantially more DEGs than B73 in

response to chilling and chilling plus high light (Fig. 3D) which may suggest a

stronger, more effective response with more genes responding to the stress. The

same type of result was reported by Zhou et al. (2022a) when they compared B73

with the chilling-tolerant Mo17 accession and Li et al. (2019) when comparing

chilling-tolerant M54 and chilling-sensitive 753F accessions. Secondly, individual

genes which have previously been reported to confer chilling tolerance showed

different patterns of expression in F7 compared with B73 (Fig. 2). The expression of

ZEP2 was significantly down-regulated in B73 in response to chilling and chilling plus

high light whereas the expression is up-regulated in F7 in these treatments (Fig. 2A).

Conversely, *VDE3b* was up-regulated in B73 but down-regulated in F7 in response

to chilling and chilling plus high light (Fig. 2B). These results suggest a greater

accumulation of zeaxanthin in B73, which is negatively associated with chilling

tolerance (Fracheboud et al., 2002). Another such example of chilling tolerance in F7

is the differences in expression of *CRR1* and *MPK8*. *CRR1*, a positive regulator of

chilling stress (Zeng et al., 2021), was up-regulated in F7 and down regulated in B73

whereas *MPK8*, a negative regulator of chilling stress in the same signalling pathway

(Zeng et al., 2021), showed the opposite expression (Fig. 2C and D). These are

some examples of possible candidates for conferring chilling tolerance, but it would

be beneficial to perform the same type of transcriptome analysis on plants exposed

to longer periods of chilling stress when the physiological differences between

accessions may become more apparent. Interestingly, comparison of the F7-specific

chilling-responsive transcriptome (i.e. DEGs in F7, but not in B73) with that of

another relatively chilling-tolerant accession, Mo17 (published by Zhou et al., 2022a)

suggested common mechanisms such as brassinosteroid modulated signalling might

help to enhance tolerance. Brassinosteroid application to chilling-sensitive species

such as rice, tomato, cucumber, Arabidopsis and maize improves plant performance

by promoting the expression of *COR* genes, master regulators of the cold response

as well as mechanisms including antioxidant mobilization, membrane fluidity and

Ca^{2+} influx (reviewed by Ramirez and Poppenberger, 2020).

To conclude, the work here provides novel insights into the role of high light during

plant exposure to chilling conditions. The addition of high light was able to alleviate

the negative impact of chilling on photosynthesis, photoinhibition and membrane

integrity induced by chilling. Additionally, of the ±2500 annotated transcription factor

genes in maize (Lin et al., 2014), five were identified as strong candidates in driving

the mitigating changes in transcription brought about by high light in combination

with chilling. Importantly the transcriptional responses associated with improved

chilling tolerance of the F7 accession are very similar to those reported for other

accessions and so we propose these reflect a common transcriptional response in

maize accessions with enhanced chilling tolerance.

Materials and methods

Plant material and growth conditions

Seeds of the maize (*Zea mays*) inbred lines B73 and F7 were obtained from the International Rice Research Institute (IRRI) and the U.S. National Plant Germplasm System (NPGS) respectively. Seeds were germinated on wet filter paper in the dark at 28°C for three days after which each germinated seed was transferred into a two litre pot containing a mixture of 2 parts nutrient-rich compost (Levington Advance M3, ICL, Ipswich, UK) to 1 part topsoil (Westland, Dungannon, Northern Ireland), 10 ml Miracle-Gro all-purpose fertiliser beads and 15 ml Miracle-Gro magnesium salt (Scotts Miracle-Gro, Marysville, OH, USA) and placed in a growth room (Conviron Ltd, Winnipeg, CA) set at 28/20°C day/night, 65% relative humidity, 400 µmol photons m^{-2} s^{-1} and a 12 hour photoperiod. 15 days post germination, the plants were transferred to a growth cabinet (Percival E-41HO, CLF Plant Climatics GmbH, Wertingen, Germany) set to the same conditions as the growth room. Plants were grown in three batches to which different experimental treatments were applied (see below). Each batch contained eight plants per accession, of which four plants per accession were randomly assigned to the experimental control and four plants per accession were exposed to the experimental treatment.

Experimental treatments and sampling

Three discrete experiments were performed within a four-week period, each with a different treatment applied to a separate batch of plants. For each of the three experiments four control plants per accession were harvested three hours after the start of the photoperiod on day 20 (post germination). At the end of day 20, the remaining four plants in the cabinet were treated with one of three treatments (i.e. treatments occurred on night 20 and three hours into the photoperiod of day 21 post germination). A chilling treatment was applied to the first batch of plants, with a temperature of 8/5°C (D/N). Plants were harvested following 12 hours chilling night (5°C) and three hours chilling day (8°C). A high light treatment of 1000 µmol photons m^{-2} s^{-1} during the three hours day was applied to the second batch of plants. This treatment was applied at the beginning of the photoperiod on day 21 (post germination) and plants were harvested following three hours high light. Finally a combined chilling and high light treatment (8/5°C; 1000 µmol photons m^{-2} s^{-1}) was applied to the third batch of plants and plants were harvested following 12 hours

chilling night and three hours combined chilling and high light day. All three independent experiments were conducted in the same growth cabinet.

Tissue harvesting was performed three hours into the photoperiod on day 20 (control plants) or day 21 (treated plants). All samples were taken from expanded leaf three (the third leaf to emerge), before expansion of leaf four. For all plants, two rectangular leaf cuttings (avoiding the midrib) were obtained and flash-frozen in liquid nitrogen for subsequent RNA extraction. Additionally, discs of 6 mm diameter were punched from each leaf for subsequent physiological measurements:

For the electrolyte leakage assay (relative injury), three discs per leaf were individually placed into 200 µL MilliQ water for a 30 minute wash period to remove electrolytes resulting from the punching of the disc. Each disc was then placed into 1 mL MilliQ water and left to soak for 20 hours.

For the chlorophyll fluorescence assay, three discs per leaf were each placed adaxial side facing down into the well of a 96-well plate and held in place with a small piece of sponge dampened in MilliQ water to prevent dehydration.

The MDA and chlorophyll assays were determined on the same sample. Three discs per leaf were pooled and flash-frozen in liquid nitrogen and stored at -80°C for subsequent analysis.

Relative injury measurement

Following the 30 minute wash period and 20 hours soaking, samples were shaken at 300 rpm for 15 minutes (Thermomixer-Mixer HC, Starlab (UK) Ltd, Milton Keynes, UK). The electrical conductivity of a 100 µl aliquot of the resultant solution was measured (LAQUAtwin EC-11 conductivity meter, Horiba Instruments, Kyoto, Japan). Following this measurement (Time 1), samples were heated at 99°C for three hours and cooled to room temperature. Each sample was vortexed and electrical conductivity was re-measured (Time 2). For each measurement of conductivity, samples were measured at least twice (a new aliquot was taken each time) until the reading was stable between aliquots. A blank (MilliQ water) was also measured at each time point.

Relative injury of the leaf material was calculated according to the following formula (Thalhammer et al., 2020):

$$Relative\ injury\ (\%) = \frac{Conductivity\ Time\ 1 - Blank\ Time\ 1}{Conductivity\ Time\ 2 - Blank\ Time\ 2}\ x\ 100$$

(Equation 1)

Chlorophyll fluorescence measurement

Chlorophyll fluorescence was used to measure F_v/F_m and NPQ (Fracheboud et al., .

1999; Baker et al., 2008). The leaf discs harvested into 96-well plates were

immediately dark adapted for one hour. For chilling treatments (both chilling and the

combined chilling and high light treatment) this dark adaptation was carried out in

chilling conditions. For the high light and control treatments this dark adaptation was

carried out at room temperature. Following dark adaptation, chlorophyll fluorescence

was measured using a FluorCam (Photon Systems Instruments, Drásov, Czech

Republic). Leaf discs were exposed to a 10 minute light period followed by a 12

minute dark relaxation period. F_v/F_m was measured on dark-adapted leaves at the

start of the measurement period. NPQ was obtained from saturating pulses in the

light and dark. In the light, saturating pulses were given every 20 seconds for the first

minute, then once per minute. In the dark, saturating pulses were given every 20

seconds for the first minute, then once per minute for two minutes, then once every

three minutes for the remaining nine minutes. For samples resulting from the chilling

treatments (the chilling treatment and the combined chilling and high light treatment)

the FluorCam measurement was carried out at a leaf temperature of 8°C, achieved

by placing a cold plate inside the FluorCam (Solid State Cold/Hot Plate AHP-

301CPV, ThermoElectric Cooling America Corporation, Chicago, IL, USA). For

samples resulting from the high light treatment the FluorCam measurement was

carried out at a leaf temperature of 20°C.

Biochemical assays

Samples harvested for MDA and chlorophyll measurements were cryo-ground

(TissueLyser II, Qiagen, Hilden, Germany). Glass beads were inserted into sampling

tubes prior to harvest to expedite the grinding process.

MDA and chlorophyll quantification

Each set of three discs harvested for MDA and chlorophyll assays was extracted in
1ml 80% ethanol (v/v) at 4°C for 72 hours. Samples were then vortexed (15 s) and
centrifuged (5 mins, 4°C, 13,000rpm).

For the chlorophyll measurement, a 150 µl aliquot of the ethanol extract was added
to a 96-well plate in duplicate, along with an ethanol blank, and absorbance was
measured at 632 nm, 649 nm, 665 nm and 696 nm (Synergy HT Plate Reader,
BioTek, Winooski, VT, USA). Chlorophyll was determined according to Ritchie et al.
(2008).

$$Chlorophyll\ a = 0.0604 \times Abs632 - 4.5224 \times Abs649$$
$$+ 13.2969 \times Abs665 - 1.7453 \times Abs696$$

732   (Equation 2)

$$Chlorophyll\ b = -4.1982 \times Abs632 + 25.7205 \times Abs649$$
$$- 7.4096\ x\ Abs665 - 2.7418 \times Abs696$$

734   (Equation 3)

$$Total\ chlorophyll = 24.1209 \times Abs632 + 11.2884 \times Abs649$$
$$+ 3.7620 \times Abs665 + 5.8338 \times Abs696$$

737   (Equation 4)

For the MDA measurement, two 250 µl aliquots were taken from each ethanol
extract. One aliquot was added to 250 µL "Minus TBA" solution and one aliquot was
added to 250 µL "Plus TBA" solution. The "Minus TBA" solution contained 200 g/L
trichloroacetic acid and 0.1 g/L butylated hydroxytoluene. The "Plus TBA solution"
consisted of "Minus TBA" with the addition of 6.5 g/L thiobarbituric acid. Samples
with plus/minus TBA solution were vortexed (15 s), heated (25 mins, 95°C), cooled
to room temperature then centrifuged (5 mins, 4°C, 13,000rpm). A 200 µl aliquot was
then added to a 96-well plate in duplicate, along with a blank ("Plus TBA" in 80%
ethanol). Absorbance was measured at 440 nm, 532 nm and 600 nm (Synergy HT

Plate Reader, BioTek, Winooski, VT, USA). MDA was determined according to the
method of Hodges et al. (1999), using the corrected equation by Landi (2017).
750

$$[(\text{Abs } 532_{+\text{TBA}}) - (\text{Abs } 600_{+\text{TBA}}) - (\text{Abs } 532_{-\text{TBA}} - \text{Abs } 600_{-\text{TBA}})] = A$$

751 (Equation 5)

$$[(\text{Abs } 440_{+\text{TBA}} - \text{Abs } 600_{+\text{TBA}}) \times 0.0571] = B$$

753 (Equation 6)

$$MDA\ equivalents\ (nmol\ ml^{-1}) = (A - B)/157000 \times 10^6$$

(Equation 7)

Total RNA extraction, cDNA library preparation and Transcriptome Sequencing
Total RNA was extracted from leaf samples of the control and treated (chilling, high
light and chilling plus high light) plants using the RNeasy Plant Mini Kit (Qiagen,
Germany) according to the manufacturer's instructions. Genomic DNA contamination
was removed from each RNA sample using the RNase-Free DNase Set (Qiagen,
Germany). RNA degradation and contamination were checked on 1% agarose gels,
then RNA quality and concentration were determined with the RNA600 Pico Assay
using the Agilent 2100 Bioanalyzer (Agilent Technologies, USA) and ND-200
nanodrop (NanoDrop Technologies Inc., USA). For each sample, a cDNA library was
prepared from 200 ng RNA starting material using the QuantSeq 3' mRNA-Seq
Library Prep kit FWD for Illumina (Lexogen, Austria) according to the manufacturer's
instructions. Library quality and concentration was determined with the High
Sensitivity DNA assay using the Agilent 2100 Bioanalyzer (Agilent Technologies,
USA) and Qubit® dsDNA HS Assay Kit (Invitrogen, USA). Finally, the libraries were
sent to the Novogene Cambridge Genomic Sequencing Centre for sequencing using
the Illumina NovaSeq 600 PE150 sequencing platform and strategy. At least 6 Gb of
raw data per sample was generated for subsequent transcriptome analysis (refer to
Supp. dataset 9 for the number of raw reads generated for each sample).

Transcriptome data processing

Due to the nature of the library preparation kit used only the forward sequencing reads were used for data analysis. The quality of raw sequencing data (.fastq format) was assessed and controlled using the FastQC platform (version 0.11.4, Andrews, S. 2010, http://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Adapter trimming and filtering of all low quality reads was performed using BBDuk (https://www.geneious.com/plugins/bbduk/) with the following parameters: k=13, ktrim=r, useshortkmers=t, mink=5, qtrim=r, trimq=20, minlength=50. The Zm-B73-REFERENCE-NAM-5.0 transcriptome was downloaded from MaizeGDB (https://www.maizegdb.org/genome/assembly/Zm-B73-REFERENCE-NAM-5.0) and used to build a Salmon reference index which was used to quantify the cleaned reads (Salmon version 1.5.2). For Salmon quantification (Patro et al., 2017), all parameters were left as default, except the --noLengthCorrection parameter was used. The Salmon alignment and quantification results were checked using MultiQC (https://multiqc.info/). Refer to Supp. dataset 9 for the number of reads remaining after BBDuk trimming and the % alignment for reads to the B73 reference genome. To check that biological replicates clustered together and to visualise how the treatments differed from controls, a principal component analysis (PCA) was performed using the ggfortify package in R (https://cran.r-project.org/web/packages/ggfortify/index.html). For this, the transcripts per million (TPM) outputs from the Salmon quantification were used. To filter out very lowly expressed genes, only genes with a total of 36 TPMs across all 48 samples were included. The filtered data were then normalised to account for library size and transformed to a log scale using variance stabilising transformation (vst) to allow for easier visualisation.

Identification of differentially expressed genes (DEGs), weighted Gene Correlation Network Analysis (WGCNA) and functional annotation of clusters and

To determine the changes in expression of individual genes, the DESeq2 package in R was used (version 4.2, Love, Huber and Anders, 2014, https://bioconductor.org/packages/release/bioc/html/DESeq2.html). The quant.sf file generated for each sample from the Salmon quantification was used as the input for the DESeq2 analysis. Independent DESeq2 analyses were performed for the three

independent experiments (i.e. samples from each treatment were compared with the

corresponding control for that experiment).

The WGCNA package in R (version 1.71, Langfelder and Horvath, 2008,

https://cran.r-project.org/web/packages/WGCNA/index.html) was used to cluster

genes which had similar patterns of expression. For the analysis the TPM outputs

from the Salmon quantification were used. To filter out genes with very low

expression levels, only genes with a total of 36 TPMs across all 48 samples were

included. The WGCNA one-step network construction and module detection function

were used to construct the gene network and identify modules using a "signed"

network type and minimum module size of 30 genes. All other parameters were left

as default. The lists of genes within each module and module eigengene values

were then extracted for downstream analysis.

To gain further biological insight into modules with interesting expression

patterns, Gene Ontology (GO) enrichment analyses were performed using AgriGO

v2 (Tian et al., 2017, http://systemsbiology.cau.edu.cn/agriGOv2/) with the Maize

AGPv4 (Maize-GAMER) genome set as the background reference.

Determination of transcription factors in gene lists and enriched transcription

factor binding sites to identify candidate transcription factors

The transcription factors present in the combined module gene lists were identified

using the Plant Transcription Factor and Protein Kinase Identifier and Classifier

(iTAK) tool (Zheng et al., 2016, http://itak.feilab.net/cgi-bin/itak/index.cgi). For this,

the unspliced gene sequences for each gene were downloaded from Biomart

(https://plants.ensembl.org/biomart/martview/) using the Ensembl Plants Gene 54

database and Zm-B73-REFERENCE-NAM-5.0 dataset. To identify the enriched

transcription factor binding sites in the promoters of genes in the combined module

gene lists, a Simple Enrichment Analysis (SEA) using the MEME Suite of Motif-

based sequence analysis tools (Bailey and Grant, 2021, https://meme-

suite.org/meme/index.html) was performed. The promoter region 1000 bp upstream

of the transcription start site of each gene was downloaded from Biomart. This was

compared to a manually curated list of control promoter sequences which were from

a list of genes which showed no significant differential expression in any of the

treatments (i.e. p-adj > 0.05 in chilling, chilling plus high light and high light

treatments). The CIS-BP 2.00 Single Species DNA motif database for *Zea mays* was

used to test for enrichment. To identify transcription factor candidates, the top 18-20
enriched TF binding motifs from the SEA were scanned to determine whether they
were present in the list of TFs in the iTAK output. Transcription factors identified
were then filtered for a module membership (kME) value greater than 0.9. The kME
values indicate the centrality of a gene within a WGCNA module.

Statistics
The adjusted *p*-values from the DESeq2 output were used to identify statistically
significantly differentially expressed genes. DESeq2 uses a Wald test for significance
testing, and Benjamini and Hochberg to adjust *p*-values for multiple comparisons.
Two-way ANOVA was performed to identify differences in relative injury and
chlorophyll fluorescence parameters, testing for effects of accession, treatment, and
an interaction between accession and treatment. All statistics were performed using
R (R Core Team, 2022).

Supplemental data
• Supplemental dataset 1: DESeq2 data for genes in figure 2, 4, 5 and 6 (excel
doc – sheet 1).
• Supplemental dataset 2: TPM data for genes in figure 2, 4, 5 and 6 (excel doc
– sheet 2).
• Supplemental dataset 3: List of DEGs common in F7 and Mo17, but not B73,
in response to chilling (excel doc – sheet 3).
• Supplemental dataset 4: Lists of genes present in each WGCNA module
(excel doc – sheet 4).
• Supplemental dataset 5: Results from GO enrichment analyses of WGCNA
modules of interest (excel doc – sheet 5).
• Supplemental dataset 6: Comparison of enriched GO terms present in
modules one versus two and modules three versus four (excel doc – sheet 6).
• Supplemental dataset 7: List of all TFs present in modules one, two, three and
four (excel doc – sheet 7).

• Supplemental dataset 8: Transcription factor binding motifs enriched in the
promoters of genes in modules one, two, three and four identified using the
MEME Suite of Motif-based sequence SEA tool (excel doc – sheet 8).
• Supplemental dataset 9: Supp. dataset 9: Illumina sequencing output (raw
reads) for each sample, reads remaining after BBDuk trimming and the % of
these reads which aligned to the B73 reference genome using Salmon
alignment (excel doc – sheet 9).
883

884　**Author contributions**
885　LC, ACB and JK conceived and designed the experiment with input from JR and
886　JMH, LC, ACB and JR performed the measurements. LC and ACB analyzed the
887　data. LC and ACB wrote the manuscript with contributions from all authors.
888

Funding

ACB, LC, JR and JK were supported by the UK Research and Innovation Future
Leaders Fellowship (UKRI-FLF) MR/T042737/1 and an Isaac Newton Trust /
Wellcome Trust ISSF / University of Cambridge Joint Research Grants Scheme
award to JK. For the purpose of open access, the authors have applied a Creative
Commons Attribution (CC BY) licence to any Author Accepted Manuscript version
arising from this submission.

References

Abdullah SNA, Azzeme AM, Yousefi K (2022) Fine-Tuning Cold Stress Response Through Regulated Cellular Abundance and Mechanistic Actions of Transcription Factors. Front Plant Sci. doi: 10.3389/fpls.2022.850216

Andrews S (2010) A quality control tool for high throughput sequence data. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Bailey TL, Grant CE (2021) SEA: Simple Enrichment Analysis of motifs. bioRxiv 2021.08.23.457422

Baker NR (2008) Chlorophyll fluorescence: a probe of photosynthesis in vivo. Annu Rev Plant Biol **59**: 89–113

Burnett AC, Kromdijk J (2022) Can we improve the chilling tolerance of maize photosynthesis through breeding? J Exp Bot **73**: 3138–3156

Capell B, Dörffling K (1993) Genotype-specific differences in chilling tolerance of maize in relation to chilling-induced changes in water status and abscisic acid accumulation. Physiol Plant **88**: 638–646

Danisman S (2016) TCP transcription factors at the interface between environmental challenges and the plant's growth responses. Front Plant Sci **7**: 1–13

Ding S, Cai Z, Du H, Wang H (2019) Genome-wide analysis of TCP family genes in Zea mays l. Identified a role for ZmTCP42 in drought tolerance. Int J Mol Sci. doi: 10.3390/ijms20112762

Dolstra O, Haalstra SR, van der Putten PEL, Schapendonk AHCM (1994) Genetic variation for resistance to low-temperature photoinhibition of photosynthesis in maize (Zea mays L.). Euphytica **80**: 85–93

Enders TA, Dennis SS, Justin O, Steven CT, Gehan MA, Miller ND, Spalding EP, Springer NM, Hirsch CD (2018) Classifying cold-stress responses of inbred maize seedlings using RGB imaging. Plant Direct 1–11

Fracheboud Y, Ribaut JM, Vargas M, Messmer R, Stamp P (2002) Identification of quantitative trait loci for cold-tolerance of photosynthesis in maize (Zea mays L.). J Exp Bot **53**: 1967–1977

Erenstein O, Jaleta M, Sonder K, Mottaleb K, Prasanna BM (2022) Global maize production, consumption and trade: trends and R&D implications. Food Secur **14**: 1295–1319

FAO (2021) World Food and Agriculture – Statistical Yearbook 2021. World Food and Agriculture – Statistical Yearbook 2021. doi: 10.4060/cb4477en

FAO (2018) The future of food and agriculture – Alternative pathways to 2050.

Farage P, Long S (1987) Damage to maize photosynthesis in field during periods when chilling is combined with high photon fluxes. *In* J Biggens, ed, Progress in photosynthesis research, Vol. IV. Martinus Nijhoff Publishers, The Netherlands, pp 139–142

Fracheboud Y, Haldimann P, Leipner J, Stamp P (1999) Chlorophyll fluorescence as a selection tool for cold tolerance of photosynthesis in maize (Zea mays L.). J Exp Bot **50**: 1533–1540

Fracheboud Y, Jompuk C, Ribaut JM, Stamp P, Leipner J (2004) Genetic analysis of
cold-tolerance of photosynthesis in maize. Plant Mol Biol **56**: 241–253

Frascaroli E, Revilla P (2019) Genomics of cold tolerance in maize. *In* J Bennetzen, S Flint-
Garcia, C Hirsch, R Tuberosa, eds, The Maize Genome. Springer Nature Switzerland, pp
287–304

Geange SR, Arnold PA, Catling AA, Coast O, Cook AM, Gowland KM, Leigh A,
Notarnicola RF, Posch BC, Venn SE, et al (2021) The thermal tolerance of photosynthetic
tissues: a global systematic review and agenda for future research. New Phytologist **229**:
2497–2513

Grzybowski M, Adamczyk J, Jończyk M, Sobkowiak A, Szczepanik J, Frankiewicz K,
Fronk J, Sowiński P (2019) Increased photosensitivity at early growth as a possible
mechanism of maize adaptation to cold springs. J Exp Bot **70**: 2887–2904

Guillaume-Schöpfer D, Jaeger KE, Geng F, Doccula FG, Costa A, Webb AAR, Wigge
PA (2020) Ribosomes act as cryosensors in plants. bioRxiv. doi:
10.1101/2020.12.07.414789

Guo J, Sun B, He H, Zhang Y, Tian H, Wang B (2021) Current understanding of bhlh
transcription factors in plant abiotic stress tolerance. Int J Mol Sci. doi:
10.3390/ijms22094921

Hodges DM, DeLong JM, Forney CF, Prange RK (1999) Improving the thiobarbituric acid-
reactive-substances assay for estimating lipid peroxidation in plant tissues containing
anthocyanin and other interfering compounds. Planta **207**: 604–611

Holcik M, Sonenberg N (2005) Translational control in stress and apoptosis. Nat Rev Mol
Cell Biol **6**: 318–327

Hüner N, Öquist G, Sarhan F (1998) Energy balance and acclimation to light and cold.
Trends Plant Sci **3**: 224–230

IPCC (2018) Global warming of 1.5° C: an IPCC special report on the impacts of global
warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission
pathways, in the context of strengthening the global response to the threat of climate
change. Geneva, Switzerland

Janowiak F, Luck E, Dörffling K (2003) Chilling tolerance of maize seedlings in the field
during cold periods in spring is related to chilling-induced increase in abscisic acid level. J
Agron Crop Sci **189**: 156–161

Janowiak F, Maas B, Dörffling K (2002) Importance of abscisic acid for chilling tolerance of
maize seedlings. J Plant Physiol **159**: 635–643

Jompuk C, Fracheboud Y, Stamp P, Leipner J (2005) Mapping of quantitative trait loci
associated with chilling tolerance in maize (Zea mays L.) seedlings grown under field
conditions. J Exp Bot **56**: 1153–1163

Landi M (2017) Commentary to: "Improving the thiobarbituric acid-reactive-substances
assay for estimating lipid peroxidation in plant tissues containing anthocyanin and other
interfering compounds" by Hodges et al., Planta (1999) 207:604–611. Planta **245**: 1067

Langfelder P, Horvath S (2008) WGCNA: An R package for weighted correlation network
analysis. BMC Bioinformatics. doi: 10.1186/1471-2105-9-559

Li H, Yue H, Xie J, Bu J, Li L, Xin X, Zhao Y, Zhang H, Yang L, Wang J, et al (2021)
Transcriptomic profiling of the high-vigour maize (Zea mays L.) hybrid variety response to
cold and drought stresses during seed germination. Sci Rep **11**: 1–16

Li Z, Liu C, Zhang Y, Wang B, Ran Q, Zhang J (2019) The bHLH family member ZmPTF1
regulates drought tolerance in maize by promoting root development and abscisic acid
synthesis. J Exp Bot **70**: 5471–5486

Liu Y, Dang P, Liu L, He C (2019) Cold acclimation by the CBF–COR pathway in a
changing climate: Lessons from Arabidopsis thaliana. Plant Cell Rep **38**: 511–519

Liu H, Able AJ, Able JA (2022) Priming crops for the future: rewiring stress memory.
Trends Plant Sci **27**: 699–716

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol **15**: 1–21

Masclaux-Daubresse C, Purdy S, Lemaitre T, Pourtau N, Taconnat L, Renou J-P,
Wingler A (2007) Genetic Variation Suggests Interaction between Cold Acclimation and
Metabolic Regulation of Leaf Senescence. Plant Physiol **143**: 434–446

Ma Y, Tan R, Zhao J (2022) Chilling Tolerance in Maize: Insights into Advances—Toward
Physio-Biochemical Responses' and QTL/Genes' Identification. Plants. doi:
10.3390/plants11162082

Malnoë A (2018) Photoinhibition or photoprotection of photosynthesis? Update on the
(newly termed) sustained quenching component qH. Environ Exp Bot **154**: 123–133

Miedema P (1982) The effects of low temperature on Zea mays. Advances in Agronomy **35**:
93–128

Mukhopadhyay P, Tyagi AK (2015) OsTCP19 influences developmental and abiotic stress
signaling by modulating ABI4-mediated pathways. Sci Rep **5**: 1–12

Muñoz A, Castellano MM (2012) Regulation of translation initiation under abiotic stress
conditions in plants: Is it a conserved or not so conserved process among eukaryotes?
Comp Funct Genomics. doi: 10.1155/2012/406357

Ortiz-Lopez A, Nie GY, Ort DR, Baker NR (1990) The involvement of the photoinhibition of
photosystem II and impaired membrane energization in the reduced quantum yield of carbon
assimilation in chilled maize. Springer-Verlag

Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C (2017) Salmon provides fast and
bias-aware quantification of transcript expression. Nat Methods **14**: 417–419

Qin F, Sakuma Y, Li J, Liu Q, Li Y, Shinozaki K, Yamaguchi-Shinozaki, Kazuko (2004)
Cloning and Functional Analysis of a Novel DREB1/CBF Transcription Factor Involved in
Cold-Responsive Gene Expression in Zea mays L. Plant Cell Physiol **45**: 1042–1052

R Core Team (2022) R: A language and environment for statistical computing. R Found.
Stat. Comput. Vienna, Austria., https://www.r-project.org/.

Ramirez VE, Poppenberger B (2020) Modes of Brassinosteroid Activity in Cold Stress
Tolerance. Front Plant Sci. doi: 10.3389/fpls.2020.583666

Revilla P, Butrón A, Cartea M, Malvar R, Ordás A (2005) Breeding for cold tolerance. *In* M
Ashraf, PJC Harris, eds, Abiotic stresses. Plant resistance through breeding and molecular
approaches. The Haworth Press, Inc., USA, pp 301–398

Ritchie RJ (2008) Universal chlorophyll equations for estimating chlorophylls a, b, c, and d
and total chlorophylls in natural assemblages of photosynthetic organisms using acetone,
methanol, or ethanol solvents. Photosynthetica **46**: 115–126

Ritonga FN, Ngatia JN, Wang Y, Khoso MA, Farooq U, Chen S (2021) AP2/ER , an
important cold stress-related transcription factor family in plants: A review. Physiol Mol Biol
Plants **27**: 1953–1968

Riva-Roveda L, Escale B, Giauffret C, Périlleux C (2016) Maize plants can enter a
standby mode to cope with chilling stress. BMC Plant Biol **16**: 1–14

Rodríguez VM, Velasco P, Garrido JL, Revilla P, Ordás A, Butrón A (2013) Genetic
regulation of cold-induced albinism in the maize inbred line A661. J Exp Bot **64**: 3657–3667

Sanghera G, Wani S, Hussain W, Singh N (2011) Engineering cold stress tolerance in crop
plants. Curr Genomics **12**: 30–43

Savitch L V., Ivanov AG, Gudynaite-Savitch L, Huner NPA, Simmonds J (2011) Cold
stress effects on PSI photochemistry in Zea mays: Differential increase of FQR-dependent
cyclic electron flow and functional implications. Plant Cell Physiol **52**: 1042–1054

Sharma P, Sharma MMM, Patra A, Vashisth M, Mehta S, Singh B, Tiwari M, Pandey V
(2020) The role of key transcription factors for cold tolerance in plants. Transcr Factors
Abiotic Stress Toler Plants. doi: 10.1016/B978-0-12-819334-1.00009-5

Sowiński P, Fronk J, Jończyk M, Grzybowski M, Kowalec P, Sobkowiak A (2020) Maize
Response to Low Temperatures at the Gene Expression Level: A Critical Survey of
Transcriptomic Studies. Front Plant Sci **11**: 1–7

Sun X, Wang Y, Sui N (2018) Transcriptional regulation of bHLH during plant response to
stress. Biochem Biophys Res Commun **503**: 397–401

Szalai G, Majláth I, Pál M, Gondor OK, Rudnóy S, Oláh C, Vanková R, Kalapos B,
Janda T (2018) Janus-faced nature of light in the cold acclimation processes of maize. Front
Plant Sci **9**: 1–17

Tenaillon MI, Charcosset A (2011) A European perspective on maize history. Comptes
Rendus - Biol **334**: 221–228
1051
Thalhammer A, Pagter M, Hincha DK, Zuther E (2020) Measuring freezing tolerance of
leaves and rosettes: electrolyte leakage and chlorophyll fluorescence assays. *In* DK Hincha,
E Zuther, eds, Plant cold acclimation methods and protocols, Second Edition. Springer
Nature, pp 9–22

Tian T, Liu Y, Yan H, You Q, Yi X, Du Z, Xu W, Su Z (2017) AgriGO v2.0: A GO analysis
toolkit for the agricultural community, 2017 update. Nucleic Acids Res **45**: W122–W129

Wang C, Dong Y (2009) Overexpression of maize ZmDBP3 enhances tolerance to drought
and cold stress in transgenic Arabidopsis plants. Biologia (Bratisl) **64**: 1108–1114

Wang C, Yang Q, Yang Y (2011) Characterization of the ZmDBP4 gene encoding a
CRT/DRE-binding protein responsive to drought and cold stress in maize. Acta Physiol Plant
33: 575–583

Wang H, Wang H, Shao H, Tang X (2016) Recent Advances in Utilizing Transcription
Factors to Improve Plant Abiotic Stress Tolerance by Transgenic Technology. Front Plant
Sci **7**: 1–13

Wang ST, Sun XL, Hoshino Y, Yu Y, Jia B, Sun ZW, Sun MZ, Duan XB, Zhu YM (2014)
MicroRNA319 positively regulates cold tolerance by targeting OsPCF6 and OsTCP21 in rice
(Oryza sativa L.). PLoS One **9**: 1–12

Waters AJ, Makarevitch I, Noshay J, Burghardt LT, Hirsch CN, Hirsch CD, Springer NM
(2017) Natural variation for gene expression responses to abiotic stress in maize. Plant
Journal **89**: 706–717

Wingler A, Roitsch T (2008) Metabolic regulation of leaf senescence: interactions of sugar
signalling with biotic and abiotic stress responses. Plant Biol **10**: 50–62

Ying J, Lee EA, Tollenaar M (2002) Response of leaf photosynthesis during the grain-filling
period of maize to duration of cold exposure, acclimation, and incident PPFD. Crop Sci **42**:
1164–1172

Yoshida S, Satake T, Mackill D (1981) High temperature stress in rice. IRRI Research
Paper Series **67**: 1–15

Zeng R, Li Z, Shi Y, Fu D, Yin P, Cheng J, Jiang C, Yang S (2021) Natural variation in a
type-A response regulator confers maize chilling tolerance. Nat Commun **12**: 1–13

Zhao W, Zhao H, Wang H (2022) Research progress on the relationship between leaf
senescence and quality, yield and stress resistance in horticultural plants. Front Plant Sci
13:1044500: 1–12

Zheng Y, Jiao C, Sun H, Rosli HG, Pombo MA, Zhang P, Banf M, Dai X, Martin GB,
Giovannoni JJ, et al (2016) iTAK: A Program for Genome-wide Prediction
and Classification of Plant Transcription Factors, Transcriptional Regulators, and Protein
Kinases. Mol Plant **9**: 1667–1670

Zhou P, Enders TA, Myers ZA, Magnusson E, Crisp PA, Noshay JM, Gomez-Cano F,
Liang Z, Grotewold E, Greenham K, et al (2022a) Prediction of conserved and variable
heat and cold stress response in maize using cis-regulatory information. Plant Cell **34**: 514–
534

Zhou X, Muhammad I, Lan H, Xia C (2022b) Recent Advances in the Analysis of Cold
Tolerance in Maize. Front Plant Sci **13**: 1–15

1094
1095

1096

1097

1098

1099

1100   **Tables**

1101

1102   **Table 1: Transcription factors present in modules one and two. The No.**

1103   **column refers to the number of transcription factors of that type present in the**

1104   **list.** For a comprehensive list of gene IDs and "Other TFs", refer to Supp. dataset 7.

1105

Type of TF	No.	Description
		Module one
HB	9	Homeobox, homeodomain, leucine-zipper containing protein
MYB-related	4	MYB domain protein
C3H	4	Zinc finger (CCCH-type) family protein
GRAS	3	GRAS family transcription factor
NF	3	Nuclear transcription factor (NF-X1, NF-YA, NF-YB, NF-YC)
C2C2-GATA	3	Class IV zinc finger transcription factor
bZIP	1	Basic -leucine zipper protein
Other TFs	12	
Total TFs	**39**	
		Module two
MYB-related	10	MYB domain protein
HB	8	Homeobox, homeodomain, leucine-zipper containing protein
bZIP	8	Basic -leucine zipper protein
C3H	7	Zinc finger (CCCH-type) family protein
C2H2	7	C2H2-type zinc finger family protein
GeBP	5	Leucine-zipper transcription factors
HSF	5	Heat shock factor
NF	3	Nuclear transcription factor (NF-X1, NF-YA, NF-YB, NF-YC)
GRAS	2	GRAS family transcription factor
C2C2-GATA	2	Class IV zinc finger transcription factor
Other TFs	41	
Total TFs	**88**	

1124

1125 **Table 2: Transcription factors present in modules three and four**. The No.

1126 column refers to the number of transcription factors of that type present in the list.

1127 For a comprehensive list of gene IDs and "Other TFs", refer to Supp. dataset 7.

1128

Type of TF	No.	Description
		Module three
AP2/EREB	24	APETALA2/Ethylene Responsive Element Binding Factor domain protein
WRKY	19	WRKY family transcription factor
NAC	18	NAM, ATAF and CUC (NAC) domain-containing protein
C3H	15	Zinc finger (CCCH-type) family protein
GRAS	13	GRAS family transcription factor
C2H2	12	C2H2-type zinc finger family protein
C2C2-Dof	10	Zinc finger transcription factor, DNA-binding with one finger
MYB-related	8	MYB domain protein
Trihelix	8	Trihelix transcription factor
bZIP	5	Basic -leucine zipper protein
Other TFs	81	
Total TFs	**213**	
		Module four
AP2/EREB	9	APETALA2/Ethylene Responsive Element Binding Factor domain protein
MYB-related	9	MYB domain protein
C2C2-Dof	4	Zinc finger transcription factor, DNA-binding with one finger
bZIP	4	Basic -leucine zipper protein
NAC	3	NAM, ATAF and CUC (NAC) domain-containing protein
C3H	3	Zinc finger (CCCH-type) family protein
GRAS	2	GRAS family transcription factor
Trihelix	2	Trihelix transcription factor
WRKY	1	WRKY family transcription factor
Other TFs	41	
Total TFs	**78**	

1129

1130

1131 **Table 3: The transcription factor binding motifs enriched in the promoters of**
1132 **genes in modules one and two.** The No. column refers to the number of genes
1133 with that specific TF binding sight enriched. For CIS-BP IDs, gene IDs and
1134 enrichment data refer to Supp. dataset 8.

TF Family name	No.	Description
Module one		
bHLH	10	Basic helix-loop-helix
bZIP	10	Basic -leucine zipper protein
SBP	3	SQUAMOSA promoter-binding protein
HSF	1	Heat shock factor
Myb/SANT	1	MYB domain protein
Other	3	
Total enriched TF motifs	**28**	
Module two		
bHLH	17	Basic helix-loop-helix
bZIP	13	Basic -leucine zipper protein
HSF	6	Heat shock factor
NAC	1	NAM, ATAF and CUC transcription factors
Other	3	
Total enriched TF motifs	**40**	

1135

1136

1137 **Table 4: The transcription factor binding motifs enriched in the promoters of**
1138 **genes in modules three and four.** The No. column refers to the number of
1139 with that specific TF binding sight enriched. For CIS-BP IDs, gene IDs and
1140 enrichment data refer to Supp. dataset 8.

TF Family name	No.	Description
Module three		
AP2/EREB	24	APETALA2/Ethylene Responsive Element Binding Factor domain protein
TCP	9	Teosinte branched1/Cincinnata/proliferating cell factor family
Myb/SANT	5	MYB domain protein
bZIP	4	Basic -leucine zipper protein
bHLH	2	Basic helix-loop-helix
WRKY	2	WRKYGQK motif and a zinc-finger region
Other	7	
Total enriched TF motifs	**53**	
Module four		
bHLH	7	Basic helix-loop-helix
Myb/SANT	4	MYB domain protein
bZIP	2	Basic -leucine zipper protein
NAC	2	NAM, ATAF and CUC transcription factors
E2F	1	E2 factor family transcription factor
TCP	1	B-box binding protein
GATA	1	Teosinte branched1/Cincinnata/proliferating cell factor family
Total enriched TF motifs	**18**	

1141

1142 **Table 5:** Information on the candidate transcription factors selected based on their
1143 presence in modules one, two, three or four and an enriched transcription factor
1144 binding motif in the promoters of the other genes in the module. The references
1145 column includes work illustrating the role of each TF, or the TF family in which they
1146 belong,

GeneID	Gene name	Functional annotation	Module	kME for module	Rank in SEA	Previous studies
Zm00001eb339360	bHLH99	Basic helix-loop-helix transcription factor	2	0.95	19/40	- Li et al ., 2019 - Li et al ., 2021
Zm00001eb301280	HSF4	Heat stress transcription factor 4	2	0.91	1/40	- Sowinski et al ., 2020 - Zhou et al ., 2021 - Lui et al ., 2022
Zm00001eb037480	TCP21	TCP-transcription factor 21	3	0.95	14/53	- Ding et al , 2019 - Danisman, 2016
Zm00001eb364110	TCP20	TCP-transcription factor 20	3	0.91	17/53	- Ding et al , 2019 - Danisman, 2016
Zm00001eb145420	EREB52	AP2-EREBP-transcription factor 52	3	0.95	4/53	- Ritonga et al ., 2021 - Zhou et al ., 2022 - Qin et al ., 2004 - Wang & Dong, 2009 - Wang, Yang & Yang, 2011

1147 during chilling and/or high light responses

1148

Figures legends

Figure 1: Global transcriptome responses to the chilling and high light treatments. A, principal component analysis (PCA) plot of all 48 samples (24 control, 8 chilling, 8 chilling plus high light (Chilling + HL), 8 high light (HL)) showing how the transcriptome data separate based on treatment and accession. **B**, PCA plot of only B73 samples (24 samples). **C**, PCA plot of only F7 samples (24 samples). For all PCA plots, transcripts per million (TPM) counts from Salmon alignments were filtered for genes with at least 36 counts across the 48 samples for **A**, and genes with at least 18 counts across the 24 samples for **B** and **C**. The filtered TPM data was then normalised using variance stabilising transformation (vst) to estimate the dispersion trend across samples. The B73 and F7 accessions are represented as a circle and triangle respectively. Chilling, chilling plus high light and high light treated samples are represented in blue, yellow, and red respectively and untreated samples in grey. **D**, number of significantly (p-adj. < 0.05) differently expressed genes (DEGs) in each treatment for each accession determined by a DESeq2 analysis. Up-regulated genes are represented by white bars, and down-regulated genes by grey bars. Numbers above each bar indicate the count of DEGs represented by the bar. **E**, Venn diagram showing the extent of overlap between DEGs in each treatment in B73. **F**, Venn diagram showing the extent of overlap between DEGs in each treatment in F7.

Figure 2: The expression of genes which indicate improved stress tolerance of F7 compared to B73. The Log2 Fold Change in expression of selected stress tolerance-related genes which show opposite patterns of expression in F7 compared to B73. Log2 Fold Change and p-adj values are from a DESeq2 analysis of the transcriptome data. Stars above bars indicate a statistically significant difference in Log2 Fold Change when compared with the untreated control (p-adj < 0.05 considered significant). Black and white bars indicate the B73 and F7 accessions respectively. DESeq2 and TPM values for each gene can be found in Supp. dataset 1 and 2. HL = High light. *ZEP2: ZEAXANTHIN EPOXIDASE 2* (Zm00001eb423570), *VDE3b: VIOLAXANTHIN DE-EPOXIDASE 3B* (Zm00001eb361500), *CRR1: CYTOKININ RESPONSE REGULATOR 1* (Zm00001eb066570), *MPK8: MAP KINASE 8* (Zm00001eb358890).

Figure 3: Eigengene expression of four modules from a WGCNA performed on the transcriptome data and GO enrichment analyses on the genes in the modules. A, **B**, **C** and **D** show the eigengene values representing the gene expression profiles of four out of the eleven modules identified by WGCNA. Numbers in brackets represent the number of

genes in each of the modules. Lists of genes in each module can be found in Supp. dataset 4. **E**, GO terms of interest enriched in genes in module one and module two. **F**, GO terms of interest enriched in genes in module three and module four. For **E** and **F**, the significance of each GO term enrichment is represented as -log10FDR, with FDR representing false discovery rate.

Figure 4: Heatmap and individual gene plots showing that the addition of high light to chilling treatment mitigates the repressive effect of chilling on gene expression in modules one and two. A, a heatmap of the Log2 Fold Change (FC) in expression of genes found in modules one and two with the "Photosynthesis" GO term (82 genes from module one and 75 genes from module two). **B**, the Log2 Fold Change in expression of selected genes found in modules one or two involved in photosynthesis (*LCHB1, LHCB2*), NPQ (*PSBS*), carbon fixation (*PPDK1, PEPC1, RBCS*), xanthophyll metabolism (*VDE1, VP7*), chlorophyll metabolism (*CHAO2*), membrane integrity (*FAD6*), ROS scavenging (*SOD14*) and sucrose metabolism (*GAPA*). Log2 Fold Change and *p*-adj values are from a DESeq2 analysis of the transcriptome data. Stars above bars indicate a statistically significant difference in Log2 Fold Change when compared with the untreated control (*p*-adj < 0.05 considered significant). Black and white bars indicate the B73 and F7 accessions respectively. DESeq2 and TPM values for each gene can be found in Supp. Dataset 1 and 2. *LCHB1: LIGHT HARVESTING CHLOROPHYLL A/B BINDING PROTEIN 1* (Zm00001eb296090)*, LHCB2: LIGHT HARVESTING CHLOROPHYLL A/B BINDING PROTEIN 2* (Zm00001eb320890)*, PSBS: PHOTOSYSTEM II SUBUNIT* (Zm00001eb146510)*, PPDK1: PYRUVATE, PHOSPHATE DIKINASE 1* (Zm00001eb287770)*, PEPC1: PHOSPHOENOLPYRUVATE CARBOXYLASE 1* (Zm00001eb383680)*, RBCS: RUBISCO SMALL SUBUNIT* (Zm00001eb092540)*, VDE1: VIOLAXANTHIN DE-EPOXIDASE 1* (Zm00001eb085840)*, VP7: VIVIPAROUS 7* (Zm00001eb234930)*, CHAO2: CHLOROPHYLLIDE A OXYGENASE 2* (Zm00001eb362510)*, FAD6: FATTY ACID DESATURASE 6* (Zm00001eb041150)*, SOD14: SUPEROXIDE DISMUTASE 14* (Zm00001eb039000)*, GAPA: GLYCERALDEHYDE 3-PHOSPHATE DEHYDROGENASE SUBUNIT A* (Zm00001eb080840).

Figure 5: Heatmap and individual gene plots showing that the addition of high light to chilling treatment mitigates the inducing effect of chilling on gene expression in modules three and four. A, a heatmap of the Log2 Fold Change (FC) in expression of genes in modules three and four with the "regulation of transcription, DNA-templated" GO term (604 genes from module three and 228 genes from module four). **B**, the Log2 Fold Change in expression of selected genes found in modules three or four involved in ABA

signalling (*SnRK2.9, CDPK4, CDPK7, PYL8, PYL13*), stress responsive transcription (*DREB1.2, DREB1.7*), regulation of translation (*AGO1A, AGO1B*) and leaf senescence (*S40-1, S40-2, S40-4*). Log2 Fold Change and *p*-adj values are from a DESeq2 analysis of the transcriptome data. Stars above bars indicate a statistically significant difference in Log2 Fold Change when compared with the untreated control (*p*-adj < 0.05 considered significant). Black and white bars indicate the B73 and F7 accessions respectively. DESeq2 and TPM values for each gene can be found in Supp. dataset 1 and 2. *SNRK2.9: SNF1-RELATED PROTEIN KINASE 2.9* (Zm00001eb051510), *CDPK4: CALCIUM DEPENDENT PROTEIN KINASE 4* (Zm00001eb428080), *CDPK7: CALCIUM DEPENDENT PROTEIN KINASE 7* (Zm00001eb187630), *PYL8: PYRABACTIN RESISTANCE-LIKE PROTEIN 8* (Zm00001eb350630), *PYL13: PYRABACTIN RESISTANCE-LIKE PROTEIN 13* (Zm00001eb204180), *DREB1.2: DEHYDRATION-RESPONSIVE ELEMENT-BINDING PROTEIN 1.2* (Zm00001eb318890), *DREB1.7: DEHYDRATION-RESPONSIVE ELEMENT-BINDING PROTEIN 1.7* (Zm00001eb269420*), AGO1A: ARGONAUTE 1A* (Zm00001eb267430), *AGO1B: ARGONAUTE LINKER 2 DOMAIN* (Zm00001eb428910), *S40-1: SENESCENCE REGULATOR S40-1* (Zm00001eb077160), *S40-2: SENESCENCE REGULATOR S40-2* (Zm00001eb083770), *S40-4: SENESCENCE REGULATOR S40-4* (Zm00001eb221600).

Figure 6: Identification of candidate transcription factors driving the mitigating effect of high light on the transcriptomic response to chilling stress. The heatmaps show the 20 most enriched transcription factor binding motifs present in the promoters of genes in modules one, two, three and four. Enriched motifs were identified using a Simple Enrichment Analysis (SEA) on sequences 1000 bp upstream of the ATG transcription start site. The TF family of each TF binding motif is indicated next to the corresponding box in each heatmap. Font colour groups TFs according to TF family. The degree of motif enrichment is depicted through -Log$_{10}$(*p-value*). The transcription factors highlighted with a black box indicate those which had enriched binding motifs identified in the SEA, were also present in the genes list for the corresponding module and had a module kME greater than 0.9. Bar plots show the expression of the identified TF candidates from a DESeq2 analyses of the transcriptome data. Stars above bars indicate a statistically significant difference in Log2 Fold Change when compared with the untreated control (*p*-adj < 0.05 considered significant). Black and white bars indicate the B73 and F7 accessions respectively. DESeq2 and TPM values for each gene can be found in Supp. dataset 1 and 2. *HSF4*: *HEAT STRESS TRANSCRIPTION FACTOR 4* (Zm00001eb301280)*, BHLH99*: *BASIC HELIX-LOOP-HELIX TRANSCRIPTION FACTOR 99* (Zm00001eb339360)*, EREB52: AP2-EREBP-TRANSCRIPTION FACTOR 52* (Zm00001eb145420)*, TCP21: TEOSINTE-BRANCHED1/CYCLOIDEA/PCF*

TRANSCRIPTION FACTOR 21 (Zm00001eb037480), *TCP20: TEOSINTE-*

BRANCHED1/CYCLOIDEA/PCF TRANSCRIPTION FACTOR 20 (Zm00001eb364110).

Figure 7: High light mitigates changes in F_v/F_m, NPQ and relative injury resulting from

chilling stress. A-I, leaf discs sampled from maize plants that had previously been given

chilling, chilling plus high light (HL), high light (HL) treatments, or corresponding control

treatments, were exposed to 10 minutes of light and 12 minutes of dark recovery. **A**, **B**, **C**,

Dark-adapted F_v/F_m; **D**, **E**, **F**, NPQ level at the end of 10 min light; **G**, **H**, **I**, NPQ level at the

end of 12 min dark recovery. **J**, **K**, **L**, relative injury of leaf discs sampled from maize plants

that had previously been given chilling, chilling plus high light, high light treatments, or

corresponding control treatments. Each treatment was analysed with respect to its own

control (n=4). Results from two-way ANOVA are shown. There were no statistically

significant interactions between accession and treatment.

1271

Figure 1: Global transcriptome responses to the chilling and high light treatments. A, principal component analysis (PCA) plot of all 48 samples (24 control, 8 chilling, 8 chilling plus high light (Chilling + HL), 8 high light (HL)) showing how the transcriptome data separate based on treatment and accession. **B**, PCA plot of only B73 samples (24 samples). **C**, PCA plot of only F7 samples (24 samples). For all PCA plots, transcripts per million (TPM) counts from Salmon alignments were filtered for genes with at least 36 counts across the 48 samples for **A**, and genes with at least 18 counts across the 24 samples for **B** and **C**. The filtered TPM data was then normalised using variance stabilising transformation (vst) to estimate the dispersion trend across samples. The B73 and F7 accessions are represented as a circle and triangle respectively. Chilling, chilling plus high light and high light treated samples are represented in blue, yellow, and red respectively and untreated samples in grey. **D**, number of significantly (p-adj. < 0.05) differently expressed genes (DEGs) in each treatment for each accession determined by a DESeq2 analysis. Up-regulated genes are represented by white bars, and down-regulated genes by grey bars. Numbers above each bar indicate the count of DEGs represented by the bar. **E**, Venn diagram showing the extent of overlap between DEGs in each treatment in B73. **F**, Venn diagram showing the extent of overlap between DEGs in each treatment in F7.

Figure 2: The expression of genes which indicate improved stress tolerance of F7 compared to B73. The Log2 Fold Change in expression of selected stress tolerance-related genes which show opposite patterns of expression in F7 compared to B73. Log2 Fold Change and *p*-adj values are from a DESeq2 analysis of the transcriptome data. Stars above bars indicate a statistically significant difference in Log2 Fold Change when compared with the untreated control (*p*-adj < 0.05 considered significant). Black and white bars indicate the B73 and F7 accessions respectively. DESeq2 and TPM values for each gene can be found in Supp. dataset 1 and 2. HL = High light. *ZEP2: ZEAXANTHIN EPOXIDASE 2* (Zm00001eb423570), *VDE3b: VIOLAXANTHIN DE-EPOXIDASE 3B* (Zm00001eb361500), *CRR1: CYTOKININ RESPONSE REGULATOR 1* (Zm00001eb066570), *MPK8: MAP KINASE 8* (Zm00001eb358890).

Figure 3: Eigengene expression of four modules from a WGCNA performed on the transcriptome data and GO enrichment analyses on the genes in the modules. A, B, C and D show the eigengene values representing the gene expression profiles of four out of the eleven modules identified by WGCNA. Numbers in brackets represent the number of genes in each of the modules. Lists of genes in each module can be found in Supp. dataset 4. E, GO terms of interest enriched in genes in module one and module two. F, GO terms of interest enriched in genes in module three and module four. For E and F, the significance of each GO term enrichment is represented as -log10FDR, with FDR representing false discovery rate.

Figure 4: Heatmap and individual gene plots showing that the addition of high light to chilling treatment mitigates the repressive effect of chilling on gene expression in modules one and two. **A**, a heatmap of the Log2 Fold Change (FC) in expression of genes found in modules one and two with the "Photosynthesis" GO term (82 genes from module one and 75 genes from module two). **B**, the Log2 Fold Change in expression of selected genes found in modules one or two involved in photosynthesis (*LCHB1, LHCB2*), NPQ (*PSBS*), carbon fixation (*PPDK1, PEPC1, RBCS*), xanthophyll metabolism (*VDE1, VP7*), chlorophyll metabolism (*CHAO2*), membrane integrity (*FAD6*), ROS scavenging (*SOD14*) and sucrose metabolism (*GAPA*). Log2 Fold Change and *p*-adj values are from a DESeq2 analysis of the transcriptome data. Stars above bars indicate a statistically significant difference in Log2 Fold Change when compared with the untreated control (*p*-adj < 0.05 considered significant). Black and white bars indicate the B73 and F7 accessions respectively. DESeq2 and TPM values for each gene can be found in Supp. Dataset 1 and 2. *LCHB1: LIGHT HARVESTING CHLOROPHYLL A/B BINDING PROTEIN 1* (Zm00001eb296090), *LHCB2: LIGHT HARVESTING CHLOROPHYLL A/B BINDING PROTEIN 2* (Zm00001eb320890), *PSBS: PHOTOSYSTEM II SUBUNIT* (Zm00001eb146510), *PPDK1: PYRUVATE, PHOSPHATE DIKINASE 1* (Zm00001eb287770), *PEPC1: PHOSPHOENOLPYRUVATE CARBOXYLASE 1* (Zm00001eb383680), *RBCS: RUBISCO SMALL SUBUNIT* (Zm00001eb092540), *VDE1: VIOLAXANTHIN DE-EPOXIDASE 1* (Zm00001eb085840), *VP7: VIVIPAROUS 7* (Zm00001eb234930), *CHAO2: CHLOROPHYLLIDE A OXYGENASE 2* (Zm00001eb362510), *FAD6: FATTY ACID DESATURASE 6* (Zm00001eb041150), *SOD14: SUPEROXIDE DISMUTASE 14* (Zm00001eb039000), *GAPA: GLYCERALDEHYDE 3-PHOSPHATE DEHYDROGENASE SUBUNIT A* (Zm00001eb080840).

Figure 5: Heatmap and individual gene plots showing that the addition of high light to chilling treatment mitigates the inducing effect of chilling on gene expression in modules three and four. **A**, a heatmap of the Log2 Fold Change (FC) in expression of genes in modules three and four with the "regulation of transcription, DNA-templated" GO term (604 genes from module three and 228 genes from module four). **B**, the Log2 Fold Change in expression of selected genes found in modules three or four involved in ABA signalling (*SnRK2.9, CDPK4, CDPK7, PYL8, PYL13*), stress responsive transcription (*DREB1.2, DREB1.7*), regulation of translation (*AGO1A, AGO1B*) and leaf senescence (*S40-1, S40-2, S40-4*). Log2 Fold Change and *p*-adj values are from a DESeq2 analysis of the transcriptome data. Stars above bars indicate a statistically significant difference in Log2 Fold Change when compared with the untreated control (*p*-adj < 0.05 considered significant). Black and white bars indicate the B73 and F7 accessions respectively. DESeq2 and TPM values for each gene can be found in Supp. dataset 1 and 2. *SNRK2.9: SNF1-RELATED PROTEIN KINASE 2.9* (Zm00001eb051510), *CDPK4: CALCIUM DEPENDENT PROTEIN KINASE 4* (Zm00001eb428080), *CDPK7: CALCIUM DEPENDENT PROTEIN KINASE 7* (Zm00001eb187630), *PYL8: PYRABACTIN RESISTANCE-LIKE PROTEIN 8* (Zm00001eb350630), *PYL13: PYRABACTIN RESISTANCE-LIKE PROTEIN 13* (Zm00001eb204180), *DREB1.2: DEHYDRATION-RESPONSIVE ELEMENT-BINDING PROTEIN 1.2* (Zm00001eb318890), *DREB1.7: DEHYDRATION-RESPONSIVE ELEMENT-BINDING PROTEIN 1.7* (Zm00001eb269420), *AGO1A: ARGONAUTE 1A* (Zm00001eb267430), *AGO1B: ARGONAUTE LINKER 2 DOMAIN* (Zm00001eb428910), *S40-1: SENESCENCE REGULATOR S40-1* (Zm00001eb077160), *S40-2: SENESCENCE REGULATOR S40-2* (Zm00001eb083770), *S40-4: SENESCENCE REGULATOR S40-4* (Zm00001eb221600).

Figure 6: Identification of candidate transcription factors driving the mitigating effect of high light on the transcriptomic response to chilling stress. The heatmaps show the 20 most enriched transcription factor binding motifs present in the promoters of genes in modules one, two, three and four. Enriched motifs were identified using a Simple Enrichment Analysis (SEA) on sequences 1000 bp upstream of the ATG transcription start site. The TF family of each TF binding motif is indicated next to the corresponding box in each heatmap. Font colour groups TFs according to TF family. The degree of motif enrichment is depicted through -Log_{10}(*p-value*). The transcription factors highlighted with a black box indicate those which had enriched binding motifs identified in the SEA, were also present in the genes list for the corresponding module and had a module kME greater than 0.9. Bar plots show the expression of the identified TF candidates from a DESeq2 analyses of the transcriptome data. Stars above bars indicate a statistically significant difference in Log2 Fold Change when compared with the untreated control (*p*-adj < 0.05 considered significant). Black and white bars indicate the B73 and F7 accessions respectively. DESeq2 and TPM values for each gene can be found in Supp. dataset 1 and 2. *HSF4: HEAT STRESS TRANSCRIPTION FACTOR 4* (Zm00001eb301280), *BHLH99: BASIC HELIX-LOOP-HELIX TRANSCRIPTION FACTOR 99* (Zm00001eb339360), *EREB52: AP2-EREBP-TRANSCRIPTION FACTOR 52* (Zm00001eb145420), *TCP21: TEOSINTE-BRANCHED1/CYCLOIDEA/PCF TRANSCRIPTION FACTOR 21* (Zm00001eb037480), *TCP20: TEOSINTE-BRANCHED1/CYCLOIDEA/PCF TRANSCRIPTION FACTOR 20* (Zm00001eb364110).

Figure 7: High light mitigates changes in F_v/F_m, NPQ and relative injury resulting from chilling stress. A-I, leaf discs sampled from maize plants that had previously been given chilling, chilling plus high light (HL), high light (HL) treatments, or corresponding control treatments, were exposed to 10 minutes of light and 12 minutes of dark recovery. A, B, C, Dark-adapted F_v/F_m; D, E, F, NPQ level at the end of 10 min light; G, H, I, NPQ level at the end of 12 min dark recovery. J, K, L, relative injury of leaf discs sampled from maize plants that had previously been given chilling, chilling plus high light, high light treatments, or corresponding control treatments. Each treatment was analysed with respect to its own control (n=4). Results from two-way ANOVA are shown. There were no statistically significant interactions between accession and treatment.