Embracing firefly flash pattern variability with data-driven species classification

Owen Martin[1,2,†], Chantal Nguyen[2,†], Raphael Sarfati[2], Murad Chowdhury[1,2], Michael L. Iuzzolino[1], Dieu My T. Nguyen[1,2], Ryan M. Layer[1,2,*], Orit Peleg[1,2,3,*]

1. Department of Computer Science, University of Colorado Boulder, Boulder, CO, USA
2. BioFrontiers Institute, University of Colorado Boulder, Boulder, CO, USA
3. Santa Fe Institute, Santa Fe, NM, USA

*Correspondence: ryan.layer@colorado.edu, orit.peleg@colorado.edu
†These authors contributed equally to this work.

Abstract

Many nocturnally active fireflies use precisely timed bioluminescent patterns to identify mates, making them especially vulnerable to light pollution. As urbanization continues to brighten the night sky, firefly populations are under constant stress, and close to half of the species are now threatened. Ensuring the survival of firefly biodiversity depends on a large-scale conservation effort to monitor and protect thousands of populations. While species can be identified by their flash patterns, current methods require expert measurement and manual classification and are infeasible given the number and geographic distribution of fireflies. Here we present the application of a recurrent neural network (RNN) for accurate automated firefly flash pattern classification. Using recordings from commodity cameras, we can extract flash trajectories of individuals within a swarm and classify their species with a precision and recall of approximately seventy percent. In addition to scaling population monitoring, automated classification provides the means to study firefly behavior at the population level. We employ the classifier to measure and characterize the variability within and between swarms, unlocking a new dimension of their behavior. Our method is open source, and deployment in community science applications could revolutionize our ability to monitor and understand firefly populations.

1 Introduction

Nocturnal fireflies (Coleoptera: Lampyridae) have evolved a visually impressive light-based communication system under simultaneous evolutionary pressures to advertise their species, accentuate their sexual fitness, and avoid predation [1–3]. Using a luciferin-luciferase reaction within their abdomen [4], fireflies broadcast their species identity via light pulses [1, 5]. In many North American genera both sexes flash and must also encode biological sex in their signals [1, 6, 7]. Sympatric species must also produce distinguishable patterns to effectively communicate species identity [1, 5, 8]. Despite temperature-dependent and individual variability in these signals, each firefly species' flash behavior is typically characterized for identification purposes by a single, imprecise pictorial representation of a flash pattern standardized at a fixed temperature [7, 8].

This unique signaling system makes fireflies particularly susceptible to human-created population threats like light pollution [9–12]. The presence of artificial light at night (ALAN) interferes with fireflies' perception of conspecific signals and disrupts their communication timings, curtailing flash signaling behavior and preventing successful mating [10, 13]. In addition to light pollution, habitat degradation, pesticide use, water pollution, and climate change comprise some of the most serious environmental stressors causing declines in firefly populations across the globe [14]. Recent Red List assessments by the International Union for Conservation of Nature (IUCN) have identified that at least 14% of 132 firefly taxa in the United States alone are in danger of extinction [12], but

this value is a likely underestimate due to a lack of information on over half the species assessed [6, 12]. Further studies and fieldwork are urgently needed for monitoring changes in firefly abundances and for evaluating and mitigating the threats imperiling fireflies worldwide [6, 12]. These environmental threats are also shared by many other insect species experiencing dramatic declines [10, 15, 16]. Because of their charismatic nature and popular appeal, fireflies can serve as a flagship symbol to foster public attention toward this conservation crisis.

The major barrier to planning effective conservation efforts is the dearth of quantitative data on firefly populations [6, 12, 14]. The high-throughput population monitoring studies that provide the necessary data have not been performed for nearly all firefly species [6]. This gap is largely due to limitations of existing monitoring methodologies, which require the presence of human expert observers [6], are often subjective [6], and fail to capture individual variability present in the swarm. To address the data deficiency in current conservation efforts, we propose a scalable, automated population monitoring method that combines recent advances in stereoscopic filming, computer vision, and machine learning to classify individuals and quantify swarm-level dynamics. Our method starts with a field recording of a swarm using two consumer-grade cameras (Fig. 1A) and following procedures described previously in [17]. From this recording, we identify individual firefly trajectories and extract their flash-pattern time series (Fig. 1B). We then trained a recurrent neural network (RNN) on these data to accurately determine species identity from nothing more than the temporal differences in each species' flash pattern, achieving precision and recall of approximately 0.70 (Table 1). We additionally visualize the distinguishability of firefly flash patterns via a dimensionality reduction of the weights of the last hidden layer of the neural network, using t-distributed stochastic neighbor embedding (t-SNE). The t-SNE embedding reveals significant clustering by species, but multiple species can coincide in the same cluster, illustrating the intra-species variability that contributes to the difficulty of the problem (Fig. 1C). To our knowledge this is the first application of machine learning to firefly behavioral biology.

Using the most probable predictions of each species, we also provide estimates of flash phrase variability, including the first-ever quantitative characterization of *Bicellonycha wickershamorum* and *Photuris forresti*, two formerly data-deficient species which may be severely threatened [18]. Automated and data-driven methods like the one proposed here are essential to accurately identifying and monitoring firefly species in order to ensure their protection and long-term survival.

2 Results

Using our precise, high-throughput data acquisition method (Fig. 1A and Methods section 4.1), we recorded multiple firefly swarms comprising seven North American firefly species and totaling 73,141 flash pattern time-series. These data are a dramatic expansion from previously published results, which were primarily single characteristic patterns per species, and represent firefly swarms filmed each summer over the past three years. This large dataset enables us to leverage machine learning for species classification and characterization of intra-swarm variability.

Using an 80:10:10 train:validation:test split of these data, we trained a recurrent neural network (RNN) to produce class label predictions based on the temporal information in each sequence (see Methods section 4.2.1). The accuracy of these predictions was compared to those obtained from several alternate classifiers utilizing signal processing metrics (Table 1 and Methods section 4.3.3). We present ensemble results evaluating the model's performance on the test set for each method in Table 1. The RNN achieves the best performance in both categories, with an increase of approximately 20% and 60% higher precision and recall, respectively, over the second best classifier.

Using the previously published, manually gathered flash patterns ("literature references") (see

Figure 1: **(A)** Fireflies in their natural habitat **(top)** are filmed with two cameras arranged in a stereoscopic vision configuration [19]. Flash streaks in the resulting videos are triangulated and concatenated into trajectories (**center**, colored by time), based on proximity and velocity, which represent the time series (truncated series shown on **bottom**) of individual fireflies flashing as they move through the space. Each firefly sequence can be parameterized with three values, annotated in the bottom figure: the number of flashes; the flash length, or the average amount of consecutive time for which a flash is detected; and the inter-flash gap, the average amount of time from the end of a flash to the start of another flash. **(B)** Four example five-second flash sequence time series for each of the seven species in our study, labeled with the location and date of the corresponding recording. Our recurrent neural network is used to classify flash sequence time series; time series shown were selected from the top 100 sequences per species with the highest classification probabilities following the filtration process outlined in Methods section 4.4. **(C)** Two-dimensional t-SNE embedding of the output of the last hidden layer of the network, just before inference. Flash patterns are clustered by similarity, and the distinguishability of each species' characteristic flash pattern can be detected by the colored clusters. However, overlapping points do occur due to the variability of species' flash signals and the precision of the data-acquisition methodology (see Methods).

Methods section 4.3.1), we constructed three different classifiers that use the Jaccard index, dot product, and dynamic time warping (see Methods section 4.3.3) to compare our time series data to the literature references. We additionally constructed a support vector machine (SVM) that classified the time series data from the three flash pattern parameters (flash length, inter-flash gap, and the number of flashes) (Fig. 1A, bottom). These literature-based classifiers perform poorly, with the SVM achieving the highest precision and recall of 0.34 and 0.30, respectively (Table 1).

We attribute the poor performance of the literature reference methods to the ineffectiveness of a single reference sequence per species at capturing the intra-species variability inherent to a large dataset. To address this, we create "population reference" sequences by aggregating sequences of each species in our dataset, aimed at capturing the intra-species variability while preserving species-specific characteristics of the flash signal. We perform an 80:20 train:test split, generate population references from the training set, and classify the remaining test data. The performance of all four classifiers improves slightly by using population references, with dynamic time warping achieving the highest precision and recall of 0.43 and 0.58, respectively (see Table 1).

Metric	Precision		Recall	
	Literature	Population reference	Literature	Population reference
SVM	0.34	0.36 ± 0.00084	0.30	0.53 ± 0.0014
Jaccard index	0.23	0.41 ± 0.0011	0.26	0.48 ± 0.0019
Dot product	0.24	0.25 ± 0.0025	0.25	0.38 ± 0.0013
Dynamic time warping	0.27	0.43 ± 0.00040	0.27	0.58 ± 0.0013
RNN	**0.70 ± 0.19**		**0.71 ± 0.19**	

Table 1: Summary of signal classification methods and their performance on the firefly flash pattern dataset. $\pm$ values indicate standard error of the mean for each statistic.

In Fig. 2 we show confusion matrices for each ensemble to further illustrate the per-class capabilities and shortcomings of each method. While the dynamic time warping and dot product classification methods occasionally produce a high scoring class (see Methods Fig. 5F), the RNN results reveal a balanced, highly accurate classifier regardless of species, and as such is our recommendation for the algorithm of choice moving forward in this space. Per-species precision and recall metrics can be found in Methods Fig. 5.

By combining our extensive data set and effective RNN, we provide deeper insights into firefly behavior with a data-driven characterization of a swarm's signaling pattern. For each swarm recording, we aggregate the one hundred most confidently classified flash patterns (see Methods section 4.4) to produce empirical distributions for flash length, inter-flash gap, and the number of flashes for each species (Fig. 3). These distributions represent the first known quantification of firefly behavioral variability from data. Table 2 compares the mean and standard deviation of flash parameters from our data with the published literature values. Encouragingly, all the previously published numbers of flashes were statistically indistinguishable from our distributions. Half of the published inter-flash gaps and all the published flash lengths differed significantly from our distributions. These differences may be due to improved measurement accuracy (see below) or could represent behavioral differences between swarms, motivated by a changing climate or other environmental factors.

These data-driven distributions have several advantages over manual timekeeping. Firstly, representing a species' flash-based communication as distributions of signal parameters rather than a single pattern allows us to explore the extent of the temporal variability present in a population or across populations. We observe, for example, high variability in the number of flashes emitted by

Figure 2: Confusion matrices for each classification method. Species 1-7 are *Bicellonycha wicker-shamorum*, *Photuris bethaniensis*, *Photinus carolinus Photuris forresti*, *Photuris frontalis*, *Photinus knulli*, and *Photinus obscurellus*, respectively. **(A-E)** Ratios of true positives and false negatives along the horizontal, true positives and false positives along the vertical. Each square in the diagonal represents the recall for the class, and brighter colors indicate higher values as shown in the colorbar. A perfect classifier would consist of yellow squares at each diagonal position and dark purple off the diagonal.

P. frontalis, while the flash trains of *P. knulli* exhibit very low variability in the number of flashes (Fig. 3A). Meanwhile, both of these species display relatively tight distributions of inter-flash gap (Fig. 3B), suggesting that these species require precise timing of flashes for communication.

Our methods are more precise than manual timing in measuring the flash length of short flashes perhaps due to the cameras' high temporal resolution (30 Hz). Our results reveal that the flash length distributions for five species are shifted considerably toward shorter flash lengths than the published literature results (Fig. 3C); this discrepancy may potentially be a consequence of the lower temporal resolution inherent in manual timekeeping due to system lag.

Overall, the filtering process that produces data-driven characterizations of flash patterns in Fig. 3 preserves the most salient characteristics of each species' signaling behavior while also illustrating the intraspecies behavioral variability. This variability may be substantial in only some of the 3 parameters: for example, *P. obscurellus* demonstrates low variability in the number of flashes but the largest variability in flash length out of the 7 species studied. In comparison, the flash lengths of *P. frontalis* sequences are all extremely similar, but the number of flashes in a train can vary greatly.

We note that these characterizations only represent the locations and conditions where the underlying data were recorded; these probability distributions may be different for data taken in different habitats, at different temperatures, or at different times. By deploying our data acquisition procedure to more sites, we will be able to better characterize the rich extent of a firefly species' communication.

Previous studies have examined the effect of environmental factors such as artificial light at night (ALAN) on firefly signaling by quantifying the reduction in flash rate or the number of flashing individuals [10]. Our method of generating data-driven characterizations will enable a more acute investigation of how ALAN and other factors affect firefly flash pattern parameters down to an individual level. In addition, we can also investigate whether responses to environmental factors are encoded in behavioral variability: it is unknown whether the variability in flash behavior changes with light pollution and the resulting implications for firefly signal processing and mating success.

Figure 3: Data-driven characterization of flash patterns. For each species, the 100 sequences with the highest classification probabilities are used to characterize flash signals. This represents a data filtering procedure that extracts the most salient properties of each species' flash behavior while retaining inherent intraspecies behavioral variability. Probability distributions of the **(A)** number of flashes, **(B)** inter-flash gap in seconds, and **(C)** length of flash in seconds are shown for each species, normalized to sum to 1 under the interval, for both the raw (transparent bars) and filtered (opaque bars) data.

Species	Number of flashes		Inter-flash gap (s)		Flash length (s)	
	Data	Literature	Data	Literature	Data	Literature
B. wickershamorum	5.3 ± 3.5	n/a	0.80 ± 0.09	n/a	0.11 ± 0.03	n/a
P. bethaniensis	2.1 ± 0.3	2	0.23 ± 0.08	0.53	0.07 ± 0.03	0.27
P. carolinus	8.8 ± 2.4	9	0.39 ± 0.09	0.37	0.10 ± 0.04	0.23
P. forresti	5.9 ± 2.4	5	0.46 ± 0.22	0.10	0.09 ± 0.04	0.16
P. frontalis	11.8 ± 5.2	10	0.64 ± 0.03	0.87	0.04 ± 0.01	0.13
P. knulli	3.0 ± 0.2	3	0.30 ± 0.02	0.32	0.10 ± 0.02	0.20
P. obscurellus	3.1 ± 1.5	3	0.32 ± 0.11	0.28	0.22 ± 0.09	0.22

Table 2: Mean and standard deviation of the number of flashes, inter-flash gap, and flash length from the filtered data for each species, along with the corresponding literature values extracted via methodology shown in Fig. 4.

3 Discussion and future directions

We have presented a new paradigm for standardized firefly data acquisition and automated species detection, using machine learning to classify firefly species from their flash patterns for the first time. Our method is robust to behavioral variability and embraces it to pursue a comprehensive data-driven characterization of firefly flash patterns. Leveraging the discriminatory power of the classifier, we can also extract the most representative sequences for each species from the dataset and provide quantitative and automatically generated characterizations of flash phrases. These characterizations align well with known descriptions, where data are available, and also provide statistical clarity for several endangered species such as *Photuris bethaniensis* and *Photinus*

knulli for which data are scarce. Additionally, our flash sequence characterization of *Bicellonycha wickershamorum* and *Photuris forresti* represent the **first known** quantitative description of the flash pattern for these vulnerable species [6]. Given that there are 171 known North American firefly species [6] - many of which have yet to have their flash patterns quantitatively described - the logical next step is a large-scale expansion of the dataset. The methodology we have developed for acquiring flash sequence data relies only on relatively affordable and extremely portable consumer camera equipment, making it readily adaptable for use by other researchers as well as citizen science volunteers [20]. This will facilitate rapid deployment of cameras across the continent to known firefly hot spots. Each deployment can add new species to the corpus, where applicable, and enhance characterization of flash phrases, expanding our knowledge of firefly behavior and empowering researchers with easy identification tools. After sufficient trajectories are acquired for a new species, the neural network can be retrained to recognize, differentiate, and classify an enlarged species set. There may be space and time dedicated to fine-tuning the training process as more and more species enter the dataset, but given the variability already encapsulated and the ability of the model to successfully classify at this stage, we predict this will be a vanishing obstacle as the process continues.

Past work has, with good reason, summarized and standardized the behavioral variability (at a particular temperature, usually 20°C [8]), resulting in discrete measurements for the relevant statistics that are used as references for species identification in tandem with physical observation of captured specimens. However, standardization is not a practice that the neural network classification can tolerate: the variability is enough that a *single* reference point will often lead the classification in the wrong direction (see Fig. SI2). These reference measurements may introduce sampling biases caused by observing different individuals within the same swarm, using different methods, and observing under different conditions. Here, we accumulate data for each species using uniform methodology, mitigating observer biases and capturing swarm level behavior from which individuals can be isolated.

Our automated trajectory detection method reliably measures the flash lengths and inter-flash gaps (Fig 3, Table 2) of firefly signals, especially as they are extracted from thousands of precise observations of the same species across multiple locations and nights of filming. We show in (Fig 3C) that human observers tend to over-estimate flash length, which could be attributed to the reaction latency in a human observer between seeing the end of a flash and pressing a stopwatch. This problem is eliminated by the camera capture procedure, which is accurate to a frame resolution of 30 Hz, a significant improvement over measured human reaction times of order 0.2 s [21]. However, some limitations with the automated method exist. For example, it is possible that a given trajectory might not capture the entire flash pattern of an individual, due to overlapping signals from nearby fireflies, prolonged dark periods between series of successive flashes, and possible visual obstruction of blinks from the environment. Hence, it can be difficult to estimate the total count of flashes in a firefly's signal, and the dark period between flash phrases is fully lost in most trajectories. This can be mitigated by generating trajectories using *a priori* assumptions of the gap lengths between flashes, but it is vital to the success of the downstream sequence classification that information about the species is withheld from the trajectory generation process. This is further mitigated in the characterization process by filtering only the classifier's most probable flash patterns into a subset of data that should represent the best trajectories in the overall dataset for any given species. Combining our method with continued observations by human experts for the most complete characterization of any given species' flashing behavior may be the optimal approach in the short term, especially in species known to have dark periods that exceed one second in length between flash phrases, or species that produce a single flash. However, the advancements presented here will indisputably accelerate this process and should only improve in effectiveness over time.

Future work concerns scalability of the project and applications of evolutionary and conservation biology. If, as the dataset gains more species, it becomes *more* difficult to discern between similar species, it is easy to pivot to an CRNN (convolutional recurrent neural network) structure [22], using spatial information extracted directly from the video recording to add additional features to the model that may serve to increase differentiability. As the capabilities increase, the classifier can be integrated into community-facing applications like iNaturalist [23], which in turn will enable faster data-training-classification loops and a widespread expansion of the model's reach. Our standardized, user-friendly data acquisition procedure can be adopted by volunteers and researchers to grow the dataset, with an eye toward recording flash sequences from data-deficient species.

The model also has utility when asking questions about firefly flash signal evolution: highly indistinguishable species may be investigated as evolutionary examples of convergent sequence design. High values off the diagonal in Fig. 2 represent relatively indistinguishable species, while those with high values on the diagonal are most distinguishable; correlating these results with geographical information can provide insight into character displacement. Those species that are indistinguishable on the basis of temporal flash information may be more likely to be geographically isolated, and can thus be easy to differentiate in our model with the additional application of location information. The evolution of sequence distinguishability, especially under competing pressures for avoiding predation, is a rich area of active research [24] where this model will certainly be useful.

However, the most important and immediate application is direct deployment of cameras to areas containing critically endangered or vulnerable firefly species, such as *P. bethaniensis* and *P. knulli*, for automatic population monitoring. Integration of the inexpensive data gathering procedure with the classification software presented here can soon take on a life of its own as a valuable tool for automatic, quantitative firefly population monitoring. Swarms of endangered species can be detected, identified, and assessed for population density or behavioral changes in the presence of encroaching light pollution or other conservation risks. If new data-driven characterizations of a species fall outside of what has been previously observed, it may be a call for alarm: additionally, if the model detects less individuals of a particular species in a known population area, that also may be a call for alarm. Data-deficient species of conservation concern can and should be targets for the next round of deployments to quickly gain a baseline characterization of their flash dynamics and enable automated population monitoring. This work constitutes a crucial advancement in the collective conservation effort to retain the amazing biodiversity within the *Lampyridae* family, helping to preserve the wonder and fascination that accompanies their bioluminescence for generations to come.

4 Methods

4.1 Acquisition of flash sequence data

To extract flash sequence data, we perform 3D reconstruction of firefly swarms based on stereoscopic video recordings. Recordings were conducted at specific locations across the country where certain species were known to emerge [20]. Field researchers placed two spherical (360°) GoPro Max cameras at known distances from each other on a level surface (Fig. 1A). Recordings started at dusk when the first flashes were seen, and filmographers performed a simultaneous catch-and-release identification process to acquire ground-truth labels from visual inspection of the individuals present in the swarm. The movies were subsequently processed as described in a previous work [19, 20] to extract the 3D locations of flash occurrences. From these locations, we apply a simple distance-based linkage method to concatenate flashes into streaks and streaks into trajectories. We consider

flashes at consecutive timesteps within a small radius to be part of the same streak; streaks occurring within both 1s and 1m of each other are assumed to come from the same individual and placed in a set of transitively connected streaks called a trajectory. To eliminate noise effects from the trajectory extraction, we threshold the trajectories to eliminate those that only contain one flash. The dataset includes ten total species before the application of the thresholding process. Following the thresholding, we also remove any species from the dataset that have fewer than one hundred total trajectories, leaving us with seven species total. Finally, from the trajectories, we extract a binary time sequence by considering the time coordinates of flashes. This process enables the capture of individual behavior from simple footage of firefly swarms of any species, provided individuals of that species flash frequently enough to meet the threshold standards of the trajectory generation. Our data acquisition procedure highlights the presence of intraspecies behavioral variability, and characterizes this variability by representing flash patterns as distributions (Fig. 3A-C).

The result of this process is 73,141 flash trajectories from the seven species. More than half of these are *P. carolinus* sequences - the majority class. About 1 percent of these are *P. forresti* and *P. bethaniensis* sequences - the minority classes. The rest of the classes are fairly evenly distributed. Once single flash sequences are removed, the dataset comprises binary sequences of between 14 and 1365 frames (0.467s to 45.5s) in length, each labeled with the corresponding firefly class.

4.2 Modeling

4.2.1 Neural network architecture

RNNs are a class of neural networks suitable for sequence learning tasks. They are characterized by feedback connectivity and the consequent ability to encode long-range temporal dependencies, such as those intrinsic to firefly flash patterns. The defining computational step of an RNN is the hidden state update, which is a function of the input at the current timestep, $x^{(t)}$, and the hidden state at the previous timestep, $h^{(t-1)}$:

$$h^{(t)} = f(W_{hh}h^{(t-1)} + W_{xh}x^{(t)} + b),$$

where f is a non-linear activation function, such as hyperbolic tangent, W_{hh} and W_{xh} are weight matrices that map hidden-to-hidden (i.e., the feedback connectivity) and input-to-hidden, respectively, and b is a bias term.

Importantly, this formulation enables a recurrent neural network to ingest variable length input sequences, as the update rule can be applied recurrently, which is suitable for firefly flash sequences that are of variable temporal duration. However, if the input duration is sufficiently large–as is the case with some of flash pattern sequences–vanishing gradients will arise when computing weight updates via backpropagation through time (BPTT) [25] due to the non-linearity in f, ultimately prohibiting learning.

To address this issue, we leverage an extension to RNNs–gated recurrent units (GRUs)–that introduces gating mechanisms that regulate which information is stored to and retrieved from the hidden state [26]. These gating mechanisms enable the model to more effectively regulate the temporal context encoded in the hidden state and enables the encoding of longer-range temporal dependencies. Additionally, GRU RNNs are computationally more efficient than other kinds of RNNs like long short-term memory networks (LSTMs), and use fewer parameters, which was a consideration due to our plans for eventual downstream application of the model in real-time population monitoring. Consequently, we implement the model in PyTorch [27] as a 2-layer GRU with 128-dimension hidden layers, no dropout, and LeakyReLU activation layers with a negative slope of 0.1.

4.2.2 Data preprocessing

To evaluate our model's predictive ability on unseen data, we perform 60-fold stratified cross validation to ensure that each sequence in the dataset is used at least once in training and at least once in testing, but never simultaneously. Each fold divides the data into ninety percent training, and ten percent testing, preserving the same class ratios as the original dataset. Due to the severe class imbalance (e.g., some species only comprise 1% of the dataset, whereas others comprise close to 50% of the dataset), we perform random undersampling on the training set of each fold to equalize the class count in the training and validation sets for each fold. This takes the form of a secondary k-fold cross validation procedure to sample from each class until classes are equalized. All the remaining data are used for testing. The reported results are thus the ensemble precision, recall, and accuracy of each model on its respective test set of approximately 30,000 sequences, averaged over the 60 model folds. The ground truth targets are species names; we performed label encoding to transform the targets into machine-readable integers representing each class.

4.2.3 Training and evaluation

We trained the model with the Adam optimizer [28] and evaluated performance via cross-entropy loss. During training, we set an early stopping callback that monitored the validation loss with a patience of 50 epochs to prevent overfitting on the training set. Additionally, to alleviate exploding gradients, we applied a gradient clipping of 0.1 to the gradient norms following the procedure recommended in [29]. We conducted a hyperparameter sweep over the batch size and learning rate, testing all combinations of batch size $\in \{8, 16, 32\}$ and learning rate $\in \{10^{-3}, 10^{-4}, 10^{-5}\}$. We selected the combination that had the highest validation set accuracy on a four-species subset of the data, which resulted in the choice of a batch size of 8 and a learning rate of 10^{-5}. No data augmentation was applied during training.

We evaluate the performance of the RNN, along with the signal processing methods described in the following section, on the test data by examining the receiver-operating characteristic (ROC) curves for each species (Fig. 5A-E). Per-species precision and recall are tabulated in Fig. 5F.

4.3 Signal processing methods

For the purposes of comparison with the RNN, we implemented four alternative classifiers which use standard signal processing algorithms to compare our dataset against ground truth references for each species. We implement these classifiers using two types of ground truth references: "literature references", which use flash patterns as previously published in the literature, and "population references", which are generated by aggregating sequences in our own dataset.

4.3.1 Literature references

"Characteristic" flash patterns for six out of the seven species analyzed in this paper, excluding *B. wickershamorum*, have been previously recorded and published in the literature. These recorded flash patterns hence served as the primary reference for researchers in identifying signals observed in the field. These reference flash patterns are typically reported pictorially; thus, we convert images to binary-valued time series by computing the relative sizes of flashes and gaps, in pixels. We determine the pixel-to-second conversion to then convert the sequence to a 30 frames per secod time series, matching the sampling frequency of our data.

These six reference time series then form the ground-truth comparisons against which our dataset is compared, using the four signal processing methods described below in section 4.3.3. We

omit *B. wickershamorum* as there is currently no published reference pattern.

4.3.2 Population references

We also generate "population references" by aggregating sequences in our own dataset. For each species, we first perform an 80:20 train:test split, similar to the preprocessing procedure performed for the RNN (see above section 4.2.2). The population references are obtained by averaging the sequences in each training set. The remaining test data is then classified using the signal processing algorithms described below in section 4.3.3. As with the RNN, we perform $N = 100$ iterations and take the ensemble average of the performance across all iterations.

4.3.3 Signal processing algorithms

Jaccard index. The Jaccard index compares the similarity between two sets by taking the ratio of the size of the intersection of the sets with the size of the union [30] and has found broad application, for example in genomics [31]. For two binary-valued sequences $(a_m)_{m=1}^M$ and $(b_n)_{n=1}^N$ of lengths M and N, respectively, with $a_m, b_n \in \{0, 1\}$ for all m and n, we define the size of the intersection as $\sum_{i=1}^{\min(M,N)} a_i b_i$, the number of 'on' (flashing) bits that occur simultaneously in both sequences. We define the union as $\sum_{m=1}^M a_m + \sum_{n=1}^N b_n$, the number of on bits for both sequences combined. The Jaccard index can also be evaluated in the same manner for two sequences that are binary-valued. Generally speaking, the intersection can also be thought of as the dot product between the two sequences. To classify a sequence using the Jaccard index, the Jaccard index between a sequence and each species reference is computed, and the softmax of the vector of Jaccard index values is computed to determine a probability of the sequence being from each species. The predicted species is then the argument maximum (arg max) of the softmax vector.

Dot product. The dot product between two sequences is given by the sum of the product of the sequences, i.e., $\sum_{i=1}^{\min(M,N)} a_i b_i$ for two sequences $(a_m)_{m=1}^M$ and $(b_n)_{n=1}^N$ of lengths M and N, respectively. Sequences are then classified by taking the arg max of the softmax of the dot product with each reference.

Dynamic time warping. Dynamic time warping (DTW) is an algorithm that computes the distance between two timeseries by locally stretching or compressing the sequences to optimize the match. DTW is useful for comparing timeseries that are qualitatively similar but vary locally in speed and has found significant application in speech recognition [32–35]. We implement DTW in MATLAB [36] to compute the distance between sequences and species references. Similarly to the other metrics, the predicted species is taken to be the arg max of the softmax of the distances, which is a vector of probabilities that maps to the probability of each label.

SVM. Each flash sequence can be parametrized by 3 values (Fig. 1A): the number of flashes in the sequence, the average duration of each flash, and the average length of spaces between flashes (inter-flash gap). We perform support vector machine (SVM) classification in this 3-dimensional space, using a radial basis kernel function.

Figure 4: Reference sequences for the firefly species examined in this paper, as previously published in the literature, with the exception of *B. wickershamorum*. **(A)** Reference pattern for *P. frontalis* from Barber, 1951 [37]. **(B)** Reference patterns for *P. obscurellus* and *P. bethaniensis* from Faust, 2017 [7]. **(C)** Reference patterns for *P. carolinus* and *P. knulli* from Stanger-Hall and Lloyd, 2015 [8]. **(D)** Reference pattern for *P. forresti* from Fallon et al., 2022 [6]. **(E)** Illustration of extracting *P. frontalis* sequence from literature pattern (top) and converting to time series (bottom).

4.4   Characterization

The data acquisition procedure is not without noise, so we perform filtering to produce accurate quantitative characterization of flash phrases that falls in alignment with previous literature observations. We leveraged the ability of the RNN to distinguish between sequences by choosing the sequences which the RNN scored highest as the top one hundred most confident classifications for each species. This subset acts as the dataset on which characterization exercises are performed for Fig. 1B and Fig. 3. The procedure is as follows:

1. Initialize the empty list c

2. for each i sequence in the test set D:

    (a) Run a forward model step

    (b) Let $p =$ the maximum probability in the resulting vector of softmax predictions

    (c) If the index of p corresponds with the correct label, add the pair (p, index) to the list c

3. Sort c by probability p and choose the top 100

4. Index into the dataset D using the associated indices of the top 100 probabilities to produce the subset

Characterizing in this way leverages the variability in the entire dataset by training the predictive classifier, then asks the predictive classifier only for what it is most confident about in order to filter out sequences that may be missing flashes or exhibiting patterns that are far from the statistical norms of the species.

5   Acknowledgements

The authors would like to thank many collaborators on this project. Avalon Owens, Paul Shaw, Richard Joyce, Cheryl Mollahan, Peggy Butler, Jason Davis, and Julie Hayes all were excellent

Figure 5: Per-species ensemble of classification results. **A-E**: Receiver operating characteristic (ROC) curves representing the true positive rate (TPR) as a function of the false positive rate (FPR) across all model thresholds of classification, labeled by method. All non-RNN classification methods are conducted using population references. **F**: Table of per-species precision and recall across all surveyed methods (N=100). Bold statistics in the table represent the highest performer for each metric and species. Precision values for *P. bethaniensis* and *P. forresti* are low because these two species represent the classes with the fewest number of samples, and so there is a very small amount of true positive values. However, these still greatly exceed what would be expected by chance (0.001 and 0.006, respectively). The high recall for these classes indicates that the true positives are correctly captured.

volunteer data gatherers following our methodology. Candace Fallon and Sara Lewis were invaluable allies in consultation about firefly behavioral ecology and conservation threats. We'd also like to thank Airy Gonzalez Peralta, Yuti Gao, and Michael Hoffert for their insight and feedback as the project developed, and Brett Melbourne for his patience and helpful guidance through endless papers related to machine learning in ecology as we moved towards the right classification approach. Finally, the authors would like to thank Golnar Gharooni Fard, Nolan Bonnie, and Sam Pugh for thoughtful discussion and insights about the paper. O.P. acknowledges support from the National Geographic Society grant NGS-84850T-21, the Research Cooperation for Science Advancement grant 28219, and the BioFrontiers Institute at the University of Colorado Boulder.

5.1 Author contributions

O.P. conceived the study and provided funds to conduct research. O.P., C.N., O.M., M.L.I., and D.M.N. designed the research. R.S. and O.P. provided flash sequence data. M.L.I. and D.M.N. prototyped preliminary models. O.M. and C.N. analyzed the data, built the final models, and performed statistical analyses. M.C. and R.L. provided signal processing and machine learning algorithmic insights. O.M., C.N., R.S., M.C., R.L., M.L.I., and O.P. wrote the paper. O.P. and R.L. supervised the project.

5.2 Competing interests

The authors declare that they have no competing interests.

5.3 Data and materials availability

The associated code and data files for this project can be found here: https://github.com/peleg-lab/FireflyClassification. Following instructions in the associated README.md should allow easy regeneration of the figures and replication of the workflows described in this paper. All data needed to evaluate the conclusions in the paper are present in the github repository, the Supplementary Information, and/or the associated dataset paper [20].

References

[1] Sara M. Lewis and Christopher K. Cratsley. Flash signal evolution, mate choice, and predation in fireflies. *Annual Review of Entomology*, 53(1):293–321, 2008. PMID: 17877452.

[2] James E Lloyd. Signals and mating behavior in several fireflies (coleoptera: Lampyridae). *The Coleopterists' Bulletin*, pages 84–90, 1966.

[3] James E Lloyd. Bioluminescence and communication in insects. *Annual Review of Entomology*, 28(1):131–160, 1983.

[4] William D. McElroy, Howard H. Seliger, and Emil H. White. Mechanism of bioluminescence, chemi-luminescence and enzyme function in the oxidation of firefly luciferin. *Photochemistry and Photobiology*, 10(3):153–170, 1969.

[5] King-Siang Goh, Chia-Ming Lee, and Tzi-Yuan Wang. Species-specific flash patterns track the nocturnal behavior of sympatric taiwanese fireflies. *Biology*, 11(1):58, 2022.

[6] Candace Fallon, Anna Walker, Sara Lewis, and Sarina Jepsen. State of the fireflies of the united states and canada: Distributions, threats, and conservation recommendations. 01 2022.

[7] Lynn Frierson Faust. *Fireflies, Glow-worms, and Lightning Bugs: Identification and Natural History of the Fireflies of the Eastern and Central United States and Canada*. University of Georgia Press, 2017.

[8] Kathrin F. Stanger-Hall and James E. Lloyd. Flash signal evolution in Photinus fireflies: Character displacement and signal exploitation in a visual communication system. *Evolution*, 69(3):666–682, March 2015.

[9] Travis Longcore and Catherine Rich. Ecological light pollution. *Frontiers in Ecology and the Environment*, 2(4):191–198, 2004.

[10] Avalon C. S. Owens and Sara M. Lewis. The impact of artificial light at night on nocturnal insects: A review and synthesis. *Ecology and Evolution*, 8:11337, 2018.

[11] Stephanie Vaz, Stella Manes, Danielle Gama-Maia, Luiz Silveira, Gustavo Mattos, Paulo C. Paiva, Marcos Figueiredo, and Maria Lucia Lorini. Light pollution is the fastest growing potential threat to firefly conservation in the atlantic forest hotspot. *Insect Conservation and Diversity*, 14(2):211–224, 2021.

[12] Candace E. Fallon, Anna C. Walker, Sara Lewis, Joseph Cicero, Lynn Faust, Christopher M. Heckscher, Cisteil X. Pérez-Hernández, Ben Pfeiffer, and Sarina Jepsen. Evaluating firefly extinction risk: Initial red list assessments for North America. *PLOS ONE*, 16(11):e0259379, November 2021.

[13] Avalon Owens, Mira Van den Broeck, Raphaël De Cock, and Sara M Lewis. Behavioral responses of bioluminescent fireflies to artificial light at night. *Frontiers in Ecology and Evolution*, page 948, 2022.

[14] Sara M Lewis, Choong Hay Wong, Avalon C.S. Owens, Candace Fallon, Sarina Jepsen, Anchana Thancharoen, Chiahsiung Wu, Raphael De Cock, Martin Novák, Tania López-Palafox, Veronica Khoo, and J Michael Reed. A global perspective on firefly extinction threats. *BioScience*, 70(2):157–167, 2020.

[15] Francisco Sánchez-Bayo and Kris A.G. Wyckhuys. Worldwide decline of the entomofauna: A review of its drivers. *Biological Conservation*, 232:8–27, 2019.

[16] Avalon C.S. Owens, Précillia Cochard, Joanna Durrant, Bridgette Farnworth, Elizabeth K. Perkin, and Brett Seymoure. Light pollution is a driver of insect declines. *Biological Conservation*, 241:108259, 2020.

[17] Raphaël Sarfati, Julie Hayes, Élie Sarfati, and Orit Peleg. Spatiotemporal reconstruction of emergent flash synchronization in firefly swarms via stereoscopic 360-degree cameras. *Journal of the Royal Society Interface*, 17:20200179, 2020.

[18] Xerces Society. Latest Update to the IUCN Red List Includes First Global Assessments for Fireflies, with a Spotlight on North America, 2021. `https://xerces.org/blog/iucn-red-list-assess-fireflies`.

[19] Raphaël Sarfati and Orit Peleg. Calibration-free 3d reconstruction of firefly trajectories from 360-degree cameras. *bioRxiv*, 2021. doi:10.1101/2021.04.07.438867.

[20] Raphaël Sarfati, Owen Martin, Julie C. Hayes, Avalon C.S. Owens, Paul Shaw, Cheryl Mollohan, Ronald L. Day Jr., Peter Mauney, Richard V. Joyce, Jason Davis, Peggy Butler, Radim Schreiber, Brian Auman, and Orit Peleg. Crowdsourced dataset of firefly trajectories obtained by automated calibration of pairs of 360-degree cameras. *In preparation*, 2023.

[21] David L. Woods, John M. Wyma, E. William Yund, Timothy J. Herron, and Bruce Reed. Factors influencing the latency of simple reaction time. *Frontiers in Human Neuroscience*, 9, 2015.

[22] Keunwoo Choi, George Fazekas, Mark Sandler, and Kyunghyun Cho. Convolutional recurrent neural networks for music classification, 2016.

[23] California Academy of Sciences. iNaturalist, 2008. `https://www.inaturalist.org`.

[24] Chantal Nguyen, Isabella Huang, and Orit Peleg. Firefly-inspired vocabulary generator for communication in multi-agent systems. *ALIFE 2022: The 2022 Conference on Artificial Life*, 2022.

[25] Michael C Mozer. A focused backpropagation algorithm for temporal pattern recognition. *Complex Systems*, 3(4), 1989.

[26] Junyoung Chung, Caglar Gulcehre, KyungHyun Cho, and Yoshua Bengio. Empirical Evaluation of Gated Recurrent Neural Networks on Sequence Modeling. *arXiv:1412.3555 [cs]*, December 2014. arXiv: 1412.3555.

[27] PyTorch. `https://pytorch.org/`.

[28] Diederik P. Kingma and Jimmy Ba. Adam: A method for stochastic optimization. 2014. doi:10.48550/arXiv.1412.6980.

[29] Jingzhao Zhang, Tianxing He, Suvrit Sra, and Ali Jadbabaie. Why gradient clipping accelerates training: A theoretical justification for adaptivity, 2019. doi:10.48550/arXiv.1905.11881.

[30] P. Jaccard. Étude comparative de la distribution florale dans une portion des alpes et des jura. *Bulletin de la Société vaudoise des sciences naturelles*, 37:547–579, 1901.

[31] Dmitry Prokopenko, Julian Hecker, Edwin K. Silverman, Marcello Pagano, Markus M. Nöthen, Christian Dina, Christoph Lange, and Heide Loehlein Fier. Utilizing the Jaccard index to reveal population stratification in sequencing data: a simulation study and an application to the 1000 Genomes Project. *Bioinformatics*, 32(9):1366–1372, 12 2015.

[32] Taras. K. Vintsyuk. Speech discrimination by dynamic programming. *Kibernetika*, 4(1), 1968.

[33] Fumitada Itakura. Minimum prediction residual principle applied to speech recognition. *IEEE Transactions on Acoustics, Speech, and Signal Processing*, ASSP-23(1), 1975.

[34] Hiroaki Sakoe and Satoshi Chiba. Dynamic programming algorithm optimization for spoken word recognition. *IEEE Transactions on Acoustics, Speech, and Signal Processing*, ASSP-26(1), 1978.

[35] Donald J. Berndt and James Clifford. Using dynamic time warping to find patterns in time series. *Proceedings of the 3rd International Conference on Knowledge Discovery and Data Mining, AAAIWS'94*, pages 359–370, 1994.

[36] Quan Wang. Dynamic time warping, 2014. `https://www.mathworks.com/matlabcentral/fileexchange/43156-dynamic-time-warping-dtw`.

[37] H. S. Barber. North American fireflies of the genus Photuris. *Smithsonian Miscellaneous Collections*, 117(1), 1951.

Supplemental Information

5.4 Batch effects

The dataset was assembled by several scientists operating at different locations across different years, so it is important to consider the possibility that the network is isolating a batch-related feature. We control for this in several ways: first, the methodology for video collection is standardized following procedural guidelines outlined in [20]. These are easy to follow, clearly delineated, and often quality checked before the data are gathered. Second, the videos are post-processed to transform them into binary sequences, eliminating extraneous factors that may normally contribute to batch effects in neural network training. Third, the resulting sequences are all transformed to the same time scale to avoid preference given to a certain frame rate. We believe these steps are sufficient to remove any noticeable effect generated by minor differences in data collection moving forward, but testing this is important to make sure the model is not learning a characteristic of the sequences that is isolated to a certain data gatherering event. To test this, we performed 5-fold cross-validation on the recurrent neural network where all of the data for a particular data gatherer was held out of the training set. For each fold, we isolated all of the *P. frontalis* and *P. carolinus* sequences in the dataset coming from R.S. [20]. We did this because *P. carolinus* and *P. frontalis* are two of the species for which there are multiple data-gathering events and camera operators present in the dataset, and R.S. took part in some of the data collection for each of these species. Holding their data out from the training dataset means the model will not be able to see any of those sequences, so if there is a batch effect, it should not be able to recognize them. Then for each of the 5 folds, we selected a different fold of the remaining *P. carolinus* and *P. frontalis* data to be part of the training set. The other species were treated normally, and the results are in Fig. S1. It is clear that there is no batch effect, as the model performs just as well during the 5-fold cross validation as it does without any data holdout. This is likely thanks to the distinctive flash length of *P. frontalis*; additionally, despite the differences *between* distributions of the inter-flash gap for *P. frontalis*, the overall distribution is still uniquely situated between 0.5s-0.8s, which is unlike any of the other species in the dataset. This analysis supports the ability for our model to scale with different volunteers acting in different locations serving as the data-gatherers, and eventually to automated camera setups, without concern.

Figure S1: Quantification of batch effects. **A-C:** Probability distributions of flash parameters from *P. frontalis* sequences collected by two different data gatherers, labeled RS (top) and PS (bottom). The distributions for number of flashes **(A)** and flash length **(C)** are similar, while the inter-flash gap **(B)** distributions are notably different due to different temperatures during data collection for the two different gatherers. **D:** Confusion matrix for 5-fold cross validation of RNN where all RS-collected *P. frontalis* data held out from training. The RNN achieves similar performance to the model described in the main text, which was trained on both RS- and PS- collected *P. frontalis* data. **E:** Table of per-species precision and recall during batch effect cross-validation.

5.5 Literature ROC results

The literature methods perform poorly compared to population reference methods, with the SVM achieving the best result (precision = 0.32, recall = 0.28). However, we observe that the SVM classifier predicts most samples as either *P. bethaniensis* or *P. forresti*, The receiver-operating characteristic (ROC) curves for the literature-based methods are shown in Fig. S2, along with the per-species precision and recall. All four methods perform poorly on *P. carolinus*. SVM and DTW result in near-perfect recall on *P. knulli*, but poor precision; conversely, these methods give near-perfect precision on *P. frontalis*, but poor recall.

These results reveal that the existing literature-based reference patterns fall short in accurately representing our collected dataset. This may be due to the different conditions under which the literature-based references; moreover, by creating population references, we can better capture the behavioral variability found in the data.

E	SVM		Jaccard index		Dot product		DTW	
Species	Precision	Recall	Precision	Recall	Precision	Recall	Precision	Recall
P. bethaniensis	0.022	**0.86**	0.0054	0.078	0.019	0.21	**0.028**	0.85
P. carolinus	0.73	0.082	0.54	0.17	0.58	**0.51**	**0.93**	0.09
P. forresti	0.019	**0.41**	0.020	0.11	**0.026**	0.13	0.025	0.13
P. frontalis	0.36	0.035	0.42	**0.45**	0.29	0.029	**0.50**	0.09
P. knulli	**0.50**	0.39	0.23	0.50	0.15	0.36	0.15	**0.45**
P. obscurellus	**0.41**	0.013	0.19	0.22	0.38	**0.24**	0.016	0.031

Figure S2: Classification results for literature-based references. A-D: Receiver operating characteristic (ROC) curves representing the true positive rate (TPR) as a function of the false positive rate (FPR) across all model thresholds of classification, labeled by method. E: Table of per-species precision and recall.