Destabilization of neuromuscular junctions and deregulation of activity-dependent signalling pathways

in Myotonic Dystrophy type I

Denis Falcetta[1-3], Sandrine Quirim[1], Ilaria Cocchiararo[1], Mélanie Cornut[1], Marine Théodore[2], Adeline

Stiefvater[2], Shuo Lin[3], Lionel Tintignac[2], Robert Ivanek[4,5], Jochen Kinter[2], Markus A. Rüegg[3], Michael

Sinnreich[2], Perrine Castets[1*]

1. Department of Cell Physiology and Metabolism, Faculty of Medicine, University of Geneva, 1 rue Michel

Servet, CH-1211 Geneva, Switzerland.

2. Neuromuscular Research Group, Departments of Neurology and Biomedicine, University and University

Hospital Basel, Klingelbergstrasse 50/70, CH-4056 Basel, Switzerland.

3. Biozentrum, University of Basel, Spitalstrasse 41, CH-4056 Basel, Switzerland.

4. Department of Biomedicine, University Hospital and University of Basel, Hebelstrasse 20, CH-4053 Basel,

Switzerland.

5. Swiss Institute of Bioinformatics, Hebelstrasse 20, CH-4053 Basel, Switzerland.

* Corresponding author: Perrine Castets, perrine.castets@unige.ch, +41 22 379 52 33

ABSTRACT

Myotonic Dystrophy type I (DM1) is the most common muscular dystrophy in adults. Previous reports have

highlighted that neuromuscular junctions (NMJs) deteriorate in skeletal muscle from DM1 patients and

mouse models thereof. However, the underlying pathomechanisms and their contribution to muscle

dysfunction remain unknown. We compared changes in NMJs and activity-dependent signalling pathways

in HSA^{LR} and $Mbnl1^{\Delta E3/\Delta E3}$ mice, two established mouse models for DM1. DM1 muscle showed major

deregulation of calcium/calmodulin-dependent protein kinases II (CaMKIIs), which are key activity sensors

regulating synaptic gene expression and acetylcholine receptor (AChR) recycling at the NMJ. Both mouse

models displayed increased fragmentation of the endplate, which preceded muscle degeneration.

Endplate fragmentation was not accompanied by changes in AChR turnover at the NMJ. However,

expression of synaptic genes was up-regulated in DM1 muscle, which may be linked to the abnormally

high activity of histone deacetylase 4 (HDAC4), a known target of CaMKII. Consistently, expression of

myosin heavy chains was deregulated as well, leading to a major switch to type IIA fibres in $Mbnl1^{\Delta E3/\Delta E3}$

muscle, and to a lesser extent in HSA^{LR} muscle. Interestingly, although HDAC4 was efficiently induced upon

nerve injury, synaptic gene up-regulation was abrogated in DM1 muscle, together with a reduced increase

in AChR turnover. This suggested that HDAC4-independent mechanisms lead to the defective response to

denervation in DM1 muscle. Our study shows that activity-dependent signalling pathways are disturbed in

DM1 muscle, which may contribute to NMJ destabilization and muscle dysfunction in DM1 patients.

INTRODUCTION

Myotonic Dystrophy type I (DM1) is a multisystemic disorder caused by expanded CTG triplet repeats in the 3'UTR of the *DMPK* (*Dystrophia Myotonica Protein Kinase*) gene, leading to muscle wasting, weakness and inability to relax (myotonia). Accumulation of toxic transcripts containing the expanded CUG repeats lead to the nuclear sequestration of splicing factors. The consecutive mis-splicing of specific genes is determinant in the pathogenesis of DM1-associated muscle alterations.[1, 2] For example, mis-splicing of the *CLCN1* gene encoding the chloride channel ClC-1 is implicated in myotonia development. Mis-splicing of several genes encoding proteins of Ca^{2+}-associated signalling pathways has also been shown to contribute to muscle dysfunction. Especially, previous reports unveiled that mis-splicing of *CAMK2* genes, encoding Ca^{2+}/calmodulin-dependent protein kinases II (CaMKIIs), is a hallmark of DM1.[3-5] However, the functional consequences of CAMKII deregulation in DM1 muscle and its contribution to DM1 pathogenesis were not explored.

CaMKIIs are important for the maintenance of neuromuscular junctions (NMJs), which are the synapses connecting motor neurons to muscle fibres. Especially, CaMKIIs enhance the recycling of acetylcholine receptors (AChRs) upon their internalization in the sub-synaptic region of muscle fibres (*i.e.* the endplate).[6] Moreover, CaMKIIs indirectly repress the expression of synaptic genes in non-synaptic regions of innervated muscle, by inhibiting the myogenic transcription factor myogenin and the histone deacetylase 4 (HDAC4).[7-9] Hence, deregulation of CaMKII signalling pathway may affect NMJ maintenance, by perturbing the expression and the dynamics of synaptic proteins.

Early studies on DM1 muscle biopsies pointed to NMJ-associated abnormalities, such as a reduced number of presynaptic vesicles, enlarged endplates or angular muscle fibres.[10-13] Lack of denervation markers (*e.g.,* non-junctional AChR clusters) and the absence of massive motor neuron loss rejected the hypothesis that spontaneous denervation is part of DM1 pathogenesis.[14, 15] NMJ alterations have also been reported in *DMSXL* mice and in mice deficient for Muscleblind-like 1/2 (Mbnl1/2), two mouse models for

DM1, as well as in C. *elegans* DM1 mutants.[16, 17] Moreover, nuclear foci, characteristic of DM1-associated

accumulation of toxic RNA, were detected at the endplate, which may alter the expression of synaptic

genes.[18] Mis-splicing of some synaptic genes has been reported in muscle cells from DM1 patients.[19]

Hence, NMJ deterioration in DM1 is likely to contribute to muscle dysfunction, but the underlying

pathomechanisms remain unknown.

Here, we analysed changes in activity-dependent signalling pathways in HSA^{LR} and $Mbnl1^{\Delta E3/\Delta E3}$ mice,

two established DM1 mouse models. Both mouse models displayed increased fragmentation of the

endplate, which preceded muscle alterations. HSA^{LR} and $Mbnl1^{\Delta E3/\Delta E3}$ mice also showed deregulation of

CaMKII / HDAC4 signalling in muscle, which was accompanied by an increased expression of synaptic genes

and a switch to slower fibre type. Interestingly, endplate remodelling upon nerve injury was hampered in

DM1 mice, despite efficient induction of HDAC4. Hence, perturbations in activity-dependent signalling

pathways may contribute to NMJ destabilization and muscle dysfunction in DM1 patients.

RESULTS

DM1-associated muscle alterations are similar in *Mbnl1*$^{\Delta E3/\Delta E3}$ and *HSA*^{LR} mice

To evaluate changes in NMJs and activity-dependent pathways in DM1, we selected the *Mbnl1*$^{\Delta E3/\Delta E3}$ and

HSA^{LR} mouse lines, which are two well-established DM1 mouse models. In *Mbnl1*$^{\Delta E3/\Delta E3}$ mice, deletion of

Mbnl1 exon 3 leads to ubiquitous depletion of the splicing factor MBNL1 (*i.e.,* including in skeletal muscle

and motor neurons).[21] In contrast, *HSA*^{LR} mice express the *HSA* transcript (*Human Skeletal Actin*) with long

(CTG) repeats only in skeletal muscle, which allowed to unveil cell-autonomous defects.[20] First, we

compared the muscle phenotype between the two mouse models. There was no dystrophic sign in muscles

from 3-months-old mutant mice (*Figure* 1A and 1B), as previously reported.[20, 21] Muscle degeneration

remained mild in 9- and 12-month-old *Mbnl1*$^{\Delta E3/\Delta E3}$ and *HSA*^{LR} mice, respectively (*Figure* 1A and 1B).

Alterations, such as internalized nuclei, vacuoles and degenerated fibres, were present in *Mbnl1*$^{\Delta E3/\Delta E3}$

muscle and in *gastrocnemius* muscle from *HSA*^{LR} mice. In contrast, alterations were rare in *tibialis anterior*

(TA) and *extensor digitorum longus* (EDL) muscles from *HSA*^{LR} mice (*Figure* 1A and 1B). Both *Mbnl1*$^{\Delta E3/\Delta E3}$

and *HSA*^{LR} muscles displayed a myotonic phenotype, as shown by the increased late relaxation time of EDL

muscle after *ex vivo* stimulation (*Figure* 1C and 1D). In contrast, specific tetanic muscle force (sP0) was not

affected in 3-month-old mutant mice (*Figure* 1E), consistent with previous reports showing late onset of

muscle weakness.[20, 21]

To evaluate DM1-associated mis-splicing, we next quantified the inclusion of exons 7a and 22 of the

Clcn1 and *Atp2a1* genes, encoding ClC-1 channel and SERCA1 (*Sarco/Endoplasmic Reticulum Ca*$^{2+}$*-ATPase*),

respectively. Inclusion of *Clcn1* exon 7a was strongly increased in *Mbnl1*$^{\Delta E3/\Delta E3}$ muscle and was

accompanied by major reduction in total transcript levels of *Clcn1* (*Figure* 1F). *Clcn1* mis-splicing and down-

regulation were comparable in *gastrocnemius* muscle from *HSA*^{LR} mice (*Figure* 1F). In contrast, TA muscle

from *HSA*^{LR} mice showed milder changes in *Clcn1* splicing and no reduction in total *Clcn1* transcript levels

(*Figure* 1F). Similarly, inclusion of *Atp2a1* exon 22 was abrogated in *Mbnl1*$^{\Delta E3/\Delta E3}$ muscle and in

gastrocnemius muscle from *HSA^{LR}* mice, while it was only reduced by half in *HSA^{LR}* TA muscle (*Figure* 1F).

These results confirm that both mouse models display mild muscle alterations, with similar DM1-

associated phenotype observed in *Mbnl1^{ΔE3/ΔE3}* muscle and *HSA^{LR}* *gastrocnemius* muscles, and milder

changes detected in TA/EDL muscles from *HSA^{LR}* mice.

DM1 muscle shows major deregulation of CaMKIIs

Previous reports pointed to mis-splicing of *CAMK2* genes in DM1 tissues.[3-5] However, it remains unclear

which CaMKII isoforms are affected in DM1 muscle, and what are the consequences of their deregulation.

CaMKIIs are encoded by the four genes *CAMK2A, 2B, 2G* and *2D*, and expressed as different splice variants.

CaMKIIβ/γ/δ isoforms are expressed in skeletal muscle. Protein levels of the muscle-specific variant of

CaMKIIβ (*i.e.,* CaMKIIβM) were strongly reduced in both *Mbnl1^{ΔE3/ΔE3}* (*Figure* 2A) and *HSA^{LR}* (*Figure* 2B)

muscles. Auto-phosphorylation of CaMKIIβM (Phospho-Ser286) was decreased as well, suggesting reduced

CaMKIIβM activity (*Figure* 2C). Of note, additional bands around the size of the ubiquitous isoforms of

CaMKIIβ and of CaMKIIγ/δ point to the expression of alternative CaMKII isoforms in mutant muscle (*Figure*

2A and 2B).

To characterize splicing events in *Camk2b, 2g* and *2d* genes in DM1 muscle, we took advantage of RNA-

seq results that we obtained in *HSA^{LR}* *gastrocnemius* muscle. We detected significant changes in *Camk2b*

transcript, with an increased exclusion of the exons 13 and 18 to 20 in *HSA^{LR}* muscle, compared to control

(*Figure* S1A and S1B). We confirmed the expression of *Camk2b* transcripts lacking exon 13 in *HSA^{LR}* muscle

by RT-PCR (*Figure* S1C and S1D). Moreover, the amplicon including exons 18 to 20, encoding the variable

inserts A, B and C of CaMKIIβM, was barely detected in *HSA^{LR}* muscle (*Figure* S1C and S1D). By quantitative

RT-PCR, total levels of *Camk2b* transcripts were increased in both *Mbnl1^{ΔE3/ΔE3}* and *HSA^{LR}* muscles (*Figure*

2D). Using primers spanning *Camk2b* exons 12-14, we confirmed that expression of *Camk2b* transcripts

without exon 13 was more than four times higher in mutant muscles than in controls (*Figure* 2E). Inversely,

levels of *Camk2b* transcripts with exons 18 to 20 were reduced by half in *Mbnl1$^{ΔE3/ΔE3}$* TA muscle and in *gastrocnemius* muscle from *HSA^{LR}* mice (*Figure* 2F) compared to controls. Of note, TA and EDL muscles from *HSA^{LR}* mice showed similar extent of *Camk2b* mis-splicing compared to *gastrocnemius* muscle (*Figure* S1E and S1F). Strikingly, the mis-splicing of exon 13 and exons 18-20 is also detected in RNA-seq data of TA muscle from DM1 patients (*Figure* S2).[28]

Transcripts encoding CaMKIIγ were also mis-spliced in *HSA^{LR}* muscle. These included increased inclusion of exons 13, 15 and 19 in *Camk2g* transcripts in *HSA^{LR}* muscle, as seen in RNA-seq reads (*Figure* S3A and S3B) and by RT-PCR (*Figure* S3C). Increased inclusion of *CAMK2G* exon 19 was also observed in TA biopsies from DM1 patients (*Figure* S3D).[28] In contrast, there was no major splicing change detected for *CAMK2D* in DM1 muscle (*Figure* S4).[28] Together, these results show that some *Camk2* transcripts are mis-spliced in DM1 muscle and that the expression pattern of CaMKIIs is altered, with a loss of the muscle-specific variant CaMKIIβM.

Endplate fragmentation is not caused by abnormal AChR dynamics in *Mbnl1$^{ΔE3/ΔE3}$* and *HSA^{LR}* mice

As CaMKIIs are key sensors of neural activity involved in NMJ maintenance, we next analysed NMJ structures in EDL, TA and *gastrocnemius* muscles from 3-month-old and 9- or 12-month-old *Mbnl1$^{ΔE3/ΔE3}$* and *HSA^{LR}* mice. Pre- and post-synaptic compartments were stained with antibodies against Neurofilament/Synaptophysin and with α-bungarotoxin (Btx), which binds specifically to AChRs, respectively. The overall organization of the NMJs was preserved in mutant mice (*Figure* 3A and 3B). In particular, denervated endplate and abnormal axonal termination were not observed in mutant muscle. However, fragmentation of the endplate was increased in muscles from both 3- and 9-month-old *Mbnl1$^{ΔE3/ΔE3}$* mice, compared to age-matched controls (*Figure* 3C). A similar increase in endplate fragmentation was observed in both *gastrocnemius* and TA/EDL muscles from *HSA^{LR}* mice (*Figure* 3D). As the *HSA^{LR}* transgene is specifically expressed in skeletal muscle, these results indicate that post-synaptic

perturbations contribute to endplate fragmentation in *HSA^LR* mice. Moreover, these NMJ alterations were

detected as soon as 3 months of age, *i.e.* before changes in muscle histology, suggesting that they are

primary deficit in DM1 mice and not a consequence of muscle degeneration/regeneration.

As CaMKIIs have been shown to regulate AChR recycling at the endplate,[6] we next evaluated AChR

turnover, by labelling "old" and newly formed receptors by two sequential injections of differently labelled

Btx (*Figure* 4A and 4B).[26, 29] AChR turnover was low in all muscles, as described before for innervated

conditions, with no change detected in *Mbnl1^ΔE3/ΔE3* muscles as compared to controls (*Figure* 4C). In

contrast, there was a trend towards increased AChR turnover in TA and EDL muscles from *HSA^LR* mice

(*Figure* 4D). We further determined the extent of AChR recycling in *Mbnl1^ΔE3/ΔE3* EDL muscle, by using Btx-

biotin and two sequential injections of fluorescent streptavidin (*Figure* 4B).[27] After 1 and 3 days, the

proportion of recycled AChRs was also unaltered in muscle from *Mbnl1^ΔE3/ΔE3* mice compared to controls

(*Figure* 4E and 4F). Together, these results indicate that CaMKII deregulation, and more specifically the

loss of CaMKIIβM, do not alter AChR dynamics in innervated DM1 muscle.

HDAC4 accumulates in DM1 muscle

Another function of CaMKIIs in skeletal muscle is to repress synaptic gene expression, by inhibiting the

nuclear import, and thereby the activity of HDACs, such as HDAC4.[30] We therefore assessed the expression

pattern of HDAC4 in DM1 muscle. Transcript levels of *Hdac4* were unchanged in DM1 mice, as compared

to controls (*Figure* 5A). In contrast, protein levels of HDAC4 were strongly increased in both *Mbnl1^ΔE3/ΔE3*

and *HSA^LR* muscles, as compared to controls (*Figure* 5B and 5C). While CaMKII-dependent phosphorylation

of HDAC4 (Ser632) remained unchanged in *Mbnl1^ΔE3/ΔE3* muscle (*Figure* 5B), it tended to decrease in *HSA^LR*

muscle, as compared to controls (*Figure* 5C). Consistent with a reduced inhibition of HDAC4 by CaMKIIs,

HDAC4 levels were increased in the nuclear fraction of *Mbnl1^ΔE3/ΔE3* and *HSA^LR gastrocnemius* muscles

(*Figure* 5D). Of note, HDAC4 accumulation remained barely detectable in mutant muscle by

immunostaining (*Figure* 5E) and it was not sufficient to increase total levels of acetylated histones, as

reported upon denervation (*Figure* S5A and S5B). Consistent with higher HDAC4 activity, transcript levels

of *Dach2*, a gene directly repressed by HDAC4 (*Figure* S5C), tended to decrease in DM1 muscles (*Figure*

5F). In contrast, expression of *Mitr* was not repressed in mutant muscles, and transcript levels of *Myog,*

encoding myogenin, remained unchanged in *Mbnl1*$^{\Delta E3/\Delta E3}$ and *HSA*^{LR} muscles, as compared to controls

(*Figures* 5F, S5C and S5D). Hence, CaMKII deregulation may trigger accumulation of HDAC4 and a moderate

increase in its activity in DM1 muscle.

Expression of synaptic genes and fibre type composition are altered in DM1 muscle

We next assessed the consequences of CaMKII/HDAC4 deregulation in DM1 muscle on known activity-

dependent signalling pathways (*Figure* S5C). We first measured the transcriptional expression of *Musk*,

Chrna1, *Chrne* and *Chrng* genes, encoding MUSK and the α, ε and γ subunits of AChR, respectively. During

muscle development, expression of *Musk* and *Chrna1* becomes restricted to sub-synaptic myonuclei upon

muscle innervation.[31] Simultaneously, *Chrng* transcripts are downregulated, while *Chrne* starts to be

expressed in sub-synaptic nuclei (AChRε subunits replace AChRγ subunits). In non-synaptic regions of

muscle fibres, synaptic gene repression is mediated by CaMKII, and dependent on HDAC4/5 inhibition

(*Figure* S5C).[8, 32] This adult gene expression pattern depends on neural activity, as denervation reverts it

back to a developmental pattern. As shown in Figure 6A, transcript levels of *Musk, Chrna1, Chrne* and

Chrng, were increased in *Mbnl1*$^{\Delta E3/\Delta E3}$ muscle and in *gastrocnemius* from *HSA*^{LR} mice. Of note, their levels

were less or not changed in EDL and TA muscles from *HSA*^{LR} mice, as compared to controls (*Figure* S6A and

S6B).

HDAC4 is also known to regulate the transcription of *Myh* genes, encoding myosin heavy chains (MHC),

and thereby to promote a switch to type IIA fibres in TA muscle upon denervation.[8] Expression of *Myh2*,

encoding MHCIIA, was strongly increased in *Mbnl1*$^{\Delta E3/\Delta E3}$ innervated muscle, as compared to control

(*Figure* 6B). In contrast, expression of *Myh4*, encoding MHCIIB, was reduced in DM1 muscle (*Figure* 6B).

Consistently, innervated TA muscle from *Mbnl1*$^{\Delta E3/\Delta E3}$ mice displayed major accumulation of type IIA fibres

and a loss of type IIB fibres compared to controls (*Figure* 6C). Similarly, we have previously unveiled a mild

switch towards type IIA fibres in TA muscle from *HSA*^{LR} mice.[23] There was also a tendency towards

increased transcript levels of *Myh2* and reduced expression of *Myh4* in *HSA*^{LR} TA muscle, as compared to

controls (*Figure* 6D). Thus, CaMKII-dependent deregulation of HDAC4 may perturb activity-dependent

signalling pathways, with increased synaptic gene expression and a switch towards slower fibres in DM1

muscle.

Mbnl1$^{\Delta E3/\Delta E3}$ muscle responds to denervation

To get further insights on the deregulation of activity-dependent signalling pathways in DM1 muscles, we

assessed their response to neural inactivity. To this end, we challenged 3-month-old *Mbnl1*$^{\Delta E3/\Delta E3}$ and *HSA*^{LR}

mice by cutting the sciatic nerve to obtain a complete denervation of hind limb muscles. Unexpectedly,

we observed that *HSA*^{LR} mice lose the expression of the *HSA* transgene after 3 days of denervation (*Figure*

S7A). Consequently, accumulation of ribonuclear foci and mis-splicing of *Clcn1* and *Camk2b* were reduced

in denervated *HSA*^{LR} muscle, as compared to innervated muscle (*Figure* S7B-D). Because of this, we limited

the analysis to the *Mbnl1*$^{\Delta E3/\Delta E3}$ mouse line.

After 3 days of denervation, we observed an efficient increase in transcript and protein levels of HDAC4

in denervated muscle from both mutant and control mice (*Figure* 7A). Consistently, there was a major

accumulation of HDAC4 in nuclei of both DM1 and control denervated muscles (*Figure* 7B). Of note, the

loss of CaMKIIβM persisted upon denervation in *Mbnl1*$^{\Delta E3/\Delta E3}$ muscle (*Figure* 7A). Expression of the direct

targets of HDAC4, *Dach2* and *Mitr*, was similarly repressed in *Mbnl1*$^{\Delta E3/\Delta E3}$ and control denervated muscles

(*Figure* 7C). Moreover, expression of the atrogenes *Fbxo32* and *Trim63*, which are indirectly induced by

HDAC4 at this stage,[33] also reached similar transcript levels in *Mbnl1*$^{\Delta E3/\Delta E3}$ and control muscles (*Figure*

S8A). We and others have reported that the anabolic pathway Akt/mTORC1 (*mammalian Target Of*

Rapamycin Complex 1) is deregulated in DM1 muscle,[23] and that it is activated upon denervation.[26] Thus,

we assessed whether Akt/mTORC1 activity is perturbed in DM1 denervated muscle. At 3 days of

denervation, there was no change in the levels of the phosphorylated active form of Akt (Akt^{P473}) in both

Mbnl1$^{\Delta E3/\Delta E3}$ and control muscles, while levels of the active form of S6 (S6^{P235}) increased similarly in both

denervated muscles (*Figure* S8B). In parallel, levels of the autophagic marker LC3II remained largely

unchanged in *Mbnl1$^{\Delta E3/\Delta E3}$* and control muscles, suggesting that autophagy is not strongly affected (*Figure*

S8B). These results suggest that both the HDAC4 and mTORC1 signalling pathways are efficiently induced

upon denervation in DM1 muscle.

Interestingly, after prolonged denervation, the loss of muscle mass was significantly less in *Mbnl1$^{\Delta E3/\Delta E3}$*

mice, as compared to control mice (*Figure* 7D). Of note, nerve injury did not aggravate muscle

degeneration in *Mbnl1$^{\Delta E3/\Delta E3}$* mice (*Figure* S8C). To get insights on this resistance to atrophy, we evaluated

muscle fibre type and size after 3 weeks of denervation. Control TA muscle shifted to type IIA fibres upon

denervation, reaching similar proportion of type IIA fibres as observed in *Mbnl1$^{\Delta E3/\Delta E3}$* innervated muscle.

Fibre type proportion remained unchanged in *Mbnl1$^{\Delta E3/\Delta E3}$* denervated muscle (*Figure* 7E). The minimum

ferret of type IIA fibres was increased in *Mbnl1$^{\Delta E3/\Delta E3}$* innervated muscle as compared to control innervated

muscle, but it significantly decreased upon denervation in mutant mice (*Figure* 7F and *Figure* S8D). In

parallel, the size of type IIX and IIB fibres was similar in *Mbnl1$^{\Delta E3/\Delta E3}$* and control innervated muscles, and

it strongly decreased upon denervation in both mutant and control mice (*Figure* 7F and *Figure* S8D). Hence,

the predominance of type IIA fibres in mutant mice and their relative resistance to denervation-induced

atrophy as compared to type IIX/B fibres (*Figure* 7F), may explain why DM1 mice showed limited loss of

muscle mass upon nerve injury.

Endplate remodelling is impaired in *Mbnl1$^{\Delta E3/\Delta E3}$* muscle upon denervation

Synaptic remodelling upon denervation includes a strong increase in AChR turnover in the sub-synaptic

region, together with major up-regulation of synaptic genes throughout the fibre.[34] Release of synaptic

gene repression in non-synaptic muscle regions is thought to be dependent on the induction of HDAC4.[8,]

[32] Importantly, despite the efficient induction of HDAC4 in *Mbnl1*$^{\Delta E3/\Delta E3}$ denervated muscle, up-regulation

of *Myog*, *Musk, Chrna1* and *Chrng* was hampered in mutant muscle, as compared to controls (*Figure* 8A).

Moreover, the expression of *Chrne* remained abnormally high in *Mbnl1*$^{\Delta E3/\Delta E3}$ muscle upon denervation

(*Figure* 8B). This suggests that HDAC4-independent mechanisms likely contribute to the defective response

of DM1 muscle to denervation.

To assess the consequences of this defect, we evaluated changes at the endplate after 2 weeks of

denervation. At this stage, endplate fragmentation increased and remained higher in EDL and TA muscles

from *Mbnl1*$^{\Delta E3/\Delta E3}$ mice, compared to controls *(Figure* 8C). To go further, we quantified AChR turnover by

labelling receptors 5 days after nerve injury and assessing their turnover 10 days later. In control muscle,

AChR turnover increased drastically after denervation (*Figure* 8D and 8E), as previously reported.[35] In

Mbnl1$^{\Delta E3/\Delta E3}$ denervated muscle, AChR turnover was reduced as compared to control denervated muscle

(*Figure* 8D and 8E). As there was no difference in AChR recycling between mutant and control denervated

muscles (*Figure* 8F), reduced AChR turnover in DM1 muscle likely arose from reduced insertion and/or

increased degradation of AChRs. Together, these results point to an impaired response of *Mbnl1*$^{\Delta E3/\Delta E3}$

mice to denervation, with reduced AChR formation, which may depend on CaMKII deficiency but unlikely

on HDAC4 deregulation.

DISCUSSION

Although mis-splicing events are at the basis of DM1 pathogenesis, it is not entirely clear how the consecutive perturbations lead to muscle alterations. NMJ deteriorations have been described in muscle biopsies from DM1 patients and in muscle from DM1 mouse models. However, it remains unknown what mechanisms underlie these perturbations and whether these defects arise from DM1-related changes in the muscle or in the nerve. Here, we showed that NMJs are affected in *Mbnl1$^{ΔE3/ΔE3}$* and *HSA^{LR}* mice, two well-characterized mouse models for DM1. We established that CaMKIIs and activity-dependent signalling pathways are disrupted in *Mbnl1$^{ΔE3/ΔE3}$* and *HSA^{LR}* muscles, which may contribute to endplate destabilization and to the abnormal response of mutant muscles to denervation.

Signs of NMJ deterioration, without denervation of muscle fibres, have been reported in muscle biopsies from DM1 patients, as well as in DMSXL and *Mbnl1/2*-deficient mice.[10-13, 16, 36] As the motor neuron is also affected in these mouse models, it is unclear whether the defects arise from pre- or post-synaptic perturbations. Here, we found endplate fragmentation in the muscles of *Mbnl1$^{ΔE3/ΔE3}$* and *HSA^{LR}* mice at different ages. As *HSA^{LR}* mice express the transgene carrying the (CTG) repeats only in muscle, it is likely that perturbations in the post-synaptic compartment, *i.e.* the muscle, are responsible for NMJ deterioration in DM1. Importantly, endplate fragmentation was similar between TA, EDL and *gastrocnemius* muscles, which are affected differentially in *HSA^{LR}* mice. Especially TA and EDL muscles showed minor signs, if not none, of muscle degeneration, suggesting that endplate destabilization is causally unrelated to DM1-associated muscle degeneration. In contrast, NMJ deterioration may contribute, and eventually precede, muscle atrophy, weakness and fatigue observed in DM1 muscle.

Seeking for pathomechanisms that may compromise NMJ integrity in DM1, we examined the potential role of CaMKII deregulation. Mis-splicing in *Camk2b, 2d* and *2g* has been reported in DM1 patients, as well as in mouse models.[3-5] In particular, abnormal exclusion of *Camk2b* exon 13 appeared as one of the most important splicing changes detected in DM1 tissues. Although pathophysiological consequences have

been investigated in the brain, the consequences of CaMKII deregulation in skeletal muscle have not yet

been analysed. We here report that the three exons specifically included in the muscle-specific CaMKIIβ

isoform are excluded in *HSA^LR* and *Mbnl1^ΔE3/ΔE3* muscles, indicating that *Camk2* splicing is MBNL1-

dependent. Of note, in *HSA^LR* mice, *Camk2b* mis-splicing was detected in both *gastrocnemius* and TA/EDL

muscles. Together with the exon 13 and 16, exons 18-20 are part of the highly variable region of the

Camk2b gene, which allows the expression of tissue-specific variants. Especially, CaMKIIβM was shown to

be the only isoform accumulating at the NMJ.[6] Consistent with the abnormal splicing of *Camk2b* in DM1

muscle, CaMKIIβM was not detected in *Mbnl1^ΔE3/ΔE3* and HSA^LR muscles. However, despite the known role

of CaMKIIs in regulating AChR recycling, we did not find perturbation in AChR dynamics at the endplate in

DM1 muscle.

CaMKIIs have also been shown to contribute to synaptic gene repression in non-synaptic regions of

muscle fibres, by inhibiting HDAC4 signalling pathway and myogenin activity.[7, 32] We unveiled that HDAC4

accumulates in myonuclei from DM1 muscle. Increased HDAC4 activity may lead to the up-regulation of

synaptic genes and to the fibre type switch observed in DM1 muscle. Of note, these perturbations did not

arise from spontaneous denervation of mutant muscle fibres. As there was no major change in the

expression of MITR/HDAC9 and myogenin, which are thought to mediate some of the effects of HDAC4,

we cannot rule out that up-regulation of activity-dependent genes in DM1 muscle involves CaMKII/HDAC4-

independent mechanisms. Alternatively, changes detected in DM1 muscle may occur predominantly at

the endplate, as expression of synaptic genes in the sub-synaptic region is independent from

HDAC4/myogenin in innervated muscle.[31] Synaptic gene up-regulation was detected in *Mbnl1^ΔE3/ΔE3*

muscle and in *HSA^LR gastrocnemius* muscle, but not in TA/EDL muscles from *HSA^LR* mice. It is therefore

unlikely that synaptic gene up-regulation causes the endplate fragmentation observed in all muscles. Of

note, the fibre type switch previously reported in *HSA^LR* muscle [23] was exacerbated in *Mbnl1^ΔE3/ΔE3* muscle.

This suggested that perturbations both in the muscle and in non-muscle tissues (*e.g.,* the nerve) contribute to these changes.

To get further insights on the capacity of DM1 muscle to regulate activity-dependent signalling and to maintain their endplates, we challenged *Mbnl1$^{\Delta E3/\Delta E3}$* and *HSA^{LR}* mice with nerve injury. Unexpectedly, transgene expression, driven by the *HSA* promoter, was lost in *HSA^{LR}* muscle upon denervation, which limited the analysis. In *Mbnl1$^{\Delta E3/\Delta E3}$* mice, the increase in AChR turnover triggered by denervation was reduced compared to control denervated muscle. Consistently, up-regulation of synaptic genes was hampered in mutant muscle, which may limit the incorporation of new receptors at the membrane. Up-regulation of synaptic genes upon denervation relies on the release of *Myog* expression, through the repression of *Mitr/Hdac9* and *Dach2* by HDAC4.[8, 37, 38] Expression of these two repressors was efficiently reduced upon denervation in DM1 muscle, which was consistent with the major accumulation of HDAC4 in mutant denervated muscle. Hence, although we cannot rule out the contribution of CaMKII deregulation in the defective up-regulation of synaptic genes in DM1 denervated muscle, it is unlikely to occur *via* deregulation of HDAC4 signalling. Deregulation of distinct CaMKII isoforms may contribute to the different changes observed in DM1 muscle in innervated and denervated conditions, respectively. Changes in the activity of other repressors of *Myog*, such as Msy3/Ybx3 [39] or CtBP1 [40], may be involved as well in the defects observed in DM1 muscle.

In conclusion, our study identified NMJ deterioration as an integral part of muscle dysfunction in DM1, which likely involves muscle perturbations in activity-dependent pathways. Especially, deregulation of CaMKII isoforms may be a key event in endplate destabilization, as well as in DM1 pathogenesis in muscle and non-muscle tissues.

METHODS

Mice

Homozygous mice of the mouse line LR20b carrying about 250 (CTG) repeats within the *HSA* transgene

(*HSA^{LR}*) were obtained from Thornton and colleagues (University of Rochester Medical Centre, Rochester,

New York, USA).[20] Mice of the corresponding background strain (FVB/N) were used as control. Mice were

genotyped for *HSA^{LR}* transgene by quantification of *ACTA1* levels normalized to endogenous actin (mouse

Acta1) in genomic DNA. Mice from the *Mbnl1ΔE3* line were obtained from Swanson and colleagues

(College of Medicine, University of Florida, Gainesville, Florida, USA).[21] *Mbnl1^{+/+}* littermates were used as

control. Mice from the *Mbnl1ΔE3* line were genotyped for exon 3 depletion in the *MBNL1* locus. Mice were

maintained in a conventional specific-pathogen-free facility with a fixed light cycle (23°C, 12-hour dark-

light cycle). All animal studies were performed in accordance with the European Union guidelines for

animal care and approved by the Veterinary Office of the Cantons of Basel city (application number 2601)

and Geneva (application number GE220).

Muscle force and relaxation

In vitro force measurement of EDL muscle and late relaxation time evaluation were conducted as

previously described.[22, 23]

Western blotting

Muscles powdered in liquid nitrogen were lysed in cold RIPA+ buffer (50 mM Tris-HCl pH 8, 150 mM NaCl,

1% NP-40, 0.5% sodium deoxycholate, 0.1% SDS, 1% Triton X, 10% glycerol, phosphatase and protease

inhibitors). Subcellular fractionation was done according to Dimauro et al. (2012).[24] Following dosage (BCA

Protein Assay, Pierce), proteins were separated on SDS-polyacrylamide gels and transferred to

nitrocellulose membrane. Blots were blocked in TBS, 3% BSA, 0.1% Tween-20, and incubated overnight at

4°C with primary antibodies, then for 2 hours with HRP-labelled secondary antibodies. Immunoreactivity

was detected using the ECL Western blot detection reagent LumiGLO (KPL) and exposed to Super RX-N

films (Fujifilm) or revealed with iBright™ Imaging System (ThermoFisher). Protein expression was

normalized to α-actinin or Gadph, or to total protein levels of the corresponding phosphorylated form.

The list of antibodies is provided in Supplementary Materials.

Polymerase chain reaction

Total RNAs were extracted with the RNeasy Mini Kit (Qiagen), reverse transcribed with the SuperScript III

First-Strand Synthesis System (Invitrogen), and amplified with the Power SYBR Green Master Mix (Applied

Biosystems) or the Hot FirePol EvaGreen qPCR Mix (Solis BioDyne). Expression of specific spliced or pan

transcripts was analysed by end-point PCR and electrophoresis, or by quantitative PCR with the Step-One

software and normalization to *Tbp* expression. The list of primers is provided in Table S1.

Histology and immunofluorescence

Muscles were frozen in liquid nitrogen-cooled isopentane. Eight-micrometre muscle sections were stained

with H&E and observed with an upright microscope (Olympus). For immunostaining, sections were fixed

with 4% paraformaldehyde (PFA) or kept unfixed, then blocked in PBS, 3% BSA, incubated sequentially

with primary and secondary fluorescent antibodies (Invitrogen, Jackson ImmunoResearch), mounted with

Vectashield medium (Vector), and observed with Leica or Zeiss fluorescent microscopes. Quantification of

fibre type and size was done as previously reported.[25]

Staining of muscle bundles

To analyse NMJ organization, muscles were bathed *ex vivo* with α-bungarotoxin-Alexa555 (2 μg/ml -

Invitrogen) for 30 min, before being washed and fixed with 4% PFA. Muscle bundles were cut,

permeabilized in PBS, 1% Triton-X100, and blocked in PBS, 1% BSA, 0.1% Triton-X100. Bundles were then successively incubated with primary antibodies against Neurofilament and Synaptophysin (to stain pre-synaptic compartment), and the corresponding secondary antibodies (Invitrogen, Jackson ImmunoResearch). Images were recorded using a Leica confocal microscope.

AChR turnover and recycling

AChR turnover was assessed by injecting α-bungarotoxin-Alexa647 and -Alexa555 (25 pmoles - Invitrogen) into TA/EDL muscles at days 1 and 10, respectively (5 and 14 days after nerve cut when combined with denervation), as previously done.[26] AChR recycling was assessed by injecting α-bungarotoxin-Biotin (25 pmoles - Invitrogen) and saturating dose of streptavidin-Alexa647 (50 pmoles - Invitrogen) into TA/EDL muscles at day 1, followed by injection of streptavidin-Alexa555 (50 pmoles - Invitrogen) at days 2 or 4. Upon AChR internalization, streptavidin-Alexa647 is released from Btx-biotin, which remains bound to AChRs.[27] Only recycled AChRs (*i.e.,* bound to free Btx-biotin) are labelled by streptavidin-Alexa555. We compared the population of recycled receptors (pixels positive for Alexa555) to the population of all receptors labelled with Btx-Biotin (pixels labelled with Alexa555 or Alexa647). For turnover and recycling quantification, images were recorded using Leica or Zeiss confocal microscopes. Pixel dominance (old *vs.* new or stable *vs.* recycled receptors) was calculated using Fiji and Matlab software. The assays could be applied to TA and EDL muscles, but not to *gastrocnemius* muscle because of its size and heterogeneity.

Statistics

Quantitative data are displayed as mean ± SEM of independent samples, with n (number of individual experiments) ≥ 3. Statistical analysis of values was performed using unpaired Student's t test or two-way ANOVA test with Tukey's multiple comparisons test correction, with a 0.05 level of confidence accepted for statistical significance.

Acknowledgements

We thank Prof. C.A. Thornton and Prof. M.S. Swanson for the generous gift of HSA^{LR} and $Mbnl1^{\Delta E3/\Delta E3}$ mice,

as well as Dr.D. Ham and M. Reischl for their help with the macros for fibre typing and AChR turnover.

MHC antibodies, developed by H.M. Blau and S. Schiaffino, were obtained from the Developmental Studies

Hybridoma Bank (University of Iowa, Iowa City, Iowa, USA). This work was supported by the Swiss

Foundation for Research on Muscle Diseases and the Swiss National Science Foundation.

Declaration of interests

The authors declare that they have no conflict of interest.

REFERENCES

Lin X, Miller JW, Mankodi A, Kanadia RN, Yuan Y, Moxley RT, et al. Failure of MBNL1-dependent post-natal splicing transitions in myotonic dystrophy. *Hum Mol Genet* 2006; 15(13):2087-2097.

Miller JW, Urbinati CR, Teng-Umnuay P, Stenberg MG, Byrne BJ, Thornton CA, et al. Recruitment of human muscleblind proteins to (CUG)(n) expansions associated with myotonic dystrophy. *Embo J* 2000; 19(17):4439-4448.

Nakamori M, Sobczak K, Puwanant A, Welle S, Eichinger K, Pandya S, et al. Splicing biomarkers of disease severity in myotonic dystrophy. *Ann Neurol* 2013; 74(6):862-872.

Suenaga K, Lee KY, Nakamori M, Tatsumi Y, Takahashi MP, Fujimura H, et al. Muscleblind-like 1 knockout mice reveal novel splicing defects in the myotonic dystrophy brain. *PLoS One* 2012; 7(3):e33218.

Sobczak K, Wheeler TM, Wang W, Thornton CA. RNA interference targeting CUG repeats in a mouse model of myotonic dystrophy. *Mol Ther* 2013; 21(2):380-387.

Martinez-Pena y Valenzuela I, Mouslim C, Akaaboune M. Calcium/calmodulin kinase II-dependent acetylcholine receptor cycling at the mammalian neuromuscular junction in vivo. *J Neurosci* 2010; 30(37):12455-12465.

Tang H, Macpherson P, Argetsinger LS, Cieslak D, Suhr ST, Carter-Su C, et al. CaM kinase II-dependent phosphorylation of myogenin contributes to activity-dependent suppression of nAChR gene expression in developing rat myotubes. *Cell Signal* 2004; 16(5):551-563.

Tang H, Macpherson P, Marvin M, Meadows E, Klein WH, Yang XJ, et al. A histone deacetylase 4/myogenin positive feedback loop coordinates denervation-dependent gene induction and suppression. *Mol Biol Cell* 2009; 20(4):1120-1131.

Macpherson P, Kostrominova T, Tang H, Goldman D. Protein kinase C and calcium/calmodulin-activated protein kinase II (CaMK II) suppress nicotinic acetylcholine receptor gene expression in

mammalian muscle. A specific role for CaMK II in activity-dependent gene expression. *J Biol Chem* 2002; 277(18):15638-15646.

Macdermot V. The histology of the neuromuscular junction in dystrophia myotonica. *Brain* 1961; 84:75-84.

Allen DE, Johnson AG, Woolf AL. The intramuscular nerve endings in dystrophia myotonica--a biopsy study by vital staining and electron microscopy. 1969; 105(Pt 1):1-26.

Coers C, Telerman-Toppet N, Gerard JM. Terminal innervation ratio in neuromuscular disease. II. Disorders of lower motor neuron, peripheral nerve, and muscle. *Arch Neurol* 1973; 29(4):215-222.

Engel A, Jerusalem F, Tsujihata M, Gomez M. The neuromuscular junction in myopathies: a quantitative ultrastructural study. In: *Recent Advances in Myology: Proceedings of the Third International Congress on Muscle Diseases, International Congress Series no 360.* Edited by Bradley W, Gardner-Medwin D, Walton J. New York: Elsevier; 1975: 132-143.

Drachman DB, Fambrough DM. Are muscle fibers denervated in myotonic dystrophy? *Arch Neurol* 1976; 33(7):485-488.

Walton JN, Irving D, Tomlinson BE. Spinal cord limb motor neurons in dystrophia myotonica. *J Neurol Sci* 1977; 34(2):199-211.

Panaite PA, Kuntzer T, Gourdon G, Lobrinus JA, Barakat-Walter I. Functional and histopathological identification of the respiratory failure in a DMSXL transgenic mouse model of myotonic dystrophy. *Dis Model Mech* 2013; 6(3):622-631.

Spilker KA, Wang GJ, Tugizova MS, Shen K. Caenorhabditis elegans Muscleblind homolog mbl-1 functions in neurons to regulate synapse formation. *Neural Dev* 2012; 7:7.

Wheeler TM, Krym MC, Thornton CA. Ribonuclear foci at the neuromuscular junction in myotonic dystrophy type 1. *Neuromuscul Disord* 2007; 17(3):242-247.

Klinck R, Fourrier A, Thibault P, Toutant J, Durand M, Lapointe E, et al. RBFOX1 cooperates with MBNL1 to control splicing in muscle, including events altered in myotonic dystrophy type 1. *PLoS One* 2014; 9(9):e107324.

Mankodi A, Logigian E, Callahan L, McClain C, White R, Henderson D, et al. Myotonic dystrophy in transgenic mice expressing an expanded CUG repeat. *Science* 2000; 289(5485):1769-1773.

Kanadia RN, Johnstone KA, Mankodi A, Lungu C, Thornton CA, Esson D, et al. A muscleblind knockout model for myotonic dystrophy. *Science* 2003; 302(5652):1978-1980.

Castets P, Lin S, Rion N, Di Fulvio S, Romanino K, Guridi M, et al. Sustained activation of mTORC1 in skeletal muscle inhibits constitutive and starvation-induced autophagy and causes a severe, late-onset myopathy. *Cell Metab* 2013; 17(5):731-744.

Brockhoff M, Rion N, Chojnowska K, Wiktorowicz T, Eickhorst C, Erne B, et al. Targeting deregulated AMPK/mTORC1 pathways improves muscle function in myotonic dystrophy type I. *J Clin Invest* 2017; 127(2):549-563.

Dimauro I, Pearson T, Caporossi D, Jackson MJ. A simple protocol for the subcellular fractionation of skeletal muscle cells and tissue. *BMC research notes* 2012; 5:513.

Ham DJ, Borsch A, Lin S, Thurkauf M, Weihrauch M, Reinhard JR, et al. The neuromuscular junction is a focal point of mTORC1 signaling in sarcopenia. *Nat Commun* 2020; 11(1):4510.

Castets P, Rion N, Theodore M, Falcetta D, Lin S, Reischl M, et al. mTORC1 and PKB/Akt control the muscle response to denervation by regulating autophagy and HDAC4. *Nat Commun* 2019; 10(1):3187.

Bruneau E, Sutter D, Hume RI, Akaaboune M. Identification of nicotinic acetylcholine receptor recycling and its role in maintaining receptor density at the neuromuscular junction in vivo. *J Neurosci* 2005; 25(43):9949-9959.

Wang ET, Treacy D, Eichinger K, Struck A, Estabrook J, Olafson H, et al. Transcriptome alterations in myotonic dystrophy skeletal muscle and heart. *Hum Mol Genet* 2019; 28(8):1312-1321.

Strack S, Khan MM, Wild F, Rall A, Rudolf R. Turnover of acetylcholine receptors at the endplate revisited: novel insights into nerve-dependent behavior. *J Muscle Res* 2015; 36(6):517-524.

Wang AH, Yang XJ. Histone deacetylase 4 possesses intrinsic nuclear import and export signals. *Mol Cell Biol* 2001; 21(17):5992-6005.

Tintignac LA, Brenner HR, Ruegg MA. Mechanisms Regulating Neuromuscular Junction Development and Function and Causes of Muscle Wasting. *Physiol Rev* 2015; 95(3):809-852.

Cohen TJ, Waddell DS, Barrientos T, Lu Z, Feng G, Cox GA, et al. The histone deacetylase HDAC4 connects neural activity to muscle transcriptional reprogramming. *J Biol Chem* 2007; 282(46):33752-33759.

Moresi V, Williams AH, Meadows E, Flynn JM, Potthoff MJ, McAnally J, et al. Myogenin and class II HDACs control neurogenic muscle atrophy by inducing E3 ubiquitin ligases. *Cell* 2010; 143(1):35-45.

Akaaboune M, Culican SM, Turney SG, Lichtman JW. Rapid and reversible effects of activity on acetylcholine receptor density at the neuromuscular junction in vivo. *Science* 1999; 286(5439):503-507.

Strack S, Petersen Y, Wagner A, Roder IV, Albrizio M, Reischl M, et al. A Novel Labeling Approach Identifies Three Stability Levels of Acetylcholine Receptors in the Mouse Neuromuscular Junction In Vivo. *PloS one* 2011; 6(6):e20524.

Lee KY, Li M, Manchanda M, Batra R, Charizanis K, Mohan A, et al. Compound loss of muscleblind-like function in myotonic dystrophy. *EMBO Mol Med* 2013; 5(12):1887-1900.

37   Cohen TJ, Barrientos T, Hartman ZC, Garvey SM, Cox GA, Yao TP. The deacetylase HDAC4 controls

myocyte enhancing factor-2-dependent structural gene expression in response to neural activity.

FASEB J 2009; 23(1):99-106.

38   Mejat A, Ramond F, Bassel-Duby R, Khochbin S, Olson EN, Schaeffer L. Histone deacetylase 9

couples neuronal activity to muscle chromatin acetylation and gene expression. *Nat Neurosci*

2005; 8(3):313-321.

39   De Angelis L, Balasubramanian S, Berghella L. Akt-mediated phosphorylation controls the activity

of the Y-box protein MSY3 in skeletal muscle. *Skelet muscle* 2015; 5:18.

40   Thomas JL, Moncollin V, Ravel-Chapuis A, Valente C, Corda D, Mejat A, et al. PAK1 and CtBP1

Regulate the Coupling of Neuronal Activity to Muscle Chromatin and Gene Expression. *Mol Cell*

Biol 2015; 35(24):4110-4120.

Figure 1

Figure 1: Muscle affection in *Mbnl1^{ΔE3/ΔE3}* and *HSA^{LR}* mice. A, B) H&E coloration reveals limited

histopathologic alterations in muscles from 3- and 9-month-old *Mbnl1^{ΔE3/ΔE3}* (A) and 3- and 12-month-old

HSA^{LR} mice (B). Asterisks and arrows point to internalized nuclei and vacuoles, respectively. Scale bar, 100

μm. C, D) Late relaxation time upon stimulation is increased in EDL muscle from 3-month-old *Mbnl1^{ΔE3/ΔE3}*

and *HSA^{LR}* mice, as compared to control mice. n = 4 Ctrl / 3 *Mbnl1^{ΔE3/ΔE3}*; 9 Ctrl / 30 *HSA^{LR}*. E) Specific tetanic

force (sP0) of EDL muscle is unchanged in 3-month-old $Mbnl1^{\Delta E3/\Delta E3}$ (n = 4 Ctrl / 3 KO) and HSA^{LR} (n= 9 Ctrl

/ 30 HSA^{LR}) mice. F-H) Quantitative RT-PCR analysis of the inclusion of $Clcn1$ exon7a (F), of total $Clcn1$

transcript (G), and of the inclusion of $Atp2a1$ exon22 (H) in TA muscle from $Mbnl1^{\Delta E3/\Delta E3}$ mice, and

$gastrocnemius$ or TA muscle from HSA^{LR} mice. Expression of spliced variants is normalised on total

expression. Total expression is normalized on Tbp expression. n = 4 Ctrl / 3 $Mbnl1^{\Delta E3/\Delta E3}$ (for $Clcn1$); 6 Ctrl

/ 5 $Mbnl1^{\Delta E3/\Delta E3}$ (for $Atp2a1$); 4 Ctrl / 4 HSA^{LR}. In all panels, data represent mean ± SEM. *p<0.05, **p<0.01,

***p<0.001, ****p<0.0001, unpaired two-tailed Student's t test.

Figure 2

Figure 2: CaMKIIβ deregulation in *Mbnl1^ΔE3/ΔE3* and *HSA^LR* muscles. A-C) Western blot analysis of CaMKII

isoforms and quantification of CaMKIIβM levels in TA muscle from 3-month-old *Mbnl1^ΔE3/ΔE3* mice (A) and

in *gastrocnemius* muscle from 3-month-old *HSA^LR* mice (B). Quantification of CaMKIIβM phosphorylated

form in *Mbnl1^ΔE3/ΔE3* and *HSA^LR* muscles is shown in C. Normalization is made on α-actinin (A, B) or on total

CaMKIIβM (C), relative to control. n = 3 (Ctrl/*Mbnl1^ΔE3/ΔE3*) and 4 (Ctrl/*HSA^LR*) per group. D-F) Quantitative

RT-PCR analysis of total levels of *Camk2b* (D), of the exclusion of exon 13 (E), and of the inclusion of exons

18-20 (F) in TA muscle from *Mbnl1^ΔE3/ΔE3* mice and in *gastrocnemius*, TA and EDL muscles from *HSA^LR* mice.

Data are normalized on levels of *Tbp* (D) or on total *Camk2b* transcripts (E, F). n = 4 Ctrl / 3 *Mbnl1^ΔE3/ΔE3*; 4

Ctrl / 4 *HSA^LR* (D, E); 3 Ctrl / 3 *HSA^LR* (F). All data are mean ± SEM; * p<0.05; ** p<0.01; **** p<0.0001; two-

tailed unpaired Student's t-test.

Figure 3

Figure 3: Altered NMJ maintenance in *Mbnl1^ΔE3/ΔE3* and *HSA^LR* mice. A, B) Fluorescent images of NMJ

regions stained with α-bungarotoxin (Btx, red), antibodies against Neurofilament/Synaptophysin (green),

and Dapi (blue) in EDL muscles from 3- and 9-month-old *Mbnl1^ΔE3/ΔE3* mice (A) and 3-and 12-month-old

HSA^LR mice (B). Scale bar, 50 μm. C, D) Quantification of the number of fragments per NMJ in EDL, TA and

gastrocnemius muscles from 3- and 9/12-month-old *Mbnl1^ΔE3/ΔE3* (C) and *HSA^LR* (D) mice. n=5/8 (EDL 3M),

3/4 (TA 3M), 5/3 (all muscles 9M) Ctrl/ *Mbnl1^ΔE3/ΔE3* (C); 11/12 (EDL 3M), 7/8 (TA 3M), 7/8 (GA 3M), 4/3

(EDL 12M), 4/4 (TA and GA 12M) Ctrl/*HSA^LR* (D), with more than 52 fibres per muscle. Data represent mean

± SEM. **p<0.01, ***p<0.001, ****p<0.0001, unpaired two-tailed Student's t test.

Figure 4

Figure 4: AChR dynamics is not altered in *Mbnl1$^{\Delta E3/\Delta E3}$* and *HSA^{LR}* muscles. A) Turnover assay in EDL and

TA muscles from *Mbnl1$^{\Delta E3/\Delta E3}$* and *HSA^{LR}* mice. Fluorescent images show "old" (green) and "new" (red)

AChRs in *Mbnl1$^{\Delta E3/\Delta E3}$* and *HSA^{LR}* muscles. Scale bar, 50 µm. B) Timeline of injections of α-bungarotoxin

(Btx) for AChR turnover and recycling assays. *d, days; Strp, streptavidin.* C, D) AChR turnover in EDL and TA

muscles from *Mbnl1$^{\Delta E3/\Delta E3}$* (C) and *HSA^{LR}* (D) mice. n = 10/11 (EDL) and 5/5 (TA) Ctrl/*Mbnl1$^{\Delta E3/\Delta E3}$* (C); 7/8

(EDL) and 3/3 (TA) Ctrl/*HSA^{LR}* (D), with more than 22 fibres per muscle. E, F) AChR recycling in EDL muscle

from *Mbnl1$^{\Delta E3/\Delta E3}$* mice. Fluorescent images show "stable" (green) and "recycled" (red) AChRs in

Mbnl1$^{\Delta E3/\Delta E3}$ muscles (E). Scale bar, 50 µm. Quantification is given in F. n = 4/5 (1d); 4/4 (3d)

Ctrl/*Mbnl1$^{\Delta E3/\Delta E3}$* per group (F), with more than 28 fibres per muscle. Data are mean ± SEM; *** p<0.001;

two-tailed unpaired Student's t-test.

Figure 5: Changes in HDAC4 signalling pathway in *Mbnl1^{ΔE3/ΔE3}* and *HSA^{LR}* muscles. A) Quantitative RT-

PCR analysis of *Hdac4* in *Mbnl1^{ΔE3/ΔE3}* and *HSA^{LR}* muscles. n = 4/3 Ctrl/*Mbnl1^{ΔE3/ΔE3}* and 4/4 Ctrl/ *HSA^{LR}*. B,

C) Western blot analysis of HDAC4 and its phosphorylated form (Ser632) in total protein lysate of

Mbnl1^{ΔE3/ΔE3} (B) and *HSA^{LR}* (C) muscles. Quantification of total and phosphorylated levels is given for

Mbnl1^{ΔE3/ΔE3} (B) and *HSA^{LR}* (C) muscles. Total and phosphorylated levels are normalized to α-actinin and

total HDAC4, respectively, and relative to control. n = 3 per group. D) Western blot analysis of HDAC4 and

acetylated histones (H3K9) in nuclear fraction of *gastrocnemius* muscle from *Mbnl1$^{\Delta E3/\Delta E3}$* mice and *HSA^{LR}*

mice. Quantifications of HDAC4 and H3K9 levels in *Mbnl1$^{\Delta E3/\Delta E3}$* and *HSA^{LR}* muscles are given in (D) and in

Figure S5, respectively. Levels are normalized to lamin, and relative to control. n = 5/4 Ctrl/*Mbnl1$^{\Delta E3/\Delta E3}$*

and 3/4 Ctrl/*HSA^{LR}*. E) Fluorescent image of *Mbnl1$^{\Delta E3/\Delta E3}$* and *HSA^{LR}* muscles stained with antibodies against

HDAC4 (red), Laminin (green) and Dapi (blue). Scale bar, 100 µm. Higher magnification panel shows

positive myonuclei. F) Quantitative RT-PCR analysis of activity-dependent genes, *Dach2*, *Mitr* and *Myog* in

TA muscle from *Mbnl1$^{\Delta E3/\Delta E3}$* mice and in *gastrocnemius* muscle from *HSA^{LR}* mice. Transcript levels are

normalized to *Tbp* and relative to control. n = 4/3 Ctrl/*Mbnl1$^{\Delta E3/\Delta E3}$* and 4/4 Ctrl/*HSA^{LR}*. All data are mean

± SEM; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; two-tailed unpaired Student's t-test.

Figure 6

Figure 6: DM1 mice display deregulation of activity-dependent signalling pathways. A) Quantitative RT-

PCR analysis of *Musk, Chrna1, Chrne* and *Chrng* in TA muscle from 3-month-old *Mbnl1$^{ΔE3/ΔE3}$* mice and in

gastrocnemius muscle from 3-month-old *HSA^{LR}* mice. Levels are normalized to *Tbp*. n=6/5 Ctrl/ *Mbnl1$^{ΔE3/ΔE3}$*

and 5/7 Ctrl/*HSA^{LR}* (except for *Chrna1*, n=4). B) Quantitative RT-PCR analysis of *Myh2* and *Myh4*, encoding

MHC2A and MHC2B, in *Mbnl1$^{ΔE3/ΔE3}$* muscle. Transcript levels are normalized to *Tbp* and relative to control.

n=6/5 Ctrl/*Mbnl1$^{ΔE3/ΔE3}$*. C) Fluorescent images of control and *Mbnl1$^{ΔE3/ΔE3}$* muscles, stained with antibodies

against MHC2A (red) and Laminin (green), and quantification of the proportion of type IIA, IIX and IIB fibres

in control and mutant muscles. Scale bar, 500 µm. D) Quantitative RT-PCR analysis of *Myh2* and *Myh4* in

HSA^{LR} TA muscle. Transcript levels are normalized to *Tbp* and relative to control. n=3/4 Ctrl/ *HSA^{LR}*. All data

are mean ± SEM; * p<0.05; ** p<0.01; *** p<0.001; two-tailed unpaired Student's t-test.

Figure 7

Figure 7: *Mbnl1^{ΔE3/ΔE3}* **muscle shows efficient HDAC4 induction and resistance to atrophy upon**

denervation. A) Quantitative RT-PCR of *Hdac4* transcript and Western blot analysis of HDAC4 protein in

innervated (In) and denervated (De, 3 days) muscles from control and *Mbnl1^{ΔE3/ΔE3}* mice. Transcript levels

are normalized to *Tbp* and relative to control innervated. Levels in the innervated control muscle are

shown by the red dotted line. Protein levels of HDAC4 are normalized to α-actinin, relative to control and

expressed as the log2(Fold Change). n=4 (qPCR) and 4/3 (WB). B) Fluorescent image of *Mbnl1^{ΔE3/ΔE3}*

denervated muscle stained with antibodies against HDAC4 (red), Laminin (green) and Dapi (blue). Scale

bar, 100 μm. Higher magnification panel shows positive myonuclei. C) Quantitative RT-PCR analysis of

Dach2 and *Mitr* in TA muscle from control and *Mbnl1$^{\Delta E3/\Delta E3}$* mice 3 days post-denervation. Levels are

normalized to *Tbp* and relative to control innervated. n=4 per group. Levels in the innervated control

muscle are shown by the red dotted line. D) Mass variation after 3 weeks of denervation in control and

Mbnl1$^{\Delta E3/\Delta E3}$ mice, for TA and *gastrocnemius* (GA) muscles. n = 4 per group. E) Fluorescent images of

denervated muscles from control and *Mbnl1$^{\Delta E3/\Delta E3}$* mice, stained with antibodies against type IIA myosin

heavy chain (MHC2A - red) and Laminin (green) and quantification of fibre type proportion in control and

mutant denervated muscles. Scale bar, 500 µm. n = 4 mice per group. F) Quantification of fibre minimum

ferret in innervated and 3-week-denervated muscles from control and *Mbnl1$^{\Delta E3/\Delta E3}$* mice. n = 4 mice per

group. All data are mean ± SEM; * p<0.05; ** p<0.01; *** p<0.001; **** p<0.0001; two-tailed unpaired

Student's t-test (D) and two-way ANOVA with a Tukey's post-hoc analysis (A, F).

Figure 8

Figure 8: Denervation-induced changes in the expression and dynamics of synaptic proteins are impaired

in *Mbnl1^{ΔE3/ΔE3}* mice. A, B) Quantitative RT-PCR analysis of activity-dependent genes, *Myog, Musk, Chrna1*

and *Chrng* (A), and *Chrne* (B), in TA muscle from control and *Mbnl1^{ΔE3/ΔE3}* mice 3 days post-denervation.

Levels are normalized to *Tbp* and relative to control innervated. n = 4 per group. Levels in the innervated

control muscle are shown by the red dotted line. C) Quantification of endplate fragmentation in EDL and

TA muscles from *Mbnl1^{ΔE3/ΔE3}* mice after 3 weeks of denervation. n = 3/4 Ctrl/KO. D-F) AChR turnover (D,

E) and recycling (F) in denervated muscle from *Mbnl1^{ΔE3/ΔE3}* and control mice. Fluorescent images of "old"

and "new" receptors are shown in (D). Scale bar, 50 μm. Quantification of AChR turnover and of AChR

recycling is giver in (E) and (F), respectively. n = 10/11 (EDL) and 5/5 (TA) Ctrl/*Mbnl1^{ΔE3/ΔE3}* (E); n=5 per

group (F). Data are mean ± SEM; ** p<0.01; *** p<0.001; two-tailed unpaired Student's t-test.