Leaf metabolic traits reveal hidden dimensions of plant form and function

Tom W. N. Walker[1,2]*, Franziska Schrodt[3], Pierre-Marie Allard[4,5], Emmanuel Defossez[2], Vincent E. J. Jassey[6], Meredith C. Schuman[23], Jake M. Alexander[1], Oliver Baines[3], Virginie Baldy[7], Richard D. Bardgett[8], Pol Capdevila[9,10], Phyllis D. Coley[12], Nicole M. van Dam[13,14,15], Bruno David[16], Patrice Descombes[1,26,27], Maria-Jose Endara[12], Catherine Fernandez[7], Dale Forrister[12], Albert Gargallo-Garriga[17,18,19], Gaëtan Gauser[2], Sue Marr[13,20,21], Steffen Neumann[13,20], Loïc Pellissier[1,26], Kristian Peters[13,20,21], Sergio Rasmann[2], Ute Roessner[22], Roberto Salguero-Gómez[11], Jordi Sardans[17,18], Wolfram Weckwerth[24,25], Jean-Luc Wolfender[4,5] & Josep Peñuelas[17,18]

[1] Department of Environmental Systems Science, ETH Zürich, 8092 Zürich, Switzerland
[2] Institute of Biology, University of Neuchâtel, 2000 Neuchâtel, Switzerland.
[3] School of Geography, University of Nottingham, Nottingham NG7 2RD, UK
[4] School of Pharmaceutical Sciences, University of Geneva, 1211 Geneva, Switzerland
[5] Institute of Pharmaceutical Sciences of Western Switzerland, University of Geneva, 1211 Geneva, Switzerland
[6] Laboratoire d'Ecologie Fonctionnelle et Environnement, Université de Toulouse, CNRS, 31062 Toulouse, France
[7] Aix Marseille Université, Avignon Université, CNRS, IRD, IMBE, Marseille, France
[8] Department of Earth & Environmental Sciences, The University of Manchester, Manchester M13 9PT, UK
[9] Departament de Biologia Evolutiva, Ecologia i Ciències Ambientals, Universitat de Barcelona, 08028 Barcelona, Spain
[10] School of Biological Sciences, University of Bristol, Bristol BS8 1TQ, UK
[11] Department of Zoology, Oxford University, Oxford OX1 3SZ, UK
[12] School of Biological Sciences, University of Utah, Salt Lake City, UT 84112, USA
[13] German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, 04103 Leipzig, Germany
[14] Institute of Biodiversity, Friedrich Schiller University Jena, 07743 Jena, Germany
[15] Leibniz Institute of Vegetable and Ornamental crops (IGZ), 14979 Großbeeren, Germany
[16] Green Mission Pierre Fabre, Institut de Recherche Pierre Fabre, 31562 Toulouse, France
[17] CSIC, Global Ecology Unit CREAF-CSIC-UAB, 08193 Bellaterra, Catalonia, Spain
[18] CREAF, 08193 Cerdanyola del Vallès, Catalonia, Spain
[19] Global Change Research Institute, Czech Academy of Sciences, 603 00 Brno, Czech Republic
[20] Leibniz Institute of Plant Biochemistry, Bioinformatics and Scientific Data, 06120 Halle, Germany
[21] Institute of Biology/Geobotany and Botanical Garden, Martin Luther University Halle Wittenberg, 06108 Halle, Germany
[22] Research School of Biology, The Australian National University, 2601 Acton, Australia
[23] Departments of Geography and Chemistry, University of Zurich, 8057 Zurich, Switzerland
[24] Molecular Systems Biology, Department of Functional and Evolutionary Ecology, 1010 University of Vienna, Austria
[25] Vienna Metabolomics Center, 1010 University of Vienna, Austria
[26] Ecosystems & Landscape Evolution, Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), 8903 Birmensdorf, Switzerland
[27] Musée et Jardins botaniques cantonaux, 1007 Lausanne, Switzerland
* Correspondence to: thomas.walker@unine.ch

Summary

The plant metabolome encompasses the biochemical mechanisms through which evolutionary and ecological processes shape plant form and function[1,2]. However, while the metabolome should thus be an important component of plant life-history variation[3], we know little about how it varies across the plant kingdom. Here, we use the plant functional trait concept[4] – a powerful framework for describing plant form and function[5–7] – to interpret leaf metabolome variation among 457 tropical and 339 temperate plant species. Distilling metabolite chemistry into five discriminant metabolic functional traits reveals that plants vary along two major axes of leaf metabolic specialization – a leaf chemical defense spectrum and an expression of leaf longevity. These axes are qualitatively consistent for tropical and temperate species, with many trait combinations being viable. However, axes of leaf metabolic specialization vary orthogonally to life-history strategies described by widely used functional traits[5–7], while being at least equally important to them. Our findings question classical trait[6] and plant defense[8] theory that predicts relationships between the leaf chemical phenotype, plant productivity, and pace of life. Moreover, we show that metabolic functional traits describe unique dimensions of plant life-history variation that are complementary to, and independent from, those captured by existing plant functional traits.

Main Text

Plants produce a staggering diversity of metabolites – upwards of one million throughout the plant kingdom and many thousands within an individual[9] (the metabolome). The plant metabolome is well known to human society, being the source of most cosmetics[10], flavors[11], and medicines[12]. Yet, little is known about how the metabolome varies across the plant kingdom. The metabolome is the biochemical basis of physiology, comprising primary metabolites involved in cellular function (e.g., carbohydrates, amino acids) and specialized metabolites produced to confront stress (e.g., flavonoids, terpenoids)[9,13,14]. The metabolome thus encompasses the biochemical mechanisms through which evolutionary and ecological processes shape plant form and function (i.e., morphology and physiology [7])[1,2]. Plant form and function vary predictably among species because evolutionary and ecological processes generally act to maximize fitness[8,15–18], thus limiting the number of morphological or physiological strategies that are viable[5–7]. It follows that the metabolome should also be constrained to a limited number of strategies that maximize fitness[3]. However, although cross-species studies are now emerging[19–22], no attempt has been made to characterize how the metabolome varies systematically throughout the plant kingdom. We thus remain unable to place the vast diversity of the plant metabolome into a coherent ecological context.

The plant functional trait concept offers a powerful framework to contextualize the plant metabolome. Plant functional traits are a standardized set of morphological and physiological characteristics that provide universal metrics of plant form and function[4]. Plant functional traits have been pivotal in identifying the life-history strategies that govern plant fitness[5–7], understanding plant impacts on population[23], community[24], and ecosystem[25] processes, and predicting global change effects on ecosystems[26]. Plant functional traits thus provide a strong reference point from which to interpret variation in the plant metabolome. However, measurements of the metabolome remain absent from the functional trait concept, in part because they yield complex data describing thousands of metabolites from hundreds of biochemical pathways[3]. Advances in chemoinformatics make it now possible to translate metabolite identities into a targeted set of chemical properties[27,28]. Doing so resonates with the functional trait concept – i.e., that quantitative characteristics provide a bridge between identity and function[4] – and is an intuitive way to contextualize the metabolome against plant functional traits. Yet, while chemoinformatics is commonplace in natural products chemistry[29–31], it remains entirely absent from ecology.

Contextualizing the plant metabolome against existing plant functional traits raises two alternative hypotheses. If the metabolome varies colinearly with functional traits, then it offers a biochemical validation of the functional trait concept. Such validation is necessary because functional traits are emergent properties of multiple biochemical processes[4,32], making them proxies for physiology that can be poor predictors of ecological processes[33–35]. Alternatively, if the metabolome varies orthogonally to functional traits, then it describes dimensions of plant life-history variation missed by existing functional traits. Such orthogonal variation would enhance the explanatory power of the functional trait concept by providing additional axes of trait specialization to examine plant form and function. Either way, integrating measurements of the plant metabolome into the plant functional trait concept will yield insight into how the metabolome varies across the plant kingdom, while also offering a high-resolution mechanistic lens with which to advance the functional trait concept itself.

105
106
Figure 1 | Variation in leaf metabolite chemistry can be captured by five chemical properties. (A) Coefficients for
Pearson correlations among 21 chemical properties (Table S1) of annotated metabolites detected in tropical and temperate
datasets (N = 4292). Properties are ordered based on correlation similarity (hierarchical clustering; Ward), with colors
separating distinct clusters of properties (maximal silhouette width: K = 5). The lower panel shows correlations between
chemical properties and scores from the first three axes of a PCA performed on a subset of five properties, as well as (B)
between subset PC scores and either scores from a PCA performed on all 21 properties (left panel) or metabolite positions
on the first three axes of ChemGPS-NP[31] space (right panel). Circle sizes indicate coefficient strengths, while fills
discriminate between positive (black) and negative (white) correlations. (C) The chemical properties selected to represent
each cluster, alongside chemical graphs of metabolites with the lowest and highest values for each trait.

Here, we performed an assessment of the leaf metabolomes of 457 tropical and 339 temperate plant
species to characterize variation in the metabolome and test whether it validates, or expands upon,
existing plant functional traits. First, we examined the chemical properties of all identified metabolites
(N = 4292) to derive a minimum set of "metabolic functional traits" that capture chemical variation in
the leaf metabolome. We then characterized how metabolic traits vary among species to identify major
axes of leaf metabolic specialization in the plant kingdom. Finally, we combined metabolic traits with
eight widely used functional traits (plant height, seed mass, stem specific density, leaf area, specific
leaf area and leaf carbon, nitrogen and phosphorus concentrations; hereafter "classical traits")[7] to
determine whether axes of metabolic specialization are colinear with, or orthogonal to, plant life-history
strategies described by classical functional traits.
126
<u>Five chemical properties capture chemical variation in the leaf metabolome</u>
We expressed the leaf metabolomes of tropical[36] and temperate[37] plant species as the presence or
absence of annotated metabolic features (i.e., metabolites) detected through liquid chromatography
mass spectrometry (tropical N = 3356; temperate N = 2227). We then characterized the chemistry of
the leaf metabolome by examining variation in the chemical properties of all unique metabolites from
both datasets (N = 4292)[31]. We focused on 21 properties describing metabolite constitution, geometry,
or topology that have relationships with (bio)chemical activity (Table S1). Pairwise correlations among
chemical properties (Fig. 1A, upper panel) reveal that properties separate into five clusters, each
representing a distinct facet of leaf metabolite chemistry. The largest cluster (Fig. 1A, green) describes
overall size and structural complexity, comprising total numbers of atoms and bonds, molecular
weight, a predictor of polarity based on the proportion of non-carbon atoms (M log P)[38], and related
topological parameters (i.e., Wiener numbers[38], Eccentric Connectivity[39]). A second cluster (Fig. 1A,
purple) describes bond conjugation and reactivity, comprising numbers of aromatic atoms and bonds,
numbers of atoms in the largest conjugated (pi-) system, and a derived complexity index indicative of
promiscuous activity in drugs (f_{MF})[40]. A third cluster (Fig. 1A, yellow) describes polar intermolecular
(non-covalent) forces, comprising total and mass-specific polar surface area and numbers of hydrogen
(H) bond donors or acceptors[41,42]. Such forces are important for the selective binding of small
molecules to DNA[43], RNA[44], and proteins[45]. A fourth cluster (Fig. 1A, blue) comprises two predictors of
metabolite polarity based on summed properties of atoms and bonds (X log P^{46}, A log P^{47}), where
lower values indicate higher polarity. A final cluster (Fig. 1A, red) describes carbon bond saturation *via*
the relative occurrence of sp^3 (unsaturated, three-dimensional) and sp^2 (saturated, planar)
hybridization[48]. Overall, these clusters indicate that leaf metabolite chemistry is highly structured and
can be described by five distinct facets: size/complexity, bond conjugation/reactivity, contributions to
polar intermolecular forces, general polarity, and carbon bond saturation.
151
The strong collinearity among chemical properties (Fig. 1A, upper panel) suggests that the five facets
of leaf chemistry can be represented by a small number of discriminant variables. Indeed, a PCA
performed on a subset of five chemical properties, one from each cluster (Fig. 1C), is equivalent to a
PCA performed on all 21 properties (Fig. 1B, left panel; PC1: r_{4290} = 0.97; PC2: r_{4290} = 0.99; PC3: r_{4290}
= 0.93; P < 0.001 in all cases). We selected molecular weight, H-bond acceptor count, aromatic atom
count, polarity (X log P), and carbon hybridization (sp^3:sp^2) as five properties that are intuitive for non-
specialists to interpret (Fig. 1C; Table S1). However, most combinations of properties covering all
facets of leaf chemistry also yield equivalent PCAs (Fig. S4). Pairwise correlations between all
properties and subset PC scores further confirm that selected properties capture observed leaf
chemical variation, in that correlations are consistent within each cluster (Fig. 1A, lower panel).

Selected properties also describe biologically relevant chemical space according to an existing
framework that classifies natural products based on size, then aromaticity, then lipophilicity (Fig. 1B,
right panel; PC1: r_{4290} = 0.99; PC2: r_{4290} = 0.87; r_{4290} = 0.81; P < 0.001 in all cases)[31]. The first axis of
the subset PCA (38% variance) separates heavy metabolites with many H-bond acceptors from light
metabolites with fewer H-bond acceptors. Alignment between metabolite size and numbers of H-bond
acceptors is expected because larger molecules usually engage in stronger intermolecular
interactions[49]. The second axis (37% variance) separates reactive metabolites with more aromatic
carbon bonds from unreactive metabolites with more saturated carbon bonds. This axis is also intuitive
because conjugated bond systems are unsaturated, reactive, and planar, whereas saturated bonds,
such as those abundant in fatty acids, are unreactive and three-dimensional[50]. The third axis (18%
variance) separates nonpolar metabolites from polar metabolites[29]. Our findings thus show that leaf
chemical variation can be represented by five chemical properties, one from each facet of leaf
chemistry.
175
<u>Five chemical properties provide more information than metabolite family</u>
The five selected chemical properties not only represent leaf metabolite chemistry (Fig. 1), but also
reflect known differences among metabolite families. Using the subset PCA to interpret variation
among metabolite families (Fig. 2), we observe that peptides are large, display average bond
saturation, and are moderately polar; carbohydrates are of average size, display high bond saturation,
and are highly polar; and fatty acids are smaller, display high bond saturation, and are of average
polarity. Lignans, alkaloids, coumarins and flavonoids, families of specialized metabolites, display low
bond saturation due to the presence of many aromatic bonds, with alkaloids and coumarins also being
relatively small. Terpenoids, a family of abundant specialized metabolites (N = 2306), vary in size,
bond saturation, and polarity, echoing their diversity across the plant kingdom[51]. Although chemical
properties generally differentiate among metabolite families, all families possess variable and
overlapping chemistries (Fig. 2, marginal boxplots). We thus suggest that chemical properties,
quantitative metrics of the chemistry that underpins metabolite composition and activity, provide more
precise information about metabolite function than metabolite family identity. Indeed, drug discovery
and medicinal chemistry rely on many of the same chemical properties to screen for bioactive
molecules[29–31]. Such a perspective resonates with the functional trait concept by reaching beyond
taxonomy to describe function[4] and provides a strong foundation for a novel set of metabolic functional
traits.

194
195
Figure 2 | Five chemical properties describe variation among and within metabolite families. Biplots of PC1 scores
and (A) PC2 scores or (B) PC3 scores from a PCA describing variation in selected chemical properties (C hybridization, H-
bond acceptors, molecular weight, polarity, aromatic atoms) among all annotated leaf metabolites (N = 4221) detected in 862
plant species. The PCA is functionally equivalent to a PCA containing all 21 chemical properties (see Main Text), with axes
separating leaf metabolites based on size (PC1), aromaticity (PC2), and a-polarity (PC3; see Fig. 1A, lower panel). Colors
separate metabolite families as defined by marginal boxplot fills/labels (center: median; box: 25% to 75% quantiles; whiskers:
1.5 × IQR; points: outliers beyond whiskers).

203
204

<u>Five metabolic functional traits describe two axes of leaf metabolic specialisation</u>
We converted selected properties into plant-level metabolic functional traits to examine how the leaf
metabolome varies among tropical and temperate plant species. We first compared the volume and
"lumpiness" (i.e., aggregation around values) of multidimensional space occupied by metabolic traits
to four null models to test whether species converge on a restricted set of interrelated trait
combinations[7]. Observed hypervolumes of metabolic traits are 98.1% to 99.8% smaller and 71.0% to
94.7% lumpier than those of three null models assuming no trait covariance (Table S2; Table S3;
always $P < 0.001$ in all cases). However, observed hypervolumes are larger (tropical: $P = 0.002$;

temperate: $P = 0.047$) than that of the null model assuming trait covariance and selection against
extreme values, while also being similarly lumpy (tropical: $P = 0.662$; temperate: $P = 0.858$). By
comparison, observed hypervolumes of eight classical traits are smaller than null models assuming no
trait covariance (Table S2; $P < 0.001$ in all cases), either the same size as (tropical species; $P =$
0.154) or smaller than (temperate species; $P = 0.018$) that of the null model assuming selection
against extreme values, and lumpier than all null models (Table S3; $P < 0.001$ in all cases).
Collectively, these findings yield three insights. First, metabolic functional traits are interrelated, so
plants are constrained to a limited number of viable metabolic trait combinations. Second, while
extreme metabolic trait values may be selected against in temperate species, this is not so for tropical
species. Expressing extreme leaf metabolome chemistry is thus a viable life-history strategy in tropical
environments, potentially due to more intense biotic interactions selecting for a diversity of defensive
leaf metabolites and thus divergence of chemical strategies[52]. Third, metabolic traits are less
aggregated around non-extreme values than classical traits. As such, although plants converge
towards one of several viable classical trait strategies (e.g., woody, non-woody[7]), they can possess a
wider array of leaf metabolome strategies.
228
229

232   **Figure 3 | Metabolic traits describe two axes of leaf metabolic specialization among plant species.** Biplots of PC1 and
233   PC2 scores for PCAs describing variation in five metabolic functional traits (arrows) between (A) tropical (yellow; N = 457)
234   and (B) temperate (blue; N = 339) plant species. Axes separate species based on the mean aromaticity *versus* polarity
235   (inverse X log P) and carbon bond saturation (PC1) and mean size/complexity (PC2) of all annotated leaf metabolites
236   present (Methods).
237

238
We performed PCAs on metabolic traits to characterize major axes of leaf metabolic specialization
among species. Metabolic trait variation is consistent among tropical and temperate species, with 92%
variation being explained by the first two PCs (Fig. 3). The first axis (tropical: 52%; temperate 49%)
separates species that produce more unsaturated, aromatic, nonpolar leaf metabolites from those that
produce more saturated, polar, leaf metabolites. This axis likely describes a spectrum of leaf chemical
defense strategies[8]. Unsaturated aromatic metabolites, such as alkaloids, coumarins, and flavonoids
(Fig. 2), are reactive and serve as toxins or antioxidants in response to stress. For instance,
conjugated bond structures can interfere with protein function by binding covalently to sidechains[53],
generate or quench oxidative stress[54], or absorb damaging wavelengths of light[55]. By contrast,

saturated non-aromatic metabolites can play different roles in leaf defense, both directly as toxins[56]
and indirectly as signaling molecules[57]. An alternative explanation is that the first axis reflects an
investment (or not) into defense metabolites[8], which could arise due to the high cost of synthesizing
aromatic metabolites[17]. However, all plants must defend their leaves against stress, and we find that
plants can maximize fitness at either end of this axis (Fig. 3). It is thus more plausible that the first axis
represents a leaf chemical defense spectrum.
254
The second PCA axis (tropical: 40%; temperate: 43%) discriminates species that produce larger
metabolites with many H-bond acceptors from those that produce smaller metabolites with fewer H-
bond acceptors. This axis likely relates to leaf longevity, since large metabolites, such as lignans,
peptides, and waxes, form the basis of long-lived physical[58,59] and storage[60] structures. Many of these
structures also depend on strong intermolecular forces to form stable macromolecules or
complexes[61,62], which are enhanced by H-bond donors and acceptors[42]. As for the first axis, plants can
maximize fitness at either end of the second axis, with species at the positive end of the axis investing
in metabolites indicative of a longer leaf lifespan. As such, the second axis is probably a manifestation
of the fast-slow continuum[63] at the leaf level. Alternatively, the second axis may reflect an investment
(or not) into structurally complex metabolites for physical defense, such as lignans or waxes[58,59].
However, such metabolites act by creating rigid structures that lengthen leaf lifespan anyway[64], so
physical protection is one component of leaf longevity. Taken together, metabolic trait PCAs reveal
two major axes of leaf metabolic specialization. These axes relate to metabolite bond saturation and
polarity (i.e., a leaf chemical defense spectrum) and metabolite size and propensity for intermolecular
interactions (i.e., leaf longevity), with many combinations yielding successful life-history strategies.
270
<u>Axes of leaf metabolic specialisation reveal hidden dimensions of plant form and function</u>
The apparent consistency of metabolic trait variation with classical life-history theory[5–8,17,63,64] raises
the possibility that metabolic traits describe aspects of plant form and function already captured by
classical traits. We thus performed a PCA on metabolic traits plus the same widely used functional
traits used above to test whether axes of leaf metabolic specialization are colinear with, or orthogonal
to, trait-based life-history strategies. Considered alone, classical trait variation supports life-history
theory[7], with the first two PCs (tropical: 49%; temperate: 51%) describing the leaf economics spectrum
(SLA, leaf nitrogen, leaf phosphorus)[6] and variation indicative of plant longevity (plant height, seed
mass)[63] and hardiness (stem density, leaf carbon)[5] (Extended Data Fig. 1). However, when combining
classical and metabolic traits in the same PCA, the first four PCs (tropical: 66%; temperate: 67%) are
equivalent to axes described by individual PCAs (Fig. 4A, lower marginal matrices). The first and third
axes describe the first two axes of leaf metabolic specialization (Fig. 3), whereas the second and
fourth axes describe the first two axes of classical trait specialization (Extended Data Fig. 1). Pairwise
correlations among all traits further show that metabolic and classical traits do not covary (Fig. 4A,B),
but separate into six clusters describing unique aspects of either metabolic or classical trait variation.
As such, major axes of leaf metabolic specialization vary orthogonally to major axes of classical trait
variation (Fig. S4) and are at least equally relevant for describing variation in plant form and function
among tropical and temperate plant species.

289
290
Figure 4 | Metabolic and classical traits describe unique dimensions of plant form and function. Correlation
coefficients for pairwise Pearson correlations among metabolic (Fig. 3) and classical functional traits (Extended Data Fig. 1)
in (A) tropical (N = 457) and (B) temperate (N = 339) species. Traits are ordered based on correlation similarity (hierarchical
clustering; Ward), with colors separating distinct clusters (maximal silhouette width: K = 6) and horizontal lines between
matrices showing correspondence between tropical and temperate traits. Lower panel shows correlations between traits and
scores from the first four axes of a PCA of the same traits (tropical: 66%; temperate: 67%; LES = leaf economics spectrum).
Circle sizes indicate coefficient strengths, while fills discriminate between positive (black) and negative (white) correlations.
298
299
Leaf metabolism underpins energy production *via* photosynthesis[65], but in doing so yields tissue that is
sensitive to light[66] and temperature[67] and is of high nutritional value to herbivores[64]. Plants are thus
under selective pressure to maximize photosynthetic rates while also protecting leaf tissue against
damage. While this trade-off is consistent with life-history trade-offs derived from classical traits[5–7], it
does not necessarily follow that individuals apply the same strategy in every organ to maintain fitness.
Indeed, we observe that species with "productive" trait values (e.g., high SLA, high leaf nitrogen) can
produce leaves containing many complex aromatic metabolites, species possessing "hardy" trait
values (e.g., high stem density, high leaf carbon) can produce leaves containing many simple polar
metabolites, and species possessing trait values typical of a fast pace of life (e.g., low stature, small
seeds) can produce leaves with many large stable or structural metabolites. Our findings thus question
classical trait[6] and plant defense[8] theory that predicts relationships between the leaf chemical
phenotype, plant productivity, and pace of life. In short, metabolic functional traits describe unique
dimensions of plant life-history variation that are complementary to, and independent from, those
captured by classical functional traits.
314
315
316

<u>Concluding remarks</u>
In summary, we show that leaf metabolite chemistry is highly structured and can be described by five
chemical properties. Using chemical properties as metabolic functional traits reveals that plants vary
on two axes of leaf metabolic specialization – a leaf chemical defense spectrum (i.e., bond saturation,
polarity) and leaf longevity (i.e., size, propensity for intermolecular interactions). Axes of metabolic
specialization are similar in tropical and temperate species, with many combinations of trait values
yielding successful life-history strategies. Axes are also consistent with expectations from life-history
theory that plants invest strategically in tissue defense[6,8,17] and between a fast *versus* slow pace of
life[7,63]. Nevertheless, we find that metabolic trait variation is orthogonal to, not colinear with, classical
trait variation. Metabolic functional traits thus reveal dimensions of plant life-history variation missed by
existing functional traits. It is important to note that metabolomics data were acquired differently for
tropical and temperate species and were matched to open functional trait databases. Nevertheless,
we find consistent variation in the leaf metabolomes of 457 tropical and 339 temperate species that
possess distinct geographic distributions and evolutionary histories (Extended Data Fig. 1A,B), as well
as that encompass most documented functional trait variation (Extended Data Fig. 1G, red ribbons).
We thus propose that distilling the leaf metabolome into metabolic functional traits captures
macroecological patterns of metabolic variation widely across the plant kingdom, enhances the
explanatory power of the functional trait concept, and offers a new set of tools for the discovery of
species or genotypes with trait combinations adapted to societal needs.

Methods

Sample collection & preparation. We used two existing datasets describing the leaf metabolomes of tropical and temperate plant species (as the unit of observation). The tropical dataset originated from a random subsample of tropical leaves from the Pierre Fabre Sample Library[68], which is an archive of ~17000 plant samples from *in situ* communities around the globe that was originally collected for drug discovery and is a registered collection of the European Union (registration code: 03-FR-2020)[69]. Samples were oven-dried (55 ºC for 3 d) and extracted using overnight maceration (8 g sample; 80 mL ethyl acetate). Extracts were dried once (Genevac, SP Industries, Warminster, PA, USA) and eluted (30 mg extract plus 200 mg silica; 1g Silica SPE cartridges; 6 mL DCM plus 85:15 DCM:MeOH), then dried again and resuspended in DMSO at a final concentration of 5 mg mL^{-1}. The temperate dataset originated from a study of landscape-scale ecological variation in phytochemical diversity[37], wherein leaves were sampled from *in situ* grassland communities across Switzerland. Briefly, leaves were dried (40 ºC for 5 d), milled (Retsch TissueLyser; Qiagen, Hilden, Germany) and extracted (20 mg tissue; 0.5 mL 80:19.5:0.5 MeOH:H_2O: H_2CO_2). Extracts were homogenized (glass beads; 30 Hz for 3 min) and centrifuged (14000 RPM for 3 min), and the supernatant was isolated for measurement.

Metabolomics measurements. Untargeted metabolomics was performed on all samples using Ultra High-Performance Liquid Chromatography Mass Spectrometry (UHPLC-MS). Tropical extracts were analyzed using a Waters Acquity UHPLC system coupled to a Q-Exactive Focus MS (Thermo Scientific, Bremen, Germany) with a heated electrospray ionization (HESI-II) source in positive mode. Extracts (2 μL) were separated on an Acquity C18 column (50 mm × 2.1 mm, 1.7 μm; Waters, Milford, MA, USA) at a flow rate of 600 μl min^{-1} using a binary solvent system (A: H_2O plus 0.1% H_2CO_2; B: acetonitrile plus 0.1% H_2CO_2) and a linear gradient elution (5%-100% B, 7 min) plus isocratic elution (100% B, 1 min). MS data were acquired in data-dependent acquisition mode, in which MS/MS scans were performed on the three most intense ions detected during repeated full MS scans (full scans = 35000 FWHM at m/z 200; MS/MS scans = 17000 FWHM; MS/MS isolation window width = 1 Da; normalized collision energy = 15, 30 and 45 units). The MS was calibrated using a quality control (QC) mix containing caffeine, methionine-arginine-phenylalanine-alanine-acetate (MRFA), sodium dodecyl sulphate, sodium taurocholate and Ultramark 1621 dissolved in acetonitrile/methanol/H_2O plus 1% H_2CO_2. Temperate extracts were analyzed using a Waters Acquity UHPLC system coupled to a Synapt G2 MS (Waters, Milford, MA, USA) with a heated electrospray ionization (HESI-II) source in positive mode. Extracts (2.5 μL) were separated on an Acquity C18 column (50 mm × 2.1 mm, 1.7 μm; Waters, Milford, MA, USA) at a flow rate of 600 μl min^{-1} using a binary solvent system (A: H_2O plus 0.05% H_2CO_2; B: acetonitrile plus 0.05% H_2CO_2) and a linear gradient elution (2%-100% B, 6 min). MS data were acquired in data-independent acquisition mode, in which all precursor ions across the full mass range (85-1200 Da) were fragmented to yield MS/MS spectra. The MS was calibrated using a QC mix derived from all sample extracts.

Metabolomics data pre-processing. Tropical MS data were treated using MZMine (version 2.53)[70]. Peak detection was performed using the centroid mass detector (MS noise = 1 × 10^4; MS/MS noise = 0) and the ADAP chromatogram builder (min. scan group size = 5, min. group intensity = 1 × 10^4; min. highest intensity = 5 × 10^5; m/z tolerance = 12 ppm)[71]. Peaks were deconvoluted using the wavelets algorithm in ADAP with an intensity window SN (S/N threshold = 10; min. feature height = 5 × 10^5;

coefficient/area threshold = 130; peak duration range = 0-5 min; retention time (RT) wavelet range =
0.01-0.03 min). Isotopes were detected using the isotopes peak grouper (m/z tolerance = 12 ppm;
absolute RT tolerance = 0.01 min absolute; max. charge = 2) and, where present, the most intense
isotope was chosen. Peaks were filtered to retain features possessing an MS/MS scan and a RT of
between 0.5 and 8 min, following which they were aligned using the join aligner method (m/z tolerance
= 40 ppm; absolute RT tolerance = 0.2 min; m/z weight = 2; RT weight = 1). Temperate MS data were
treated using MS-DIAL[72]. Peaks were detected with a minimum height of 300 and data were collected
with an MS1 tolerance of 0.05 Da and an MS2 tolerance of 0.01 Da from 100–1200 Da and 0.5 to 12
minutes. Peaks were deconvoluted with a sigma window value of 0.5 and an MS/MS abundance cut-
off of 50. Finally, raw features were aligned using the QC mix with a retention time tolerance of 0.1
minute. Following this, we separately clustered tropical and temperate metabolic features into sets of
spectrally similar "consensus" features using molecular networks constructed on the Global Natural
Products Social (GNPS; precursor and MS/MS fragment ion mass tolerances = 0.02 Da)[73,74]. We
retained edges with a cosine score of more than 0.7 and with more than six matched peaks, permitted
edges that connected two nodes only if each node appeared in the neighbour's "top 10" most similar
node list and, where necessary, limited the total size of a cluster to the 100 highest scoring edges.
Finally, we converted peak height/area data to presence-absence data by summing peak
heights/areas at the consensus metabolic feature level and expressing them as a one (present) or
zero (absent) (tropical N = 7343, temperate N = 6682).
399
Metabolite annotation. We annotated all metabolic features with chemical information by *in silico*
spectral matching and taxonomically-informed scoring using a theoretically fragmented version of the
Dictionary of Natural Products[75] following[76,77], retaining matches with a minimum score of 0.7 and at
least six matched peaks. We assigned consensus metabolic features with a single consensus
chemical classification, which we derived by taking the most common chemical classification across
individual features contributing to a consensus feature. We filtered data to retain those features that
were annotated with a metabolite identifier (i.e., SMILES notation; tropical N = 3356 (46%), temperate
N = 2227 (33%)). We then derived the chemistry of annotated metabolites by using SMILES identifiers
to query the Chemistry Development Kit[27] with the package "rcdk"[78]. The CDK comprises 51 chemical
descriptors, but we focused on 21 chemical properties that describe metabolite constitution, geometry,
and topology (Table S1). We also derived values for annotated metabolites from the first three
dimensions of the ChemGPS-NP tool[28], which is an existing framework for classifying natural products
that separates molecules based on size, then aromaticity, then lipophilicity[31].
413
Classical functional traits. All further data processing, analysis and visualization were undertaken in
R[79]. For core functionality, we used the R packages "cowplot"[80], "data.table"[81], "drake"[82], "furrr"[83],
"future"[84], "parallel"[79], "psych"[85] and "tidyverse"[86] (see below for packages pertaining to specific steps).
Plant species names were cleaned against the Catalogue of Life and Encyclopedia of Life using the
global name resolver in the package "taxize"[87], following which we added a taxonomic hierarchy using
"taxonlookup"[88]. We removed 14 tropical (ferns, cycads, conifers) and nine temperate (ferns, trees)
species that were phylogenetically distinct from others within their respective datasets to prevent a
small number of divergent species from overwhelming downstream analyses (Fig. S1). We obtained
data for classical functional traits with proven relevance to plant form and function from the open-
access version of the TRY database (version 5)[89] – namely plant height (m), seed mass (mg), stem

specific density (mg mm^{-3}), leaf area (mm^2), specific leaf area (mm^2 mg^{-1}) and leaf carbon, nitrogen
and phosphorus concentrations (mg g^{-1})[6,7,90]. After removing clear outliers (reported error risk ≥ 4;
values falling outside of documented limits), we calculated species-wise means and gap-filled missing
trait values for all species present in the dataset (N = 46905 species) using a two-step process. First,
we performed a Bayesian hierarchical probabilistic matrix factorization imputation (package
"BHPMF"[91]), which is a taxonomically-constrained gap-filling approach that has been regularly applied
to the TRY database[7,24]. The imputation was repeated 90 times, each time starting with a different
combination of parameters (per-fold samples = 900-1000; cross-validation steps = 10-20; burn-in
steps = 10% data length) and filtering out extreme (>1.5 times the maximum observed value for a trait)
or highly uncertain (coefficient of variation >1) gap-filled values. We then calculated means of
remaining gap-filled values across all iterations and used these to replace missing cases in original
trait data. Second, we performed five iterations of multivariate imputation by chained equations
(package "mice"[92]) on the partially gap-filled dataset and replaced remaining missing cases with the
mean values from all iterations. Following this, we filtered fully gap-filled datasets for selected tropical
and temperate species (see Table S4 for details about missing data), using genus-level mean trait
values where species-level data were not available (215 tropical species, 5 temperate species). It is
important to note that trait data coverage differed substantially between tropical and temperate
species (Table S4). While this undoubtedly reduced the strength of observed trait variation in tropical
relative to temperate data, it is unlikely that the reduced resolution of the tropical data changed the
nature of functional trait variation, since artificially reducing the resolution of trait datasets to genus
level did not change resulting PC axes (Fig. S2). Finally, we accounted for non-normal distributions in
plant height, seed mass and leaf area data using log$_{10}$-transformations and summarized trait variation
among tropical and temperate species separately using scores from the first two components of
varimax-rotated PCAs performed on all traits (Extended Data Fig. 1).

Species distributions. We matched species names to Global Biodiversity Information Facility (GBIF)
IDs using "rgbif"[93] and gathered all available species occurrences from the GBIF website
(www.gbif.org; tropical N = 336632, temperate N = 1847095). Working with each species individually,
we filtered occurrences to include human observations and living or preserved specimens recorded
from the year 1945 to present day and accompanied by either no individual counts or counts with a
reasonable value (>100)[94]. We used the packages "CoordinateCleaner"[95] to discard records falling in
the sea, outside the stated country or within 10 km of capital cities, GBIF headquarters or known
biodiversity institutions, records lying more than five times the interquartile range of the minimum
distances to the nearest neighbor and records containing known issues with longitude/latitude
information (i.e., zeros, rounding errors or conversion errors from other coordinate systems)[94,95].
Following this, we manually removed a further 20 tropical and four temperate records representing
single occurrences within a continent, leaving a total of 239223 and 1251500 cleaned occurrences for
tropical and temperate species, respectively. We plotted cleaned occurrences to visualize the
geographical extents of tropical and temperate plant species (Extended Data Fig. 1AB), although it is
important to emphasize that samples for metabolomics analyses were taken from within natural (not
ornamental) distributions.

Leaf metabolite chemistry. We collated all unique annotated metabolites detected in any tropical or
temperate species into a single dataset capturing overall variation in the chemical properties of the

leaf metabolome (N = 4292). We tested for interrelatedness between metabolite chemical properties
using pairwise Pearson correlations coupled with hierarchical clustering (Ward; Fig. 1, upper panel),
and estimated the number of discrete clusters of chemical properties using the maximal average
silhouette width[96] (Fig. 1A, dendrogram). We described major dimensions of leaf chemical variation
using a varimax-rotated PCA performed on a subset of five representative chemical properties (one
from each cluster: C-hybridization ratio, H-bond acceptor count, molecular weight, polarity (X log P),
aromatic atom count). The first three PC axes described 97% variation, which we anchored back to
individual chemical properties using pairwise Pearson correlations (Fig. 1A, lower panel). We tested
whether the subset PCA was functionally equivalent to a full PCA performed on all 21 properties using
two approaches. First, we performed pairwise Pearson correlations between subset *versus* full PC
scores to determine whether the two PCAs separated metabolic features similarly on the same
dimensions of chemical variation (Fig. 1B, left panel; see Main Text for statistics). Second, we used a
Procrustes Rotation test (package "vegan"[97]; 999 permutations) to determine whether maximally-
rotated configurations of full *versus* subset PCAs were significantly (i.e., non-randomly) associated
(m_{12} = 0.04, r^2 = 0.98, P < 0.001). We also confirmed that the similarity among subset and full PCAs
was not sensitive to which chemical properties were selected from each cluster. This was achieved by
repeating Procrustes Rotation tests and PC score correlations for PCAs performed on every
combination of five chemical properties that covered all five clusters and plotting distributions of
resulting coefficients (Fig. S4). We performed pairwise Pearson correlations between subset PC
scores and values from the first three dimensions of the ChemGPS-NP framework[28] to test for
correspondence between the subset PCA and existing descriptors of biologically relevant chemical
space (Fig. 1B, right panel; see Main Text for statistics). We visualized chemical variation among
metabolites using biplots of PC scores from the subset PCA, grouped by biochemical class (Fig. 2).
We also illustrated the range of chemical structures captured by the five selected chemical properties
by using the PubChem sketcher (https://pubchem.ncbi.nlm.nih.gov//edit3/index.html) to predict
chemical graphs of metabolites with the lowest and highest values for each property (Fig. 1C).
494
Plant-level metabolic functional traits. We selected C-hybridization ratio, H-bond acceptor count,
molecular weight, X log P, and aromatic atom count to convert into five plant-level metabolic functional
traits. For each species (i.e., unit of observation), we calculated the mean value of a chemical property
considering all the metabolites present. In doing so, we generated datasets akin those capturing
classical functional trait variation among tropical or temperate species, but instead quantifying
variation in the mean C-hybridization ratio, H-bond acceptor count, molecular weight, polarity, and
aromatic atom count of the annotated leaf metabolome among tropical or temperate species. We
considered tropical and temperate datasets as separate entities for data analysis of trait variation, for
four reasons. First, while *in silico* data treatment was identical for aligned MS data, sample collection
and LC-MS measurements were not (see above), making it challenging to combine data on species-
level metabolic variation without inflating differences between them. Second, unlike genomics and
proteomics analyses, where only one type of molecule is isolated, untargeted metabolomics aims to
simultaneously isolate thousands of molecules with a wide range of chemical properties[3,98]. Technical
decisions made during sample extraction and analysis discriminate for or against certain types of
metabolites, which filters the resulting view of the metabolome. There is still no single accepted
method for extracting samples or acquiring metabolomics data[99], so we chose here to use two
datasets generated using different approaches to confirm that findings are not sensitive to the
methods chosen. Third, classical functional trait and species occurrence data were sparser for tropical
than temperate species (see *Classical Functional Traits*, above), so we chose to keep datasets

separate rather than artificially reduce the resolution of the temperate dataset. Fourth, maintaining a
separation between the two datasets increased our interpretive power by providing the opportunity to
characterize coupling between the metabolome and functional traits that held true not only within, but
also among, tropical and temperate plant species.
518
Plant multidimensional space. Following[7], we explored constraints on the multidimensional space
occupied by metabolic and classical functional traits separately for tropical and temperate species
using an observed *versus* simulated hypervolume approach. Briefly, we calculated an *n*-dimensional
convex hull volume (package "geometry"[100]) representing the observed multidimensional trait space
occupied by species, where *n* is the number of metabolic (n = 5) or classical (n = 8) functional traits.
We compared the observed hypervolume to the same four simulated null hypervolumes used by ref. [7],
which tested the following null hypotheses: (i) traits vary independently and have uniform distributions
(i.e., each trait is a unique axis and no selection against extreme trait values); (ii) traits vary
independently but have normal distributions (i.e., each trait is a unique axis and selection against
extreme trait values); (iii) traits vary independently but distributions are as observed (i.e., each trait is a
unique axis but no assumptions about selection on values); and (iv) traits covary as but have normal
distributions (i.e., no assumptions about trait interdependence but selection against extreme values).
We simulated null hypervolumes 999 times and used permutation tests (package "ade4"[100]) to test for
differences between the overall size of observed *versus* simulated hypervolumes, and presented
percent changes between observed and mean simulated hypervolume sizes (where positive =
observed > simulated; Table S2). We also calculated the "lumpiness" (i.e., aggregation around certain
values) of species in multidimensional space by splitting *n*-dimensional space into 10 bins (i.e., 10^n
cells), counting the number of species present in each cell, and calculating the minimum number of
cells required to capture 10% of species[7]. Again, we tested for differences between observed and
simulated null hypervolumes using permutation tests and presented percent changes between
observed and mean simulated hypervolumes (Table S3).
540
Axes of leaf metabolic specialization. We characterized major axes of leaf metabolic specialization
separately among tropical or temperate species using varimax-rotated PCAs performed on metabolic
functional traits. The first two PCs explained 92% variation for both tropical and temperate species,
respectively, which we displayed using biplots of PC scores (Fig. 3). We explored whether axes of leaf
metabolic specialization (Fig. 3) were orthogonal to, or colinear with, axes of classical functional trait
variation (Fig. 4) by doing varimax-rotated PCAs containing metabolic plus classical functional traits,
again separately among tropical and temperate species. Four PCs were needed to explain 66% and
67% variation among tropical and temperate species, respectively, which we displayed in two ways.
First, we created a biplot matrix of all pairwise combinations of the first four axes (Fig. S3) to visualize
the lack of overlap between metabolic and classical functional traits. Second, we performed Pearson
correlations between PC scores and individual functional traits (Fig. 4, lower panels). Finally, we
examined interdependence within and among metabolic and classical functional traits by performing
pairwise Pearson correlations (Fig. 4, upper panels), clustering correlation coefficients (hierarchical
clustering; Ward), and identifying discrete clusters of traits using the maximal silhouette width[96] (Fig. 4,
dendrograms). Dendrograms are organized based on the outcome of a tanglegram that tested for
correspondence between the clustering of tropical and temperate traits (Fig. 4, central connectors).
557

558   **Data and materials availability.** Raw tropical metabolomics data are available under the DOI

559   10.25345/C59J97[36]. Raw temperate species metabolomics data are available under the DOI

560   10.6084/m9.figshare.13032740[37]. Flat data and R code is available under the DOI

561   10.17605/OSF.IO/YZR4C.

References

1. Wink, M. Evolution of secondary metabolites from an ecological and molecular phylogenetic perspective. *Phytochemistry* **64**, 3–19 (2003).

2. Schuman, M. C., van Dam, N. M., Beran, F. & Harpole, W. S. How does plant chemical diversity contribute to biodiversity at higher trophic levels? *Curr. Opin. Insect Sci.* **14**, 46–55 (2016).

3. Walker, T. W. N. *et al.* Functional Traits 2.0: The power of the metabolome for ecology. *Journal of Ecology* **110**, 4–20 (2022).

4. Violle, C. *et al.* Let the concept of trait be functional! *Oikos* **116**, 882–892 (2007).

5. Grime, J. P. *et al.* Integrated screening validates primary axes of specialisation in plants. *Oikos* **79**, 259 (1997).

6. Wright, I. J. *et al.* The worldwide leaf economics spectrum. *Nature* **428**, 821–827 (2004).

7. Díaz, S. *et al.* The global spectrum of plant form and function. *Nature* **529**, 167–171 (2016).

8. Feeny, P. Plant apparency and chemical defense. in *Biochemical interaction between plants and insects* (eds. Wallace, J. W. & Mansell, R. L.) 1–40 (Springer US, 1976). doi:10.1007/978-1-4684-2646-5_1.

9. Wang, S., Alseekh, S., Fernie, A. R. & Luo, J. The structure and function of major plant metabolite modifications. *Mol. Plant* **12**, 899–919 (2019).

10. Faccio, G. Plant complexity and cosmetic innovation. *iScience* **23**, 101358 (2020).

11. Klee, H. J. & Tieman, D. M. The genetics of fruit flavour preferences. *Nat. Rev. Genet.* **19**, 347–356 (2018).

12. Schmidt, B. M., Ribnicky, D. M., Lipsky, P. E. & Raskin, I. Revisiting the ancient concept of botanical therapeutics. *Nat. Chem. Biol.* **3**, 360–366 (2007).

13. Sardans, J. *et al.* Ecometabolomics for a better understanding of plant responses and acclimation to abiotic factors linked to global change. *Metabolites* **10**, (2020).

14. Sardans, J. *et al.* Ecometabolomics of plant–herbivore and plant–fungi interactions: a synthesis study. *Ecosphere* **12**, (2021).

15. Zanne, A. E. *et al.* Three keys to the radiation of angiosperms into freezing environments. *Nature* **506**, 89–92 (2014).

16. Hutchinson, G. E. *The Ecological Theater and the Evolutionary Play*. (Yale University Press, 1965).

17. Coley, P. D., Bryant, J. P. & Chapin, F. S. Resource availability and plant antiherbivore defense. *Science* **230**, 895–899 (1985).

18. Capdevila, P., Walker, T. W. N., Schrodt, F., Rodriguez Caro, R. C. & Salguero-Gomez, R. Global patterns of plant form and function are strongly determined by evolutionary relationships. *BioRxiv* (2023) doi:10.1101/2023.01.13.523963.

19. Sedio, B. E., Parker, J. D., McMahon, S. M. & Wright, S. J. Comparative foliar metabolomics of a tropical and a temperate forest community. *Ecology* **99**, 2647–2653 (2018).

20. Forrister, D. L., Endara, M.-J., Younkin, G. C., Coley, P. D. & Kursar, T. A. Herbivores as drivers of negative density dependence in tropical forest saplings. *Science* **363**, 1213–1216 (2019).

21. Gargallo-Garriga, A. *et al.* Different "metabolomic niches" of the highly diverse tree species of the French Guiana rainforests. *Sci. Rep.* **10**, 6937 (2020).

22. Scherling, C., Roscher, C., Giavalisco, P., Schulze, E.-D. & Weckwerth, W. Metabolomics unravel contrasting effects of biodiversity on the performance of individual plant species. *PLoS ONE* **5**, e12569 (2010).

23. Adler, P. B. *et al.* Functional traits explain variation in plant life history strategies. *Proc Natl Acad Sci USA* **111**, 740–745 (2014).

24. Bruelheide, H. *et al.* Global trait-environment relationships of plant communities. *Nat. Ecol. Evol.* **2**, 1906–1917 (2018).

25. De Deyn, G. B., Cornelissen, J. H. C. & Bardgett, R. D. Plant functional traits and soil carbon sequestration in contrasting biomes. *Ecol. Lett.* **11**, 516–531 (2008).

26. Lavorel, S. *et al.* A novel framework for linking functional diversity of plants with other trophic levels for the quantification of ecosystem services. *J. Veg. Sci.* **24**, 942–948 (2013).

27. Willighagen, E. L. *et al.* The Chemistry Development Kit (CDK) v2.0: atom typing, depiction, molecular formulas, and substructure searching. *J. Cheminform.* **9**, 33 (2017).

28. Rosén, J. *et al.* ChemGPS-NP(Web): chemical space navigation online. *J. Comput. Aided Mol. Des.* **23**, 253–259 (2009).

29. Lipinski, C. A., Lombardo, F., Dominy, B. W. & Feeney, P. J. Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. *Adv. Drug Deliv. Rev.* **46**, 3–26 (2001).

30. Lovering, F., Bikker, J. & Humblet, C. Escape from flatland: increasing saturation as an approach to improving clinical success. *J. Med. Chem.* **52**, 6752–6756 (2009).

31. Larsson, J., Gottfries, J., Muresan, S. & Backlund, A. ChemGPS-NP: tuned for navigation in biologically relevant
chemical space. *J. Nat. Prod.* **70**, 789–794 (2007).
32. Funk, J. L. *et al.* Revisiting the Holy Grail: using plant functional traits to understand ecological processes. *Biol. Rev.
Camb. Philos. Soc.* **92**, 1156–1173 (2017).
33. Firn, J. *et al.* Leaf nutrients, not specific leaf area, are consistent indicators of elevated nutrient inputs. *Nat. Ecol. Evol.*
3, 400–406 (2019).
34. van der Plas, F. *et al.* Plant traits alone are poor predictors of ecosystem properties and long-term ecosystem
functioning. *Nat. Ecol. Evol.* **4**, 1602–1611 (2020).
35. Walker, T. W. N. *et al.* Plastic and genetic responses of a common sedge to warming have contrasting effects on
carbon cycle processes. *Ecol. Lett.* **22**, 159–169 (2019).
36. Allard, P.-M. *et al.* Open and reusable annotated mass spectrometry dataset of a chemodiverse collection of 1,600
plant extracts. *Gigascience* **12**, (2022).
37. Defossez, E. *et al.* Spatial and evolutionary predictability of phytochemical diversity. *Proc Natl Acad Sci USA* **118**,
(2021).
38. Wiener, H. Structural determination of paraffin boiling points. *J. Am. Chem. Soc.* **69**, 17–20 (1947).
39. Sharma, V., Goswami, R. & Madan, A. K. Eccentric connectivity index: A novel highly discriminating topological
descriptor for structure–property and structure–activity studies. *J. Chem. Inf. Comput. Sci.* **37**, 273–282 (1997).
40. Yang, Y., Chen, H., Nilsson, I., Muresan, S. & Engkvist, O. Investigation of the relationship between topology and
selectivity for druglike molecules. *J. Med. Chem.* **53**, 7709–7714 (2010).
41. Kuhn, B., Mohr, P. & Stahl, M. Intramolecular hydrogen bonding in medicinal chemistry. *J. Med. Chem.* **53**, 2601–
2611 (2010).
42. Zeegers-Huyskens, Th. & Huyskens, P. Intermolecular Forces. in *Intermolecular Forces* (eds. Huyskens, P. L., Luck,
W. A. P. & Zeegers-Huyskens, T.) 1–30 (Springer Berlin Heidelberg, 1991). doi:10.1007/978-3-642-76260-4_1.
43. Dervan, P. B. Molecular recognition of DNA by small molecules. *Bioorg. Med. Chem.* **9**, 2215–2235 (2001).
44. Padroni, G., Patwardhan, N. N., Schapira, M. & Hargrove, A. E. Systematic analysis of the interactions driving small
molecule-RNA recognition. *RSC Med. Chem.* **11**, 802–813 (2020).
45. Gao, M. & Skolnick, J. A comprehensive survey of small-molecule binding pockets in proteins. *PLoS Comput. Biol.* **9**,
e1003302 (2013).
46. Wang, R., Fu, Y. & Lai, L. A New Atom-Additive Method for Calculating Partition Coefficients. *J. Chem. Inf. Comput.
Sci.* **37**, 615–621 (1997).
47. Ghose, A. K. & Crippen, G. M. Atomic Physicochemical Parameters for Three-Dimensional Structure-Directed
Quantitative Structure-Activity Relationships I. Partition Coefficients as a Measure of Hydrophobicity. *J. Comput.
Chem.* **7**, 565–577 (1986).
48. Clemons, P. A. *et al.* Small molecules of different origins have distinct distributions of structural complexity that
correlate with protein-binding profiles. *Proc Natl Acad Sci USA* **107**, 18787–18792 (2010).
49. Israelachvili, J. N. The nature of van der waals forces. *Contemp. Phys.* **15**, 159–178 (1974).
50. Van Vranken, D. & Weiss, G. A. *Introduction to Bioorganic Chemistry and Chemical Biology*. 504 (Garland Science,
2012).
51. Pichersky, E. & Raguso, R. A. Why do plants produce so many terpenoid compounds? *New Phytol.* **220**, 692–702
(2018).
52. Forrister, D. L. *et al.* Diversity and Divergence: Evolution of secondary metabolism in the tropical tree genus Inga.
New Phytol. (2022) doi:10.1111/nph.18554.
53. LoPachin, R. M., Barber, D. S. & Gavin, T. Molecular mechanisms of the conjugated alpha,beta-unsaturated carbonyl
derivatives: relevance to neurotoxicity and neurodegenerative diseases. *Toxicol. Sci.* **104**, 235–249 (2008).
54. Widomska, J. & Subczynski, W. K. Mechanisms enhancing the protective functions of macular xanthophylls in the
retina during oxidative stress. *Exp. Eye Res.* **178**, 238–246 (2019).
55. Tran, M. H., Phan, D.-P. & Lee, E. Y. Review on lignin modifications toward natural UV protection ingredient for lignin-
based sunscreens. *Green Chem.* **23**, 4633–4646 (2021).
56. Tan, N.-H. & Zhou, J. Plant cyclopeptides. *Chem. Rev.* **106**, 840–895 (2006).
57. Lim, G.-H., Singhal, R., Kachroo, A. & Kachroo, P. Fatty Acid- and Lipid-Mediated Signaling in Plant Defense. *Annu.
Rev. Phytopathol.* **55**, 505–536 (2017).
58. Yeats, T. H. & Rose, J. K. C. The formation and function of plant cuticles. *Plant Physiol.* **163**, 5–20 (2013).
59. Liu, Q., Luo, L. & Zheng, L. Lignins: biosynthesis and biological functions in plants. *Int. J. Mol. Sci.* **19**, (2018).

60.   Staswick, P. E. Storage proteins of vegetative plant tissues. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **45**, 303–322
(1994).
61.   Sandberg, T. O., Weinberger, C. & Smått, J.-H. Molecular Dynamics on Wood-Derived Lignans Analyzed by
Intermolecular Network Theory. *Molecules* **23**, (2018).
62.   Pollard, M., Beisson, F., Li, Y. & Ohlrogge, J. B. Building lipid barriers: biosynthesis of cutin and suberin. *Trends Plant*
Sci. **13**, 236–246 (2008).
63.   Salguero-Gómez, R. *et al.* Fast-slow continuum and reproductive strategies structure plant life-history variation
worldwide. *Proc Natl Acad Sci USA* **113**, 230–235 (2016).
64.   Coley, P. D. Herbivory and defenses: a temperate/tropic comparison. in *Plant-Animal Interactions: Evolutionary*
Ecology in Tropical and Temperate Regions (eds. Price, P. W., Lewison, T. M., Fernandes, G. W. & Benson, W. W.)
54–69 (John Wiley and Sons, 1991).
65.   Sulpice, R. & McKeown, P. C. Moving toward a comprehensive map of central plant metabolism. *Annu. Rev. Plant*
Biol. **66**, 187–210 (2015).
66.   Powles, S. B. Photoinhibition of photosynthesis induced by visible light. *Annu. Rev. Plant Physiol.* **35**, 15–44 (1984).
67.   Berry, J. & Bjorkman, O. Photosynthetic response and adaptation to temperature in higher plants. *Annu. Rev. Plant*
Physiol. **31**, 491–543 (1980).
68.   David, B. & Aussiel, F. High throughput screening of vegetal natural substances. in *Handbook of Chemical and*
Biological Plant Analytical Methods (ed. Hostettmann, K.) 1176 (Wiley, 2014).
69.   David, B. & Ausseil, F. High-Throughput Screening of Plant Chemodiversity. in *Encyclopedia of analytical chemistry:*
applications, theory and instrumentation (ed. Meyers, R. A.) 1–24 (John Wiley & Sons, Ltd, 2006).
doi:10.1002/9780470027318.a9944.
70.   Pluskal, T., Castillo, S., Villar-Briones, A. & Oresic, M. MZmine 2: modular framework for processing, visualizing, and
analyzing mass spectrometry-based molecular profile data. *BMC Bioinformatics* **11**, 395 (2010).
71.   Myers, O. D., Sumner, S. J., Li, S., Barnes, S. & Du, X. One Step Forward for Reducing False Positive and False
Negative Compound Identifications from Mass Spectrometry Metabolomics Data: New Algorithms for Constructing
Extracted Ion Chromatograms and Detecting Chromatographic Peaks. *Anal. Chem.* **89**, 8696–8703 (2017).
72.   Tsugawa, H. *et al.* MS-DIAL: data-independent MS/MS deconvolution for comprehensive metabolome analysis. *Nat.*
Methods **12**, 523–526 (2015).
73.   Frank, A. M. *et al.* Clustering millions of tandem mass spectra. *J. Proteome Res.* **7**, 113–122 (2008).
74.   Wang, M. *et al.* Sharing and community curation of mass spectrometry data with Global Natural Products Social
Molecular Networking. *Nat. Biotechnol.* **34**, 828–837 (2016).
75.   Chassagne, F., Cabanac, G., Hubert, G., David, B. & Marti, G. The landscape of natural product diversity and their
pharmacological relevance from a focus on the Dictionary of Natural Products®. *Phytochem. Rev.* 1–22 (2019)
doi:10.1007/s11101-019-09606-2.
76.   Allard, P.-M. *et al.* Integration of Molecular Networking and In-Silico MS/MS Fragmentation for Natural Products
Dereplication. *Anal. Chem.* **88**, 3317–3323 (2016).
77.   Rutz, A. *et al.* Taxonomically informed scoring enhances confidence in natural products annotation. *Front. Plant Sci.*
10, 1329 (2019).
78.   Guha, R. Chemical Informatics Functionality in *R. J. Stat. Softw.* **18**, (2007).
79.   R Core Team. *R: A language and environment for statistical computing.* (R Foundation for Statistical Computing,
2014).
80.   Wilke, C. O. *cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2."* (R Project, 2019).
81.   Dowle, M. & Srinivasan, A. *data.table: Extension of `data.frame`.* (R Project, 2021).
82.   Michael Landau, W. The drake R package: a pipeline toolkit for reproducibility and high-performance computing.
JOSS **3**, 550 (2018).
83.   Vaughan, D. & Dancho, M. *furrr: Apply Mapping Functions in Parallel using Futures.* (R Project, 2021).
84.   Bengtsson, H. A unifying framework for parallel and distributed processing in R using futures. *arXiv* (2021).
85.   Revelle, W. R. *psych: Procedures for Personality and Psychological Research.* (Northwestern University, 2019).
86.   Wickham, H. *et al.* Welcome to the tidyverse. *JOSS* **4**, 1686 (2019).
87.   Chamberlain, S. A. & Szöcs, E. taxize: taxonomic search and retrieval in R. [version 2; peer review: 3 approved].
F1000Res. **2**, 191 (2013).
88.   Cornwell, W., FitzJohn, R. & Pennell, M. *taxonlookup: A dynamically-updating versioned taxonomic resource for*
vascular plants. (R Project, 2016).

89.   Kattge, J. *et al.* TRY plant trait database - enhanced coverage and open access. *Glob. Chang. Biol.* **26**, 119–188
(2020).
90.   Grime, J. P. Vegetation classification by reference to strategies. *Nature* **250**, 26–31 (1974).
91.   Schrodt, F. *et al.* BHPMF - a hierarchical Bayesian approach to gap-filling and trait prediction for macroecology and
functional biogeography. *Global Ecology and Biogeography* **24**, 1510–1521 (2015).
92.   van Buuren, S. & Groothuis-Oudshoorn, K. mice : Multivariate Imputation by Chained Equations in *R. J. Stat. Softw.*
45, (2011).
93.   Chamberlain, S. *et al. rgbif: Interface to the Global Biodiversity Information Facility API.* (R Project, 2021).
94.   Maldonado, C. *et al.* Estimating species diversity and distribution in the era of Big Data: to what extent can we trust
public databases? *Glob. Ecol. Biogeogr.* **24**, 973–984 (2015).
95.   Zizka, A. *et al.* CoordinateCleaner: standardized cleaning of occurrence records from biological collection databases.
Methods Ecol. Evol. (2019) doi:10.1111/2041-210X.13152.
96.   Maynard, D. S. *et al.* Global relationships in tree functional traits. *Nat. Commun.* **13**, 3185 (2022).
97.   Oksanen, J. *et al. vegan: Community Ecology Package.* (R Project, 2019).
98.   Weckwerth, W. Metabolomics in systems biology. *Annu. Rev. Plant Biol.* **54**, 669–689 (2003).
99.   Uthe, H. *et al.* A practical guide to implementing metabolomics in plant ecology and biodiversity research. in *Plant*
Metabolomics in full swing vol. 98 163–203 (Elsevier, 2021).
100.   Dray, S. & Dufour, A.-B. The ade4 Package: Implementing the Duality Diagram for Ecologists. *J. Stat. Softw.* **22**,
(2007).
101.   Salguero-Gómez, R. Applications of the fast-slow continuum and reproductive strategy framework of plant life
histories. *New Phytol.* **213**, 1618–1624 (2017).

Acknowledgements

Special thanks to Jeanne Clément and Ghislain Vieilledent for building bespoke functionality into the R package "jSDM" for exploratory analyses. This study was part funded by a grant from the Synthesis Centre (sDiv) of the German Centre for Integrative Biodiversity Research (iDiv), awarded to TWNW, FS and NMvD. Plant material collection and metabolomics analysis were financed by Swiss National Science Foundation grants awarded to LP (grant no. 31003A-162604), SR (grant no. 31003A-131956), and JLW (grant no. CRSII5-189921/1). JMA is funded by the Swiss National Science Foundation (grant no. 31003A-176044). FS acknowledges funding by the University of Nottingham Anne McLaren Fellowship. ED and SR are funded by the Swiss National Science Foundation (grant no. 31003A_179481). VEJJ is funded by the French National Research Agency (grant no. ANR-17-CE01-0007: Mixxopeat). VB is supported by the French Agence Nationale pour la Recherche (grant no. ANR-12-BSV7-0016-01: SecPriMe2), the BioDivMeX Mistrals program, the Aix-Marseille University 'Investissements d'Avenir' program and the Labex OT-Med programs. PC is supported by a Ramón Areces Foundation Postdoctoral Scholarship. PDC is funded by the US National Science Foundation (grant no. DEB-1135733). NMvD, SM and KP are supported by iDiv Halle-Jena-Leipzig, which is funded by the German Research Foundation (grant nos DFG–FZT 118, 202548816). SN and KP are additionally funded by the German Network for Bioinformatics Infrastructure (de.NBI) and acknowledge BMBF funding (grant no. 031L0107). DLF is supported by the US National Science Foundation Graduate Research Fellowships Program. RSG is funded by a UK Natural Environment Research Council Independent Research Fellowship (grant no. NE/M018458/1). JP and JS are funded by the Spanish Government (grant no. CGL2016-79835), Catalan Government (grant no. SGR 2017-1005) and a European Research Council Synergy Grant (grant no. ERC-SyG-2013-610028: IMBALANCE-P). MCS is supported by a NOMIS Foundation project ("Remotely Sensing Ecological Genomics") and by the University of Zürich Research Priority Program on Global Change and Biodiversity.

Author contributions

Conceptualization: TWNW in collaboration with PMA, RDB, PC, BD, ED, AG-G, VEJJ, JP, SR, RSG, JS, MCS, FS, NMvD and WW, with support from all co-authors. Funding acquisition: FS, NMvD, and TWNW. Data collection: PMA, ED, and BD. Data curation: TWNW, with support from PMA, OB, PC, and ED. Formal analysis: TWNW, with support from PC, ED, MJE, DF, and VEJJ. Writing: TWNW, with support from MCS. Reviewing and editing: all co-authors.

Competing interests

The authors declare no competing interests.

Materials & correspondence

Requests for materials and correspondence should be sent to the corresponding author.

     Extended Data Figures

786
787
Extended Data Figure 1 | Tropical and temperate species capture major trade-offs in plant functional traits. (A,B) The
global geographic extents of (A) tropical (yellow; N = 457) and (B) temperate (blue; N = 405) plant species used in this study.
Points are processed GBIF records representing native and non-native occurrences, with sample collection being restricted
to *in situ* native populations. (C-F) Loadings for the first two PCs of PCAs performed on eight classical functional traits
separately for (C,E) tropical and (D,F) temperate species. Axes collectively capture well-established life-history trade-offs
(i.e., leaf economics spectrum[6], plant size/longevity[7,101], plant hardiness[15]; see Main Text), although the coupling of these
trade-offs and the ordering of PC axes differs between tropical and temperate species. (G) Kernel density estimates showing
the distributions of classical traits used in PCAs (height, seed mass (SM), stem specific density (SSD), leaf area (LA) and leaf
carbon, nitrogen and phosphorus) for tropical (blue) and temperate (yellow) species, as well as for all species in the TRY
database (red ribbon[89]). Text shows group means, with values in brackets being back-transformed to original units (log$_{10}$-
transformed traits; see Methods). Asterisks illustrate significant differences (*P* < 0.001; linear models) between tropical and
temperate species means.