1 **Title:**
2 **Leaf metabolic traits reveal hidden dimensions of plant form and function**
3

Authors:
Tom W. N. Walker[1,2]*, Franziska Schrodt[3], Pierre-Marie Allard[4,5], Emmanuel Defossez[2], Vincent E. J.
Jassey[6], Meredith C. Schuman[23], Jake M. Alexander[1], Oliver Baines[3], Virginie Baldy[7], Richard D.
Bardgett[8], Pol Capdevila[9,10], Phyllis D. Coley[12], Nicole M. van Dam[13,14,15], Bruno David[16], Patrice
Descombes[1,26,27], Maria-Jose Endara[12], Catherine Fernandez[7], Dale Forrister[12], Albert Gargallo-
Garriga[17,18,19], Gaëtan Gauser[2], Sue Marr[13,20,21], Steffen Neumann[13,20], Loïc Pellissier[1,26], Kristian
Peters[13,20,21], Sergio Rasmann[2], Ute Roessner[22], Roberto Salguero-Gómez[11], Jordi Sardans[17,18],
Wolfram Weckwerth[24,25], Jean-Luc Wolfender[4,5] & Josep Peñuelas[17,18]
12

Affiliations:
[1] Department of Environmental Systems Science, ETH Zürich, 8092 Zürich, Switzerland
[2] Institute of Biology, University of Neuchâtel, 2000 Neuchâtel, Switzerland.
[3] School of Geography, University of Nottingham, Nottingham NG7 2RD, UK
[4] School of Pharmaceutical Sciences, University of Geneva, 1211 Geneva, Switzerland
[5] Institute of Pharmaceutical Sciences of Western Switzerland, University of Geneva, 1211 Geneva, Switzerland
[6] Laboratoire d'Ecologie Fonctionnelle et Environnement, Université de Toulouse, CNRS, 31062 Toulouse, France
[7] Aix Marseille Université, Avignon Université, CNRS, IRD, IMBE, Marseille, France
[8] Department of Earth & Environmental Sciences, The University of Manchester, Manchester M13 9PT, UK
[9] Departament de Biologia Evolutiva, Ecologia i Ciències Ambientals, Universitat de Barcelona, 08028 Barcelona, Spain
[10] School of Biological Sciences, University of Bristol, Bristol BS8 1TQ, UK
[11] Department of Zoology, Oxford University, Oxford OX1 3SZ, UK
[12] School of Biological Sciences, University of Utah, Salt Lake City, UT 84112, USA
[13] German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, 04103 Leipzig, Germany
[14] Institute of Biodiversity, Friedrich Schiller University Jena, 07743 Jena, Germany
[15] Leibniz Institute of Vegetable and Ornamental crops (IGZ), 14979 Großbeeren, Germany
[16] Green Mission Pierre Fabre, Institut de Recherche Pierre Fabre, 31562 Toulouse, France
[17] CSIC, Global Ecology Unit CREAF-CSIC-UAB, 08193 Bellaterra, Catalonia, Spain
[18] CREAF, 08193 Cerdanyola del Vallès, Catalonia, Spain
[19] Global Change Research Institute, Czech Academy of Sciences, 603 00 Brno, Czech Republic
[20] Leibniz Institute of Plant Biochemistry, Bioinformatics and Scientific Data, 06120 Halle, Germany
[21] Institute of Biology/Geobotany and Botanical Garden, Martin Luther University Halle Wittenberg, 06108 Halle, Germany
[22] Research School of Biology, The Australian National University, 2601 Acton, Australia
[23] Departments of Geography and Chemistry, University of Zurich, 8057 Zurich, Switzerland
[24] Molecular Systems Biology, Department of Functional and Evolutionary Ecology, 1010 University of Vienna, Austria
[25] Vienna Metabolomics Center, 1010 University of Vienna, Austria
[26] Ecosystems & Landscape Evolution, Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), 8903
Birmensdorf, Switzerland
[27] Musée et Jardins botaniques cantonaux, 1007 Lausanne, Switzerland
* Correspondence to: thomas.walker@unine.ch
43

44 **One sentence summary:**
45 Leaf metabolic traits describe novel dimensions of plant life-history variation that expand the functional
46 trait concept.
47
Abstract:

The metabolome is the biochemical basis of plant form and function, but we know little about its
macroecological variation across the plant kingdom. Here, we used the plant functional trait concept to
interpret leaf metabolome variation among 457 tropical and 339 temperate plant species. Distilling
metabolite chemistry into five metabolic functional traits reveals that plants vary on two major axes of
leaf metabolic specialization – a leaf chemical defense spectrum and an expression of leaf longevity.
Axes are similar for tropical and temperate species, with many trait combinations being viable.
However, metabolic traits vary orthogonally to life-history strategies described by widely used
functional traits. The metabolome thus expands the functional trait concept by providing novel axes of
metabolic specialization for examining plant form and function.

Main Text:

Plants produce a staggering diversity of metabolites – upwards of one million throughout the plant kingdom and many thousands within an individual (*1*) (the metabolome). The plant metabolome is well known to human society, being the source of most cosmetics (*2*), flavors (*3*), and medicines (*4*). Yet, little is known about how the metabolome varies across the plant kingdom. The metabolome is the biochemical basis of physiology, comprising primary metabolites involved in cellular function (e.g., carbohydrates, amino acids) and specialized metabolites produced to confront stress (e.g., flavonoids, terpenoids) (*1*, *5*, *6*). The metabolome thus encompasses the biochemical mechanisms through which evolutionary and ecological processes shape plant form and function (i.e., morphology and physiology (*7*)) (*8*, *9*). Plant form and function vary predictably among species because evolutionary and ecological processes generally act to maximize fitness (*10–14*), thus limiting the number of morphological or physiological strategies that are viable (*7*, *15*, *16*). It follows that the metabolome should also be constrained to a limited number of strategies that maximize fitness (*17*). However, although cross-species studies are now emerging (*18–21*), no attempt has been made to characterize how the metabolome varies systematically throughout the plant kingdom. We thus remain unable to place the vast diversity of the plant metabolome into a coherent ecological context.

The plant functional trait concept offers a powerful framework to contextualize the plant metabolome. Plant functional traits are a standardized set of morphological and physiological characteristics that provide universal metrics of plant form and function (*22*). Plant functional traits have been pivotal in identifying the life-history strategies that govern plant fitness (*7*, *15*, *16*), understanding plant impacts on population (*23*), community (*24*), and ecosystem (*25*) processes, and predicting global change effects on ecosystems (*26*). Plant functional traits thus provide a strong reference point from which to interpret variation in the plant metabolome. However, measurements of the metabolome remain absent from the functional trait concept, in part because they yield complex data describing thousands of metabolites from hundreds of biochemical pathways (*17*). Advances in chemoinformatics make it now possible to translate metabolite identities into a targeted set of chemical properties (*27*, *28*). Doing so resonates with the functional trait concept – i.e., that quantitative characteristics provide a bridge between identity and function (*22*) – and is an intuitive way to contextualize the metabolome against plant functional traits. Yet, while chemoinformatics is commonplace in natural products chemistry (*29–31*), it remains entirely absent from ecology.

Contextualizing the plant metabolome against existing plant functional traits raises two alternative hypotheses. If the metabolome varies colinearly with functional traits, then it offers a biochemical validation of the functional trait concept. Such validation is necessary because functional traits are emergent properties of multiple biochemical processes (*22*, *32*), making them proxies for physiology that can be poor predictors of ecological processes (*33–35*). Alternatively, if the metabolome varies orthogonally to functional traits, then it describes dimensions of plant life-history variation missed by existing functional traits. Such orthogonal variation would enhance the explanatory power of the functional trait concept by providing additional axes of trait specialization to examine plant form and function. Either way, integrating measurements of the plant metabolome into the plant functional trait concept will yield insight into how the metabolome varies across the plant kingdom, while also offering a high-resolution mechanistic lens with which to advance the functional trait concept itself.

102
103
Figure 1 | Variation in leaf metabolite chemistry can be captured by five chemical properties. (A) Coefficients for
Pearson correlations among 21 chemical properties (Table S1) of annotated metabolites detected in tropical and temperate
datasets (N = 4292). Properties are ordered based on correlation similarity (hierarchical clustering; Ward), with colors
separating distinct clusters of properties (maximal silhouette width: K = 5). The lower panel shows correlations between
chemical properties and scores from the first three axes of a PCA performed on a subset of five properties, as well as (B)
between subset PC scores and either scores from a PCA performed on all 21 properties (left panel) or metabolite positions
on the first three axes of ChemGPS-NP (*31*) space (right panel). Circle sizes indicate coefficient strengths, while fills
discriminate between positive (black) and negative (white) correlations. (C) The chemical properties selected to represent
each cluster, alongside chemical graphs of metabolites with the lowest and highest values for each trait.

Here, we performed an assessment of the leaf metabolomes of 457 tropical and 339 temperate plant
species to characterize variation in the metabolome and test whether it validates, or expands upon,
existing plant functional traits. First, we examined the chemical properties of all identified metabolites
(N = 4292) to derive a minimum set of "metabolic functional traits" that capture chemical variation in
the leaf metabolome. We then characterized how metabolic traits vary among species to identify major
axes of leaf metabolic specialization in the plant kingdom. Finally, we combined metabolic traits with
eight widely used functional traits (plant height, seed mass, stem specific density, leaf area, specific
leaf area and leaf carbon, nitrogen and phosphorus concentrations; hereafter "classical traits") (7) to
determine whether axes of metabolic specialization are colinear with, or orthogonal to, plant life-history
strategies described by classical functional traits.
123
<u>Five chemical properties capture chemical variation in the leaf metabolome</u>
We expressed the leaf metabolomes of tropical (36) and temperate (37) plant species as the presence
or absence of annotated metabolic features (i.e., metabolites) detected through liquid chromatography
mass spectrometry (tropical N = 3356; temperate N = 2227). We then characterized the chemistry of
the leaf metabolome by examining variation in the chemical properties of all unique metabolites from
both datasets (N = 4292) (31). We focused on 21 properties describing metabolite constitution,
geometry, or topology that have relationships with (bio)chemical activity (Table S1). Pairwise
correlations among chemical properties (Fig. 1A, upper panel) reveal that properties separate into five
clusters, each representing a distinct facet of leaf metabolite chemistry. The largest cluster (Fig. 1A,
green) describes overall size and structural complexity, comprising total numbers of atoms and bonds,
molecular weight, a predictor of polarity based on the proportion of non-carbon atoms (M log P; (38)),
and related topological parameters (i.e., Wiener numbers (39), Eccentric Connectivity (40)). A second
cluster (Fig. 1A, purple) describes bond conjugation and reactivity, comprising numbers of aromatic
atoms and bonds, numbers of atoms in the largest conjugated (pi-) system, and a derived complexity
index indicative of promiscuous activity in drugs (f_{MF}; (41)). A third cluster (Fig. 1A, yellow) describes
polar intermolecular (non-covalent) forces, comprising total and mass-specific polar surface area and
numbers of hydrogen (H) bond donors or acceptors (42, 43). Such forces are important for the
selective binding of small molecules to DNA (44), RNA (45), and proteins (46). A fourth cluster (Fig.
1A, blue) comprises two predictors of metabolite polarity based on summed properties of atoms and
bonds (X log P (47), A log P (48)), where lower values indicate higher polarity. A final cluster (Fig. 1A,
red) describes carbon bond saturation via the relative occurrence of sp^3 (unsaturated, three-
dimensional) and sp^2 (saturated, planar) hybridization (49). Overall, these clusters indicate that leaf
metabolite chemistry is highly structured and can be described by five distinct facets: size/complexity,
bond conjugation/reactivity, contributions to polar intermolecular forces, general polarity, and carbon
bond saturation.
149
The strong collinearity among chemical properties (Fig. 1A, upper panel) suggests that the five facets
of leaf chemistry can be represented by a small number of discriminant variables. Indeed, a PCA
performed on a subset of five chemical properties, one from each cluster (Fig. 1C), is equivalent to a
PCA performed on all 21 properties (Fig. 1B, left panel; PC1: r_{4290} = 0.97; PC2: r_{4290} = 0.99; PC3: r_{4290}
= 0.93; P < 0.001 in all cases). We selected molecular weight, H-bond acceptor count, aromatic atom
count, polarity (X log P), and carbon hybridization (sp^3:sp^2) as five properties that are intuitive for non-
specialists to interpret (Fig. 1C; Table S1). However, most combinations of properties covering all
facets of leaf chemistry also yield equivalent PCAs (Fig. S5). Pairwise correlations between all
properties and subset PC scores further confirm that selected properties capture observed leaf

chemical variation, in that correlations are consistent within each cluster (Fig. 1A, lower panel).
Selected properties also describe biologically relevant chemical space according to an existing
framework that classifies natural products based on size, then aromaticity, then lipophilicity (Fig. 1B,
right panel; PC1: r_{4290} = 0.99; PC2: r_{4290} = 0.87; r_{4290} = 0.81; P < 0.001 in all cases) (*31*). The first axis
of the subset PCA (38% variance) separates heavy metabolites with many H-bond acceptors from
light metabolites with fewer H-bond acceptors. Alignment between metabolite size and numbers of H-
bond acceptors is expected because larger molecules usually engage in stronger intermolecular
interactions (*50*). The second axis (37% variance) separates reactive metabolites with more aromatic
carbon bonds from unreactive metabolites with more saturated carbon bonds. This axis is also intuitive
because conjugated bond systems are unsaturated, reactive, and planar, whereas saturated bonds,
such as those abundant in fatty acids, are unreactive and three-dimensional (*51*). The third axis (18%
variance) separates nonpolar metabolites from polar metabolites (*29*). Our findings thus show that leaf
chemical variation can be represented by five chemical properties, one from each facet of leaf
chemistry.
173
<u>Five chemical properties provide more information than metabolite family</u>
The five selected chemical properties not only represent leaf metabolite chemistry (Fig. 1), but also
reflect known differences among metabolite families. Using the subset PCA to interpret variation
among metabolite families (Fig. 2), we observe that peptides are large, display average bond
saturation, and are moderately polar; carbohydrates are of average size, display high bond saturation,
and are highly polar; and fatty acids are smaller, display high bond saturation, and are of average
polarity. Lignans, alkaloids, coumarins and flavonoids, families of specialized metabolites, display low
bond saturation due to the presence of many aromatic bonds, with alkaloids and coumarins also being
relatively small. Terpenoids, a family of abundant specialized metabolites (N = 2306), vary in size,
bond saturation, and polarity, echoing their diversity across the plant kingdom (*52*). Although chemical
properties generally differentiate among metabolite families, all families possess variable and
overlapping chemistries (Fig. 2, marginal boxplots). We thus suggest that chemical properties,
quantitative metrics of the chemistry that underpins metabolite composition and activity, provide more
precise information about metabolite function than metabolite family identity. Indeed, drug discovery
and medicinal chemistry rely on many of the same chemical properties to screen for bioactive
molecules (*29–31*). Such a perspective resonates with the functional trait concept by reaching beyond
taxonomy to describe function (*22*) and provides a strong foundation for a novel set of metabolic
functional traits.
192
193

Figure 2 | Five chemical properties describe variation among and within metabolite families. Biplots of PC1 scores
and (A) PC2 scores or (B) PC3 scores from a PCA describing variation in selected chemical properties (C hybridization, H-
bond acceptors, molecular weight, polarity, aromatic atoms) among all annotated leaf metabolites (N = 4221) detected in 862
plant species. The PCA is functionally equivalent to a PCA containing all 21 chemical properties (see Main Text), with axes
separating leaf metabolites based on size (PC1), aromaticity (PC2), and a-polarity (PC3; see Fig. 1A, lower panel). Colors
separate metabolite families as defined by marginal boxplot fills/labels (center: median; box: 25% to 75% quantiles; whiskers:
1.5 × IQR; points: outliers beyond whiskers).

<u>Five metabolic functional traits describe two axes of leaf metabolic specialisation</u>
We converted selected properties into plant-level metabolic functional traits to examine how the leaf
metabolome varies among tropical and temperate plant species. We first compared the volume and
"lumpiness" (i.e., aggregation around values) of multidimensional space occupied by metabolic traits
to four null models to test whether species converge on a restricted set of interrelated trait
combinations (7). Observed hypervolumes of metabolic traits are 98.1% to 99.8% smaller and 71.0%
to 94.7% lumpier than those of three null models assuming no trait covariance (Table S2; Table S3;
always $P < 0.001$ in all cases). However, observed hypervolumes are larger (tropical: $P = 0.002$;

temperate: P = 0.047) than that of the null model assuming trait covariance and selection against
extreme values, while also being similarly lumpy (tropical: P = 0.662; temperate: P = 0.858). By
comparison, observed hypervolumes of eight classical traits are smaller than null models assuming no
trait covariance (Table S2; P < 0.001 in all cases), either the same size as (tropical species; P =
0.154) or smaller than (temperate species; P = 0.018) that of the null model assuming selection
against extreme values, and lumpier than all null models (Table S3; P < 0.001 in all cases).
Collectively, these findings yield three insights. First, metabolic functional traits are interrelated, so
plants are constrained to a limited number of viable metabolic trait combinations. Second, while
extreme metabolic trait values may be selected against in temperate species, this is not so for tropical
species. Expressing extreme leaf metabolome chemistry is thus a viable life-history strategy in tropical
environments, potentially due to more intense biotic interactions selecting for a diversity of defensive
leaf metabolites and thus divergence of chemical strategies (*53*). Third, metabolic traits are less
aggregated around non-extreme values than classical traits. As such, although plants converge
towards one of several viable classical trait strategies (e.g., woody, non-woody; (*7*)), they can possess
a wider array of leaf metabolome strategies.
228
229

232 **Figure 3 | Metabolic traits describe two axes of leaf metabolic specialization among plant species.** Biplots of PC1 and
233 PC2 scores for PCAs describing variation in five metabolic functional traits (arrows) between (A) tropical (yellow; N = 457)
234 and (B) temperate (blue; N = 339) plant species. Axes separate species based on the mean aromaticity *versus* polarity
235 (inverse X log *P*) and carbon bond saturation (PC1) and mean size/complexity (PC2) of all annotated leaf metabolites
236 present (Methods).
237
238
We performed PCAs on metabolic traits to characterize major axes of leaf metabolic specialization
among species. Metabolic trait variation is consistent among tropical and temperate species, with 92%
variation being explained by the first two PCs (Fig. 3). The first axis (tropical: 52%; temperate 49%)
separates species that produce more unsaturated, aromatic, nonpolar leaf metabolites from those that
produce more saturated, polar, leaf metabolites. This axis likely describes a spectrum of leaf chemical
defense strategies (*12*). Unsaturated aromatic metabolites, such as alkaloids, coumarins, and
flavonoids (Fig. 2), are reactive and serve as toxins or antioxidants in response to stress. For instance,
conjugated bond structures can interfere with protein function by binding covalently to sidechains (*54*),
generate or quench oxidative stress (*55*), or absorb damaging wavelengths of light (*56*). By contrast,
saturated non-aromatic metabolites can play different roles in leaf defense, both directly as toxins (*57*)

and indirectly as signaling molecules (*58*). An alternative explanation is that the first axis reflects an
investment (or not) into defense metabolites (*12*), which could arise due to the high cost of
synthesizing aromatic metabolites (*13*). However, all plants must defend their leaves against stress,
and we find that plants can maximize fitness at either end of this axis (Fig. 3). It is thus more plausible
that the first axis represents a leaf chemical defense spectrum.
254
The second PCA axis (tropical: 40%; temperate: 43%) discriminates species that produce larger
metabolites with many H-bond acceptors from those that produce smaller metabolites with fewer H-
bond acceptors. This axis likely relates to leaf longevity, since large metabolites, such as lignans,
peptides, and waxes, form the basis of long-lived physical (*59*, *60*) and storage (*61*) structures. Many
of these structures also depend on strong intermolecular forces to form stable macromolecules or
complexes (*62*, *63*), which are enhanced by H-bond donors and acceptors (*43*). As for the first axis,
plants can maximize fitness at either end of the second axis, with species at the positive end of the
axis investing in metabolites indicative of a longer leaf lifespan. As such, the second axis is probably a
manifestation of the fast-slow continuum (*64*) at the leaf level. Alternatively, the second axis may
reflect an investment (or not) into structurally complex metabolites for physical defense, such as
lignans or waxes (*59*, *60*). However, such metabolites act by creating rigid structures that lengthen leaf
lifespan anyway (*65*), so physical protection is one component of leaf longevity. Taken together,
metabolic trait PCAs reveal two major axes of leaf metabolic specialization. These axes relate to
metabolite bond saturation and polarity (i.e., a leaf chemical defense spectrum) and metabolite size
and propensity for intermolecular interactions (i.e., leaf longevity), with many combinations yielding
successful life-history strategies.
271
<u>Axes of leaf metabolic specialisation reveal hidden dimensions of plant form and function</u>
The apparent consistency of metabolic trait variation with classical life-history theory (*7*, *12*, *13*, *15*, *16*,
64, *65*) raises the possibility that metabolic traits describe aspects of plant form and function already
captured by classical traits. We thus performed a PCA on metabolic traits plus the same widely used
functional traits used above to test whether axes of leaf metabolic specialization are colinear with, or
orthogonal to, trait-based life-history strategies. Considered alone, classical trait variation supports life-
history theory (*7*), with the first two PCs (tropical: 49%; temperate: 51%) describing the leaf economics
spectrum (SLA, leaf nitrogen, leaf phosphorus; (*16*)) and variation indicative of plant longevity (plant
height, seed mass; (*64*)) and hardiness (stem density, leaf carbon; (*15*)) (Fig. S1). However, when
combining classical and metabolic traits in the same PCA, the first four PCs (tropical: 66%; temperate:
67%) are equivalent to axes described by individual PCAs (Fig. 4A, lower marginal matrices). The first
and third axes describe the first two axes of leaf metabolic specialization (Fig. 3), whereas the second
and fourth axes describe the first two axes of classical trait specialization (Fig. S1). Pairwise
correlations among all traits further show that metabolic and classical traits do not covary (Fig. 4A,B),
but separate into six clusters describing unique aspects of either metabolic or classical trait variation.
As such, major axes of leaf metabolic specialization vary orthogonally to major axes of classical trait
variation (Fig. S4) and are at least equally relevant for describing variation in plant form and function
among tropical and temperate plant species.
290
291

Figure 4 | Metabolic and classical traits describe unique dimensions of plant form and function. Correlation
coefficients for pairwise Pearson correlations among metabolic (Fig. 3) and classical functional traits (Fig. S1) in (A) tropical
(N = 457) and (B) temperate (N = 339) species. Traits are ordered based on correlation similarity (hierarchical clustering;
Ward), with colors separating distinct clusters (maximal silhouette width: K = 6) and horizontal lines between matrices
showing correspondence between tropical and temperate traits. Lower panel shows correlations between traits and scores
from the first four axes of a PCA of the same traits (tropical: 66%; temperate: 67%; LES = leaf economics spectrum). Circle
sizes indicate coefficient strengths, while fills discriminate between positive (black) and negative (white) correlations.

301
302
Leaf metabolism underpins energy production *via* photosynthesis (*66*), but in doing so yields tissue
that is sensitive to light (*67*) and temperature (*68*) and is of high nutritional value to herbivores (*65*).
Plants are thus under selective pressure to maximize photosynthetic rates while also protecting leaf
tissue against damage. While this trade-off is consistent with life-history trade-offs derived from
classical traits (*7, 15, 16*), it does not necessarily follow that individuals apply the same strategy in
every organ to maintain fitness. Indeed, we observe that species with "productive" trait values (e.g.,
high SLA, high leaf nitrogen) can produce leaves containing many complex aromatic metabolites,
species possessing "hardy" trait values (e.g., high stem density, high leaf carbon) can produce leaves
containing many simple polar metabolites, and species possessing trait values typical of a fast pace of
life (e.g., low stature, small seeds) can produce leaves with many large stable or structural
metabolites. Our findings thus question classical trait (*16*) and plant defense (*12*) theory that predicts
relationships between the leaf chemical phenotype, plant productivity, and pace of life. In short,
metabolic functional traits describe unique dimensions of plant life-history variation that are
complementary to, and independent from, those captured by classical functional traits.

<u>Concluding remarks</u>
In summary, we show that leaf metabolite chemistry is highly structured and can be described by five
chemical properties. Using chemical properties as metabolic functional traits reveals that plants vary

on two axes of leaf metabolic specialization – a leaf chemical defense spectrum (i.e., bond saturation,
polarity) and leaf longevity (i.e., size, propensity for intermolecular interactions). Axes of metabolic
specialization are similar in tropical and temperate species, with many combinations of trait values
yielding successful life-history strategies. Axes are also consistent with expectations from life-history
theory that plants invest strategically in tissue defense (*12*, *13*, *16*) and between a fast *versus* slow
pace of life (*7*, *64*). Nevertheless, we find that metabolic trait variation is orthogonal to, not colinear
with, classical trait variation. Metabolic functional traits thus reveal dimensions of plant life-history
variation missed by existing functional traits. It is important to note that metabolomics data were
acquired differently for tropical and temperate species and were matched to open functional trait
databases. Nevertheless, we find consistent variation in the leaf metabolomes of 457 tropical and 339
temperate species that possess distinct geographic distributions and evolutionary histories (Fig.
S1A,B), as well as that encompass most documented functional trait variation (Fig. S1G, red ribbons).
We thus propose that distilling the leaf metabolome into metabolic functional traits captures
macroecological patterns of metabolic variation widely across the plant kingdom, enhances the
explanatory power of the functional trait concept, and offers a new set of tools for the discovery of
species or genotypes with trait combinations adapted to societal needs.

References and Notes

1. S. Wang, S. Alseekh, A. R. Fernie, J. Luo, The structure and function of major plant metabolite modifications. *Mol. Plant*. **12**, 899–919 (2019).
2. G. Faccio, Plant complexity and cosmetic innovation. *iScience*. **23**, 101358 (2020).
3. H. J. Klee, D. M. Tieman, The genetics of fruit flavour preferences. *Nat. Rev. Genet.* **19**, 347–356 (2018).
4. B. M. Schmidt, D. M. Ribnicky, P. E. Lipsky, I. Raskin, Revisiting the ancient concept of botanical therapeutics. *Nat. Chem. Biol.* **3**, 360–366 (2007).
5. J. Sardans, A. Gargallo-Garriga, O. Urban, K. Klem, T. W. N. Walker, P. Holub, I. A. Janssens, J. Peñuelas, Ecometabolomics for a better understanding of plant responses and acclimation to abiotic factors linked to global change. *Metabolites*. **10** (2020), doi:10.3390/metabo10060239.
6. J. Sardans, A. Gargallo-Garriga, O. Urban, K. Klem, P. Holub, I. A. Janssens, T. W. N. Walker, A. Pesqueda, J. Peñuelas, Ecometabolomics of plant–herbivore and plant–fungi interactions: a synthesis study. *Ecosphere*. **12** (2021), doi:10.1002/ecs2.3736.
7. S. Díaz, J. Kattge, J. H. C. Cornelissen, I. J. Wright, S. Lavorel, S. Dray, B. Reu, M. Kleyer, C. Wirth, I. C. Prentice, E. Garnier, G. Bönisch, M. Westoby, H. Poorter, P. B. Reich, A. T. Moles, J. Dickie, A. N. Gillison, A. E. Zanne, J. Chave, S. J. Wright, S. N. Sheremet'ev, H. Jactel, C. Baraloto, B. Cerabolini, S. Pierce, B. Shipley, D. Kirkup, F. Casanoves, J. S. Joswig, A. Günther, V. Falczuk, N. Rüger, M. D. Mahecha, L. D. Gorné, The global spectrum of plant form and function. *Nature*. **529**, 167–171 (2016).
8. M. Wink, Evolution of secondary metabolites from an ecological and molecular phylogenetic perspective. *Phytochemistry*. **64**, 3–19 (2003).
9. M. C. Schuman, N. M. van Dam, F. Beran, W. S. Harpole, How does plant chemical diversity contribute to biodiversity at higher trophic levels? *Curr. Opin. Insect Sci.* **14**, 46–55 (2016).
10. A. E. Zanne, D. C. Tank, W. K. Cornwell, J. M. Eastman, S. A. Smith, R. G. FitzJohn, D. J. McGlinn, B. C. O'Meara, A. T. Moles, P. B. Reich, D. L. Royer, D. E. Soltis, P. F. Stevens, M. Westoby, I. J. Wright, L. Aarssen, R. I. Bertin, A. Calaminus, R. Govaerts, F. Hemmings, M. R. Leishman, J. Oleksyn, P. S. Soltis, N. G. Swenson, L. Warman, J. M. Beaulieu, Three keys to the radiation of angiosperms into freezing environments. *Nature*. **506**, 89–92 (2014).
11. G. E. Hutchinson, *The Ecological Theater and the Evolutionary Play* (Yale University Press, 1965).
12. P. Feeny, in *Biochemical interaction between plants and insects*, J. W. Wallace, R. L. Mansell, Eds. (Springer US, Boston, MA, 1976), pp. 1–40.
13. P. D. Coley, J. P. Bryant, F. S. Chapin, Resource availability and plant antiherbivore defense. *Science*. **230**, 895–899 (1985).
14. P. Capdevila, T. W. N. Walker, F. Schrodt, R. C. Rodriguez Caro, R. Salguero-Gomez, Global patterns of plant form and function are strongly determined by evolutionary relationships. *BioRxiv* (2023), doi:10.1101/2023.01.13.523963.
15. J. P. Grime, K. Thompson, R. Hunt, J. G. Hodgson, J. H. C. Cornelissen, I. H. Rorison, G. A. F. Hendry, T. W. Ashenden, A. P. Askew, S. R. Band, R. E. Booth, C. C. Bossard, B. D. Campbell, J. E. L. Cooper, A. W. Davison, P. L. Gupta, W. Hall, D. W. Hand, M. A. Hannah, S. H. Hillier, D. J. Hodkinson, A. Jalili, Z. Liu, J. M. L. Mackey, N. Matthews, M. A. Mowforth, A. M. Neal, R. J. Reader, K. Reiling, W. Ross-Fraser, R. E. Spencer, F. Sutton, D. E. Tasker, P. C. Thorpe, J. Whitehouse, Integrated screening validates primary axes of specialisation in plants. *Oikos*. **79**, 259 (1997).
16. I. J. Wright, P. B. Reich, M. Westoby, D. D. Ackerly, Z. Baruch, F. Bongers, J. Cavender-Bares, T. Chapin, J. H. C. Cornelissen, M. Diemer, J. Flexas, E. Garnier, P. K. Groom, J. Gulias, K. Hikosaka, B. B. Lamont, T. Lee, W. Lee, C. Lusk, J. J. Midgley, M.-L. Navas, U. Niinemets, J. Oleksyn, N. Osada, H. Poorter, P. Poot, L. Prior, V. I. Pyankov, C. Roumet, S. C. Thomas, M. G. Tjoelker, E. J. Veneklaas, R. Villar, The worldwide leaf economics spectrum. *Nature*. **428**, 821–827 (2004).
17. T. W. N. Walker, J. M. Alexander, P. Allard, O. Baines, V. Baldy, R. D. Bardgett, P. Capdevila, P. D. Coley, B. David, E. Defossez, M. Endara, M. Ernst, C. Fernandez, D. Forrister, A. Gargallo-Garriga, V. E. J. Jassey, S. Marr, S. Neumann, L. Pellissier, J. Peñuelas, K. Peters, S. Rasmann, U. Roessner, J. Sardans, F. Schrodt, M. C. Schuman, A. Soule, H. Uthe, W. Weckwerth, J. Wolfender, N. M. Dam, R. Salguero-Gómez, Functional Traits 2.0: The power of the metabolome for ecology. *Journal of Ecology*. **110**, 4–20 (2022).
18. B. E. Sedio, J. D. Parker, S. M. McMahon, S. J. Wright, Comparative foliar metabolomics of a tropical and a temperate forest community. *Ecology*. **99**, 2647–2653 (2018).
19. D. L. Forrister, M.-J. Endara, G. C. Younkin, P. D. Coley, T. A. Kursar, Herbivores as drivers of negative density dependence in tropical forest saplings. *Science*. **363**, 1213–1216 (2019).
20. A. Gargallo-Garriga, J. Sardans, V. Granda, J. Llusià, G. Peguero, D. Asensio, R. Ogaya, I. Urbina, L. Van Langenhove, L. T. Verryckt, J. Chave, E. A. Courtois, C. Stahl, O. Grau, K. Klem, O. Urban, I. A. Janssens, J. Peñuelas, Different "metabolomic niches" of the highly diverse tree species of the French Guiana rainforests. *Sci. Rep.* **10**, 6937 (2020).
21. C. Scherling, C. Roscher, P. Giavalisco, E.-D. Schulze, W. Weckwerth, Metabolomics unravel contrasting effects of biodiversity on the performance of individual plant species. *PLoS ONE*. **5**, e12569 (2010).
22. C. Violle, M.-L. Navas, D. Vile, E. Kazakou, C. Fortunel, I. Hummel, E. Garnier, Let the concept of trait be functional! *Oikos*. **116**, 882–892 (2007).
23. P. B. Adler, R. Salguero-Gómez, A. Compagnoni, J. S. Hsu, J. Ray-Mukherjee, C. Mbeau-Ache, M. Franco, Functional traits explain variation in plant life history strategies. *Proc Natl Acad Sci USA*. **111**, 740–745 (2014).
24. H. Bruelheide, J. Dengler, O. Purschke, J. Lenoir, B. Jiménez-Alfaro, S. M. Hennekens, Z. Botta-Dukát, M. Chytrý, R. Field, F. Jansen, J. Kattge, V. D. Pillar, F. Schrodt, M. D. Mahecha, R. K. Peet, B. Sandel, P. van Bodegom, J.

Altman, E. Alvarez-Dávila, M. A. S. Arfin Khan, F. Attorre, I. Aubin, C. Baraloto, J. G. Barroso, M. Bauters, E. Bergmeier, I. Biurrun, A. D. Bjorkman, B. Blonder, A. Čarni, L. Cayuela, T. Černý, J. H. C. Cornelissen, D. Craven, M. Dainese, G. Derroire, S. De Sanctis, S. Díaz, J. Doležal, W. Farfan-Rios, T. R. Feldpausch, N. J. Fenton, E. Garnier, G. R. Guerin, A. G. Gutiérrez, S. Haider, T. Hattab, G. Henry, B. Hérault, P. Higuchi, N. Hölzel, J. Homeier, A. Jentsch, N. Jürgens, Z. Kącki, D. N. Karger, M. Kessler, M. Kleyer, I. Knollová, A. Y. Korolyuk, I. Kühn, D. C. Laughlin, F. Lens, J. Loos, F. Louault, M. I. Lyubenova, Y. Malhi, C. Marcenò, M. Mencuccini, J. V. Müller, J. Munzinger, I. H. Myers-Smith, D. A. Neill, Ü. Niinemets, K. H. Orwin, W. A. Ozinga, J. Penuelas, A. Pérez-Haase, P. Petřík, O. L. Phillips, M. Pärtel, P. B. Reich, C. Römermann, A. V. Rodrigues, F. M. Sabatini, J. Sardans, M. Schmidt, G. Seidler, J. E. Silva Espejo, M. Silveira, A. Smyth, M. Sporbert, J.-C. Svenning, Z. Tang, R. Thomas, I. Tsiripidis, K. Vassilev, C. Violle, U. Jandt, Global trait-environment relationships of plant communities. *Nat. Ecol. Evol.* **2**, 1906–1917 (2018).

25. G. B. De Deyn, J. H. C. Cornelissen, R. D. Bardgett, Plant functional traits and soil carbon sequestration in contrasting biomes. *Ecol. Lett.* **11**, 516–531 (2008).

26. S. Lavorel, J. Storkey, R. D. Bardgett, F. de Bello, M. P. Berg, X. Le Roux, M. Moretti, C. Mulder, R. J. Pakeman, S. Díaz, R. Harrington, A novel framework for linking functional diversity of plants with other trophic levels for the quantification of ecosystem services. *J. Veg. Sci.* **24**, 942–948 (2013).

27. E. L. Willighagen, J. W. Mayfield, J. Alvarsson, A. Berg, L. Carlsson, N. Jeliazkova, S. Kuhn, T. Pluskal, M. Rojas-Chertó, O. Spjuth, G. Torrance, C. T. Evelo, R. Guha, C. Steinbeck, The Chemistry Development Kit (CDK) v2.0: atom typing, depiction, molecular formulas, and substructure searching. *J. Cheminform.* **9**, 33 (2017).

28. J. Rosén, A. Lövgren, T. Kogej, S. Muresan, J. Gottfries, A. Backlund, ChemGPS-NP(Web): chemical space navigation online. *J. Comput. Aided Mol. Des.* **23**, 253–259 (2009).

29. C. A. Lipinski, F. Lombardo, B. W. Dominy, P. J. Feeney, Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. *Adv. Drug Deliv. Rev.* **46**, 3–26 (2001).

30. F. Lovering, J. Bikker, C. Humblet, Escape from flatland: increasing saturation as an approach to improving clinical success. *J. Med. Chem.* **52**, 6752–6756 (2009).

31. J. Larsson, J. Gottfries, S. Muresan, A. Backlund, ChemGPS-NP: tuned for navigation in biologically relevant chemical space. *J. Nat. Prod.* **70**, 789–794 (2007).

32. J. L. Funk, J. E. Larson, G. M. Ames, B. J. Butterfield, J. Cavender-Bares, J. Firn, D. C. Laughlin, A. E. Sutton-Grier, L. Williams, J. Wright, Revisiting the Holy Grail: using plant functional traits to understand ecological processes. *Biol. Rev. Camb. Philos. Soc.* **92**, 1156–1173 (2017).

33. J. Firn, J. M. McGree, E. Harvey, H. Flores-Moreno, M. Schütz, Y. M. Buckley, E. T. Borer, E. W. Seabloom, K. J. La Pierre, A. M. MacDougall, S. M. Prober, C. J. Stevens, L. L. Sullivan, E. Porter, E. Ladouceur, C. Allen, K. H. Moromizato, J. W. Morgan, W. S. Harpole, Y. Hautier, N. Eisenhauer, J. P. Wright, P. B. Adler, C. A. Arnillas, J. D. Bakker, L. Biederman, A. A. D. Broadbent, C. S. Brown, M. N. Bugalho, M. C. Caldeira, E. E. Cleland, A. Ebeling, P. A. Fay, N. Hagenah, A. R. Kleinhesselink, R. Mitchell, J. L. Moore, C. Nogueira, P. L. Peri, C. Roscher, M. D. Smith, P. D. Wragg, A. C. Risch, Leaf nutrients, not specific leaf area, are consistent indicators of elevated nutrient inputs. *Nat. Ecol. Evol.* **3**, 400–406 (2019).

34. F. van der Plas, T. Schröder-Georgi, A. Weigelt, K. Barry, S. Meyer, A. Alzate, R. L. Barnard, N. Buchmann, H. de Kroon, A. Ebeling, N. Eisenhauer, C. Engels, M. Fischer, G. Gleixner, A. Hildebrandt, E. Koller-France, S. Leimer, A. Milcu, L. Mommer, P. A. Niklaus, Y. Oelmann, C. Roscher, C. Scherber, M. Scherer-Lorenzen, S. Scheu, B. Schmid, E.-D. Schulze, V. Temperton, T. Tscharntke, W. Voigt, W. Weisser, W. Wilcke, C. Wirth, Plant traits alone are poor predictors of ecosystem properties and long-term ecosystem functioning. *Nat. Ecol. Evol.* **4**, 1602–1611 (2020).

35. T. W. N. Walker, W. Weckwerth, L. Bragazza, L. Fragner, B. G. Forde, N. J. Ostle, C. Signarbieux, X. Sun, S. E. Ward, R. D. Bardgett, Plastic and genetic responses of a common sedge to warming have contrasting effects on carbon cycle processes. *Ecol. Lett.* **22**, 159–169 (2019).

36. P.-M. Allard, A. Gaudry, L.-M. Quirós-Guerrero, A. Rutz, M. Dounoue-Kubo, T. W. N. Walker, E. Defossez, C. Long, A. Grondin, B. David, J.-L. Wolfender, Open and reusable annotated mass spectrometry dataset of a chemodiverse collection of 1,600 plant extracts. *Gigascience*. **12** (2022), doi:10.1093/gigascience/giac124.

37. E. Defossez, C. Pitteloud, P. Descombes, G. Glauser, P.-M. Allard, T. W. N. Walker, P. Fernandez-Conradi, J.-L. Wolfender, L. Pellissier, S. Rasmann, Spatial and evolutionary predictability of phytochemical diversity. *Proc Natl Acad Sci USA*. **118** (2021), doi:10.1073/pnas.2013344118.

38. R. Mannhold, G. I. Poda, C. Ostermann, I. V. Tetko, Calculation of molecular lipophilicity: State-of-the-art and comparison of log P methods on more than 96,000 compounds. *J. Pharm. Sci.* **98**, 861–893 (2009).

39. H. Wiener, Structural determination of paraffin boiling points. *J. Am. Chem. Soc.* **69**, 17–20 (1947).

40. V. Sharma, R. Goswami, A. K. Madan, Eccentric connectivity index: A novel highly discriminating topological descriptor for structure−property and structure−activity studies. *J. Chem. Inf. Comput. Sci.* **37**, 273–282 (1997).

41. Y. Yang, H. Chen, I. Nilsson, S. Muresan, O. Engkvist, Investigation of the relationship between topology and selectivity for druglike molecules. *J. Med. Chem.* **53**, 7709–7714 (2010).

42. B. Kuhn, P. Mohr, M. Stahl, Intramolecular hydrogen bonding in medicinal chemistry. *J. Med. Chem.* **53**, 2601–2611 (2010).

43. Th. Zeegers-Huyskens, P. Huyskens, in *Intermolecular Forces*, P. L. Huyskens, W. A. P. Luck, T. Zeegers-Huyskens, Eds. (Springer Berlin Heidelberg, Berlin, Heidelberg, 1991), pp. 1–30.

44. P. B. Dervan, Molecular recognition of DNA by small molecules. *Bioorg. Med. Chem.* **9**, 2215–2235 (2001).

45. G. Padroni, N. N. Patwardhan, M. Schapira, A. E. Hargrove, Systematic analysis of the interactions driving small molecule-RNA recognition. *RSC Med. Chem.* **11**, 802–813 (2020).

46. M. Gao, J. Skolnick, A comprehensive survey of small-molecule binding pockets in proteins. *PLoS Comput. Biol.* **9**,
e1003302 (2013).
47. R. Wang, Y. Fu, L. Lai, A New Atom-Additive Method for Calculating Partition Coefficients. *J. Chem. Inf. Comput. Sci.*
37, 615–621 (1997).
48. A. K. Ghose, G. M. Crippen, Atomic Physicochemical Parameters for Three-Dimensional Structure-Directed
Quantitative Structure-Activity Relationships I. Partition Coefficients as a Measure of Hydrophobicity. *J. Comput.*
Chem. **7**, 565–577 (1986).
49. P. A. Clemons, N. E. Bodycombe, H. A. Carrinski, J. A. Wilson, A. F. Shamji, B. K. Wagner, A. N. Koehler, S. L.
Schreiber, Small molecules of different origins have distinct distributions of structural complexity that correlate with
protein-binding profiles. *Proc Natl Acad Sci USA*. **107**, 18787–18792 (2010).
50. J. N. Israelachvili, The nature of van der waals forces. *Contemp. Phys.* **15**, 159–178 (1974).
51. D. Van Vranken, G. A. Weiss, *Introduction to Bioorganic Chemistry and Chemical Biology* (Garland Science, ed. 1,
2012).
52. E. Pichersky, R. A. Raguso, Why do plants produce so many terpenoid compounds? *New Phytol.* **220**, 692–702
(2018).
53. D. L. Forrister, M.-J. Endara, A. J. Soule, G. C. Younkin, A. G. Mills, J. Lokvam, K. G. Dexter, R. T. Pennington, C. A.
Kidner, J. A. Nicholls, O. Loiseau, T. A. Kursar, P. D. Coley, Diversity and Divergence: Evolution of secondary
metabolism in the tropical tree genus Inga. *New Phytol.* (2022), doi:10.1111/nph.18554.
54. R. M. LoPachin, D. S. Barber, T. Gavin, Molecular mechanisms of the conjugated alpha,beta-unsaturated carbonyl
derivatives: relevance to neurotoxicity and neurodegenerative diseases. *Toxicol. Sci.* **104**, 235–249 (2008).
55. J. Widomska, W. K. Subczynski, Mechanisms enhancing the protective functions of macular xanthophylls in the retina
during oxidative stress. *Exp. Eye Res.* **178**, 238–246 (2019).
56. M. H. Tran, D.-P. Phan, E. Y. Lee, Review on lignin modifications toward natural UV protection ingredient for lignin-
based sunscreens. *Green Chem.* **23**, 4633–4646 (2021).
57. N.-H. Tan, J. Zhou, Plant cyclopeptides. *Chem. Rev.* **106**, 840–895 (2006).
58. G.-H. Lim, R. Singhal, A. Kachroo, P. Kachroo, Fatty Acid- and Lipid-Mediated Signaling in Plant Defense. *Annu. Rev.*
Phytopathol. **55**, 505–536 (2017).
59. T. H. Yeats, J. K. C. Rose, The formation and function of plant cuticles. *Plant Physiol.* **163**, 5–20 (2013).
60. Q. Liu, L. Luo, L. Zheng, Lignins: biosynthesis and biological functions in plants. *Int. J. Mol. Sci.* **19** (2018),
doi:10.3390/ijms19020335.
61. P. E. Staswick, Storage proteins of vegetative plant tissues. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **45**, 303–322
(1994).
62. T. O. Sandberg, C. Weinberger, J.-H. Smått, Molecular Dynamics on Wood-Derived Lignans Analyzed by
Intermolecular Network Theory. *Molecules*. **23** (2018), doi:10.3390/molecules23081990.
63. M. Pollard, F. Beisson, Y. Li, J. B. Ohlrogge, Building lipid barriers: biosynthesis of cutin and suberin. *Trends Plant*
Sci. **13**, 236–246 (2008).
64. R. Salguero-Gómez, O. R. Jones, E. Jongejans, S. P. Blomberg, D. J. Hodgson, C. Mbeau-Ache, P. A. Zuidema, H.
de Kroon, Y. M. Buckley, Fast-slow continuum and reproductive strategies structure plant life-history variation
worldwide. *Proc Natl Acad Sci USA*. **113**, 230–235 (2016).
65. P. D. Coley, in *Plant-Animal Interactions: Evolutionary Ecology in Tropical and Temperate Regions*, P. W. Price, T. M.
Lewison, G. W. Fernandes, W. W. Benson, Eds. (John Wiley and Sons, New York, 1991), pp. 54–69.
66. R. Sulpice, P. C. McKeown, Moving toward a comprehensive map of central plant metabolism. *Annu. Rev. Plant Biol.*
66, 187–210 (2015).
67. S. B. Powles, Photoinhibition of photosynthesis induced by visible light. *Annu. Rev. Plant Physiol.* **35**, 15–44 (1984).
68. J. Berry, O. Bjorkman, Photosynthetic response and adaptation to temperature in higher plants. *Annu. Rev. Plant*
Physiol. **31**, 491–543 (1980).
69. B. David, F. Aussiel, in *Handbook of Chemical and Biological Plant Analytical Methods*, K. Hostettmann, Ed. (Wiley,
Chichester, West Sussex, ed. 1, 2014), p. 1176.
70. B. David, F. Ausseil, in *Encyclopedia of analytical chemistry: applications, theory and instrumentation*, R. A. Meyers,
Ed. (John Wiley & Sons, Ltd, Chichester, UK, 2006), pp. 1–24.
71. T. Pluskal, S. Castillo, A. Villar-Briones, M. Oresic, MZmine 2: modular framework for processing, visualizing, and
analyzing mass spectrometry-based molecular profile data. *BMC Bioinformatics*. **11**, 395 (2010).
72. O. D. Myers, S. J. Sumner, S. Li, S. Barnes, X. Du, One Step Forward for Reducing False Positive and False
Negative Compound Identifications from Mass Spectrometry Metabolomics Data: New Algorithms for Constructing
Extracted Ion Chromatograms and Detecting Chromatographic Peaks. *Anal. Chem.* **89**, 8696–8703 (2017).
73. H. Tsugawa, T. Cajka, T. Kind, Y. Ma, B. Higgins, K. Ikeda, M. Kanazawa, J. VanderGheynst, O. Fiehn, M. Arita, MS-
DIAL: data-independent MS/MS deconvolution for comprehensive metabolome analysis. *Nat. Methods*. **12**, 523–526
(2015).
74. A. M. Frank, N. Bandeira, Z. Shen, S. Tanner, S. P. Briggs, R. D. Smith, P. A. Pevzner, Clustering millions of tandem
mass spectra. *J. Proteome Res.* **7**, 113–122 (2008).
75. M. Wang, J. J. Carver, V. V. Phelan, L. M. Sanchez, N. Garg, Y. Peng, D. D. Nguyen, J. Watrous, C. A. Kapono, T.
Luzzatto-Knaan, C. Porto, A. Bouslimani, A. V. Melnik, M. J. Meehan, W.-T. Liu, M. Crüsemann, P. D. Boudreau, E.
Esquenazi, M. Sandoval-Calderón, R. D. Kersten, L. A. Pace, R. A. Quinn, K. R. Duncan, C.-C. Hsu, D. J. Floros, R.
G. Gavilan, K. Kleigrewe, T. Northen, R. J. Dutton, D. Parrot, E. E. Carlson, B. Aigle, C. F. Michelsen, L. Jelsbak, C.
Sohlenkamp, P. Pevzner, A. Edlund, J. McLean, J. Piel, B. T. Murphy, L. Gerwick, C.-C. Liaw, Y.-L. Yang, H.-U.

Humpf, M. Maansson, R. A. Keyzers, A. C. Sims, A. R. Johnson, A. M. Sidebottom, B. E. Sedio, A. Klitgaard, C. B.
Larson, C. A. B. P, D. Torres-Mendoza, D. J. Gonzalez, D. B. Silva, L. M. Marques, D. P. Demarque, E. Pociute, E. C.
O'Neill, E. Briand, E. J. N. Helfrich, E. A. Granatosky, E. Glukhov, F. Ryffel, H. Houson, H. Mohimani, J. J. Kharbush,
Y. Zeng, J. A. Vorholt, K. L. Kurita, P. Charusanti, K. L. McPhail, K. F. Nielsen, L. Vuong, M. Elfeki, M. F. Traxler, N.
Engene, N. Koyama, O. B. Vining, R. Baric, R. R. Silva, S. J. Mascuch, S. Tomasi, S. Jenkins, V. Macherla, T.
Hoffman, V. Agarwal, P. G. Williams, J. Dai, R. Neupane, J. Gurr, A. M. C. Rodríguez, A. Lamsa, C. Zhang, K.
Dorrestein, B. M. Duggan, J. Almaliti, N. Bandeira, Sharing and community curation of mass spectrometry data with
Global Natural Products Social Molecular Networking. *Nat. Biotechnol.* **34**, 828–837 (2016).

76.   F. Chassagne, G. Cabanac, G. Hubert, B. David, G. Marti, The landscape of natural product diversity and their
pharmacological relevance from a focus on the Dictionary of Natural Products®. *Phytochem. Rev.*, 1–22 (2019).

77.   P.-M. Allard, T. Péresse, J. Bisson, K. Gindro, L. Marcourt, V. C. Pham, F. Roussi, M. Litaudon, J.-L. Wolfender,
Integration of Molecular Networking and In-Silico MS/MS Fragmentation for Natural Products Dereplication. *Anal.*
Chem. **88**, 3317–3323 (2016).

78.   A. Rutz, M. Dounoue-Kubo, S. Ollivier, J. Bisson, M. Bagheri, T. Saesong, S. N. Ebrahimi, K. Ingkaninan, J.-L.
Wolfender, P.-M. Allard, Taxonomically informed scoring enhances confidence in natural products annotation. *Front.*
Plant Sci. **10**, 1329 (2019).

79.   R. Guha, Chemical Informatics Functionality in *R. J. Stat. Softw.* **18** (2007), doi:10.18637/jss.v018.i05.

80.   R Core Team, *R: A language and environment for statistical computing* (R Foundation for Statistical Computing,
Vienna, Austria, 2014).

81.   C. O. Wilke, *cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2"* (R Project, 2019).

82.   M. Dowle, A. Srinivasan, *data.table: Extension of `data.frame`* (R Project, 2021).

83.   W. Michael Landau, The drake R package: a pipeline toolkit for reproducibility and high-performance computing.
JOSS. **3**, 550 (2018).

84.   D. Vaughan, M. Dancho, *furrr: Apply Mapping Functions in Parallel using Futures* (R Project, 2021).

85.   H. Bengtsson, A unifying framework for parallel and distributed processing in R using futures. *arXiv* (2021).

86.   W. R. Revelle, *psych: Procedures for Personality and Psychological Research* (Northwestern University, 2019).

87.   H. Wickham, M. Averick, J. Bryan, W. Chang, L. McGowan, R. François, G. Grolemund, A. Hayes, L. Henry, J.
Hester, M. Kuhn, T. Pedersen, E. Miller, S. Bache, K. Müller, J. Ooms, D. Robinson, D. Seidel, V. Spinu, K.
Takahashi, D. Vaughan, C. Wilke, K. Woo, H. Yutani, Welcome to the tidyverse. *JOSS.* **4**, 1686 (2019).

88.   S. A. Chamberlain, E. Szöcs, taxize: taxonomic search and retrieval in R. [version 2; peer review: 3 approved].
F1000Res. **2**, 191 (2013).

89.   W. Cornwell, R. FitzJohn, M. Pennell, *taxonlookup: A dynamically-updating versioned taxonomic resource for
vascular plants* (R Project, 2016).

90.   J. Kattge, G. Bönisch, S. Díaz, S. Lavorel, I. C. Prentice, P. Leadley, S. Tautenhahn, G. D. A. Werner, T. Aakala, M.
Abedi, A. T. R. Acosta, G. C. Adamidis, K. Adamson, M. Aiba, C. H. Albert, J. M. Alcántara, C. Alcázar C, I. Aleixo, H.
Ali, B. Amiaud, C. Ammer, M. M. Amoroso, M. Anand, C. Anderson, N. Anten, J. Antos, D. M. G. Apgaua, T.-L.
Ashman, D. H. Asmara, G. P. Asner, M. Aspinwall, O. Atkin, I. Aubin, L. Baastrup-Spohr, K. Bahalkeh, M. Bahn, T.
Baker, W. J. Baker, J. P. Bakker, D. Baldocchi, J. Baltzer, A. Banerjee, A. Baranger, J. Barlow, D. R. Barneche, Z.
Baruch, D. Bastianelli, J. Battles, W. Bauerle, M. Bauters, E. Bazzato, M. Beckmann, H. Beeckman, C. Beierkuhnlein,
R. Bekker, G. Belfry, M. Belluau, M. Beloiu, R. Benavides, L. Benomar, M. L. Berdugo-Lattke, E. Berenguer, R.
Bergamin, J. Bergmann, M. Bergmann Carlucci, L. Berner, M. Bernhardt-Römermann, C. Bigler, A. D. Bjorkman, C.
Blackman, C. Blanco, B. Blonder, D. Blumenthal, K. T. Bocanegra-González, P. Boeckx, S. Bohlman, K. Böhning-
Gaese, L. Boisvert-Marsh, W. Bond, B. Bond-Lamberty, A. Boom, C. C. F. Boonman, K. Bordin, E. H. Boughton, V.
Boukili, D. M. J. S. Bowman, S. Bravo, M. R. Brendel, M. R. Broadley, K. A. Brown, H. Bruelheide, F. Brumnich, H. H.
Bruun, D. Bruy, S. W. Buchanan, S. F. Bucher, N. Buchmann, R. Buitenwerf, et al., TRY plant trait database -
enhanced coverage and open access. *Glob. Chang. Biol.* **26**, 119–188 (2020).

91.   J. P. Grime, Vegetation classification by reference to strategies. *Nature.* **250**, 26–31 (1974).

92.   F. Schrodt, J. Kattge, H. Shan, F. Fazayeli, J. Joswig, A. Banerjee, M. Reichstein, G. Bönisch, S. Díaz, J. Dickie, A.
Gillison, A. Karpatne, S. Lavorel, P. Leadley, C. B. Wirth, I. J. Wright, S. J. Wright, P. B. Reich, BHPMF - a
hierarchical Bayesian approach to gap-filling and trait prediction for macroecology and functional biogeography.
Global Ecology and Biogeography. **24**, 1510–1521 (2015).

93.   S. van Buuren, K. Groothuis-Oudshoorn, mice : Multivariate Imputation by Chained Equations in *R. J. Stat. Softw.* **45**
(2011), doi:10.18637/jss.v045.i03.

94.   C. Maldonado, C. I. Molina, A. Zizka, C. Persson, C. M. Taylor, J. Albán, E. Chilquillo, N. Rønsted, A. Antonelli,
Estimating species diversity and distribution in the era of Big Data: to what extent can we trust public databases?
Glob. Ecol. Biogeogr. **24**, 973–984 (2015).

95.   A. Zizka, D. Silvestro, T. Andermann, J. Azevedo, C. Duarte Ritter, D. Edler, H. Farooq, A. Herdean, M. Ariza, R.
Scharn, S. Svanteson, N. Wengström, V. Zizka, A. Antonelli, CoordinateCleaner: standardized cleaning of occurrence
records from biological collection databases. *Methods Ecol. Evol.* (2019), doi:10.1111/2041-210X.13152.

96.   D. S. Maynard, L. Bialic-Murphy, C. M. Zohner, C. Averill, J. van den Hoogen, H. Ma, L. Mo, G. R. Smith, A. T. R.
Acosta, I. Aubin, E. Berenguer, C. C. F. Boonman, J. A. Catford, B. E. L. Cerabolini, A. S. Dias, A. González-Melo, P.
Hietz, C. H. Lusk, A. S. Mori, Ü. Niinemets, V. D. Pillar, B. X. Pinho, J. A. Rosell, F. M. Schurr, S. N. Sheremetev, A.
C. da Silva, Ê. Sosinski, P. M. van Bodegom, E. Weiher, G. Bönisch, J. Kattge, T. W. Crowther, Global relationships in
tree functional traits. *Nat. Commun.* **13**, 3185 (2022).

97.   J. Oksanen, F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R. Minchin, R. B. O'Hara, G. L.
596   Simpson, P. Solymos, M. H. H. Stevens, E. Szoecs, H. Wagner, *vegan: Community Ecology Package* (R Project,
597   2019).
598   98.   W. Weckwerth, Metabolomics in systems biology. *Annu. Rev. Plant Biol.* **54**, 669–689 (2003).
599   99.   H. Uthe, N. M. van Dam, M. R. Hervé, M. Sorokina, K. Peters, A. Weinhold, in *Plant Metabolomics in full swing*
600   (Elsevier, 2021), vol. 98 of *Advances in botanical research*, pp. 163–203.
601   100.   K. Habel, R. Grasman, R. B. Gramacy, P. Mozharovskyi, D. C. Sterrat, *geometry: Mesh Generation and Surface
602   Tessellation* (R, 2022).
603   101.   S. Dray, A.-B. Dufour, The ade4 Package: Implementing the Duality Diagram for Ecologists. *J. Stat. Softw.* **22** (2007),
604   doi:10.18637/jss.v022.i04.
605   102.   R. Salguero-Gómez, Applications of the fast-slow continuum and reproductive strategy framework of plant life
606   histories. *New Phytol.* **213**, 1618–1624 (2017).
607   103.   H. Kusano, H. Li, H. Minami, Y. Kato, H. Tabata, K. Yazaki, Evolutionary developments in plant specialized
608   metabolism, exemplified by two transferase families. *Front. Plant Sci.* **10**, 794 (2019).
609   104.   R. T. Bush, F. A. McInerney, Leaf wax n-alkane distributions in and across modern plants: Implications for
610   paleoecology and chemotaxonomy. *Geochim. Cosmochim. Acta*. **117**, 161–179 (2013).
611   105.   L. F. Maia, V. E. De Oliveira, H. G. M. Edwards, L. F. C. De Oliveira, The diversity of linear conjugated polyenes and
612   colours in nature: raman spectroscopy as a diagnostic tool. *ChemPhysChem*. **22**, 231–249 (2021).
613   106.   J. K. Hoober, L. L. Eggink, M. Chen, Chlorophylls, ligands and assembly of light-harvesting complexes in chloroplasts.
614   *Photosyn. Res.* **94**, 387–400 (2007).
615   107.   D. F. Veber, S. R. Johnson, H.-Y. Cheng, B. R. Smith, K. W. Ward, K. D. Kopple, Molecular properties that influence
616   the oral bioavailability of drug candidates. *J. Med. Chem.* **45**, 2615–2623 (2002).
617   108.   P. Ertl, B. Rohde, P. Selzer, Fast calculation of molecular polar surface area as a sum of fragment-based
618   contributions and its application to the prediction of drug transport properties. *J. Med. Chem.* **43**, 3714–3717 (2000).
619   109.   K. Palm, P. Stenberg, K. Luthman, P. Artursson, Polar molecular surface properties predict the intestinal absorption of
620   drugs in humans. *Pharm. Res.* **14**, 568–571 (1997).
621   110.   G. A. Jeffrey, W. Saenger, *Hydrogen bonding in biological structures* (Springer Berlin Heidelberg, Berlin, Heidelberg,
622   1991).

Acknowledgements

Special thanks to Jeanne Clément and Ghislain Vieilledent for building bespoke functionality into the R package "jSDM" for exploratory analyses.

Funding: This study was part funded by a grant from the Synthesis Centre (sDiv) of the German Centre for Integrative Biodiversity Research (iDiv), awarded to TWNW, FS and NMvD. Plant material collection and metabolomics analysis were financed by Swiss National Science Foundation grants awarded to LP (grant no. 31003A-162604), SR (grant no. 31003A-131956), and JLW (grant no. CRSII5-189921/1). JMA is funded by the Swiss National Science Foundation (grant no. 31003A-176044). FS acknowledges funding by the University of Nottingham Anne McLaren Fellowship. ED and SR are funded by the Swiss National Science Foundation (grant no. 31003A_179481). VEJJ is funded by the French National Research Agency (grant no. ANR-17-CE01-0007: Mixxopeat). VB is supported by the French Agence Nationale pour la Recherche (grant no. ANR-12-BSV7-0016-01: SecPriMe2), the BioDivMeX Mistrals program, the Aix-Marseille University 'Investissements d'Avenir' program and the Labex OT-Med programs. PC is supported by a Ramón Areces Foundation Postdoctoral Scholarship. PDC is funded by the US National Science Foundation (grant no. DEB-1135733). NMvD, SM and KP are supported by iDiv Halle-Jena-Leipzig, which is funded by the German Research Foundation (grant nos DFG–FZT 118, 202548816). SN and KP are additionally funded by the German Network for Bioinformatics Infrastructure (de.NBI) and acknowledge BMBF funding (grant no. 031L0107). DLF is supported by the US National Science Foundation Graduate Research Fellowships Program. RSG is funded by a UK Natural Environment Research Council Independent Research Fellowship (grant no. NE/M018458/1). JP and JS are funded by the Spanish Government (grant no. CGL2016-79835), Catalan Government (grant no. SGR 2017-1005) and a European Research Council Synergy Grant (grant no. ERC-SyG-2013-610028: IMBALANCE-P). MCS is supported by a NOMIS Foundation project ("Remotely Sensing Ecological Genomics") and by the University of Zürich Research Priority Program on Global Change and Biodiversity.

Authors contributions: Conceptualization: TWNW in collaboration with PMA, RDB, PC, BD, ED, AG-G, VEJJ, JP, SR, RSG, JS, MCS, FS, NMvD and WW, with support from all co-authors. Funding acquisition: FS, NMvD, and TWNW. Data collection: PMA, ED, and BD. Data curation: TWNW, with support from PMA, OB, PC, and ED. Formal analysis: TWNW, with support from PC, ED, MJE, DF, and VEJJ. Writing: TWNW, with support from MCS. Reviewing and editing: all co-authors.

Competing interests: The authors declare no competing interests.

Data and materials availability: Raw tropical metabolomics data are available under the DOI 10.25345/C59J97 (*36*). Raw temperate species metabolomics data are available under the DOI 10.6084/m9.figshare.13032740 (*37*). Flat data and R code is available under the DOI 10.17605/OSF.IO/YZR4C.

Supplementary Materials
Materials and Methods
Figs. S1 to S5
Tables S1 to S3