LeishMANIAdb: a comparative resource for *Leishmania* proteins

Gábor E. Tusnády[1,2], **András Zeke**[1], **Zsófia E. Kálmán**[3], **Marie Fatoux**[4,5], **Sylvie Ricard-Blum**[4], **Toby J. Gibson**[6], **Laszlo Dobson**[1,6,*]

[1]Protein Bioinformatics Research Group, Institute of Enzymology, Research Centre for Natural Sciences, Magyar Tudósok körútja 2, Budapest, 1117, Hungary; [2]Department of Bioinformatics, Semmelweis University, Tűzoltó u. 7, Budapest, 1094, Hungary; [3]Faculty of Information Technology and Bionics, Pázmány Péter Catholic University, Práter u. 50/A, Budapest, 1083 Hungary; [4]ICBMS UMR CNRS 5246, University Lyon 1, Rue Victor Grignard, Villeurbanne, Lyon, 69622, France; [5]The work was done in UMR CNRS 5086, University Lyon 1, 7 Passage du Vercors, Lyon, 69367, France; [6]Structural and Computational Biology Unit, European Molecular Biology Laboratory, Meyerhofstraße 1, Heidelberg, 69117, Germany

*For correspondence:
laszlo.dobson@embl.de (FMS);
dobson.laszlo@ttk.hu (FS)

Abstract   Leishmaniasis is a detrimental disease causing serious changes in quality of life and some forms lead to death. The disease is spread by the parasite *Leishmania* transmitted by sandfly vectors and their primary hosts are vertebrates including humans. The pathogen penetrates host cells and secretes proteins (the secretome) to repurpose cells for pathogen growth and to alter cell signaling via host-pathogen Protein-Protein Interactions (PPIs). Here we present LeishMANIAdb, a database specifically designed to investigate how *Leishmania* virulence factors may interfere with host proteins. Since the secretomes of different *Leishmania* species are only partially characterized, we collected various experimental evidence and used computational predictions to identify *Leishmania* secreted proteins to generate a user-friendly unified web resource allowing users to access all information available on experimental and predicted secretomes. In addition, we manually annotated host-pathogen interactions of 211 proteins, and the localization/function of 3764 transmembrane (TM) proteins of different *Leishmania* species. We also enriched all proteins with automatic structural and functional predictions that can provide new insights in the molecular mechanisms of infection. Our database, available at https://leishmaniadb.ttk.hu may provide novel insights into *Leishmania* host-pathogen interactions and help to identify new therapeutic targets for this neglected disease.

Introduction

Leishmaniasis is a neglected tropical disease causing severe symptoms, affecting around 1 million new people yearly, with annual deaths estimated to be around 60,000 *Torres et al.* (*2017*). Although over 90% of cases occur in poor regions south of the Equator, due to climatic changes it also appears in new areas, and it has already shown up in Mediterranean European countries *Gianchecchi and Montomoli* (*2020*) and Texas, USA *McIlwee et al.* (*2018*). To this date no approved human vaccine is available and treatment is most effective at an early stage of the infection. *Leishmania* parasites are unicellular, flagellated trypanosomatids, belonging to the class Kinetoplastea. Upon infection, the amastigote stage pathogen (with reduced flagella) is engulfed by phagocytes, where it ends up in a stable parasitophorous vacuole that protects it *Arango Duque et al.* (*2019*). *Leishmania*

cells then proliferate unhindered within host cells until egress and spreading to nearby phagocytes *Real et al.* (*2014*). The parasite secretes proteins that enter various parts of the cell *Atayde et al.* (*2015*). The secreted virulence factors can then interfere with cell signaling by interacting with the host proteins: they increase glycolytic metabolism *Ohms et al.* (*2021*), perturb microbicidal pathways *Matheoud et al.* (*2013*), escape the innate immune response, and repurpose macrophages for parasite replication *Atayde et al.* (*2016*) by disturbing with cellular protein-protein interactions (PPIs). Interestingly, these mechanisms are somewhat unique to *Leishmania* among trypanosomes, which are usually extracellular pathogens and do not enter host cells. In contrast, *Leishmania* secretes proteins which are critical for host cell subjugation, but how they enter the cytoplasm of host cells is still poorly understood.

The host targeted interactions are often mediated via Short Linear Motifs (SLiMs) in many distant, unrelated intracellular pathogens, ranging from viruses and bacteria to unicellular eukaryotes *Davey et al.* (*2011*). SLiMs are flexible protein segments composed of a restricted number of residues (between 3-10), that usually bind to structured protein domains. Their short length and structural flexibility enable them to bind to a wide range of domains. Cellular SLiMs typically bind their targets with low micromolar affinity. These weak and transient interactions enable SLiMs to work in cooperative regulatory systems *Van Roey et al.* (*2014*). Pathogens mimic host SLiMs to interact with host cell proteins *Davey et al.* (*2011*). Pathogen SLiMs often bind with higher affinities than the cellular ones, outcompeting the native interactions, permanently re-wiring the host regulation network. A few modulatory SLiMs have already been discovered in eukaryotic pathogens, such as the Toxoplasma gondii MapK docking motif *Pellegrini et al.* (*2017*) and the stage-specific (promastigote-amastigote) phosphorylation motifs from *Leishmania Tsigankov et al.* (*2013*). In addition, several putative SLiMs were recently detected in *Leishmania* such as heparin-binding sequences or RGD integrin-binding motifs but their function has not been confirmed yet *Peysselon et al.* (*2013*).

Numerous studies investigated *Leishmania* secretomes. Most of them expose promastigotes to a heat shock and pH change (attempting to emulate the conditions that promote promastigote-to-amastigote stage transition) and then analyze the *Leishmania* conditioned medium by proteomics to identify secreted proteins *Cuervo et al.* (*2009*), and measure their protein abundance or by transcriptomics to detect mRNA levels *Lahav et al.* (*2011*). While high-throughput experiments inherently suffer from a certain level of noise, experiments on individual proteins may be more reliable - in the case of *Leishmania* the vast majority focuses on leishmanolysin (GP63), a surface-anchored protease important for pathogenesis *Gregory et al.* (*2008*); *Guay-Vincent et al.* (*2022*). Furthermore, data were collected on different *Leishmania* species/strains identified via names and identifiers varying from one source to another, making a unified overview challenging. Another key step towards understanding the infection mechanism would be the identification of *Leishmania* surface proteins that can mediate the attachment of the pathogen to the host cell. Some surfaceome experiments were carried out on *Leishmania*-related species, and human host proteins binding to the surface of 24 strains of intact *Leishmania* have been identified *Fatoux-Ardore et al.* (*2014*). Beside the characterization of *Leishmania* secretomes, the identification of host-*Leishmania* PPIs is needed to narrow down virulence factors perturbing the host cell regulation to modules interfering with host proteins. SLiMs have low information content and simply scanning them in *Leishmania* secretomes may yield many false positives. Their structural and functional context, such as accessibility, conservation and localization, are all key elements to successfully identify those that may have a role in rewiring the host cell regulation. Notably, SLiMs also play a key role in maintaining housekeeping processes in *Leishmania*, therefore to find candidate SLiMs that may alter the host regulation, we need to discriminate SLiMs of proteins that reach the host cytoplasm or nucleus but limited information about these proteins are available. Currently the only publicly available database dealing with *Leishmania* proteins is TriTrypDB *Shanmugasundram et al.* (*2023*), which is part of the VEuPathDB *Amos et al.* (*2022*). TriTrypDB is a functional genomic resource for Trypanosomatidae, offering proteomic datasets, however it does not focus on protein structure,

₉₃ protein motif search and interactions.

₉₄ We developed LeishMANIAdb to expedite *Leishmania* research by unifying scattered informa-
₉₅ tion from the literature in a user-friendly way and to extend available resources by adding protein
₉₆ level information. We collected high-throughput experiments and interaction studies on individual
₉₇ proteins, and used various prediction methods to enrich proteins with structural information.

₉₈ Results

₉₉ Selection of *Leishmania* proteomes and homology mapping of various kinetoplastid
₁₀₀ proteins

₁₀₁ We selected 5 *Leishmania* species (reference proteomes: *L. brazliensis, L. donovani, L. infantum, L.*
₁₀₂ *major, L. mexicana*), 13 Leishmania strains (*Lbraziliensis MHOMBR75M2903*, *Lbraziliensis MHOMBR75M2904*,
₁₀₃ *Lbraziliensis MHOMBR75M29042019*, *Ldonovani BPK282A1*, *LdonovaniCL-SL*, *Ldonovani HU3*, *Ldono-*
₁₀₄ *vani LV*, *Linfantum JPCM5*, *LmajorFriedlin*, *Lmajor Friedlin2021*, *Lmajor LV39c5*, *Lmajor SD75.1*, *Lmexi-*
₁₀₅ *cana MHOMGT2001U1103*), and 6 related species (reference proteomes: *Bodo saltans, Leptomonas*
₁₀₆ *seymouri, Trypansoma brucei, Trypansoma cruzi, Trypansoma rangeli, Trypanosoma theileri*) as an out-
₁₀₇ group. The 5 *Leishmania* proteomes and the 6 related kinetoplastid proteomes were selected
₁₀₈ based on their quality (i.e. number of fragments and missing proteins) and were downloaded
₁₀₉ from UniProtKB *UniProt Consortium* (*2023*). *Leishmania* proteins were also cross-referenced to
₁₁₀ TriTrypDB *Shanmugasundram et al.* (*2023*). Around 30% of the cross-referenced proteins have dif-
₁₁₁ ferent sequences deposited into these resources, and in most cases the difference is due to the
₁₁₂ position of the initiator methionine. For data compatibility we always use the UniProt sequence
₁₁₃ version but the conflicts are highlighted in LeishMANIAdb. We also performed a similarity search
₁₁₄ between these proteins and linked close homologs (see Methods) so annotations and predictions
₁₁₅ can be easily compared between them. All manual annotations and experimental data from dif-
₁₁₆ ferent sources were mapped to these proteins. The 13 *Leishmania* strain proteomes were down-
₁₁₇ loaded from TriTrypDB. Altogether LeishMANIAdb contains 40 537 searchable *Leishmania* proteins
₁₁₈ from reference proteomes, 108 766 proteins from different strains and 68 924 other kinetoplastid
₁₁₉ proteins to strengthen predictions.

₁₂₀ Manual annotation of host-pathogen PPIs and TM protein localization

₁₂₁ We manually curated hundreds of proteins, using two strategies. The first type of annotation was
₁₂₂ the collection of host-pathogen PPI experiments on individual proteins, with the majority of them
₁₂₃ involving leishmanolysin (GP63). We collected 29 papers reporting 82 *Leishmania* PPIs with dif-
₁₂₄ ferent hosts. Although experiments were mapped back to specific proteins, the results are also
₁₂₅ displayed on close homologs (with a note that the experimental data is derived from a different
₁₂₆ protein) resulting in 211 proteins that contain PPI data. Interactions were reported using the Mini-
₁₂₇ mum Information required for reporting a Molecular Interaction eXperiment MIMIx *Orchard et al.*
₁₂₈ (*2007*) community standard description. The second type of manually curated data was the local-
₁₂₉ ization and functional annotation of TM proteins. The aim was to find surface proteins that may
₁₃₀ facilitate the infection, but we annotated hundreds of other TM proteins with their localizations
₁₃₁ too. For this task, we used all close homologous proteins defined in the previous step. Altogether
₁₃₂ 342 protein families were annotated and these annotations were shared between 3764 proteins
₁₃₃ (which is 45.11% percent of the predicted TM proteomes, and 9.28% of all proteins of the 5 species
₁₃₄ combined).

₁₃₅ The definition of *Leishmania* secretome and protein localization is still incomplete

₁₃₆ *Leishmania* not only exploits host-secretory pathways to distribute effectors but also utilizes an
₁₃₇ unusual mechanism to deliver proteins to the cytosol of infected cells by releasing exosomes into
₁₃₈ the parasitophorous vesicle, which might fuse with the vesicular membrane to release their pro-
₁₃₉ tein content *Silverman et al.* (*2010*). Therefore computational methods based on signal peptides

140 and localization predictions are not sufficient to predict *Leishmania* secretomes. To overcome this
141 limitation we also used high-throughput experiments *Silverman et al.* (*2008*); *Cuervo et al.* (*2009*);
142 *Hassani et al.* (*2011*); *Forrest et al.* (*2020*); *Pissarra et al.* (*2022*) to increase the coverage of *Leishma-*
143 *nia* secretomes. Strikingly, the number of proteins in these secretomes varies to a large extent, and
144 some proteins cannot be identified by mass spectrometry. Other datasets include proteins found
145 in glycosomes *Jamdhade et al.* (*2015*), stage-dependent (promastigote/amastigote) phosphopro-
146 teomics *Tsigankov et al.* (*2013*), housekeeping gene localizations *Jardim et al.* (*2018*), exosome con-
147 tent *Silverman et al.* (*2010*), protein and mRNA abundance data *Lahav et al.* (*2011*); *Pescher et al.*
148 (*2011*). When we mapped back all secretome and abundance experiments to *Leishmania infantum*
149 (from orthologous proteins of other *Leishmania* species), the number of identified proteins ranges
150 from 10 to 2,000 (Figure 1/A), and even when experimental conditions were similar they yielded
151 highly different amounts of proteins. For example, pioneer secretome studies only provided a few
152 hundred hits, while the latest ones are more inclusive with thousands of hits. Gene duplication
153 is often acting on protein families responsible for host-pathogen PPIs, therefore we also collected
154 proteins that are highly expanded. Notably, as all kinetoplastids have a polycistronic transcrip-
155 tion system, the main way to amplify expression of critical proteins is through gene duplication.
156 Thereby highly expanded gene families can be directly mapped to functions critical for these para-
157 sites *Jackson et al.* (*2016*). In this case we could discriminate between proteins with already many
158 paralogs within kinetoplastids and *Leishmania*-exclusive amplified proteins. When we searched
159 for homologs of *Leishmania infantum* proteins, we found distinct amino acid transporter and co-
160 factor families already expanded in all kinetoplastids including *Leishmania*. In contrast, amastins,
161 leishmanolysin, 3'A2-related proteins, kinase-containing putative receptor proteins (and several
162 uncharacterized proteins) seemed to be highly abundant in *Leishmania* proteomes compared to
163 all kinetoplastids (Figure 1/B). Comparing complete proteomic datasets yielded only a small over-
164 lap. We defined 1) *Leishmania*_novelty proteins, which are proteins without close homologs in
165 SwissProt, without characterized Pfam domains, and expanded in *Leishmania infantum* (compared
166 to other kinetoplastids); 2) abundant proteins, which are proteins showing increased abundance
167 upon infection; 3) secreted proteins experimentally identified in at least two secretome experi-
168 ments. These definitions provided markedly different protein sets, with some overlap between
169 secreted and abundant proteins (611 proteins) and with only 22 proteins contained in all datasets
170 (Figure 1/C).

AlphaFold2 provides an alternative way estimate structural features

172 We used different methods to predict the structural features of proteins. Classical sequence-based
173 methods can detect globular domains *Paysan-Lafosse et al.* (*2023*), TM regions *Dobson et al.* (*2015*)
174 and intrinsically disordered regions (IDRs) *Erdős et al.* (*2021*). However, the use of AlphaFold2 (AF2)
175 *Jumper et al.* (*2021*) provides alternative ways to obtain structural information. In LeishMANIAdb
176 we used structures available in the AlphaFold database *Varadi et al.* (*2022b*) (however we could
177 not find 3192 proteins (6% of all *Leishmania* proteins in LeishMANIAdb)). We not only displayed
178 the predicted 3D structure of the proteins, but also information derived from the AF2 models such
179 as the secondary structures and the position of the lipid bilayer for membrane proteins using the
180 method introduced in the TMAlphaFold database *Dobson et al.* (*2023*). Although AF2 was originally
181 built to predict protein structure, the scientific community quickly realized it is as much (if not more)
182 efficient at predicting protein disorder *Akdel et al.* (*2022*). To analyze IDRs we displayed predicted
183 local distance difference test (pLDDT) values and relative surface accessibility from AF2. For IDR
184 prediction in TM proteins, we tailored MemDis *Dobson and Tusnády* (*2021*) to incorporate features
185 from AF2 instead of sequence-based predictors (see Methods).

Short linear motif candidates that may hijack host cell regulation

187 We scanned *Leishmania* proteins for SLiMs using the regular expressions stored in the Eukaryotic
188 Linear Motif (ELM) resource *Kumar et al.* (*2022*). Scanning SLiMs alone would mostly yield false pos-

Figure 1. LeishMANIAdb content. All data were calculated on *Leishmania infantum*. A: Number of proteins in different proteomic datasets (purple: promastigote secretome, red: amastigote secretome, orange: exosome, green: housekeeping genes, blue: higher protein abundance level upon infection). B: Number of kinetoplastid and *Leishmania* close homologs. Each dot represents a protein (red: at least 80% of close homologs are in *Leishmania*, blue: other proteins). Green circles represent distinctive groups. C: Overlap between abundant, secreted and *Leishmania* novelty proteins (for more detail see text). D: Distribution of all predicted SLiMs with different scores. Red marks candidate motifs above 0.85 cutoff (for more details see text).

itive hits, so we developed a scoring system that ranges from 0 to 1, and that takes into account most information we collected. We aimed to develop a scoring system where conservation and accessibility/disorder has a reasonably high weight, while keeping in mind that proteomic experiments and localization information are a good way to narrow down the potentially large number of false positive hits. Unfortunately, due to the lack of data, in the case of *Leishmania* it is not possible to construct a benchmark set to evaluate motif scores. We can still assume that a good starting point can be when most predictions and proteomic data agree. Considering *Leishmania infantum* alone, we detected over a million putative motifs, from which 1.21% had a score above 0.85, on 343 proteins (Figure 1/D).

The LeishMANIAdb web resource

To visualize all the collected and calculated information we developed an open-access resource. In LeishMANIAdb users can search for proteins using their UniProt Accession (AC), Entry name (formerly ID), gene name and protein name. We also provide several protein sets as examples to help users browsing the database. Currently proteins are sorted based on 1) species: L. braziliensis, L. donovani, L. infantum, L. major, and L. mexicana; 2) manual curation data; 3) experimental data: secreted proteins, protein abundance/mRNA level data, proteins with any kind of experimental data listed above; 4) computationally predicted information: proteins expanded in *Leishmania* (score >= 0.8 - see Methods, Supplementary Material), transmembrane proteins, proteins with high disordered content (at least 70% predicted disorder), proteins with high-scoring linear motifs (score >= 0.85) and novel kinetoplastid proteins (proteins without SwissProt close homologs or Pfam domains). After searching (or selecting a protein set) users can further narrow their selection by choosing any other criterion (Figure 2/A).

The entry page for proteins consists of 10 sections, which are only visible if they contain data. The "Quick info" displays the protein name, species, cross-references and its number of amino acid residues. Data curation appears under the "Annotations" section. PPI (curated at the MIMIx level), localization and function annotations are mirrored from close homologous proteins. We also display functional annotations by Jardim et al. *Jardim et al.* (*2018*). The "Localization" section contains high-throughput experiment data - promastigote and amastigote secretion, an exosome experiment and the glycosome. Protein localization, signal peptide and glycosylphosphatidyl (GPI) anchor predictions are also displayed here. Since the reliability of both predictions and experiments may vary, we also display all this data for close homologous proteins, so users can quickly check the robustness of information. Furthermore, we also collected Gene_Ontology (GO) *The Gene Ontology Consortium* (*2019*) annotations for cellular compartments. In this case, the specificity of the term (how deep it is on the tree) is shown in the level column. GO annotations are collected for all close homologous proteins too, and the number of occurrences of each term is displayed. We highlighted terms that are associated with the inspected protein itself that is displayed on the page. The "Abundance" module can display the mRNA and protein level experiments: static/single point (upregulated or not) or time-course experiments (e.g., mRNA and protein levels available for 7 timesteps across 120 hours, Figure 2/B). In the "Expansion" section the number of close homologs are displayed by species with a color-code to identify *Leishmania* intracellular and extracellular/free-living relatives (Figure 2/C). The "Sequence features" displays various information (Figure 2/D). On the top, the gapless multiple sequence alignment (MSA) of proteins from the reference proteomes is visible. In this alignment gaps from the entry protein were removed (the original alignment with all strains can be downloaded) so other protein features could be visualized. Protein disorder, secondary structures, transmembrane topology prediction, domains, signal peptides and GPI anchors and stage-dependent phosphorylation are also displayed. Predicted SLiMs are shown with a color-coded score (see Methods, Supplementary Material). In the "Structure" section the AF2 predicted structure is available (with the position of the membrane domain for TM proteins). The "Putative motif mimicry" section is the table format version of SLiMs from the "Sequence features" module (Figure 2/E). The "Function" section contains GO Molecular Function and Biological Process terms.

Figure 2. Figure 2: Layout of LeishMANIAdb: A: the search/browse result menu. B: Expression section of the entry page. C: Expansion section. D: Sequence features section. E: Putative motif mimicry section. F: Function section.

As done for the Cellular Component, terms are mirrored from close homologous proteins, and they can be sorted based on their specificity (how deep they are located in the tree) and occurrences considering homologs (Figure 2/F). Finally, each result from a BLAST search against SwissProt and kinetoplastids relatives are listed in the "Homologs" table.

For each protein the full MSA, high-throughput experiments, annotation and predicted sequential features can be downloaded from the bottom of the page. Batch download is also available to download the full database or different protein sets.

Discussion

Reliability of data

In LeishMANIAdb we aimed to collect high-throughput experimental data, PPI data on individual proteins, predictions and localization information based on distant homologs. We noticed that the amount of data from MS experiments differs highly, and therefore likely the quality also varies. Not unexpected from high-throughput techniques applied to less-studied organisms, this can be attributed to the quality of sample preparation, MS experimental techniques and most likely to the sequences in the background databases. One striking finding was that the secretory datasets contain a large number of proteins that are likely to take part in the housekeeping processes of

²⁵⁵ *Leishmania* cells, such as cytoskeletal proteins, nuclear histones and metabolic enzymes. Exosomes
²⁵⁶ are known to contain a relatively high amount of „background" proteins leaking from the cytosol
²⁵⁷ of cells. Another explanation is that several housekeeping genes (such as intracellular chaperones
²⁵⁸ and enzymes) are moonlighting proteins, they are generally constitutively expressed and have high
²⁵⁹ levels of expression, while they are fulfilling other functions outside the cells *Jeffery* (*2018*). Due
²⁶⁰ to the lack of comparative studies, we cannot assess the enrichment ratios of secreted molecules,
²⁶¹ to see if there is selective exosomal packaging of a well-defined subset of *Leishmania* proteins.
²⁶² However, *Leishmania* exosomal-like secretion also differs from the typical exosomal sorting seen
²⁶³ in other eukaryotic organisms because budding primarily initiates at the cell membrane, and not
²⁶⁴ inside multivesicular bodies (endosomes). Therefore it is equally possible that in Leishmaniids, the
²⁶⁵ budding is non-selective for its cytoplasmic cargos. Instead it would be initiated by cell surface
²⁶⁶ receptors and primarily serve as a defense mechanism against membrane-attached host comple-
²⁶⁷ ment and other immune complexes, removing them before they could damage the parasite mem-
²⁶⁸ brane. Currently, testing of the latter hypothesis is impossible, since only soluble components but
²⁶⁹ not the integral membrane proteins of *Leishmania* exosomes have been studied in depth in the
²⁷⁰ above cited studies.

²⁷¹ From a computational point of view, predicting any features on *Leishmania* proteins might be
²⁷² highly challenging, as methods established were mostly trained on sequences that show little or
²⁷³ no similarity to *Leishmania* proteins. The 5 *Leishmania* reference proteomes contain 10 267 unchar-
²⁷⁴ acterized proteins combined, which is 25% of LeishMANIAdb. TMAlphaFold provides an objective
²⁷⁵ quality measurement option for alpha-helical membrane proteins. When we compared the TM
²⁷⁶ proteome of Homo sapiens and *Leishmania infantum* we noticed that the ratio of good and excel-
²⁷⁷ lent quality structures was much lower in *Leishmania*, probably caused by the different coverage
²⁷⁸ of kinetoplastid and human structures deposited into the PDB (Figure 3/A).

Case studies

²⁸⁰ LeishMANIAdb can be utilized for different purposes, and can be a good starting point for various
²⁸¹ analyses. We selected three examples that highlight some use cases of the resource.

²⁸² Using the Browse menu, after selecting a category users can further narrow down their search
²⁸³ for proteins selecting additional categories to refine the results. For instance, if users are looking
²⁸⁴ for *Leishmania* SLiMs that may alter or rewire host cell regulation network, they can look for pro-
²⁸⁵ teins that were experimentally proven to be secreted, and then select proteins with disordered re-
²⁸⁶ gions because SLiMs are mostly located in IDRs. "Kinetoplastid novelty" selection ensures that the
²⁸⁷ protein and its domains are not present in other organisms, while *Leishmania* novelty/expansions
²⁸⁸ select proteins that are new or highly expanded in *Leishmania* species. Last, by selecting high-
²⁸⁹ scoring motifs users get a list of proteins where the motif is most likely to be functional (Figure 3/B
²⁹⁰ shows the Venn diagram of the selection). These proteins may be an interesting starting point for
²⁹¹ further analyses.

²⁹² When performing systematic searches to identify possible parasite hits of integrin ligand motifs
²⁹³ (that only functions in the host, as kinetoplastids have no integrins), we identified a striking set of
²⁹⁴ examples in a family of poorly-known *Leishmania* genes called 3'A2 related ORFs. This kinetoplastid-
²⁹⁵ specific family of genes is actually expanded in *Leishmania* species together with the canonically se-
²⁹⁶ creted A2 proteins, which are known pathogenicity factors *Zhang and Matlashewski* (*2001*). While
²⁹⁷ the actual sequences of these proteins are poorly conserved and very little is known about their
²⁹⁸ subcellular location, the *Leishmania* versions have at least one transmembrane region and a C-
²⁹⁹ terminal cytoplasmic tail, with an N-terminal signal peptide (or possibly another TM segment).
³⁰⁰ Nevertheless, in the predicted, largely disordered extracellular segment we observed multiple,
³⁰¹ short, conserved stretches that may have amyloidogenic properties (high Val, Ala and Gly content,
³⁰² upon visual inspection), presumably capable of oligomerization and amphiphilic interaction with
³⁰³ membranes (Figure 3/C). A highly conserved cysteine residue preceding the first amyloidogenic
³⁰⁴ sequence might help the homodimerization by forming a disulphide bridge with neighboring 3'A2

related protein. Strikingly, in *Leishmania infantum* and *Leishmania donovani* (both species capable of causing visceral leishmaniasis), the N-terminus of these proteins carries canonical RGD (Arg-Gly-Asp) sequences, immediately after the putative signal peptide cleavage site. In addition, *Leishmania donovani* and *Leishmania infantum* proteins contain an NGR motif where asparagine deamidation might yield an isoDGR motif. If these proteins are expressed on the cell surface, they might bind to host integrins in an oligomeric state, and might even attack the host membrane as if it were a beta-barrel pore-forming toxin. However, much more experiments are needed to test any of these hypotheses.

Amastins are a large family of kinetoplastid-specific membrane proteins that belong to the broader claudin-like superfamily, implicated in the maintenance of parasitophorous vacuoles *de Paiva et al.* (*2015*). Accordingly, the majority of amastins have 4 tightly packed TM segments, with cytosolic tail regions. Similarly to their vertebrate counterparts that form tight cell-cell junctions by complex oligomerization processes, amastins might also engage in a variety of interactions with internal as well as external, host proteins. Although their exact function is not known, among the 221 identified amastins with 4 TM regions we looked for SLiMs that occur in multiple proteins. Predicted SLiMs (within disordered regions) were packed in their cytoplasmic tail regions (Figure 3/D). Since these regions face inward the parasite, we further narrowed hits based on their binding domain to be present in *Leishmania*. We identified multiple potential phosphorylation sites and protein-protein interaction motifs, such as SH3 ligands (*Leishmania* species do encode SH3 domain proteins) as well as vesicular trafficking signals. The tail region of amastins seem to be highly variable, likely acting as a hotspot in the pathogen-host arms race.

Comparison with other resources

In the past decades several databases were built to investigate *Leishmania*, however they are unfortunately often offline and no longer updated by now. LeishCyc *Saunders et al.* (*2012*) focused on biochemical pathways. LeishDB *Torres et al.* (*2017*) included coding genes and non-RNAs and provided new annotation to them. The cysteine protease database in *Leishmania* species *Rana et al.* (*2012*) was designed to find data related to cysteine protease and LeishBase was a structural database. There are a few active databases: Leish-ExP (http://www.hpppi.iicb.res.in/Leish-ex) (which has not so far been published in a peer-reviewed journal) contains proteins exclusively present in *Leishmania*. Leish-ExP incorporates localization tools, includes GO annotations and calculates physico-chemical properties of proteins. LmSmdB *Patel et al.* (*2016*) focuses on metabolic and biosynthetic pathways. TriTrypDB *Shanmugasundram et al.* (*2023*) is a kinetoplastid database that is part of the VEuPathDB resource *Amos et al.* (*2022*). These databases contain a lot of experimental data and various tools to analyze eukaryotic pathogens, but they are mostly focused on genomic data - although proteomic datasets, and some protein prediction algorithms are also incorporated.

There are also a handful of databases that include information on host-pathogen interactions: HPIDB *Ammari et al.* (*2016*), PHIDIAS *Xiang et al.* (*2007*) and PHI-base *Urban et al.* (*2022*) contain information about PPIs between the host and pathogen, while ImitateDB *Tayal et al.* (*2022*) specifically focuses on motif mimicry. These resources contain no or very little data about *Leishmania*.

In LeishMANIAdb our main goal was to include protein information relevant to the infection and to complement previously established and still available resources. We included several proteomic datasets, and enriched experimental information with state-of-the-art prediction tools. Still, the most powerful way to explore uncharted proteomes is to inspect MSAs and check for conserved residues and regions - LeishMANIAdb contains precalculated alignments for all proteins. We also added hundreds of annotations to thousands of proteins, including localization and interaction information. While several databases seem to be shut down after a couple of years, our laboratory hosts several resources and we routinely update them. We plan to do so with LeishMANIAdb as well as to expand its repertoire to host-*Leishmania* interactions involving glycans and glycolipids, which play major roles in the infection.

Figure 3. Figure 3: A: Distribution of membrane protein quality levels of AlphaFold structure in Homo sapiens and *Leishmania infantum*. B: Venn diagram of proteins that are 1) secreted 2) novel kinetoplastid 3) expanded (or new) in *Leishmania* 4) disordered 5) contain candidate SLiMs. C: left: Multiple Sequence Alignment of 3'A2 related proteins (alignment is available under UniProt AC: E9AGZ3). Amyloidogenic regions, conserved cysteine and Integrin-binding motifs are highlighted; right: proposed topology of 3'A2 related proteins D: Frequent SLiMs in the cytoplasmic tail regions of amastins (the numbers denote the unique/total occurrences).

Methods

Resources

Protein sequences were retrieved from UniProtKB (release 2022_05) *UniProt Consortium* (*2023*) and from TriTrypDB *Shanmugasundram et al.* (*2023*) based on the UniProt cross-references (*L. braziliensis, L. donovani, L. infantum, L. major, L. mexicana, Bodo saltans, Leptomonas seymouri, Trypansoma brucei, Trypansoma cruzi, Trypansoma rangeli, Trypanosoma theileri*). Homologs in other kinetoplastids and in SwissProt were searched with BLAST using e-value: 10-5; sequence identity>20%; coverage>50%. In the "Homologs" section all results are displayed, however for most other sections (and calculation) we only used homologous proteins until the first non-kinetoplastid SwissProt hit considering sequence identity (termed as "close-homologs"); Further similar kinetoplastid proteins were therefore considered as a different homology group. Furthermore, we downloaded strains belonging to the 5 selected *Leishmania* species from TriTrypDB. In this case a more stringent condition was used in BLAST, by setting E-value: 10-5; sequence identity>80%; coverage>80%. All kinetoplastid species and strains were used to calculate motif conservation.

We prepared three different type of MSAs using ClustalOmega *Sievers et al.* (*2011*): 1) "non-redundant" MSA using homologous proteins from kinetoplastid reference proteomes; 2) the same MSA but with gaps removed from the "reference" protein that is currently displayed on the webpage; 3) a more redundant MSA using homologous kinetoplastid proteins in all species and strains (used to calculate motif conservation).

High-throughput experiments were first mapped to the corresponding protein using the identifier provided in the original paper, then mirrored to close *Leishmania* homologs if their sequence were identical.

IDRs were predicted using IUPred3 *Erdős et al.* (*2021*) and using the AF2 models' pLDDT and accessibility values - the latter was calculated by DSSP *Joosten et al.* (*2011*), normalized using maximum values calculated as in Tien et al. *Tien et al.* (*2013*), the exposed value threshold defined as suggested by Rost et al. *Rost and Sander* (*1994*). In the case of TM proteins, IDRs were also predicted by MemDis *Dobson and Tusnády* (*2021*). In this in-house modified version, the Position-Specific Scoring Matrices (PSSMs) were generated using kinetoplastid sequence library and secondary structure and accessibility were calculated using AlphaFold2. Topology was predicted by CCTOP *Dobson et al.* (*2015*), however to minimise sporadic erroneous predictions, after an initial prediction we performed a constrained iteration where the topologies of homologous proteins were used as a constraint. Using this approach, closely related proteins will likely have the same topology. Secondary structure elements derived from AF2 structures are also displayed. Pfam domains were identified using InterPro *Paysan-Lafosse et al.* (*2023*). Protein localization was assigned by GO *The Gene Ontology Consortium* (*2019*), predicted by DeepLoc *Almagro Armenteros et al.* (*2017*) and SignalP6.0 *Teufel et al.* (*2022*). NetGPI *Gíslason et al.* (*2021*) was used to predict GPI-anchors (all prediction results are visible, therefore in case of a contradiction it is up to the user to judge the results).

To detect SLiMs that may alter or rewire host cell regulation, we used the regular expressions from ELM *Kumar et al.* (*2022*) on all *Leishmania* sequences. We defined different context filters and merged them into a single score to rank motifs (for more details see Supplementary Material): 1) Disordered: The score is the average of the IUPred3, AF2-based pLDDT and accessibility values. These disordered scores were first transformed so they range from 0 to 1, with 0.5 being the threshold, before calculating their mean; 2) Conservation of the motif was checked among close homologs with some permission for slight misalignment, and penalizing motifs that are present across all kinetoplastids - notably, in this case proteins from different *Leishmania* strains were also considered; 3) Localization: we used a simplified (intracellular/extracellular) distinction. Motif localization was determined using ELM GO annotations, secretion information and CCTOP prediction, while the domain localization was determined from TOPDOM (Varga et al., 2016). We looked for motif-domain pairs where they both have the same simplified (in/out) localization; 4) mRNA level:

using transcriptomic experiments about expression data; 5) protein level: from experiments about
protein abundance; 6) Secretion score based on secretome experiments; 7) Expansion score: re-
flecting how much the protein is expanded in *Leishmania* species (strains not included) compared
to all kinetoplastids; 8) Outgroups score favoring proteins without homologs in SwissProt. Struc-
ture data reflects structure data deposited in the PDB *Varadi et al.* (*2022a*) before 26.03.2023 and
the AlphaFold database (v3). All other data was downloaded in October, 2022 from the source
databases.

Manual curation

We manually curated hundreds of proteins, using two strategies. First, we searched PubMed and
Google scholar for "*Leishmania* host-pathogen protein interaction" and manually processed the re-
sults. Each protein in the experiments was mapped to the corresponding UniProt entry. Then we
mapped interaction data to the 5 *Leishmania* proteomes. When the experiment was performed on
a protein from different species, we mirrored it to the closest homology group in LeishMANIAdb,
and we also indicated on the webpage that the experiment is from a different protein. All interac-
tions were reported according to the community standard MIMIx level *Orchard et al.* (*2007*).

Next we searched for possible surface proteins. For this task we considered all homologs. We
collected topology prediction, GPI-anchor prediction, GO terms and also searched for homologous
proteins that were measured *Bausch-Fluck et al.* (*2018*); *Langó et al.* (*2017*) or predicted to be on
the surface. We manually processed the entries using this approach, taking distant homologues,
domain architectures and conservation patterns into consideration.

Website design

The LeishMANIAdb website is written in PHP (v8.0) using the Laravel (v9.19) framework. All down-
loaded, predicted or calculated data are stored in a local MySQL (v8.0) database. To visualize se-
quence features over amino acid sequences, we developed a javascript package using React (18.2),
while 3D structures are visualized using the original (for non-TM proteins) or a locally modified ver-
sion of Mol* *Sehnal et al.* (*2021*) for TM proteins. The modified version can display the membrane
as two planes around the investigated TM protein using the results of TMDET *Tusnády et al.* (*2005*).

Acknowledgments

This project has received funding from the European Union's Horizon 2020 research and innova-
tion programme under the Marie Skłodowska-Curie grant agreement No 101028908; Ministry of
Innovation and Technology of Hungary from the National Research, Development and Innovation
Fund [K132522]. The authors would like to thank the help of the Gibson team at the European
Molecular Biology Laboratory, especially Jesus Alvarado-Valverde for helpful suggestions and dis-
cussions, and Philippe Esterre (Pasteur Network, Institut Pasteur, Paris) for his help in selecting a
number of *Leishmania* interactions to be curated.

References

Akdel M, Pires DEV, Pardo EP, Jänes J, Zalevsky AO, Mészáros B, Bryant P, Good LL, Laskowski RA, Pozzati G,
Shenoy A, Zhu W, Kundrotas P, Serra VR, Rodrigues CHM, Dunham AS, Burke D, Borkakoti N, Velankar S, Frost
A, et al. A structural biology community assessment of AlphaFold2 applications. Nat Struct Mol Biol. 2022
Nov; 29(11):1056–1067.

Almagro Armenteros JJ, Sønderby CK, Sønderby SK, Nielsen H, Winther O. DeepLoc: prediction of protein
subcellular localization using deep learning. Bioinformatics. 2017 Nov; 33(21):3387–3395.

Ammari MG, Gresham CR, McCarthy FM, Nanduri B. HPIDB 2.0: a curated database for host-pathogen inter-
actions. Database. 2016 Jul; 2016.

Amos B, Aurrecoechea C, Barba M, Barreto A, Basenko EY, Bażant W, Belnap R, Blevins AS, Böhme U, Brestelli
J, Brunk BP, Caddick M, Callan D, Campbell L, Christensen MB, Christophides GK, Crouch K, Davis K, DeBarry

J, Doherty R, et al. VEuPathDB: the eukaryotic pathogen, vector and host bioinformatics resource center.
Nucleic Acids Res. 2022 Jan; 50(D1):D898–D911.

Arango Duque G, Jardim A, Gagnon É, Fukuda M, Descoteaux A. The host cell secretory pathway mediates
the export of Leishmania virulence factors out of the parasitophorous vacuole. PLoS Pathog. 2019 Jul;
15(7):e1007982.

Atayde VD, Aslan H, Townsend S, Hassani K, Kamhawi S, Olivier M. Exosome Secretion by the Parasitic Proto-
zoan Leishmania within the Sand Fly Midgut. Cell Rep. 2015 Nov; 13(5):957–967.

Atayde VD, Hassani K, da Silva Lira Filho A, Borges AR, Adhikari A, Martel C, Olivier M. Leishmania exosomes
and other virulence factors: Impact on innate immune response and macrophage functions. Cell Immunol.
2016 Nov; 309:7–18.

Bausch-Fluck D, Goldmann U, Müller S, van Oostrum M, Müller M, Schubert OT, Wollscheid B. The in silico
human surfaceome. Proceedings of the National Academy of Sciences. 2018; 115(46).

Cuervo P, De Jesus JB, Saboia-Vahia L, Mendonça-Lima L, Domont GB, Cupolillo E. Proteomic characterization
of the released/secreted proteins of Leishmania (Viannia) braziliensis promastigotes. J Proteomics. 2009
Nov; 73(1):79–92.

Davey NE, Travé G, Gibson TJ. How viruses hijack cell regulation. Trends in Biochemical Sciences. 2011;
36(3):159–169.

Dobson L, Reményi I, Tusnády GE. CCTOP: a Consensus Constrained TOPology prediction web server. Nucleic
Acids Res. 2015 Jul; 43(W1):W408–12.

Dobson L, Szekeres LI, Gerdán C, Langó T, Zeke A, Tusnády GE. TmAlphaFold database: membrane localization
and evaluation of AlphaFold2 predicted alpha-helical transmembrane protein structures. Nucleic Acids Res.
2023 Jan; 51(D1):D517–D522.

Dobson L, Tusnády GE. MemDis: Predicting Disordered Regions in Transmembrane Proteins. Int J Mol Sci.
2021 Nov; 22(22).

Erdős G, Pajkos M, Dosztányi Z. IUPred3: prediction of protein disorder enhanced with unambiguous experi-
mental annotation and visualization of evolutionary conservation. Nucleic Acids Res. 2021 Jul; 49(W1):W297–
W303.

Fatoux-Ardore M, Peysselon F, Weiss A, Bastien P, Pratlong F, Ricard-Blum S. Large-scale investigation of Leish-
mania interaction networks with host extracellular matrix by surface plasmon resonance imaging. Infect
Immun. 2014 Feb; 82(2):594–606.

Forrest DM, Batista M, Marchini FK, Tempone AJ, Traub-Csekö YM. Proteomic analysis of exosomes derived
from procyclic and metacyclic-like cultured Leishmania infantum chagasi. J Proteomics. 2020 Sep; 227.

Gianchecchi E, Montomoli E. The enemy at home: leishmaniasis in the Mediterranean basin, Italy on the focus.
Expert Rev Anti Infect Ther. 2020 Jun; 18(6):563–577.

Gíslason MH, Nielsen H, Armenteros JJA, Johansen AR. Prediction of GPI-Anchored proteins with pointer neural
networks. Current Research in Biotechnology. 2021; 3:6–13.

Gregory DJ, Godbout M, Contreras I, Forget G, Olivier M. A novel form of NF-kappaB is induced by Leishmania
infection: involvement in macrophage gene expression. Eur J Immunol. 2008 Apr; 38(4):1071–1081.

Guay-Vincent MM, Matte C, Berthiaume AM, Olivier M, Jaramillo M, Descoteaux A. Revisiting Leishmania GP63
host cell targets reveals a limited spectrum of substrates. PLoS Pathog. 2022 Oct; 18(10):e1010640.

Hassani K, Antoniak E, Jardim A, Olivier M. Temperature-induced protein secretion by Leishmania mexicana
modulates macrophage signalling and function. PLoS One. 2011 May; 6(5).

Jackson AP, Otto TD, Aslett M, Armstrong SD, Bringaud F, Schlacht A, Hartley C, Sanders M, Wastling JM, Dacks JB,
Acosta-Serrano A, Field MC, Ginger ML, Berriman M. Kinetoplastid Phylogenomics Reveals the Evolutionary
Innovations Associated with the Origins of Parasitism. Curr Biol. 2016 Jan; 26(2):161–172.

Jamdhade, Pawar H, Chavan S, Sathe G, Umasankar PK, Mahale KN, Dixit T, Madugundu AK, Prasad TS, Gowda
H, Pandey A, Patole MS. Comprehensive proteomics analysis of glycosomes from Leishmania donovani.
OMICS. 2015 Mar; 19(3).

499 **Jardim A**, Hardie DB, Boitz J, Borchers CH. Proteomic Profiling of Leishmania donovani Promastigote Subcellular
500 Organelles. J Proteome Res. 2018 Mar; 17(3):1194–1215.

501 **Jeffery CJ**. Protein moonlighting: what is it, and why is it important? Philos Trans R Soc Lond B Biol Sci. 2018
502 Jan; 373(1738).

503 **Joosten RP**, te Beek TAH, Krieger E, Hekkelman ML, Hooft RWW, Schneider R, Sander C, Vriend G. A series of
504 PDB related databases for everyday needs. Nucleic Acids Res. 2011 Jan; 39(Database issue):D411–9.

505 **Jumper J**, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A,
506 Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J,
507 Back T, et al. Highly accurate protein structure prediction with AlphaFold. Nature. 2021 Aug; 596(7873):583–
508 589.

509 **Kumar M**, Michael S, Alvarado-Valverde J, Mészáros B, Sámano-Sánchez H, Zeke A, Dobson L, Lazar T, Örd M,
510 Nagpal A, Farahi N, Käser M, Kraleti R, Davey NE, Pancsa R, Chemes LB, Gibson TJ. The Eukaryotic Linear
511 Motif resource: 2022 release. Nucleic Acids Res. 2022 Jan; 50(D1):D497–D508.

512 **Lahav T**, Sivam D, Volpin H, Ronen M, Tsigankov P, Green A, Holland N, Kuzyk M, Borchers C, Zilberstein D,
513 Myler PJ. Multiple levels of gene regulation mediate differentiation of the intracellular pathogen Leishmania.
514 FASEB J. 2011 Feb; 25(2):515–525.

515 **Langó T**, Róna G, Hunyadi-Gulyás É, Turiák L, Varga J, Dobson L, Várady G, Drahos L, Vértessy BG, Medzihradszky
516 KF, Szakács G, Tusnády GE. Identification of Extracellular Segments by Mass Spectrometry Improves Topology
517 Prediction of Transmembrane Proteins. Sci Rep. 2017 Feb; 7:42610.

518 **Matheoud D**, Moradin N, Bellemare-Pelletier A, Shio MT, Hong WJ, Olivier M, Gagnon E, Desjardins M, De-
519 scoteaux A. Leishmania evades host immunity by inhibiting antigen cross-presentation through direct cleav-
520 age of the SNARE VAMP8. Cell Host Microbe. 2013 Jul; 14(1):15–25.

521 **McIlwee BE**, Weis SE, Hosler GA. Incidence of Endemic Human Cutaneous Leishmaniasis in the United States.
522 JAMA Dermatology. 2018; 154(9):1032.

523 **Ohms M**, Ferreira C, Busch H, Wohlers I, Guerra de Souza AC, Silvestre R, Laskay T. Enhanced Glycolysis
524 Is Required for Antileishmanial Functions of Neutrophils Upon Infection With. Front Immunol. 2021 Mar;
525 12:632512.

526 **Orchard S**, Salwinski L, Kerrien S, Montecchi-Palazzi L, Oesterheld M, Stümpflen V, Ceol A, Chatr-aryamontri A,
527 Armstrong J, Woollard P, Salama JJ, Moore S, Wojcik J, Bader GD, Vidal M, Cusick ME, Gerstein M, Gavin AC,
528 Superti-Furga G, Greenblatt J, et al. The minimum information required for reporting a molecular interaction
529 experiment (MIMIx). Nat Biotechnol. 2007 Aug; 25(8):894–898.

530 **de Paiva RMC**, Grazielle-Silva V, Cardoso MS, Nakagaki BN, Mendonça-Neto RP, Canavaci AMC, Souza Melo N,
531 Martinelli PM, Fernandes AP, daRocha WD, Teixeira SMR. Amastin Knockdown in Leishmania braziliensis
532 Affects Parasite-Macrophage Interaction and Results in Impaired Viability of Intracellular Amastigotes. PLoS
533 Pathog. 2015 Dec; 11(12):e1005296.

534 **Patel P**, Mandlik V, Singh S. LmSmdB: an integrated database for metabolic and gene regulatory network in
535 Leishmania major and Schistosoma mansoni. Genom Data. 2016 Mar; 7:115–118.

536 **Paysan-Lafosse T**, Blum M, Chuguransky S, Grego T, Pinto BL, Salazar GA, Bileschi ML, Bork P, Bridge A, Colwell
537 L, Gough J, Haft DH, Letunić I, Marchler-Bauer A, Mi H, Natale DA, Orengo CA, Pandurangan AP, Rivoire C,
538 Sigrist CJA, et al. InterPro in 2022. Nucleic Acids Res. 2023 Jan; 51(D1):D418–D427.

539 **Pellegrini E**, Palencia A, Braun L, Kapp U, Bougdour A, Belrhali H, Bowler MW, Hakimi MA. Structural Basis for
540 the Subversion of MAP Kinase Signaling by an Intrinsically Disordered Parasite Secreted Agonist. Structure.
541 2017 Jan; 25(1):16–26.

542 **Pescher P**, Blisnick T, Bastin P, Späth GF. Quantitative proteome profiling informs on phenotypic traits that
543 adapt Leishmania donovani for axenic and intracellular proliferation. Cell Microbiol. 2011 Jul; 13(7).

544 **Peysselon F**, Launay G, Lisacek F, Duclos B, Ricard-Blum S. Comparative analysis of Leishmania exoproteomes:
545 Implication for host–pathogen interactions. Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics.
546 2013; 1834(12):2653–2662.

547 **Pissarra J**, Pagniez J, Petitdidier E, Séveno M, Vigy O, Bras-Gonçalves R, Lemesre JL, Holzmuller P. Proteomic
548 Analysis of the Promastigote Secretome of Seven Leishmania Species. J Proteome Res. 2022 Jan; 21(1).

549 **Rana S**, Dikhit MR, Rani M, Moharana KC, Sahoo GC, Das P. CPDB: cysteine protease annotation database in
550 Leishmania species. Integr Biol. 2012 Nov; 4(11):1351–1357.

551 **Real F**, Florentino PTV, Reis LC, Ramos-Sanchez EM, Veras PST, Goto H, Mortara RA. Cell-to-cell transfer of Leish-
552 mania amazonensis amastigotes is mediated by immunomodulatory LAMP-rich parasitophorous extrusions.
553 Cell Microbiol. 2014 Oct; 16(10):1549–1564.

554 **Rost B**, Sander C. Conservation and prediction of solvent accessibility in protein families. Proteins. 1994 Nov;
555 20(3):216–226.

556 **Saunders EC**, MacRae JI, Naderer T, Ng M, McConville MJ, Likić VA. LeishCyc: a guide to building a metabolic
557 pathway database and visualization of metabolomic data. Methods Mol Biol. 2012; 881:505–529.

558 **Sehnal D**, Bittrich S, Deshpande M, Svobodová R, Berka K, Bazgier V, Velankar S, Burley SK, Koča J, Rose AS.
559 Mol* Viewer: modern web app for 3D visualization and analysis of large biomolecular structures. Nucleic
560 Acids Res. 2021 Jul; 49(W1):W431–W437.

561 **Shanmugasundram A**, Starns D, Böhme U, Amos B, Wilkinson PA, Harb OS, Warrenfeltz S, Kissinger JC, Mc-
562 Dowell MA, Roos DS, Crouch K, Jones AR. TriTrypDB: An integrated functional genomics resource for kineto-
563 plastida. PLoS Negl Trop Dis. 2023 Jan; 17(1):e0011058.

564 **Sievers F**, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M, Söding J, Thompson
565 JD, Higgins DG. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal
566 Omega. Mol Syst Biol. 2011 Oct; 7(1):539.

567 **Silverman JM**, Chan SK, Robinson DP, Dwyer DM, Nandan D, Foster LJ, Reiner NE. Proteomic analysis of the
568 secretome of Leishmania donovani. Genome Biol. 2008; 9(2).

569 **Silverman JM**, Clos J, de'Oliveira CC, Shirvani O, Fang Y, Wang C, Foster LJ, Reiner NE. An exosome-based
570 secretion pathway is responsible for protein export from Leishmania and communication with macrophages.
571 J Cell Sci. 2010 Mar; 123(Pt 6).

572 **Tayal S**, Bhatia V, Mehrotra T, Bhatnagar S. ImitateDB: A database for domain and motif mimicry incorporating
573 host and pathogen protein interactions. Amino Acids. 2022 Jun; 54(6):923–934.

574 **Teufel F**, Almagro Armenteros JJ, Johansen AR, Gíslason MH, Pihl SI, Tsirigos KD, Winther O, Brunak S, von
575 Heijne G, Nielsen H. SignalP 6.0 predicts all five types of signal peptides using protein language models. Nat
576 Biotechnol. 2022 Jan; 40(7):1023–1025.

577 **The Gene Ontology Consortium**. The Gene Ontology Resource: 20 years and still GOing strong. Nucleic Acids
578 Res. 2019 Jan; 47(D1):D330–D338.

579 **Tien MZ**, Meyer AG, Sydykova DK, Spielman SJ, Wilke CO. Maximum Allowed Solvent Accessibilites of Residues
580 in Proteins. PLoS ONE. 2013; 8(11):e80635.

581 **Torres F**, Arias-Carrasco R, Caris-Maldonado JC, Barral A, Maracaja-Coutinho V, De Queiroz ATL. LeishDB: a
582 database of coding gene annotation and non-coding RNAs in Leishmania braziliensis. Database. 2017 Jan;
583 2017.

584 **Tsigankov P**, Gherardini PF, Helmer-Citterich M, Späth GF, Zilberstein D. Phosphoproteomic analysis of differ-
585 entiating Leishmania parasites reveals a unique stage-specific phosphorylation motif. J Proteome Res. 2013
586 Jul; 12(7):3405–3412.

587 **Tusnády GE**, Dosztányi Z, Simon I. TMDET: web server for detecting transmembrane regions of proteins by
588 using their 3D coordinates. Bioinformatics. 2005 Apr; 21(7):1276–1277.

589 **UniProt Consortium**. UniProt: the Universal Protein Knowledgebase in 2023. Nucleic Acids Res. 2023 Jan;
590 51(D1):D523–D531.

591 **Urban M**, Cuzick A, Seager J, Wood V, Rutherford K, Venkatesh SY, Sahu J, Iyer SV, Khamari L, De Silva N, Martinez
592 MC, Pedro H, Yates AD, Hammond-Kosack KE. PHI-base in 2022: a multi-species phenotype database for
593 Pathogen-Host Interactions. Nucleic Acids Res. 2022 Jan; 50(D1):D837–D847.

594 **Van Roey K**, Uyar B, Weatheritt RJ, Dinkel H, Seiler M, Budd A, Gibson TJ, Davey NE. Short linear motifs: ubiq-
595 uitous and functionally diverse protein interaction modules directing cell regulation. Chem Rev. 2014 Jul;
596 114(13):6733–6778.

Varadi M, Anyango S, Appasamy SD, Armstrong D, Bage M, Berrisford J, Choudhary P, Bertoni D, Deshpande M, Leines GD, Ellaway J, Evans G, Gaborova R, Gupta D, Gutmanas A, Harrus D, Kleywegt GJ, Bueno WM, Nadzirin N, Nair S, et al. PDBe and PDBe-KB: Providing high-quality, up-to-date and integrated resources of macromolecular structures to support basic and applied research and education. Protein Sci. 2022 Oct; 31(10):e4439.

Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, Yuan D, Stroe O, Wood G, Laydon A, Žídek A, Green T, Tunyasuvunakool K, Petersen S, Jumper J, Clancy E, Green R, Vora A, Lutfi M, Figurnov M, et al. AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-sequence space with high-accuracy models. Nucleic Acids Res. 2022 Jan; 50(D1):D439–D444.

Xiang Z, Tian Y, He Y. PHIDIAS: a pathogen-host interaction data integration and analysis system. Genome Biol. 2007; 8(7):R150.

Zhang WW, Matlashewski G. Characterization of the A2-A2rel gene cluster in Leishmania donovani: involvement of A2 in visceralization during infection. Mol Microbiol. 2001 Feb; 39(4):935–948.