Bridging the gap between the evolutionary dynamics and the molecular mechanisms of meiosis : a model based exploration of the *PRDM9* intra-genomic Red Queen

Alice Genestier[1], Laurent Duret[1], Nicolas Lartillot[1,*]

1 Laboratoire de Biometrie et Biologie Evolutive, UMR CNRS 5558, Universite Lyon 1, Villeurbanne, France

* nicolas.lartillot@univ-lyon1.fr

Abstract

Molecular dissection of meiotic recombination in mammals, combined with population-genetic and comparative studies, have revealed a complex evolutionary dynamic characterized by short-lived recombination hotspots. Hotspots are chromosome positions containing DNA sequences where the protein PRDM9 can bind and cause crossing-over. To explain these fast evolutionary dynamic, a so-called intra-genomic Red Queen model has been proposed, based on the interplay between two antagonistic forces: biased gene conversion, mediated by double-strand breaks, resulting in hotspot extinction (the hotspot conversion paradox), followed by positive selection favoring mutant *PRDM9* alleles recognizing new sequence motifs. Although this model predicts many empirical observations, the exact causes of the positive selection acting on new *PRDM9* alleles is still not well understood. In this direction, experiment on mouse hybrids have suggested that, in addition to targeting double strand breaks, *PRDM9* has another role during meiosis. Specifically, PRDM9 symmetric binding (simultaneous binding at the same site on both homologues) would facilitate homology search and, as a result, the pairing of the homologues. Although discovered in hybrids, this second function of *PRDM9* could also be involved in the evolutionary dynamic observed within populations. To address this point, here, we present a theoretical model of the evolutionary dynamic of meiotic recombination integrating current knowledge about the molecular function of PRDM9. Our modeling work gives important insights into the selective forces driving the turnover of recombination hotspots. Specifically, the reduced symmetrical binding of PRDM9 caused by the loss of high affinity binding sites induces a net positive selection eliciting new *PRDM9* alleles recognizing new targets. The model also offers new insights about the influence of the gene

dosage of PRDM9, which can paradoxically result in negative selection on new *PRDM9* alleles entering the population, driving their eviction and thus reducing standing variation at this locus.

Author summary

Meiosis is an important step in the eukaryotic life cycle, leading to the formation of gametes and implementing genetic mixing by recombination of paternal and maternal genomes. A key step of meiosis is the pairing of homologous chromosomes, which is required in order to distribute them evenly into the gametes. Chromosome pairing will also determine the exact position at which paternal and maternal chromosomes will exchange material. Research on the molecular basis of meiosis has revealed the role of a key gene, *PRDM9*. The protein encoded by *PRDM9* binds to specific DNA sequences, by which it determines the location of recombination points. Symmetric binding of the protein (at the same position on the homologous chromosomes) also facilitates chromosome pairing. This molecular mechanism, however, has paradoxical consequences, among which the local destruction of the DNA sequences recognized by PRDM9, leading to their rapid loss at the level of the population over a short evolutionary time. In order to better understand why recombination is maintained over time despite this process, we have developed a simulation program implementing a model taking into account these molecular mechanisms. Our model makes realistic predictions about recombination evolution and confirms the important role played by *PRDM9* during meiosis.

Introduction

In eukaryotes, meiosis is a fundamental step in the reproduction process, allowing the formation of haploid cells from a diploid cell. This process requires the success of a key step, namely, the pairing of homologous chromosomes. Correct pairing is essential for proper segregation of chromosomes into daughter cells. In addition, it allows for the formation of cross-overs, thus implementing recombination and generating new combinations of alleles. Finally, meiosis and recombination are at the heart of the questions of hybrid sterility and speciation [1–3]. However, the evolutionary dynamics of meiotic recombination still remains poorly understood. A correct understanding of this dynamics requires explicit description of the population genetics processes, on one side, and the molecular mechanisms of meiosis, on the other side. In the present work, we present an attempt in this direction, using theoretical and simulation models.

In mammals and many other eukaryotes, recombination points are not uniformly distributed along the chromosomes [4]. Instead, crossovers frequently occur at the same positions in independent meioses, into regions of the genome called recombination hotspots, where the frequency of crossing-over occurrence is 10

to 100 times higher than in the rest of the genome [5, 6]. Recombination hotspots are typically 1 to 2 kb long, and they are often located outside of genes. More than 30,000 hotspots were found in humans [4, 7], and around 40,000 in mice [8]. These hotspots are characterized in humans by the presence of a sequence motif (13-mer) determining up to 40% of crossing-overs [9]. This motif has helped to identify the *PRDM9* gene as the gene responsible for hotspot location, [9, 10]. *PRDM9* encodes a DNA-binding protein, and the hotspots therefore correspond to strong binding sites of this protein.

Once bound to its target site, the PRDM9 protein trimethylates surrounding histones (H3K4me3 and H3K36me3 [11, 12]), inducing the recruitment of the double strand break (DSB) machinery near the target site [13] (for a review see [14]). DSB induction, followed by DNA resection, produces a single-stranded end that searches for its homologous sequence on the other chromosome and is then repaired, using the sequence at the same locus on the homologous chromosome as the template. This repair leads to the conversion of the sequence located near the break (often overlapping the site targeted by PRDM9 [15]). Some of these gene conversion events lead to the formation of crossing-overs (CO), while others are repaired without exchange of flanking regions (non crossing-overs, NCO). In some cases, repair is done not with the homologue but with the sister chromatid [15].

Crucially, when allelic variation exists at a given target motif that modulates the binding affinity for PRDM9, DSBs will form more frequently on the allele for which PRDM9 has the highest affinity (the 'hot' allele). Given that DSBs are repaired by using the intact homologue as a template, this process leads to the preferential replacement of 'hot' alleles by alleles for which PRDM9 has a lower affinity. This mechanism of biased gene conversion favors the transmission of mutations that inactivate hotspots, leading to an increase in the frequency of these mutant inactive versions of binding sites, as well as their fixation in the population. This self-destruction phenomenon, commonly called the "hotspot conversion paradox" [16], leads to the progressive inactivation, hereafter called *erosion*, of PRDM9 binding sites at the genome scale, over short evolutionary times (in the order of 10,000 to 100,000 generations [17]), which therefore raises the question of the maintenance of recombination in the long term.

As a solution to the paradox, a model has been proposed [18], which works as follows: the disappearance of recombination hotspots compromises the proper functioning of meiosis and greatly reduces the fertility of individuals. In this context, new *PRDM9* alleles, recognizing new target sites already present by chance in the genome and thus restoring recombination, would be positively selected. They would eventually replace the alleles currently segregating in the population. By analogy with the so-called Red Queen dynamics [19] typically displayed by prey-predator or host-pathogen systems, this model has been called the intragenomic Red Queen model of recombination hotspot evolution [18]. It predicts a rapid evolutionary dynamic of

recombination landscapes, as well as a strong positive selection on the DNA binding domain of *PRDM9*. [89]

Several empirical observations support the Red Queen model. First, fine scale recombination landscapes [90] differ between closely related species, like between humans and chimpanzees [10, 20], or between modern [91] humans and Denisovan [17], suggesting a fast turnover of recombination hotspots. Second, the DNA binding [92] domain of PRDM9 is a zinc finger domain, consisting of an array of 7 to 10 zinc fingers. This domain is [93] encoded by a minisatellite, which mutates rapidly [21] by a combination of point mutations and unequal [94] crossing over between the sequence repeats. This allows for the rapid accumulation of new combinations of [95] zinc fingers, and thus of new alleles recognizing different hotspots, providing the necessary mutational input [96] for the Red Queen to run. In part because of this high mutation rate, *PRDM9* is typically characterized by [97] a high genetic diversity in natural populations [22–26] . Finally, non-synonymous substitutions in the zinc [98] finger domain of *PRDM9* are more frequent than synonymous substitutions, suggesting the presence of a [99] strong positive selection acting on the DNA binding domain [3, 24]. [100]

Studies combining computer simulations and theoretical analyses have also given many arguments in favor [101] of the Red Queen model [18], [27]. However, current theoretical models remain silent on two essential points. [102] First, what exact mechanism explains the positive selection acting on *PRDM9* to restore recombination? [103] Current models invoke a decline in fertility caused by the erosion of recombination, but without providing a [104] precise explanation of this point. Second, these models do not provide any explanation, at this stage, of the [105] link between the intra-genomic Red Queen and the hybrid sterility seen when mice are heterozygous for [106] certain *PRDM9* alleles. [1, 28]. [107]

In this respect, a series of molecular and comparative analyses conducted on the mouse provide some [108] clues. First, the differential erosion of PRDM9 binding sites between mouse subspecies (due to different [109] *PRDM9* alleles eroding their respective target sites within each subspecies) leads to asymmetrical binding [110] configurations in hybrid individuals [29]. Specifically, in a F1 hybrid, each of the two *PRDM9* alleles has [111] eroded its targets in the genome of its parental strain, but not in the other strain's genome. Each *PRDM9* [112] allele will therefore tend to bind preferentially to the still active target sites present on the chromosomes [113] inherited from the other parent, but not to the homologous but eroded sites on the chromosome from [114] its own parent. These asymmetrical binding patterns of PRDM9 across the whole genome are suspected [115] to be involved in the sterility phenotype. Indeed, Chip-Seq experiments have uncovered a correlation [116] between PRDM9 binding asymmetry rates in a variety of mouse hybrids for different pairs of *PRDM9* alleles [117] and hybrid sterility [30, 31]. Cytogenetic observations have also shown that chromosomes that are more [118] asymmetrically bound are less often correctly paired during metaphase I [31]. Finally, the introduction by [119] transgenesis in mice of a *PRDM9* allele possessing the human zinc finger domain, which has never been [120]

present in the mouse species complex and has then not eroded its targets asymmetrically in either of the two populations, restores both a good binding symmetry and high levels of fertility [1].

In the light of these empirical results, a model has been proposed [1], according to which *PRDM9* would in fact have a dual role during meiosis: in addition to being responsible for the recruitment of the DSB machinery, it would also facilitate the pairing between the homologous chromosomes, thanks to its symmetrical binding, that is, its simultaneous binding to the same target site on both homologues. More precisely, PRDM9 is thought to act to bring the sites on which it is bound to the chromosomal axis. The mechanisms involved in this step are not very clear, although it would seem that H3K4me3 markers [32] and the SSXRD and KRAB domains of PRDM9 are implicated [33]. This is where symmetry would play a key role (Fig 1): when PRDM9 is bound symmetrically, the two homologous loci are each brought closer to the chromosomal axis, which would, by some yet unknown mechanism, make the complementary sequence of the homologue more directly accessible to the single-stranded end produced by resection of the DSB [34, 35]. In contrast, in the case where PRDM9 binds only on one of the two homologous loci, a possible DSB would be repaired only later on, either by the homologue as a non CO event or by the sister chromatid [15].

This mechanistic model based on the symmetrical binding of PRDM9 provides a globally coherent explanation of the hybrid sterility phenotype. A last question remains open however: could this dual role of *PRDM9* also be implicated in the Red Queen evolutionary dynamic of recombination observed within a single population? Of note, in the hybrid, failure of meiosis is a consequence of macroscopic asymmetric sequence patterns due to differential erosion in the two parental lineages. Such macroscopic asymmetric sequence patterns are unlikely within a single population. On the other hand, statistical binding asymmetries might nevertheless occur and play a role.

To investigate this question, in this paper, we revisit the theoretical modeling of the intra-genomic Red Queen of *PRDM9*-dependent recombination. In contrast to previous work [18, 27] (but see the recent work of Baker *et al.* [36]), we explicitly model the molecular mechanism of meiosis and, more specifically, the function currently hypothesized for *PRDM9*. Our specific aim was to investigate whether the combined effects of biased gene conversion and symmetry provide sufficient ingredients for running an intra-genomic Red Queen able to explain current empirical observations.

Results

To investigate the evolutionary dynamic of *PRDM9*-dependent recombination, we developed a simulation program modeling the evolution of a randomly mating population of diploid individuals and accounting

Figure 1: **A model of how symmetrical PRDM9 binding facilitates chromosome pairing.** Upon binding DNA at a specific target motif, PRDM9 (orange oval) brings the DNA segment close to the chromosomal axis. Some of the sites bound by PRDM9 may then undergo a DSB (red star). Resection of the DSB generates a single-stranded end, which will search for a complementary sequence to use as a template for repair. In the case where PRDM9 is symmetrically bound (i.e. on both homologues, case on the left-hand side), and assuming that homology search is restricted to the axis region, the templates provided by the two sister chromatids of the homologue are more directly accessible, thus facilitating homology search and pairing with the homologue. The break can then be repaired either as a CO or a NCO event, in both cases, implementing gene conversion at the broken site. In the case of asymmetrical PRDM9 binding (case shown on the right-hand side), the homologue is less directly accessible, preventing efficient homologue engagement. The broken site is assumed to be repaired later on, once the homologues have synapsed (and this, thanks to other DSBs occurring at symmetrically bound sites somewhere else on the same pair of chromosomes), as NCO events. In the case where the homologue bears an inactive binding site at the position corresponding to the DSB, the NCO will effectively implement biased gene conversion in favor of the inactive version.

for the key molecular processes involved in meiotic recombination (Fig 2). The genome consists of a single chromosome, bearing a locus encoding the *PRDM9* gene (Fig 2A). The locus mutates at a rate u, producing new alleles recognizing different sequence motifs (Fig 2B). Each allele recognizes a set of binding sites randomly scattered across the genome, of varying binding affinity. Binding sites undergo inactivating mutations at a rate v (Fig 2B). At each generation, sexual reproduction entails the production of gametes that are themselves obtained by explicitly implementing the process of meiosis on randomly chosen individuals (Fig 2C-F).

An individual is randomly chosen to attempt meiosis and reproduce. Meiosis is modeled step by step. First, the PRDM9 proteins bind to their target sites, according to a simple chemical equilibrium (Fig 2C). The probability of occupation of a site will thus depend on the binding affinity of this hotspot site for PRDM9. It may also depend on PRDM9 concentration, which can itself depend on the genotype of the individual (either homozygous for single *PRDM9* allele, or heterozygous for two distinct *PRDM9* alleles), through gene dosage effects. Then, a small number of double strand breaks are induced at some of the sites bound by PRDM9 (Fig 2D). At that step, a key assumption of the model is that meiosis will succeed only if at least one of those DSBs occurs at a site that is also bound by PRDM9 on at least one of the two chromatids of the homologous chromosome, in which case a single cross-over is performed at the site of one of the DSBs fulfilling this requirement. Finally, all DSBs are repaired using the homologue (Fig 2E), upon which meiosis is assumed to proceed successfully, producing four gametes, one of which is chosen at random for reproduction (Fig 2F).

Of note, because of the symmetry requirement, the probability of success of meiosis will depend on the genotype of the individuals on which it is attempted, which will thus induce differences in fertility between individuals. In addition, explicit repair of the DSBs by the homologue effectively implements the process of gene conversion at the binding sites, causing hotspot erosion in the population. The main question is then to what extent these two aspects of the molecular mechanism will influence the evolutionary dynamic.

The overall procedure is run for several tens of thousands of generations, during which several summary statistics are monitored: the frequency of each *PRDM9* allele and the resulting genetic diversity at the *PRDM9* locus, the mean proportion of binding sites of *PRDM9* alleles that are currently active, their mean affinity across the genome, the mean probability of symmetrical binding and the mean probability of success of meiosis. The model parameters and the monitored statistics are summarized in Table 1

The results section is divided in three main parts. First, a simple version of the model is presented, in which PRDM9 gene dosage is ignored (that is, assuming that the number of proteins produced by a given *PRDM9* allele is the same in a homozygote and in a heterozygote for this allele). Of note, this model is

Figure 2: **Diagram summarizing the main features of the model and the successive steps of the simulation cycle.** (A) The model assumes a population of N diploid individuals ($2N$ chromosomes, vertical lines). Each chromosome has a *PRDM9* locus (filled oval, with a different color for each allele) and, for each *PRDM9* allele, a set of target sites (filled rectangles, with color matching their cognate *PRDM9* allele) of variable binding affinity (variable width of the filled rectangles). (B) Mutations at the *PRDM9* locus create new alleles (here, purple allele), while mutations at the target sites inactivate PRDM9 binding (grey sites with a cross). (C-E) meiosis (here in a heterozygous individual); (C) Chromosome replication and binding of PRDM9 to its target sites (filled circles). (D) Induction of DSBs at a small number of randomly chosen sites bound by PRDM9 (red stars) and search for DSBs at symmetrically bound sites (symmetrical DSBs); if no symmetrical DSB is found, meiosis fails; otherwise, one symmetrical DSB is chosen uniformly at random (here, on the yellow binding site). (E) Completion of the crossing-over (red curved arrow) and repair of all DSBs using the homologous chromosome (red box on top); (F) random choice of one of the 4 gametes of the tetrad, which will fuse with another gamete and become part of the next generation.

not empirically plausible. It implies that the total concentration of PRDM9 protein in meiocytes is twice 183

as high in heterozygotes compared to homozygotes, something that would be possible only if there was 184

Table 1: **Description of input parameters and output variables**

Parameters	Description	Value
u	Mutation rate at the *PRDM9* locus	2×10^{-6} to 5×10^{-3}
v	Mutation rate at the targets	2×10^{-6} to 5×10^{-3}
N	Population size	$5,000$
h	Number of targets recognised by a new allele	400
d	Mean number of DSB per chromosome pair per meiocyte	6
y_i	Affinity of target i	variable
g	Gene conversion rate	variable
n_{mei}	Number of meioses allowed per individual before reproduction failure	1 to 5

Variables	Description
f_i	Frequency of allele i [a]
θ_i	Proportion of target sites still active for allele i; erosion level is then defined as $1 - \theta_i$ [a]
q_i	Mean probability of symmetrical binding of allele i [a,b]
w_i	Mean fertility of allele i ($\bar{w}$ is the mean over all of the alleles) [a,b]
ρ	Net rate of erosion per generation
x_j	Occupancy probability at site j at equilibrium
z	Intrinsic age of a generic allele (Eq 2 and 3)[a]
$\bar{z}$	Mean intrinsic age – also an analytical estimate of mean erosion level (Eq 3 and 4)[a]
α	Linear response of the log-fitness as a function of erosion
τ	Mean time between successive invasions of the population by new *PRDM9* alleles
D	*PRDM9* diversity in the population at equilibrium
s_0	mean selection coefficient acting on new *PRDM9* alleles at equilibrium
σ_0	Relative difference in fertility between homo- and heterozygotes (or equivalently, between hemi- and heterozygotes) for young alleles
$\bar{\sigma}$	Mean relative difference in fertility between homozygotes and hemizygotes in the population

[a] These variables also change over time
[b] the mean is over all individuals carrying this allele (with a weight of 1 for heterozygotes and a weight of 2 for homozygotes).

an allele-specific transcriptional feedback. The fast turnover of *PRDM9* alleles, each with its own binding [185] specificity, does not seem to be compatible with such a mechanism. Nevertheless, in spite of its lack of [186] empirical relevance, this model represents a useful stepping stone for understanding the consequences of [187] erosion and of the requirement of symmetrical binding, before considering the additional complications [188] brought about by gene dosage. Then, in a second part, we introduce PRDM9 gene dosage and work out its [189] consequences, by systematically comparing the outcome of the simulations run with and without dosage, [190] under otherwise identical settings. Finally, we attempt an empirical calibration of the full model (including [191] dosage) based on current knowledge in the mouse, so as to test its predictions. [192]

Intragenomic Red Queen [193]

We first ran the model without gene dosage, and with a low mutation rate at the *PRDM9* locus ($u = 5 \times 10^{-6}$). [194] Note that the parameter values used here are not empirically relevant. Instead, the aim is to illustrate the [195]

different regimes produced by the model. An empirical calibration of the model will be presented below.   196

Assuming a low mutation rate for *PRDM9* results in few, rarely more than one, alleles segregating at   197
any given time in the population (Fig 3A). We call this a *monomorphic* regime, after Latrille *et al.* [27].   198
The dynamic is as follows. First, an allele appears in the population and invades, until reaching a frequency   199
close to 1. Meanwhile, the proportion θ of active binding sites for this allele decreases, due to inactivating   200
mutations that are then driven to fixation by biased gene conversion Once the allele has eroded a fraction of   201
around 20% of its sites, it is quickly replaced by a newly arisen *PRDM9* allele that recognizes a different   202
hotspot sequence motif. This rapid replacement clearly suggests the presence of strong positive selection.   203
The newly invading allele then erodes its target until being replaced by a new allele which in turn follows a   204
similar trajectory, and so on.   205

A key aspect of the model is that it does not explicitly invoke a fitness function. Instead, the positive   206
selection that seems to be operating on new alleles (Fig 3) is an emerging property of the mechanism of   207
meiosis. This positive selection can be more precisely understood by examining how the probability of   208
PRDM9 symmetrical binding (defined by Eq (13) and Eq (14), see Materials and methods section) and the   209
mean fertility of a typical allele evolve over the lifespan of this allele (Fig 3D and 3E), and how this relates   210
to the level of erosion (defined as the fraction of targets that have been inactivated), or equivalently the   211
proportion of still-active targets (Fig 3B) and the mean affinity of the remaining non-eroded sites (Fig 3C).   212

First, we observe a clear correlation between allele frequency and proportion of active (non-eroded) target   213
sites. When the frequency of an allele increases in the population, the proportion of still active sites for   214
this allele decreases. This erosion (Fig 3B) seems to occur at a rate which is directly proportional to the   215
frequency of the allele in the population (Fig 3A). This is expected: the more frequent an allele, the more   216
often it will be implicated in a meiosis and the more opportunities its target sites will have to undergo gene   217
conversion events.   218

Second, erosion results in a decrease of the mean affinity of the sites that are still active (Fig 3C).   219
This reflects the fact that sites of high affinity are more often bound by PRDM9, and thus are more often   220
converted by inactive mutant versions.   221

Third, sites of lower affinity are also less often symmetrically bound (i.e. bound simultaneously on both   222
homologues). The key quantity that captures this effect is the conditional rate of symmetrical binding   223
(q). Since DSBs are chosen uniformly at random among all bound sites by PRMD9, q corresponds to the   224
probability that a DSB will occur in a symmetrically bound site and will thus contribute to a successful   225
pairing of the homologues. This probability is a monotonous function of the mean affinity of the binding   226
sites for the PRDM9 protein (Fig 3D). The mean value of q over all target sites of a given *PRDM9* allele is   227

Figure 3: **A simulation trajectory showing a typical evolutionary dynamic, under a monomorphic regime (N = 5000, u = 5 × 10⁻⁶, v = 5 × 10⁻⁵).** In all panels, each color corresponds to a different allele. Note that a given color can be reassigned to a new allele later in the simulation. Each panel represents the variation through time of (A) the frequency of *PRDM9* alleles and their corresponding (B) proportion of active sites, (C) mean affinity, (D) probability of symmetrical binding and (E) fertility. The thick line singles out the trajectory of a typical allele.

thus lower for older alleles (Fig 3D).   228

Finally, since, in our model, meiosis requires at least one DSB in a symmetrically bound site, the mean   229

fertility of an older allele across all the meiosis it participates in is lower (Fig 3E). Hence, new alleles (young   230

alleles) will be positively selected at the expense of old alleles and will ultimately replace them in the population.

To assess to what extent the requirement of symmetrical binding provides the selective force driving the evolutionary dynamics of *PRDM9*, we performed additional simulations with a simpler model, in which PRDM9 is required for the formation of DSBs, but symmetric binding is not required for chromosome pairing. With this setting, the model still predicts a turnover of *PRDM9* alleles, but with high levels of erosion (36% of mean target inactivation under the control simulation, versus 12% under the model requiring symmetrical binding). Fundamentally, *PRDM9* alleles persist in the population until they have no more sites to bind (S2 Fig), at which point they cannot anymore recruit DSBs and are thus selectively eliminated. Thus, a key result obtained here is that the requirement of symmetrical binding for chromosome pairing, combined with preferential erosion of high affinity sites, is sufficient for creating differences in fitness (fertility) between young alleles and older alleles having intermediate levels of erosion, ultimately providing the selective force behind the turnover of *PRDM9* alleles.

Of note some stochastic deviations from this typical life-cycle for a *PRDM9* allele are sometimes observed, such as an allele being first outcompeted by a subsequent allele but then showing a rebound in frequency when the competitor has itself eroded a large fraction of its target sites. Such deviations are relatively rare and do not seem to fundamentally change the overall regime. It should also be noted that we observe an increase in the rate of symmetrical binding and in the mean fertility at the beginning and at the end of the life of the alleles. The reason for this is that these two summary statistics are defined, for each allele, as a mean over all diploid genotypes segregating in the population that are carrying this allele. As a result, when old alleles have declined to a low frequency, they often find themselves in a heterozygous state with new alleles, which restores the rate of symmetrical binding and thus the fertility of the corresponding diploid individual. Likewise, when a new allele appears in the population, it is in a heterozygous state with an older allele, which gives a lower rate of symmetrical binding and fertility than being in the homozygous state.

The simulation shown above (Fig 3) was run under a low mutation rate, hence resulting in a *monomorphic* regime. Running the simulation under higher mutation rates for *PRDM9* ($u = 5 \times 10^{-4}$) results in a *polymorphic* regime, where many alleles segregate together at any given time. In this regime, of which a typical simulation is shown in Fig 4, the Red Queen process is also operating, except that many alleles are simultaneously segregating (Fig 4A), at a lower frequency. As in the previously shown monomorphic regime, each allele undergoes erosion (Fig 4B), primarily of its higher affinity sites (Fig 4C), again causing a decrease in symmetrical binding rate (Fig 4D) and fertility (Fig 4E). Owing to the high mutation rate, however there is a much lower erosion, leading to a lower decrease in fertility than in a monomorphic regime.

Note that in Fig 4, the scale of the axes of the ordinates are not the same as in Fig 3 (see S1 Fig for a figure
with same scale on the y-axis).

Figure 4: **A simulation trajectory showing a typical evolutionary dynamics, under a polymorphic regime (**$N = 5000$**,** $\mathbf{u = 5 \times 10^{-4}}$**,** $\mathbf{v = 5 \times 10^{-5}}$ **).** In all panels, each color corresponds to a different allele. Note that a given color can be reassigned to a new allele later in the simulation. Successive panels represent the variation through time of (A) the frequency of *PRDM9* alleles and their corresponding (B) proportion of active sites, (C) mean affinity, (D) probability of symmetrical binding and (E) mean fertility. The thick line singles out the trajectory of a typical allele. Note that the scale of the y-axis for all panels is not the same as in Fig 3.

Control simulations (without the requirement of symmetry) run in the polymorphic case result in erosion

levels of around 15% of the targets inactivated on average (Supplementary S2 Fig), thus not as high as in the control simulations in the monomorphic regime, although still much higher than in the simulations in which symmetrical binding is required (around 4%, Fig 4). Without the requirement of symmetry, and in a context where the number of sites bound by PRDM9 is in excess compared to DSBs (thus unlike the control shown above for a low mutation rate u), there is no selective difference between individuals. The overall dynamic under this control is therefore neutral, with a turnover caused by the mutational input of new alleles and loss of old alleles by drift. This is in contrast with the fundamentally selective regime observed under the requirement of symmetrical binding (with a difference in fertility between young and old alleles of the order of 10^{-3}, Fig 4, bottom panel, resulting in a mean scaled selection coefficient in favor of new alleles of $4Ns_0 \simeq 34$). Thus, here also, as in the monomorphic regime considered above (Fig 3), the requirement of symmetrical binding, and more precisely the differential success of new and old alleles in fulfilling this requirement, is ultimately the mechanism mediating the selective force behind the turnover of *PRDM9* alleles.

Taking into account *PRDM9* gene dosage

The previous results were obtained with a model assuming the same concentration of the PRDM9 protein product of a given allele in individuals that are either homozygous or heterozygous for this allele. Yet, as mentioned above, this model is not realistic. Instead, gene dosage is likely to be an important aspect of the regulation of *PRDM9* expression [37]. To account for this fact, gene dosage was introduced in the simulation model. This was done by assuming that the amount of protein produced by a given allele is proportional to its copy number (i.e. twice as high in a homozygote than in a heterozygote). The main consequence of introducing gene dosage is that, in homozygotes, the occupancy of a site of a given affinity is increased, compared to a heterozygote, leading to a higher probability of symmetrical binding and fertility. This turns out to have an important impact on the Red Queen dynamics.

To illustrate this point, Fig 5 shows the results obtained with gene dosage in an identical parameter configuration as in the simulation without gene dosage shown in Fig 4. We observe a drastic change of regime between the two settings (compare Fig 4 and Fig 5). While the simulation without gene dosage gives a polymorphic regime with an average maximum frequecy of 0.2, the simulation with gene dosage gives a more heterogeneous dynamic. Indeed, in this regime, some alleles can partially or even completely invade the population for certain period of time. During this period of domination by one allele, new alleles apparently cannot invade the population.

What happens here is that, due to gene dosage, homozygotes have a fitness advantage over heterozygotes.

Figure 5: **Example of a *PRDM9* dynamic with gene dosage ($N = 5000$, mutation rates $u = 5 \times 10^{-4}$ at the *PRDM9* locus, $v = 5 \times 10^{-5}$ at the target sites).** In all panels, each color corresponds to a different allele. Note that a given color can be reassigned to a new allele later in the simulation. Successive panels represent the variation through time of (A) the frequency of *PRDM9* alleles and their corresponding (B) proportion of active sites and (C) selection coefficient acting on new alleles.

Since alleles at high frequency are more often present in a homozygous diploid genotype, they have an advantage over low frequency alleles. And so, paradoxically, old (but not too old) alleles at high frequency can have a higher mean fitness than new alleles that just appeared in the population, and this, in spite of their higher levels of erosion. This last point is confirmed by measuring the selection coefficient associated with new alleles (Fig 5C): during a phase of domination of one allele, new alleles are strongly counter-selected. When the dominant allele becomes too old, its homozygote advantage is no longer strong enough to compensate for its erosion. At this point, new alleles become positively selected (e.g. at around $15,000$ generations on Fig 5C). The old allele then quickly disappears and all other alleles competes for invasion. This transient polymorphic regime is unstable however: as soon as one of the young alleles reaches a higher frequency than its competitors, it benefits from a homozogous advantage, ultimately eliminating all competitors from the population while still rare and thus becoming the new dominant allele. In the following, we call *eviction* this phenomenon fundamentally induced by gene dosage.

Running a control experiment without the requirement of symmetry (see S3 Fig) in the present case gives a regime characterized by higher levels of diversity ($D = 12$, compared to $D = 7.5$ when symmetrical binding is required), and an equilibrium erosion (15% of the targets inactivated on average) that is more pronounced than when symmetrical binding is required (8%). Again, as in the case without dosage, the regime is essentially characterized by a neutral turnover of *PRDM9* alleles. Thus, in the presence of dosage, the requirement of symmetrical binding to complete meiosis provides the selective force that drives both the selective replacement of old alleles by new ones (as in the case without dosage) and the eviction of all alleles except one (the specific contribution of dosage).

Quantitative assessment of the equilibrium regime

The simulations shown in the previous section have helped to tease out the molecular details of the implication of *PRDM9* in the Red Queen dynamics of recombination. However, it remains to be understood how the equilibrium regime quantitatively depends on the parameters of the model. Another important question is to better understand the conditions under which gene dosage is expected to have an impact on the qualitative regime of the Red Queen, such as observed in the example shown above (Fig 5). To address these issues, scaling experiments were performed (see methods and S4 Fig), systematically varying the parameters of the model so as to determine how the equilibrium changes accordingly, and systematically contrasting the models with and without dosage.

A first experiment was performed, varying the mutation rate u at the *PRDM9* locus and the rate v of target inactivation. The standing diversity at the *PRDM9* locus is strongly dependent on u, with higher diversity levels under higher values of u (Fig 6A). This merely reflects the general fact that genetic diversity directly depends on the fresh mutational input at the locus. Diversity depends much more weakly on v, showing a variation of at most 20% over 4 orders of magnitude for v, except in a regime with otherwise intermediate values of u ($4Nu = 10$, 6C), where the equilibrium shows a sharp transition from a monomorphic ($D \simeq 1$) to a polymorphic ($D \simeq 10$) regime as v is increased (Fig 6A).

Importantly, this sharp transition is critically dependent on the presence of gene dosage, being absent in the control conditions without gene dosage (Fig 6B). More generally, diversity tends to be lower, and the monomorphic regime more prevalent, in the presence of gene dosage (compare panels A and B). This confirms and generalizes the observations made above (Fig 5), while giving a better characterization of the conditions under which introducing gene dosage under otherwise constant parameter values can result in a transition from a polymorphic to a monomorphic regime: essentially for values of $4Nu$ of the order of 10 and low values of v (Fig 6C).

Figure 6: **Scaling experiment as a function of the mutation rates u and v ($N = 5 \times 10^3$).** A and B: equilibrium levels of *PRDM9* diversity with (A) and without (B) genetic dosage. D and E : equilibrium levels of *PRDM9* erosion level with (D) and without (E) dosage. C and F: summary statistics that are predictive of the type of regime observed in the presence of dosage, namely: (C) $4Nu$, and (F) $\sigma_0\tau$ (selection coefficient associated to dosage multiplied by the time between successive invasions by *PRDM9* alleles). The white horizontal line separates the graph into two parts: $4Nu \geq 10$ above and $4Nu < 10$ below. The regime is predicted to be polymorphic in the presence of dosage if $4Nu > 10$ and $\sigma_0\tau < 3$

When in the polymorphic regime (i.e. for high values of u, such that $4Nu \geq 10$, the mutation rates u and v have opposing effects on erosion levels (Fig 6D). This can be understood as follows. The equilibrium set point of the Red Queen is fundamentally the result of a balance between erosion and invasion. If u increases, this also increases the rate of replacement of old alleles shifting the equilibrium towards weaker equilibrium erosion. Conversely, the rate of erosion is directly proportional to v, such that increasing v

will result in an increase of the rate of erosion at equilibrium and, correlatively, lower rates of symmetrical binding and fertility.

However, these opposing effects of u and v on equilibrium erosion levels are only valid for high values of u. For lower values of u ($4Nu \leq 10$), on the other hand, as v is increased, the equilibrium erosion level first shows a decrease and then stabilizes. This somewhat paradoxical response of erosion to changes in the rate of inactivation of the targets of PRDM9 is dependent on gene dosage: in the control experiment in which dosage is deactivated, the equilibrium erosion level always increases as a function of v, except for the lowest values of u (Fig 6E).

Characterization of the impact of gene dosage

The observations just presented show that gene dosage has a critical impact on the observed equilibrium. Specifically, for intermediate values of u and low values of v, dosage tends to create a monomorphic regime characterized by high levels of erosion.

To better understand this point, an analytical approximation was derived, using the mean-field approach introduced in Latrille *et al*. The complete analytical developments can be found in S1 Appendix sections 1 to 3. Here, a simplified summary of the main result is given. This analytical derivation relies several assumptions: a polymorphic regime, strong selection at the *PRDM9* locus, weak erosion, and a weak impact of gene dosage. It is thus not valid over the full range of parameter values explored here, in particular given the strong impact of dosage on the standing diversity of *PRDM9* observed above. However, it will give the conditions on the model parameters for which gene dosage is in fact not weak and is thus expected to have a strong impact on the observed dynamic.

First, to a good approximation, the rate of erosion of the targets of an allele depends on its frequency in the population :

$$\frac{d\theta}{dt} \approx -\rho f \theta, \tag{1}$$

where $\rho = \frac{Nvd}{2h} = 4Nvg$ and $g = \frac{d}{8h}$ is the net rate of erosion per generation. As a result, an analytical proxy for the cumulated erosion of an allele of age t is given by:

$$z(t) = \rho \int_0^t f(t') \, dt', \tag{2}$$

where $f(t')$ is the frequency of the allele at time t'. The quantity z can thus be seen as the *intrinsic age* of the allele (i.e. an allele ages more rapidly if more frequent in the population). Because of selection, the

frequency of an allele changes as a function of its erosion, or intrinsic age z.

Assuming weak erosion ($z << 1$) allows one to linearize the differential equation describing the evolution of the frequency of an allele, which gives:

$$\frac{d\ln f}{dt} \simeq -\frac{\alpha}{2}(z - \bar{z}) + \sigma_0(f - \bar{f}). \tag{3}$$

Here, $\bar{z}$ is the mean intrinsic age of all alleles currently segregating in the population, weighted by their frequency, and α is the slope of the log fitness as a function of erosion (how much fertility decreases as a function of the erosion of the targets of the allele). Thus, the first term of this equation captures the age effect: everything else being equal, older (and thus more eroded) alleles tend to bind less often symmetrically and therefore tend to have a lower fitness than younger allele. As a result, they will tend to undergo a decrease in their frequency in the population.

The second term of the equation is the specific contribution of gene dosage. In this second term, f and $\bar{f}$ are the current frequency of the allele and the mean frequency of all other segregating in the population. As for σ_0, it is defined as the relative difference in fertility between homozygotes and heterozygotes for young alleles (with $z = 0$). It thus captures the homozygous advantage induced by gene dosage. The homozygous advantage depends on the age of the allele, however, under the weak erosion limit considered here, it can to good approximation be taken as constant and equal to that calculated for new alleles. Of note, under the assumptions of the model, σ_0 also measures the relative difference in fertility between a homozygote and a hemizgote for a new allele, that is, the haplo-insufficency of *PRDM9*.

This second term of the equation captures the frequency effect: everything being equal, more frequent alleles benefit from a homozygous advantage and will therefore tend to become even more frequent. Importantly, this term systematically acts against diversity. It captures the self-reinforcing nature of the homozygous advantage, and thus gives the key for explaining the origin of the eviction phenomenon observed in Fig 5.

Based on Eq 3, we can determine when dosage will play an important role. To address this point, we adopt a perturbative approach. Here, perturbative means that we first work out the equilibrium in the simpler situation where gene dosage is absent, thus ignoring the second term of Eq 3, and then characterize the conditions under which this term should in fact have an important impact.

In the absence of dosage, a self-consistent mean-field approximation for the mean equilibrium erosion

can be obtained (Supp Material S1 Appendix section 2.4):

$$1 - \bar{\theta} = \bar{z} \approx \sqrt{\frac{vg}{u\alpha}},$$

(4)

as well as the equilbirium diversity D:

$$D \quad \approx \quad 24Nu.$$

(5)

Also of interest is the mean time τ between successive invasions by new *PRDM9* alleles:

$$\tau = \frac{1}{2N} \frac{1}{\sqrt{u\alpha vg}}.$$

(6)

Note that these equations are the same as those presented in Latrille *et al.* [27]. However, in [27], α and g were phenomenological parameters, set arbitrarily, while in our mechanistic model, they emerge directly from the molecular mechanisms of meiosis. As such, they can be expressed as functions of the model parameters (d, h, N or v, see S1 Appendix Eq 56 and Eq 34).

In Eq 4 and 5, we recover a quantitative account of the main trends observed in the absence of dosage, such as explored above, namely, the opposing effects of u (at the denominator) and v (at the numerator) on the equilibrium erosion level (equation 4), and the fact that the genetic diversity at the *PRDM9* locus mostly depends on Nu.

Based on these analytical results under the model without dosage, we can now determine when dosage will play an important role. First, based on Eq 3, σ_0 should be large. Second, there should be enough time between successive invasions for eviction to take place. Thus, τ should be small. Altogether we predict a qualitative change induced by gene dosage on *PRDM9* standing diversity when Nu is large (i.e. the regime would have been polymorphic without dosage) and $\sigma_0\tau$ is also large (i.e. eviction due to homozygous advantage is sufficiently strong and has enough time to occur). Conversely, we predict a polymorphic regime when Nu is large and when the dosage effects are negligible, i.e. when $\sigma_0\tau$ is small.

These predictions were verified by systematically computing the mean value of $\sigma_0\tau$ across the scaling experiment on u and v. In all cases where a pronounced difference is observed between 6A and B (essentially for large u and small v), the criterion that $\sigma_0\tau$ should be large is met (*i.e.* is always greater than 3, Fig 6F). In the present case, σ_0 is not dependent on u and v, only τ depends on these parameters. This explains why, for constant u, gene dosage has a stronger impact for low v: a low v would imply a longer time between successive invasions (equation 6), giving more time for the frequency effect (second term of Eq 3) to operate

and thus pushing the Red Queen into the eviction regime. 422

In order to get additional insights on the impact of dosage, and more specifically on the role of σ_0, another 423
bi-dimensional scaling experiment was performed, this time allowing for the variation of the parameters d, 424
which is the mean number of DSBs per meiosis per chromosome pair, and $\bar{y}$, corresponding to the mean 425
value of the binding affinity distribution of all the *PRDM9* target sites (Fig 7). Here, a change of regime 426
is essentially found when these two parameters are low. It is also well predicted by the criterion that $\sigma_0\tau$ 427
should not be large (Fig 7C). Of note, both parameters, d and $\bar{y}$, play on σ_0, rather than on τ. Thus, if 428
DSBs are limiting or sites tend to be of low affinity, a change in dosage can make a big difference in the 429
chance of having at least one DSB in a symmetrically bound site. 430

Figure 7: **Scaling experiment showing the change of regime (from polymorphic to monomorphic) caused by the introduction of genetic dosage $N = 5 \times 10^3$.** A and B: equilibrium levels of *PRDM9* diversity without (A) and with (B) genetic dosage. C: summary statistic that is predictive of the type of regime observed in the presence of dosage, here $\sigma_0\tau$ (selection coefficient associated to dosage multiplied by the time between successive invasions by *PRDM9* alleles). The regime is predicted to be polymorphic in the presence of dosage if $\sigma_0\tau < 3$

These results show that gene dosage has important consequences on the Red Queen regime, by susbtantially 431
restricting the conditions under which a high *PRDM9* diversity can be observed: not just a high mutation 432
rate at this locus, but also a sufficiently short time between two invasions of alleles as well as a fairly weak 433
heterozygous disadvantage (or, equivalently, a weak haplo-insufficiency) in young alleles. 434

Empirical calibrations of the model 435

Empirical input parameters 436

Finally an empirical calibration of the model was attempted. The idea of this calibration is to try to match 437
the parameters of the model to known empirical values, based on current knowledge in mammals and, more 438
specifically, in the mouse, so as to see whether the model is able to at least roughly reproduce key empirical 439
observations, such as the typical erosion levels or genetic diversity at the *PRDM9* locus. 440

Focussing on the mouse, the effective population size (N_e) of different *Mus musculus* subspecies such as *Mus m. musculus* or *Mus m. domesticus* is around 10^5 [38]. The number of hostspots recognized per *PRDM9* alleles across the genome (h) is estimated between $15,000$ and $20,000$ [39]. Knowing that the mouse possesses 20 pairs of chromosomes, this corresponds approximately to $h \approx 1000$ sites per chromosome. The mean number of DSBs (d) performed per individual is estimated between 200 and 250 per meiosis [40], which correspond to approximately 10 DSBs per chromosome pair per meiosis. An exponential distribution for the affinities has a good fit to observed distribution obtained from Chip-seq experiments [28] (S5 Fig). The mean, however, is unknown but can be determined by assuming an average of $5,000$ targets bound per meiocyte [41] out of $20,000$ available targets. We obtain a mean of approximately 0.2.

Concerning the target mutation rate, considering that the mutation rate per nucleotide per generation across the mouse genome is $5.4.10^{-9}$ [42] and that the motifs characterising most of the hotspots in the mouse are several tens of nucleotides long [29], the inactivating mutation rate per generation (v) at the targets can be estimated at 10^{-7} [27]. The other mutation rate to determine is the one at the *PRDM9* locus (u). Owing to the mutational instability of the minisatellite encoding the zinc finger domain, this mutation rate is high and has been estimated at 10^{-5} in humans [21]. However, this is the raw mutation rate (including mutation that lead to either non-functional or functionally equivalent alleles). In contrast, the mutation rate u of the model is the net functional mutation rate (probability of creating a new functional allele recognizing entirely different targets), and the net rate is likely to be smaller than the raw experimental estimate. Accordingly, and as in Latrille *et al.* [27], we used $u = 3.10^{-6}$.

Finally, since a population size of $N = 10^5$ is too large for running the simulator, a standard scaling argument was used, by setting $N = 5.10^3$ and multiplying u and v by a factor 20 (i.e. using between $u = 6.10^{-5}$ to $u = 6.10^{-4}$ and $v = 2.10^{-6}$). The other parameters are set for the smallest chromosome in the mouse, so with lower h and d than the mean in the entire genome ($h = 800$, $d = 8$, $\bar{y} = 0.2$). This rescaling leaves approximately invariant the following quantities: the PRDM9 diversity D, the erosion level z or, equivalently, the proportion of active targets θ, the homozygous advantage σ and the selection coefficient experienced by new alleles s_0. The settings just presented represent our reference for empirical confrontation. Based on this reference, several variations of the model were also explored, which are described below.

Model predictions

The simulator was calibrated based on these rough empirical estimates for its parameters, then run and monitored for several key summary statistics, specifically, the genetic diversity of *PRDM9* (D), the proportion of eroded sites ($1-\bar{\theta}$), the mean haplo-insufficiency at the birth of the allele (σ_0), the mean haplo-insufficiency

over alleles sampled at the equilibrium regime ($\bar{\sigma}$) and the selection coefficient experienced by new *PRDM9*

alleles (s_0). The predictions are shown in Table 2 and examples of Red Queen dynamics are shown in Figs 8

and 9.

Table 2: **Empirical calibration experiments.**

u [a]	$\bar{y}$	d	c_{hom}	n_{mei}	D	$1-\bar{\theta}$	σ_0 [b]	$\bar{\sigma}$ [b]	s_0
3×10^{-6}	0.2	8	2	1	1	0.25	2.9×10^{-2}	5.2×10^{-2}	-2.2×10^{-2}
3×10^{-6}	0.44	8	1	1	2.8	0.05	1.5×10^{-5}	-4.9×10^{-7}	5.5×10^{-4}
3×10^{-6}	0.3	8	1.5	1	1.1	0.22	1×10^{-2}	1.5×10^{-2}	-6×10^{-3}
3×10^{-5}	0.2	8	2	1	1.2	0.23	2.9×10^{-2}	4.6×10^{-2}	-2.2×10^{-2}
3×10^{-6}	2	8	2	1	2.1	0.23	5.1×10^{-4}	7.8×10^{-4}	-8.5×10^{-5}
3×10^{-6}	0.2	8	2	5	2.2	0.22	2.9×10^{-2}	6.9×10^{-2}	6.5×10^{-6}
3×10^{-6}	0.2	24	2	1	2.5	0.26	5.7×10^{-5}	3.3×10^{-3}	1.5×10^{-4}

Equilibrium values of output summary statistics at equilibrium (*PRDM9* diversity D,mean level of erosion $\bar{z}$, mean haplo-insufficiency at the birth of the allele (σ_0), mean haplo-insufficiency at the equilibrium ($\sigma_{\bar{z}}$) and the mean selection coefficient experienced by new *PRDM9* alleles (s_0) as a function of the input parameters (mutation rate at *PRDM9* locus (u), mean of the affinity distribution ($\bar{y}$) , number of DSB par meiosis (d), dosage coefficient (c_{hom}) and maximum number of meiosis allowed for each individual), and with fixed values for the other parameters ($d = 8$, $h = 800$ and $v = 10^{-7}$)
[a] The values reported are before the down-scaling of the model, for the original population size of $N = 10^5$ (thus $4Nu = 1.2$ when $u = 3 \times 10^{-6}$ and $4Nu = 12$ when $u = 3 \times 10^{-5}$)
[b] Haplo-insufficiency is here formally defined as the relative difference in success rate of meiosis between homozygotes and hemizygotes. This is equivalent to the relative difference in fertility, except in the case where $n_{mei} = 5$, where fitness and rate of successful meiosis are not proportional.

Figure 8: **An example evolutionary trajectory under the fitness scheme allowing for only one meiosis per individual ($n_{mei} = 1$, and with parameters $u = 3 \times 10^{-6}$, $N = 10^5$ and $\bar{y} = 0.2$).** Variation through time of the frequency of each *PRDM9* allele (A) and their proportion of active sites (B).

The first line of Table 2 reports the results obtained from simulations run under the parameter values

corresponding to our reference for empirical comparison (see also Fig 8 for a typical simulation trajectory). ₄₇₆

The level of erosion $(1 - \bar{\theta})$ predicted by the model is consistent with what is known in the mouse (between ₄₇₇ 20% and 50% of erosion [1, 28, 29]). However, the predicted *PRDM9* diversity appears to be too low: these ₄₇₈ simulations result in a monomorphic regime (*i.e.* $D \sim 1$), whereas the *PRDM9* diversity observed in natural ₄₇₉ populations of mice is of the order of $D = 6$ to $D = 18$ [28] (See S1 Text). Of note, the diversity predicted ₄₈₀ by the model is the diversity of functionally different alleles (owing to the assumption made by our model of ₄₈₁ non-overlapping sets of targets for different alleles). In reality, closely related alleles can share a substantial ₄₈₂ fraction of their targets [28]. However, even accounting for this redundancy, the empirical functional diversity ₄₈₃ is still typically greater than 1, of the order of 2 to 6 in mouse subspecies [28] (See S1 Text). Thus, the ₄₈₄ monomorphic regime obtained here seems to be in contradiction with empirical observations. ₄₈₅

This low diversity can be explained by the fact that we are in a range of parameters involving an eviction ₄₈₆ regime due to gene dosage effects (compare Fig 8A and Fig 5A,B,C), such that one allele dominates in ₄₈₇ the population, while all the other alleles are counter-selected except during the short phases of allelic ₄₈₈ replacement (on average, the mean selection coefficient experienced by new alleles, s_0, is negative, Table 1). ₄₈₉

The fact that eviction is caused by gene dosage can be further verified by running the model without ₄₉₀ gene dosage (i.e. $c = 1$, Table 2, line 2), in which case higher levels of diversity are produced. Intermediate ₄₉₁ levels of gene dosage ($c = 1.5$, Table 2, line 3), on the other hand, give results that are essentially identical to ₄₉₂ those obtained with a dosage directly proportional to the number of gene copies ($c = 2$). Gene dosage also ₄₉₃ results in a non-negligible haplo-insufficiency, especially in old alleles (Table 2 column $\sigma_{\bar{z}}$), with a reduction ₄₉₄ of a few percents in the success of meiosis in hemizygotes (or, equivalently, heterozygotes with two alleles of ₄₉₅ same age), compared to homozygotes. Interestingly, such levels of haplo-insufficiency are comparable to ₄₉₆ those observed in old alleles in the mouse (B6, C3H [37]). The predicted haplo-insufficiency of young alleles ₄₉₇ σ_0 is weaker (Table 2). Such an age-dependency for the impact of gene dosage was previously suggested [1]. ₄₉₈ Nevertheless, at least in its current form and under those parameter values, the model does not predict an ₄₉₉ empirically reasonable regime. ₅₀₀

Increasing the *PRDM9* functional mutation rate (u) by as much as a factor 10 is not sufficient to get out ₅₀₁ of the eviction regime, resulting instead in a predicted *PRDM9* diversity still very close to one (fourth row ₅₀₂ of Table 2). Alternatively, increasing the mean affinity (or, equivalently, the concentration of PRDM9 in the ₅₀₃ cell) allows for higher levels of *PRDM9* equilibrium diversity while maintaining empirically reasonable levels ₅₀₄ of erosion (fifth row of Table 2). However, the mean number of sites bound by PRDM9 in a meiocyte would ₅₀₅ then predicted to be too large ($\sim 20,000$), compared to current empirical knowleldge ($\sim 5,000$ [41]). In ₅₀₆ addition, in this parameter regime, the model predicts very low levels of haplo-insufficiency ($\sigma_{\bar{z}} < 1 \times 10^{-3}$ ₅₀₇

for old alleles), substantially lower than empirically observed levels in the mouse ($\sigma_{\bar{z}} > 1 \times 10^{-2}$ [37]).   508

The model is naive in several other respects. First, gametes may often not be limiting and, as a result,   509
the fitness of an individual may not be proportional to the probability of success of meiosis, such as assumed   510
by the model thus far. A less-than-linear relation between fitness and success of meiosis can be modeled by   511
increasing the number of meiosis that an individual can attempt before being declared sterile. Allowing   512
for 5 attempts, thus mimicking a situation where an up to 80% reduction in the number of gametes would   513
not substantially affect the reproductive success of individuals (sixth row of Table 2), we observe a higher   514
functional diversity with a still reasonable level of erosion. The predicted reduction in meiosis success in   515
hemizygotes is also consistent with the ones observed empirically in mice (~ 2 to 5% [37]). However, the   516
fitness now reacts more weakly to variation in the success of meiosis, and as a result, the model is running   517
in an nearly-neutral regime, with a mean scaled selection coefficient acting on new alleles entering the   518
population (s_0) smaller than $1/N_e = 10^{-5}$, such that the turnover of recombination landscapes is mostly   519
driven by the neutral mutational turnover at the *PRDM9* locus. Although this could be seen as a possible   520
working regime for the evolutionary dynamics of recombination given the high mutation rate at this locus, it   521
is incompatible with the empirical support previously found for a positive selection acting on *PRDM9* [3,24].   522

Alternatively, the version of the model considered thus far assumes that the total number of DSBs   523
induced along the chromosome does not depend on the subsequent steps of the process. In reality, DSBs are   524
tightly regulated, in a way that may entail a negative feedback inhibiting the creation of further DSBs once   525
the chromosome has managed to synapse. For that reason, the mean number of DSBs per successful meiosis,   526
which is what is empirically measured [13], may be substantially smaller than the maximum number of DSBs   527
allowed before a meiocyte undergoes apoptosis. Yet, the success rate of meiosis depends on the maximum,   528
not on the mean number. As a way to indirectly account for this, we performed a last simulation allowing   529
for $d = 24$ DSBs (instead of 8, last row of Table 2, Fig 9). Running the model under this configuration   530
results in an evolutionary dynamics which is not in the eviction regime, with moderate levels of diversity   531
($D = 2.5$), reasonable levels of erosion ($\bar{z} = 0.26$), and strong positive selection on new *PRDM9* alleles   532
($s_0 > 10^{-4}$). On the other hand, the haplo-insuffiency predicted for old alleles is now weaker than that   533
observed empirically [37].   534

Altogether, these results show that the requirement of symmetric binding can result in a Red Queen   535
maintaining erosion at moderate levels under empirically reasonable parameter values. On the other hand,   536
there is only a narrow window for which eviction due to dosage is avoided and a sufficiently high *PRDM9*   537
diversity is maintained, while still having haplo-insufficiency for old alleles and strong selection acting on   538
PRDM9.   539

Figure 9: **An example evolutionary trajectory with 24 DSBs (d = 24, u = 3 × 10⁻⁶, N = 10⁵ and ȳ = 0.2).** Variation through time of the frequency of each *PRDM9* allele (A) and their proportion of active sites (B).

Discussion 540

Fundamental role of the symmetrical binding of PRDM9 541

The first studies on the function of *PRDM9* uncovered its role in targeting DSBs at specific sites, by its 542

capacity to set histone marks to recruit the DSB machinery [11, 43–45]. More recently, it was discovered that 543

this gene plays another important role during meiosis, namely, by facilitating chromosome pairing [1, 31]. In 544

this process, the symmetrical binding of PRDM9 to its target sites plays a key role. By bringing to the 545

chromosomal axis the sites on the homologue, PRDM9 presents them to the single-stranded DNA produced 546

by the resection around the DSB, thereby facilitating the search for the complementary sequence (Fig 547

1). This second function of *PRDM9*, combined with the differential erosion of the target sites between 548

sub-species (which is itself a direct consequence of the first function of *PRDM9*, the recruitment of the DSB 549

machinery), was found to be sufficient for explaining most aspects of the hybrid sterility pattern [1, 31]. 550

Here, we show that, similarly, these two aspects of the function of *PRDM9* during meiosis provide 551

the necessary ingredients for explaining the selective force promoting the turnvover of *PRDM9* alleles in 552

the context of a single population – thus giving a globally coherent picture bridging the gap between the 553

molecular mechanisms and the evolutionary dynamics of recombination. 554

Impact of gene dosage of *PRDM9* on the Red Queen dynamics

Gene dosage of *PRDM9* implies that a homozygote has a PRDM9 concentration for its allele which is twice that of each of the two alleles of a heterozygote. By the law of mass action, everything else being equal, increased dosage results in an increased occupancy of the targets and therefore an increased probability of symmetrical binding and a higher fertility. This clearly impacts the Red Queen dynamics for certain combinations of parameters, by acting against diversity and leading to a monomorphic regime characterized by the eviction of minor alleles (mostly found in a heterozygous state) by the currently dominant allele (mostly present in a homozygous state). The higher the selection coefficient associated to the homozygous advantage (captured here by σ_0), the stronger the effect against diversity. This impact of gene dosage on the evolutionary regime had been overseen thus far in the verbal and mathematical models of the Red Queen [18, 27], except in the more recent work of Baker *et al.* ([36], see below).

The impact of gene dosage is mitigated when PRDM9 is in high concentration in the cell. Indeed, a large concentration of PRDM9 makes the binding probability at a target of a given affinity less responsive to gene dosage (S6 Fig), thus reducing the fertility gap between a homo- and a heterozygote (S7 Fig). Based on these observations, we therefore hypothesize that the *PRDM9* expression level has been selected at a sufficiently high level so as to limit the impact of gene dosage. This is an interesting hypothesis that could be studied in the future.

Comparison with Baker *et al.*'s model

In a recent article, Baker *et al.* [36] also propose a mechanistic model of the *PRDM9* Red Queen. Several of their conclusions agree with ours. Most notably, Baker *et al.* also recognize that the symmetric binding of PRDM9 provides the key ingredient for mediating positive selection on PRDM9 in the context of the Red Queen, and they incorporate the consequences of gene dosage in their model. However, the two studies differ on some important points.

First, in our model, the amount of PRDM9 protein is not limiting (that is, most PRDM9 molecules are not bound to the target sites), whereas Baker *et al.* assume strong competition between targets for PRDM9 proteins (most molecules are bound). Of note, gene dosage and competition are two distinct aspects of the steady-state equilibrium of PRDM9 binding. Gene dosage is a direct consequence of the law of mass action (site occupancy is an increasing function of the free concentration of PRDM9, see S1 Fig), a phenomenon that can happen even when PRDM9 is not limiting, as shown here. Competition between targets, on the other hand, relates to the fact that the inactivation of some of the binding sites results in extra PRDM9 molecules available for other sites to bind, an effect that is negligible if PRDM9 is in excess (as in our model),

but important otherwise (as in Baker *et al.*'s model [36]). Whether PRDM9 is limiting ultimately depends 586 on its affinity for its targets, which is know to be high [46], but also on the total concentration of PRDM9 in 587 the cell, which, to our knowledge, is still unknown, and finally, on the total number of targets in the genome. 588 Altogether, the question thus still remains open. In any case, our model shows that competition between 589 targets is not a necessary feature to drive the evolutionary dynamics of *PRDM9*. 590

The second difference concerns the affinity distribution. Baker *et al.* use a two-heat distribution. Here 591 we use an exponential distribution (S5 Fig), which is motivated by results from Chip-seq experiments [28], 592 although with some uncertainty regarding the shape of the distribution in the low affinity range, which is 593 not captured by Chip-seq. The use of these different affinity distributions has clear consequences on the 594 behavior of models with respect to dosage. Indeed, increasing the gene dosage can have two opposing effects. 595 On the one hand, a higher dosage increases the symmetrical binding at sites of intermediate affinity. On the 596 other hand, it also results in more sites of low affinity being bound, and this, most often asymmetrically. 597 Depending on the balance between these two effects, different outcomes are obtained. In our case, where the 598 affinity distribution has a moderate variance, the increased symmetrical binding of sites of intermediate 599 affinity always wins. As a result, there is always an advantage to increasing dosage. That is, homozygotes 600 always have an advantage over heterozygotes, which creates an eviction regime that tends to play against 601 diversity. A contrario, in Baker's model, the two-heat affinity distribution results in a non-monotonous 602 dependency, with a turning point, such that the dosage is in favor of homozygotes for young alleles, but in 603 favor of heterozygotes for old alleles. This turning point is one potential solution to the problem of eviction. 604 Here, we present an alternative solution, which is captured in our model by the statistics $\sigma_0 \tau$. Intuitively, 605 eviction does not take place if the homozygote advantage is sufficiently weak and erosion sufficiently rapid, 606 such that ageing alleles are replaced before eviction has enough time to take place. 607

As it turns out, it is not so easy to find empirically reasonable parameter configurations such that the 608 eviction regime is avoided, although this could be the consequence of other aspects of the biology of meiosis 609 and reproduction being missed by the model. On the other hand, our model predicts that old alleles should 610 be haplo-insufficient, thus in agreement with what is observed in the mouse (for allele B6 and C3H [37]). This 611 haplo-insufficiency of old alleles is not observed systematically in Baker's *et al.* model, but only occasionally 612 and in small populations. 613

Altogether it would be useful to empirically investigate the haplo-insufficiency for young and old alleles 614 over a broader range of alleles and species, in order to determine whether a lifelong homozygous advantage is 615 systematic or occasional. More globally, it will be important to unravel the exact roles of affinity distribution, 616 PRDM9 concentration and competition between targets in the evolutionary dynamic. Finally, all this raises 617

the question of a possible evolution of the *PRDM9* expression level, the affinity (whether in its distribution or in its mean) and the number of targets recognized per allele.

Current limitations and perspectives

In addition to those discussed above, the model introduced here has other limitations. First, in its current form, the model implements only limited variance among alleles in their strength at birth. Yet, in reality, some alleles are dominant over others [47], such that, in a heterozygote for a strong and a weak allele, DSBs are more often produced at the target sites of the stronger allele. Of note, although some dominance is expected to emerge purely as a consequence of erosion (with younger alleles being on average dominant over older ones), a substantial part of it appears to be instead related to intrinsic differences between alleles regardless of their age [15, 48]. If stronger alleles have an advantage over weaker alleles, either because they promote a higher binding symmetry or just because of their higher penetrance, then weaker alleles should be less likely to invade. As a result, the population would tend to be dominated by stronger alleles. In good approximation, this would amount to running the model under a lower effective mutation rate (now to be taken as the rate at which new functional and sufficiently strong *PRDM9* alleles are being produced by mutation) but otherwise would not fundamentally change the evolutionary dynamic.

Second, the model currently assumes that all DSBs are repaired with the homologue. In reality, some are repaired with the sister chromatid [15]. This simplifying assumption, however, should have a moderate impact on our conclusions, as it essentially amounts to a change in the rate of erosion, which can be accounted for by considering a lower mutation rate v at the target sites. Our experiments suggest that small changes in v do not fundamentally impact the dynamic.

Third, in addition to implementing the idea that symmetrical binding of PRDM9 is required for successful pairing of chromosomes, our model also more specifically assigns the (obligate and unique) CO to one of the DSBs occurring in symmetrically bound sites. In hindsight, this specific choice is not warranted by current empirical evidence, which only suggests that repair of DSBs with the homologue, and thus ultimately chromosome pairing, is facilitated by symmetrical binding of PRDM9, but this, independently of the choice of CO versus NCO among all DSBs. However, this specific feature of our model should have a minor impact on our conclusions. Choosing instead the CO uniformly at random among all DSBs (but still requiring at least one DSB in a symmetrically bound site) would merely change the patterns of linkage dissipation, and this, in a context where linkage is unlikely to be a major factor.

Although deserving more careful examination, the limitations just mentioned are therefore probably minor. Perhaps a more fundamental issue is to fully understand how the eviction regime induced by gene

dosage effects can be avoided, so as to predict reasonable levels of *PRDM9* diversity, while having a Red 649
Queen regime driven by strong positive selection on *PRDM9*. As it stands, the model appears to be stuck in 650
a dilemma, such that either dosage has a substantial impact on fertility, but then eviction takes place and 651
the model fails at explaining empirically observed levels of *PRDM9* diversity, or the effects of dosage are 652
made weaker by assuming globally less stringent conditions for achieving a good fertility (such as allowing 653
for an excess in PRDM9 protein, or in gametes or in DSBs), but then the fitness differences between old 654
and new alleles also become small and the model approaches a nearly-neutral regime, in which the turnover 655
of *PRDM9* alleles is primarily driven by the mutational turnover at the locus. Although the latter regime 656
may not be so unreasonable from an evolutionary point of view, it fails at explaining the positive selection 657
observed on *PRDM9*, or its haplo-insufficiency, depending on the detailed model configuration (last two 658
rows of Table 2). 659

This dilemma is still in need of a robust and convincing explanation. In this respect, as mentioned above, 660
the questions of the affinity distribution and the concentration of PRDM9 relative to the affinity of the 661
target sites should be explored in depth. Ultimately, a more mechanistically motivated model of PRDM9 662
binding, accounting for the combinatorial sequence features susceptible to determine the affinity, could 663
provide useful clues as to what would be a reasonable affinity distribution. It would also naturally lead to a 664
more realistic model in which point mutations may only partially inactivate the targets [17], as opposed to 665
completely inactivate them as in our current model. 666

Alternatively, the empirical regime of the Red Queen may possibly alternate between long nearly-neutral 667
epochs, with occasional bouts of positive selection whenever the current alleles become too eroded. Such 668
occasional epsiodes of positive selection could be sufficient to induce the empirically observed patterns of 669
accelerated evolution of the zinc finger domain at the non-synonymous level. The impact of Hill-Robertson 670
interference and its dissipation on the evolutionary dynamic of *PRDM9* could also contribute to maintaining 671
a high diversity at the *PRDM9* locus, a point that may also need to be investigated. Finally, population 672
structure could play a role, by maintaining different pools of alleles in different sub-populations connected 673
by recurrent migration, thus resulting in a large diversity at the metapopulation level, and this, in spite of 674
non-negligible gene dosage effects. Population structure is also pointing toward the other big question still 675
in need of a model-based exploration: the potential role of *PRDM9* in hybrid sterility and speciation. 676

Materials and methods

The model

The simulation model is graphically summarized in Fig 2, and its key parameters are listed in Table 1.
This model assumes of a population of N diploid individuals, whose genetic map is composed of a single
chromosome. A *PRDM9* locus is located on this chromosome. Each *PRDM9* allele has a set of h binding
sites, each of which has an intrinsic binding affinity for its cognate PRDM9 protein. The positions of binding
sites along the chromosome are drawn uniformly at random and their affinities are drawn according to an
exponential law of parameter $\bar{y}$ (S5 Fig). Each target site is associated to a unique allele and different sites
cannot overlap. In practice, this is implemented by encoding the chromosome as an array of L slots, such
that, upon creation of a new *PRDM9* allele by mutation, each binding site of this allele chooses one of the
available slots uniformly at random. Given the composition of the population of the current generation, the
next generation, of the same population size, is generated as follows.

Mutations

The *PRDM9* locus mutates with a probability u per gene copy. Given a mutation, a new functional *PRDM9*
allele is created to which are associated h new sites along the genome, according to the procedure just
described. Next, each target site recognized by each allele currently present in the population mutates with a
probability v. This type of mutation results in a complete inactivation of the target site. Of note, all target
sites are assumed to be monomorphic for the active variant at the time of the birth of the corresponding
PRDM9 allele. As a result, the loss of target sites is entirely contributed by gene conversion acting on
inactivating mutations that have occurred after the birth of the allele. Also, we neglect new target sites that
might arise by mutation, which is reasonable since hotspot alleles newly arisen by mutation would be at a
very high risk of being lost by conversion.

Meiosis and reproduction

A meiosis is attempted on a randomly selected individual, according to the following steps. Of note, the
meiosis can fail (in which case we assume that the meiocyte undergoes apoptosis) for multiple reasons, all of
which will be described below. In the simulations presented here, unless stated otherwise, whenever meiosis
fails, then a new individual is chosen at random from the population and a new meiosis is attempted.

First, the two homologous chromosomes are replicated, thus creating a set of 4 chromatids. Then, PRDM9
binds to its target site according to an occupation probability determined by the chemical equilibrium. A

binding site i has an affinity

$$K_i = \frac{[PS_i]}{[P]_{free}[S_i]} = \frac{x_i}{[P]_{free}(1 - x_i)},$$ (7)

where $[PS_i] = x_1$ and $[S_i] = 1 - x_i$ are the proportions of target site i (across meiocytes) which are occupied by PRDM9 or free, respectively. Thus:

$$x_i = \frac{[P]_{free}K_i}{1 + [P]_{free}K_i}.$$ (8)

We assume that PRDM9 is not limiting, meaning that most PRDM9 molecules are free $[P]_{free} \approx [P]_{tot}$; there is therefore a total absence of competition between binding sites. Thus, if we define the rescaled affinity as $y_i = [P]_{tot}K_i$, then the occupancy probability can be re-written :

$$x_i = \frac{y_i}{1 + y_i}.$$ (9)

Based on this equation, for each target site, binding is randomly determined, by drawing a Bernoulli random variable of parameter x_i for site i. Of note, at a given target locus, there are four instances of the binding site (one on each of the two sister chromatids for each of the two homologues), and binding is determined independently for each of those instances.

Once the occupation status of all binding sites has been determined, the total number k of sites bound by PRDM9 over the four chromatids, is calculated. If $k = 0$, meiosis fails. Otherwise, each site bound by PRDM9 undergoes a DSB with a probability equal to $p = \min\left(1, \frac{d}{k}\right)$, with d being a parameter of the model. In most experiments, we use $d = 6$. Thus, in the most frequent case where $k > d$, an average number of d DSBs are being produced. This procedure aims to model the regulation of the total number of DSBs through the genome, which in mammals seems to be independent from PRDM9 binding [1, 13].

Next, the four chromatids are scanned for symmetrically bound sites, which the model assumes are essential for chromosome pairing (see introduction). If no such site is detected, meiosis fails. Otherwise, one of the symmetrically bound sites is uniformly picked at random and becomes the initiation site for a CO. Thus, only one CO is performed per meiosis (and this, in order to model CO interference), and all other DSBs are repaired as NCO events. Note that in our model we do not allow for the possibility of DSB repair by the sister chromatid.

A successful meiosis therefore produces 4 chromatids, with exactly one CO between two of them, and some events of gene conversion at all sites that have undergone a DSB. Of note, in the presence of inactive

versions of the binding sites (created by mutations, see above), these gene conversion events are effectively 730
implementing the hotspot conversion paradox [16]. Finally one chromatid is randomly picked among the 731
tetrad to become a gamete. The whole procedure is repeated, starting from the choice of a new individual 732
until another gamete is obtained. Then these two gametes merge and create a new diploid individual of the 733
next generation. All these steps, from the choice of an individual until the choice of a chromatid among the 734
tetrad, are performed as many time as needed until the complete fill of the next generation. 735

Gene dosage 736

In the case where the gene dosage of PRDM9 is taken into account, it is assumed that, for a given *PRDM9* 737
allele, a homozygote will produce twice the concentration of the corresponding protein compared to a 738
heterozygote. By assuming that $[P]_{tot}^{homo} = 2[P]_{tot}^{het}$, the probability that a PRDM9 protein binds to a site i 739
of rescaled affinity y_i is 740

$$x_i = \frac{cy_i}{1 + cy_i} \tag{10}$$

where $c = 1$ for a heterozygote and $c = 2$ for a homozygote. 741

Summary statistics 742

Several summary statistics were monitored during the run of the simulation program. They were used, first, 743
to evaluate the time until convergence to the equilibrium regime (burn-in). The burn-in was taken in excess, 744
so as to be adapted to all simulation settings. In practice, the burn-in is set at 10000 generations over a total 745
of $50,000$ generations. Second, the summary statistics were averaged over the entire run (burn-in excluded), 746
thus giving a quantitative characterization of the equilibrium regime, as a function of the model parameters. 747
The main statistics are the *PRDM9* diversity in the population (D), the mean proportion of binding sites 748
that are still active per allele θ, the mean symmetrical binding probability (q) and the mean fertility (w). 749

Diversity is defined as the effective number of *PRDM9* alleles, or in other words as the inverse of the 750
homozygosity [27]. At time t: 751

$$D_t = \frac{1}{\sum_i f_{i,t}^2} \tag{11}$$

where $f_{i,t}$ is the frequency of allele i at time t. With this definition, when K alleles segregate each at 752
frequency $\frac{1}{K}$ in the population, the diversity D is equal to K. Averaging Eq 11 over the simulation run 753

gives the average diversity D.

The mean proportion of binding sites that are still active per allele is defined as:

$$\theta_t = \sum_i f_{i,t} \theta_{i,t} \tag{12}$$

where $\theta_{i,t}$ is the proportion of sites that are still active for allele i at time t. Averaging θ_t over the simulation trajectory gives $\bar{\theta}$.

The probability of symmetrical binding q corresponds to the mean probability of having a PRDM9 protein bound on at least one of the two chromatids of the homologous chromosome at a certain position, given that a DSB has occured at this very position. In the case of a homozygous individual for allele i, this quantity can be obtained analytically for a given complete diploid genotype and is given by:

$$q_{i,t}^{hom} = \frac{2 < x^2 > - < x^3 >}{< x >}, \tag{13}$$

where $x = \frac{cy}{1+cy}$ is the occupancy of a binding site of affinity y and, for any k, $< x^k >$ is the mean of x^k over all sites, taking into account their affinity distribution. In the case of a heterozygous individual, with two *PRDM9* alleles i and j, the symmetrical binding rate is:

$$q_{i,j,t}^{het} = \frac{2 < x^2 >_i - < x^3 >_i + 2 < x^2 >_j - < x^3 >_j}{< x >_i + < x >_j}, \tag{14}$$

where the subscripts i and j correspond to averages over sites of allele i and j, respectively. For allele i, these statistics were then averaged over all diploid backgrounds for this allele (either homo- or heterozygotes), thus against each allele j present in the population at a given generation, yielding the mean symmetrical binding rate for allele i at time t, $q_{i,t}$. Finally, this was averaged over all alleles (weighted by their frequency), and over the simulation run, yielding the overall population-level equilibrium mean value of q.

For a given genotype, the fertility can be computed analytically. Here we assumed that fertility in proportional to the rate of success of meiosis. Thus, it is equivalent to 1-(the mean probability of failure of meiosis) characterized by the absence of a DSB in a symmetric site. In turn, the number of DSBs in a symmetric site follows a Poisson law with parameter dq where d is the average number of DSBs per meiocyte and q is the mean probability of symmetrical binding for this allele. In the case of a homozygous individual for allele i, this quantity can be obtained analytically for a given complete diploid genotype and is given by:

$$w_{i,t}^{hom} = 1 - e^{-dq_{i,t}^{hom}} \tag{15}$$

In the case of a heterozygous individual, with two *PRDM9* alleles i and j, the fertility rate is:

$$w_{i,j,t}^{het} = 1 - e^{-dq_{i,j,t}^{het}} \tag{16}$$

where the subscripts i and j correspond to averages over sites of allele i and j, respectively. As for the symmetrical binding rate, for each allele i, these statistics were averaged over all diploid backgrounds for this allele (either homo- or heterozygotes), against all other alleles present in the population at a given generation, yielding the fertility rate for allele i at time t, $w_{i,t}$. Finally, this was averaged over all alleles (weighted by their frequency), and over the simulation run, yielding the overall population-level equilibrium mean value of w.

The haplo-insufficiency coefficient for an allele i at time t $\sigma_{i,t}$ is defined as the relative difference in fertility between homozygotes (w^{hom}) and hemizygotes (w^{hemi}) for this specific allele:

$$\sigma_{i,t} = \frac{w_{i,t}^{hom} - w_{i,t}^{hemi}}{w_{i,t}^{hemi}}. \tag{17}$$

Here, the hemizygote fertility is given by:

$$w_{i,t}^{hemi} = 1 - e^{-dq_{i,t}^{hemi}} \tag{18}$$

and

$$q_{i,t}^{hemi} = \frac{2 <x^2> - <x^3>}{<x>}, \tag{19}$$

Of note, although Eq 19 and 13 look identical, the site occupancies x are different, owing to the differing concentrations of the protein corresponding to allele i when in one or two copies. This can again be averaged over all alleles in the population and over the run, yielding $\bar{\sigma}$.

The haplo-insufficiency coefficient of a new allele, σ_0, is defined similarly as for segregating alleles (Eq 17), although its actual computation proceeds differently. Taking advantage of the fact that this coefficient does not entail any average over the genetic backgrounds segregating in the population, it is obtained independently of the simulation run, by randomly creating new alleles (and their corresponding arrays of binding sites over the genome) and then computing the symmetrical binding rate, and then the fertility, assuming one or two copies for this allele. This is repeated 100 times, and the average over these 100 alleles yields an estimate of σ_0.

Similarly, the selection coefficient for new alleles arising in the population, s_0, is defined, and computed, as the difference between the log fitness of a new allele (created on the fly and combined with all genetic backgrounds segregating in the population) and the log fitness averaged over the whole population. The resulting time-dependent estimate is then averaged over the simulation run.

Scaling experiments

In order to visualize how the equilibrium regime varies according to the model parameters, first, a central parameter configuration was chosen, which will represent a fixed reference across all scaling experiments. Then only one parameter at a time (or two for bi-dimensional scaling) is systematically varied over a range of regularly spaced values on both sides of the reference configuration. This parameter is fixed along the entire simulation and is variable only between simulations. For each parameter value over this range, a complete simulation is run. Once the equilibrium is reached for a given simulation setting, the summary statistics described above are averaged over the entire trajectory, excluding the burn-in. These mean equilibrium values, which characterize and quantify the stationary regime of the model, were finally plotted as a function of the varying parameter(s).

The central parameters were chosen as follows. First, the parameters of population size N was set to $5,000$, corresponding to the maximum population size that can be afforded computationally. Then the number of sites recognized as target sites by each PDRM9 allele's protein was set to $h = 400$ in order to get closer to the average number of target sites found on the smallest chromosome of the mouse (around 800 sites) while limiting the memory requirements. The mean for the affinity distribution of the target was set to $\bar{y} = 6$. The parameter d representing the number average of DSBs in a meiocyte is set at 6 which corresponds to the approximate number of DSBs found on the smallest chromosome of the mouse. For uni-dimensional scaling, the parameter v, the mutation rate at target sites, was set to 2×10^{-6} and the parameter u was set to 5×10^{-5}. For the bi-dimensional scaling of d and $\bar{y}$, $u = 2 \times 10^{-4}$ and $v = 5 \times 10^{-5}$.

Data accessibility

The model, and the codes to generate the figures can be found at `https://github.com/alicegenestier/` `Red_Queen_PRDM9_Panmictic.git` or on Zenodo with all the datasets [49]. The model was implemented in C++ and all figures were created using Python.

Acknowledgments 824

We would like to thank Corinne Grey and Frédéric Baudat for their comments on the manuscript, as well as 825 Zachary Baker for helpful discussions. All simulations of this work were performed using the computing 826 facilities of the CC LBBE/PRABI. 827

Competing interest 828

The authors have declared that no competing interests exist. 829

Funding 830

This work was funded by the Agence Nationale de la Recherche, Grant ANR-19-CE12-0019 / HotRec. This 831 funder covered the publication costs and had no role in study design, data collection and analysis, decision 832 to publish, or preparation of the manuscript. Alice Genestier has a PhD scholarship contributed by the 833 French Ministry of Research and Higher Education. 834

References

[1] Davies B, Hatton E, Altemose N, Hussin JG, Pratto F, Zhang G, et al. Re-engineering the zinc fingers of PRDM9 reverses hybrid sterility in mice. Nature. 2016;530(7589):171–176. doi:10.1038/nature16931.

[2] Mihola O, Trachtulec Z, Vlcek C, Schimenti JC, Forejt J. A Mouse Speciation Gene Encodes a Meiotic Histone H3 Methyltransferase. Science. 2009;323(5912):373–375. doi:10.1126/science.1163601.

[3] Oliver PL, Goodstadt L, Bayes JJ, Birtle Z, Roach KC, Phadnis N, et al. Accelerated Evolution of the Prdm9 Speciation Gene across Diverse Metazoan Taxa. PLOS Genetics. 2009;5(12):e1000753. doi:10.1371/journal.pgen.1000753.

[4] Myers S, Bottolo L, Freeman C, McVean G, Donnelly P. A Fine-Scale Map of Recombination Rates and Hotspots Across the Human Genome. Science. 2005;310(5746):321–324. doi:10.1126/science.1117196.

[5] Jeffreys AJ, Kauppi L, Neumann R. Intensely punctate meiotic recombination in the class II region of the major histocompatibility complex. Nature Genetics. 2001;29(2):217–222. doi:10.1038/ng1001-217.

[6] Kauppi L, Jeffreys AJ, Keeney S. Where the crossovers are: recombination distributions in mammals. Nature Reviews Genetics. 2004;5(6):413–424. doi:10.1038/nrg1346.

[7] McVean GAT, Myers SR, Hunt S, Deloukas P, Bentley DR, Donnelly P. The Fine-Scale Structure of Recombination Rate Variation in the Human Genome. Science. 2004;304(5670):581–584. doi:10.1126/science.1092500.

[8] Brunschwig H, Levi L, Ben-David E, Williams RW, Yakir B, Shifman S. Fine-Scale Maps of Recombination Rates and Hotspots in the Mouse Genome. Genetics. 2012;191(3):757–764. doi:10.1534/genetics.112.141036.

[9] Baudat F, Buard J, Grey C, Fledel-Alon A, Ober C, Przeworski M, et al. PRDM9 Is a Major Determinant of Meiotic Recombination Hotspots in Humans and Mice. Science. 2010;327(5967):836–840. doi:10.1126/science.1183439.

[10] Myers S, Bowden R, Tumian A, Bontrop RE, Freeman C, MacFie TS, et al. Drive Against Hotspot Motifs in Primates Implicates the PRDM9 Gene in Meiotic Recombination. Science. 2010;327(5967):876–879. doi:10.1126/science.1182363.

[11] Hayashi K, Yoshida K, Matsui Y. A histone H3 methyltransferase controls epigenetic events required for meiotic prophase. Nature. 2015;438(7066):374–378. doi:10.1038/nature04112.

[12] Powers NR, Parvanov ED, Baker CL, Walker M, Petkov PM, Paigen K. The Meiotic Recombination Activator PRDM9 Trimethylates Both H3K36 and H3K4 at Recombination Hotspots In Vivo. PLOS Genetics. 2016;12(6):e1006146. doi:10.1371/journal.pgen.1006146.

[13] Lange J, Yamada S, Tischfield SE, Pan J, Kim S, Zhu X, et al. The Landscape of Mouse Meiotic Double-Strand Break Formation, Processing, and Repair. Cell. 2016;167(3):695–708.e16. doi:10.1016/j.cell.2016.09.035.

[14] Grey C, Baudat F, Massy Bd. PRDM9, a driver of the genetic map. PLOS Genetics. 2018;14(8):e1007479. doi:10.1371/journal.pgen.1007479.

[15] Li R, Bitoun E, Altemose N, Davies RW, Davies B, Myers SR. A high-resolution map of non-crossover events reveals impacts of genetic diversity on mammalian meiotic recombination. Nature Communications. 2019;10:3900. doi:10.1038/s41467-019-11675-y.

[16] Boulton A, Myers RS, Redfield RJ. The hotspot conversion paradox and the evolution of meiotic recombination. Proceedings of the National Academy of Sciences. 1997;94(15):8058–8063. doi:10.1073/pnas.94.15.8058.

[17] Lesecque Y, Glémin S, Lartillot N, Mouchiroud D, Duret L. The Red Queen Model of Recombination Hotspots Evolution in the Light of Archaic and Modern Human Genomes. PLOS Genetics. 2014;10(11):e1004790. doi:10.1371/journal.pgen.1004790.

[18] Úbeda F, Wilkins JF. The Red Queen theory of recombination hotspots. Journal of Evolutionary Biology. 2011;24(3):541–553. doi:10.1111/j.1420-9101.2010.02187.x.

[19] Van Valen L. Molecular evolution as predicted by natural selection. Journal of Molecular Evolution. 1974;3(2):89–101. doi:10.1007/BF01796554.

[20] Auton A, Fledel-Alon A, Pfeifer S, Venn O, Ségurel L, Street T, et al. A Fine-Scale Chimpanzee Genetic Map from Population Sequencing. Science. 2012;336(6078):193–198. doi:10.1126/science.1216872.

[21] Jeffreys AJ, Cotton VE, Neumann R, Lam KWG. Recombination regulator PRDM9 influences the instability of its own coding sequence in humans. Proceedings of the National Academy of Sciences. 2013;110(2):600–605. doi:10.1073/pnas.1220813110.

[22] Berg IL, Neumann R, Sarbajna S, Odenthal-Hesse L, Butler NJ, Jeffreys AJ. Variants of the protein PRDM9 differentially regulate a set of human meiotic recombination hotspots highly active in African populations. Proceedings of the National Academy of Sciences of the United States of America. 2011;108(30):12378–12383. doi:10.1073/pnas.1109531108.

[23] Kono H, Tamura M, Osada N, Suzuki H, Abe K, Moriwaki K, et al. Prdm9 Polymorphism Unveils Mouse Evolutionary Tracks. DNA Research. 2014;21(3):315–326. doi:10.1093/dnares/dst059.

[24] Buard J, Rivals E, Segonzac DDd, Garres C, Caminade P, Massy Bd, et al. Diversity of Prdm9 Zinc Finger Array in Wild Mice Unravels New Facets of the Evolutionary Turnover of this Coding Minisatellite. PLOS ONE. 2014;9(1):e85021. doi:10.1371/journal.pone.0085021.

[25] Vara C, Capilla L, Ferretti L, Ledda A, Sánchez-Guillén RA, Gabriel SI, et al. PRDM9 Diversity at Fine Geographical Scale Reveals Contrasting Evolutionary Patterns and Functional Constraints in Natural Populations of House Mice. Molecular Biology and Evolution. 2019;36(8):1686–1700. doi:10.1093/molbev/msz091.

[26] Alleva B, Brick K, Pratto F, Huang M, Camerini-Otero RD. Cataloging Human PRDM9 Allelic Variation Using Long-Read Sequencing Reveals PRDM9 Population Specificity and Two Distinct Groupings of Related Alleles. Frontiers in Cell and Developmental Biology. 2021;9.

[27] Latrille T, Duret L, Lartillot N. The Red Queen model of recombination hot-spot evolution: a theoretical investigation. Philosophical Transactions of the Royal Society B: Biological Sciences. 2017;372(1736):20160463. doi:10.1098/rstb.2016.0463.

[28] Smagulova F, Brick K, Pu Y, Camerini-Otero RD, Petukhova GV. The evolutionary turnover of recombination hot spots contributes to speciation in mice. Genes & Development. 2016;30(3):266–280. doi:10.1101/gad.270009.115.

[29] Baker CL, Kajita S, Walker M, Saxl RL, Raghupathy N, Choi K, et al. PRDM9 Drives Evolutionary Erosion of Hotspots in Mus musculus through Haplotype-Specific Initiation of Meiotic Recombination. PLOS Genetics. 2015;11(1):e1004916. doi:10.1371/journal.pgen.1004916.

[30] Gregorová S, Forejt J. PWD/Ph and PWK/Ph Inbred Mouse Strains of Mus

cyrchar

cyrt. musculus Subspecies-a Valuable Resource of Phenotypic Variations and Genomic Polymorphisms. Folia biologica. 2000;46:31–41.

[31] Gregorova S, Gergelits V, Chvatalova I, Bhattacharyya T, Valiskova B, Fotopulosova V, et al. Modulation of Prdm9-controlled meiotic chromosome asynapsis overrides hybrid sterility in mice. eLife. 2018;7:e34282. doi:10.7554/eLife.34282.

[32] Grey C, Clément JAJ, Buard J, Leblanc B, Gut I, Gut M, et al. In vivo binding of PRDM9 reveals interactions with noncanonical genomic sites. Genome Research. 2017;27(4):580–590. doi:10.1101/gr.217240.116.

[33] Parvanov ED, Tian H, Billings T, Saxl RL, Spruce C, Aithal R, et al. PRDM9 interactions with other proteins provide a link between recombination hotspots and the chromosomal axis in meiosis. Molecular Biology of the Cell. 2017;28(3):488–499. doi:10.1091/mbc.E16-09-0686.

[34] Borde V, de Massy B. Programmed induction of DNA double strand breaks during meiosis: setting up communication between DNA and the chromosome structure. Current Opinion in Genetics & Development. 2013;23(2):147–155. doi:10.1016/j.gde.2012.12.002.

[35] Biot M, Toth A, Brun C, Guichard L, Massy Bd, Grey C. Fundamental principles of chromosome organization for meiotic recombination; 2024. Available from: `https://www.biorxiv.org/content/10.1101/2024.03.06.583329v1`.

[36] Baker Z, Przeworski M, Sella G. Down the Penrose stairs, or how selection for fewer recombination hotspots maintains their existence. eLife. 2023;12:e83769. doi:10.7554/eLife.83769.

[37] Baker CL, Petkova P, Walker M, Flachs P, Mihola O, Trachtulec Z, et al. Multimer Formation Explains Allelic Suppression of PRDM9 Recombination Hotspots. PLoS Genetics. 2015;11(9):e1005512. doi:10.1371/journal.pgen.1005512.

[38] Salcedo T, Geraldes A, Nachman MW. Nucleotide Variation in Wild and Inbred Mice. Genetics. 2007;177(4):2277–2291. doi:10.1534/genetics.107.079988.

[39] Brick K, Smagulova F, Khil P, Camerini-Otero RD, Petukhova GV. Genetic recombination is directed away from functional genomic elements in mice. Nature. 2012;485(7400):642–645. doi:10.1038/nature11089.

[40] Kauppi L, Barchi M, Lange J, Baudat F, Jasin M, Keeney S. Numerical constraints and feedback control of double-strand breaks in mouse meiosis. Genes & Development. 2013;27(8):873–886. doi:10.1101/gad.213652.113.

[41] Baker CL, Walker M, Kajita S, Petkov PM, Paigen K. PRDM9 binding organizes hotspot nucleosomes and limits Holliday junction migration. Genome Research. 2014;24(5):724–732. doi:10.1101/gr.170167.113.

[42] Uchimura A, Higuchi M, Minakuchi Y, Ohno M, Toyoda A, Fujiyama A, et al. Germline mutation rates and the long-term phenotypic effects of mutation accumulation in wild-type laboratory mice and mutator mice. Genome Research. 2015;25(8):1125–1134. doi:10.1101/gr.186148.114.

[43] Wu H, Mathioudakis N, Diagouraga B, Dong A, Dombrovski L, Baudat F, et al. Molecular Basis for the Regulation of the H3K4 Methyltransferase Activity of PRDM9. Cell Reports. 2013;5(1):13–20. doi:10.1016/j.celrep.2013.08.035.

[44] Eram MS, Bustos SP, Lima-Fernandes E, Siarheyeva A, Senisterra G, Hajian T, et al. Trimethylation of Histone H3 Lysine 36 by Human Methyltransferase PRDM9 Protein. The Journal of Biological Chemistry. 2014;289(17):12177–12188. doi:10.1074/jbc.M113.523183.

[45] Grey C, Barthès P, Friec GCL, Langa F, Baudat F, Massy Bd. Mouse PRDM9 DNA-Binding Specificity Determines Sites of Histone H3 Lysine 4 Trimethylation for Initiation of Meiotic Recombination. PLOS Biology. 2011;9(10):e1001176. doi:10.1371/journal.pbio.1001176.

[46] Striedner Y, Schwarz T, Welte T, Futschik A, Rant U, Tiemann-Boege I. The long zinc finger domain of PRDM9 forms a highly stable and long-lived complex with its DNA recognition sequence. Chromosome Research. 2017;25(2):155–172. doi:10.1007/s10577-017-9552-1.

[47] Diagouraga B, Clément JAJ, Duret L, Kadlec J, Massy Bd, Baudat F. PRDM9 Methyltransferase Activity Is Essential for Meiotic DNA Double-Strand Break Formation at Its Binding Sites. Molecular Cell. 2018;69(5):853–865.e6. doi:10.1016/j.molcel.2018.01.033.

[48] Hinch AG, Zhang G, Becker PW, Moralli D, Hinch R, Davies B, et al. Factors Influencing Meiotic Recombination Revealed by Whole Genome Sequencing of Single Sperm. Science (New York, NY). 2019;363(6433):eaau8861. doi:10.1126/science.aau8861.

[49] Genestier A, Duret L, Lartillot N Datasets and codes used in the paper "Bridging the gap between the evolutionary dynamics and the molecular mechanisms of meiosis : a model based exploration of the PRDM9 intra-genomic Red Queen". Zenodo. 2024. doi:10.5281/zenodo.10840159