Dialect differences correlate with environment in migratory coastal White-crowned Sparrows

Jiaying Yang[1], Bryan C. Carstens[1], Kaiya L. Provost[1*]

1: Department of Evolution, Ecology and Organismal Biology, The Ohio State University, Columbus, Ohio, USA.

* Corresponding author: provost.27@osu.edu

ACKNOWLEDGEMENTS
We thank the Carstens Lab past and present, as well as D. Nicholson and A. Delgado for helpful feedback.
Funding statement: KLP was funded by NSF DEB-2016189. BCC was funded by NSF DBI-1661029, DEB-2016189, NSF DEB-1831319, and DBI-1910623. JY was funded by the Undergraduate Diversity at Evolution program.
Author contributions: Experimental design: JY, KLP, BCC. Data collection: JY. Methods development: KLP, BCC. Analyses: JY, KLP. Writing: JY. Figures: JY, KLP. Editing: JY, KLP, BCC.
Data depository: Data will be available upon acceptance on Dryad. Scripts will be available on GitHub.

ABSTRACT
Vocalization behavior in birds, especially songs, strongly affects reproduction, with differences in vocalizations between populations often interpreted as evidence of divergence. Previous work has demonstrated that the White-crowned Sparrow (*Zonotrichia leucophrys*) has multiple dialects, including between neighboring (and genetically distinct) subspecies *nuttalli* and *pugetensis*. However, it is unknown whether the divergence in their songs correlates to environmental or geographical factors. Previous work has relied on limited amounts of data because traditional methods to study bird songs rely on the manual annotation of song spectrograms into individual syllables. Here we explore the performance of automated machine learning methods of song annotation, which can process large datasets more efficiently, paying particular attention to the question of subspecies differences. We utilize a recently published artificial neural network to automatically annotate hundreds of White-crowned Sparrow vocalizations across two subspecies. By analyzing differences in syllable usage and composition, we find that *nuttalli* and *pugetensis* have significantly different dialects. Our results are consistent with the interpretation that these differences are caused by the reduction in the number of shared syllables in the White-crowned Sparrow repertoire, as reported by some previous research. Our large sample size enables us to demonstrate that divergence in dialects is correlated with environmental difference, but not with geographical distance, and is also associated with migratory status. Our findings support the hypothesis that the evolution of vocalization behavior is affected by both biotic and abiotic environments, in addition to population structure.

Keywords: Bird songs; Bioacoustics; Machine learning; *Zonotrichia leucophrys*.

LAY SUMMARY
- Bird song is an important behavior because it is important in bird communication and reproduction.

•   White-crowned Sparrows in western North America are known to use different songs
along their range, but it is unknown if those songs vary due to the environment.
•   We used machine learning to analyze these songs and found that populations of White-
crowned Sparrows can be differentiated based on their song dialects.
•   Environmental factors during the summer months exert a greater influence on song
evolution in migratory subspecies.
53
INTRODUCTION
Vocalization is the primary method of bird communication (Catchpole and Slater 2008, Brumm
and Zollinger 2013). Unlike other sensory communication channels (e.g., visual, chemical,
tactile), vocalizations can transmit long distances and have a fast rate of evolutionary change,
due in part to cultural transmission (Slater 1986, Slater 1989, Slabbekoorn and Smith 2002a,
Podos and Warren 2007, Derryberry 2009; but see Mason et al. 2017). In birds, vocalization
behavior can be distinguished into two types: calls and songs (Catchpole and Slater 2008). Bird
calls are short and simple notes that are produced by both sexes year-round and are usually used
for alarms and within-flock contact (Evans et al 1993, Marler 2004, Kondo and Watanabe 2009).
In contrast to calls, bird songs are more complex and melodious. While it was previously thought
that songs were mostly produced by males during the breeding seasons, with common functions
of territory defense and mating (Kroodsma and Byers 1991), more recent observations and
research shows that females also produce songs, and songs are being produced also in non-
breeding contexts (DeWolfe et al 1989, Baptista et al 1993, DeWolfe and Baptista 1995, Odom
et al 2014, Webb et al 2016, Riebel et al 2019).
Song evolution is influenced by biotic and abiotic factors. Vocal behaviors vary among
species and play a role in species recognition (Beer 1971, Emlen 1972, Baker et al. 1981b,
Freeberg et al 1995). Even closely related populations can have different songs (i.e., dialects)
that they can use to tell each other apart (Cunningham and Baker 1983, Thompson et al. 1993,
MacDougall-Shackleton 2001). Many taxa are functionally reproductively isolated due to song
differentiation (e.g., Provost et al. 2021), potentially contributing to speciation of distinct
populations. These differences can also arise due to evolution of other traits like body size and
beak shape, which indirectly influence the sounds a bird can produce (Nowicki et al. 1992;
Podos and Nowicki 2004a, 2004b, Derryberry 2009). Within species, sexual selection via same-
sex competition and mate choice can lead to multiple different outcomes (reviewed by Searcy
and Andersson 1986, Nowicki et al 1998). Another biotic factor influencing song is song
learning, which is present in three orders of birds (Apodiformes, Psittaciformes, and
Passeriformes). In the aptly named oscine songbirds (Order: Passeriformes, Suborder: Passeri),
songs are learned by juvenile individuals listening to the singing adults immediately around
them. As such, geographically restricted song variants emerge through time (Podos and Warren
2007), resulting in a pattern analogous to isolation-by-distance (Nelson et al 2004). Rarely, older
individuals can learn new variants, for example mimics (Robinson 1974) and brood parasites
(Rothstein and Fleischer 1987, Warren 2002)
In contrast to these biotic factors, the abiotic environment can impact the evolution of song
by influencing the efficacy of its transmission. Lower frequencies and slower notes transmit
more effectively through denser vegetation, and birds living in those habitats have songs with
those characteristics (Slabbekoorn and Smith 2002a, Naguib 2003, Kopuchian et al. 2004,
Seddon 2005, Baker 2006, van Dongen and Mulder 2006, Derryberry 2009). Likewise, birds in
urbanized habitats sing at higher frequencies to prevent overlap with low-frequency

anthropogenic noise (Derryberry et al. 2016, Derryberry et al. 2020). In sum, differences in song
between bird taxa can result from any combination of these biotic and abiotic factors.
95
White-crowned Sparrow
One well-studied songbird is the White-crowned Sparrow (*Zonotrichia leucophrys*), which has
multiple dialects across North America (Baptista 1989). This species is known to sing year-
round, not just during the breeding season, making them good candidates for understanding
environmental impacts on their sounds (Blanchard 1941, Baptista 1974, Brenowitz et al. 1998).
Non-breeding singing in *pugetensis* appears to be associated with flock hierarchy formation as
well as hormone stimulation (DeWolfe and Baptista 1995), in contrast with *nuttalli* which sings
due to a protracted breeding season.
They are known to have divergent dialects across multiple different populations, particularly
at the subspecies level (Petrinovich and Patterson 1981, Baker et al. 1984, Lien and Corbin
1990). We focus here on *Zonotrichia leucophrys nuttalli* and *Z. l. pugetensis* (hereafter *nuttalli*
and *pugetensis*, respectively). These subspecies have divergent dialects and population genetic
differentiation (Lipshutz et al. 2017), suggesting that they are distinct evolutionary units. In
addition, previous estimates show that dialects present in *pugetensis* are more wide-ranging than
in *nuttalli* (DeWolfe and Baptista 1995). Songs help to identify these two subspecies, both for
researchers and for the birds themselves. The dialects of these birds vary such that the subspecies
can tell each other apart, suggesting that reproductive isolation is evolving. The subspecies have
an area of hybridization in Mendocino County, CA, where intermediate individuals are found
(Lipshutz et al 2016). Generally, subspecies *nuttalli* (Figure 1A) and *pugetensis* (Figure 1B)
differ in the length of introductory whistle notes, presence of buzz notes, and length/presence of
terminal trill notes (Dunn et al. 1995; Baptista and King 1980), though variation exists within
populations and through time. Song is directly implicated in reducing gene flow between these
populations, which have been at least partially genetically isolated for ~45,000 years (Lipshutz et
al 2016). The different dialects of both *nuttalli* and *pugetensis* also show a geographic pattern
within the subspecies, which are proposed to have been formed by dialect formation, either
limited or biased dispersal, morphological variation, and/or habitat (Marler and Tamura 1962,
1964, Barker and Mewaldt 1978, Nelson 2000, Nelson et al. 2004, Kroodsma et al. 1985,
reviewed by Podos et al. 2007, Derryberry 2009).
124
125

Figure 1. Spectrogram of (A) *nuttalli,* BLB11552, and (B) *pugetensis,* BLB07718, songs with manually annotation in red boxes. The X axis shows the duration time, in seconds, and the Y axes shows the frequency, in kilohertz. Higher frequencies represent higher pitch; and darker color represents higher amplitude i.e., louder sound. Each red box represents one annotated syllable, with the width representing the duration while the height indicates the differences of maximum and minimum frequency i.e., bandwidth. Note that background sounds are not annotated. Above certain notes are annotated the note type: W = whistle, BZ = buzz, and TR = trill of simple notes.

Though the two subspecies are phenotypically similar, with only slight differences in body size (Dunn et al. 1995), they do vary in their migratory behavior and geographic distributions. Subspecies *nuttalli* is a non-migratory subspecies that is distributed in coastal California (from ~40.4 to 34.3 degrees latitude; Banks 1963; Figure 2). In contrast, *pugetensis* is a migratory subspecies that is distributed from British Columbia to northwest California during summer, both coastal and inland, and expands to the southern part of its range through southern California during winter (Banks 1963, Patten et al. 2003, Baptista 1974, Heinemann 1981; Figure 2). Subspecies *pugetensis* thus shares its wintering grounds with three other subspecies: *nuttalli*, which is phenotypically similar, as well as *oriantha* and *gambelii*, which are distinguished by

differences in color of the beak and the lores (Welke et al 2021, Lisovski et al 2019). Moreover, *pugetensis* has clinal dialects along its range during the breeding season, with populations having the same dialect tending to migrate to the same location in the non-breeding season (Nelson 2000, Nelson et al 2004, Baptista 1974, Heinemann 1981) despite those locations not being segregated by song dialect (DeWolfe and Baptista 1995).

Figure 2. The geographical range of *Zonotrichia leucophrys* recordings used in this study. The sedentary *nuttalli* individuals are blue circles and the migratory *pugetensis* individuals are red triangles. Individuals in western California and Utah were outliers (see main text). Map below shows species distribution in all year (purple) and non-breeding season (light blue) (BirdLife International and Handbook of the Birds of the World 2021; Welke et al. 2021, Weckstein et al. 2001, Grinnell and Miller 1944, Taylor et al 2021, Cortopassi and Mewaldt 1965). Note that this map includes distributions for other subspecies (*gambelii* and *oriantha*). Individuals are separated by Summer (May to August, left), Winter (September to April, center) and season unknown (right).

Though there is compelling evidence that the divergence of White-crowned Sparrow songs is caused by sexual selection, there is less evidence on whether the evolution of vocalization behavior is correlated to the abiotic environment, such as climate, or geographic distance as a proxy for cultural evolution. Distinct differences in the histories of these two subspecies may be mediating environmental adaptation. Traditionally, *pugetensis'* breeding and wintering grounds were thought to be distinct, with the breeding not occurring south of Oregon and wintering only occurring in California (DeWolfe and Baptista 1995). However, *pugetensis* experienced a range shift east through the Cascade Mountains and North into Alaska because of habitat change and logging regimes (Welke et al 2021, Hunn and Beaudette 2014, Gibson et al 2013), and birds identified as *pugetensis* are found north of their supposed breeding grounds during the winter, though typically the reverse is not true (Sullivan et al, 2009; GBIF 2023). As such, it is likely that these shifts in location may be due to *pugetensis* being sensitive to environmental changes. In this paper, we use these White-crowned Sparrow subspecies to investigate whether the divergence in songs within and between subspecies is correlated to these abiotic factors. We

177 hypothesize that the different dialects are correlated to climates and geographical differences
178 between the two subspecies. We also hypothesize that *pugetensis* may have a greater impact of
179 environment on their songs compared to *nuttalli*, and that impact should be more apparent during
180 the summer months.
181
182 **METHODS**
183 All work was done on MacBook Pro running macOS Ventura (13.0) with an 8 core Apple M1
184 chip and 8 GB of RAM.
185
Recording data and metadata
We downloaded our sound data from two sources: the Borror Laboratory of Bioacoustics (BLB;
blb.osu.edu) directly from the database and the Xeno-Canto database (X-C; xeno-canto.org) by
using *query_xc* function in the *warbleR* package version 1.1.26 (Araya-Salas and Smith-Vidaurre
2017) in R version 4.0.3 (R Core Team 2020). We used all recordings of *Z. l. pugetensis* and *Z. l.*
nuttalli in the BLB. On X-C, we restricted recordings to be graded as Quality "A." We
transformed the X-C recordings from MP3 format to WAV format by using the *readMP3*
function in the *tuneR* package version 1.3.3 (Ligges et al. 2018) in R. We also converted
recordings to have a sample rate of 48,000 Hz if needed using the *resamp* function in the
seewave package version 2.2.0 (Suer et al 2008) in R. This resulted in 315 *nuttalli* recordings and
360 *pugetensis* recordings from BLB, plus four *nuttalli* recordings and 38 *pugetensis* recordings
from X-C. Recordings were made between 1965 and 2005 and included metadata subspecies
identification, date, locality, and recordist.
Both databases contained recordings from the region of interest that were not identified to
subspecies; we assigned them using latitude as a criterion for breeding-season birds. The
subspecies identification followed Lipshutz et al (2017). We assigned all summertime
"unknown" individuals north of 40.544 and south of 48.461 to be *pugetensis*, and all
summertime "unknown" individuals south of 38.419 and north of 37.803 to be *nuttalli*, because
they do not have overlapping distribution within these regions during summer. Based on the
distribution, we reassigned some mis-identified birds in the original datasets. We did not assign
"unknown" individuals between the two regions because the two subspecies intergrade in that
area. Moreover, "unknown" individuals that were further than ~110km (one decimal degree)
from the coast were not classified as either subspecies as they likely belonged to subspecies
oriantha or *gambelii*. For non-breeding seasons birds, we did not assign "unknown" individuals
due to substantial amounts of overlap between *nuttalli, pugetensis, gambelii,* and *oriantha* which
all overwinter through the region.
Our recordings were mostly restricted to the west coast of the United States and Canada.
However, there were two localities that were further inland and at higher elevation (above 1900
m). As these localities were likely to influence our environmental analyses, we removed them
from our downstream analyses (but see Supplemental). Our final dataset included 238 recordings
from summer (111 *pugetensis*, 127 *nuttalli*), 461 recordings from winter (276 *pugetensis*, 185
nuttalli), and 16 with no date (11 *pugetensis*, 5 *nuttalli*) for a total of 715.
218
Manual Annotation
We followed the methodology of Provost et al. (2022) to annotate our samples. 201 (28%) of
these samples were annotated for that study. For manual annotation, we input the recording to
Raven Pro version 1.6.1 (K. Lisa Yang Center for Conservation Bioacoustics at the Cornell Lab

of Ornithology 2019) and generated a spectrogram using a Fast Fourier Transform size of 512
samples. We annotated the spectrogram by drawing boxes that show the maximum and minimum
values of the frequency and time of each syllable. All selections were output as TXT files. We
then split the original recordings and their corresponding annotation files into subsets that had
approximately equal amounts of annotated syllables and silence using previously generated
scripts (Provost et al. 2022). Although TweetyNet can identify and segment multiple kinds of
syllables, we annotated all syllables as "1".
230
Using Machine Learning for Bioacoustics
Bird songs have been studied for a long time with many methods for qualitatively and
quantitatively describing their sounds having been developed. Traditional methods require
researchers to analyze and measure song parameters manually. Since the mid 20th century, the
most popular methods of analysis use spectrograms, which display frequencies through time
(Borror and Reese 1953; Figure 1). In the digital era, these spectrograms are generated digitally
by using fast Fourier transforms (e.g., via Raven; K. Lisa Yang Center for Conservation
Bioacoustics Cooley and Tukey 1965). Many analyses require every individual syllable or note
on the spectrogram to be identified and annotated, or segmented, for downstream processing.
Segmentation is time intensive and the amount of data that can be processed is limited (Swiston
et al. 2009).
Recently, machine learning and artificial intelligence techniques have been used broadly in
biological science (e.g., Pelletier and Carstens 2018, Cushman et al 2017, Viener et al 2022,
Schrider and Kern 2017, Buxton et al 2018, Clapham et al 2020). Machine learning has also been
applied to syllable segmentation with high accuracy via the neural network TweetyNet (Cohen et
al. 2020). Originally trained on single species, the algorithm has since been applied to a suite of
birds (Provost et al 2022) with similarly high performance, allowing for the use of larger
datasets. Although the segmentation itself has been sped up, it is unknown whether this and other
automatic segmentation methods effectively recapitulate known differences in populations. Here,
we examine whether we can apply automatic syllable recognition to the different dialects of
White-crowned Sparrow songs.
To automatically annotate the White-crowned Sparrow songs, we used the TweetyNet model
trained on *Zonotrichia leucophrys* from Provost et al. (2022). The latter analyses included
examples from all five subspecies. We then fine-tuned this model by continuing to train it on just
data from *nuttalli* and *pugetensis*. We used vak version 0.6.0 and TweetyNet version 0.8.0 for
that and we used identical parameters to (Provost et al. 2022). After being fine-tuned, we had the
model predict the locations of syllables for all data, including data that TweetyNet had never
seen before. We assessed model performance (syllable error rates, accuracy, precision, recall,
and f-score) for each subspecies.
260
Summary statistics
Once we had syllables segmented for all recordings, we extracted bioacoustic data for each
annotation. Data were calculated using the *spectro_analysis* function in the *warbleR* package in
R. The data extracted from syllables include bandwidth (the difference between highest
frequency and lowest frequency of the syllable), time duration (the length of the syllable), center
frequency (central frequency between the upper and lower cutoff frequencies), average slope
(slope of the change in dominant frequency), and inflection points (the local extrema). For each
recording, we calculated the centroid value across all syllables. We then calculated a distance

between centroids between all individuals within and between subspecies, which we used as a
representation of song distance.
271
Principal Components Analysis and ANOVAs
We performed a principal components analysis (PCA) on bandwidth, time duration, center
frequency, average slope, and inflection points of syllables. To do this, we used the *prcomp*
function in the *stats* package version 4.1.2 in R. We centered and scaled all data. The PCA
results were used to compare differences between the subspecies. We used an ANOVA on both
the raw statistics and results of PCA to test the differences between the two subspecies and two
seasons (winter and summer) using the *aov* function in the *stats* package in R. We analyzed
significance for this and all other analyses using an alpha value of 0.05.
280
Multiple Matrix Regression with Randomization
To determine whether differences in songs were being driven by differences in either the abiotic
environment or geographic distance, we used multiple matrix regression with randomization
analysis (MMRR) in R (Wang 2020). This technique allows one to correct correlations between
environmental features and geography. To calculate geographic distances as a proxy for cultural
evolution, we calculated the Euclidean distance between individuals with respect to latitude and
longitude. For environmental differences, we downloaded 19 climate variables from WorldClim
(Fick and Hijmans 2017) and extracted values at each locality in our dataset. We then calculated
Euclidean distance between individuals with respect to those values. MMRR models were
created such that we predicted song divergence from geographic distances and environmental
differences simultaneously (i.e., song divergence ~ geographic distance + environmental
distance). Note that there were several *nuttalli* individuals across these analyses for whom either
geographic or environmental information was lacking; as such, our dataset for the MMRR
analyses included 398 *pugetensis* and 317 *nuttalli*.
Because *pugetensis* is a subspecies that migrates through the year, we separated our data by
seasons: summer (May to August, when *pugetensis* are in their breeding range) and winter
(September to April, when *pugetensis* are in their non-breeding range; Chilton et al. 2020). We
performed these MMRR analyses described above with only the winter and summer datasets
separately because the different ranges due to migration of *pugetensis* may result in variation in
geographical distance and ecological distance.
301
RESULTS
First, we analyzed how the machine learning model performed in syllable annotation. These
models had high performance across both subspecies. The syllable error rate of our trained
machine learning model's annotations was 15% for *nuttalli* and 4% for *pugetensis* (Table 1),
such that for both subspecies, the model predicted more syllables than were present. Accuracy,
precision, recall, and F-score were all above 0.91 for both subspecies, with *nuttalli* having 2–3%
better performance than *pugetensis* (Table 1).
309
310 **Table 1**. The machine learning model showed a high accuracy in annotating the syllables of both
311 *pugetensis* and *nuttalli* for all data.

Subspecies	Syllable Error Rate	Accuracy	Precision	Recall	F-score
nuttalli	15%	0.96	0.96	0.93	0.95
pugetensis	4%	0.93	0.94	0.91	0.92

312
PCA
We found that our PCA on syllable variables explained a high proportion of the data, with the five PCs explaining 45%, 20%, 17%, 13%, and 3% of variation, respectively (Table 2). Syllables with high PC1 had wide bandwidth, short time, low frequency, few inflection points, and negative slopes (Table 2). Subspecies *nuttalli* and *pugetensis* used different dialects and songs were different across seasons (Table 3). Subspecies varied in PC1 (mean *pugetensis* = -0.16, mean *nuttalli* = 0.21) and PC2 (mean *pugetensis* = 0.25, mean *nuttalli* = -0.32) reflecting significant differences in center frequency (mean *pugetensis* = 3.12, mean *nuttalli* = 2.92), inflection points (mean *pugetensis* = 13.13, mean *nuttalli* = 10.85), and slope (mean *pugetensis* = -1.06, mean *nuttalli* = -18.35). Across seasons, songs varied in PC1 (mean winter = 0.13, mean summer = -0.23, mean unknown = -0.36) and PC4 (mean winter = 0.03, mean summer = -0.10, mean unknown = 0.41), reflecting differences in duration (mean winter = 0.16, mean summer = 0.18, mean unknown = 0.16) and inflection points (mean winter = 11.50, mean summer = 13.26, mean unknown = 13.07). PC3, PC5, and bandwidth were not different across season or subspecies.

Table 2: Principal component importance and rotation of individual PC axes for song across five summary statistics of song properties.

		PC1	PC2	PC3	PC4	PC5
Importance	Standard deviation	1.50	1.01	0.92	0.80	0.43
	Proportion of variance	0.45	0.20	0.17	0.13	0.03
	Cumulative proportion	0.45	0.65	0.83	0.96	1.00
Rotation	Bandwidth	0.37	0.34	0.65	-0.55	0.12
	Duration	-0.52	-0.05	-0.15	-0.68	-0.48
	Center	-0.42	-0.07	0.7	0.42	-0.35
	Inflection	-0.61	0.08	0.07	-0.09	0.77
	Slope	-0.14	0.93	-0.19	0.2	-0.17

Table 3: Differences between season (winter-summer) and subspecies (*pugetensis-nuttalli*) using ANOVAs while controlling for both. Df = 1 for all comparisons. Note that unknown season individuals were removed. SSQ = sum of squares. MSQ = mean sum of squares. F = F-statistic. P = p-value.

		Winter-Summer				*pugetensis-nuttalli*			
		SSQ	MSQ	F	P	SSQ	MSQ	F	P
Raw	Bandwidth	0.1	0.091	1.08	0.74	1.4	1.36	1.61	0.20
	Center	1.7	1.715	2.52	0.11	7.2	7.17	10.54	**0.00***
	Duration	0.094	0.093	13.12	**0.00***	0.003	0.00	0.45	0.49
	Inflection	482	482.3	10.08	**0.00***	865	865	18.09	**0.00***
	Slope	167	167	0.207	0.64	53005	5305	65.57	**0.00***
PCA	PC1	21.1	21.1	9.8	**0.00***	24.1	41.1	11.30	**0.00***
	PC2	0.1	0.06	0.059	0.80	59.8	59.8	63.4	**0.00***
	PC3	0.3	0.26	0.3	0.58	3.2	3.18	3.74	0.05
	PC4	3	3.02	4.73	**0.03***	2	1.97	3.07	0.07
	PC5	0	0	0	0.99	0.25	0.25	1.33	0.24

MMRR

We found that year-round, environmental (climate) distance explained song divergence when looking at both subspecies together, but geographic distance did not (R^2 = 0.01, climate p = 0.02, geography p = 0.69; Table 4). Neither climate nor geography significantly explained song divergence in winter alone (R^2 = 0.00, climate p = 0.13, geography p = 0.13) or summer alone (R^2 = 0.00, climate p = 0.75, geography p = 0.68) for both subspecies together. Subspecies *nuttalli* showed no relationship between song divergence and climate or geography for any season (climate p > 0.20, geography p > 0.26). In contrast, subspecies *pugetensis* showed significant impact of geography year-round (R^2 = 0.03, climate p = 0.00, geography p = 0.23) and both geography and climate in summer only (R^2 = 0.14, climate p = 0.01, geography p = 0.02). There was no relationship in winter alone (R^2 = 0.00, climate p = 0.64, geography p = 0.52). These patterns cannot be explained solely by differences in year recorded (see Supplemental).

Table 4: MMRR analysis shows that divergence in songs of *pugetensis* and *nuttalli* is strongly correlated with environmental differences.

Subspecies	Season	N	R²	F	T-values Geo.	T-values Env.	Coefficients Geo.	Coefficients Env.	P-Values Geo.	P-Values Env.	P-Values Overall
Both	Year	694	0.01	980	3.48	25.22	0.01	0.08	0.69	**0.02***	**0.00***
	Winter	456	0.00	92.5	-13.13	12.89	-0.07	0.07	0.13	0.13	0.18
	Summer	234	0.00	3.83	2.77	-2.02	0.02	-0.02	0.68	0.75	0.91
N	Year	305	0.00	61.7	-4.88	10.60	-0.03	0.07	0.60	0.20	0.44
	Winter	181	0.01	82.8	6.65	1.31	0.09	0.02	0.28	0.85	0.13
	Summer	124	0.00	15.7	5.48	-5.11	0.11	-0.10	0.26	0.25	0.43
P	Year	389	0.03	1207	37.46	-9.94	0.22	-0.06	**0.00***	0.23	**0.00***
	Winter	275	0.00	12.2	-4.95	3.72	-0.04	0.03	0.52	0.64	0.82
	Summer	110	0.14	504	-20.67	30.96	-0.42	0.62	**0.02***	**0.01***	**0.02***

DISCUSSION

Our analysis demonstrates that the trained machine learning model by TweetyNet has high
accuracy with low error rates and can annotate *nuttalli* and *pugetensis* songs. Our results suggest
that the annotation by the machine learning model is as reliable as manual annotations and
produces comparable data. Given its computational efficiency, machine learning can be a useful
tool to automatically annotate bird songs and extract data because it can be used to process large
datasets quickly and accurately.

We found significant differences in the syllables used in the songs of *nuttalli* and the songs of
pugetensis, which corroborates previous evidence that these two taxa have divergent dialects
(e.g., Lipshutz et al. 2017). Our bioacoustic metrics found that on average, *nuttalli* syllables were
lower in pitch, had fewer inflections, and had more negative slopes compared to *pugetensis*. At
present our analyses cannot distinguish whether this is due to subspecies using different syllables
or the same syllables at different frequencies. Preliminary analyses of syllable shape suggest that
the differences are mediated by use of different syllables (KLP pers. comm.) but further
investigation is needed to confirm.

We also found significant differences between the seasons. Syllables used in summer were
on average longer and had more inflection points, suggesting higher complexity. Although more
complex songs have been found to be sexually selected for in other species (e.g., Catchpole
1987), complexity itself is likely a result of many competing selective pressures within and
between species including migratory behavior (Read and Weary 1992). In *Zonotrichia*
leucophrys, different parts of the song appear to have different function (Nelson and Poesel
2007): introductory whistles (as well as buzzes) are highly conserved and serve as species
identifiers, terminal trills are moderately conserved and indicate dialect, and other notes in the
song are highly variable and indicate individual identity. At the same time, this species is known
to modulate its song with respect to the environment, being especially impacted by noise
transmission (Derryberry et al 2009, Luther et al 2016, Derryberry et al 2020). Our findings
indicate that White-crowned Sparrows use different aspects of their repertoire for different
functions during the breeding and non-breeding seasons.

The differences in dialect between subspecies and seasons are compounded by the fact that
the two subspecies have different responses to the environment, and that these are dependent on
the time of year analyzed. The migratory *pugetensis* shows a high impact of environment and
geographic distance on songs recorded in summer, but not winter. Year-round, geographic
distances remain significant predictors of song divergence in this subspecies, suggesting that
winter and summer songs may be different. In contrast, the sedentary *nuttalli* shows no
relationships between environment or geographic distance with respect to song. The evidence
suggests that overall, the migratory subspecies *pugetensis* shows higher impact of environment
during the breeding season. This may be driving the patterns where both species are combined in
those seasons; alternatively, the two subspecies may still be adapted to their relatively different
environments. It is likely that these factors (i.e., migration and seasonality) have contributed to
the evolution of dialect differences.
We performed a *post-hoc* analysis of the environments that each of the subspecies occupy
during different seasons (see Supplemental) and found that these environments are significantly
different based on climate data. Year-round, *nuttalli* occupies warmer and drier areas than
pugetensis does in the summer. In contrast, wintering *pugetensis* experience nearly the full
yearly range of temperature and precipitation across both subspecies, and overall experience
more variable and more seasonal climates than *nuttalli*. This evidence indicates that *pugetensis*
has a higher sensitivity to the environment. A synthesis across multiple birds hypothesizes that
migration occurs as a mechanism to avoid harsh winters and exploit resource surpluses
(Somerville et al 2015, Global Ecology and Biogeography); given that the range of *pugetensis* is
wide-spread and has been changing in response to habitat disruption (Welke et al 2021, Hunn
and Beaudette 2014, Gibson et al 2013), we hypothesize that *pugetensis* as a subspecies may be
more sensitive to fluctuations in habitat and climate, leading to more sensitivity in their songs. In
addition, because *pugetensis* is known to have dialects ranging over larger areas that encapsulate
more variable climates (DeWolfe and Baptista, 1995), it is possible that the selection pressures to
adapt song to environment is higher across the range of this subspecies.
410
Caveats
Several caveats are worth documenting to place our results in the appropriate context. All
analyses shows that geographic distance is not correlated to the divergence in song across these
two subspecies, which is not consistent with our hypothesis. Some explanations are as follows.
Firstly, although the number of recordings of *pugetensis* in non-breeding habitats is large, they
do not cover the entire projected breeding range. Second, since our recordings are all from the
BLB and X-C database from 1965 to 2005, we do not know if songs have changed during this
period and so we may have quite different songs in the same area. Over the time scale of the
study, White-crowned Sparrows are known to have demonstrably changed their song dialects
(e.g., Derryberry 2009). However, even if we control for the year of recording, we still find
significant relationships with spatial factors like geography and environment (see Supplemental).
Third, our analysis only used each syllable's characteristics but did not analyze the type, syntax,
or order of syllables. It is possible that they used the same syllables but different orders in some
dialects. Furthermore, because White-crowned Sparrows learn their songs not only directly from
their parents but also from neighboring community (Cunningham and Baker 1983) that with
same or similar songs, it is unknown that whether some juveniles learned songs from the other
subspecies, especially in the hybrid zone, or some late-hatched *nuttalli* may have chance to
expose to the environment that migrated *pugetensis* already arrived in their winter habitats. It is

hard to answer these questions with only the recordings and data of their locality and habitats.
We need more research and specific data to solve the problems. However, the strong positive
relationship between ecological differences and song dialects in winter corresponds with our
original hypothesis. However, in this research, we only analyzed the numeric data of the features
of bird songs and the climate of each locality, but we did not investigate the other properties of
each habitat, such as the habitats' type. We still need further research on whether the abiotic
environments directly or indirectly affect the evolution of bird songs and what is the mechanism
of it.
In conclusion, using machine learning for automatic analyses, we evaluated divergence
within and between two subspecies of the White-crowned Sparrow. Our results indicated that
Zonotrichia leucophrys pugetensis and *Z. l. nuttalli* have significantly different songs from each
other, and both sing different songs in breeding and non-breeding contexts. The divergence in
dialects correlate with spatial factors such as geographic distance and climate, but subspecies
identity and season mediate the response.
443
LITERATURE CITED

Araya-Salas, M., and G. Smith-Vidaurre (2017). "warbleR: an r package to streamline analysis
of animal acoustic signals." https://dx.doi.org/10.1111/2041-210X.12624.
Baker, M. C. (2006). Differentiation of mating vocalizations in birds: Acoustic features in
mainland and island populations and evidence of habitat-dependent selection on songs.
Ethology 112:757-771. https://doi.org/10.1111/j.1439-0310.2006.01212.x
Baker, M. C., D. B. Thompson, and G. L. Sherman (1981b). Neighbor-stranger song
discrimination in White-crowned Sparrows. Condor 83:265-267.
Baker, M. C., M. A. Cunningham, and A. D. Thompson Jr. (1984). Cultural and genetic
differentiation of two subspecies of White-crowned Sparrow. Condor 86:359-367.
Baker, M. C., and M. A. Cunningham (1985). The Biology of Bird-Song Dialects. Behavioral
and Brain Sciences 8:85-100. https://doi.org/10.1017/S0140525X00019750
Baker, M. C., and L. R. Mewaldt (1979). The use of space by White-crowned Sparrows: juvenile
and adult ranging patterns and home range versus body size comparisons in an avian
granivore community. Behavioral Ecology and Sociobiology 6:45-52.
Banks, R. C. (1963c). Geographic variation in the White-crowned Sparrow *Zonotrichia
leucophrys*. University of California Publications in Zoology 70:1-123.
Baptista, L. F. (1974). The Effects of Songs of Wintering White-crowned Sparrows on Song
Development in Sedentary Populations of the Species. Zeitschrift Für Tierpsychologie,
34:147-171. https://doi.org/10.1111/j.1439-0310.1974.tb01794.x
Baptista, L. F. (1989). The best studied songbird: White-crowned Sparrows in the wild and in the
laboratory. Birds International 1:10-21.
Beer, C. G. (1971). Individual Recognition of Voice in the Social Behavior of Birds. Advances
in the Study of Behavior, 3:27-74. https://doi.org/10.1016/S0065-3454(08)60154-0
Baptista, L. F. (1989). The best studied songbird: White-crowned Sparrows in the wild and in the
laboratory. Birds International 1:10-21.
BirdLife International and Handbook of the Birds of the World (2021) Bird species distribution
maps of the world. Available at http://datazone.birdlife.org/species/requestdis.
Borror, D. J., and C. R. Reese (1953). The analysis of bird songs by means of a vibralyzer. The
Wilson Bulletin 65:271-276.

Brenowitz, E. A., L. F. Baptista, K. Lent, and J. C. Wingfield (1998). Seasonal plasticity of the
song control system in wild Nuttall's white☐crowned sparrows. Journal of neurobiology,
34:69-82.
Brumm, H., and S. A. Zollinger (2013). Avian Vocal Production in Noise. In Animal
Communication and Noise. (Brumm, H., Editor). Animal Signals and Communication 2:187-
227. https://doi.org/10.1007/978-3-642-41494-7_7
Catchpole, C. K., and P. J. B. Slater (2008). Bird Song: Biological Themes and Variations.
Cambridge University Press, Cambridge, UK.
Catchpole, C. K. (1987). Bird song, sexual selection and female choice. In Trends in Ecology
and Evolution 2:94-97. https://doi.org/10.1016/0169-5347(87)90165-0
Chilton, G., M. C. Baker, C. D. Barrentine, and M. A. Cunningham (2020). White-crowned
Sparrow *(Zonotrichia leucophrys)*, version 1.0. In Birds of the World (A. F. Poole and F. B.
Gill, Editors). Cornell Lab of Ornithology, Ithaca, NY, USA.
https://doi.org/10.2173/bow.whcspa.01
Cohen, Y., D. A. Nicholson, A. Sanchioni, E. K. Mallaber, V. Skidanova, and T. J. Gardner
(2022) Automated annotation of birdsong with a neural network that segments spectrograms.
eLife 11:e63853. https://doi.org/10.7554/eLife.63853
Cooley, J. W., and J. W. Tukey (1965). An algorithm for the machine calculation of complex
Fourier series. Mathematics of computation 19:297-301.
Cortopassi, A. J., and L. R. Mewaldt (1965). The circumannual distribution of White-crowned
Sparrows. Bird-banding 36:141-169.
Cunningham, M. A., and M. C. Baker (1983). Vocal learning in White-crowned Sparrows:
sensitive phase and song dialects. Behavioral Ecology and Sociobiology 13:259-269.
https://doi.org/10.1007/BF00299673
Derryberry, E. P. (2009). Ecology shapes birdsong evolution: Variation in morphology and
habitat explains variation in white-crowned sparrow song. American Naturalist 174:24-33.
https://doi.org/10.1086/599298
Derryberry, E. P., R. M. Danner, J. E. Danner, G. E. Derryberry, J. N. Phillips, S. E. Lipshutz, K.
Gentry, and D. A. Luther (2016). Patterns of song across natural and anthropogenic
soundscapes suggest that White-crowned Sparrows minimize acoustic masking and
maximize signal content. PLoS ONE 11:e0154456.
Derryberry, E. P., J. N. Phillips, G. E. Derryberry, M. J. Blum, and D. Luther (2020). Singing in
a silent spring: Birds respond to a half-century soundscape reversion during the COVID-19
shutdown. Science 370:575-579.
Dunn, J. L., K. L. Garrett and J. K. Alderfer (1995). White-crowned Sparrow subspecies:
identification and distribution. Birding 27:182-200.
Evans, C. S., L. Evans, and P. Marler (1993). On the meaning of alarm calls: functional reference
in an avian vocal system. Animal Behaviour 46:23-38.
Fick, S.E., and R.J. Hijmans (2017). WorldClim 2: new 1 km spatial resolution climate surfaces
for global land areas. International Journal of Climatology 37:4302-4315.
Fitch, W. T. (2006). The biology and evolution of music: A comparative perspective. Cognition
100:173-215. https://doi.org/10.1016/j.cognition.2005.11.009
Freeberg, T. M., A. P. King, and M. J. West (1995). Social Malleability in Cowbirds (*Molothrus
ater artemisiae*): Species and Mate Recognition in the First 2 Years of Life. Journal of
Comparative Psychology 109:357-367. https://doi.org/10.1037/0735-7036.109.4.357

Freeman, B. G., J. Rolland, G. A. Montgomery, and D. Schluter (2022). Faster evolution of a
premating reproductive barrier is not associated with faster speciation rates in New World
passerine birds. Proceedings of the Royal Society B: Biological Sciences 289:20211514.
https://doi.org/10.1098/rspb.2021.1514

GBIF.org (28 February 2023) GBIF Occurrence Download https://doi.org/10.15468/dl.vg5xnc

Gil, D., and M. Gahr (2002). The honesty of bird song: Multiple constraints for multiple traits. In
Trends in Ecology and Evolution 17:133-141. https://doi.org/10.1016/S0169-5347(02)02410-
2

Grinnell, J., and A.H. Miller (1944). The distribution of the birds of California. Pacific Coast
Avifauna 27:523-524. Cooper Ornithological Club, Berkeley, CA, USA.

Heinemann, D. (1981). Song Dialects, Migration, and Population Structure of Puget Sound
White-Crowned Sparrows. The Auk 98:512-521.

K. Lisa Yang Center for Conservation Bioacoustics at the Cornell Lab of Ornithology. (2019).
Raven Pro: Interactive Sound Analysis Software (Version 1.6.1) [Computer software]. The
Cornell Lab of Ornithology, Ithaca, NY, USA. Available from
https://ravensoundsoftware.com/.

Kondo, N., and S. Watanabe (2009). Contact calls: Information and social function. In Japanese
Psychological Research 51:197-208. https://doi.org/10.1111/j.1468-5884.2009.00399.x

Kopuchian, C., D. Alejandro Lijtmaer,P. Luis Tubaro, and P. Handford (2004). Temporal
stability and change in a microgeographical pattern of song variation in the rufous-collared
sparrow. Animal Behaviour 68:551-559. https://doi.org/10.1016/j.anbehav.2003.10.025

Kroodsma, D., and J. Verner (1978). Complex Singing Behaviors among *Cistothorus* Wrens.
The Auk 95:703-716.

Kroodsma, D. E., M. C. Bake, L. F. Baptista, and L. Petrinovich (1985). Vocal "Dialects" in
Nuttall's White-Crowned Sparrow. Current Ornithology 2:103-133.
https://doi.org/10.1007/978-1-4613-2385-3_3

Kroodsma, D. E., P. W. Houlihan, P. A. Falleon, and J. A. Wells (1997). Song development by
grey catbirds. Animal Behaviour 54:457-464. https://doi.org/10.1006/anbe.1996.0387

Lein, M. R., and K. W. Corbin (1990). Song and plumage phenotypes in a contact zone between
subspecies of the White-crowned Sparrow (*Zonotrichia leucophrys*). Canadian Journal of
Zoology 68:2625-2629.

Lewis, R. N., M. Soma, S. R. de Kort, and R. T. Gilman (2021). Like Father Like Son: Cultural
and Genetic Contributions to Song Inheritance in an Estrildid Finch. Frontiers in Psychology
12:654198. https://doi.org/10.3389/fpsyg.2021.654198

Lipshutz, S. E., I. A. Overcast, M. J. Hickerson, R. T. Brumfield, and E. P. Derryberry (2017).
Behavioural response to song and genetic divergence in two subspecies of white-crowned
sparrows (*Zonotrichia leucophrys*). Molecular Ecology 26:3011-3027.
https://doi.org/10.1111/mec.14002

Ligges, U., S. Krey, O. Mersmann, and S. Schnackenberg (2018). tuneR: Analysis of Music and
Speech. https://CRAN.R-project.org/package=tuneR.

MacDougall-Shackleton, E. A., and S. A. MacDougall-Shackleton (2001). Cultural and genetic
evolution in mountain White-crowned Sparrows: Song dialects are associated with
population structure. Evolution 55:2568-2575. https://doi.org/10.1111/j.0014-
3820.2001.tb00769.x

Mahler, P. (1970). A comparative approach to vocal learning: song development in White-
crowned Sparrows. Journal of Comparative and Physiological Psychology 71:1-25.

Marler, P. (2004). Bird calls: their potential for behavioral neurobiology. Annals of the New York Academy of Sciences 1016:31-44.

Marler, P., and M. Tamura (1962). Song "dialects" in three populations of White-crowned Sparrows. The Condor 64:368-377. https://www.jstor.org/stable/1365545

Marler, P., and M. Tamura (1964). Culturally transmitted patterns of vocal behavior in sparrows. Science, 146:1483-1486.

Marler, P., P. Mundinger, M. S. Waser, and A. N. N. Lutjen (1972). Effects of acoustical stimulation and deprivation on song development in red-winged blackbirds (*Agelaius phoeniceus*). Animal Behaviour 20:586-606.

Mason, N. A., K. J. Burns, J. A. Tobias, S. Claramunt, N. Seddon, and E. P. Derryberry (2017). Song evolution, speciation, and vocal learning in passerine birds. Evolution 71:786-796. https://doi.org/10.1111/evo.13159

Naguib, M. (2003). Reverberation of rapid and slow trills: implications for signal adaptations to long-range communication. The Journal of the Acoustical Society of America, 113:1749-1756.

Nelson, D. A. (2000). Song overproduction, selective attrition and song dialects in the white-crowned sparrow. Animal Behaviour 60:887–898. https://doi.org/10.1006/anbe.2000.1560

Nelson, D. A., K. I. Hallberg, and J. A. Soha (2004). Cultural evolution of Puget Sound white-crowned sparrow song dialects. Ethology 110:879-908. https://doi.org/10.1111/j.1439-0310.2004.01025.x

Nice, M. M., and B. D. Blanchard (1942). The White-Crowned Sparrows (*Zonotrichia leucophrys*) of the Pacific Seaboard: Environment and Annual Cycle. Bird-Banding 13:96-97. https://doi.org/10.2307/4509737

Nowicki, S., S. Peters, and J. Podos (1998). Song learning, early nutrition and sexual selection in. American Zoologist 38:179-190. https://doi.org/10.1093/icb/38.1.179

Nowicki, S., M. Westneat, and W. Hoese (1992). Birdsong: motor function and the evolution of communication. Seminars in Neuroscience 4:385-390.

Otter, K. A., A. Mckenna, S. E. LaZerte, and S. M. Ramsay (2020). Continent-wide Shifts in Song Dialects of White-Throated Sparrows. Current Biology 30:3231-3235. https://doi.org/10.1016/j.cub.2020.05.084

Päckert, M. (2018). Song: The Learned Language of Three Major Bird Clades. In Bird Species (D. T. Tietze, Editor). Springer, Cham. https://doi.org/10.1007/978-3-319-91689-7_5

Patten, M. A., G. McCaskie, and P. Unitt (2003). Birds of the Salton Sea: Status, biogeography, and ecology. University of California Press, Berkeley, CA, USA.

Petrinovich, L., and T. L. Patterson (1981). The response of White-crowned Sparrows to songs of different dialects and subspecies. Zeitschrift für Tierpsychologie 57:1-14.

Podos, J., and S. Nowicki (2004a). Performance limits on birdsong. Nature's Music 318-342.

Podos, J., and S. Nowicki (2004b). Beaks, adaptation, and vocal evolution in Darwin's finches. Bioscience 54:501-510.

Podos, J., and P. S. Warren (2007). The Evolution of Geographic Variation in Birdsong. In Advances in the Study of Behavior 37:403-458. https://doi.org/10.1016/S0065-3454(07)37009-5

Provost, K. L., J. Yang, and B. C. Carstens (2022). The impacts of transfer learning, phylogenetic distance, and sample size on big-data bioacoustics. PLoS ONE, 17:e0278522. https://doi.org/10.1371/journal.pone.0278522

Provost, K. L., E. A. Myers, and B. T. Smith (2021). Community phylogeographic patterns
reveal how a barrier filters and structures taxa in North American warm deserts. Journal of
Biogeography 48:1267-1283.
Robinson, F. N. (1975). Vocal Mimicry and the Evolution of Bird Song. Emu 75: 23–
27. doi:10.1071/mu9750023
Rothstein, S. I., and R. C. Fleischer (1987). Vocal dialects and their possible relation to honest
status signalling in the Brown-headed Cowbird. The Condor 89:1-23.
R Core Team. (2020). R: a language and environment for statistical computing, version 4.0.3. R
Foundation for Statistical Computing.
Searcy, W. A., and M. Andersson (1986). Sexual selection and the evolution of song. Annual
Review of Ecology and Systematics 17:507-533.
https://doi.org/10.1146/annurev.es.17.110186.002451
Seddon, N. (2005). Ecological adaptation and species recognition drives vocal evolution in
neotropical suboscine birds. Evolution 59:200-215. https://doi.org/10.1111/j.0014-
3820.2005.tb00906.x
Slabbekoorn, H., and T. B. Smith (2002). Bird song, ecology and speciation. Philosophical
Transactions of the Royal Society B: Biological Sciences 357:493–503.
https://doi.org/10.1098/rstb.2001.1056
Slater, P. J. B. (1986). The cultural transmission of bird song. Trends in Ecology and Evolution,
1:94-97. https://doi.org/10.1016/0169-5347(86)90032-7
Slater, P. J. B. (1989). Bird song learning: causes and consequences. Ethology Ecology &
Evolution 1:19-46.
Swiston, K. A., and D. J. Mennill (2009) Comparison of manual and automated methods for
identifying target sounds in audio recordings of Pileated, Pale□billed, and putative
Ivory□billed woodpeckers. Journal of Field Ornithology 80: 42-50.
Thompson, Jr., A. D., and M. C. Baker (1993). Song dialect recognition by male White-crowned
Sparrows: effects of manipulated song components. Condor 95:414-421.
Taylor, R. S., A. C. Bramwell, R. Clemente-Carvalho, N. A. Cairns, F. Bonier, K. Dares, and S.
C. Lougheed (2021). Cytonuclear discordance in the crowned-sparrows, *Zonotrichia*
atricapilla and *Zonotrichia leucophrys*. Molecular Phylogenetics and Evolution 162:107216.
Van Dongen, W. F., and R. A. Mulder (2006). Habitat density, song structure and dialects in the
Madagascar paradise flycatcher Terpsiphone mutata. Journal of Avian Biology 37:349-356.
Wang I.J. (2020) Topographic path analysis for modeling dispersal and functional connectivity:
calculating topographic distances using the topoDistance R package. Methods in Ecology and
Evolution 11:265-272.
Weckstein, J. D., R. M. Zink, R. C. Blackwell-Rago, and D. A. Nelson (2001). Anomalous
variation in mitochondrial genomes of White-crowned (*Zonotrichia leucophrys*) and Golden-
crowned (*Z. atricapilla*) sparrows: Pseudogenes, hybridization, or incomplete lineage
sorting? The Auk 118:231-236.
Welke, C. A., B. Graham, R. R. Conover, J. W. Rivers, and T. M. Burg (2021). Habitat□linked
genetic structure for white□crowned sparrow (*Zonotrichia leucophrys*): Local factors shape
population genetic structure. Ecology and evolution 11:11700-11717.
652
653
654